Discovery of re-purposed drugs that slow SARS-CoV-2 replication in human cells

*Adam Pickard, Ben C. Calverley, Joan Chang, Richa Garva, Yinhui Lu, and *Karl E. Kadler

Wellcome Centre for Cell-Matrix Research, Faculty of Biology, Medicine & Health, University of Manchester, Manchester Academic Health Science Centre, Manchester, M13 9PT UK

* Co-corresponding authors: AP email: adam.pickard@manchester.ac.uk (orcid.org/0000-0001-9757-143X); KEK email: karl.kadler@manchester.ac.uk (orcid.org/0000-0003-4977-4683)

Abstract

Effective vaccines against COVID-19 based on the Spike protein of SARS-CoV-2 have been developed. However, a complementary approach would be to identify drugs that target replication of the virus. The identification of a safe drug that could be repurposed to stop SARS-CoV-2 replication would offer protection against COVID-19 for those awaiting vaccination and may also be effective against emerging vaccine escape variants. Here, using a nano-luciferase tagged version of the virus (SARS-CoV-2-ΔOrf7a-NLuc) to quantitate viral load, we evaluated a range of human cell types for their ability to be infected and support replication of the virus, and performed a screen of 1971 FDA-approved drugs. Hepatocytes, kidney glomerulus, and proximal tubule cells were particularly effective in supporting SARS-CoV-2 replication, which is in-line with reported proteinuria and liver damage in patients with COVID-19. We identified 35 drugs that reduced viral replication in Vero and human hepatocytes when treated prior to SARS-CoV-2 infection and found amodiaquine, atovaquone, bedaquiline, ebastine, LY2835219, manidipine, panobinostat, and vitamin D3 to be effective in slowing SARS-CoV-2 replication in human cells when used to treat infected cells. In conclusion, our study has identified strong candidates for drug repurposing, which could prove powerful additions to the treatment of COVID-19.

INTRODUCTION

The COVID-19 pandemic caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus is having a widespread impact on global health with substantial loss of life. SARS-CoV-2 infection in patients with COVID-19 can result in pulmonary distress, inflammation, and broad tissue tropism. Some infected individuals are asymptomatic whilst others mount an exaggerated immune response, or 'cytokine storm'(Chatenoud et al., 1991; Fajgenbaum and June, 2020), which correlates with multiple organ failure and poor outcome (Ruan et al., 2020). Vaccines have been developed to help protect people from COVID-19 but the emergence of mutational variants, some with increased transmissibility, calls for additional research into alternative treatment and prevention strategies.

An early step in the infection process is interaction of the Spike protein on the surface of the virus with angiotensin-converting enzyme 2 (ACE2) on the surface of the host cell (Li et al., 2003). Using this information, first-generation vaccines have been generated against the Spike protein. SARS-CoV-2 entry can be potentiated by additional host factors including neuropilin-1 (Cantuti-Castelvetri et al., 2020). Once inside, the virus uses components of the host cell to replicate and secrete viral particles (Maier et al., 2016) and disrupt RNA handling and protein translation to suppress host defenses (Banerjee et al., 2020). The different stages of the disease, from the initial infection of host cells through to virus replication and the response (normal or extreme) of the immune system, offer opportunities to identify drugs, treatments and therapies to help stop disease progression. Furthermore, large proportions of the world's population remain at risk of contracting COVID-19 as they wait to be vaccinated, which has prompted governments to encourage people to continue to self-isolate, remain at home, and maintain social distancing. The identification of safe and easily distributed medications that can target the different stages of virus infection and replication, could reduce the spread of SARS-CoV-2 and reduce the cases of COVID-19.

The Food and Drug Administration (FDA) and the European Medicines Agency (EMA) work with pharmaceutical companies to develop safe and effective drugs for the benefit of public health. Using a library of 1971 FDA-approved compounds, here we used a traceable clone of SARS-Cov-2 to identify well-characterized drugs that could potentially slow the infection and replication of the SARS-CoV-2 virus in human cells.

We had previously developed an approach to quantitate collagen synthesis by using CRISPR-Cas9 to insert the gene encoding NanoLuciferase (NLuc) into the Col1a2 gene in fibroblasts (Calverley et al., 2020). Here, we adapted the approach to tag the SARS-CoV-2 virus with NLuc, and then used the recombinant NLuc-tagged virus to monitor virus replication and to identify drugs that slow the replication process. SARS-CoV-2 was originally recovered by culturing human specimens in the African green monkey kidney cell line, Vero (Zhou et al., 2020), and the virus readily replicates in this cell type. We have performed initial screening in the Vero cell line, and in order to draw reliable conclusions from our studies we performed a replicate screen in a human hepatocyte cell line, HUH7. We evaluated a panel of human cell types for their ability to be infected by SARS-CoV-2 and to support virus replication, and optimized culture conditions for each cell type to provide a robust system for screening in human cells. We found that hepatocytes and kidney epithelial cells were proficient in supporting SARS-CoV-2 replication whereas fibroblasts and lung epithelial cells supported only minimal replication.

Through applying stringent criteria in our hit selection and confirming efficacy of these compounds in suppressing replication in cells already infected with SARS-CoV-2, we identified a shortlist of nine drugs that were effective in inhibiting SARS-CoV-2 replication. The drugs identified included anti-cancer and anti-viral drugs, but also less toxic drugs such as atovoquone, ebastine and vitamin D3. The multiple steps involved in virus infection and proliferation, means that each of these drugs may have different molecular targets, and for some, additional targets are still being elucidated. As these drugs are FDA-approved and safe, and dosimetry has been established for use in people, clinical trials could be initiated for these drugs.

RESULTS

Generation of a traceable SARS-CoV-2 virus

High throughput screens have been developed to identify drugs suitable for re-purposing for treatment of COVID19 (e.g., (Chen et al., 2020; Riva et al., 2020)). These screens have been performed in non-human cell lines, such as Vero, and rely on secondary factors such as cell viability to identify candidates, or markers that indicate virus infection. To monitor the replication of SARS-CoV-2 we have generated a modified traceable virus where Orf7a is replaced with the sequence encoding NanoLuciferase (SARS-Cov-2-ΔOrf7a-NLuc, **Fig. 1A**). NanoLuciferase (NLuc) is an enzyme that produces light when supplied with its substrate (coelenterazine) and is readily detectable even at low quantities (Calverley et al., 2020). Orf7a has previously been removed in SARS-CoV and SARS-CoV-2 and yielded infectious and replicative virus particles (Thi Nhu Thao et al., 2020; Xie et al., 2020a; Xie et al., 2020b). We have used a reverse-genetics approach to generate virus particles (**Supplemental Fig. 1**), and plaque forming assays demonstrated that the replication of SARS-CoV-2-ΔOrf7a-NLuc viruses were equivalent to the wild-type SARS-CoV-2 (Wuhan-Hu-1, NC_045512.2) (**Fig. 1B**). Viral RNAs (**Supplemental Fig. 1**), and electron microscopy (**Fig. 1C-D, Supplementary Fig. 2**) confirmed virus production. NLuc activity was readily detected in infected cultures upon addition of the substrate coelentrazine (**Fig. 1E**).

NLuc activity as a marker of virus replication

To monitor replication of the SARS-CoV-2-ΔOrf7a-NLuc virus, Vero cells were exposed to increasing numbers of replicative virus particles (plaque forming units, PFU) and luminescence was used as a measure of virus replication. Luminescence measurements were compared to SARS-CoV-2-ΔOrf7a-NLuc virus in the absence of cells as background (**Fig. 2A**), 24, 48 and 72 hours post infection (h.p.i.). A lag in virus replication of at least 48 hrs was observed but replication was readily observed at 72 hrs, when as little as 2 PFU per well (MOI 0.0004) was added. Using these optimization experiments 72 h.p.i. was chosen as the time point at which to assess virus replication in

subsequent assays, the average Z' factor (Zhang et al., 1999) for our replication assay in 96 well format was 0.8 indicating suitability for library screening. Amplification of NLuc signal upon virus infection was consistent for all passages of the traceable virus (**Supplementary Fig. 3**) and with varying cell number (**Supplementary Fig. 4**).

Earlier studies have demonstrated interferon alpha 2 (IFNA2) is able to reduce SARS-CoV-2 replication (Xie et al., 2020a). We confirmed this observation using our assay (**Fig. 2C**) but noted that with higher virus load (100 PFU), IFNA2 had little effect on virus replication (**Fig. 2D**). These results provided confidence that the traceable virus could be used to quantify the effects of drugs on virus replication.

SARS-CoV-2 replication screen validation
High throughput screens have been developed to identify drugs suitable for re-purposing for treatment of COVID19 (e.g., (Jeon et al., 2020; Riva et al., 2020; Touret et al., 2020)). These screens have been performed in non-human cell lines, such as Vero, and rely on secondary factors such as cell viability to identify candidates, or markers that indicate virus infection. A first step in validating our viral replication screen was to measure virus replication after treatment with interferon alpha 2 (IFNA2) which has previously demonstrated to reduce virus replication (Xie et al., 2020a). When Vero cells were infected with a single PFU, pre-treatment of IFNA2 was able to reduce replication of SARS-CoV-2-ΔOrf7a-NLuc in a dose dependent manner. However, with higher virus load (100 PFU), IFNA2 had little effect on virus replication (**Fig. 2C and D**). These results provided confidence that the NLuc-tagged virus system could be used to quantify the effects of drugs on virus replication, especially if reduction of bioluminescence was detected with high PFU.

Drug repurposing to target SARS-CoV-2 replication.
A 1971 FDA-approved compound library was used to pretreat Vero cells before infection with either 1PFU (MOI 0.0002) or 100 PFU (MOI 0.02) SARS-CoV-2-ΔOrf7a-NLuc virus as outlined in **Fig. 3A**. Infected Vero cells that were untreated and background (virus only wells) were used as controls. The Z' factor was assessed for each plate of the screen averaging 0.867± 0.140 SD and 0.915± 0.058 SD, for the 1PFU and 100PFU screen, respectively demonstrating a robust assay. More hits were identified in the 1PFU screen and therefore we focused our hit selection on compounds that reduced virus replication by at least 85% in the 100PFU screen (**Fig. 3B** and **Supplementary Table 3**). We identified 69 compounds that reduced replication (FDR<0.1 compared to untreated and DMSO controls, 55 showed no viral replication). These hits were also identified in the duplicate screen performed with 1PFU per well (**Fig. 3C**). Five of these compounds were false positive hits due to direct inhibition of NLuc activity (**Fig. 3C**, **Supplemental Table 4**).

Drug repurposing screen for SARS-CoV-2 replication in human cells.
Having established an effective screening strategy in Vero cells, we sought to establish a similar approach in a human cell line capable of supporting SARS-CoV-2 replication. SARS-CoV-2 has been reported to infect multiple cell lines, and replication of the viral genome has been identified (Chu et al., 2020). We tested the replicative capacity of SARS-CoV-2-ΔOrf7a in the lung epithelial cell lines Calu3, A549, and 16HBEo and in lung fibroblasts. Using real-time PCR and western blotting for the SARS-CoV-2 nucleocapsid protein, we observed virus replication in these cell lines, and we confirmed that SARS-CoV-2 WT and SARS-CoV-2-ΔOrf7a-NLuc viruses infected these lung cell lines (**Supplemental Fig. 5**). However the Z' factor suggested that they were not suitable for high throughput screening (**Fig. 4A**). We assessed additional cell lines reported to be infected by SARS-CoV-2 including Caco-2 (Chu et al., 2020; Touret et al., 2020) and HUH7 (Zhou et al., 2020), as well as additional kidney cell lines. These were chosen as we were able to recover virus particles from 293T cells and kidney is reported to express high levels of ACE2 (Wysocki et al., 2020). Replication in monocytic cells and fibroblastic cell lines was also monitored due to the systemic and fibrotic nature of COVID-19. We identified virus replication in HUH7 cells, microvascular cells of the kidney glomerulus, and proximal tubule cells of the kidney and THP1 monocytic cells (**Fig. 4A**).

As each cell line has a specific culture medium for optimal growth, we set out to test if different media would support cell growth and virus replication in lung epithelial cells, as this is the primary site of Sars-CoV2 infection. We used 15 different medium conditions to grow lung epithelial cells prior to infection with SARS-CoV-2-ΔOrf7a-NLuc; for comparison, HUH7 cells were also included. Changes in growth conditions improved assay performance in lung epithelial and fibroblasts but signal-to-noise ratios remained low. In contrast, for HUH7 cells, these optimized conditions significantly improved the 96 well format assay as indicated by Z' factor >0.5 and signal-to-noise ratios >20 (**Fig. 4B and C**). Using the 1971 FDA-compound library we identified 223 compounds that suppressed SARS-CoV-2 replication by greater than 85% whilst maintaining cell viability. We refined these hits by overlapping with positive hits from the screen in Vero cells (**Fig. 4E**, **Supplemental Table 5**). We identified 35 inhibitory compounds and 2 compounds that increased NLuc activity, that were common to the two screens (**Fig. 4F**). The intended clinical use or targets of the 35 inhibitory compounds included anti-virals, antibiotics, modifiers of dopamine and estrogen receptor activity, calcium ion channel inhibitors and HMG-CoA reductase inhibitors (**Fig. 5A**). The hits also identified vitamin D3, which is available over the counter. Multiple vitamin D related compounds were present in the screen and these too suppressed SARS-CoV-2-ΔOrf7a-NLuc although these did not all meet our criteria for both Vero and human cell lines (**Fig. 5B**).

These initial hits were further tested to determine if treatment could also *prevent* replication in cells already infected with SARS-CoV-2-ΔOrf7a-NLuc. We infected Vero cells with SARS-CoV-2 for 24 hrs (which is sufficient time for the virus to begin replication) then added each of the 35 inhibitor compounds. The cells were incubated for 48 hrs (i.e. a total of 72 h.p.i) prior to bioluminescence being measured. The majority of the compounds had no impact on virus replication; but 9 of the 35 compounds reduced replication relative to DMSO controls (Table 1, **Supplemental Fig. 6**). The effective doses for these compounds were then determined in HUH7 cells to provide a future reference for selecting dose after comparison to pharmacokinetic data established in human trials (**Fig. 5**, **Table 1**). Whilst these are pre-clinical *in vitro* data, they demonstrate the efficacy of the compounds in reducing virus replication post infection and warrant further investigation to determine if these could ease the burden of the virus in patients.

DISCUSSION

In this study we have shown that the SARS-CoV-2 virus infects and replicates in a range of human cells especially hepatocytes, kidney glomerulus, and proximal tubule cells of the kidney, and, that 9 drugs that have previously been shown to be safe in humans and approved by the FDA for clinical use are effective in inhibiting SARS-CoV-2 infection and replication.

The identification that liver and kidney cells are infection and replication competent for SARS-CoV-2 aligns with the observed liver and kidney abnormalities in patients with COVID-19. Liver comorbidities have been reported in 2-11% of patients with COVID-19, and 14-53% of cases reported abnormal levels of liver enzymes (Huang et al., 2020), with liver injury being more prevalent in severe cases (reviewed by (Zhang et al., 2020)). Another study reported that patients with chronic liver disease, especially African Americans, were at increased risk of COVID-19 (Wang et al., 2020). It has been suggested that liver damage in patients with COVID-19 might be caused by viral infection of liver cells, which is supported by the presence of SARS-CoV-2 RNA in stool (Yeo et al., 2020). On a similar note, acute kidney injury is a common complication of COVID-19 and has been associated with increased morbidity and mortality (reviewed by (Braun et al., 2020)). Our data showing SARS-CoV-2 infection and replication in kidney cells helps to explain kidney pathology associated with COVID-19.

RECOVERY (Group) has set out to identify therapeutics that could be re-purposed for treatment of COVID patients. For example, dexamethasone, which is a broad spectrum immunosuppressor, has been applied clinically to inhibit the destructive effects of the cytokine storm and shown to reduce mortality and decrease the length of hospitalization (Group et al., 2021), did not demonstrate any effect on viral replication in our study. It may be possible

in the future to combine the therapeutics identified in our study with dexamethasone so that both the cytokine storm and virus replication are targeted.

Of interest, other therapeutics that have been tested in the RECOVERY trial include ritonavir and lopinavir (antiretroviral protease inhibitors), which failed to show clinical benefit in the treatment of COVID-19 (Group). These drugs were present in the DiscoveryProbe library used in our study but showed no significant effect on SARS-CoV-2 replication, in our assays.

The high costs and lengthy lead-in times associated with new drug development, make repurposing of existing drugs for the treatment of common and rare diseases an increasingly attractive idea (for review see (Pushpakom et al., 2019)). The approaches used include hypothesis driven, preclinical trials including computational (e.g., computational molecular docking) and experimental (e.g., biochemical or cell-based assays of drug interactions) methods, and evaluation of efficacy in phase II clinical trials. As noted by Pushpakom *et al.*, of these three steps, the identification of the right drug for an indication of interest (in our case SARS-CoV-2 and COVID-19) is critical.

The nine drugs that we identified here have been approved for use in the treatment of a variety of diseases. Panobinostat, for example, is a HDAC inhibitor that blocks DNA replication, and has been used to inhibit cell growth in the management of cancer. Panobinostat had the strongest effect on limiting SARS-CoV-2 replication whilst maintaining cell viability, and completely blocked replication of SARS-CoV-2 at all doses tested (**Figure 5**); however, if cells were infected prior to treatment a more modest effect on replication were observed. This difference may be related to the recent observations that panobinostat can suppress ACE2 expression (He and Garmire, 2020), which would explain its beneficial effect prior to entry of the virus into cells and not thereafter. Abemaciclib (LY2835219) is another cell cycle inhibitor, suppressor of DNA replication and anti-cancer drug that emerged from the screen however these are likely to be detrimental to recovery from SARS-CoV-2 infection.

Atovaquone is of particular interest because it has been identified in other studies of SARS-CoV-2 in the context of COVID-19. It is a hydroxynaphthoquinone approved by NICE for the treatment of mild to moderate pneumocystis pneumonia and as a prophylaxis against pneumocystis pneumonia. It also has been used in combination with Proguanil (Malarone) as an antimalarial. The mechanism of action of atovaquone has been widely studied; it is a competitive inhibitor of ubiquinol, and against *P. falciparum*, it acts by inhibiting the electron transport chain at the level of the cytochrome bc1 complex (Fry and Pudney, 1992). A more recent study showed that atovaquone inhibits Zika and Dengue virus infection by blocking envelop protein-mediated membrane fusion (Yamamoto et al., 2020). Furthermore, in silico molecular docking strategies suggested potential binding of atovaquone to the SARS-CoV-2 spike protein (Farag et al., 2020; Ramirez-Salinas et al., 2020). These two mechanisms of action help to explain how atovaquone slows both infection and replication of SARS-CoV-2 in cells.

Our observation that bedaquiline inhibits SARS-CoV-2 replication supports evidence from an in silico repurposing study in which bedaquiline was proposed to be a promising inhibitor of the main viral protease of SARS-CoV-2 (Ferraz et al., 2020). The main protease cleaves pp1a and pp1ab polypeptides, which encode nonstructural proteins to form the replication-transcription complex (Snijder et al., 2006), and help explain how inhibitors to the main protease are effective in inhibiting replication of SARS-CoV-2 in Vero 76 cells (Ma et al., 2020). Manidipine (a calcium ion channel blocker approved for the use in treating hypertension) is proposed to reduce the activity of the main protease of SARS-CoV-2 (Ghahremanpour et al., 2020) and limit the availability of calcium ions, which are required for insertion of the coronavirus fusion peptide into the host lipid bilayers during viral fusion (Lai et al., 2017).

Our screen also identified ebastine (a second generation H1 receptor antagonist that has been approved for the treatment of allergic rhinitis and chronic idiopathic urticaria) and vitamin D3, which is a health supplement available over the counter. Ebastine is a second generation H1-antihistamine with proven efficacy for treating seasonal allergic rhinitis. However, histamine antagonists exhibit effects in addition to blockage of the histamine receptor. For

example, ebastine blocks the release of anti-IgE-induced prostaglandin D2 (Campbell et al., 1996). The vitamin D receptor (VDR, a member of the nuclear hormone receptor superfamily) is proposed to be essential for liver lipid metabolism because its deficiency in mice protects against hepatosteatosis (Bozic et al., 2016). Once bound to VDR, vitamin D plays a major role in hepatic pathophysiology, regulation of innate and adaptive immune responses, and might contribute to anti-proliferative, anti-inflammatory and anti-fibrotic activates (reviewed by (Triantos et al., 2021)). Thus far, whether vitamin D supplementation reduces the risk of SARS-CoV-2 infection or COVID-19 severity is unclear (The Lancet Diabetes, 2021). Whilst vitamin D3 met the stringent cut-offs of 85% virus reduction, vitamin D2 and other Vitamin D related therapeutics also reduced virus replication but did not meet all criteria across both Vero and HUH7 cell lines.

In conclusion, our study has identified compounds that are safe in humans and show effectiveness in reducing SARS-CoV-2 infection and replication in human cells, especially hepatocytes. Their potency in stopping SARS-CoV-2 replicating in human cells in the face of the COVID pandemic, warrants further study.

MATERIALS AND METHODS

Cell culture
Cell lines maintained in growth medium are shown in Supplementary Table 1.

Generation of functional SARS-CoV-2 virus
DNA encoding the genome of SARS-CoV-2 and SARS-CoV-2-ΔOrf7a-NLuc were purchased from Vectorbuilder Inc. (Chicago, US). Transfection of virus encoding the DNA failed to generate replicative virus when electroporated into 293T cells. We therefore produced RNA molecules which encoded the virus *in vitro.* Briefly, virus encoding DNA (1 µg) was transcribed using T7 mMessenger mMachine (Thermo) with a GTP:Cap ratio of 2:1 used in a 20 µL reaction. In addition, RNA encoding the SARS-CoV-2 nucleocapsid was also generated by PCR using primers P1 and P2 (Supplementary Table 2). It has been reported that this aids the recovery of replicative virus (Insert ref 3). Viral RNA genomes (10 µl) and 2.5 µL nucleocapsid RNA were electroporated into 293T cells within the BSL3 laboratory. Viral RNAs were electroporated (5,000,000 cells, 1100 V, 20 ms and 2 pulses) and grown in T75 cm^2 flasks and 24 well plates. Cells grown in 24 well plates for 24-120 hrs were used to monitor changes in NLuc activity.

Virus production, maintenance and assessment of titer
Culture medium was collected from 293T cells 6 days after electroporation. This virus (P0) was used to infect cells of interest. As virus replication was slow in 293T cells, virus stocks were maintained by passage in Vero cells grown in DMEM supplemented with 2% FBS. Medium (1 mL) was used to infect Vero cells in order to generate P1 virus. Replication was assessed by measuring NLuc activity over 10 days. For subsequent passage of the virus, Vero cells in T75 cm^2 flasks were infected with 2 mL of medium containing virus, after 3 days the medium was collected and passed through 0.45 µm filters using Luer-loc syringes.

To titre the virus, 200,000 Vero cells were seeded in 6-well plates overnight in growth medium. After removing growth medium Vero cells were infected with 200 µL of serially diluted virus containing medium at 37 °C. After 1 hour, wells were overlaid with 2 mL of 0.3% low melt agarose in 2xDMEM containing 1% FBS and grown for 3 days. After 3 days infected cells were fixed with 10% PFA overnight and then stained with crystal violet. Plaques were identified by imaging plates on BioDoc-It gel documentation system (UVP, Upland, US).

To detect viral RNA in medium, 0.25 mL of virus containing medium was collected 3 days post infection, 0.75 mL TriPure LS reagent (Sigma-Aldrich St. Louis, US) was added, and RNA isolated according to the manufacturer's recommendations, in the final step RNA was dissolved in 15 µL DNAse/RNAse free water. For detection of SARS-COV-2 nucleocapsid transcripts in lung epithelial cells, 200,000 cells were infected at the indicated MOI for 3 days, monolayers were lysed directly in 1 mL Trizol (Invitrogen, Paisley UK). For assessing expression of ACE2, TMPRSS2 and NLP1 expression in lung epithelial cells RNA was isolated from the cells using Trizol and RNA isolated according

to the manufacturer's recommendations. For cDNA generation and real-time PCR we used conditions previously described (Pickard et al., 2019) using primers detailed in Supplementary Table 2.

Electron microscopy

After 3 days infection with P4 virus, Vero cells were scrapped and pelleted. Cell pellets were fixed using 2.5% glutaraldehyde and 4% paraformaldehyde in 0.1 M cacodylate buffer for 24 h, washed in ddH$_2$O three times, 30 mins for each wash. Cell pellets were incubated in freshly made 2% (vol/vol) osmium tetroxide and 1.5% (wt/vol) potassium ferrocyanide in cacodylate buffer (100 mM, pH 7.2) for 1 hr at room temperature. Samples were washed in ddH$_2$O five times each for 3 minutes. Specimens were transferred to freshly made 1% (wt/vol) tannic acid in 100 mM cacodylate buffer (pH 7.2) for 40 mins at RT and washed in ddH$_2$O five times for 3 mins each at RT. The specimen was incubated with 1% (vol/vol) osmium tetroxide in ddH$_2$O for 30 minutes at room temperature and washed in ddH$_2$O three times for 5 min each at room temperature. The specimen was then incubated with 1% (wt/vol) uranyl acetate (aqueous) at 4 °C for 16 hrs (overnight) and then washed in ddH$_2$O three times for 5 mins each time at room temperature.

Specimens were dehydrated in graded ethanol: 30, 50, 70, 90% (vol/vol) ethanol in ddH$_2$O for 10 mins at each step. Then samples were washed four times for 10 mins each time in 100% ethanol at room temperature. Samples were transferred to propylene oxide for 10 min at room temperature.

The specimen was finally infiltrated in a graded series of Agar100Hard in propylene oxide at room temperature: first for 1 hour in 30% (vol/vol) Agar100Hard, 1 hr in 50% (vol/vol) Agar100Hard then overnight in 75% (vol/vol) Agar100Hard, and then 100% (vol/vol) Agar100Hard for 5 hrs. After transferring samples to freshly made 100% Agar100 Hard in labelled molds and allowed to cure at 60 °C for 72 hrs. Sections were imaged on an FEI Tecnai12 BioTwin.

NLuc activity assay

Vero cells grown in 24-well plates (Corning, 3526) were assayed for NLuc activity by adding 1 µL of coelenterazine (final concentration 1.5 µM). For 96 well formats, cell lines were seeded in white walled microwell plates (Nunc™ MicroWell™ 96-Well, Nunclon Delta-Treated, Flat-Bottom Microplate, Thermo Fisher Scientific, Paisley, UK# 136101). To measure NLuc activity, 0.5 µL coelenterazine was added per well (final concentration 3 µM). Light production was measured using filter cubes #114 and #3 on the Synergy Neo2 Multi-Mode Reader (Biotek), readings for each well were integrated over 200 ms with 4 replicate measurements per well (Gain 135 and read height 6 mm). For viability measurements, 2.5 µL Prestoblue (Thermo Fisher Scientific, Paisley UK) was added per well incubated for 10 mins before reading fluorescence at Excitation: 555/20 nm, Emission: 596/20 nm (Xenon flash, Lamp energy low, Gain 100 and read height 4.5 mm, 10 measurements per data point).

Drug screens

The DiscoveryProbe FDA-approved library of 1971 compounds (L1021, APExBIO Boston, US) was prepared as follows. After thawing the library for 4 hours at room temperature the library was arrayed into 96 well plates at 1 mM in DMSO and stored at -20 °C. Stocks (1 mM) were thawed at room temperature for 2 hrs before compounds were added to cells.

For all drug screens and validation, 5000 cells were seeded in 50 µL of growth medium for 24 hrs in white walled microwell plates. DMEM containing 2% FBS was used for Vero cells and alphaMEM containing 10% FBS was used for HUH7 cells. The following day 0.5 µL of each compound (final concentration 10 µM) was added per well and incubated for 24 hours. Eighty-eight compounds were tested per plate and each plate contained the following controls: two untreated wells, two DMSO treated wells, and one well treated with 10 µM puromycin to kill cells. SARS-CoV-2-ΔOrf7a-NLuc virus was added to all wells. In addition, two wells did not contain any cells but were infected with the SARS-CoV-2-ΔOrf7a-NLuc virus as a measure of background NLuc activity. A final well which contained cells but were uninfected were also included. Twenty-four hours after drug treatment, cells were infected with 100 PFU per well

SARS-CoV-2-ΔOrf7a-NLuc virus in 50 μL of the indicated growth medium for 72 hrs. To assess virus replication and viability, 2.5 μL of Prestoblue was added to each well and plates incubated for 10 mins at 37 °C. Coelenterazine was then added to a final concentration of 3 μM. Plates were sealed prior to reading luciferase activity and viability as described above. Validation of hits were performed using the same procedures described for the drug screen.

Study design and statistical analysis

To identify hit compounds, raw luciferase luminescence reads (x) were normalised relative to the virus-infected drug-untreated controls (u) and plate minimum read (m) on each plate by the formula:

$$x_{norm} = \frac{x_{raw} - m}{u - m}$$

This meant that a normalized luciferase value of 1 implied no difference from untreated virus replication, and a value of 0 represented total inhibition of viral replication.

The PrestoBlue reads (p) were normalised relative to the virus-infected DMSO-treated controls (d) and plate minimum read (n) on each plate by the formula:

$$p_{norm} = \frac{p_{raw} - n}{d - n}$$

This meant that a normalized PrestoBlue value of 1 implied no difference in cell viability from DMSO-treated virus infected cells, and a value of 0 representing maximal reduction in cell viability.

Z-scores were calculated relative to the mean log$_2$(fold change) for each plate.

Compounds were categorized as either inhibitors or enhancers of NLuc-SARS-CoV-2 activity. Inhibitors were compounds where normalized NLuc-SARS-CoV-2 levels (x_{norm}) was reduced such that $x_{norm} < 0.15$, and cell viability as measured by normalised PrestoBlue levels (p_{norm}) wasn't affected by more than 50%, such that $p_{norm} > 0.5$.

Where indicated one-tailed Student's T-test were performed to evaluate significance in changes to NLuc activity.

Contributors

AP co-conceived the project, performed the experiments, interpretated the data, drafted the manuscript. BCC performed experiments and interpreted the data. JC performed experiments. RG performed experiments. YL performed experiments. KEK co-conceived the project, raised the funding, drafted the manuscript. All authors approved the final manuscript.

Declaration of interests

The authors have no conflicts of interest.

Acknowledgements

The authors thank Dr. Jennifer Cavet for providing assistance with working at containing level 3 and use of the facilities. The project was approved by the COVID-19 Rapid Response Group, the R3G Research Operations Group and the R3G Executive Committee at the University of Manchester. We would also like to thank staff and students at the University of Manchester for the provision of cell lines for this study: Dr Stuart Cain, Dr Jonathan Humphries, Dr Shiu-wan Chan, Dr Chris Smith, Mrs Rachel Compton, Dr Rogerio Almeida, Dr Sara Gago, Dr Andrew Higham, and Dr Jeremy Herrera. Also thanks to Mrs Nikki-Maria Koudis, Mrs Maryline Fresquet, Dr Bernard Davenport and Dr Richard Naylor from Professor Rachel Lennon's laboratory for help sourcing kidney cell lines and T7 mMessenger mMachine reagents. We would also like to thank Thermo Fisher Scientific for their help in providing microwell plates for this

study, with special thanks to Charlotte Connor and Claire Marshall. We would like to thank Miss Anna Pickard for her illustration of the coronavirus.

Table 1: Approved uses of the 9 compounds of interest that inhibit SARS-CoV-2 infection and replication in HUH7 cells.

Compound	HUH7 IC50	Vero IC50	Post-infection at IC50	Approved Use	Target
Panobinostat	<0.2 µM	<0.2 µM	-25%	Multiple myeloma	HDAC inhibitor
LY2835219	0.2 µM	1 µM	-40%	Abemaciclib for advanced breast cancer	CDK4/6 inhibitor
Manidipine	2 µM	7.5 µM	-50%	Anti-hypertensive	Calcium channel blocker
Manidipine 2HCl	2.5 µM	7.5 µM	-50%	Anti-hypertensive	Calcium channel blocker
Ebastine	0.5 µM	5 µM	-25%	Anti-histamine for allergic rhinitis and chronic idiopathic urticaria	H_1 receptor blocker
Atovaquone	7.5 µM	3 µM	-25%	antimicrobial (anti-malarial)	Selective inhibitor of parasite mitochondrial electron transport
Bedaquiline	5 µM	10 µM	-30%	pulmonary tuberculosis	mycobacterial ATP synthase
Vitamin D3	3 µM	10 µM	-30%	Common supplement	Transcriptional/calcium regulation
Amodiaquine	1 µM	5 µM	-25%	antimicrobial (anti-malarial)	Inhibitor heme polymerase activity

Figure legends

Figure 1: Nanoluciferase (NLuc) modified SARS-CoV-2 virus as a reporter for virus replication.

A) Diagram of the SARS-CoV-2 genome highlighting the insert site for the reporter NLuc in place of Orf7a, (SARS-CoV-2-ΔOrf7a-Nluc).

B) Virus particles recovered following transfection of wild-type (WT) or NLuc modified SARS-CoV-2 encoding RNA into 293T cells were used to infect Vero cells. The recovered virus (P1 virus) was then titred in Vero cells to assess virus replication and plaque forming potential.

C) Electron microscopy of the SARS-Cov-2 virus 72 h.p.i. of Vero cells (Passage 4, P4), both intra- and extra-cellular virus particles were identified.

D) Measurement of diameter for the wild-type (WT) and NLuc modified SARS-CoV-2 particles.

Figure 2: Timings of SARS-CoV-2-ΔOrf7a-NLuc virus replication.

A) Schematic showing how SARS-CoV-2 ΔOrf7a-NLuc was used to assess virus replication. During virus replication as virus particles are released from the cell, or as a result of cell death, NLuc activity is detected in the conditioned medium together with virus particles. In subsequent assays the NLuc activity is recorded in the absence of cells, and NLuc signals above background signal are used to indicate the level of virus replication.

B) The replication of SARS-CoV-2-ΔOrf7a-Nluc was monitored in a 96 well assay. Using 5000 Vero cells per well, increasing numbers of virus particles were added and monitored 24, 48 and 72 hours post infection (h.p.i). Substantial and significant increases in NLuc activity were observed when only two plaque forming units (PFU) were added. Higher NLuc activity was associated with higher viral input. Even with higher viral inputs the substantial increases in NLuc activity occurred only after 72 h.p.i.

C) Five thousand Vero cells were pre-treated with increasing doses of IFNα2 for 24 hours prior to infection with 1PFU SARS-CoV-2-ΔOrf7a-Nluc. NLuc activity and viability were assessed 72 h.p.i. demonstrating effective inhibition of virus replication. The blue line indicates the luminescence counts per second for DMSO treated controls. N=3 replicate samples

D) As in C, 5000 Vero cells were treated with increasing doses of IFNα2 for 24 hours prior to infection with 100PFU SARS-CoV-2-ΔOrf7a-Nluc. With this higher titre, virus replication was not inhibited. The blue line indicates the luminescence counts per second for DMSO treated controls. N=3 replicate samples.

Figure 3: Screen of 1971 FDA-approved compounds to identify therapeutics that inhibit SARS-CoV-2-ΔOrf7a-NLuc virus replication.

A) Schematic of the screening procedure used to assess whether FDA-approved compounds alter SARS-CoV-2 ΔOrf7a-Nluc virus replication.

B) Scatterplot showing the effects of 1971 compounds on Sars-CoV-2 ΔOrf7a-NLuc replication and cell viability.

C) Scatterplot showing the effects of 1971 compounds on Sars-CoV-2 ΔOrf7a-NLuc replication in replicate screens using 100 PFU or 1 PFU per well. The Z-score for luciferase activity is plotted.

D) Scatterplot of 50 compounds that reduced NLuc activity in C. The luciferase activities relative to untreated controls (indicated by the blue dashed line) are shown. Highlighted in green are compounds that inhibit the NLuc enzyme directly.

Figure 4: Replication of SARS-CoV-2-ΔOrf7a-Nluc in human cell lines.

A) SARS-CoV-2 replication in human cell lines. Five thousand cells per well were seeded and infected as in Figure 3A. All cells were grown in the recommended growth medium, detailed in Table 1. The NLuc activity after 3 days was used to assess the degree of virus replication. NLuc activity above the baseline activity is shown as a heat map. Each box is the average of 2 biological repeats of each cell line at each virus dose, with the same findings observed when 2000, or 10,000 cells were seeded.

B) Lung epithelial, fibroblasts, HUH7 and 293T cell lines were grown in 15 different growth medium compositions. NLuc activity over background three days after infection is shown. Each row represents a different growth medium as indicated in C). Each box represents 4 replicate measures of replication. Similar findings were observed in n=6 biological repeats. Z' factors were calculated for each virus dose using 'virus only' wells as negative controls, the

average Z' factor for each growth medium is shown. The signal-to-noise ratio of infected cells relative to 'virus only' wells are also shown.

C) NLuc activity above background three days after infection in HUH7 cells. Each box represents 4 replicate measures of replication. Similar findings were observed in n=6 biological repeats.

D) The effects of 1971 FDA-approved compounds on the replication of SARS-CoV-2-ΔOrf7a-NLuc (100 PFU/well, MOI 0.02) in HUH7 cells grown in alpha-MEM supplemented with 10% FBS. Five thousand cells were treated with 10 µM of each compound for 24 hours prior to infection with 100 PFU of virus. Virus replication progressed for 72 hours as in Figure 3A.

E) Comparison of compounds identified by screening for HUH7 and Vero cells Z-scores for luciferase activity are shown

F) Comparison and overlap of inhibitors identified in the compound screening for HUH7 and Vero cells.

Figure 5: Dose response of SARS-CoV-2 inhibitors.

A) Summary of the inhibitor class of the 35 compounds identified in both the Vero and HUH7 screens.

B) Effects of vitamin D related compounds from the APExBIO DiscoveryProbe library on NLuc activity in the Vero and HUH7 screens.

C) Dose responses to the 9 compounds in HUH7 cells. Cells were treated and infected as described in Figure 4D. NLuc activity relative to DMSO controls are shown in blue, and viability relative to DMSO are shown in black. N=2 independent repeats are shown.

Supplementary Figure 1: Recovery of replication competent SARS-CoV-2 from synthetic DNA constructs.

A) Schematic for the recovery of SARS-CoV-2 virus particles from DNA encoding the wild type and NLuc modified SARS-CoV-2 genome. RNA was transcribed from the synthetic DNA constructs and electroporated into 293T cells, the cells were monitored daily for NLuc activity and evidence of the cytopathic effects of virus replication. Virus containing medium was collected 6 days post electroporation.

B) NLuc activity in 293T cells electroporated with wild type and NLuc modified SARS-CoV-2 RNA transcripts.

C) Real-time PCR primer validation using SARS-CoV-2 encoding DNA. NLuc coding sequences were inserted in place of the Orf7a gene. Primers designed to amplify sequences within the DNA constructs confirmed primer specificity. The number of cycles for each gene were normalized to those of the WT virus, or for NLuc, to ΔOrf7a NLuc. For each DNA construct the qPCR data was also normalized to the number of cycles obtained for the N gene (N=3 technical repeats).

D) Real-time qPCR detection of viral RNAs in the medium of infected Vero cells 72 h.p.i. Vero cells were infected with Sars-CoV-2 WT (MOI 0.1) or Sars-CoV-2 ΔOrf7a-Nluc (MOI 1).

E) Initial assessment of viral replication in Vero cells. Twenty four h.p.i. NLuc activity was readily detected in SARS-CoV-2 ΔOrf7a-Nluc infected samples (N=3 replicate samples).

F) NLuc activity in Vero cells infected with wild type SARS-CoV-2 or SARS-CoV-2 ΔOrf7a-Nluc modified virus. NLuc activity was significantly elevated in SARS-CoV-2 ΔOrf7a-Nluc infected cultures across all time points, *** represents $p<0.001$ in a Students T-Test, N=3 replicate samples. With prolonged culture NLuc activity within infected samples increased approximately 30-fold.

Supplementary Figure 2: Detection of SARS-CoV-2 virus particles in Vero cells infected with passage 4 virus stocks.

A) After 4 passages of recovered wild type SARS-CoV-2 virus particles in Vero cells, naïve Vero cells were infected and fixed 72 h.p.i. Viral particles could be observed inside and outside of Vero cells indicated by red arrows. Scale bar, 2 µm.

B) Higher magnification image of an independent region of the sample in A. Scale bar, 0.5 μm.

C) After 4 passages of recovered SARS-CoV-2-ΔOrf7a-NLuc virus particles in Vero cells, naïve Vero cells were infected and fixed 72 h.p.i. Viral particles could be observed inside and outside of Vero cells indicated by blue arrows. Scale bar, 2 μm.

D) Higher magnification image of an independent region of the sample in C. Scale bar, 0.5 μm.

Supplementary Figure 3: Replication assays for different passages of SARS-CoV-2-ΔOrf7a-NLuc virus.

A) The plots show 5000 Vero cells infected with different volumes of virus harvested from repeat passages of the virus (P) in Vero cells. With increasing viral load, there is an increase in background as more NLuc activity is added to each well (Blue). The background signal arises when NLuc, expressed as the virus replicates, is released from infected cells that have lysed upon liberation of the virus particles. N=2 independent experiments.

B) After subtracting the background noise observed in wells without cells (blue in A), the additional NLuc activity generated through viral replication showed comparable signals for each virus passage. N=2 independent experiments.

C) The signal-to-noise ratio for each batch was used to identify the optimal conditions for virus replication for subsequent screening experiments.

Supplementary Figure 4: SARS-CoV-2-ΔOrf7a-Nluc replication optimisation

NLuc activity at different times after infection with increasing numbers of SARS-CoV-2 virus particles. Enhanced NLuc signals were observed 72 h.p.i. when either A) 2000, B) 5000 or C) 10,000 cells per well were used. Maximal signals were generated with 5000 cells per well, as seeding 10,000 cells tended to reduce NLuc activity. Five thousand cells were used for all subsequent assays.

Supplemental Figure 5: Detection of SARS-CoV-2 infection in lung epithelial cells.

A) Real-time PCR detection of SARS-CoV-2 nucleocapsid (N) RNA in cells 72 h.p.i. Viral RNAs were readily detected in cells infected with WT or ΔOrf7a-Nluc virus particles.

B) Western blot detection of SARS-CoV-2 nucleocapsid (N) protein in cells 72 hours post infection. Lower levels of the N protein are detected in fibroblasts and lung epithelial cells when compared to 293T and Vero cells suggesting the virus can infect but not replicate in these cell lines.

C) Real-time detection of known mediators of SARS-CoV-2 entry into cells. ACE2, NLP1 and TMPRSS2 RNA levels are shown relative to RPLP0. Infectious virus in lung cell lines but no replication.

Supplemental Figure 6: Effective compounds that suppress SARS-CoV-2 replication with treatment commencing after infection.

A) Schematic for assessing whether any compounds that suppressed virus replication prior to infection could also affect replication post-infection.

B) Fifty compounds identified to suppress SARS-CoV-2 replication in Figure 3 were assessed to identify dose-dependent effects on SARS-CoV-2 replication with treatment progressing after infection (as outlined in A). Nine compounds were found to suppress replication in a dose dependent manner.

References

Banerjee, A.K., Blanco, M.R., Bruce, E.A., Honson, D.D., Chen, L.M., Chow, A., Bhat, P., Ollikainen, N., Quinodoz, S.A., Loney, C., et al. (2020). SARS-CoV-2 Disrupts Splicing, Translation, and Protein Trafficking to Suppress Host Defenses. Cell. 183(5), 1325-1339 e1321. Published online 2020/10/21 DOI: 10.1016/j.cell.2020.10.004.

Bozic, M., Guzman, C., Benet, M., Sanchez-Campos, S., Garcia-Monzon, C., Gari, E., Gatius, S., Valdivielso, J.M., and Jover, R. (2016). Hepatocyte vitamin D receptor regulates lipid metabolism and mediates experimental diet-induced steatosis. J Hepatol. 65(4), 748-757. Published online 2016/06/02 DOI: 10.1016/j.jhep.2016.05.031.

Braun, F., Lutgehetmann, M., Pfefferle, S., Wong, M.N., Carsten, A., Lindenmeyer, M.T., Norz, D., Heinrich, F., Meissner, K., Wichmann, D., et al. (2020). SARS-CoV-2 renal tropism associates with acute kidney injury. Lancet. 396(10251), 597-598. Published online 2020/08/21 DOI: 10.1016/S0140-6736(20)31759-1.

Calverley, B.C., Kadler, K.E., and Pickard, A. (2020). Dynamic High-Sensitivity Quantitation of Procollagen-I by Endogenous CRISPR-Cas9 NanoLuciferase Tagging. Cells. 9(9). Published online 2020/09/16 DOI: 10.3390/cells9092070.

Campbell, A., Michel, F.B., Bremard-Oury, C., Crampette, L., and Bousquet, J. (1996). Overview of allergic mechanisms. Ebastine has more than an antihistamine effect. Drugs. 52 Suppl 1, 15-19. Published online 1996/01/01 DOI: 10.2165/00003495-199600521-00005.

Cantuti-Castelvetri, L., Ojha, R., Pedro, L.D., Djannatian, M., Franz, J., Kuivanen, S., van der Meer, F., Kallio, K., Kaya, T., Anastasina, M., et al. (2020). Neuropilin-1 facilitates SARS-CoV-2 cell entry and infectivity. Science. 370(6518), 856-860. Published online 2020/10/22 DOI: 10.1126/science.abd2985.

Chatenoud, L., Ferran, C., and Bach, J.F. (1991). The anti-CD3-induced syndrome: a consequence of massive in vivo cell activation. Curr Top Microbiol Immunol. 174, 121-134. Published online 1991/01/01 DOI: 10.1007/978-3-642-50998-8_9.

Chen, Y., Tao, H., Shen, S., Miao, Z., Li, L., Jia, Y., Zhang, H., Bai, X., and Fu, X. (2020). A drug screening toolkit based on the -1 ribosomal frameshifting of SARS-CoV-2. Heliyon. 6(8), e04793. Published online 2020/09/02 DOI: 10.1016/j.heliyon.2020.e04793.

Chu, H., Chan, J.F., Yuen, T.T., Shuai, H., Yuan, S., Wang, Y., Hu, B., Yip, C.C., Tsang, J.O., Huang, X., et al. (2020). Comparative tropism, replication kinetics, and cell damage profiling of SARS-CoV-2 and SARS-CoV with implications for clinical manifestations, transmissibility, and laboratory studies of COVID-19: an observational study. Lancet Microbe. 1(1), e14-e23. Published online 2020/08/25 DOI: 10.1016/S2666-5247(20)30004-5.

Fajgenbaum, D.C., and June, C.H. (2020). Cytokine Storm. N Engl J Med. 383(23), 2255-2273. Published online 2020/12/03 DOI: 10.1056/NEJMra2026131.

Farag, A., Wang, P., Boys, I.N., L. Eitson, J., Ohlson, M.B., and Fan, W.e.a. (2020). Identification of Atovaquone, Ouabain and Mebendazole as FDA Approved Drugs Tar-geting SARS-CoV-2 (Version 4). ChemRxiv. DOI: https://doi.org/10.26434/chemrxiv.12003930.v4

Ferraz, W.R., Gomes, R.A., AL, S.N., and Goulart Trossini, G.H. (2020). Ligand and structure-based virtual screening applied to the SARS-CoV-2 main protease: an in silico repurposing study. Future Med Chem. 12(20), 1815-1828. Published online 2020/08/14 DOI: 10.4155/fmc-2020-0165.

Fry, M., and Pudney, M. (1992). Site of action of the antimalarial hydroxynaphthoquinone, 2-[trans-4-(4'-chlorophenyl) cyclohexyl]-3-hydroxy-1,4-naphthoquinone (566C80). Biochem Pharmacol. 43(7), 1545-1553. Published online 1992/04/01 DOI: 10.1016/0006-2952(92)90213-3.

Ghahremanpour, M.M., Tirado-Rives, J., Deshmukh, M., Ippolito, J.A., Zhang, C.H., Cabeza de Vaca, I., Liosi, M.E., Anderson, K.S., and Jorgensen, W.L. (2020). Identification of 14 Known Drugs as Inhibitors of the Main Protease of SARS-CoV-2. ACS Med Chem Lett. 11(12), 2526-2533. Published online 2020/12/17 DOI: 10.1021/acsmedchemlett.0c00521.

Group, R.C. Lopinavir–ritonavir in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. The Lancet. DOI: https://doi.org/10.1016/S0140-6736(20)32013-4.

Group, R.C. A randomised trial of treatments to prevent death in patients hospitalised with COVID-19 (coronavirus). ISRCTN registry. DOI: https://doi.org/10.1186/ISRCTN50189673.

Group, R.C., Horby, P., Lim, W.S., Emberson, J.R., Mafham, M., Bell, J.L., Linsell, L., Staplin, N., Brightling, C., Ustianowski, A., et al. (2021). Dexamethasone in Hospitalized Patients with Covid-19. N Engl J Med. 384(8), 693-704. Published online 2020/07/18 DOI: 10.1056/NEJMoa2021436.

He, B., and Garmire, L. (2020). Prediction of repurposed drugs for treating lung injury in COVID-19. F1000Res. 9, 609. Published online 2020/09/18 DOI: 10.12688/f1000research.23996.2.

Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., et al. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet. 395(10223), 497-506. Published online 2020/01/28 DOI: 10.1016/S0140-6736(20)30183-5.

Jeon, S., Ko, M., Lee, J., Choi, I., Byun, S.Y., Park, S., Shum, D., and Kim, S. (2020). Identification of Antiviral Drug Candidates against SARS-CoV-2 from FDA-Approved Drugs. Antimicrob Agents Chemother. 64(7). Published online 2020/05/06 DOI: 10.1128/AAC.00819-20.

Lai, A.L., Millet, J.K., Daniel, S., Freed, J.H., and Whittaker, G.R. (2017). The SARS-CoV Fusion Peptide Forms an Extended Bipartite Fusion Platform that Perturbs Membrane Order in a Calcium-Dependent Manner. J Mol Biol. 429(24), 3875-3892. Published online 2017/10/24 DOI: 10.1016/j.jmb.2017.10.017.

Li, W., Moore, M.J., Vasilieva, N., Sui, J., Wong, S.K., Berne, M.A., Somasundaran, M., Sullivan, J.L., Luzuriaga, K., Greenough, T.C., et al. (2003). Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. Nature. 426(6965), 450-454. Published online 2003/12/04 DOI: 10.1038/nature02145.

Ma, C., Sacco, M.D., Hurst, B., Townsend, J.A., Hu, Y., Szeto, T., Zhang, X., Tarbet, B., Marty, M.T., Chen, Y., et al. (2020). Boceprevir, GC-376, and calpain inhibitors II, XII inhibit SARS-CoV-2 viral replication by targeting the viral main protease. Cell Res. 30(8), 678-692. Published online 2020/06/17 DOI: 10.1038/s41422-020-0356-z.

Maier, H.J., Neuman, B.W., Bickerton, E., Keep, S.M., Alrashedi, H., Hall, R., and Britton, P. (2016). Extensive coronavirus-induced membrane rearrangements are not a determinant of pathogenicity. Sci Rep. 6, 27126. Published online 2016/06/04 DOI: 10.1038/srep27126.

Pickard, A., Chang, J., Alachkar, N., Calverley, B., Garva, R., Arvan, P., Meng, Q.J., and Kadler, K.E. (2019). Preservation of circadian rhythms by the protein folding chaperone, BiP. FASEB J. 33(6), 7479-7489. Published online 2019/03/20 DOI: 10.1096/fj.201802366RR.

Pushpakom, S., Iorio, F., Eyers, P.A., Escott, K.J., Hopper, S., Wells, A., Doig, A., Guilliams, T., Latimer, J., McNamee, C., et al. (2019). Drug repurposing: progress, challenges and recommendations. Nat Rev Drug Discov. 18(1), 41-58. Published online 2018/10/13 DOI: 10.1038/nrd.2018.168.

Ramirez-Salinas, G.L., Martinez-Archundia, M., Correa-Basurto, J., and Garcia-Machorro, J. (2020). Repositioning of Ligands That Target the Spike Glycoprotein as Potential Drugs for SARS-CoV-2 in an In Silico Study. Molecules. 25(23). Published online 2020/12/03 DOI: 10.3390/molecules25235615.

Riva, L., Yuan, S., Yin, X., Martin-Sancho, L., Matsunaga, N., Pache, L., Burgstaller-Muehlbacher, S., De Jesus, P.D., Teriete, P., Hull, M.V., et al. (2020). Discovery of SARS-CoV-2 antiviral drugs through large-scale compound repurposing. Nature. 586(7827), 113-119. Published online 2020/07/25 DOI: 10.1038/s41586-020-2577-1.

Ruan, Q., Yang, K., Wang, W., Jiang, L., and Song, J. (2020). Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. Intensive Care Med. 46(5), 846-848. Published online 2020/03/04 DOI: 10.1007/s00134-020-05991-x.

Snijder, E.J., van der Meer, Y., Zevenhoven-Dobbe, J., Onderwater, J.J., van der Meulen, J., Koerten, H.K., and Mommaas, A.M. (2006). Ultrastructure and origin of membrane vesicles associated with the severe acute respiratory syndrome coronavirus replication complex. J Virol. 80(12), 5927-5940. Published online 2006/05/30 DOI: 10.1128/JVI.02501-05.

The Lancet Diabetes, E. (2021). Vitamin D and COVID-19: why the controversy? Lancet Diabetes Endocrinol. 9(2), 53. Published online 2021/01/15 DOI: 10.1016/S2213-8587(21)00003-6.

Thi Nhu Thao, T., Labroussaa, F., Ebert, N., V'Kovski, P., Stalder, H., Portmann, J., Kelly, J., Steiner, S., Holwerda, M., Kratzel, A., et al. (2020). Rapid reconstruction of SARS-CoV-2 using a synthetic genomics platform. Nature. 582(7813), 561-565. Published online 2020/05/05 DOI: 10.1038/s41586-020-2294-9.

Touret, F., Gilles, M., Barral, K., Nougairede, A., van Helden, J., Decroly, E., de Lamballerie, X., and Coutard, B. (2020). In vitro screening of a FDA approved chemical library reveals potential inhibitors of SARS-CoV-2 replication. Sci Rep. 10(1), 13093. Published online 2020/08/06 DOI: 10.1038/s41598-020-70143-6.

Triantos, C., Aggeletopoulou, I., Thomopoulos, K., and Mouzaki, A. (2021). Vitamin D - liver disease association: Biological basis and mechanisms of action. Hepatology. Published online 2021/01/07 DOI: 10.1002/hep.31699.

Wang, Q., Davis, P., and Xu, R. (2020). COVID-19 risk, disparities and outcomes in patients with chronic liverdisease in the United States. EClinicalMedicine. DOI: ,https://doi.org/10.1016/j.eclinm.2020.100688.

Wysocki, J., Lores, E., Ye, M., Soler, M.J., and Batlle, D. (2020). Kidney and Lung ACE2 Expression after an ACE Inhibitor or an Ang II Receptor Blocker: Implications for COVID-19. J Am Soc Nephrol. 31(9), 1941-1943. Published online 2020/07/17 DOI: 10.1681/ASN.2020050667.

Xie, X., Muruato, A., Lokugamage, K.G., Narayanan, K., Zhang, X., Zou, J., Liu, J., Schindewolf, C., Bopp, N.E., Aguilar, P.V., et al. (2020a). An Infectious cDNA Clone of SARS-CoV-2. Cell Host Microbe. 27(5), 841-848 e843. Published online 2020/04/15 DOI: 10.1016/j.chom.2020.04.004.

Xie, X., Muruato, A.E., Zhang, X., Lokugamage, K.G., Fontes-Garfias, C.R., Zou, J., Liu, J., Ren, P., Balakrishnan, M., Cihlar, T., et al. (2020b). A nanoluciferase SARS-CoV-2 for rapid neutralization testing and screening of anti-infective drugs for COVID-19. Nat Commun. 11(1), 5214. Published online 2020/10/17 DOI: 10.1038/s41467-020-19055-7.

Yamamoto, M., Ichinohe, T., Watanabe, A., Kobayashi, A., Zhang, R., Song, J., Kawaguchi, Y., Matsuda, Z., and Inoue, J.I. (2020). The Antimalarial Compound Atovaquone Inhibits Zika and Dengue Virus Infection by Blocking E Protein-Mediated Membrane Fusion. Viruses. 12(12). Published online 2020/12/30 DOI: 10.3390/v12121475.

Yeo, C., Kaushal, S., and Yeo, D. (2020). Enteric involvement of coronaviruses: is faecal-oral transmission of SARS-CoV-2 possible? Lancet Gastroenterol Hepatol. 5(4), 335-337. Published online 2020/02/23 DOI: 10.1016/S2468-1253(20)30048-0.

Zhang, C., Shi, L., and Wang, F.S. (2020). Liver injury in COVID-19: management and challenges. Lancet Gastroenterol Hepatol. 5(5), 428-430. Published online 2020/03/08 DOI: 10.1016/S2468-1253(20)30057-1.

Zhang, J.H., Chung, T.D., and Oldenburg, K.R. (1999). A Simple Statistical Parameter for Use in Evaluation and Validation of High Throughput Screening Assays. J Biomol Screen. 4(2), 67-73. Published online 2000/06/06 DOI: 10.1177/108705719900400206.

Zhou, P., Yang, X.L., Wang, X.G., Hu, B., Zhang, L., Zhang, W., Si, H.R., Zhu, Y., Li, B., Huang, C.L., et al. (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature. 579(7798), 270-273. Published online 2020/02/06 DOI: 10.1038/s41586-020-2012-7.

Figure 1 (Pickard et al.)

A

CMV | T7 | UTR | | | | | | | | Envelope | M | 7a 8 | 6 7b | N | 10 | Poly A T7term

ORF1a/b | S | 3

ΔOrf7a NLuc

B

P1 virus

Sars Wt

Sars ΔOrf7a NLuc

C

Sars-Cov-2 ΔOrf7a – Nluc (Vero) P4

500 nm

500 nm

100 nm

D

p=0.35

Virus diameter (nm)

Sars-WT   Sars-ΔOrf7a-NLuc

E

Luciferase counts per second

● - Coelentrazine
■ + Coelentrazine

Sars-WT   Sars-ΔOrf7a-NLuc

Figure 2 (Pickard et al.)

Figure 3 (Pickard et al.)

Figure 4 (Pickard et al.)

Figure 5 (Pickard et al.)

Supplemental Figure 1 (Pickard et al.)

Supplemental Figure 2 (Pickard et al.)

A) SARS-CoV-2 (wild type)

B) SARS-CoV-2 (wild type)

C) SARS-Cov-2-ΔOrf7a–NLuc

D) SARS-Cov-2-ΔOrf7a–NLuc

Supplemental Figure 3 (Pickard et al.)

Supplemental Figure 4 (Pickard et al.)

Supplemental Figure 5 (Pickard et al.)

Supplemental Figure 6 (Pickard et al.)

Supplementary Table 1: Source and growth conditions for cell lines used in this study.

Cell line	Growth medium	Source
293T	DMEM + 10% FBS	ATCC
HUH7	DMEM + 10% FBS + NEAA	Accegen #ABC-TC0437
THP1	RPMI + 10%	Sigma-Aldrich, #88081201
SW1353	DMEM + 10% FBS	ATCC® HTB-94™
TC28a	DMEM + 10% FBS	Reference[1]
Hacat	DMEM + 10% FBS	Caltag Medsystems #T0020001
HFF	DMEM + 10% FBS	Cascade Biologics/Thermo Fisher C0045C
Caco2	DMEM + 10% FBS	ECACC #86010202
16HBE14o	AlphaMEM + 10% FBS	Reference[2]
Calu3	AlphaMEM + 10% FBS	ATCC® HTB-55™
A549	DMEM + 10% FBS	ATCC® CCL-185™
HLF (CCD-19Lu)	DMEM + 10% FBS	ATCC® CCL-210™
PodoTERT	PodoUP3	Evercyte #PODO/TERT256
MVGenC	Endothelial Basal medium-2 kit (without VEGF)	NeoBiotech #NB-11-0002
Ab Podocytes	RPMI + 10% FBS + ITS	Reference[3]
HK-2	DMEM:F12 + 10% FBS	ATCC® CRL 2190™
Vero	DMEM + 10% FBS	ATCC® CCL 81™

FBS= Foetal bovine serum, NEAA=Non-essential amino acids, ITS= Insulin-Transferrin-Selenium, VEGF=Vascular endo

References

1 **7989586**
2 **7507342**
3 11856766

thelial growth factor.

Supplementary Table 2: Primers used in this study

Target	Forward	Reverse
Nucleocapsid for RNA generation (P1 and P2)	TACTGTAATACGACTCACTATAGGATGTCTGATAATGGACCCCAAAATC	TTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTAGGCCTGAGTTGAGTCAGCAC
Sars-Cov2 N	CAACATTGCCAAAAGGCTTC	ACGAGAAGAGGCTTGACTGC
Sars-Cov-2 E	CCATCCTTACTGCGCTTCGA	AAGAAGGTTTTACAAGACTCACGTT
Sars-Cov2 Orf7a	TGGCACTGATAACACTCGCT	TGCCCTCGTATGTTCCAGAAG
NLuc	GACGAGCGCCTGATCAAC	GGTCACTCCGTTGATGGTTAC
ACE2	TTCTGTCACCCGATTTTCAA	TCCCAACAATCGTGAGTGC
TMPRSS2	CGCTGGCCTACTCTGGAA	CTGAGGAGTCGCACTCTATCC
NLP1	TACCCTGAGAATGGGTGGAC	CGTGACAAAGCGCAGAAG
RPLP0	ACTGGTCTAGGACCCGAGAAG	CTCCCACCTTGTCTCCAGTC

Supplemental Table 3: Inhibitors of Sars-Cov2-ΔOrf7a-Nluc replication in Vero cells

Plate	Row	Column	Drug name	Mean relati	SD	PrestoBlue	Replication greater than
4	A	1	Sitagliptin	0.000141	0.00015	2.560703	1.095235
4	C	1	Belinostat (	0.000366	0.000141	0.89498	1.095828
1	B	10	Cilnidipine	0.00039	0.000274	0.801674	1
9	G	1	Lacidipine	0.000439	0.000302	0.86	1.003471
4	B	1	MK-2048	0.000491	0.000218	2.266647	1.09346
4	H	1	Apixaban	0.00124	0.000138	2.563682	1.098206
4	D	1	LY2784544	0.001449	0.000197	0.785938	1.094642
4	F	1	Rosiglitazo	0.001834	0.000162	3.110979	1.09761
4	G	1	Pioglitazon	0.002045	0.000181	3.48488	1.096421
4	E	1	Lenalidomi	0.002184	0.000236	2.058394	1.094051
9	F	4	Benidipine	0.003673	0.000674	0.808649	1.003968
10	A	5	KPT-330	0.005454	0.002884	0.581033	1.019134
2	A	4	Vortioxetir	0.007391	0.002635	0.508971	1.048162
8	B	2	Fingolimoc	0.007569	0.002924	0.691059	1.112639
12	A	10	Ebastine	0.007931	0.001473	0.605394	1.00847
8	B	3	LY335979 (	0.008621	0.00215	0.586608	1.116935
15	C	1	Atovaquon	0.009591	0.002484	1.549669	1.057471
16	A	2	Eltrombopa	0.009908	0.003764	0.719096	1.072602
17	D	5	Bedaquilin	0.010156	0.002018	0.878372	1.050337
17	D	4	Bedaquilin	0.010526	0.001862	1.113109	1.047077
20	A	9	Chlorproth	0.0137	0.002098	0.829945	1.101788
22	A	4	Homoharri	0.014405	0.002412	0.506763	1.060797
1	H	1	Fluoxetine	0.014733	0.000426	0.760391	1.000989
1	H	4	Idarubicin	0.014922	0.000762	0.540586	1.006965
18	A	1	Ospemifen	0.015744	0.004355	0.686275	1.080042
1	H	3	Pamidrona	0.016484	0.000473	0.545607	1.001484
6	A	7	Panobinost	0.018674	0.003673	0.602074	1.07317
3	E	5	Salinomyci	0.018851	0.00162	0.612145	1.043837
16	G	5	Daunorubi	0.021008	0.004246	0.696494	1.089341
14	F	7	Probucol	0.022348	0.00316	0.753645	1.083511
8	F	11	Anidulafun	0.02321	0.002285	0.545309	1.12002
2	G	3	CO-1686 (A	0.023392	0.004377	0.509569	1.075451
10	D	10	Raloxifene	0.026277	0.001894	0.578958	1.018621
1	E	3	Cediranib (	0.026739	0.00104	0.558996	1.022222
1	D	8	LY2835219	0.026845	0.001383	0.551185	1.086418
1	H	2	Epirubicin	0.027999	0.00022	0.912692	1.000494
13	G	1	Cyclospori	0.031121	0.002339	0.524669	1.087002
9	E	3	Amiodaron	0.03182	0.002961	0.837838	1.007466
6	H	11	Dinaciclib (	0.032924	0.002786	0.663984	1.078891
13	F	3	Chloroxine	0.035536	0.003397	0.577688	1.107189
7	D	1	Isradipine (	0.039218	0.00404	0.615461	1.091102
3	G	5	Vortioxetir	0.042108	0.002727	0.608762	1.090515
5	A	5	Fluvastatin	0.050054	0.005574	0.555306	1.110813
1	C	9	Posaconazo	0.051682	0.00386	0.767643	1.125232
9	D	11	Ciclopirox	0.054363	0.003008	0.517297	1.009476
15	C	10	Doxercalcif	0.055609	0.005686	0.541842	0.544259
9	E	10	Trifluopera	0.056782	0.004016	0.699459	1.01352
4	E	10	Abirateron	0.064772	0.003719	0.564874	1.119405
15	A	1	Tioconazol	0.071615	0.015376	1.172185	0.072057
9	E	7	Nitrendipir	0.072115	0.003323	0.864865	1.012

5	A	7	Lovastatin	0.082796	0.006712	0.547887	0.051421
16	E	5	Paroxetine	0.090967	0.015031	1.087638	0.01146
15	B	1	Vitamin D3	0.096781	0.00863	1.048766	0.001255
13	B	5	Triclabenda	0.09736	0.005545	0.620029	9.97E-06
7	A	9	Duloxetine	0.097981	0.013281	0.991389	1.036885
18	F	9	Amodiaqui	0.100704	0.011124	0.640441	0.002692
16	D	1	Aripiprazol	0.100735	0.013726	0.627768	0.004359
9	F	2	Cabozantin	0.102637	0.003553	0.511892	1.015554
18	A	11	Thonzoniur	0.103504	0.004362	0.604902	3.06E-06
20	C	2	Triclosan	0.106199	0.010286	0.523446	0.001269
9	G	2	Manidipine	0.108566	0.004229	1.035676	1.017085
9	G	3	Manidipine	0.109555	0.003831	0.812973	1.016064
9	F	7	Flunarizine	0.113322	0.008289	2.629189	1.021191
13	E	6	Butoconazo	0.116023	0.007817	0.692194	0.000274
9	F	6	Amlodipine	0.124006	0.008264	1.385405	1.021706
2	C	4	Masitinib (/	0.133853	0.020828	1.308014	0.007066
6	H	1	Clotrimazo	0.138853	0.021956	1.057112	0.00397
17	C	9	Ciclesonide	0.140872	0.016726	0.512541	0.001288
18	B	8	Desogestre	0.145321	0.013193	0.665686	0.000563

n background q-values

Compound	Reads Counts per second				Relative Recombinant l	% activity in Vero 100PFU screen
(+)-Catechi	362209	357899	366108	367703	0.9620636	160
17-AAG (K	339450	347084	343744	348179	0.912130114	209
Abiraterone	279631	289523	288897	289231	0.759160759	6
Alogliptin E	316906	332493	331161	323679	0.863019788	166
Alosetron H	328498	338671	337783	339698	0.889759897	177
Alvimopan	338215	350063	350082	352328	0.920223412	169
Amiodaron	364410	365102	365644	366409	0.967122986	5
Amodiaqui	351271	347518	344113	339282	0.914596284	12
AN-2690	329173	329953	332892	324160	0.870919869	171
Anidulafun	365229	376574	364202	366012	0.974039114	5
Ascomycin	312887	325360	316950	314498	0.840161895	171
AT13387	289397	302902	309451	303901	0.797783743	191
Atovaquon	224278	226697	224646	216287	0.590178835	4
Baicalin	353023	358945	357158	357276	0.943855499	164
Baricitinib p	337511	341167	346435	349331	0.909474697	199
Bedaquilin	348601	355872	347079	351646	0.928501326	4
Bedaquilin	351974	352561	351293	338630	0.922718035	4
Belinostat	351877	362403	352055	355423	0.940782547	5
Benidipine	50508	49563	49185	49177	0.131303853	2
Bivalirudin	309590	311058	307409	301908	0.813872407	172
Boceprevir	341877	348889	346623	349417	0.917654679	178
Cabozantir	333265	320522	329153	328687	0.867908455	12
Caffeic aci	390077	320522	399891	403178	1.053990126	173
Cediranib (	350524	320522	354116	362844	0.943901818	3
Cetirizine	170134	320522	168670	171706	0.448561006	161
Chloroxine	380665	320522	374596	374884	0.997545476	6
Chlorprothi	359664	320522	361391	361186	0.957434982	4
Ciclopirox e	358391	320522	365653	361119	0.961032665	7
Cilnidipine	28828	320522	28106	27971	0.075210567	1
CO-1686 (/	327566	320522	324163	319872	0.855087284	3
Cyclospori	373363	320522	381885	373048	1.000139355	6
Daidzein	339091	320522	350216	346126	0.915372462	161
Daunorubi	319222	320522	327409	328432	0.86150581	5
Dinaciclib (	345056	320522	342151	346073	0.913362207	5
Doxercalcif	361602	320522	370984	356613	0.961268894	8
Duloxetine	297459	320522	293652	301826	0.789374812	12
Ebastine	323732	320522	324560	326603	0.857585216	4
Eltrombopa	331875	320522	331553	322913	0.866618795	4
Epirubicin I	298948	320522	302560	310338	0.802456032	3
Ethamsylat	388357	320522	391647	386776	1.028650844	167
Fingolimod	354171	320522	356587	351555	0.939334077	3
Fluoxetine	317425	320522	314163	307707	0.830595645	2
Fluvastatin	344916	320522	339204	335899	0.899431358	6
Idarubicin I	327538	320522	325691	318222	0.860993651	2
Isradipine (	19176	320522	18561	18431	0.049588738	6
KPT-330	378225	320522	389989	381493	1.021585172	4
Lacidipine	61246	320522	60310	59492	0.160064805	1
Levodopa	304150	320522	322065	316577	0.839019133	233
L-Glutamin	349150	320522	361792	350740	0.941896194	204

Lithium Citr	343238	320522	350985	343490	0.921697688	205
Lithocholic	350340	320522	350189	353302	0.934238297	205
Lovastatin	351803	320522	353101	349914	0.931862781	9
LY283521!	345999	320522	345910	348685	0.918872213	3
LY335979	352212	320522	356783	351420	0.936199587	3
Manidipine	249924	320522	250350	251205	0.661459544	12
Manidipine	242466	320522	242968	239469	0.638229111	12
Meloxicam	308460	320522	309204	312953	0.824020956	178
Mepenzola	330221	320522	339233	336213	0.889583222	175
Methyldopa	361429	320522	364798	367357	0.966613475	175
NAD+	385544	320522	386401	376264	1.016345802	164
Nevirapine	371882	320522	379254	373316	0.993240432	187
Nitrendipin	26271	320522	26096	26141	0.069146715	9
Noscapine	358232	320522	362883	354994	0.950175431	213
Ospemifen	356755	320522	358779	352991	0.940726302	4
Oxethazair	330534	320522	331517	321724	0.873612341	167
Pamidrona	309519	320522	313471	313941	0.831273891	2
Panobinos	330994	320522	343634	341728	0.896255842	4
Paroxetine	360169	320522	369849	368856	0.973172282	12
Posaconaz	325406	320522	324458	315077	0.856451055	6
Probucol	337686	320522	341300	339611	0.902269406	5
Raloxifene	334037	320522	331016	335545	0.88121267	6
Ranolazine	359507	320522	367539	367791	0.967918354	167
Ruxolitinib	314540	320522	317488	316727	0.838138405	169
Salinomyci	365050	320522	373342	367850	0.977863099	4
Tioconazol	349867	320522	345556	349383	0.922076183	10
Triclabenda	335937	320522	336143	339544	0.892308118	12
Trifluopera:	358281	320522	348422	360671	0.939701323	7
Vitamin D3	370629	320522	360711	364312	0.968153921	12
Vortioxetin	326546	320522	329264	332965	0.87317297	6
Vortioxetin	280784	320522	284246	284019	0.747903855	1
DMSO	356875	320522	360022	355819	0.945032008	100
DMSO1	365885	320522	366954	370334	0.969202059	100
DMSO2	387195	320522	397670	396896	1.042738516	100
DMSO3	382401	320522	382402	379669	1.012664744	100
DMSO4	390492	320522	388004	391251	1.030362672	100

Compounds that inhibited SARS-Cov2-ΔOrf7a-Nluc replication in Vero and HUH7 screens

HUH7	Row	Column	Drug name	Mean relati	SD	PrestoBlue
9	E	3	Amiodarone HCl	0.034035	0.002111	0.963298
9	F	6	Amlodipine Besylate	0.033811	0.002081	0.68594
18	F	9	Amodiaquine dihydrochlorid	0.012584	0.000224	0.778309
16	D	1	Aripiprazole	0.008694	0.001695	0.741567
15	C	1	Atovaquone	0.058244	0.010118	1.150489
17	D	4	Bedaquiline fumarate	0.108424	0.006709	0.522411
9	*F*	*4*	*Benidipine HCl*	*0.00382*	*0.0012*	*0.66087*
9	F	2	Cabozantinib malate (XL184)	0.03412	0.003109	0.801129
20	A	9	Chlorprothixene (hydrochlori	0.031761	0.003566	0.762326
9	D	11	Ciclopirox ethanolamine	0.046013	0.000597	0.502315
1	*B*	*10*	*Cilnidipine*	*0.00182*	*0.00039*	*0.79704*
13	G	1	Cyclosporin A	0.080205	0.007228	0.506995
16	G	5	Daunorubicin HCl	0.006058	0.001727	0.518582
6	H	11	Dinaciclib (SCH727965)	0.062944	0.000228	0.593445
7	A	9	Duloxetine HCl	0.028451	0.00042	0.504807
12	A	10	Ebastine	0.015939	0.00032	1.613674
9	F	7	Flunarizine 2HCl	0.109125	0.006023	0.968944
1	H	1	Fluoxetine HCl	0.080571	0.001764	0.91596
5	A	5	Fluvastatin Sodium	0.05448	0.002919	1.301701
22	A	4	Homoharringtonine	0.013304	0.003262	0.679424
7	*D*	*1*	*Isradipine (Dynacirc)*	*0.00135*	*0.00094*	*0.67618*
10	A	5	KPT-330	0.058589	0.000707	0.710223
9	*G*	*1*	*Lacidipine*	*0.02662*	*0.00129*	*0.92468*
5	A	7	Lovastatin	0.087582	0.002308	1.075407
1	D	8	LY2835219	0.072422	0.002963	0.604272
8	B	3	LY335979 (Zosuquidar 3HCL)	0.082311	0.009091	0.620331
9	G	2	Manidipine	0.100031	0.002327	0.539582
9	G	3	Manidipine 2HCl	0.044466	0.002178	0.572558
2	C	4	Masitinib (AB1010)	0.039172	0.000431	0.715617
9	*E*	*7*	*Nitrendipine*	*0.00073*	*0.00059*	*0.65455*
18	A	1	Ospemifene	0.144859	0.002836	1.203434
6	A	7	Panobinostat (LBH589)	0.058971	0.000949	0.753019
10	D	10	Raloxifene HCl	0.036459	0.000553	0.5413
15	A	1	Tioconazole	0.124683	0.020143	1.128639
15	B	1	Vitamin D3	0.025671	0.009357	0.641766

Red indicates Nluc inhibitors