Territory-level temperature influences settlement decisions, breeding phenology and productivity

in three forest passerine birds

Jack D Shutt[1,2,*], Sophie C Bell[1], Fraser Bell[3], Joan Castello[4], Myriam El Harouchi[1], and Malcolm D

Burgess[1,4,5]

1. Piedfly.Net, Yarner Wood, Bovey Tracey, Devon TQ13 9LN

2. Department of Natural Sciences, Manchester Metropolitan University, Manchester, M1 5GD, UK

3. Centre for Ecology and Conservation, Cornwall Campus, University of Exeter, Penryn, Cornwall,

TR10 9FE, UK

4. Centre for Research in Animal Behaviour, University of Exeter, Perry Road, Exeter, Devon, EX4

4QG

5. RSPB Centre for Conservation Science, The Lodge, Sandy, Bedfordshire, SG19 2DL, UK

* Corresponding author: jackshutt8@gmail.com

Abstract

Temperature is a key determinant of breeding phenology in many temperate zone species, with

increasing spring temperature across years or large geographic gradients known to advance the

timing of breeding. The effect of localised, territory-scale temperature variations is less well

understood, primarily due to a lack of fine-scale temperature data. Here, we interpolate spring

temperatures recorded at intersections of a $150m^2$ grid encompassing 417 ha of forest in South-

West England 2015-18 to examine the influence of territory-scale temperature, and its interaction

with mean annual temperature, on territory selection, breeding phenology and productivity of three

co-occurring forest passerine bird species using data from 672 nests. Two of these species are long-

distance migrants (pied flycatcher *Ficedula hypoleuca* and wood warbler *Phylloscopus sibilatrix*) and

one is resident (blue tit *Cyanistes caeruleus*); all species are dependent upon seasonal invertebrate

food for reproductive success. All three species exhibited significant trends in reproductive traits

associated with territory-scale temperature. Pied flycatchers settled in cooler territories first, where

they raised more fledglings. Blue tits laid larger clutches in warmer territories in warm years and

always laid earlier at warmer territories irrespective of annual mean temperature. Contrastingly,

migrant pied flycatcher and wood warbler breeding phenology was earlier at warmer territories in

cool years and cooler territories in warm years, with wood warbler clutch size responding similarly to

this interaction. Greater previous breeding experience and increased territory quality also predicted

earlier breeding phenology and higher productivity for pied flycatchers. We suggest that the two

migrant species may be best synchronised with their prey requirements in cooler years at a

population level while resident blue tits matched local phenology at lower trophic levels across all

years, which is potentially advantageous under warmer predicted climate change scenarios.

Keywords: Local temperature, trophic mismatch, blue tit, pied flycatcher, wood warbler, breeding

experience, territory quality, nestbox

<u>**Introduction**</u>

Temperature plays a key role in determining the timing of breeding of temperate zone bird species

(Crick and Sparks 1999, Dunn and Møller 2013). In warmer years, and at generally warmer locations,

breeding phenology occurs earlier, and under recent climate change average breeding phenology

has advanced over time for many species (Parmesan and Yohe 2003, Both et al. 2004, McLean et al.

2016, Phillimore et al. 2016). An increase of 1°C has been demonstrated to advance breeding

phenology by 3-5 days in a range of birds (McLean et al. 2016, Phillimore et al. 2016, Socolar et al.

2017). Functionally, this trend to breed earlier is to maintain temporal synchrony with key prey

resources (Burgess et al. 2018). Prey synchrony is particularly important for single-brooded

insectivorous birds (e.g. tits *Paridae*, flycatchers *Ficedulidae*, leaf warblers *Phylloscopidae*) breeding

in highly seasonal environments such as deciduous forests (Visser et al. 1998, Burgess et al. 2018).

Asynchrony with this prey peak can reduce fledgling number and quality, negatively impacting

fitness, with such asynchrony termed trophic mismatch (Both et al. 2006, Miller-Rushing et al. 2010).

While the effect of average temperature on breeding phenology across years and geographic

gradients is well established, effects of subtler localised intra-annual temperature variation is less

understood. This is primarily due to a lack of studies collecting fine-scale within-site temperature

data alongside bird breeding data, despite this being the scale relevant to individual birds (Hinks et

al. 2015). It is therefore unknown how territory settlement, breeding phenology and breeding

outcome are influenced by territory-scale temperature. It has been shown that within-nest

temperature can modify breeding phenology (Simmonds et al. 2017), and that fine-scale

temperature is a strong predictor of localised species distribution (Frey et al. 2016). Phenological

timings that may be influenced by territory-scale temperature variation include the date of arrival,

date of nest building initiation and first egg laying date, while breeding parameters that may be

sensitive include clutch size and productivity. In the same breeding locations resident species with

similar ecological requirements to long-distance migrants often commence breeding earlier

(Källander et al. 2017, Samplonius et al. 2018). With a warming spring climate this difference can

become more divergent (Samplonius et al. 2018, Haest et al. 2020) suggesting fine-scale

temperature may determine breeding onset differently between migrants and residents as well as

differing patterns of territory occupancy at a local scale.

Understanding the influence of temperature on territory occupation, breeding phenology and

outcome could indicate how optimally timed a population is. As tree and caterpillar phenology is

well synchronised with fine-scale temperature (Buse et al. 1999, van Asch et al. 2012), locally

warmer territories will show advanced phenology in lower trophic levels than cool territories (Cole

et al. 2015, Hinks et al. 2015). If breeding always commences earlier in warmer territories this could

indicate successful thermal niche tracking, where a well-timed population is able to track the local

phenology of lower trophic levels (Dobrowski 2011, Socolar et al. 2017). However, if within a

woodland breeding occurred earlier in cooler territories relative to warmer territories with earlier

prey phenology, then the average population consequence would be later than optimal timing as

late breeders (either through late arrival, lower quality or competition) will be forced to occupy the

remaining warmer territories with the greatest levels of mismatch. If breeding commences earlier in

warmer territories in cool years, and in cooler territories in warm years, this would suggest that

average population timing is optimal in cool years but suboptimal (late) in warm years.

Examining a population's breeding phenology across a temperature gradient requires few

assumptions to be made about precise optimal dates or specific prey items compared to studies

focussed on dates. Optimal synchrony between bird species breeding in the same locality may vary

due to different prey requirements and developmental needs of young. The ephemeral invertebrate

peak in deciduous woodland that several single-brooded passerine species are commonly assumed

to depend on are moth caterpillars (Visser et al. 1998, Burgess et al. 2018). However, the most well

studied species of this system (blue tit *Cyanistes caeruleus*, great tit *Parus major* and *Ficedula*

flycatchers) breed at overlapping but different times to one another (Burgess et al. 2018, Samplonius

et al. 2018), and so within any single year and locality breeding timing across these bird species will

encompass peaks of different prey species at different nestling developmental stages. Timing

breeding to match peak demands of nestlings to the caterpillar peak can have little impact on

fledging mass or productivity in some species (Maziarz and Wesołowski 2010, Mallord et al. 2017,

Wesołowski and Neubauer 2017), while it does in others (Visser et al. 1998, Both et al. 2005),

suggesting differences in prey requirements. Whether breeding is timed to the same or different

optima, a preference for breeding in cooler territories first would indicate local population-level

mistiming. It would also suggest that the birds were adjusting territory selection and breeding

phenology due to the local phenological stage (Hušek et al. 2014, Hinks et al. 2015, Shutt et al.

2019), in addition to purely relying on temperature as a causal cue of breeding time (Caro et al.

2013, Verhagen et al. 2020).

In this study, we investigate how territory-scale temperature variation over four years within a

417ha woodland affects territory settlement, breeding phenology, clutch size and productivity of

three single-brooded co-occurring insectivorous woodland passerine bird species: pied flycatcher

Ficedula hypoleuca, blue tit and wood warbler *Phylloscopus sibilatrix*. All three are similarly sized,

primarily single-brooded, spring-breeding species. We hypothesise that as these migrants tend to

breed later than residents (Slagsvold 1976, Samplonius et al. 2018), and later breeding in warm years

increases mismatching (Burgess et al. 2018), that in warm years migratory species will be less

optimally timed compared to resident species, selecting cooler territories first and showing lower

productivity in warm territories.

Material and methods

This study was conducted at East Dartmoor National Nature Reserve in Devon, South-West England,

UK (50.60°N -3.73°E) over four breeding seasons 2015-2018. Habitat across the reserve is relatively

uniform and dominated by sessile oak *Quercus petraea* with a European holly *Ilex aquifolium,* and

rowan *Sorbus aucuparia* understory and a European bilberry *Vaccinium myrtillus* field layer. The

study area extends linearly approximately 5 km along a steep-sided river valley in the north, with a

150ha woodland containing two shallow stream valleys in the south, together resulting in slopes at

different aspects and an altitudinal range of 60-300m.

Breeding bird data

Throughout the study area, 309 wooden nestboxes with a 32 mm entrance hole were available each

year and monitored annually throughout the bird breeding season April-June. Nestbox checks were

made at least weekly and more frequently at some nest stages to determine species occupancy,

date of first nesting material, date of first egg laying, completed clutch size and the number of

successful fledglings (productivity) for each breeding attempt. Date of first nesting material was

taken as the date of the first nest check that found fresh nesting material for nests that ultimately

had a clutch of eggs laid in, with nests found half complete (cup shape evident) or more over the

maximum 7 days inspection interval assigned the mid-point between the two visits. All species were

assumed to lay one egg per day (Perrins 1979, Lundberg and Alatalo 2010), with the first egg date

back-calculated from incomplete clutches recorded at nest inspections. Clutch size records the total

number of eggs laid that were incubated, and the number fledged was the number alive at the last

visit pre-fledging minus the number found dead at the first visit post-fledging.

The date of spring arrival by territorial male pied flycatchers to nestboxes was also recorded from

the first week in April. Territorial activity, such as singing, alarm calling and inspection of nestboxes,

was observed on alternate mornings, and the presence of, colour and position of any rings and

variable plumage characteristics such as forehead patch size were recorded. These were cross-

referenced against males captured at the same nestboxes later in the breeding season. Male arrival

date to a territory was taken as the mid-point between the date of the survey prior to the first

observation of the male and the date of the survey when the male was first observed. This is a

common method used for pied flycatchers (Bell et al. 2017, Potti et al. 2018) and its accuracy has

been verified with geolocator tracking data (Both et al. 2016). Nearly all breeding male and female

pied flycatchers were captured annually and are marked with a unique metal numbered ring upon

first capture, enabling the frequency of breeding over an individuals' lifetime to be determined,

including for years prior to 2014 using data collected prior to the present study. Between 2015-2018,

320 pied flycatcher and 315 blue tit nesting attempts were monitored.

Wood warbler nests were also monitored throughout the breeding seasons 2015-2018. Wood

warblers are ground nesting and were located by observing parental behaviour, with the location of

each nest determined by GPS and visited every 3-4 days. First egg-laying dates were determined

from visits during egg laying assuming that one egg is laid per day. A total of 37 wood warbler nests

were monitored during the study period.

Temperature data

Between 2015-2018 96 temperature dataloggers (Thermachron iButton DS1921G) were placed

throughout the study area at each intersection of a 150m grid. Each was attached to a tree 1.5m

above ground level in a waterproof standard sized plastic container with three drilled holes to

provide air circulation (Shutt et al. 2019). Temperature dataloggers recorded temperature hourly to

an accuracy of 0.5°C. Additionally, 22 higher-sensitivity temperature dataloggers (Thermachron

iButton DS1922L) that record temperature to an accuracy of 0.0625°C were attached alongside 22 of

the lower resolution dataloggers on the same trees with the same aspect in a non-random way to

maximise coverage of the entire study area.

We used temperatures recorded between Julian days 97 and 132, which encapsulates the spring

temperature sensitivity window previously estimated for blue tit and pied flycatcher egg laying for

the study area (Samplonius et al. 2018). Data from temperature dataloggers with more than 16

missing hourly data points within the focal period within a year were discarded as incomplete (n=25,

from a total of 333). Additionally, data from temperature dataloggers recording at least one

measurement beyond the extreme credible temperature bounds within the focal period (<-8°C or

>35°C) were discarded for the year in question as unreliable or faulty (n=9). As all faulty dataloggers

were the higher sensitivity model, and data from high sensitivity dataloggers are not directly

comparable with data from low sensitivity dataloggers, the unaffected higher resolution datalogger

data were used to determine the accuracy of their paired low sensitivity datalogger only and were

excluded from all other analyses (n=56). Paired high sensitivity and low sensitivity dataloggers gave

very similar readings (interquartile range <0.2°C despite recording on different sensitivities with a

possible correct error range of 0.25°C), whereas the variation across the entire study area was much

higher (interquartile range >0.6°C in each year), providing confidence that local temperature

differentials across the study area correspond to genuine local temperature variation rather than

datalogger inaccuracy. Temperature means across the focal period were used and a total of 257

datalogger mean readings remained over the four-year period after this quality control (87 in 2015,

54 in 2016, 56 in 2017, and 60 in 2018).

All spatial analysis was conducted within R version 3.6.3 (R Core Team, 2018) using the raster, gstat

and tmap libraries. The data from each remaining low sensitivity temperature datalogger within

each year was then projected to its geographical location and temperature across the study area

interpolated using inverse distance weighting (Gräler et al. 2016) raised to the power two. The

inverse distance squared weighting power was decided upon after investigating the ability of the

data to predict known temperature points in a 'leave one out' validation routine of powers between

one and four (root-mean-squared-error of P1 = 0.467, P2 = 0.447, P3 = 0.453, P4 = 0.461).

Temperature differentials for each nestbox and wood warbler nest location were then extracted

from the resultant heatmap raster (Gräler et al. 2016). On inspection, the heatmap differential

rasters for each year were highly consistent in both spatial pattern and variance and so an

interannual differential mean was used which we term the local temperature differential. This allows

determination of whether territories were in a relatively cool or warm area within the overall study

area and afforded greater differential consistency as not every temperature datalogger position

recorded data in all years.

Statistical analysis

General linear models (GLM) were constructed to assess the effects of local temperature differences

on measures of breeding success for this system. Not all breeding parameters were assessed as a

response variable for each of the three bird species due to lack of data. Five breeding parameters

were used: date of male arrival to breeding territory and date of nest building initiation (both pied

flycatcher only), date of first egg laying, clutch size and productivity (number of fledglings) (all three

species). All dates are reported as April days (days from the 1st April, e.g. 25th April = day 25). For

wood warblers, as nests were often found discovered after eggs had been laid, if date of first egg

laying was not recorded but date of hatching was, first egg date was inferred by subtracting the

mean number of days (n=18) between first egg date and hatching date across those nests where

both were recorded, from hatching date where first egg date was not recorded. All analyses were

conducted in R version 3.6.3 (R Core Team 2018).

Response variables were tested by Bayesian GLMs (Hadfield 2010) with annual mean temperature,

local temperature differential, territory quality, previous breeding experience and the interaction

between annual and local temperature as fixed predictor variables. Continuous predictor variables

were mean centred for ease of interpretation and model functioning (Schielzeth 2010). Each model

was run for 100,000 iterations with the first 10,000 discarded as burn-in with a Gaussian error

structure, with both response data and model traces visually inspected to ensure the fit was

acceptable. Year was not included as a random predictor variable as it covaries with annual mean

temperature for each data point, however it was included for additional, unreported models for

each response to ensure that its' omission did not significantly influence results.

Territory quality was inferred from the proportion of years each nestbox had been occupied by

nesting pied flycatchers or blue tits out of the total number of years each nestbox was available

since 1980 and until 2018. Long-term nestbox/territory occupancy patterns have previously been

shown to provide a clear correlate of territory quality, with more frequently occupied territories of

higher quality (Sergio and Newton 2003, Potti et al. 2018). We used the occupancy rate since 1980

as both nestbox number and bird population size had by this year stabilised after nest box

provisioning started incrementally from 1955. Territory quality could not be included as a predictor

variable for wood warbler as this species does not use nestboxes and therefore territories could not

be ascribed long term occupancy rates. Individual breeding experience was defined as the number of

years an individual had previously been captured breeding in the population. Previous breeding

experience could not be included as a predictor variable for blue tit or wood warbler due to

incomplete data on parental identity. For pied flycatcher date of arrival, male breeding experience

was included as a fixed effect, whereas in all other models female breeding experience was used, as

females build the nests, lay the eggs and provide the majority of parental care in pied flycatchers

(Lundberg and Alatalo 2010). Pairwise correlation between fixed effects was assessed and as all

were <0.18 this suggested no issue with colinearity within the models, with the most significant

correlation being that higher quality territories tended to have cooler local temperature

differentials.

Results

Local spring temperature varied by around 2°C from the coldest to warmest areas across the study

area with colder areas at the higher elevations of Yarner Wood in the south and Neadon Cleave in

the north, and warmer areas at lower elevations along the River Bovey in Lustleigh Cleave (Fig 1).

This pattern was spatially consistent across all four years. Variation in mean spring temperature

between years was observed, with 2018 the warmest (mean = 10.7°C) and 2016 the coolest (mean =

9.2°C).

Male pied flycatchers returning from migration in spring occupied the coolest territories first,

irrespective of the relative mean annual temperature, with our model predicting arrival to the

coolest territories on 12th April and warmest territories on 20th April (Table 1, Fig 2a). There was also

an interaction effect of local temperature on the timing of nest building initiation; in cool years nests

were initiated significantly earlier at warmer territories, with this pattern reversed in warm years

with nests initiated earlier at cooler territories (Table 1, Fig 2b). This temperature effect on the

timing of nest building did not translate into an effect on the timing of egg laying or clutch size

(Table 1). Productivity varied by 1.7 chicks per nest attempt across the temperature gradient, with

productivity higher at the coolest territories compared to the warmest territories (5.6:3.9 fledglings

per attempt respectively) (Table 1, Fig 2c). This latter effect was strongest is cool years, while in

warm years productivity was similar irrespective of the local temperature (Table 1, Fig 2d).

For resident blue tits, higher local temperatures predicted a significant advancement of five days in

the timing of egg laying between the coolest and warmest territories (Table 1, Fig 2e). There was

also an interaction effect of local temperature on clutch size, with cooler territories predicted to

have substantially larger clutches in cool years, while in warm years warmer territories were

predicted to have slightly larger clutches (Table 1, Fig 2f). There was no significant effect of local

temperature on blue tit productivity, although the interaction effect was in the same direction as for

clutch size (Table 1). In warm years, blue tits laid eggs earlier (-2.22 days/°C) and had higher

productivity (+0.78 fledglings/°C) (Table 1).

There were significant effects of local temperature interactions on migratory wood warbler breeding

parameters. In cooler years, eggs were predicted to be laid substantially earlier at warmer territories

than cooler territories, whereas in warmer years the opposite was true, with eggs laid earlier at

cooler territories (Table 1, Fig 2g). Wood warbler clutches were also significantly larger at cooler

territories in warmer years, and at warmer territories in cooler years (Table 1, Fig 2h). There was no

significant effect of local temperature on wood warbler productivity (Table 1).

Previous breeding experience also affected pied flycatcher breeding parameters. Males with more

experience arrived earlier (-0.81 days/year experience, Table 1, Fig 3a) and females with more

experience laid larger clutches (+0.17 eggs/year experience, Table 1, Fig 3b) earlier in the year (-0.93

days/year experience, Table 1, Fig 3C). In higher quality territories pied flycatchers also initiated

earlier nest building (21st April: 1st May, Table 1, Fig 3d) and egg laying (6th May: 12th May, Table 1,

Fig 3e) and produced more fledglings (3.9: 5.4 fledglings per attempt, Table 1, Fig 3f), with figures at

the two extremes of territory quality found in our study. Territory quality showed no significant

relationship with blue tit breeding parameters (Table 1).

Discussion

This study demonstrates multiple relationships resulting from territory-level spatial variation in

temperature to differences in territory occupancy, breeding phenology, clutch size and productivity

for three insectivorous woodland birds. The patterns show that within a 417ha woodland both

migratory and resident species either select breeding territories where territory temperature best

matches breeding phenology, with these locations differing between cold and warm years, and/or

show fitness benefits in relation to the temperature of the territory occupied. In cold years, when

vegetation and invertebrate phenology is expected to be latest (Burgess et al. 2018), pied flycatcher

nest building and wood warbler egg laying commenced earlier in warmer territories, with larger

clutch sizes for wood warbler and blue tit. In warm years, when vegetation and invertebrate

phenology is earlier (Burgess et al. 2018), cooler territories were preferred by the migratory species',

being associated with earlier nest building by pied flycatchers, and earlier egg laying and larger

clutch sizes in wood warblers. Territory-scale temperature also influenced pied flycatcher

productivity across years, with 1.7 more fledglings predicted per nest in the coolest territories

compared to the warmest. This pattern was driven by considerably higher productivity in cool

territories in cooler years, suggestive of a negative effect on productivity from trophic mismatch in

relatively warmer territories in cool years.

Temperature impacts on breeding phenology

Male pied flycatchers returning from Africa settled in cooler territories first, with nest building also

beginning first in cooler territories in warm years, with this pattern reversed in cold years. Our

measure of arrival was only for males as these are more conspicuous, but our method of estimating

nest building initiation is frequently used as a proxy for female arrival date (Potti and Montalvo

1991, Visser et al. 2015) as pairing and nest building can commence on the day of female arrival

(Potti and Montalvo 1991). Making that assumption, our results suggest that both male and female

pied flycatchers may partly base territory selection on territory temperature, with the territories

selected first by the earliest arriving birds those most likely to have a better match with prey

phenology. In warmer years migration timings may constrain earlier arrival (Both and Visser 2001)

with the implication that in warm years later arrivals are constrained to breed in the warmest

territories and be more likely mismatched with prey. In cooler years the local population overall is

therefore better phenologically matched.

Territory and female mate selection is well studied, with males establishing territories first and

females selecting breeding territories held by males based on either habitat quality (Alatalo et al.

1986), or male traits (Sirkiä and Laaksonen 2009). Territory choice can be further influenced by

information obtained from heterospecifics, for example from cues related to occupancy and density

patterns of resident species (Forsman et al. 2008) or inspecting multiple nest sites upon arrival

(Schuett et al. 2017). Our study suggests fine-scale temperature is another key determinant of

territory choice in pied flycatchers.

Egg laying date in our three study species is constrained by photoperiod (Lambrechts and Perret

2000, Caro et al. 2007) and environmental factors (Shutt et al. 2019), and is strongly related to

spring temperature at larger spatial scales (Both and Visser 2001, Schaper et al. 2011, Phillimore et

al. 2016), with laying starting later at cooler locations such as higher latitudes and elevations (Sanz

1998, Burgess et al. 2018), but examination at local scales is rare. We found spatial variation in laying

date with regards to local temperature for both wood warbler and blue tit. Wood warblers started

egg laying substantially earlier at warmer territories in cooler years and cool territories in warmer

years. Previous work on wood warblers showed a strong relationship between the temperature of

the pre-laying period and laying date at larger spatial scales, with earlier laying associated with

higher pre-laying temperatures (Wesołowski and Maziarz 2009). We find the same pattern at the

territory scale in cool and average years, providing further support that temperature is a cue to

wood warbler laying. However, in warm years the opposite pattern occurs, which may indicate

phenological mismatch within the local population in warm years as later breeders are occupying the

warmest (and thus most phenologically advanced) territories, possibly due to these being the only

ones remaining once earlier breeders have occupied the better-matched, cooler territories.

Higher territory temperature linearly predicted earlier egg laying in resident blue tits, similar to that

found across longer gradients of space and time but with a slightly shallower slope to other

estimates (McLean et al. 2016, Phillimore et al. 2016). Our results are consistent with temperature

being a direct cue for blue tit breeding phenology (Visser et al. 2009, Shutt et al. 2019, Verhagen et

al. 2020) and that this cue applies at a very localised scale. No significant temperature relationship

for lay date was found for pied flycatcher, suggesting temperature may not be a direct cue to egg

laying in this species. Pied flycatcher egg laying date shows less spatial and inter-annual variation

compared with arrival phenology (Nicolau et al. 2021), indicating that in years with early arrival,

individual birds can nest build well in advance of egg laying. Our results on the effects of territory

scale temperature on breeding phenology suggest that warming springs may be more favourable to

resident blue tits than migrant pied flycatchers and wood warblers at this locality.

Temperature impacts on clutch size

We found spatial variation related to temperature in clutch size for wood warbler and blue tit, but

not for pied flycatcher. Wood warbler clutch size followed a similar pattern to the breeding

phenology of this species with regards to local temperature, being larger at cool territories in warm

years, and at warm territories in cool years, with clutch size therefore larger in territories selected

earlier for breeding. The effect of territory temperature on clutch size may therefore help to explain

why wood warbler clutch size declines seasonally (Grendelmeier et al. 2015). Although wood

warblers do not time breeding to match caterpillar peaks (Wesołowski and Maziarz 2009, Maziarz

and Wesołowski 2010), if higher clutch sizes are indicative of better phenological synchrony to their

own invertebrate prey peaks these results imply that wood warblers may be late matched in warm

years and search for their optimal territory level temperature locally (Frey et al. 2016), refined within

territories by nest site preferences such as by slope and distance from forest edge (Pasinelli et al.

2016). Blue tit clutch size showed the opposite interaction, with larger clutches in warm territories in

warm years and cool territories in cool years. Blue tits time breeding to match caterpillar

phenological peaks (their principle prey) (Wesołowski and Rowiński 2014, Shutt et al. 2020), and our

results indicate that blue tit breeding phenology might be early in cool years and better matched

birds breed in cooler territories. As blue tit clutch size does not vary with woodland habitat or

geography but by annual variation (Shutt et al. 2018), annual temperature could be an important

factor determining the quality of the breeding year for blue tits.

Temperature impacts on productivity

There were no territory-level temperature effects on productivity in either blue tit or wood warbler,

but pied flycatchers showed a strong response, fledging 1.7 more chicks in cool compared to warm

territories, with this effect strongest in cool years. Higher productivity from cooler territories

suggests cooler territories have a better match with invertebrate prey abundance in this species.

Cooler territories conferring greater productivity may also explain why these territories are occupied

first by both male and female pied flycatchers upon arrival. There may be a greater risk of mistiming

breeding in warmer springs due to lower levels of local variation in tree budburst (Denéchère et al.

2019).

Non-temperature impacts: territory quality and individual experience

For pied flycatcher our models also controlled for other factors that may influence nest building:

clutch size and productivity. Higher territory quality predicted earlier nest initiation and egg laying,

and higher productivity, supporting the general validity of using the simple indirect proxy of long-

term occupation trends to indicate territory quality (Sergio and Newton 2003, Potti et al. 2018),

particularly for measuring territory quality in relatively homogeneous habitats when habitat

composition and structural preferences can be hard to determine or differentiate. The spatial

pattern of temperature across the study area was consistent, and the territories that were relatively

cool in 2015-2018 had the highest pied flycatcher occupancy rates since 1980, showing that cool

territories have been preferentially selected for almost 40 years. The models also indicated that

adult flycatchers with greater previous breeding experience (i.e. older individuals) tended to arrive

earlier in spring (male experience), lay eggs earlier and lay larger clutches (both female experience).

This is concordant with other studies (Potti 1998, Both et al. 2016), and is indicative of more

experienced parents having higher nesting success.

Conclusions

Territory-level temperature variations within a single site have profound effects on the breeding

decisions and outcomes of all three passerine bird species studied. Resident blue tit breeding

phenology responds linearly to increasing territory temperatures similar to responses across large-

scale geographic and annual temperature gradients (Phillimore et al. 2016). Migrant pied flycatchers

and wood warblers, on the other hand, may be responding to small-scale phenological cues at lower

trophic levels, and local populations may be late timed in warm years. Current climate change

scenarios potentially favour resident blue tits that track local phenology well across all years, over

the two migrant species which may be best synchronised in cool years. As avian distributions are

limited by thermal breeding niches (Socolar et al. 2017), this may contribute to shaping breeding

distribution and abundance of these species, with southern populations possibly impacted to a

greater extent by climate change (Both et al. 2006). For populations of declining long-distance

migrant species, including pied flycatcher and wood warbler in the UK (Massimino et al. 2019),

knowledge of territory and thermal preferences and their influence on fecundity and productivity

could be used to help maintain distribution and population size by informing woodland management

and the locations of conservation interventions.

<u>**Author contributions**</u>

JDS, SCB and MDB conceived the study; SCB, FB, JC, MEH and MDB collected the field data; JDS

curated and analysed the data; JDS and MDB wrote the first draft of the manuscript, with SCB and FB

also contributing critically to later drafts; MDB managed the project.

Acknowledgements

We thank Ed Shephard, Amy Vanstone, Mary Wallis, Matt Parkins and Emma Inzani for further

assistance in the field, Steve Cramer for assistance managing the temperature data, and David

Rickwood (Woodland Trust) and Simon Lee (Natural England) for logistical help.

Funding statement

This project was funded by Devon Birds, Kevin Cox and through a Heritage Lottery Fund grant. Wood

warbler fieldwork was undertaken as part of a RSPB project joint funded by Natural England within

the Action for Birds in England partnership.

449
450 **References**
451

Alatalo, R. V. et al. 1986. Female pied flycatchers choose territory quality and not male
characteristics. - Nature 323: 152–153.
Bell, S. C. et al. 2017. No short- or long-term effects of geolocator attachment detected in Pied
Flycatchers Ficedula hypoleuca. - Ibis (Lond. 1859). in press.
Both, C. and Visser, M. E. 2001. Adjustment to climate change is constrained by arrival date in a long-
distance migrant bird. - Nature 411: 296–8.
Both, C. et al. 2004. Large-scale geographical variation confirms that climate change causes birds to
lay earlier. - Proc. R. Soc. B Biol. Sci. 271: 1657–62.
Both, C. et al. 2005. Climatic effects on timing of spring migration and breeding in a long-distance
migrant , the pied flycatcher Ficedula hypoleuca. - J. Avian Biol. 36: 368–373.
Both, C. et al. 2006. Climate change and population declines in a long-distance migratory bird. -
Nature 441: 81–83.
Both, C. et al. 2016. Repeatability in Spring Arrival Dates in Pied Flycatchers Varies Among Years and
Sexes. - Ardea 104: 3–21.
Burgess, M. D. et al. 2018. Tritrophic phenological match–mismatch in space and time. - Nat. Ecol.
Evol. in press.
Buse, A. et al. 1999. Effects of elevated temperature on multi-species interactions: the case of
Pedunculate Oak, Winter Moth and Tits. - Funct. Ecol. 13: 74–82.
Caro, S. P. et al. 2007. Non-photoperiodic factors and timing of breeding in blue tits: Impact of
environmental and social influences in semi-natural conditions. - Behav. Processes 75: 1–7.
Caro, S. P. et al. 2013. The case of the missing mechanism: how does temperature influence seasonal
timing in endotherms? - PLoS Biol. 11: e1001517.
Cole, E. F. et al. 2015. Predicting bird phenology from space: Satellite-derived vegetation green-up
signal uncovers spatial variation in phenological synchrony between birds and their
environment. - Ecol. Evol. 5: 5057–5074.
Crick, H. Q. P. and Sparks, T. H. 1999. Climate change related to egg-laying trends. - Nature 399: 423–
424.
Denéchère, R. et al. 2019. The within-population variability of leaf spring and autumn phenology is
influenced by temperature in temperate deciduous trees. - Int. J. Biometeorol. in press.
Dobrowski, S. Z. 2011. A climatic basis for microrefugia: The influence of terrain on climate. - Glob.

Chang. Biol. 17: 1022–1035.

Dunn, P. O. and Møller, A. P. 2013. Changes in breeding phenology and population size of birds. - J.
Anim. Ecol.: 729–739.

Forsman, J. T. et al. 2008. Competitor density cues for habitat quality facilitating habitat selection
and investment decisions. - Behav. Ecol. 19: 539–545.

Frey, S. J. K. et al. 2016. Microclimate predicts within-season distribution dynamics of montane
forest birds. - Divers. Distrib. 22: 944–959.

Gräler, B. et al. 2016. Spatio-temporal interpolation using gstat. - R J. 8: 204–218.

Grendelmeier, A. et al. 2015. Reproductive performance of a declining forest passerine in relation to
environmental and social factors: Implications for species conservation. - PLoS One 10: 1–23.

Hadfield, J. D. 2010. MCMC methods for multi-response generalized linear mixed models: The
MCMCglmm R package. - J. Stat. Softw. 33: 1–22.

Haest, B. et al. 2020. Weather at the winter and stopover areas determines spring migration onset,
progress, and advancements in Afro-Palearctic migrant birds. - Proc. Natl. Acad. Sci. U. S. A. in
press.

Hinks, A. E. et al. 2015. Scale-dependent phenological synchrony between songbirds and their
caterpillar food source. - Am. Nat. 186: 84–97.

Hušek, J. et al. 2014. Natal dispersal based on past and present environmental phenology in the pied
flycatcher (Ficedula hypoleuca). - Oecologia 174: 1139–49.

Källander, H. et al. 2017. Variation in laying date in relation to spring temperature in three species of
tits (Paridae) and pied flycatchers Ficedula hypoleuca in southernmost Sweden. - J. Avian Biol.
48: 83–90.

Lambrechts, M. M. and Perret, P. 2000. A long photoperiod overrides non-photoperiodic factors in
blue tits' timing of reproduction. - Proc. R. Soc. B Biol. Sci. 267: 585–588.

Lundberg, A. and Alatalo, R. 2010. The Pied Flycatcher. - A & C Black.

Mallord, J. W. et al. 2017. Diet flexibility in a declining long-distance migrant may allow it to escape
the consequences of phenological mismatch with its caterpillar food supply. - Ibis (Lond. 1859).
159: 76–90.

Massimino, D. et al. 2019. BirdTrends 2019: trends in numbers, breeding success and survival for UK
breeding birds.

Maziarz, M. and Wesołowski, T. 2010. Timing of breeding and nestling diet of Wood Warbler
Phylloscopus sibilatrix in relation to changing food supply. - Bird Study 57: 540–552.

McLean, N. et al. 2016. Predicting when climate-driven phenotypic change affects population
dynamics. - Ecol. Lett. 19: 595–608.

Miller-Rushing, A. J. et al. 2010. The effects of phenological mismatches on demography. - Philos.
Trans. R. Soc. Lond. B. Biol. Sci. 365: 3177–86.

Nicolau, P. et al. 2021. Citizen science data reveal highly variable intervals between arrival and
breeding for a long-distance migratory bird. - J. Avian Biol. in press.

Parmesan, C. and Yohe, G. 2003. A globally coherent fingerprint of climate change impacts across
natural systems. - Nature 421: 37–42.

Pasinelli, G. et al. 2016. Rodent-avoidance, topography and forest structure shape territory selection
of a forest bird. - BMC Ecol. 16: 1–18.

Perrins, C. M. 1979. British Tits.

Phillimore, A. B. et al. 2016. Passerines may be sufficiently plastic to track temperature-mediated
shifts in optimum lay date. - Glob. Chang. Biol. 22: 3259–3272.

Potti, J. 1998. Arrival Time from Spring Migration in Male Pied Flycatchers: Individual Consistency
and Familial Resemblance. - Condor 100: 702–708.

Potti, J. and Montalvo, S. 1991. Male Arrival and Female Mate Choice in Pied Flycatchers Ficedula
hypoleuca in Central Spain. - Ornis Scand. 22: 45–54.

Potti, J. et al. 2018. Long-term occupancy of nest boxes as a measure of territory quality for Pied
Flycatchers. - J. F. Ornithol. 89: 337–347.

R Core Team 2018. R Core Team (2018). R: A language and environment for statistical computing. - R
Found. Stat. Comput. Vienna, Austria. URL http//www.R-project.org/.: R Foundation for
Statistical Computing.
Samplonius, J. M. et al. 2018. Phenological sensitivity to climate change is higher in resident than in
migrant bird populations among European cavity breeders. - Glob. Chang. Biol. 24: 3780–3790.
Sanz, J. J. 1998. Effects of Geographic Location and Habitat on Breeding Parameters of Great Tits. -
Auk 115: 1034–1051.
Schaper, S. V et al. 2011. Spring phenology does not affect timing of reproduction in the great tit
(Parus major). - J. Exp. Biol. 214: 3664–71.
Schielzeth, H. 2010. Simple means to improve the interpretability of regression coefficients. -
Methods Ecol. Evol. 1: 103–113.
Schuett, W. et al. 2017. Nosy neighbours: large broods attract more visitors. A field experiment in
the pied flycatcher, Ficedula hypoleuca. - Oecologia 184: 115–126.
Sergio, F. and Newton, I. 2003. Occupancy as a measure of habitat quality. - J. Anim. Ecol. 72: 857–
865.
Shutt, J. D. et al. 2018. The effects of woodland habitat and biogeography on blue tit Cyanistes
caeruleus territory occupancy and productivity along a 220 km transect. - Ecography (Cop.). 41:
1967–1978.
Shutt, J. D. et al. 2019. The environmental predictors of spatiotemporal variation in the breeding
phenology of a passerine bird. - Proc. R. Soc. B Biol. Sci. in press.
Shutt, J. D. et al. 2020. Gradients in richness and turnover of a forest passerine's diet prior to
breeding: A mixed model approach applied to faecal metabarcoding data. - Mol. Ecol.: 1199–
1213.
Simmonds, E. G. et al. 2017. Incubation behavior adjustments, driven by ambient temperature
variation, improve synchrony between hatch dates and caterpillar peak in a wild bird
population. - Ecol. Evol. 7: 9415–9425.
Sirkiä, P. M. and Laaksonen, T. 2009. Distinguishing between male and territory quality: females
choose multiple traits in the pied flycatcher. - Anim. Behav. 78: 1051–1060.
Slagsvold, T. 1976. Annual and Geographical Variation in the Time of Breeding of the Great Tit Parus
major and the Pied Flycatcher Ficedula hypoleuca in Relation to Environmental Phenology and
Spring Temperature. - Ornis Scand. 7: 127–145.
Socolar, J. B. et al. 2017. Phenological shifts conserve thermal niches in North American birds and
reshape expectations for climate-driven range shifts. - Proc. Natl. Acad. Sci. U. S. A. 114:
12976–12981.
van Asch, M. et al. 2012. Evolutionary response of the egg hatching date of a herbivorous insect
under climate change. - Nat. Clim. Chang. 3: 244–248.
Verhagen, I. et al. 2020. Temperature has a causal and plastic effect on timing of breeding in a small
songbird. - J. Exp. Biol.: jeb.218784.
Visser, M. E. et al. 1998. Warmer springs lead to mistimed reproduction in great tits (Parus major). -
Proc. R. Soc. B Biol. Sci. 265: 1867–1870.
Visser, M. E. et al. 2009. Temperature has a causal effect on avian timing of reproduction. - Proc. R.
Soc. B Biol. Sci. 276: 2323–31.
Visser, M. E. et al. 2015. Effects of Spring Temperatures on the Strength of Selection on Timing of
Reproduction in a Long-Distance Migratory Bird. - PLoS Biol. 13: 1–17.
Wesołowski, T. and Maziarz, M. 2009. Changes in breeding phenology and performance of Wood
Warblers phylloscopus sibilatrix in a primeval forest: A thirty-year perspective. - Acta Ornithol.
44: 69–80.
Wesołowski, T. and Rowiński, P. 2014. Do Blue Tits Cyanistes caeruleus synchronize reproduction
with caterpillar peaks in a primeval forest? - Bird Study 61: 231–245.
Wesołowski, T. and Neubauer, G. 2017. Diet of Marsh Tit Poecile palustris Nestlings in a Primeval
Forest in Relation to Food Supply and Age of Young . - Acta Ornithol. 52: 105–118.

 Figures and tables

586
587   **Figure 1** Map of the study area (East Dartmoor National Nature Reserve, Devon, UK), showing its
588   location within Great Britain (inset, red star), the location of nestboxes (black points) and wood
589   warbler nests (white squares), the local names of sections of the reserve, and the relative local
590   temperature differential throughout.

Table 1 The effects of temperature, territory quality and breeding experience on reproductive phenology and outcomes in three forest passerine species. All results are taken from Bayesian GLM's, with slope estimates and credible intervals for each fixed term, with significance asterisks for significant terms (pMCMC ≤0.05* ≤0.01** ≤0.001***). Yearly temperature and territory quality are mean centred, breeding experience relates to males for arrival date but females for all other measures, and the temperature interaction is between yearly temperature and territory temperature.

Fixed Term	Arrival Date	Nest Initiation	Egg Laying Date	Clutch Size	Productivity
			Pied Flycatcher		
Intercept	15.97 (15.18 – 16.77)	25.16 (24.25 – 25.99)	38.59 (37.90 – 39.37)	6.80 (6.70 – 6.90)	4.75 (4.49 – 5.01)
Yearly Temperature	1.22 (-0.08 – 2.54)	-0.69 (-2.09 – 0.70)	-1.01 (-2.22 – 0.23)	-0.11 (-0.28 – 0.05)	-0.09 (-0.53 – 0.36)
Territory Temperature	4.27 (1.99 – 6.51) ***	0.94 (-1.46 – 3.36)	0.37 (-1.63 – 2.37)	-0.18 (-0.47 – 0.08)	-0.93 (-1.65 - -0.21) *
Temperature Interaction	2.84 (-0.60 – 6.23)	4.38 (0.36 – 8.39) *	-0.49 (-4.11 – 2.74)	0.16 (-0.30 – 0.63)	1.19 (0.00 – 2.38) *
Territory Quality	-4.18 (-9.22 – 1.37)	-11.34 (-17.28 - -5.58) ***	-6.57 (-11.45 - -1.68) *	0.28 (-0.36 – 0.96)	1.76 (0.00 – 3.49) *
Breeding Experience	-0.81 (-1.55 - -0.07) *	-0.49 (-1.24 – 0.24)	-0.93 (-1.55 - -0.28) **	0.17 (0.08 – 0.25) ***	0.02 (-0.20 – 0.24)
			Blue Tit		
Intercept			24.45 (23.47 – 25.50)	8.46 (8.23 – 8.69)	3.14 (2.76 – 3.56)
Yearly Temperature			-2.22 (-3.89 - -0.48) *	0.20 (-0.19 – 0.60)	0.78 (0.10 – 1.48) *
Territory Temperature			-3.02 (-5.89 - -0.24) *	-0.57 (-1.21 – 0.06)	-0.77 (-1.93 – 0.28)
Temperature Interaction			0.10 (-4.71 – 5.12)	1.22 (0.17 – 2.33) *	1.46 (-0.47 – 3.35)
Territory Quality			-2.33 (-8.21 – 3.35)	0.34 (-0.95 – 1.65)	1.31 (-0.95 – 3.73)
			Wood Warbler		
Intercept			49.92 (45.12 – 55.08)	5.59 (5.23 – 5.94)	2.99 (2.02 – 3.94)
Yearly Temperature			2.57 (-6.54 – 11.40)	0.35 (-0.28 – 0.97)	0.09 (-1.56 – 1.89)
Territory Temperature			-3.53 (-25.31 – 17.68)	0.11 (-1.33 – 1.40)	-1.68 (-5.41 – 2.10)
Temperature Interaction			51.59 (13.58 – 91.09) *	-2.82 (-5.23 - -0.50) *	-0.44 (-7.06 – 5.86)

Figure 2 Illustrating the significant effects of territory temperature on the breeding parameters of three woodland passerine species: (a-d) pied flycatcher (e-f) blue tit (g-h) wood warbler. Figures a, c and e show significant single predictor linear effects (solid black line) along with 95% credible intervals (grey shading). Figures b, d and f-h show significant interaction linear effects (solid lines) and 95% confidence intervals (shading) in a cold (blue, -0.7°C below mean interannual temperature), average (green, mean interannual temperature) and warm (red, +0.7°C above mean interannual temperature) example year. Bird drawings reproduced with permission from Mike Langman, RSPB Images.

Figure 3 Illustrating the significant effects (solid black lines) of breeding experience (a male and b-c female) and territory quality (d-f) on the breeding parameters of pied flycatcher, along with 95% credible intervals (grey shading).