Territory-level temperature influences settlement decisions, breeding phenology and productivity

in three forest passerine birds

Jack D Shutt[1,2,*], Sophie C Bell[1], Fraser Bell[3], Joan Castello[4], Myriam El Harouchi[1], and Malcolm D

Burgess[1,4,5]

1. Piedfly.Net, Yarner Wood, Bovey Tracey, Devon TQ13 9LN

2. Department of Natural Sciences, Manchester Metropolitan University, Manchester, M1 5GD, UK

3. Centre for Ecology and Conservation, Cornwall Campus, University of Exeter, Penryn, Cornwall,

TR10 9FE, UK

4. Centre for Research in Animal Behaviour, University of Exeter, Perry Road, Exeter, Devon, EX4

4QG

5. RSPB Centre for Conservation Science, The Lodge, Sandy, Bedfordshire, SG19 2DL, UK

* Corresponding author: jackshutt8@gmail.com

Keywords: Local temperature, trophic mismatch, blue tit, pied flycatcher, wood warbler, breeding

experience, territory quality, nestbox

Abstract

1. Temperature is a key determinant of breeding phenology in temperate zone birds, with increasing

spring temperature across years or large geographic gradients known to advance timing of breeding.

It is less well understood how localised, territory-scale temperature variations effect territory

occupation, breeding phenology and productivity, primarily due to a lack of fine-scale temperature

data.

2. We examine the influence of temperature at the territory-scale, and its interaction with mean

annual temperature, on territory selection and breeding outcomes of three co-occurring forest

passerine bird species; two long-distance migrants (pied flycatcher *Ficedula hypoleuca* and wood

warbler *Phylloscopus sibilatrix*) and one resident (blue tit *Cyanistes caeruleus*); all species dependent

on seasonal invertebrate food for reproductive success.

3. Spring temperature was recorded at intersections of a 150m^2 grid encompassing 417 ha of forest

in South-West England 2015-18, with temperature across the study area interpolated from this point

data. Breeding phenology and productivity from 672 nests across the three species was quantified,

and general linear models used to examine the influence of territory temperature on breeding

phenology and productivity.

4. All three species exhibited significant trends in reproductive traits associated with territory-scale

temperature. Pied flycatchers settled in cooler territories first, where they raised more fledglings.

Blue tits laid larger clutches in warmer territories in warm years and always laid earlier at warmer

territories irrespective of annual mean temperature. Contrastingly, pied flycatcher and wood

warbler breeding phenology was earlier at warmer territories in cool years and cooler territories in

warm years, with wood warbler clutch size responding similarly to this interaction. Greater previous

breeding experience and increased territory quality also predicted earlier breeding phenology and

higher productivity for pied flycatchers.

5. Our results show that local-scale spatial variation in temperature is associated with spatial

variation in territory occupancy, breeding phenology and productivity in forest passerines. We

suggest that the two migrant species may be best synchronised with their prey requirements in

cooler years at a local population level while resident blue tits match local phenology at lower

trophic levels across all years, potentially advantageous under warmer predicted climate change

scenarios.

Introduction

Temperature plays a key role in determining the timing of breeding of temperate zone bird species

(Crick & Sparks, 1999; Dunn & Møller, 2013). In warmer years, and at generally warmer locations,

breeding phenology occurs earlier, and under recent climate change average breeding phenology

has advanced over time for many species (Both et al., 2004; McLean, Lawson, Leech, & van de Pol,

2016; Parmesan & Yohe, 2003; Phillimore, Leech, Pearce-Higgins, & Hadfield, 2016). An increase of

1°C has been demonstrated to advance breeding phenology by 3-5 days in a range of birds (McLean

et al., 2016; Phillimore et al., 2016; Socolar, Epanchin, Beissinger, & Tingley, 2017). Functionally, this

trend to breed earlier is to maintain temporal synchrony with key prey resources (Burgess et al.,

2018). Prey synchrony is particularly important for single-brooded insectivorous birds (e.g. tits

Paridae, flycatchers *Ficedulidae*, leaf warblers *Phylloscopidae*) breeding in highly seasonal

environments such as deciduous forests (Burgess et al., 2018; Visser, van Noordwijk, Tinbergen, &

Lessells, 1998). Asynchrony with this prey peak can reduce fledgling number and quality, negatively

impacting fitness, with such asynchrony termed trophic mismatch (Both, Bouwhuis, Lessells, &

Visser, 2006; Miller-Rushing, Høye, Inouye, & Post, 2010).

While the effect of average temperature on breeding phenology across years and geographic

gradients is well established, effects of subtler localised intra-annual temperature variation is less

understood. This is primarily due to a lack of studies collecting fine-scale within-site temperature

data alongside bird breeding data, despite this being the scale relevant to individual birds (Hinks et

al., 2015). It is therefore unknown how territory settlement, breeding phenology and breeding

outcome are influenced by territory-scale temperature. It has been shown that within-nest

temperature can modify breeding phenology (Simmonds, Sheldon, Coulson, & Cole, 2017), and that

fine-scale temperature is a strong predictor of localised species distribution (Frey, Hadley, & Betts,

2016). Phenological timings that may be influenced by territory-scale temperature variation include

the date of arrival, date of nest building initiation and first egg laying date, while breeding

parameters that may be sensitive include clutch size and productivity. In the same breeding

locations resident species with similar ecological requirements to long-distance migrants often

commence breeding earlier (Källander et al., 2017; Samplonius et al., 2018). With a warming spring

climate this difference can become more divergent (Haest, Hüppop, & Bairlein, 2020; Samplonius et

al., 2018) suggesting fine-scale temperature may determine breeding onset differently between

migrants and residents as well as differing patterns of territory occupancy at a local scale.

Understanding the influence of temperature on territory occupation, breeding phenology and

outcome could indicate how optimally timed a population is. As tree and caterpillar phenology is

well synchronised with fine-scale temperature (Buse, Dury, Woodburn, Perrins, & Good, 1999; van

Asch, Salis, Holleman, van Lith, & Visser, 2012), locally warmer territories will show advanced

phenology in lower trophic levels than cool territories (Cole, Long, Zelazowski, Szulkin, & Sheldon,

2015; Hinks et al., 2015). If breeding always commences earlier in warmer territories this could

indicate successful thermal niche tracking, where a well-timed population is able to track the local

phenology of lower trophic levels (Dobrowski, 2011; Socolar et al., 2017). However, if within a

woodland breeding occurred earlier in cooler territories relative to warmer territories with earlier

prey phenology, then the average population consequence would be later than optimal timing as

late breeders (either through late arrival, lower quality or competition) will be forced to occupy the

remaining warmer territories with the greatest levels of mismatch. If breeding commences earlier in

warmer territories in cool years, and in cooler territories in warm years, this would suggest that

average population timing is optimal in cool years but suboptimal (late) in warm years.

Examining a population's breeding phenology across a temperature gradient requires few

assumptions to be made about precise optimal dates or specific prey items compared to studies

focussed on dates. Optimal synchrony between bird species breeding in the same locality may vary

due to different prey requirements and developmental needs of young. The ephemeral invertebrate

peak in deciduous woodland that several single-brooded passerine species are commonly assumed

to depend on are moth caterpillars (Burgess et al., 2018; Visser et al., 1998). However, the most well

studied species of this system (blue tit *Cyanistes caeruleus*, great tit *Parus major* and *Ficedula*

flycatchers) breed at overlapping but different times to one another (Burgess et al., 2018;

Samplonius et al., 2018), and so within any single year and locality breeding timing across these bird

species will encompass peaks of different prey species at different nestling developmental stages.

Timing breeding to match peak demands of nestlings to the caterpillar peak can have little impact on

fledging mass or productivity in some species (Mallord et al., 2017; Maziarz & Wesołowski, 2010;

Wesołowski & Neubauer, 2017), while it does in others (Both, Bijlsma, & Visser, 2005; Visser et al.,

1998), suggesting differences in prey requirements. Whether breeding is timed to the same or

different optima, a preference for breeding in cooler territories first would indicate local population-

level mistiming. It would also suggest that the birds were adjusting territory selection and breeding

phenology due to the local phenological stage (Hinks et al., 2015; Hušek, Lampe, & Slagsvold, 2014;

Shutt et al., 2019), in addition to purely relying on temperature as a causal cue of breeding time

(Caro, Schaper, Hut, Ball, & Visser, 2013; Verhagen, Tomotani, Gienapp, & Visser, 2020).

In this study, we investigate how territory-scale temperature variation over four years within a

417ha woodland affects territory settlement, breeding phenology, clutch size and productivity of

three single-brooded co-occurring insectivorous woodland passerine bird species: pied flycatcher

Ficedula hypoleuca, blue tit and wood warbler *Phylloscopus sibilatrix*. All three are similarly sized,

primarily single-brooded, spring-breeding species. We hypothesise that as these migrants tend to

breed later than residents (Samplonius et al., 2018; Slagsvold, 1976), and later breeding in warm

years increases mismatching (Burgess et al., 2018), that in warm years migratory species will be less

optimally timed compared to resident species, selecting cooler territories first and showing lower

productivity in warm territories.

Materials and methods

This study was conducted at East Dartmoor National Nature Reserve in Devon, South-West England,

UK (50.60°N -3.73°E) over four breeding seasons 2015-2018. Habitat across the reserve is relatively

uniform and dominated by sessile oak *Quercus petraea* with a European holly *Ilex aquifolium,* and

rowan *Sorbus aucuparia* understory and a European bilberry *Vaccinium myrtillus* field layer. The

study area extends linearly approximately 5 km along a steep-sided river valley in the north, with a

150ha woodland containing two shallow stream valleys in the south, together resulting in slopes at

different aspects and an altitudinal range of 60-300m.

Breeding bird data

Throughout the study area, 309 wooden nestboxes with a 32 mm entrance hole were available each

year and monitored annually throughout the bird breeding season April-June. Nestbox checks were

made at least weekly and more frequently at some nest stages to determine species occupancy,

date of first nesting material, date of first egg laying, completed clutch size and the number of

successful fledglings (productivity) for each breeding attempt. Date of first nesting material was

taken as the date of the first nest check that found fresh nesting material for nests that ultimately

had a clutch of eggs laid in, with nests found half complete (cup shape evident) or more over the

maximum 7 days inspection interval assigned the mid-point between the two visits. All species were

assumed to lay one egg per day (Lundberg & Alatalo, 2010; Perrins, 1979), with the first egg date

back-calculated from incomplete clutches recorded at nest inspections. Clutch size records the total

number of eggs laid that were incubated, and the number fledged was the number alive at the last

visit pre-fledging minus the number found dead at the first visit post-fledging.

The date of spring arrival by territorial male pied flycatchers to nestboxes was also recorded from

the first week in April. Territorial activity, such as singing, alarm calling and inspection of nestboxes,

was observed on alternate mornings, and the presence of, colour and position of any rings and

variable plumage characteristics such as forehead patch size were recorded. These were cross-

referenced against males captured at the same nestboxes later in the breeding season. Male arrival

date to a territory was taken as the mid-point between the survey prior to the first observation of a

male and the survey when first observed. This is a common method used for pied flycatchers (Bell,

Harouchi, Hewson, & Burgess, 2017; Potti, Camacho, Canal, & Martinez-Padilla, 2018) and its

accuracy has been verified with geolocator tracking data (Both, Bijlsma, & Ouwehand, 2016). Nearly

all breeding male and female pied flycatchers were captured annually and are marked with a unique

metal numbered ring upon first capture, enabling the frequency of breeding over an individuals'

lifetime to be determined, including for years prior to 2014 using data collected prior to the present

study. Between 2015-2018, 320 pied flycatcher and 315 blue tit nesting attempts were monitored.

Wood warbler nests were also monitored throughout the breeding seasons 2015-2018. Wood

warblers are ground nesting and were located by observing parental behaviour, with the location of

each nest determined by GPS and visited every 3-4 days. First egg-laying dates were determined

from visits during egg laying assuming that one egg is laid per day. A total of 37 wood warbler nests

were monitored during the study period.

Temperature data

Between 2015-2018 96 temperature dataloggers (Thermachron iButton DS1921G) were placed

throughout the study area at each intersection of a 150m grid. Each was attached to a tree 1.5m

above ground level in a waterproof standard sized plastic container with three drilled holes to

provide air circulation (Shutt et al., 2019). Temperature dataloggers recorded temperature hourly to

an accuracy of 0.5°C. Additionally, 22 higher-sensitivity temperature dataloggers (Thermachron

iButton DS1922L) that record temperature to an accuracy of 0.0625°C were attached alongside 22 of

the lower resolution dataloggers on the same trees with the same aspect in a non-random way to

maximise coverage of the entire study area.

We used temperatures recorded between Julian days 97 and 132, which encapsulates the spring

temperature sensitivity window previously estimated for blue tit and pied flycatcher egg laying for

the study area (Samplonius et al., 2018). Data from temperature dataloggers with more than 16

missing hourly data points within the focal period within a year were discarded as incomplete (n=25,

from a total of 333). Additionally, data from temperature dataloggers recording at least one

measurement beyond the extreme credible temperature bounds within the focal period (<-8°C or

>35°C) were discarded for the year in question as unreliable or faulty (n=9). As all faulty dataloggers

were the higher sensitivity model, and data from high sensitivity dataloggers are not directly

comparable with data from low sensitivity dataloggers, the unaffected higher resolution datalogger

data were used to determine the accuracy of their paired low sensitivity datalogger only and were

excluded from all other analyses (n=56). Paired high sensitivity and low sensitivity dataloggers gave

very similar readings (interquartile range <0.2°C despite recording on different sensitivities with a

possible correct error range of 0.25°C), whereas the variation across the entire study area was much

higher (interquartile range >0.6°C in each year), providing confidence that local temperature

differentials across the study area correspond to genuine local temperature variation rather than

datalogger inaccuracy. Temperature means across the focal period were used and a total of 257

datalogger mean readings remained over the four-year period after this quality control (87 in 2015,

54 in 2016, 56 in 2017, and 60 in 2018).

All spatial analysis was conducted within R version 3.6.3 (R Core Team, 2018) using the raster, gstat

and tmap libraries. The data from each remaining low sensitivity temperature datalogger within

each year was then projected to its geographical location and temperature across the study area

interpolated using inverse distance weighting (Gräler, Pebesma, & Heuvelink, 2016) raised to the

power two. The inverse distance squared weighting power was decided upon after investigating the

ability of the data to predict known temperature points in a 'leave one out' validation routine of

powers between one and four (root-mean-squared-error of P1 = 0.467, P2 = 0.447, P3 = 0.453, P4 =

0.461). Temperature differentials for each nestbox and wood warbler nest location were then

extracted from the resultant heatmap raster (Gräler et al., 2016). On inspection, the heatmap

differential rasters for each year were highly consistent in both spatial pattern and variance and so

an interannual differential mean was used which we term the local temperature differential. This

allows determination of whether territories were in a relatively cool or warm area within the overall

study area and afforded greater differential consistency as not every temperature datalogger

position recorded data in all years.

Statistical analysis

General linear models (GLM) were constructed to assess the effects of local temperature differences

on measures of breeding success for this system. Not all breeding parameters were assessed as a

response variable for each of the three bird species due to lack of data. Five breeding parameters

were used: date of male arrival to breeding territory and date of nest building initiation (both pied

flycatcher only), date of first egg laying, clutch size and productivity (number of fledglings) (all three

species). All dates are reported as April days (days from the 1st April, e.g. 25th April = day 25). For

wood warblers, as nests were often found discovered after eggs had been laid, if date of first egg

laying was not recorded but date of hatching was, first egg date was inferred by subtracting the

mean number of days (n=18) between first egg date and hatching date across those nests where

both were recorded, from hatching date where first egg date was not recorded. All analyses were

conducted in R version 3.6.3 (R Core Team, 2018).

Response variables were tested by Bayesian GLMs (Hadfield, 2010) with annual mean temperature,

local temperature differential, territory quality, previous breeding experience and the interaction

between annual and local temperature as fixed predictor variables. Continuous predictor variables

were mean centred for ease of interpretation and model functioning (Schielzeth, 2010). Each model

was run for 100,000 iterations with the first 10,000 discarded as burn-in with a Gaussian error

structure, with both response data and model traces visually inspected to ensure the fit was

acceptable. Year was not included as a random predictor variable as it correlated perfectly with

annual mean temperature for each data point, however it was included in additional, unreported

models for each response to ensure its omission did not significantly influence results.

Territory quality was inferred from the proportion of years each nestbox had been occupied by

nesting pied flycatchers or blue tits out of the total number of years each nestbox was available

since 1980 and until 2018. Long-term nestbox/territory occupancy patterns have previously been

shown to provide a clear correlate of territory quality, with more frequently occupied territories of

higher quality (Potti et al., 2018; Sergio & Newton, 2003). We used the occupancy rate since 1980 as

both nestbox number and bird population size had by this year stabilised after nest box provisioning

started incrementally from 1955. Territory quality could not be included as a predictor variable for

wood warbler as this species does not use nestboxes and therefore territories could not be ascribed

long term occupancy rates. Individual breeding experience was defined as the number of years an

individual had previously been captured breeding in the population. Previous breeding experience

could not be included as a predictor variable for blue tit or wood warbler due to incomplete data on

parental identity. For pied flycatcher date of arrival, male breeding experience was included as a

fixed effect, whereas in all other models female breeding experience was used, as females build the

nests, lay the eggs and provide the majority of parental care in pied flycatchers (Lundberg & Alatalo,

2010). Pairwise correlation between fixed effects was assessed and as all were <0.18 this suggested

no issue with colinearity within the models, with the most significant correlation being that higher

quality territories tended to have cooler local temperature differentials.

Results

Local spring temperature varied by around 2°C from the coldest to warmest areas across the study

area with colder areas at the higher elevations of Yarner Wood in the south and Neadon Cleave in

the north, and warmer areas at lower elevations along the River Bovey in Lustleigh Cleave (Fig 1).

This pattern was spatially consistent across all four years. Variation in mean spring temperature

between years was observed, with 2018 the warmest (mean = 10.7°C) and 2016 the coolest (mean =

9.2°C).

Male pied flycatchers returning from migration in spring occupied the coolest territories first,

irrespective of the relative mean annual temperature, with our model predicting arrival to the

coolest territories on 12th April and warmest territories on 20th April (Table 1, Fig 2a). There was also

an interaction effect of local temperature on the timing of nest building initiation; in cool years nests

were initiated significantly earlier at warmer territories, with this pattern reversed in warm years

with nests initiated earlier at cooler territories (Table 1, Fig 2b). This temperature effect on the

timing of nest building did not translate into an effect on the timing of egg laying or clutch size

(Table 1). Productivity varied by 1.7 chicks per nest attempt across the temperature gradient, with

productivity higher at the coolest territories compared to the warmest territories (5.6:3.9 fledglings

per attempt respectively) (Table 1, Fig 2c). This latter effect was strongest is cool years, while in

warm years productivity was similar irrespective of the local temperature (Table 1, Fig 2d).

For resident blue tits, higher local temperatures predicted a significant advancement of five days in

the timing of egg laying between the coolest and warmest territories (Table 1, Fig 2e). There was

also an interaction effect of local temperature on clutch size, with cooler territories predicted to

have substantially larger clutches in cool years, while in warm years warmer territories were

predicted to have slightly larger clutches (Table 1, Fig 2f). There was no significant effect of local

temperature on blue tit productivity, although the interaction effect was in the same direction as for

clutch size (Table 1). In warm years, blue tits laid eggs earlier (-2.22 days/°C) and had higher

productivity (+0.78 fledglings/°C) (Table 1).

There were significant effects of local temperature interactions on migratory wood warbler breeding

parameters. In cooler years, eggs were predicted to be laid substantially earlier at warmer territories

than cooler territories, whereas in warmer years the opposite was true, with eggs laid earlier at

cooler territories (Table 1, Fig 2g). Wood warbler clutches were also significantly larger at cooler

territories in warmer years, and at warmer territories in cooler years (Table 1, Fig 2h). There was no

significant effect of local temperature on wood warbler productivity (Table 1).

Previous breeding experience also affected pied flycatcher breeding parameters. Males with more

experience arrived earlier (-0.81 days/year experience, Table 1, Fig 3a) and females with more

experience laid larger clutches (+0.17 eggs/year experience, Table 1, Fig 3b) earlier in the year (-0.93

days/year experience, Table 1, Fig 3C). In higher quality territories pied flycatchers also initiated

earlier nest building (21st April: 1st May, Table 1, Fig 3d) and egg laying (6th May: 12th May, Table 1,

Fig 3e) and produced more fledglings (3.9: 5.4 fledglings per attempt, Table 1, Fig 3f), with figures at

the two extremes of territory quality found in our study. Territory quality showed no significant

relationship with blue tit breeding parameters (Table 1).

Discussion

This study demonstrates multiple relationships resulting from territory-level spatial variation in

temperature to differences in territory occupancy, breeding phenology, clutch size and productivity

for three insectivorous woodland birds. The patterns show that within a 417ha woodland both

migratory and resident species either select breeding territories where territory temperature best

matches breeding phenology, with these locations differing between cold and warm years, and/or

show fitness benefits in relation to the temperature of the territory occupied. In cold years, when

vegetation and invertebrate phenology is expected to be latest (Burgess et al., 2018), pied flycatcher

nest building and wood warbler egg laying commenced earlier in warmer territories, with larger

clutch sizes for wood warbler and blue tit. In warm years, when vegetation and invertebrate

phenology is earlier (Burgess et al., 2018), cooler territories were preferred by the migratory

species', being associated with earlier nest building by pied flycatchers, and earlier egg laying and

larger clutch sizes in wood warblers. Territory-scale temperature also influenced pied flycatcher

productivity across years, with 1.7 more fledglings predicted per nest in the coolest territories

compared to the warmest. This pattern was driven by considerably higher productivity in cool

territories in cooler years, suggestive of a negative effect on productivity from trophic mismatch in

relatively warmer territories in cool years.

Temperature impacts on breeding phenology

Male pied flycatchers returning from Africa settled in cooler territories first, with nest building also

beginning first in cooler territories in warm years, with this pattern reversed in cold years. Our

measure of arrival was only for males as these are more conspicuous, but our method of estimating

nest building initiation is frequently used as a proxy for female arrival date (Potti & Montalvo, 1991;

Visser et al., 2015) as pairing and nest building can commence on the day of female arrival (Potti &

Montalvo, 1991). Making that assumption, our results suggest that both male and female pied

flycatchers may partly base territory selection on territory temperature, with the territories selected

first by the earliest arriving birds those most likely to have a better match with prey phenology. In

warmer years migration timings may constrain earlier arrival (Both & Visser, 2001) with the

implication that in warm years later arrivals are constrained to breed in the warmest territories and

be more likely mismatched with prey. In cooler years the local population overall is therefore better

phenologically matched.

Territory and female mate selection is well studied, with males establishing territories first and

females selecting breeding territories held by males based on either habitat quality (Alatalo,

Lundberg, & Glynn, 1986), or male traits (Sirkiä & Laaksonen, 2009). Territory choice can be further

influenced by information obtained from heterospecifics, for example from cues related to

occupancy and density patterns of resident species (Forsman, Hjernquist, Taipale, & Gustafsson,

2008) or inspecting multiple nest sites upon arrival (Schuett, Järvistö, Calhim, Velmala, & Laaksonen,

2017). Our study suggests fine-scale temperature is another key determinant of territory choice in

pied flycatchers.

Egg laying date in our three study species is constrained by photoperiod (Caro, Lambrechts,

Balthazart, & Perret, 2007; Lambrechts & Perret, 2000) and environmental factors (Shutt et al.,

2019), and is strongly related to spring temperature at larger spatial scales (Both & Visser, 2001;

Phillimore et al., 2016; Schaper, Rueda, Sharp, Dawson, & Visser, 2011), with laying starting later at

cooler locations such as higher latitudes and elevations (Burgess et al., 2018; Sanz, 1998), but

examination at local scales is rare. We found spatial variation in laying date with regards to local

temperature for both wood warbler and blue tit. Wood warblers started egg laying substantially

earlier at warmer territories in cooler years and cool territories in warmer years. Previous work on

wood warblers showed a strong relationship between the temperature of the pre-laying period and

laying date at larger spatial scales, with earlier laying associated with higher pre-laying temperatures

(Wesołowski & Maziarz, 2009). We find the same pattern at the territory scale in cool and average

years, providing further support that temperature is a cue to wood warbler laying. However, in

warm years the opposite pattern occurs, which may indicate phenological mismatch within the local

population in warm years as later breeders are occupying the warmest (and thus most

phenologically advanced) territories, possibly due to these being the only ones remaining once

earlier breeders have occupied the better-matched, cooler territories.

Higher territory temperature linearly predicted earlier egg laying in resident blue tits, similar to that

found across longer gradients of space and time but with a slightly shallower slope to other

estimates (McLean et al., 2016; Phillimore et al., 2016). Our results are consistent with temperature

being a direct cue for blue tit breeding phenology (Shutt et al., 2019; Verhagen et al., 2020; Visser,

Holleman, & Caro, 2009) and that this cue applies at a very localised scale. No significant

temperature relationship for lay date was found for pied flycatcher, suggesting temperature may not

be a direct cue to egg laying in this species. Pied flycatcher egg laying date shows less spatial and

inter-annual variation compared with arrival phenology (Nicolau et al., 2021), indicating that in years

with early arrival, individual birds can nest build well in advance of egg laying. Our results on the

effects of territory scale temperature on breeding phenology suggest that warming springs may be

more favourable to resident blue tits than migrant pied flycatchers and wood warblers at this

locality.

Temperature impacts on clutch size

We found spatial variation related to temperature in clutch size for wood warbler and blue tit, but

not for pied flycatcher. Wood warbler clutch size followed a similar pattern to the breeding

phenology of this species with regards to local temperature, being larger at cool territories in warm

years, and at warm territories in cool years, with clutch size therefore larger in territories selected

earlier for breeding. The effect of territory temperature on clutch size may therefore help to explain

why wood warbler clutch size declines seasonally (Grendelmeier, Arlettaz, Gerber, & Pasinelli, 2015).

Although wood warblers do not time breeding to match caterpillar peaks (Maziarz & Wesołowski,

2010; Wesołowski & Maziarz, 2009), if higher clutch sizes are indicative of better phenological

synchrony to their own invertebrate prey peaks these results imply that wood warblers may be late

matched in warm years and search for their optimal territory level temperature locally (Frey et al.,

2016), refined within territories by nest site preferences such as by slope and distance from forest

edge (Pasinelli, Grendelmeier, Gerber, & Arlettaz, 2016). Blue tit clutch size showed the opposite

interaction, with larger clutches in warm territories in warm years and cool territories in cool years.

Blue tits time breeding to match caterpillar phenological peaks (their principle prey) (Shutt et al.,

2020; Wesołowski & Rowiński, 2014), and our results indicate that blue tit breeding phenology might

be early in cool years and better matched birds breed in cooler territories. As blue tit clutch size does

not vary with woodland habitat or geography but by annual variation (Shutt, Bolton, Cabello,

Burgess, & Phillimore, 2018), annual temperature could be an important factor determining the

quality of the breeding year for blue tits.

Temperature impacts on productivity

There were no territory-level temperature effects on productivity in either blue tit or wood warbler, but pied flycatchers showed a strong response, fledging 1.7 more chicks in cool compared to warm territories, with this effect strongest in cool years. Higher productivity from cooler territories suggests cooler territories have a better match with invertebrate prey abundance in this species. Cooler territories conferring greater productivity may also explain why these territories are occupied first by both male and female pied flycatchers upon arrival. There may be a greater risk of mistiming breeding in warmer springs due to lower levels of local variation in tree budburst (Denéchère et al., 2019).

Non-temperature impacts: territory quality and individual experience

For pied flycatcher our models also controlled for other factors that may influence nest building: clutch size and productivity. Higher territory quality predicted earlier nest initiation and egg laying, and higher productivity, supporting the general validity of using the simple indirect proxy of long-term occupation trends to indicate territory quality (Potti et al., 2018; Sergio & Newton, 2003), particularly for measuring territory quality in relatively homogeneous habitats when habitat composition and structural preferences can be hard to determine or differentiate. The spatial pattern of temperature across the study area was consistent, and the territories that were relatively cool in 2015-2018 had the highest pied flycatcher occupancy rates since 1980, showing that cool territories have been preferentially selected for almost 40 years. The models also indicated that adult flycatchers with greater previous breeding experience (i.e. older individuals) tended to arrive earlier in spring (male experience), lay eggs earlier and lay larger clutches (both female experience).

This is concordant with other studies (Both et al., 2016; Potti, 1998), and is indicative of more

experienced parents having higher nesting success.

Conclusions

Territory-level temperature variations within a single site have profound effects on the breeding

decisions and outcomes of all three passerine bird species studied. Resident blue tit breeding

phenology responds linearly to increasing territory temperatures similar to responses across large-

scale geographic and annual temperature gradients (Phillimore et al., 2016). Migrant pied flycatchers

and wood warblers, on the other hand, may be responding to small-scale phenological cues at lower

trophic levels, and local populations may be late timed in warm years. Current climate change

scenarios potentially favour resident blue tits that track local phenology well across all years, over

the two migrant species which may be best synchronised in cool years. As avian distributions are

limited by thermal breeding niches (Socolar et al., 2017), this may contribute to shaping breeding

distribution and abundance of these species, with southern populations possibly impacted to a

greater extent by climate change (Both et al., 2006). For populations of declining long-distance

migrant species, including pied flycatcher and wood warbler in the UK (Massimino et al., 2019),

knowledge of territory and thermal preferences and their influence on fecundity and productivity

could be used to help maintain distribution and population size by informing woodland management

and the locations of conservation interventions.

Acknowledgements

We thank Ed Shephard, Amy Vanstone, Mary Wallis, Matt Parkins and Emma Inzani for further

assistance in the field, Steve Cramer for assistance managing the temperature data, and David

Rickwood (Woodland Trust) and Simon Lee (Natural England) for logistical help. This project was

funded by Devon Birds, Kevin Cox and through a Heritage Lottery Fund grant. Wood warbler

fieldwork was undertaken as part of a RSPB project joint funded by Natural England within the

Action for Birds in England partnership.

Author contributions

JDS, SCB and MDB conceived the study; SCB, FB, JC, MEH and MDB collected the field data; JDS

curated and analysed the data; JDS and MDB wrote the first draft of the manuscript, with SCB and FB

also contributing critically to later drafts; MDB managed the project.

Data availability statement

Data will be made publicly available in the Dryad Digital Repository upon manuscript acceptance.

References

Alatalo, R. V., Lundberg, A., & Glynn, C. (1986). Female pied flycatchers choose territory quality and
not male characteristics. *Nature*, *323*, 152–153. doi: 10.1038/323152a0

Bell, S. C., Harouchi, M. E. L., Hewson, C. M., & Burgess, M. D. (2017). No short- or long-term effects
of geolocator attachment detected in Pied Flycatchers Ficedula hypoleuca. *Ibis*. doi:
10.1111/ibi.12493

Both, C., Artemyev, A. V, Blaauw, B., Cowie, R. J., Dekhuijzen, A. J., Eeva, T., … Visser, M. E. (2004).
Large-scale geographical variation confirms that climate change causes birds to lay earlier.
Proceedings of the Royal Society B: Biological Sciences, *271*(1549), 1657–1662. doi:
10.1098/rspb.2004.2770

Both, C., Bijlsma, R. G., & Ouwehand, J. (2016). Repeatability in Spring Arrival Dates in Pied
Flycatchers Varies Among Years and Sexes. *Ardea*, *104*(1), 3–21. doi: 10.5253/arde.v104i1.a1

Both, C., Bijlsma, R. G., & Visser, M. E. (2005). Climatic effects on timing of spring migration and
breeding in a long-distance migrant , the pied flycatcher Ficedula hypoleuca. *Journal of Avian*
Biology, *36*, 368–373.

Both, C., Bouwhuis, S., Lessells, C. M., & Visser, M. E. (2006). Climate change and population declines
in a long-distance migratory bird. *Nature*, *441*(7089), 81–83. doi: 10.1038/nature04539

Both, C., & Visser, M. E. (2001). Adjustment to climate change is constrained by arrival date in a long-
distance migrant bird. *Nature*, *411*(6835), 296–298. doi: 10.1038/35077063

Burgess, M. D., Smith, K. W., Evans, K. L., Leech, D., Pearce-Higgins, J. W., Branston, C. J., …
Phillimore, A. B. (2018). Tritrophic phenological match–mismatch in space and time. *Nature*
Ecology & Evolution. doi: 10.1038/s41559-018-0543-1

Buse, A., Dury, S. J., Woodburn, R. J. W., Perrins, C. M., & Good, J. E. G. (1999). Effects of elevated

temperature on multi-species interactions: the case of Pedunculate Oak, Winter Moth and Tits. *Functional Ecology*, *13*(SUPPL. 1), 74–82. doi: 10.1046/j.1365-2435.1999.00010.x

Caro, S. P., Lambrechts, M. M., Balthazart, J., & Perret, P. (2007). Non-photoperiodic factors and timing of breeding in blue tits: Impact of environmental and social influences in semi-natural conditions. *Behavioural Processes*, *75*(1), 1–7. doi: 10.1016/j.beproc.2007.02.011

Caro, S. P., Schaper, S. V, Hut, R. a, Ball, G. F., & Visser, M. E. (2013). The case of the missing mechanism: how does temperature influence seasonal timing in endotherms? *PLoS Biology*, *11*(4), e1001517. doi: 10.1371/journal.pbio.1001517

Cole, E. F., Long, P. R., Zelazowski, P., Szulkin, M., & Sheldon, B. C. (2015). Predicting bird phenology from space: Satellite-derived vegetation green-up signal uncovers spatial variation in phenological synchrony between birds and their environment. *Ecology and Evolution*, *5*, 5057–5074. doi: 10.1002/ece3.1745

Crick, H. Q. P., & Sparks, T. H. (1999). Climate change related to egg-laying trends. *Nature*, *399*, 423–424.

Denéchère, R., Delpierre, N., Apostol, E. N., Berveiller, D., Bonne, F., Cole, E. F., … Vincent, G. (2019). The within-population variability of leaf spring and autumn phenology is influenced by temperature in temperate deciduous trees. *International Journal of Biometeorology*.

Dobrowski, S. Z. (2011). A climatic basis for microrefugia: The influence of terrain on climate. *Global Change Biology*, *17*(2), 1022–1035. doi: 10.1111/j.1365-2486.2010.02263.x

Dunn, P. O., & Møller, A. P. (2013). Changes in breeding phenology and population size of birds. *The Journal of Animal Ecology*, 729–739. doi: 10.1111/1365-2656.12162

Forsman, J. T., Hjernquist, M. B., Taipale, J., & Gustafsson, L. (2008). Competitor density cues for habitat quality facilitating habitat selection and investment decisions. *Behavioral Ecology*, *19*(3), 539–545. doi: 10.1093/beheco/arn005

Frey, S. J. K., Hadley, A. S., & Betts, M. G. (2016). Microclimate predicts within-season distribution dynamics of montane forest birds. *Diversity and Distributions*, *22*(9), 944–959. doi: 10.1111/ddi.12456

Gräler, B., Pebesma, E., & Heuvelink, G. (2016). Spatio-temporal interpolation using gstat. *R Journal*, *8*, 204–218. doi: 10.32614/rj-2016-014

Grendelmeier, A., Arlettaz, R., Gerber, M., & Pasinelli, G. (2015). Reproductive performance of a declining forest passerine in relation to environmental and social factors: Implications for species conservation. *PLoS ONE*, *10*(7), 1–23. doi: 10.1371/journal.pone.0130954

Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: The MCMCglmm R package. *Journal of Statistical Software*, *33*(2), 1–22. doi: 10.1002/ana.22635

Haest, B., Hüppop, O., & Bairlein, F. (2020). Weather at the winter and stopover areas determines spring migration onset, progress, and advancements in Afro-Palearctic migrant birds. *Proceedings of the National Academy of Sciences of the United States of America*. doi: 10.1073/pnas.1920448117

Hinks, A. E., Cole, E. F., Daniels, K. J., Wilkin, T. A., Nakagawa, S., & Sheldon, B. C. (2015). Scale-dependent phenological synchrony between songbirds and their caterpillar food source. *The American Naturalist*, *186*(1), 84–97. doi: 10.1086/681572

Hušek, J., Lampe, H. M., & Slagsvold, T. (2014). Natal dispersal based on past and present environmental phenology in the pied flycatcher (Ficedula hypoleuca). *Oecologia*, *174*(4), 1139–1149. doi: 10.1007/s00442-013-2842-1

Källander, H., Hasselquist, D., Hedenström, A., Nord, A., Smith, H. G., & Nilsson, J.-Å. (2017). Variation in laying date in relation to spring temperature in three species of tits (Paridae) and pied flycatchers Ficedula hypoleuca in southernmost Sweden. *Journal of Avian Biology*, *48*, 83–90. doi: 10.1111/jav.01287

Lambrechts, M. M., & Perret, P. (2000). A long photoperiod overrides non-photoperiodic factors in blue tits' timing of reproduction. *Proceedings of the Royal Society B: Biological Sciences*, *267*(1443), 585–588. doi: 10.1098/rspb.2000.1041

Lundberg, A., & Alatalo, R. (2010). *The Pied Flycatcher*. A & C Black.

Mallord, J. W., Orsman, C. J., Cristinacce, A., Stowe, T. J., Charman, E. C., & Gregory, R. D. (2017). Diet
flexibility in a declining long-distance migrant may allow it to escape the consequences of
phenological mismatch with its caterpillar food supply. *Ibis*, *159*(1), 76–90. doi:
10.1111/ibi.12437

Massimino, D., Woodward, I., Hammond, M., Harris, S., Leech, D. I., Noble, D., … Robinson, R. A.
(2019). *BirdTrends 2019: trends in numbers, breeding success and survival for UK breeding
birds*. Thetford.

Maziarz, M., & Wesołowski, T. (2010). Timing of breeding and nestling diet of Wood Warbler
Phylloscopus sibilatrix in relation to changing food supply. *Bird Study*, *57*(4), 540–552. doi:
10.1080/00063657.2010.512954

McLean, N., Lawson, C. R., Leech, D. I., & van de Pol, M. (2016). Predicting when climate-driven
phenotypic change affects population dynamics. *Ecology Letters*, *19*(6), 595–608. doi:
10.1111/ele.12599

Miller-Rushing, A. J., Høye, T. T., Inouye, D. W., & Post, E. (2010). The effects of phenological
mismatches on demography. *Philosophical Transactions of the Royal Society of London. Series
B, Biological Sciences*, *365*(1555), 3177–3186. doi: 10.1098/rstb.2010.0148

Nicolau, P., Burgess, M. D., Leech, D. I., Marques, T., Baillie, S. R., Moran, N., & Johnson, A. (2021).
Citizen science data reveal highly variable intervals between arrival and breeding for a long-
distance migratory bird. *Journal of Avian Biology*, (In press).

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across
natural systems. *Nature*, *421*(6918), 37–42. doi: 10.1038/nature01286

Pasinelli, G., Grendelmeier, A., Gerber, M., & Arlettaz, R. (2016). Rodent-avoidance, topography and
forest structure shape territory selection of a forest bird. *BMC Ecology*, *16*(1), 1–18. doi:
10.1186/S12898-016-0078-8

Perrins, C. M. (1979). British Tits. In *William Collins Sons & Co Ltd.*

Phillimore, A. B., Leech, D. I., Pearce-Higgins, J. W., & Hadfield, J. D. (2016). Passerines may be
sufficiently plastic to track temperature-mediated shifts in optimum lay date. *Global Change
Biology*, *22*(10), 3259–3272. doi: 10.1111/gcb.13302

Potti, J. (1998). Arrival Time from Spring Migration in Male Pied Flycatchers: Individual Consistency
and Familial Resemblance. *The Condor*, *100*(4), 702–708. doi: 10.2307/1369752

Potti, J., Camacho, C., Canal, D., & Martinez-Padilla, J. (2018). Long-term occupancy of nest boxes as
a measure of territory quality for Pied Flycatchers. *Journal of Field Ornithology*, *89*(4), 337–347.
doi: 10.1111/jofo.12266

Potti, J., & Montalvo, S. (1991). Male Arrival and Female Mate Choice in Pied Flycatchers Ficedula
hypoleuca in Central Spain. *Ornis Scandinavica*, *22*(1), 45–54.

R Core Team. (2018). R Core Team (2018). R: A language and environment for statistical computing.
R Foundation for Statistical Computing, Vienna, Austria. URL Http://Www.R-Project.Org/., R
Foundation for Statistical Computing.

Samplonius, J. M., Bartošová, L., Burgess, M. D., Bushuev, A. V., Eeva, T., Ivankina, E. V., … Both, C.
(2018). Phenological sensitivity to climate change is higher in resident than in migrant bird
populations among European cavity breeders. *Global Change Biology*, *24*(8), 3780–3790. doi:
10.1111/gcb.14160

Sanz, J. J. (1998). Effects of Geographic Location and Habitat on Breeding Parameters of Great Tits.
The Auk, *115*(4), 1034–1051.

Schaper, S. V, Rueda, C., Sharp, P. J., Dawson, A., & Visser, M. E. (2011). Spring phenology does not
affect timing of reproduction in the great tit (Parus major). *The Journal of Experimental Biology*,
214(Pt 21), 3664–3671. doi: 10.1242/jeb.059543

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients.
Methods in Ecology and Evolution, *1*, 103–113. doi: 10.1111/j.2041-210X.2010.00012.x

Schuett, W., Järvistö, P. E., Calhim, S., Velmala, W., & Laaksonen, T. (2017). Nosy neighbours: large

broods attract more visitors. A field experiment in the pied flycatcher, Ficedula hypoleuca. *Oecologia*, *184*(1), 115–126. doi: 10.1007/s00442-017-3849-9

Sergio, F., & Newton, I. (2003). Occupancy as a measure of habitat quality. *Journal of Animal Ecology*, *72*, 857–865. doi: 10.1046/j.1365-2656.2003.00758.x

Shutt, J. D., Bolton, M., Cabello, I. B., Burgess, M. D., & Phillimore, A. B. (2018). The effects of woodland habitat and biogeography on blue tit Cyanistes caeruleus territory occupancy and productivity along a 220 km transect. *Ecography*, *41*(12), 1967–1978. doi: 10.1111/ecog.03573

Shutt, J. D., Cabello, I. B., Keogan, K., Leech, D. I., Samplonius, J. M., Whittle, L., … Phillimore, A. B. (2019). The environmental predictors of spatiotemporal variation in the breeding phenology of a passerine bird. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1908). doi: 10.1098/rspb.2019.0952

Shutt, J. D., Nicholls, J. A., Trivedi, U. H., Burgess, M. D., Stone, G. N., Hadfield, J. D., & Phillimore, A. B. (2020). Gradients in richness and turnover of a forest passerine's diet prior to breeding: A mixed model approach applied to faecal metabarcoding data. *Molecular Ecology*, (October 2019), 1199–1213. doi: 10.1111/mec.15394

Simmonds, E. G., Sheldon, B. C., Coulson, T., & Cole, E. F. (2017). Incubation behavior adjustments, driven by ambient temperature variation, improve synchrony between hatch dates and caterpillar peak in a wild bird population. *Ecology and Evolution*, *7*(July), 9415–9425. doi: 10.1002/ece3.3446

Sirkiä, P. M., & Laaksonen, T. (2009). Distinguishing between male and territory quality: females choose multiple traits in the pied flycatcher. *Animal Behaviour*, *78*(5), 1051–1060. doi: 10.1016/j.anbehav.2009.06.022

Slagsvold, T. (1976). Annual and Geographical Variation in the Time of Breeding of the Great Tit Parus major and the Pied Flycatcher Ficedula hypoleuca in Relation to Environmental Phenology and Spring Temperature. *Ornis Scandinavica*, *7*(2), 127–145.

Socolar, J. B., Epanchin, P. N., Beissinger, S. R., & Tingley, M. W. (2017). Phenological shifts conserve thermal niches in North American birds and reshape expectations for climate-driven range shifts. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(49), 12976–12981. doi: 10.1073/pnas.1705897114

van Asch, M., Salis, L., Holleman, L. J. M., van Lith, B., & Visser, M. E. (2012). Evolutionary response of the egg hatching date of a herbivorous insect under climate change. *Nature Climate Change*, *3*(10), 244–248. doi: 10.1038/nclimate1717

Verhagen, I., Tomotani, B. M., Gienapp, P., & Visser, M. E. (2020). Temperature has a causal and plastic effect on timing of breeding in a small songbird. *The Journal of Experimental Biology*, (March), jeb.218784. doi: 10.1242/jeb.218784

Visser, M. E., Gienapp, P., Husby, A., Morrisey, M., de la Hera, I., Pulido, F., & Both, C. (2015). Effects of Spring Temperatures on the Strength of Selection on Timing of Reproduction in a Long-Distance Migratory Bird. *PLoS Biology*, *13*(4), 1–17. doi: 10.1371/journal.pbio.1002120

Visser, M. E., Holleman, L. J. M., & Caro, S. P. (2009). Temperature has a causal effect on avian timing of reproduction. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1665), 2323–2331. doi: 10.1098/rspb.2009.0213

Visser, M. E., van Noordwijk, A. J., Tinbergen, J. M., & Lessells, C. M. (1998). Warmer springs lead to mistimed reproduction in great tits (Parus major). *Proceedings of the Royal Society B: Biological Sciences*, *265*(1408), 1867–1870. doi: 10.1098/rspb.1998.0514

Wesołowski, T., & Maziarz, M. (2009). Changes in breeding phenology and performance of Wood Warblers phylloscopus sibilatrix in a primeval forest: A thirty-year perspective. *Acta Ornithologica*, *44*(1), 69–80. doi: 10.3161/000164509X464902

Wesołowski, T., & Neubauer, G. (2017). Diet of Marsh Tit Poecile palustris Nestlings in a Primeval Forest in Relation to Food Supply and Age of Young . *Acta Ornithologica*, *52*(1), 105–118. doi: 10.3161/00016454ao2017.52.1.010

Wesołowski, T., & Rowiński, P. (2014). Do Blue Tits Cyanistes caeruleus synchronize reproduction

648   with caterpillar peaks in a primeval forest? *Bird Study*, *61*(2), 231–245. doi:

649   10.1080/00063657.2014.899307

 Figures and tables

654
655 **Figure 1** Map of the study area (East Dartmoor National Nature Reserve, Devon, UK), showing its
656 location within Great Britain (inset, red star), the location of nestboxes (black points) and wood
657 warbler nests (white squares), the local names of sections of the reserve, and the relative local
658 temperature differential throughout.

Table 1 The effects of temperature, territory quality and breeding experience on reproductive phenology and outcomes in three forest passerine species. All results are taken from Bayesian GLM's, with slope estimates and credible intervals for each fixed term, with significance asterisks for significant terms (pMCMC ≤0.05* ≤0.01** ≤0.001***). Yearly temperature and territory quality are mean centred, breeding experience relates to males for arrival date but females for all other measures, and the temperature interaction is between yearly temperature and territory temperature.

Fixed Term	Arrival Date	Nest Initiation	Egg Laying Date	Clutch Size	Productivity
Pied Flycatcher					
Intercept	15.97 (15.18 – 16.77)	25.16 (24.25 – 25.99)	38.59 (37.90 – 39.37)	6.80 (6.70 – 6.90)	4.75 (4.49 – 5.01)
Yearly Temperature	1.22 (-0.08 – 2.54)	-0.69 (-2.09 – 0.70)	-1.01 (-2.22 – 0.23)	-0.11 (-0.28 – 0.05)	-0.09 (-0.53 – 0.36)
Territory Temperature	4.27 (1.99 – 6.51) ***	0.94 (-1.46 – 3.36)	0.37 (-1.63 – 2.37)	-0.18 (-0.47 – 0.08)	-0.93 (-1.65 - -0.21) *
Temperature Interaction	2.84 (-0.60 – 6.23)	4.38 (0.36 – 8.39) *	-0.49 (-4.11 – 2.74)	0.16 (-0.30 – 0.63)	1.19 (0.00 – 2.38) *
Territory Quality	-4.18 (-9.22 – 1.37)	-11.34 (-17.28 - -5.58) ***	-6.57 (-11.45 - -1.68) *	0.28 (-0.36 – 0.96)	1.76 (0.00 – 3.49) *
Breeding Experience	-0.81 (-1.55 - -0.07) *	-0.49 (-1.24 – 0.24)	-0.93 (-1.55 - -0.28) **	0.17 (0.08 – 0.25) ***	0.02 (-0.20 – 0.24)
Blue Tit					
Intercept			24.45 (23.47 – 25.50)	8.46 (8.23 – 8.69)	3.14 (2.76 – 3.56)
Yearly Temperature			-2.22 (-3.89 - -0.48) *	0.20 (-0.19 – 0.60)	0.78 (0.10 – 1.48) *
Territory Temperature			-3.02 (-5.89 - -0.24) *	-0.57 (-1.21 – 0.06)	-0.77 (-1.93 – 0.28)
Temperature Interaction			0.10 (-4.71 – 5.12)	1.22 (0.17 – 2.33) *	1.46 (-0.47 – 3.35)
Territory Quality			-2.33 (-8.21 – 3.35)	0.34 (-0.95 – 1.65)	1.31 (-0.95 – 3.73)
Wood Warbler					
Intercept			49.92 (45.12 – 55.08)	5.59 (5.23 – 5.94)	2.99 (2.02 – 3.94)
Yearly Temperature			2.57 (-6.54 – 11.40)	0.35 (-0.28 – 0.97)	0.09 (-1.56 – 1.89)
Territory Temperature			-3.53 (-25.31 – 17.68)	0.11 (-1.33 – 1.40)	-1.68 (-5.41 – 2.10)
Temperature Interaction			51.59 (13.58 – 91.09) *	-2.82 (-5.23 - -0.50) *	-0.44 (-7.06 – 5.86)

Figure 2 Illustrating the significant effects of territory temperature on the breeding parameters of three woodland passerine species: (a-d) pied flycatcher (e-f) blue tit (g-h) wood warbler. Figures a, c and e show significant single predictor linear effects (solid black line) along with 95% credible intervals (grey shading). Figures b, d and f-h show significant interaction linear effects (solid lines) and 95% confidence intervals (shading) in a cold (blue, -0.7°C below mean interannual temperature), average (green, mean interannual temperature) and warm (red, +0.7°C above mean interannual temperature) example year. Bird drawings reproduced with permission from Mike Langman, RSPB Images.

Figure 3 Illustrating the significant effects (solid black lines) of breeding experience (a male and b-c female) and territory quality (d-f) on the breeding parameters of pied flycatcher, along with 95% credible intervals (grey shading).