<u>**Faecal metabarcoding reveals pervasive long-distance impacts of garden bird feeding**</u>

Jack D Shutt [1,2,*], Urmi H Trivedi [3], James A Nicholls [2,4]

1. Department of Natural Sciences, Manchester Metropolitan University, Manchester, M1 5GD, UK

2. Institute of Evolutionary Biology, The King's Buildings, University of Edinburgh, Edinburgh, EH9

3FL, UK

3. Edinburgh Genomics, University of Edinburgh, Edinburgh, EH9 3FL, UK

4. Australian National Insect Collection, CSIRO, Acton, ACT, 2601, Australia

* Corresponding author: <u>jackshutt8@gmail.com</u>

ORCIDs: JDS 0000-0002-4146-8748, JAN 0000-0002-9325-563X

Keywords: human-wildlife interaction, diet, phenology, population change, supplementary

provisioning, breeding density

Abstract

Supplementary feeding of wildlife is widespread, being undertaken by more than half of households

in many countries. However, the impact that these supplemental resources have is unclear, with

impacts assumed to be restricted to urban ecosystems. We reveal the pervasiveness of

supplementary foodstuffs in the diet of a wild bird using metabarcoding of blue tit (*Cyanistes*

caeruleus) faeces collected in early spring from a 220km transect in Scotland with a large

urbanisation gradient. Supplementary foodstuffs were present in the majority of samples, with

peanut (*Arachis hypogaea*) the single commonest (either natural or supplementary) dietary item.

Consumption rates exhibited a distance decay from human habitation but remained high at several

hundred metres from the nearest household and continued to our study limit of 1.4km distant.

Supplementary food consumption was associated with a near quadrupling of blue tit breeding

density and a five-day advancement of breeding phenology. We show that woodland bird species

using supplementary food have increasing UK population trends, while species that don't, and/or are

outcompeted by blue tits, are likely to be declining. We suggest that the impacts of supplementary

feeding are larger and more spatially extensive than currently appreciated and could be disrupting

population and ecosystem dynamics.

Introduction

Supplementary feeding of garden wildlife is the most common active form of human-wildlife

interaction and occurs globally [1,2]. It is particularly widespread in the Western world, with over

half of all households participating in many Northern European and North American countries,

providing an ever-increasing variety of foodstuffs and feeder designs targeting more diverse species

each year [3,4]. Many mammal and insect species are intentionally provided with supplementary

food, but bird feeding is the commonest activity [2,5]. In the UK, for example, the wild birdfood

market is estimated to be worth £241 million and supply around 150,000 tonnes of supplementary

food annually [6] while in the USA over 500,000 tonnes are supplied annually [2,7]. In the UK there is

estimated to be one supplementary bird feeder per 9 feeder-using birds [5], providing enough

resources nationally to feed three times the entire breeding populations of the ten commonest

feeder-using species year-round if they consumed nothing else [8]. Many mammal species such as

squirrels and rats also use these resources incidentally but at high frequencies [2,9,10]. While garden

wildlife feeding is actively and enthusiastically encouraged by conservation organisations in a

majority of countries, including the UK and USA [11,12], such an enormous resource addition is likely

to have profound effects on both the organisms benefitting from it and their natural competitors

and prey, and these effects are far from well understood [1,13,14].

To date, research into the direct effects of supplementary garden wildlife feeding on the species

utilising it has developed a rather contradictory and mixed evidence base. While some studies have

found that supplementary feeding advances breeding phenology and improves reproductive success

due to increased resources [15–17], others have found the opposite, possibly due to poor nutrition

[18,19]. Similarly, some studies have found benefits to individual user health [20] while others have

found detrimental effects [21]. Population- or species-level health is also at risk as promoting

artificial long-term aggregations of novel individual and species interactions has facilitated disease

spread and crossover, causing large declines in some susceptible species [22,23]. There is consensus

as to overwinter survival benefits, to such an extent that migration patterns can be altered due to

novel year-round resources [24,25]. Such a large-scale change in diet and feeding behaviour is also

likely to have further effects that are just being realised, such as changes to blood chemistry [26] and

evolutionary traits [27]. Elucidating any effects on the breeding ecology of feeder-using species is

particularly important due to the immediate fitness and population impacts.

One reason why the evidence is conflicting may lie in many studies not being able to account

accurately for supplementary food uptake rates in their study organisms due to difficulties in diet

detection, and without this critical information it is impossible to assess large-scale impacts and

background consumption rates [1,13]. The advent of faecal metabarcoding provides a mechanism

whereby this can in part be overcome [28]. This method detects fragments of prey DNA contained

within faeces non-invasively, and while the technique is in its infancy and primarily applied to insect

prey DNA [29,30], many food types can be distinguished, including plant DNA [31] which are

traditionally the commonest supplementary foods provisioned for garden wildlife [1].

Most studies to date have provided additional experimental supplementary food and assumed a

distance decay in uptake [18,19], however this has two major caveats. Firstly, it does not account for

background supplementary feeding rates from resources provided by the local human population

unconnected to the study itself, with such cross-contamination rates probably high due to the

ubiquity of supplementary provisioning [1,5]. Secondly, the distance decay rate is unknown in most

species and therefore supplementary feeding may be impacting over a greater spatial scale than

imagined [17,32]. Assessing diet composition directly through faecal metabarcoding without

providing additional experimental resources addresses these caveats.

The largely unknown distance over which the impacts of supplementary feeding occur is also

evidenced by the majority of research explicitly assuming that impacts are only, or overwhelmingly,

encountered in the urban environment, even altering community structure there [33–35], without

considering the wider rural environment. This ignores the ability of many provisioned bird, mammal

and insect species to move long distances in search of reliable feeding opportunities [32], and that

rural human dwellings are likely to provide more supplementary food per household than urban

dwellings [4]. Additionally, research has focussed solely on the species that utilise supplementary

feeding without considering those that do not. It is likely that increases in populations of

supplementary food-using species [33] and individual competitiveness will have a negative effect on

their competitor and prey species that do not benefit from supplementary feeding, as background

habitat availability is unchanged. Furthermore, if the effects of supplementary feeding are felt over a

wider area than solely urban environments, the impacts on community composition and

conservation could be far-reaching [16]. Therefore, it is crucial to understand over what distances

feeder-using species are travelling to make use of supplementary food resources and what impact

this is having upon their ecology, fitness and populations.

In this study, we analyse data from a widespread and common European avian supplementary food

user, the blue tit (*Cyanistes caeruleus*), across a 220km transect of Scotland [36] with a large

gradient in distance to human habitation and therefore access to supplementary feeding. We use

faecal metabarcoding to uncover what proportion of faeces contain supplementary garden bird food

immediately prior to breeding and over what distance supplementary food use is occurring,

predicting that use will decline with increasing distance. We then use site average supplementary

food use to determine effects on breeding ecology. Finally, we use long-term UK-wide survey data to

address the broader implications by assessing whether the utilisation of supplementary food is

affecting recent population trends in blue tits and their competitors (insectivorous forest bird

species) across the UK, hypothesising that if supplementary feeding is supporting higher populations

of those species using it, these inflated populations may be having detrimental effects on the

populations of competitor species that do not, contributing to human-mediated homogenising

impacts on biodiversity [37,38]. We believe that this focal study system is highly representative of

many supplementary feeding systems and that insights garnered should extrapolate across many

systems.

Methods

Field data collection

Field data were collected from a transect of 39 predominantly deciduous Scottish woodlands during the springs of 2014-2016 [36]. At each site there were six Schwegler 1B 26 mm hole nestboxes distributed at approximately 40 m intervals. From mid-March in both 2014 and 2015 the base of each nestbox was lined with greaseproof paper which was replaced when damaged or heavily soiled and removed at the onset of nest building or once a bird had attempted removal by pulling it through the hole [29]. Each nestbox was visited on alternate days and all faeces on the greaseproof paper were removed with sterilised tweezers, with up to a maximum of three faeces collected into a 2 ml Eppendorf tube containing pure ethanol, and the number of faeces collected recorded (with the exception of samples in early 2014). Faecal samples were stored at −18°C within a day of collection and transferred to a −20°C freezer at the end of each field season. Samples were collected from 19 March in 2014 and 18 March in 2015 until nest building, giving a median sampling range of 20 days per site in 2014 and 24 days in 2015, and a maximum sampling range at a site of 34 days. Faecal samples were not collected in 2016.

The date of first egg laying was recorded for each nestbox (taken as the previous day if two eggs found, as blue tits lay one egg daily [39]) and nestboxes were designated as occupied in a particular year if at least one egg was laid in a nest by a blue tit. Clutch size was counted once all eggs were laid and incubation had begun. All nestlings were fitted with a metal identification ring under license and productivity was defined as the number of nestlings successfully fledged (number of nestlings alive at day 12 after hatching minus number of dead nestlings found in nestbox post-fledging). Parent birds of both sexes were also captured and fitted with metal identification rings under license and their mass, sex and age (first year breeder or second year plus) recorded. Latitude (site range 55.98 - 57.88°N) and elevation (10 - 433 m) were obtained for each nestbox [36] and the Euclidian distance to nearest human habitation (33 – 1384m) was calculated for each nestbox after finding the

coordinates of the nearest human dwelling via Google maps [40]. Due to the high incidence of

supplementary bird feeding in the UK [2,4] this should be a good predictor of feeder availability,

although we note it provides just the lower limit to the potential distance moved so is a conservative

estimate.

Molecular protocol and bioinformatics

Of the total 959 faecal samples collected, 793 were used for metabarcoding, selected by balancing

subsampling across nestboxes and dates and enforcing an upper limit of ten samples per nestbox

per year [29]. If multiple faeces were present in the sample tube, part of each was used for DNA

extraction. Thirty samples were also processed in duplicate by dividing the faecal sample into two to

assess repeatability. Twenty-four experimental controls were also included (six extraction negatives,

nine PCR negatives and nine PCR positives using *Inga pezizifera* as a non-native plant PCR positive).

DNA extraction was performed using the QIAamp DNA Stool Mini kit, following a modified protocol

that improved yields [29]. PCR amplification of three loci (COI, 16S and rbcL) was performed for the

broader project; of particular importance to this study was the rbcL 'minibarcode' designed to detect

184 base pairs of plant DNA. A second PCR subsequently added indexed Illumina adaptors to the

amplicons from each sample; amplicons were then multiplexed in three pools and each pool

sequenced on an Illumina MiSeq using 150 base pair paired-end reads.

Sequencing reads were demultiplexed, trimmed and clustered into molecular operational taxonomic

units (MOTUs) as per the bioinformatics protocol detailed in [29]. The taxonomic identity of MOTUs

was determined using a BLAST search of the reference set of MOTU sequences against the GenBank

and BOLD public databases.

Samples were initially screened for the presence of blue tit sequence at the 16S locus and those with

fewer than 100 reads of blue tit were excluded from further analyses (n=9) following [29]. No non-

blue tit avian DNA was found in any sample. All nine PCR positive control samples contained MOTUs

attributable to *Inga pezizifera* (range of reads = 4,007 – 12,697) and no more than 19 reads of

another MOTU. All nine PCR negative control samples and three of the six extraction negative

control samples contained no more than 22 reads of any MOTU. The remaining three extraction

negative control samples showed high numbers of reads (n = 991 – 6,302) from contaminating

tomato but nothing else. Systematic contamination at the rbcL locus was investigated by assessing

row and column MOTU correlations [29] but no systematic contamination was found. As there were

few cases where a control had >20 reads for any nontarget MOTU, we adopted 20 reads as the cut-

off for classifying a MOTU as being present.

MOTUs with <90% match to their best BLAST hit were then discarded as inconclusively identified.

Remaining MOTUs were amalgamated based on their genus-level identification, as identification to

species was seen to be unreliable, consistent with previous assessment of this section of rbcL [41]. A

total of 185 plant genera were identified and compared with common supplementary garden bird

foods to extract relevant genera. All further analyses were carried out only on these identified

supplementary food taxa within focal samples (excluding experimental replicate and control

samples, and those not confirmed to be from blue tit). Although the detection of plant DNA could be

due either to direct ingestion of that plant taxon by a blue tit itself or that plant being secondarily

present in the gut of ingested invertebrate prey, our focus on supplementary food taxa not present

in the general Scottish environment means our inference almost certainly reflects solely the direct

diet of blue tits and not those of their animal prey.

Statistical analyses

The first model examined how supplementary food consumption varied with respect to

environmental factors. A binary value of presence/absence of supplementary food in a faecal sample

was used as the response variable of a Bayesian generalised linear mixed model (GLMM) [42], with

distance to nearest human habitation, date, elevation, latitude, year and number of faeces in sample

as fixed predictor variables. Year and number of faeces (1-3 and unknown) were categorical factors,

while the remainder were continuous variables; the latter were mean centred for ease of

interpretation [43] and to facilitate model convergence. Distance to human habitation was analysed

on the logarithmic scale due to right skewed data and consistent with a decay model. Date was

coded as a deviation from the respective sample site mean per year, as different sites and years

have different blue tit breeding phenology. Site and nestbox were included as random effects and

the model was run for ten million iterations, removing the first 100,000 as burn-in and thinning

every 100. A binomial error structure was used along with parameter expanded priors for the

variance terms with residual variance fixed at 0.5. Repeatability in the detection of supplementary

food consumption was analysed by calculating a Jaccard similarity index for 29 replicate pairs of

samples (one was removed during quality control steps above).

The second set of models aimed to infer whether supplementary food consumption affected the

breeding parameters and adult condition of blue tits. For this analysis the mean supplementary food

consumption at each site was calculated. Site-level mean consumption was used rather than nestbox

level consumption for two reasons: i) blue tits often do not nest in a nestbox they are roosting in

prior to breeding, but rather nearby, precluding direct attribution, ii) faeces were only produced in

certain nestboxes so most nesting attempts are not in a nestbox from which faeces were collected.

In addition, if we assume that the mean supplementary food consumption at a site is representative

of all individuals nesting there then framing the analysis at the site level benefits the sample size and

power. Firstly, nestbox occupancy was treated as the response variable in a Bayesian GLMM [42]

containing mean supplementary food consumption (varying 0-1), elevation, latitude and year (as a

factor) as fixed predictor variables, and site and nestbox as random effects, with all numeric

predictor variables mean-centred. A binomial error structure was used with similar priors to the first

model. Similar models were then run with first egg laying date, clutch size, productivity and adult

blue tit mass as response variables with Gaussian error structures and no fixed residual variance. In

addition to the standard fixed predictor variables mentioned above, the mass model also contained

the age and sex of the bird.

To gain an indication of whether supplementary feeder usage and competition with blue tits may be

affecting UK woodland bird populations over time, the 25-year population trends of potential

competitor forest bird species were analysed. 21 species were included based on the following

criteria: average body length less than twice a blue tit (<24cm), foraging substantially on foliage-

gleaned invertebrates during the breeding season, occupying wooded habitats, and with a

substantial enough UK population to have a 25-year BTO BirdTrends population trend estimate [44].

Four categorical variables were coded for each species: their population trend (1 = > -50%, 2 = -11 to

-50%, 3 = -10 to +10%, 4 = +11 to +50%, 5 = > +50%; Massimino et al., 2019), supplementary garden

bird feeder usage (1 = rare or never, 2 = frequent (>5% occurrence in 2020 RSPB Big Garden

Birdwatch www.rspb.org.uk/get-involved/activities/birdwatch/results/)), competition status versus

blue tit (1 = outcompeted (average lower mass and/or published evidence of outcompetition for

food or breeding sites; Massimino et al., 2019; Samplonius & Both, 2019; Smith & Smith, 2020), 2 =

not outcompeted (all others)), and competition type (1 = food, 2 = food and breeding site (if nesting

in small cavities; Holden & Cleeves, 2010)). Three Welch's Two-sample T-tests were then conducted

to analyse how population trend varied with regards to i) supplementary feeder usage, ii)

competition status, and iii) competition type.

Results

Out of 785 faecal samples, 53% (n = 414) contained evidence of supplementary food consumption.

Five supplementary plant foodstuffs were identified, with peanut (*Arachis*) by far the most common,

present in 49% of total samples. Sunflower (*Helianthus*) was also highly prevalent (17%), with maize

(*Zea*) (9%), barley (*Hordeum*) (5%) and millet (*Panicum*) (1%) all rarer. 63% of samples containing

supplementary food contained only one type, with 37% containing more than one supplementary

foodstuff, and two samples containing all five. Experimental repeatability was high both for

detecting peanuts (Jaccard similarity = 0.923) or any supplementary food (Jaccard similarity = 0.923)

between replicate samples.

Increasing distance to nearest human habitation predicted a significant reduction in supplementary

food consumption (Table 1, Fig 1a). Different years also had significantly different supplementary

food consumption rates. In 2014 the model predicted a 93% chance of a faecal sample containing

supplementary foodstuffs at the shortest distances examined in our study (33m) reducing to 29% at

200m, 6% at 500m, and 1% at the furthest site distances examined (1384m), while in 2015 these

figures were higher, with 97% chance at 33m, 51% chance at 200m, 15% chance at 500m, and 2%

chance at 1384m to the nearest human habitation (Fig 1a). The faeces collected at the site furthest

from human habitation did however show supplementary food consumption in 75% of samples in

2015 (Fig 1a).

Supplementary food use also significantly declined through the sampling period, in the run-up to

breeding (Table 1, Fig 1b). In 2014 the model predicted a 65% chance of a faecal sample containing

supplementary food at the earliest sampling times (70 days before mean first egg laying), declining

to 24% by 30 days to egg laying and 7% by egg laying (Fig 1b). For 2015 these figures were elevated

to 83% at the earliest times, 44% in the mid time frame and 17% at egg laying (Fig 1b). Elevation and

latitude showed no significant effect on supplementary food consumption, and combining more

faeces per sample increased the likelihood of supplementary food detection (Table 1). Site and

nestbox random effects explained similar amounts of variance (Table 1).

Increased supplementary food consumption significantly predicted a large increase in nestbox

occupation, from a 20% likelihood with no supplementary food consumption to a 75% likelihood

with supplementary food present in every faecal sample (Table 2, Fig 2a). Supplementary food

consumption also significantly advanced egg laying date by five days (from day 122 to day 117, Table

2, Fig 2b). However, it did not significantly affect clutch size, productivity or the mass of either male

or female parent blue tits (Table 2).

The t-tests showed that the population trends of feeder-using and non-feeder-using bird species

were significantly different (t = -2.3, df = 18.0, p = 0.03, Fig 3), with feeder-using species increasing

on average and non-feeder-using species decreasing. Similarly, the population trends of species

outcompeted by blue tits and those not outcompeted by blue tits were significantly different (t = -

2.4, df = 17.8, p = 0.03, Fig 3), with those outcompeted on average declining. While competition type

did not significantly predict population trends (t = 0.9, df = 17.7, p = 0.4), a non-significant suggestion

that those species competing with blue tits for both food and nesting sites declining more than those

only competing for food was observed (Fig 3).

Discussion

This study reveals just how prevalent and ubiquitous supplementary food is in the diet of a wild bird

species in a country with high provisioning rates [2,4,32]. Supplementary foodstuffs were shown by

faecal metabarcoding to be present in the majority (53%) of blue tit faecal samples immediately

prior to breeding, with peanuts identified in more faecal samples (49%) than any other single dietary

item, natural or supplementary. For comparison, the most frequent natural prey item, the moth

Argyresthia goedartella, was present in 34% of the same samples [29]. We show blue tits can travel

almost 1.4km to use supplementary bird feeders during a time of year when movement is thought to

be restricted around breeding territories [39]; however, we note our study measured just the

distance to the closest human habitation so birds may be moving even further than this. Indeed, as

the study area incorporates some of the more remote parts of the UK, these results reveal it likely

that supplementary food is available to almost every blue tit in the UK (and other feeder-using bird

species, as blue tits are relatively sedentary and short-winged compared to other feeder users [39]),

with implications likely to extrapolate across large parts of the western world due to similarly high

supplementary feeding rates [11,12]. We infer from this that any impacts from supplementary

feeding will be felt far wider than solely in urban environments as has hitherto been assumed [1,33].

As we find that supplementary food usage is strongly associated with a dramatic increase in nestbox

occupation (a proxy of breeding density) and an advance in lay date, it is perhaps unsurprising then

that we find the national population trends of supplementary feeder-using woodland bird species

are increasing on average while the populations of competitor species not benefitting from

supplementary feeders are decreasing.

As predicted, supplementary food use declined with increasing distance to nearest human

habitation. While this relationship has previously been assumed [18,19], we believe our

quantification of it to be the first in a natural situation, made possible by the use of a highly

repeatable faecal metabarcoding procedure. Supplementary food usage was still considerable at

several hundred metres from the closest potential feeder, yet this distance is greater than the cut-

off distance used between treatments and/or nearby human habitation in previous supplementary

feeding experiments [19]. Therefore, widespread feeder usage may contribute a background or even

confounding effect that was inadequately accounted for in many previous experimental contexts.

The distance travelled to supplementary food, and overall usage rates, differed markedly between

the two years in our study, with 2015 having higher uptake rates than 2014. This may be due to 2015

being considerably colder across our study region, as natural food levels are lower in these

conditions [29,48], and benefits from supplementary feeding larger due to natural nutrient

limitation, concurring with previous studies [15]. In addition to a distance decay, supplementary

food usage also declined over a temporal gradient throughout our study period. This is presumably

due to large increases in natural invertebrate prey as spring progresses [29,49] alongside individuals

being more restricted to breeding territories as nesting commences [39]. There was no impact of the

geographic gradients of latitude and elevation, which vary substantially over the study region,

indicating a spatially widespread similarity in feeder usage.

Previous research has developed a mixed picture of the benefits and costs that supplementary

feeding confers on the species, including blue tits, using these extra resources. Using faecal

metabarcoding to identify definite rather than assumed supplementary food intake without the

need for additional experimentation has allowed us to demonstrate major fitness benefits conferred

upon blue tits at sites with higher supplementary food uptake. A change in supplementary food use

between the observed lowest and highest values predicted an almost four-fold increase in nestbox

occupation, an accurate proxy for breeding density in our system due to sites having equal numbers

of equally spaced nestboxes [36]. We expect increasing breeding densities to extrapolate to other

feeder-using species as feeder presence increases local abundances of feeder-using species [34],

providing an explanation for bird breeding densities co-varying with human household densities

[50]. The five-day advancement in egg laying we identify is very similar to that found in previous

food supplementation studies [15,19] and may represent the limit of the plastic phenological

response to the lifting of an energetic constraint, with earlier laying associated with higher

productivity [51]. Perhaps this is why individual nest productivity didn't decline due to density

effects as might be imagined [52], but instead showed a minor increase. Clutch size not being

significantly predicted by supplementary feeding agrees with previous studies [19] and reinforces

that environmental aspects seem to have little effect on clutch size [36].

Many species utilising supplementary feeding, such as blue tits, are common, adaptable and already

at population carrying capacity [38,53]. Boosting the productivity, survival, fitness and breeding

densities of such species without any increase in available habitat or natural resources is likely to

affect their competitors negatively, particularly those not utilising the new supplementary resources

[38]. This may be particularly evident in woodland species compared to farmland species, as rather

than replace natural resources that have been lost to all species due to landscape intensification

[54], supplementary feeding is providing additional resources solely to certain species. To this end,

we demonstrate that populations of UK woodland bird species that don't use supplementary feeders

are likely to be declining whereas those that do are likely to be increasing over the last 25 years. In

addition, species that are less behaviourally dominant than blue tits are likely to be declining

whereas more dominant species are likely to be increasing. Supplementary feeding is therefore likely

a driver of population change, in line with other recent evidence [33], however at a much larger

scale than solely urbanised environments.

While we do not analyse a causal link between supplementary feeding and the declines of these

competitor species, the mechanisms whereby increased blue tit densities could impact other species

are clear. For example, blue (and great) tits are frequently known to evict species such as willow tit

and lesser spotted woodpecker from nest holes that they have excavated [45,55], kill pied

flycatchers when claiming nesting sites [46], and dominate subordinate marsh and willow tits at food

resources [39]. Abundant and permanent feeding might also eliminate any competitive advantage

other species (such as marsh tits) exhibit in finding and exploiting natural resources first [56], before

being outcompeted by dominant species like blue tits, or migrating to warmer climes for winter to

avoid winter starvation, as for pied flycatchers [57]. Supplementary feeding therefore, although well-

intentioned and beneficial to the species partaking, may be shifting the competitive balance of

natural ecosystems and the structures enabling community coexistence, favouring certain species at

the expense of others, and contributing to human-mediated ecological homogenisation [37,38].

In conclusion, we reveal through faecal metabarcoding the pervasiveness of supplementary

foodstuffs in the diet of a wild bird and the large benefits using these substantial additional

resources confer on its breeding density and phenology. We also show that the distances travelled

to utilise these resources are further than previously imagined, even in a largely sedentary species at

a time of year when movement is thought to be restricted. This indicates that the effects of

supplementary feeding on ecosystems are likely to extend far beyond just urban environments as

has hitherto been assumed. Finally, we demonstrate that species making use of supplementary

resources are likely to have increasing populations while those that do not are likely declining,

possibly due to shifting competition balances and ecosystem dynamics. As supplementary

provisioning of wildlife (both intentional and incidental) is hugely prevalent and increasing [2], it may

have large and widespread ramifications for biodiversity conservation, and we urge caution upon

policy makers advocating supplementary feeding for wildlife engagement.

Acknowledgements

We wish to thank Ally Phillimore for commenting on an earlier draft of the manuscript and providing

laboratory funding, Jarrod Hadfield for additional laboratory funding, Irene Benedicto Cabello and Ed

Ivimey-Cook for assistance with fieldwork, and all the landowners and managers who allowed us

access to their land. This work was funded by a NERC Doctoral Training Studentship (NE/1338530) to

JDS and faecal metabarcoding was funded by NERC grant NE/I020598/1 to Ally Phillimore.

Authors' Contributions

JDS conceived and managed the study, conducted the fieldwork, contributed to data curation,

analysed the data, and wrote the manuscript. UHT contributed to data curation. JAN devised and

conducted all laboratory protocols and contributed to later manuscript editing.

Data Availability Statement

Data will be made publicly available via the Dryad digital repository.

411
412 **References**
413

1. Jones DN, Reynolds SJ. 2008 Feeding Birds in Our Towns and Cities: A Global Research
Opportunity. *J. Avian Biol.* **39**, 265–271. (doi:10.1111/j.2008.0908-8857.04271.x)
2. Cox DTC, Gaston KJ. 2018 Human–nature interactions and the consequences and drivers of
provisioning wildlife. *Philos. Trans. R. Soc. B Biol. Sci.* **373**, 20170092.
(doi:10.1098/rstb.2017.0092)
3. Hanmer HJ, Thomas RL, Fellowes MDE. 2017 Provision of supplementary food for wild birds
may increase the risk of local nest predation. *Ibis (Lond. 1859).* **159**, 158–167.
(doi:10.1111/ibi.12432)
4. Davies ZG, Fuller RA, Dallimer M, Loram A, Gaston KJ. 2012 Household factors influencing
participation in bird feeding activity: A national scale analysis. *PLoS One* **7**.
(doi:10.1371/journal.pone.0039692)
5. Davies ZG, Fuller RA, Loram A, Irvine KN, Sims V, Gaston KJ. 2009 A national scale inventory of
resource provision for biodiversity within domestic gardens. *Biol. Conserv.* **142**, 761–771.
(doi:10.1016/j.biocon.2008.12.016)
6. PFMA. 2018 PFMA's pet data report 2018.
7. Robb GN, McDonald RA, Chamberlain DE, Bearhop S. 2008 Food for thought: supplementary
feeding as a driver of ecological change in avian populations. *Front. Ecol. Environ.* **6**, 476–484.
(doi:10.1890/060152)
8. Orros ME, Fellowes MDE. 2015 Wild Bird Feeding in an Urban Area: Intensity, Economics and
Numbers of Individuals Supported. *Acta Ornithol.* **50**, 43–58.
(doi:10.3161/00016454ao2015.50.1.006)
9. Hanmer HJ, Thomas RL, Fellowes MDE. 2018 Introduced Grey Squirrels subvert
supplementary feeding of suburban wild birds. *Landsc. Urban Plan.* **177**, 10–18.
(doi:10.1016/j.landurbplan.2018.04.004)
10. Reed JH, Bonter DN. 2018 Supplementing non-target taxa: bird feeding alters the local
distribution of mammals. *Ecol. Appl.* **28**, 761–770. (doi:10.1002/eap.1683)
11. Baverstock S, Weston MA, Miller KK. 2019 A global paucity of wild bird feeding policy. *Sci.*
Total Environ. **653**, 105–111. (doi:10.1016/j.scitotenv.2018.10.338)
12. Jones DN. 2011 An appetite for connection: Why we need to understand the effect and value
of feeding wild birds. *Emu* **111**. (doi:10.1071/MUv111n2_ED)
13. Robb GN, McDonald RA, Inger R, Reynolds SJ, Newton J, McGill RAR, Chamberlain DE,
Harrison TJE, Bearhop S. 2011 Using stable-isotope analysis as a technique for determining
consumption of supplementary foods by individual birds. *Condor* **113**, 475–482.
(doi:10.1525/cond.2011.090111)

14.   Orros ME, Thomas RL, Holloway GJ, Fellowes MDE. 2015 Supplementary feeding of wild birds
indirectly affects ground beetle populations in suburban gardens. *Urban Ecosyst.* **18**, 465–
475. (doi:10.1007/s11252-014-0404-x)

15.   Nager RG, Ruegger C, van Noordwijk AJ. 1997 Nutrient or energy limitation on egg formation:
A feeding experiment in great tits. *J. Anim. Ecol.* **66**, 495–507.

16.   Robb GN, McDonald RA, Chamberlain DE, Reynolds SJ, Harrison TJEE, Bearhop S. 2008 Winter
feeding of birds increases productivity in the subsequent breeding season. *Biol. Lett.* **4**, 220–
223. (doi:10.1098/rsbl.2007.0622)

17.   Peach WJ, Sheehan DK, Kirby WB. 2014 Supplementary feeding of mealworms enhances
reproductive success in garden nesting House Sparrows Passer domesticus. *Bird Study* **61**,
378–385. (doi:10.1080/00063657.2014.918577)

18.   Plummer KE, Bearhop S, Leech DI, Chamberlain DE, Blount JD. 2013 Winter food provisioning
reduces future breeding performance in a wild bird. *Sci. Rep.* **3**, 2002.
(doi:10.1038/srep02002)

19.   Harrison TJE, Smith J a, Martin GR, Chamberlain DE, Bearhop S, Robb GN, Reynolds SJ. 2010
Does food supplementation really enhance productivity of breeding birds? *Oecologia* **164**,
311–20. (doi:10.1007/s00442-010-1645-x)

20.   Knutie SA. 2020 Food supplementation affects gut microbiota and immunological resistance
to parasites in a wild bird species. *J. Appl. Ecol.* **57**, 536–547. (doi:10.1111/1365-2664.13567)

21.   Plummer KE, Bearhop S, Leech DI, Chamberlain DE, Blount JD. 2013 Fat provisioning in winter
impairs egg production during the following spring: a landscape-scale study of blue tits. *J.*
Anim. Ecol. **82**, 673–82. (doi:10.1111/1365-2656.12025)

22.   Lawson B, Robinson RA, Toms MP, Risely K, Macdonald S, Cunningham AA. 2018 Health
hazards to wild birds and risk factors associated with anthropogenic food provisioning. *Philos.*
Trans. R. Soc. B Biol. Sci. **373**. (doi:10.1098/rstb.2017.0091)

23.   Moyers SC, Adelman JS, Farine DR, Thomason CA, Hawley DM. 2018 Feeder density enhances
house finch disease transmission in experimental epidemics. *Philos. Trans. R. Soc. B Biol. Sci.*
373. (doi:10.1098/rstb.2017.0090)

24.   Greig EI, Wood EM, Bonter DN. 2017 Winter range expansion of a hummingbird is associated
with urbanization and supplementary feeding. *Proc. R. Soc. B Biol. Sci.* **284**, 20170256.
(doi:10.1098/rspb.2017.0256)

25.   Plummer KE, Siriwardena GM, Conway GJ, Risely K, Toms MP. 2015 Is supplementary feeding
in gardens a driver of evolutionary change in a migratory bird species? *Glob. Chang. Biol.* **21**,
4353–4363. (doi:10.1111/gcb.13070)

26.   Ishigame G, Baxter GS, Lisle AT. 2006 Effects of artificial foods on the blood chemistry of the
Australian magpie. *Austral Ecol.* **31**, 199–207. (doi:10.1111/j.1442-9993.2006.01580.x)

27.   Bosse M *et al.* 2017 Recent natural selection causes adaptive evolution of an avian polygenic
trait. *Science (80-.).* **358**, 365–368. (doi:10.1126/science.aal3298)

28.   Pompanon F, Deagle BE, Symondson WOC, Brown DS, Jarman SN, Taberlet P. 2012 Who is
eating what: diet assessment using next generation sequencing. *Mol. Ecol.* **21**, 1931–1950.
(doi:10.1111/j.1365-294X.2011.05403.x)

29.   Shutt JD, Nicholls JA, Trivedi UH, Burgess MD, Stone GN, Hadfield JD, Phillimore AB. 2020
Gradients in richness and turnover of a forest passerine's diet prior to breeding: A mixed
model approach applied to faecal metabarcoding data. *Mol. Ecol.* , 1199–1213.
(doi:10.1111/mec.15394)

30.   Krüger F, Clare EL, Symondson WOC, Keišs O, Petersons G. 2014 Diet of the insectivorous bat
Pipistrellus nathusii during autumn migration and summer residence. *Mol. Ecol.* **23**, 3672–
3683. (doi:10.1111/mec.12547)

31.   Groom C, White NE, Mitchell N, Roberts JD, Mawson P. 2017 Assessing the spatial ecology
and resource use of a mobile and endangered species in an urbanized landscape using
satellite telemetry and DNA faecal metabarcoding. *Ibis (Lond. 1859).* **159**, 390–405.

(doi:10.1111/ibi.12442)

32.   Milligan ND, Radersma R, Cole EF, Sheldon BC. 2017 To graze or gorge: consistency and
flexibility of individual foraging tactics in tits. *J. Anim. Ecol.* **86**, 826–836. (doi:10.1111/1365-
2656.12651)

33.   Plummer KE, Risely K, Toms MP, Siriwardena GM. 2019 The composition of British bird
communities is associated with long-term garden bird feeding. *Nat. Commun.* **10**, 2088.
(doi:10.1038/s41467-019-10111-5)

34.   Fuller RA, Warren PH, Armsworth PR, Barbosa O, Gaston KJ. 2008 Garden bird feeding
predicts the structure of urban avian assemblages. *Divers. Distrib.* **14**, 131–137.
(doi:10.1111/j.1472-4642.2007.00439.x)

35.   Galbraith JA, Beggs JR, Jones DN, Stanley MC. 2015 Supplementary feeding restructures
urban bird communities. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E2648–E2657.
(doi:10.1073/pnas.1501489112)

36.   Shutt JD, Bolton M, Cabello IB, Burgess MD, Phillimore AB. 2018 The effects of woodland
habitat and biogeography on blue tit Cyanistes caeruleus territory occupancy and
productivity along a 220 km transect. *Ecography (Cop.).* **41**, 1967–1978.
(doi:10.1111/ecog.03573)

37.   Newbold T *et al.* 2019 Climate and land-use change homogenise terrestrial biodiversity, with
consequences for ecosystem functioning and human well-being. *Emerg. Top. Life Sci.*
(doi:10.1042/etls20180135)

38.   Oro D, Genovart M, Tavecchia G, Fowler MS, Martínez-Abraín A. 2013 Ecological and
evolutionary implications of food subsidies from humans. *Ecol. Lett.* **16**, 1501–1514.
(doi:10.1111/ele.12187)

39.   Perrins CM. 1979 *British Tits.*

40.   Nychka D, Furrer R, Paige J, Sain S. 2017 fields: Tools for spatial data.
(doi:10.5065/D6W957CT)

41.   Little DP. 2014 A DNA mini-barcode for land plants. *Mol. Ecol. Resour.* **14**, 437–446.
(doi:10.1111/1755-0998.12194)

42.   Hadfield JD. 2010 MCMC methods for multi-response generalized linear mixed models: The
MCMCglmm R package. *J. Stat. Softw.* **33**, 1–22. (doi:10.1002/ana.22635)

43.   Schielzeth H. 2010 Simple means to improve the interpretability of regression coefficients.
Methods Ecol. Evol. **1**, 103–113. (doi:10.1111/j.2041-210X.2010.00012.x)

44.   Massimino D *et al.* 2019 BirdTrends 2019: trends in numbers, breeding success and survival
for UK breeding birds.

45.   Smith KW, Smith L. 2020 Long-term trends in the nest survival and productivity of the Lesser
Spotted Woodpecker Dryobates minor in Britain. *Bird Study* **67**, 109–118.
(doi:10.1080/00063657.2020.1780195)

46.   Samplonius JM, Both C. 2019 Climate Change May Affect Fatal Competition between Two
Bird Species. *Curr. Biol.* **29**, 327-331.e2. (doi:10.1016/j.cub.2018.11.063)

47.   Holden P, Cleeves T. 2010 *RSPB Handbook of British Birds.* 3rd edn. London: Christopher
Helm.

48.   Southwood TRE, Wint GRW, Kennedy CEJ, Greenwood SR. 2004 Seasonality, abundance,
species richness and specificity of the phytophagous guild of insects on oak (Quercus)
canopies. *Eur. J. Entomol.* **101**, 43–50. (doi:10.14411/eje.2004.011)

49.   Shutt JD, Burgess MD, Phillimore AB. 2019 A spatial perspective on the phenological
distribution of the spring woodland caterpillar peak. *Am. Nat.* **194**, E109–E121.
(doi:10.1086/705241)

50.   Tratalos J, Fuller RA, Evans KL, Davies RG, Newson SE, Greenwood JJD, Gaston KJ. 2007 Bird
densities are associated with household densities. *Glob. Chang. Biol.* **13**, 1685–1695.
(doi:10.1111/j.1365-2486.2007.01390.x)

51.   Visser ME, van Noordwijk AJ, Tinbergen JM, Lessells CM. 1998 Warmer springs lead to

mistimed reproduction in great tits (Parus major). *Proc. R. Soc. B Biol. Sci.* **265**, 1867–1870.
(doi:10.1098/rspb.1998.0514)

52.   Dhondt AA, Kempenaers B, Adriaensen F. 1992 Density-Dependent Clutch Size Caused by
Habitat Heterogeneity. *J. Anim. Ecol.* **61**, 643–648.

53.   Galbraith JA, Jones DN, Beggs JR, Parry K, Stanley MC. 2017 Urban bird feeders dominated by
a few species and individuals. *Front. Ecol. Evol.* **5**. (doi:10.3389/fevo.2017.00081)

54.   Pierret P, Jiguet F. 2018 The potential virtue of garden bird feeders: More birds in citizen
backyards close to intensive agricultural landscapes. *Biol. Conserv.* **222**, 14–20.
(doi:10.1016/j.biocon.2018.03.033)

55.   Parry W, Broughton RK. 2018 Nesting behaviour and breeding success of Willow Tits Poecile
montanus in north-west England. *Ringing Migr.* **33**, 75–85.
(doi:10.1080/03078698.2018.1631610)

56.   Farine DR, Aplin LM, Sheldon BC, Hoppitt W. 2015 Interspecific social networks promote
information transmission in wild songbirds. *Proc. R. Soc. B Biol. Sci.* **282**.
(doi:10.1098/rspb.2014.2804)

57.   Lundberg A, Alatalo R. 2010 *The Pied Flycatcher*. A & C Black.

 Figures and Tables

Table 1 Environmental predictors of the consumption of supplementary food by blue tits. Results are
taken from a Bayesian GLMM with categorical error structure and logit link function, showing slope
estimates and credible intervals for each fixed and random term, with significance asterisks for
significant and near-significant terms (pMCMC ≤0.1 ° ≤0.05* ≤0.01** ≤0.001***). Numeric predictor
variables are mean centred, distance to habitation is log transformed, and date has been adjusted
for phenology by representing days before mean first egg laying at a given site within a given year.
The intercept value for year is 2014 and for number of faeces is one.

Fixed Effects	Coefficient (C.I.'s)
Intercept	-1.12 (-2.64 − 0.44)
Distance to habitation	-1.97 (-3.10 − -0.80) ***
Days before laying	0.04 (0.02 − 0.07) ***
Elevation	0.0005 (-0.0096 − 0.0104)
Latitude	-0.81 (-2.73 − 1.08)
Year 2015	0.95 (-0.12 − 2.03) °
Faeces = 2	-0.07 (-1.21 − 1.07)
Faeces = 3	1.37 (0.42 − 2.36) **
Faeces = unknown	2.25 (0.88 − 3.67) ***
Random Effects	
Site	5.85 (1.40 − 11.44)
Nestbox	5.01 (2.27 − 8.24)

577
578

Figure 1: Effects of a) distance to nearest human habitation and b) sampling date (adjusted for the phenology of the site within year) on the probability of supplementary food consumption by blue tits. The predicted response of each is shown in 2014 (gold) and 2015 (green). Asterisks in a) show the proportion of faeces containing supplementary foodstuff per site per year, not the presence/absence response analysed in the model.

Table 2 The effects of supplementary food intake and other environmental variables on a range of blue tit breeding parameters. Results are taken from Bayesian GLMMs, showing slope estimates and credible intervals for each fixed and random term, with significance asterisks for significant terms (pMCMC ≤0.05* ≤0.01** ≤0.001***). Occupancy is presented on a binomial scale with the other response variables gaussian. Numeric predictor variables are mean centred, and the intercept year is 2014, with intercept values of first year adult and female for the mass model. Supplementary food intake was calculated as a per site per year proportion of faeces containing supplementary food which was applied to all nests at that site in that year.

	Occupancy	Egg Laying	Clutch Size	Productivity	Mass
Fixed Effects	Coefficient (C.I.'s)	Coefficient (C.I.'s)	Coefficient (C.I.'s)	Coefficient (C.I.'s)	Coefficient (C.I.'s)
Intercept	0.06 (-0.44 – 0.60)	119.33 (117.57 – 121.06)	8.70 (8.28 – 9.12)	6.96 (6.25 – 7.63)	10.71 (10.57 – 10.84)
Supplementary Food	2.47 (1.14 – 3.70) ***	-4.39 (-8.50 – -0.17) *	-0.15 (-0.96 – 0.62)	0.87 (-0.47 – 2.11)	0.07 (-0.12 – 0.25)
Elevation	-0.01 (-0.01 – -0.01) ***	0.02 (0.01 – 0.03) **	0.001 (-0.002 – 0.004)	0.002 (-0.003 – 0.006)	-0.0006 (-0.0012 – 0.0002)
Latitude	-1.30 (-2.11 – -0.61) ***	0.79 (-1.82 – 3.20)	-0.55 (-0.99 – -0.09) *	-0.77 (-1.52 – -0.07) *	-0.13 (-0.25 – -0.03) *
Year 2015	1.20 (0.61 – 1.74) ***	4.41 (2.89 – 6.11) ***	-1.13 (-1.59 – -0.66) ***	-3.41 (-4.17 – -2.61) ***	0.003 (-0.14 – 0.12)
Year 2016	0.60 (0.08 – 1.18) *	8.19 (6.64 – 9.75) ***	-0.64 (-1.14 – -0.18) **	-1.54 (-2.34 – 0.74) ***	-0.13 (-0.27 – -0.002)
2nd Year +					0.06 (-0.04 – 0.16)
Sex Male					0.05 (-0.04 – 0.14)
Random Effects					
Site	0.71 (0 – 1.57)	12.41 (4.29 – 22.54)	0.24 (0 – 0.58)	0.60 (0 – 1.42)	0.008 (0 – 0.023)
Nestbox	0.74 (0 – 1.69)	2.76 (0 – 7.16)	0.15 (0 – 0.47)	0.15 (0 – 0.58)	0.003 (0 – 0.052)
Residual	0.5	30.98 (25.00 – 37.09)	2.90 (2.41 – 3.42)	8.19 (6.93 – 9.52)	0.23 (0.20 – 0.27)

Figure 2 Effects of site-level supplementary food intake on a) probability of nestbox occupancy and b) first egg laying date. Asterisks in a) depict site per year occupancy rates rather than the 0/1 occupied response analysed in the model. Predictions correspond to 2014.

Figure 3 Differing population trends of 21 insectivorous woodland bird species in the UK with regards to i) supplementary garden feeder usage ii) behavioural dominance in comparison to blue tits and iii) whether the species competes for nest sites with blue tits, with values depicting mean ± standard error with asterisks above significant differences as determined by a t-test. A trend value of 1 represents a large population decline (> -50%) and 5 a large population increase (>50%) over 25 years, with a stable population (trend value = 3) shown by a grey dashed line.