Dendritic domain-specific sampling of long-range axons shapes feedforward and feedback connectivity of L5 neurons

Alessandro R. Galloni[1,2], Zhiwen Ye[1] and Ede Rancz[1*]

[1]The Francis Crick Institute, London, United Kingdom; [2]University College London, United Kingdom.

*For correspondence: ede.rancz@crick.ac.uk

Abstract

Feedforward and feedback pathways interact in specific dendritic domains to enable cognitive functions such as predictive inference and learning. Based on axonal projections, hierarchically lower areas are thought to form synapses primarily on dendrites in middle cortical layers, while higher-order areas are posited to target dendrites in layer 1 and in deep layers. However, the extent to which functional synapses form in regions of axo-dendritic overlap has not been extensively studied. Here, we use viral tracing in the visual cortex of mice to map brain-wide inputs to a genetically-defined population of layer 5 pyramidal neurons. Furthermore, we provide a comprehensive map of input locations through subcellular optogenetic circuit mapping. We show that input pathways target distinct dendritic domains with far greater specificity than appears from their axonal branching, often deviating substantially from the canonical patterns. Common assumptions regarding the dendrite-level interaction of feedforward and feedback inputs may thus need revisiting.

Introduction

One of the key organizing principles of connectivity within the neocortex is thought to be hierarchy between cortical areas. This notion was originally proposed by Hubel & Wiesel (Hubel & Wiesel 1962) to account for the increasing receptive field complexity in areas progressively further from the retina. In purely feedforward (FF) networks, such as artificial neural nets used successfully in computer vision (LeCun et al 2015), hierarchy is generally defined by synaptic distance from the sensory periphery. However, in highly recurrent networks like the cortex, it is not possible to apply this definition consistently beyond the initial levels. Instead, the laminar patterns of axonal projections are often used to deduce relative levels of hierarchy. For example, the projection from primary to secondary visual cortex, which is classically defined as FF, is characterised by dense axon terminations in middle cortical layers (particularly L4). Meanwhile, the projection from secondary to primary visual cortex, feedback (FB) by definition, primarily targets L1, and to a lesser extent deeper layers (Rockland & Pandya 1979). This pattern of FF and FB projections also appears in many other brain regions, and has been used as a proxy to describe the hierarchical relationships between a large number of areas across the brain (D'Souza et al 2016, D'Souza et al 2020, Felleman & Van Essen 1991, Harris et al 2019, Wang et al 2020b, Zeng 2018).

Projections, however, do not guarantee functional connections. The link between the two is called Peters' rule, which postulates that the probability of synaptic connections can be predicted from the overlap between axonal and dendritic arbours (Rees et al 2017). While synapses can only form where there is overlap between axons and dendrites, the converse is not necessarily true. The link between axo-dendritic overlap and connection probability may thus be overly simplistic and requires further scrutiny. While some studies have found support for Peters' rule at the level of functional synaptic connectivity for interneurons (Fino & Yuste 2011, Packer et al 2013, Rieubland et al 2014), its general applicability has been refuted, at least for local networks, by dense anatomical reconstructions of retinal (Briggman et al 2011, Helmstaedter et al 2013, Kim et al 2014, Krishnaswamy et al 2015) and cortical circuits (Kasthuri et al 2015, Lee et al 2016). To investigate how this principle applies to long-range projections, a technique that has become widely adopted is subcellular channelrhodopsin-assisted circuit mapping (sCRACM). Here optogenetics is combined with spatially targeted optical stimulation to map the distribution of synaptic currents for a given input (Petreanu et al 2009). While this has been used to show that different presynaptic populations target dendritic subdomains with high specificity (Collins et al 2018, Hooks et al 2013, Little & Carter 2012, Yamawaki et al 2019), the extent to which this can be explained and predicted by the distribution of axons and dendrites remains largely an open question. This is particularly relevant for the case of layer 5 pyramidal neurons (L5PN), given their central role in several theories of cortical computation (Aru et al 2020, Guerguiev et al 2017, Larkum 2013, Richards et al 2019). Furthermore, the interaction between FF and FB information streams across cortical layers (Larkum et al 2018) is thought to underlie the brain's ability to modulate sensory perception (Larkum 2013, Takahashi et al 2020, Takahashi et al 2016) and implement global inference algorithms such as predictive coding (Shipp 2016).

All these approaches assume Peters' rule, according to which FF connectivity to L5PNs should target
primarily basal dendrites, while FB connectivity preferentially synapses on the apical tuft. The evidence
for this remains circumstantial and a direct examination of Peters' rule across a large number of inputs to
individual L5PNs remains to be done.

Here we present a comprehensive description of the functional input connectivity to a population of thick-
tufted layer 5 (ttL5) pyramidal neurons in the visual cortex. In particular, we set out to do three things:
determine the source of input connectivity to ttL5 neurons using monosynaptically restricted rabies tracing
(Kim et al 2015, Reardon et al 2016), create a census of subcellular input maps using sCRACM, and test
Peters' rule directly by comparing synaptic input maps to the respective axonal projection maps. As
higher order cortical regions are likely to receive a broader diversity of long-range inputs, we recorded
from a genetically defined population (Colgalt2-Cre; Groh et al 2010, Kim et al 2015) with sparse and
ttL5-specific labelling spanning the mouse medial secondary visual areas (VISam, VISpm and RSPagl,
collectively referred to as V2m following Franklin & Paxinos 2007).

71 Results

72 Brain-wide input map to V2m ttL5 pyramidal neurons

We employed a monosynaptically restricted rabies virus approach (Reardon et al 2016, Wickersham et al
2007) to generate a presynaptic input map of V2m ttL5 neurons. Briefly, a mix of adeno-associated
viruses carrying floxed N2c G-protein, or TVA-receptor and EGFP genes were injected into V2m of
Colgalt2-Cre mice under stereotaxic guidance. Five to seven days later, mCherry expressing EnvA-CVS-
N2c-ΔG rabies virus was injected at the same location. After a further 10-12 days, the brains (n = 3) were
fixed and imaged using serial section 2-photon tomography. These were then registered to the Allen
common coordinate framework (CCFv3) and presynaptic cell bodies were detected and counted using an
automated pipeline (see Methods for details). Names and abbreviation for brain areas follow the Allen
CCFv3 (Wang et al 2020a) and are given in Figure 1—figure supplement 1.

Cell density maps for an example experiment are shown in Figure 1. The fifteen starter cells scattered
across V2m (Figure 1A) revealed presynaptic input neurons in a broad range of cortical and subcortical
areas, with the vast majority located in the ipsilateral cortex and thalamus (94 ± 1%; n = 3 mice; Figure
1B,C). We have grouped the most prominent input areas into proximal cortex (53%), distal cortex (8%)
and thalamus (19%; Figure 1E) and calculated total input proportion as the number of presynaptic cells in
an input area divided by the total number of presynaptic cells (Figure 1—supplementary table 1).

The majority of input was provided by VISp (10.6 ± 3.2 %) and the three RSP areas (29 % in total), while

orbitofrontal cortex (ORB; 2.7 ± 1.1 %) and the anterior cingulate area (ACA; 1.9 ± 1 %) provided the

most significant (albeit still modest) distal cortical input. Interestingly, while the majority of cortical input

cells were located in the infragranular layers, especially layer 5, input from ORB was almost exclusively

from layer 2/3 (Figure 1D,E). Prominent thalamic inputs were also observed, originating mainly in the

lateral posterior nucleus (LP, 10.3 ± 1.7 %) and anterior thalamic nuclei (ATN, composed of areas LD, 4.9
± 0.8 %, and AM, 2.2 ± 0.5 %). Cell counts for individual experiments and all brain areas can be found in
Figure 1—supplementary table 1.

To understand the organization of inputs onto ttL5 neurons in V2m, we chose to focus our efforts on 7
prominent input areas. VISp and V2m for FF input; RSPg (RSPv, RSPd), ACA and ORB for cortical FB
input; and LP and ATN for thalamic FB connections. These areas in total represent 72% of all presynaptic
input neurons.

Axonal projection patterns in V2m define putative synaptic regions

To determine axonal projection patterns of the previously selected areas and subcellular synaptic
connectivity to ttL5 neurons in V2m, we injected AAVs expressing Chronos-eGFP into input areas. Acute
brain slices were prepared after a minimum of 3 weeks of survival to obtain electrophysiological
recordings. The brain slices were then fixed, and axonal projections imaged using confocal microscopy.

Figure 2. Axonal projection densities from different input areas. A-C. Top: coronal brain pictograms indicating
the injection sites. Middle: example confocal images from V2m showing axonal projections (cyan) and Colgalt2-Cre
cell bodies (red). Bottom: projection density profiles averaged across 3-5 injections.

The spatial distribution of axons followed three basic patterns. Axons from VISp and ORB were densest
in layer (L) 2/3 and L5 while little projection was apparent in L1, reminiscent of the classical FF projection
pattern (Figure 2A). In contrast, axons from RSPg and ACA showed an FB-like pattern with dense
labelling in the middle part of L1 followed by sparse labelling in L2 and diffuse axons in layers 3, 5 and 6
(Figure 2B). The final group, which consists of the thalamic projections from LP and ATN, showed the
classical FB pattern strongly innervating the external part of L1, with a secondary peak in L3, but little or
no projections in layers 2, 5 and 6 (Figure 2C).

Having established where it is possible for synaptic connections to form, we next sought to determine the
dendritic location of synaptic inputs from the selected areas.

Subcellular optogenetic input mapping reveals diverse targeting of dendritic domains by input areas

To determine the spatial distribution of synaptic inputs to ttL5 neurons in V2m, we performed sCRACM
experiments from selected input areas identified by rabies tracing. Following expression of the
optogenetic activator Chronos, we made voltage-clamp recordings (at -70 mV) from ttL5 (Colgalt2-Cre)
neurons in acute brain slices. Optical stimulation with a 463 nm laser was spatially targeted using a digital
micromirror device (Figure 3A). Sodium and potassium channels were blocked using TTX (1μm) and 4-
AP (100 μm) to ensure that evoked currents were restricted to directly stimulated nerve terminals and to
enhance presynaptic release, respectively. The stimulus consisted of 24 x 12 spots of light in a 1000 x
500 μm grid aligned to the axis of the apical dendrite of the recorded neuron and covering the whole
depth of cortex. For each input area, we recorded 14-27 cells from 3-9 animals.

To estimate synaptic strength at each location, we measured the area of evoked synaptic currents
(corresponding to charge; Figure 3B) and made normalized 2D maps of the spatial distribution of inputs
(Figure 3C). Individual maps were then aligned (either to the pia or soma) and averaged. To quantify the
spatial location of inputs, we projected the average 2D maps in directions parallel (Figure 3C) or
perpendicular to the apical dendrite (Figure 3D). Furthermore, we defined the spatial distribution of the
three main dendritic compartments - the apical tuft, oblique, and basal dendrites - on the basis of 11
morphologically reconstructed Colgalt2-Cre neurons. The horizontal projection of the average morphology
was used to separate the contribution of each dendritic domain to the total synaptic input (Figure 3D; see
Methods for details).

To test the detectability of distal inputs regardless of their synaptic strength, we examined the correlation
between the location of synaptic input and the total synaptic charge evoked by full-field stimulation
(Figure 3—figure supplement 1). No correlation was found for any of the recorded areas, suggesting no
detection bias for distal inputs.

Figure 3. Using sCRACM to map subcellular connectivity from proximal cortical input areas. A. Experimental
setup and micrograph showing a brain slice with Chronos expression in RSPg and recording pipette in V2m. The
stimulation grid is overlaid, and an example spot is highlighted in blue. **B**. example sCRACM recording of excitatory
synaptic currents (red > 7 x S.D.) for an example neuron. **C.** Charge heatmap corresponding to recording in B with
the morphology of the recorded neuron overlaid. **D.** Normalized vertical profile of input map in C. Right: average
morphology profile used for dendritic domain deconvolution.

Primary visual cortex

We first recorded optically evoked synaptic responses arising from VISp axons (n = 14 cells from 7
animals, average soma depth 493 ± 18 µm; Figure 4A). The apical tuft received 57% of the input, with a
peak input located 198 µm from the pia (Figure 4—supplementary table 1). The remaining input was
spread between the oblique compartment, receiving 25%, and basal dendrites, receiving 18% of the total
input. The majority of recorded neurons had the peak input in the apical compartment (n = 10 / 14) while a
few cells lacked apical input (Figure 4—figure supplement 1A). Soma depth for individual recordings did
not correlate with either tuft or perisomatic input location, suggesting that VISp synapses on ttL5 neurons
target cortical layers rather than specific dendritic domains (Figure 4—figure supplement 1A). The
horizontal input distribution showed a slight medial skew (towards RSPg), most prominent in the oblique
and basal compartments (-42 µm for both; -21 µm in the tuft; Figure 4—figure supplement 1B). The total
synaptic charge measured via the somatic recording following full-field stimulation was 1.46 ± 0.33 pC

(Figure 4A). VISp thus provides moderate direct input to ttL5 neurons in V2m, primarily targeting the

proximal part of the apical tuft (0.84 pC) with smaller input arriving to the oblique (0.36 pC) and basal

(0.26 pC) compartments.

Figure 4. Subcellular connectivity maps of FF input areas. A. *i*: confocal image of a representative brain slice

(blue = DAPI) showing the injection site in VISp (green) and recorded neurons in V2m (red). *ii*: pia-aligned average

sCRACM heatmap for VISp inputs. Triangles represent soma locations. The vertical profile indicates the normalized

average and SEM of the input distributions across all recorded neurons. *iii*: Same as in ii but aligned on the soma

location. Dots indicate pia locations. *iv*: Normalized input magnitude deconvolved with the average morphology.

Dotted line indicates soma location. *v*: Box plot showing total input charge recorded during full-field stimulation. **B.**

Same as in A but for Cre-off Chronos injections into V2m.

Local input from V2m

To estimate the distribution of local input we used a Cre-off viral strategy, limiting Chronos expression to

non-Colgalt2-Cre neurons (n = 23 cells from 4 animals, average soma depth 482 ± 12 μm; Figure 4B).

When testing this strategy using the much denser Rbp4-Cre line, we found that only a very small

proportion of Cre-positive cells expressed Chronos (3%, Figure 4—figure supplement 2). The peak input

was located close to the soma, at 427 μm from the pia. The oblique compartment received the majority

(62%) of this input, with the basal dendrites and apical tuft receiving 23% and 16%, respectively, of the

total input (Figure 4—supplementary table 1). For the majority of recorded neurons, the peak input

occurred perisomatically (n = 22 / 23; Figure 4—figure supplement 3A). There was significant correlation

between soma depth and perisomatic input location, suggesting compartment-specific targeting of

individual ttL5 neurons rather than generic targeting of the cortical layer (Figure 4—figure supplement

3A). The horizontal input distribution showed a very slight lateral bias (-21 μm for all peaks; Figure 4—

figure supplement 3B). The total synaptic charge triggered by full-field stimulation was 12.61 ± 1.17 pC

(Figure 4B). Local neurons thus provide large direct input to ttL5 neurons in V2m, primarily targeting the

oblique (7.78 pC) compartment, with smaller input arriving to the basal (2.85 pC) and tuft (1.98 pC)

compartments.

Granular retrosplenial area

Next, we recorded optically evoked synaptic responses arising from RSPg axons (n = 26 cells from 9

animals, average soma depth 491 ± 13 μm; Figure 5A). The overall input displayed a bimodal distribution

peaking at 135 and 510 μm from the pia. The apical tuft received 25% of the input, with the oblique

compartment receiving 42% and basal dendrites 32% of the total input (Figure 4—supplementary table

1). For the majority of recorded neurons, the peak input targeted the perisomatic dendrites (n = 24 / 26;

Figure 5—figure supplement 1A). While there was no correlation between soma depth and tuft input, the

perisomatic input location followed the soma location to a significant level, suggesting that RSPg axons

target ttL5 dendrites at a single-cell level with specificity going beyond the general cortical layer (Figure

5—figure supplement 1A). The horizontal input distribution showed no lateral bias (Figure 5—figure

supplement 1B). Total synaptic charge triggered by full-field stimulation was 3.56 ± 0.58 pC (Figure 5A).

RSPg thus provides a relatively moderate direct input to ttL5 neurons in V2m, primarily targeting the

oblique (1.5 pC) and basal (1.15 pC) compartments with smaller input arriving to the more distal part of

the apical tuft (0.9 pC).

Anterior cingulate area

Next, we recorded optically evoked synaptic responses arising from ACA axons (n = 27 cells from 5

animals, average soma depth 461 ± 8 μm; Figure 5B). The overall input was bimodal, peaking at 94 μm

and 448 μm from the pia. The apical tuft received 27% of the input, with the oblique compartment

receiving 43% and basal dendrites 30% of the total input (Figure 4—supplementary table 1). The majority

of recorded neurons had the peak input located perisomatically (n = 25 / 27; Figure 5-figure supplement

1A). There was no correlation between soma depth and tuft input, while the perisomatic input location

followed the soma to a significant degree, suggesting single-cell sublayer specificity of ACA axons (Figure

5—figure supplement 2A). The horizontal input distribution showed a slight lateral bias (tuft 21 μm,

oblique -21 μm; Figure 5—figure supplement 2B). The total synaptic charge triggered by full-field

stimulation was 8.75 ± 1.18 pC (Figure 5B). ACA thus provides a large direct input to ttL5 neurons in

V2m, primarily targeting the oblique (3.77 pC) compartment with smaller input arriving to the basal (2.64

pC) and most distal part of the apical tuft (2.34 pC).

Figure 5. Subcellular connectivity maps of cortical FB areas. A. *i:* confocal image of a representative brain slice (blue = DAPI) showing the injection site in RSPg (green). *ii:* pia-aligned average sCRACM heatmap for RSPg inputs. Triangles represent soma locations. The vertical profile indicates the normalized average and SEM of the input distributions across all recorded neurons. *iii:* Same as in *ii* but aligned on the soma location. Dots indicate pia locations. *iv:* Normalized input magnitude deconvolved with the average morphology. Dotted line indicates soma location. *v:* Box plot showing total input charge recorded during full-field stimulation. **B.** Same as in A but for Chronos injections into ACA. **C.** Same as in A but for Chronos injections into ORB.

Orbitofrontal cortex

Optically evoked synaptic responses arising from ORB axons (n = 18 cells from 3 animals, average soma depth 502 ± 18 µm; Figure 5C) showed a strong perisomatic bias, with a peak at 510 µm from the pia. The apical tuft received only 8% of all input, with the oblique compartment receiving 55% and basal dendrites 37% of the total input (Figure 4—supplementary table 1). This distribution was also highly homogeneous across neurons, with almost all recorded neurons having their peak input in the perisomatic region (n = 17/18; Figure 5—figure supplement 3A). Interestingly, the input location of individual cells followed their soma depth significantly, suggesting ORB axons are targeting specific dendrites of individual ttL5 neurons rather than the cortical layer in general (Figure 5—figure supplement

240   3A). The horizontal input distribution showed no lateral bias (Figure 5—figure supplement 3B). The total

241   synaptic charge triggered by full-field stimulation was 7.11 ± 1.36 pC (Figure 5C). ORB thus provides a

242   large direct input to ttL5 neurons in V2m, primarily targeting the oblique (3.94 pC) and basal (2.63 pC)

243   compartments, with slight input arriving to the proximal part of the apical tuft (0.55 pC).

Anterior thalamic nuclei

Finally, we recorded optically evoked synaptic responses from thalamic axons, starting with the ATN (n =

16 cells from 5 animals, average soma depth 458 ± 11 µm; Figure 6A). This input had peaks at both 115

µm and 385 µm from the pia, with the apical tuft receiving the majority (68%) of the input, while the

oblique compartment received 25% and basal dendrites a mere 7% of the total input (Figure 4—

supplementary table 1). The majority of recorded neurons had the peak input in the tuft compartment (n =

9 / 16) and while all cells had some tuft input, in 7/16 cells the input peak was located perisomatically

(Figure 6—figure supplement 1A). There was no correlation between soma depth and either tuft or

perisomatic input location for individual recordings, suggesting ATN axons are targeting cortical layers

rather than specific ttL5 dendrites directly (Figure 6—figure supplement 1A). The horizontal input

distribution showed a slight lateral bias (21 µm for tuft and -21 µm for obliques and basal peaks; Figure

6—figure supplement 1B). The total synaptic charge triggered by full-field stimulation was 2.55 ± 0.3 pC

(Figure 6A). ATN thus provides a moderate direct input to ttL5 neurons in V2m, primarily targeting the

more distal part of the apical tuft (1.73 pC) while the oblique (0.65 pC) and the basal (0.17 pC)

compartment received less input.

Figure 6. Subcellular connectivity maps of thalamic input areas. A. *i:* confocal image of a representative brain

slice (blue = DAPI) showing the injection site in ATN (green). *ii:* pia-aligned average sCRACM heatmap for ATN

inputs. Triangles represent soma locations. The vertical profile indicates the normalized average and SEM of the

input distributions across all recorded neurons. *iii:* Same as in *ii* but aligned on the soma location. Dots indicate pia

locations. *iv:* Normalized input magnitude deconvolved with the average morphology. Dotted line indicates soma

location. *v:* Box plot showing total input charge recorded during full-field stimulation. **B.** Same as in A but for Chronos

injections in LP.

Lateral posterior nucleus of the thalamus

Lastly, we recorded optically evoked synaptic responses arising from LP axons (n = 13 cells from 5

animals, average soma depth 502 ± 19 µm; Figure 6B). Due to excessive retrograde labelling resulting in

direct photocurrent in the recorded V2m cells, a 1:10 dilution of virus was used for these injections. The

absolute value of the evoked input is thus likely an underestimate. As with ATN axons, the LP input was

strongly biased towards the most superficial part of the cortex and peaked at 73 µm from the pia. The

apical tuft received the vast majority (68%) of the input, with the oblique compartment receiving 18% and

basal dendrites 13 % of the total input (Figure 4—supplementary table 1). The majority of recorded

neurons had the peak input in the tuft compartment (n = 11/13; Figure 6—figure supplement 2A). There

was no correlation between soma depth and either tuft or perisomatic input location for individual

recordings, suggesting LP axons are targeting cortical layers rather than specific ttL5 dendrites directly

(Figure 6—figure supplement 2A). The horizontal input distribution showed strong lateral bias (-83 µm for

tuft and 63 µm for the oblique peak; Figure 6—figure supplement 2B). The total synaptic charge triggered

by full-field stimulation was 0.97 ± 0.15 pC (Figure 6B). LP thus provides small direct input to ttL5 neurons

in V2m, primarily targeting the most distal part of the apical tuft (0.67 pC) compartment with smaller input

arriving to the oblique (0.18 pC) and the basal (0.13 pC) compartments.

Comparison of anatomical and functional connectivity maps

To accurately estimate morphological overlap between axons and dendrites, we convolved the axonal

projection maps with the average dendritic morphology, resulting in the predicted input distribution one

would expect to see based on Peters' rule. When overlaying this with the pia-aligned vertical sCRACM

maps, the alignments between functional synapses and the axo-dendritic maps were diverse (Figure 7A).

For some regions, like ORB perisomatic and LP tuft inputs, a clear correspondence could be seen

between predicted and measured input distributions. A lesser degree of overlap can also be seen in the

VISp perisomatic or ACA tuft inputs. For other inputs, however, strong functional input could be detected

where there is very little overlap between dendrites and axons, such as at VISp tuft inputs. This stood in

stark contrast to the ORB projection, for which the opposite was true, and apical regions of dense

morphological overlap of axons and dendrites resulted in no functional input.

Next we examined the correspondence between the anatomical input connectivity results obtained from

rabies tracing and the functional connectivity results obtained by sCRACM. The proportion of rabies-

labelled inputs (only taking into account the 7 areas investigated by sCRACM) showed a strong

contribution from RSPg and modest input cell proportions for the more distal cortical regions (e.g. ACA,

ORB). The proportional sCRACM input calculated from the full-field stimuli however shows an altogether

different and effectively anti-correlated picture (Figure 7B), with modest input from RSPg and most input

arriving from V2m, ACA and ORB.

Figure 7. Comparing different input maps. A. Axonal density distribution convolved with dendritic morphology

(dotted black lines) overlaid with pia-aligned synaptic input distribution (coloured lines). Six input areas are shown.

Triangles represent soma location in average morphology. **B.** Proportion of input cells across 7 areas established by

rabies tracing (black) and proportion of input magnitude established by sCRACM (red). **C.** Excitatory synaptic input

map to the three different dendritic domains of ttL5 pyramidal neurons. The height of ellipses represents dendritic

location while width represents input strength. The tuft input map was generated from pia aligned maps while oblique

and basal maps from soma (dashed line) aligned maps. Area names are ordered by relative peak location; asterisks

mark FF areas. Only inputs > 0.6 pC are shown for clarity. NB: LP input magnitude is likely underestimated due to

lower virus titer.

Taken together, these results show a remarkable specificity of synaptic targeting by brain-wide

connections, with only a loose adherence to Peters' rule for the majority of inputs. This highlights that the

location of many long-range inputs cannot be surmised on the basis of axonal branching and may not

follow clear rules associated with their position above or below the target area in a cortical hierarchy.

Furthermore, the discrepancy between the numbers of rabies-labelled neurons and measured synaptic

currents indicates that the former should be taken as a guide for anatomical connectivity and identity, but

not the postsynaptic effects, of these inputs.

319   Discussion

Using an array of techniques for long-range circuit dissection, we have comprehensively mapped the

location and dendritic targets of inputs to ttL5 neurons in the medial secondary visual cortex in mice. This

allowed us to determine the dendritic targets of FF and FB connections and to make a direct assessment

of Peters' rule for brain-wide connections.

The whole-brain input map generated via rabies tracing was qualitatively similar to previous results from

the primary visual cortex (Kim et al 2015). Axonal projections from the rabies-identified input regions

broadly followed the expected pattern, with FB projections being biased toward L1 and FF areas toward

the deeper layers (D'Souza et al 2020, Harris et al 2019, Rockland & Pandya 1979). Accordingly, L1 was

densely innervated by higher-order cortical areas like RSPg and ACA as well as the secondary thalamic

nuclei (LP, ATN). Interestingly, ORB, a higher-order cortical region, displayed a projection pattern

associated with FF areas. Additionally, the majority of ORB projection neurons were found in L2/3,

another feature of FF connectivity. ORB is thus an exemption in terms of FB axonal projections.

In comparison to the strong similarities in axonal projections patterns, synaptic input maps showed a

remarkable degree of heterogeneity. We used a novel approach to allocate sCRACM input to specific

dendritic compartments by deconvolving the synaptic input maps with morphological ttL5 reconstructions,

in which tuft, oblique, and basal compartments had been manually labelled. This revealed several

unexpected results. The connection from VISp, which is a textbook example of FF connections and is

thus generally assumed to primarily target perisomatic dendrites, was instead heavily biased towards the

apical tuft. Conversely, the axons from ORB, which had a very similar, FF-like projection pattern,

synapsed almost exclusively with basal and oblique dendrites. Furthermore, the location of synaptic input

to individual neurons showed a strong correlation with soma depth, indicating that beyond the large

overlap of axons and dendrites, ORB input targets each cell so as to preferentially form synapses at

locations proximal to the soma. Other recorded areas (e.g. RSPg, ACA, ATN) showed some degree of

conformity to Peters' rule, yet still with significant differences in the proportion of synapses generated in

the different regions of high axonal projections. The only area where the correspondence was remarkable

and Peter's rule held fully was LP.

Comparing anatomical connectivity obtained by rabies tracing to functional connectivity obtained by full-

field optogenetic stimulation of axons revealed large and unexpected differences. However, several

technical caveats potentially biasing these results needs to be taken into account. First, the total input
proportion to a population of starter neurons does not necessarily represent faithfully the input proportions
received by single neurons, due to varying degrees of convergence from different input areas. We
attempted to minimize this effect by using a sparse Cre driver line to keep the number of starter cells low.
Second, the magnitude of optogenetically evoked input depends on the number of presynaptic cells
covered by viral injection. To mitigate this, we sought to cover a reasonably large fraction of each region
containing presynaptic cells by making several injections. Third, L5PNs can also be infected retrogradely
via their axonal projections to input areas, albeit with much lower probability. Recorded neurons with
direct photocurrent can be readily excluded based on the latency of sCRACM currents. However, likely
due to the dense reciprocal connectivity, LP injections resulted in a substantial number of retrogradely
labelled cells in V2m. This in turn could distort the LP synaptic input map by giving rise to unwanted local
input. To avoid this direct retrograde labelling, we injected a lower titer of virus into LP, which may have
caused an underestimation of LP input magnitude.

Despite these technical caveats, the detected discrepancy between anatomical and functional input
proportions estimated from the rabies tracing and sCRACM experiments is remarkable. There are several
possible explanations for this. First, there may be differing convergence of connectivity. For example,
inputs with strong rabies labelling but relatively weaker sCRACM input (like VISp and RSPg) may have
lower convergence, with a unique mapping of selected pre- and postsynaptic cells, perhaps carrying more
specific and detail-rich information to individual V2m neurons. Meanwhile, areas with strong synaptic
currents relative to their small rabies-labelled populations (like ORB and ACA) may have higher
convergence and be less discerning of their targets in order to convey more general contextual or state-
specific information. Second, there may be a difference in the strength of individual synapses not
reflected in rabies efficiency, with sparse input areas having disproportionately strong synapses.
However, the increase in the strengths of individual synapses required to overcome such a large
discrepancy seems rather unlikely. A third possible contributing factor could be the recently reported
activity-dependence of rabies transmission (Beier et al 2017). This would suggest that sparse input areas
(like ORB and ACA) have much lower activity despite their strong synaptic connections. Conversely, VISp
and RSPg should provide high activity input for the relatively modest synaptic input to results in extensive
rabies labelling.

Our results provide a complex picture regarding the possible interaction of FF and FB inputs (Figure 7C).
Functionally, V2m has been linked to visual motion processing (Sun et al 2009) and is thought to take
part in navigation and spatial processing as part of the dorsal stream (Glickfeld & Olsen 2017). Thalamic
FB input, which targets almost exclusively the apical tuft, arrives from multiple higher-order nuclei. Parts
of ATN receive strong vestibular input (Rancz et al 2015) and, together with RSP form a central part of
the head-direction system (Taube 2007, Velez-Fort et al 2018). It is thus likely that spatial and
multisensory contextual information carried by LP inputs (Roth et al 2016) interacts with FF visual input in
the tuft compartment. However, the role of tuft integration is likely to differ from primary sensory cortices,

considering that ttL5 neurons in the secondary visual cortex can have substantially different integrative

properties (Galloni et al 2020). Contrary to thalamic input, cortical FB inputs target all three dendritic

domains. Perhaps surprisingly, the strongest of these are ORB and ACA, which can interact with the local

FF input at the level of oblique and basal dendrites. Frontal cortices in general are involved in decision

making and executive control of behavior (Hamilton & Brigman 2015), and ORB in particular has been

shown to encode spatial goals (Feierstein et al 2006). ACA, meanwhile, can directly regulate visually

evoked responses and sensory discrimination in the primary visual cortex (Zhang et al 2014), and

contributes to learning to predict sensory input to primary visual cortex (Fiser et al 2016). Their precise

roles in the functioning of V2m, however, remains unknown.

A surprisingly large proportion (around half) of all the synaptic input, arrives at the oblique dendrites. This

is important as oblique dendrites are known to strongly affect L5 excitability (Schaefer et al 2003) and can

gate information flow from the apical dendrites (Jarsky et al 2005). While FB inputs targeting the apical

tuft have been suggested to act as a general gain control for ttL5 neurons, it is thus possible that ORB

and ACA perform a similar gating function, but perhaps with more specificity regarding input identity or

dendritic branches. Indeed, their particularly strong targeting of oblique dendrites might allow both fine-

level control of plasticity for FF synapses at these dendrites, while simultaneously allowing them to exert a

more binary gating control over input from both the thalamic nuclei and VISp, which all predominantly

arrive at the tuft. Furthermore, the large amount of cortical FB input to basal dendrites suggests a strong

amplification of local FF input close to the AP initiation zone.

In general, our results show that while the classification of areas as FF or FB can be based on axonal

projections (albeit with exceptions, such as ORB), macroscopic projectomes do not predict cell-type level

input location. We found that FF and FB inputs synapse on dendritic domains not purely determined by

projection and dendritic density, indicating that Peters' rule on the level of cell types is at best a weak

guideline. Similarly, while rabies tracing from a population of starter cells is an effective tool to study the

general wiring diagram, the proportion of input neurons thus estimated is likely to give a poor prediction of

functional input strength. Finally, the location and possible interactions between FF and the broad range

of FB inputs as well as their specific information content suggests that ttL5 neurons may be adopting a

multitude of integrative strategies that are different from those previously suggested.

Materials and Methods

Animals

All animal experiments were prospectively approved by the local ethics panel of the Francis Crick Institute (previously National Institute for Medical Research) and the UK Home Office under the Animals (Scientific Procedures) Act 1986 (PPL: 70/8935). All surgery was performed under isoflurane anesthesia, and every effort was made to minimize suffering. Transgenic mice were used: Tg(Colgalt2-Cre)NF107Gsat (RRID:MMRRC_036504-UCD, also known as Glt25d2-Cre) were crossed with the Ai14 reported line expressing tdTomato (RRID:IMSR_JAX:007908). Additionally, Tg(Rbp4-Cre)KL100Gsat/Mmucd (RRID:MMRRC_031125-UCD) mice were used to establish the efficacy of the cre-off approach. As only male mice are transgenic in the Colgalt2-Cre line, all experiments were done on male animals. Animals were housed in individually ventilated cages under a 12 hr light/dark cycle.

Viruses

EnvA-CVS-N2c$^{\Delta G}$-mCherry rabies virus, and adeno associated viruses expressing TVA and EGFP (AAV8-EF1a-flex-GT), N2c glycoprotein (AAV1-Syn-flex-H2B-N2CG), or Cre-OFF Chronos-GFP (AAV1-EF1-CreOff-Chronos-GFP) were a generous gift of Molly Strom and Troy Margrie. Chronos-GFP (also called ShChR) expressing adeno associated virus (rAAV1-Syn-Chronos-GFP) was obtained from UNC Vector Core.

Surgical procedures

Surgeries were performed on mice aged 3–8 weeks using aseptic technique under isoflurane (2–4%) anesthesia and analgesia (meloxicam 2 mg/kg and buprenorphine 0.1 mg/kg). The animals were head-fixed in a stereotaxic frame and a small hole (0.5–0.7 mm) was drilled in the skull above the injection site. Virus was loaded into a glass microinjection pipette (pulled to a tip diameter of around 20 µm) and pressure injected into the target region at a rate of 0.4 nl/s using a Nanoject III delivery system (Drummond Scientific). To reduce backflow, the pipette was left in the brain for approximately 5 min after completion of each injection.

For rabies virus tracing experiments, a 1:2 mixture of TVA and N2c glycoprotein expressing cre-dependent AAVs (10-20 nL) was injected at stereotaxic brain coordinates (λ - 0.8 mm, ML 1.6 mm, DV 0.6 mm). Rabies virus (50-100 nL) was injected 5-7 days later at the same site. Ten to twelve days later, animals were transcardially perfused under terminal anesthesia with cold phosphate-buffer (PB, 0.1 M) followed by 4% paraformaldehyde (PFA) in PB (0.1 M).

For the sCRACM experiments, Chronos-GFP expressing AAV was injected into one of the identified presynaptic regions. The virus was allowed to express for at least 3 weeks before acute brain slice

preparation. The range of stereotaxic coordinates for each region are listed in Figure 2--supplementary
table 1. The injected virus was Chronos-GFP for every region except V2m, where Cre-OFF Chronos-GFP
was instead used to avoid expression in the recorded Colgalt2-Cre neurons. For some of the injections in
LP, the Chronos-GFP virus was diluted by 10-fold in sterile cortex buffer before injection.

Data acquisition and analysis for rabies tracing experiments

Brain samples were embedded in 4-5% agarose (Sigma-Aldrich: 9012-36-6) in 0.1M PB and imaged
using serial two-photon tomography (Han et al 2018, Osten & Margrie 2013, Ragan et al 2012). Eight
optical sections were imaged every 5 μm with 1.2 μm x 1.2 μm lateral resolution, after which a 40μm
physical section was removed. Excitation was provided by a pulsed femto-second laser at 800 nm
wavelength (MaiTai eHP, Spectraphysics). Images were acquired through a 16X, 0.8 NA objective (Nikon
MRP07220) in three channels (green, red, blue) using photomultiplier tubes. Image tiles for each channel
and optical plane were stitched together with an open-source software written in MATLAB
(https://github.com/SainsburyWellcomeCentre/StitchIt). For cell detection, full resolution images were first
filtered with a Gaussian blur (sigma = 1) using Fiji (ImageJ 1.52e) to reduce imaging noise. The open-
source package cellfinder (DOI: 10.5281/zenodo.3891338; (Tyson et al 2020) was used for cell candidate
detection then classification. Automated mouse atlas propagation (Niedworok et al 2016) was used for
registration and segmentation on brain samples down-sampled to 10 μm voxels (to match the resolution
of the Allen CCFv3; (Wang et al 2020a). Cell coordinates were similarly down-sampled to 10 μm and the
number of cells was counted for each segmented area. For cell density visualisation, cell coordinates
were reverse-transformed onto the Allen CCFv3 space using the open source registration tool, Elastix
(Klein et al 2010) and were projected onto a 2D matrix in 10 μm / pixel resolution. For each unique cell
position, a cell density value was calculated as total cell counts in a nearby 9x9 pixel grid, normalized to
the sum of all starter or presynaptic cells. Data is reported as the percentage of all presynaptic cells
averaged across 3 experiments. Starter cell numbers were 7, 15, and 36.

Acute slice preparation and electrophysiological recordings

Adult mice were deeply anaesthetised with isoflurane and decapitated. The brain was rapidly removed
and placed in oxygenated ice-cold slicing solution containing (in mM): 125 sucrose, 62.5 NaCl, 2.5 KCl,
1.25 NaH_2PO_4, 26 $NaHCO_3$, 2 $MgCl_2$, 1 $CaCl_2$, 25 dextrose; osmolarity 340–350 mOsm. The cerebellum
and frontal cortex were removed manually with a coronal cut using a single-edged razor blade and the
rostral surface was affixed to a metal platform with cyanoacrylate glue. Coronal slices (300 μm thick)
between 2.6 and 3.5 mm posterior to bregma were prepared using a vibrating blade microtome (Leica
VT1200S). Slices were kept submerged in artificial cerebrospinal fluid (ACSF, containing in mM: 125
NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 26 $NaHCO_3$, 1 $MgCl_2$, 2 $CaCl_2$, 25 dextrose; osmolarity 308–312 mOsm) at
35°C for the first 30–60 min after slicing, then at room temperature (22°C). All solutions and chambers
were continuously bubbled with carbogen (95% O2 / 5% CO2).

For recordings, individual slices were perfused in the recording chamber at a rate of approximately 6

mL/min with ACSF at room temperature (22°C), continuously bubbled with carbogen. To prevent axonal

spike propagation and enhance responses to optical stimulation, 1 μM tetrodotoxin (TTX) and 100 μM 4-

aminopyridin (4-AP) were added to the recording ACSF. This ensured that any light-evoked responses

were direct monosynaptic responses resulting from stimulation of Chronos-expressing axon terminals,

rather than from passing axons terminating in unknown locations on the dendrites.

Filamented borosilicate thick-walled glass micropipettes were pulled and heat-polished using a two-stage

horizontal puller (Zeitz DMZ Universal Electrode Puller) to obtain an electrode resistance of 3–6 MΩ. The

glass electrodes were filled with internal solution optimized for voltage clamp recordings, containing (in

mM): 120 CsMeSO$_3$ (CH$_3$O$_3$SCs), 3 CsCl, 10 HEPES, 1 EGTA, 4 Na$_2$ATP, 0.3 NaGTP, 5 Na$_2$-

phosphoreatine (C$_4$H$_8$N$_3$O$_5$PNa$_2$), 3.5 QX-314 chloride, 0.5 % (w/v) biocytin hydrochloride, 50 μM Alexa

Fluor 488 hydrazide; osmolarity 290–295 mOsm; pH adjusted to 7.3 with CsOH.

Visually guided whole-cell patch-clamp recordings from tdTomato-labelled Colgalt2-Cre neurons in V2m

were performed using a Scientifica SliceScope Pro 3000 microscope equipped with a 40x/0.8 NA

objective and an infrared (IR) Dodt Gradient Contrast system. The epifluorescence system used to

visualize fluorescent neurons was a CoolLED pE-4000 illumination system with a 550 nm peak excitation

wavelength. To avoid stimulating Chronos-expressing axons, epifluorescent illumination was kept to a

minimum during selection of cells to record. Recordings were made with a Multiclamp 700B amplifier

(Molecular Devices) in voltage-clamp configuration with a holding potential of -70 mV. Filtered signals

(8kHz low-pass) were digitized at 20 kHz with a National Instruments DAQ board (PCIe-6323). Acquisition

and stimulus protocols were generated in Igor Pro (Wavemetrics) with the NeuroMatic software package

(Rothman & Silver 2018). Throughout each recording, series resistance compensation was applied and

set to the highest value possible without inducing oscillations in the cell (typically between 40 and 75%).

Recordings wtih series resistance larger than 40 MΩ were excluded.

Patterned optogenetic stimulation

Optical stimulation was implemented using a digital micromirror device (DMD) with a 463 nm laser (laser-

coupled Polygon 400, Mightex Systems). The stimulus consisted of a 1000 x 500 μm grid divided into 24

x 12 spots of light (41.7 μm x 41.7 μm square) delivered through a 5x/0.15 NA dry objective (Olympus

MPlanFL N). The grid was approximately centered on the soma being recorded from, aligned to the pia

orthogonal to the apical dendrite. For each individual spot, the laser power was measured at the

specimen plane using a PM100D (Thorlabs) optical power meter equipped with a S121C sensor

(Thorlabs). The laser output associated with each spot was adjusted to obtain a measured power of

approximately 300 μW (173 mW/mm2).

Optical stimuli were delivered for 1 ms at 0.1 Hz in a pseudo-random sequence designed to maximise the

distance between consecutive spots and the time between stimulation of neighbouring spots. Each

recording trial consisted of a single repetition of all 288 stimuli followed by a full-field stimulus, in which all

stimulation spots were illuminated simultaneously for 1 ms. For each cell, 5-20 trials were recorded, with

30s pauses between trials, making the interval between consecutive stimulation of the same spot

approximately 60s. Following each recording, an image was taken to record the location of the recorded

cell (filled with Alexa Fluor 488) relative to the stimulation grid. This was used during analysis to align the

recorded sCRACM heatmap with the location of the pia or soma.

Immunohistochemistry & morphological reconstructions

After recording, slices were fixed overnight at 4°C in a 4% paraformaldehyde solution and were

subsequently kept in PBS. Slices were stained with DAPI (5 µg/mL) for 10 min, mounted on glass slides

and images were acquired with either a confocal microscope for high-resolution images (Leica SP5;

objective: 20x/0.7NA or 10x/0.4NA; pinhole size: 1 airy unit) or a slide scanner for visualizing injection

sites (Olympus VS120, objective: 4x/0.16NA). Image processing was done with the FIJI software package

(Schindelin et al 2012).

Data analysis

Analysis and data visualization were performed with custom macros and scripts written in Igor Pro and

MATLAB (Mathworks). Unless otherwise specified, all reported data values refer to the mean ± standard

error (SEM). Recordings were not corrected for liquid junction potential.

Recordings were baselined in a 40 ms window before each stimulus and averaged across trials, and the

peak and area of the evoked currents were measured in a 50 ms window after the stimulus. Stimulus

spots for which the peak current was lower than seven times the standard deviation of the baseline noise

were scored as zero.

For some of the injection sites, a degree of retrograde transport of the Chronos virus was noted in areas

outside of the primary injection site, including in V2m. A few recorded Colgalt2-Cre neurons in V2m were

thus found to be intrinsically expressing Chronos. This was easily detected in the recordings by an

instantaneous inward current at the onset of laser stimulation, in contrast to the 4-5 ms delay between

stimulus onset and sCRACM current observed normally. Any cells with no such delay were excluded from

the analysis.

Because peak response amplitudes varied between cells and preparations, to obtain average input

distributions from a presynaptic population, the heatmap for each cell was normalised to the peak EPSC

value for that cell. Heatmaps were then aligned horizontally by soma location and then vertically by either

soma or pia location before averaging each pixel across cells. For each cell, the soma could be localised

to within one quadrant of a given stimulation spot. When averaging across cells, the effective sampling

resolution (i.e. the pixel dimension) for the average sCRACM heatmaps and related horizontal and

vertical projections were thus approximately 20.8 µm (equal to half of the stimulus spot size, i.e. 1000/48

µm). All values reported for the locations of sCRACM inputs from different presynaptic regions are in
multiples of this number. Note, however, that the actual resolution with which synapses can be localised
in the heatmaps is likely to be lower than this, as it is limited by both light scattering in the tissue and by
the spread of voltage along stimulated axons, which is determined by the length constant of presynaptic
axons. Previous studies have indicated that these factors limit the actual sCRACM resolution to
approximately 60 µm (Petreanu et al 2009).

To estimate the proportion of sCRACM input targeting different dendritic domains, the recorded input map
for each cell was convolved with the average ttL5 morphology obtained from 11 reconstructed Colgalt2-
Cre neurons in V2m. This was done by manually separating the apical tuft, oblique (including the apical
trunk), and basal dendrites of the reconstructions in Neurolucida 360 and quantifying the total dendritic
length in 10 µm thick sections perpendicular to the main axis of the apical dendrite. The resulting dendrite
profiles were then aligned by the soma and averaged. Using this average morphology, at each distance
from the soma the proportion of dendrites belonging to each domain was calculated relative to the total
dendritic length within that section. For each sCRACM recording, these proportions were calculated using
an average morphology that was scaled to the soma-pia distance of the recorded cell. Each pixel of the
heatmap for that cell was then multiplied by these dendritic proportions to obtain the proportion of evoked
current assigned to each dendritic domain.

A similar calculation was performed to quantify the expected input profile on the basis of axon and
dendrite densities at each distance from the pia. In this case, the reconstructed morphologies were
aligned by the apical tuft before averaging. Both the axon and dendrite distributions were then normalised
to the peak of each curve and multiplied, resulting in large values at locations containing both axons and
dendrites.

573 Acknowledgments

574 We thank Troy Margrie, Molly Strom, Thomas Mrsic-Flogel and Charlie Rousseau for help with data
575 acquisition and analysis of rabies tracing experiments. We are grateful to Florencia Iacaruso and Zoltán
576 Kisvárday for helpful comments on the manuscript.

577 **Competing interests:** The authors declare that no competing interests exist.

References

Aru J, Suzuki M, Larkum ME. 2020. Cellular Mechanisms of Conscious Processing. *Trends Cogn Sci* 24: 814-25

Beier KT, Kim CK, Hoerbelt P, Hung LW, Heifets BD, et al. 2017. Rabies screen reveals GPe control of cocaine-triggered plasticity. *Nature* 549: 345-50

Briggman KL, Helmstaedter M, Denk W. 2011. Wiring specificity in the direction-selectivity circuit of the retina. *Nature* 471: 183-8

Collins DP, Anastasiades PG, Marlin JJ, Carter AG. 2018. Reciprocal Circuits Linking the Prefrontal Cortex with Dorsal and Ventral Thalamic Nuclei. *Neuron* 98: 366-79 e4

D'Souza RD, Meier AM, Bista P, Wang Q, Burkhalter A. 2016. Recruitment of inhibition and excitation across mouse visual cortex depends on the hierarchy of interconnecting areas. *Elife* 5

D'Souza RD, Wang Q, Ji W, Meier AM, Kennedy H, et al. 2020. Canonical and noncanonical features of the mouse visual cortical hierarchy. *biorxiv.org*

Feierstein CE, Quirk MC, Uchida N, Sosulski DL, Mainen ZF. 2006. Representation of spatial goals in rat orbitofrontal cortex. *Neuron* 51: 495-507

Felleman DJ, Van Essen DC. 1991. Distributed hierarchical processing in the primate cerebral cortex. *Cereb Cortex* 1: 1-47

Fino E, Yuste R. 2011. Dense inhibitory connectivity in neocortex. *Neuron* 69: 1188-203

Fiser A, Mahringer D, Oyibo HK, Petersen AV, Leinweber M, Keller GB. 2016. Experience-dependent spatial expectations in mouse visual cortex. *Nat Neurosci* 19: 1658-64

Franklin KBJ, Paxinos G. 2007. *Paxinos and Franklin's The mouse brain in stereotaxic coordinates*. Amsterdam: Academic Press, an imprint of Elsevier. 1 volume (unpaged) pp.

Galloni AR, Laffere A, Rancz E. 2020. Apical length governs computational diversity of layer 5 pyramidal neurons. *Elife* 9

Glickfeld LL, Olsen SR. 2017. Higher-Order Areas of the Mouse Visual Cortex. *Annu Rev Vis Sci* 3: 251-73

Groh A, Meyer HS, Schmidt EF, Heintz N, Sakmann B, Krieger P. 2010. Cell-type specific properties of pyramidal neurons in neocortex underlying a layout that is modifiable depending on the cortical area. *Cereb Cortex* 20: 826-36

Guerguiev J, Lillicrap TP, Richards BA. 2017. Towards deep learning with segregated dendrites. *Elife* 6

Hamilton DA, Brigman JL. 2015. Behavioral flexibility in rats and mice: contributions of distinct frontocortical regions. *Genes Brain Behav* 14: 4-21

Han Y, Kebschull JM, Campbell RAA, Cowan D, Imhof F, et al. 2018. The logic of single-cell projections from visual cortex. *Nature* 556: 51-56

Harris JA, Mihalas S, Hirokawa KE, Whitesell JD, Choi H, et al. 2019. Hierarchical organization of cortical and thalamic connectivity. *Nature* 575: 195-202

Helmstaedter M, Briggman KL, Turaga SC, Jain V, Seung HS, Denk W. 2013. Connectomic reconstruction of the inner plexiform layer in the mouse retina. *Nature* 500: 168-74

Hooks BM, Mao T, Gutnisky DA, Yamawaki N, Svoboda K, Shepherd GM. 2013. Organization of cortical and thalamic input to pyramidal neurons in mouse motor cortex. *J Neurosci* 33: 748-60

Hubel DH, Wiesel TN. 1962. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. *J Physiol* 160: 106-54

Jarsky T, Roxin A, Kath WL, Spruston N. 2005. Conditional dendritic spike propagation following distal synaptic activation of hippocampal CA1 pyramidal neurons. *Nat Neurosci* 8: 1667-76

Kasthuri N, Hayworth KJ, Berger DR, Schalek RL, Conchello JA, et al. 2015. Saturated Reconstruction of a Volume of Neocortex. *Cell* 162: 648-61

Kim EJ, Juavinett AL, Kyubwa EM, Jacobs MW, Callaway EM. 2015. Three Types of Cortical Layer 5 Neurons That Differ in Brain-wide Connectivity and Function. *Neuron* 88: 1253-67

Kim JS, Greene MJ, Zlateski A, Lee K, Richardson M, et al. 2014. Space-time wiring specificity supports direction selectivity in the retina. *Nature* 509: 331-36

Klein S, Staring M, Murphy K, Viergever MA, Pluim JP. 2010. elastix: a toolbox for intensity-based medical image registration. *IEEE Trans Med Imaging* 29: 196-205

Krishnaswamy A, Yamagata M, Duan X, Hong YK, Sanes JR. 2015. Sidekick 2 directs formation of a retinal circuit that detects differential motion. *Nature* 524: 466-70

Larkum M. 2013. A cellular mechanism for cortical associations: an organizing principle for the cerebral
cortex. *Trends Neurosci* 36: 141-51
Larkum ME, Petro LS, Sachdev RNS, Muckli L. 2018. A Perspective on Cortical Layering and Layer-
Spanning Neuronal Elements. *Front Neuroanat* 12: 56
LeCun Y, Bengio Y, Hinton G. 2015. Deep learning. *Nature* 521: 436-44
Lee WC, Bonin V, Reed M, Graham BJ, Hood G, et al. 2016. Anatomy and function of an excitatory
network in the visual cortex. *Nature* 532: 370-4
Little JP, Carter AG. 2012. Subcellular synaptic connectivity of layer 2 pyramidal neurons in the medial
prefrontal cortex. *J Neurosci* 32: 12808-19
Niedworok CJ, Brown AP, Jorge Cardoso M, Osten P, Ourselin S, et al. 2016. aMAP is a validated
pipeline for registration and segmentation of high-resolution mouse brain data. *Nat Commun* 7:
11879
Osten P, Margrie TW. 2013. Mapping brain circuitry with a light microscope. *Nat Methods* 10: 515-23
Packer AM, McConnell DJ, Fino E, Yuste R. 2013. Axo-dendritic overlap and laminar projection can
explain interneuron connectivity to pyramidal cells. *Cereb Cortex* 23: 2790-802
Petreanu L, Mao T, Sternson SM, Svoboda K. 2009. The subcellular organization of neocortical excitatory
connections. *Nature* 457: 1142-5
Ragan T, Kadiri LR, Venkataraju KU, Bahlmann K, Sutin J, et al. 2012. Serial two-photon tomography for
automated ex vivo mouse brain imaging. *Nat Methods* 9: 255-8
Rancz EA, Moya J, Drawitsch F, Brichta AM, Canals S, Margrie TW. 2015. Widespread vestibular
activation of the rodent cortex. *J Neurosci* 35: 5926-34
Reardon TR, Murray AJ, Turi GF, Wirblich C, Croce KR, et al. 2016. Rabies Virus CVS-N2c(DeltaG)
Strain Enhances Retrograde Synaptic Transfer and Neuronal Viability. *Neuron* 89: 711-24
Rees CL, Moradi K, Ascoli GA. 2017. Weighing the Evidence in Peters' Rule: Does Neuronal Morphology
Predict Connectivity? *Trends Neurosci* 40: 63-71
Richards BA, Lillicrap TP, Beaudoin P, Bengio Y, Bogacz R, et al. 2019. A deep learning framework for
neuroscience. *Nat Neurosci* 22: 1761-70
Rieubland S, Roth A, Hausser M. 2014. Structured connectivity in cerebellar inhibitory networks. *Neuron*
81: 913-29
Rockland KS, Pandya DN. 1979. Laminar origins and terminations of cortical connections of the occipital
lobe in the rhesus monkey. *Brain Res* 179: 3-20
Roth MM, Dahmen JC, Muir DR, Imhof F, Martini FJ, Hofer SB. 2016. Thalamic nuclei convey diverse
contextual information to layer 1 of visual cortex. *Nat Neurosci* 19: 299-307
Rothman JS, Silver RA. 2018. NeuroMatic: An Integrated Open-Source Software Toolkit for Acquisition,
Analysis and Simulation of Electrophysiological Data. *Front Neuroinform* 12: 14
Schaefer AT, Larkum ME, Sakmann B, Roth A. 2003. Coincidence Detection in Pyramidal Neurons Is
Tuned by Their Dendritic Branching Pattern. *Journal of Neurophysiology* 89: 3143-54
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, et al. 2012. Fiji: an open-source platform
for biological-image analysis. *Nat Methods* 9: 676-82
Shipp S. 2016. Neural Elements for Predictive Coding. *Front Psychol* 7: 1792
Sun B, Lv B, Dong Q, Wang S, Chai Z. 2009. Watching moving images specifically promotes
development of medial area of secondary visual cortex in rat. *Dev Neurobiol* 69: 558-67
Takahashi N, Ebner C, Sigl-Glockner J, Moberg S, Nierwetberg S, Larkum ME. 2020. Active dendritic
currents gate descending cortical outputs in perception. *Nat Neurosci* 23: 1277-85
Takahashi N, Oertner TG, Hegemann P, Larkum ME. 2016. Active cortical dendrites modulate perception.
Science 354: 1587-90
Taube JS. 2007. The head direction signal: origins and sensory-motor integration. *Annu Rev Neurosci* 30:
181-207
Tyson AL, Rousseau CV, Niedworok CJ, Keshavarzi S, Tsitoura C, Margrie TW. 2020.
Velez-Fort M, Bracey EF, Keshavarzi S, Rousseau CV, Cossell L, et al. 2018. A Circuit for Integration of
Head- and Visual-Motion Signals in Layer 6 of Mouse Primary Visual Cortex. *Neuron* 98: 179-91
e6
Wang Q, Ding SL, Li Y, Royall J, Feng D, et al. 2020a. The Allen Mouse Brain Common Coordinate
Framework: A 3D Reference Atlas. *Cell* 181: 936-53 e20
Wang XJ, Pereira U, Rosa MG, Kennedy H. 2020b. Brain connectomes come of age. *Curr Opin*
Neurobiol 65: 152-61

Wickersham IR, Lyon DC, Barnard RJ, Mori T, Finke S, et al. 2007. Monosynaptic restriction of
691   transsynaptic tracing from single, genetically targeted neurons. *Neuron* 53: 639-47
692   Yamawaki N, Li X, Lambot L, Ren LY, Radulovic J, Shepherd GMG. 2019. Long-range inhibitory
693   intersection of a retrosplenial thalamocortical circuit by apical tuft-targeting CA1 neurons. *Nat*
694   *Neurosci* 22: 618-26
695   Zeng H. 2018. Mesoscale connectomics. *Curr Opin Neurobiol* 50: 154-62
696   Zhang S, Xu M, Kamigaki T, Hoang Do JP, Chang WC, et al. 2014. Selective attention. Long-range and
697   local circuits for top-down modulation of visual cortex processing. *Science* 345: 660-5

699 # Supplementary material

bregma - 0.94 mm

bregma - 2.46 mm

Cortex

1	Main olfactory bulb	MOB	20	Posterior parietal association areas	PTLp
2	Orbital area	ORB	20a	Rostrolateral visual area	VISrl
3	Prelimbic area	PL	20b	Anterior area	VISa
4	Anterior cingulate area	ACA	21	Postrhinal area	VISpor
5	Primary motor area	MOp	22	Lateral visual area	VISl
6	Secondary motor area	MOs	23	Posterolateral visual area	VISpl
7	Supplemental somatosensory area	SSs	24	Anterolateral visual area	VISal
8	Primary somatosensory area, nose	SSp-n	25	Ectorhinal area	ECT
9	Primary somatosensory area, mouth	SSp-m	26	Entorhinal area	ENT
10	Primary somatosensory area, barrel field	SSp-bfd	27	Presubiculum	PRE
11	Primary somatosensory area, lower limb	SSp-ll	28	Postsubiculum	POST
12	Primary somatosensory area, trunk	SSp-tr	29	Area prostriata	Apr
13	Primary somatosensory area, upper limb	SSp-ul	30	Primary visual area	VISp
14	Temporal association areas	TEa	31	Anteromedial visual area	VISam
15	Laterointermediate area	VISli	32	posteromedial visual area	VISpm
16	Primary auditory area	AUDp	33	Retrosplenial area, dorsal part	RSPd
17	Ventral auditory area	AUDv	34	Retrosplenial area, ventral part	RSPv
18	Dorsal auditory area	AUDd	35	Retrosplenial area, lateral agranular part	RSPagl
19	Posterior auditory area	AUDpo			

Thalamus

36	Lateral posterior nucleus of the thalamus	LP	39	Anteroventral nucleus of thalamus	AV
37	Dorsal part of the lateral geniculate complex	LGd	40	Anterodorsal nucleus	AD
38	Lateral dorsal nucleus of thalamus	LD	41	Anteromedial nucleus	AM

702 **Figure 1—figure supplement 1.** Brain parcellation and nomenclature according to CCFv3.

total input proportion (area / whole brain)								
n = 3 experiments	average	SD	range					
number of starter cells	19	15	7-36					
total input cells	11622	2235	9059-13161					
% contralateral input cells	6.30%	0.90%						

thalamus	average	sd	L1	L2/3	L4	L5	L6a	L6b
LP	10.30%	1.70%						
LD	4.90%	0.80%						
AM	2.20%	0.50%						
LGd	1.70%	1.40%						
proximal cortex	**53.50%**							
VISp	10.60%	3.20%	1.83%	39.36%	8.10%	42.97%	5.97%	1.77%
VISpm	6.60%	1.80%	0.28%	16.63%	14.62%	55.98%	9.68%	2.80%
VISam	3.40%	1.50%	0.00%	15.12%	10.75%	60.37%	12.43%	1.32%
VISl	2.90%	1.50%	0.17%	25.97%	20.55%	46.06%	5.92%	1.33%
VISal	1.00%	0.80%	0.00%	20.08%	23.55%	50.62%	5.36%	0.40%
RSPv	14.70%	4.70%	0.29%	11.40%	n/a	70.43%	17.87%	0.01%
RSPd	8.50%	1.00%	0.29%	9.97%	n/a	65.15%	24.45%	0.14%
RSPagl	5.80%	1.70%	1.14%	24.25%	n/a	64.46%	9.96%	0.19%
distal cortex	**7.90%**							
ORB	2.70%	1.10%	19.25%	72.86%	n/a	6.22%	1.67%	0.00%
ACA	1.90%	1.00%	0.96%	12.20%	n/a	73.28%	13.56%	0.00%
PTLp (VISrl+VISa)	1.90%	1.00%	0.00%	11.16%	10.62%	65.23%	12.60%	0.39%
AUD	1.40%	0.50%	1.58%	17.60%	10.83%	60.17%	8.18%	1.64%

704 **Figure 1—supplementary table 1.** Results of presynaptic input tracing experiments. Areas highlighted in red are
705 chosen for further study.

Area	Distance from bregma (mm)	Mediolateral distance (mm)	Depth from pia (mm)	Number & volume of injections
VISp	[−3.5 : −2.8]	[1.8 : 2.7]	[0.5 : 0.6]	3 x 100 nL
RSPg	[−3.2 : −2.7]	0.5	[0.5 : 0.7]	2 x 100 nL
V2m	[−3.2 : −2.7]	1	[0.2 : 0.5]	2 x 100 nL
ACA	[0 : 1]	0.5	[1.2 : 1.5]	3 x 100 nL
ORB	[2 : 2.8]	1	[1.5 : 2.3]	3 x 100 nL
ATN	[−0.5 : −1.2]	[0.5 : 0.7]	[3.2 : 3.3]	2 x 100 nL
LP	[−2.5 : −1.7]	[1 : 1.5]	[2.4 : 2.6]	3 x 100 nL

708
709
710 **Figure 2—supplementary table 1.** Stereotaxic coordinates and volumes of viral injections.

711
712
713 **Figure 3—figure supplement 1. Input location does not correlate with input strength. A.** Horizontal projections
714 of VISp input to individual cells sorted by full-field stimulation response. Right: Location of the largest input peak
715 versus full-field response. Dashed line is a linear fit. **B-G.** Same as in A, but for RSPg, V2m, ACA, ORB, ATN and LP
716 input.

input area	parameter	basal	oblique	tuft*		
VISp	peak location (µm)	-63	21	198	total input (pC)	1.46
N = 7	input proportion (%)	18%	25%	57%	total input SEM	0.33
n = 14	proportional input (pC)	0.26	0.36	**0.84**	soma depth (µm)	493 ± 18
	horizontal bias (µm) **	41.7	41.7	20.8	cells with peak in tuft	10/14
V2m	peak location (µm)	-42	42	177	total input (pC)	12.61
N = 4	input proportion (%)	22%	61%	17%	total input SEM	1.17
n = 23	proportional input (pC)	**2.76**	**7.71**	**2.14**	soma depth (µm)	482 ± 12
	horizontal bias (µm) **	20.8	20.8	20.8	cells with peak in tuft	1/23
RSPg	peak location (µm)	-42	63	135	total input (pC)	3.56
N = 9	input proportion (%)	32%	42%	25%	total input SEM	0.58
n = 26	proportional input (pC)	**1.16**	**1.51**	**0.90**	soma depth (µm)	491 ± 13
	horizontal bias (µm) **	0	0	0	cells with peak in tuft	2/26
ACA	peak location (µm)	-42	21	94	total input (pC)	8.75
N = 5	input proportion (%)	30%	43%	27%	total input SEM	1.18
n = 27	proportional input (pC)	**2.64**	**3.77**	**2.34**	soma depth (µm)	461 ± 8
	horizontal bias (µm) **	0	20.8	-20.8	cells with peak in tuft	2/27
ORB	peak location (µm)	-42	21	N/A	total input (pC)	7.11
N = 3	input proportion (%)	37%	55%	8%	total input SEM	1.36
n = 18	proportional input (pC)	**2.63**	**3.94**	0.55	soma depth (µm)	502 ± 18
	horizontal bias (µm) **	0	0	0	cells with peak in tuft	0/18
ATN	peak location (µm)	-21	63	115	total input (pC)	2.55
N = 5	input proportion (%)	7%	25%	68%	total input SEM	0.30
n = 16	proportional input (pC)	0.17	**0.65**	**1.73**	soma depth (µm)	458 ± 11
	horizontal bias (µm) **	20.8	20.8	20.8	cells with peak in tuft	9/16
LP	peak location (µm)	-42	21	73	total input (pC)	0.97
N = 5	input proportion (%)	13%	18%	68%	total input SEM	0.15
n = 13	proportional input (pC)	0.13	0.18	**0.67**	soma depth (µm)	502 ± 19
	horizontal bias (µm) **	-41.7	62.5	-83.3	cells with peak in tuft	11/13
	total proportional input (pC)	**9.75**	**18.12**	**9.16**	**sum total input (pC)**	**37.03**

*tuft from pia, rest from soma
** negative means lateral, positive medial

718
719   **Figure 4—supplementary table 1.** Results of all sCRACM experiments.

Figure 4—figure supplement 1. Detailed data analysis for VISp. A. Vertical projections of individual input maps sorted by the location of the peak input. Right: Distance of tuft (green, defined as above the apical trough in the vertical projection) and perisomatic (blue, defined as below the apical trough in the vertical projection) input peaks from pia versus soma depth. Dashed lines are linear fits. **B.** Horizontal projections of individual input maps and their average for all, and for domain-separated inputs.

Figure 4—figure supplement 2. Efficacy of Cre-off virus. A. Example confocal images showing cre expression in

L5 pyramidal neurons (red), GFP expression following cre-off virus injection (cyan) and overlap (blue = DAPI) on two

magnifications. **B.** Quantification of overlap between cre and GFP expression.

Figure 4—figure supplement 3. Detailed data analysis for V2m. A. Vertical projections of individual input maps
sorted by the location of the strongest input. *Right:* Distance of tuft (green) and perisomatic (blue) input peaks from
pia versus soma depth. Dashed lines are linear fits. **B.** Horizontal projections of individual input maps and their
average for all, and for domain-separated inputs.

Figure 5—figure supplement 1. Detailed data analysis for RSPg. A. Vertical projections of individual input maps sorted by the location of the strongest input. *Right:* Distance of tuft (green) and perisomatic (blue) input peaks from pia versus soma depth. Dashed lines are linear fits. **B.** Horizontal projections of individual input maps and their average for all, and for domain-separated inputs.

Figure 5—figure supplement 2. Detailed data analysis for ACA. A. Vertical projections of individual input maps sorted by the location of the strongest input. *Right:* Distance of tuft (green) and perisomatic (blue) input peaks from pia versus soma depth. Dashed lines are linear fits. **B.** Horizontal projections of individual input maps and their average for all, and for domain-separated inputs.

746 **Figure 5—figure supplement 3. Detailed data analysis for ORB. A.** Vertical projections of individual input maps

747 sorted by the location of the strongest input. *Right:* Distance of the perisomatic input peak from pia versus soma

748 depth. Dashed line is linear fit. **B.** Horizontal projections of individual input maps and their average for all, and for

749 domain-separated inputs.

Figure 6—figure supplement 1. Detailed data analysis for ATN. A. Vertical projections of individual input maps sorted by the location of the strongest input. *Right:* Distance of tuft (green) and perisomatic (blue) input peaks from pia versus soma depth. Dashed lines are linear fits. **B.** Horizontal projections of individual input maps and their average for all, and for domain-separated inputs.

Figure 6—figure supplement 2. Detailed data analysis for LP. A. Vertical projections of individual input maps

sorted by the location of the strongest input. *Right:* Distance of tuft (green) and perisomatic (blue) input peaks from

pia versus soma depth. Dashed lines are linear fits. **B.** Horizontal projections of individual input maps and their

average for all, and for domain-separated inputs.