β-importins Tnpo-SR and Cadmus and the small GTPase Ran promote ovarian cyst formation in *Drosophila*

Allison N. Beachum, Taylor D. Hinnant, Anna E. Williams, Amanda M. Powell, and Elizabeth T. Ables*

Department of Biology, East Carolina University, Greenville, NC 27858.

*Correspondence: ablese@ecu.edu; 1001 E. 10th St. Mailstop 551, Greenville, NC 27858; Phone: 252-328-9770; Fax: 252-328-4178

RUNNING TITLE
importins promote cyst formation

KEYWORDS
Oocyte, oogenesis, germline, cyst, germline stem cell

CORRESPONDING AUTHOR:
Elizabeth T. Ables
East Carolina University
Department of Biology
1001 E. 10th St., Mailstop 551
553 Science & Technology Building
Greenville, NC 27858
Office phone: 252-328-9770
e-mail: ablese@ecu.edu

SUMMARY STATEMENT

Ran and two β-importins function independently to promote oocyte selection and cyst

development in the *Drosophila* ovary, likely by regulating distinct cargos critical for cell cycle

progression.

ABSTRACT

Germ cells undergo mitotic expansion via incomplete cytokinesis, forming cysts of

undifferentiated cells that remain interconnected prior to meiotic initiation, through mechanisms

that are not well-defined. In somatic cells, an actomyosin contractile network controls cleavage

furrow formation and abscission and is spatiotemporally regulated by Ras-related nuclear protein

(Ran). Here, we identify Ran and β-importins as critical regulators of cyst development in the

Drosophila ovary. Depletion of *Ran* or the β-importins *Tnpo-SR* and *cadmus* disrupts oocyte

selection and results in egg chambers with variable numbers of germ cells, suggesting abnormal

cyst development and cyst fragmentation. We demonstrate that Ran, Tnpo-SR, and Cadmus

regulate key cellular processes during cyst formation, including cell cycle dynamics, fusome

biogenesis, and ring canal stability, yet do so independently of mitotic spindle assembly. Further,

Tnpo-SR and Cadmus control cyclin accumulation and suppress cytokinesis independent of Ran-

GTP, suggesting that β-importins sequester protein cargos that normally promote the mitotic-to-

meiotic transition. Our data demonstrates that Ran and β-importins are critical for germ cell cyst

formation, a role that is likely conserved in other organisms.

INTRODUCTION

Germ cells develop as cysts of undifferentiated cells that remain that remain interconnected

during mitotic division (Lu et al., 2017; Matova and Cooley, 2001; Pepling and Lei, 2018). Cysts

arise via incomplete cytokinesis, manifested by cleavage furrow arrest and the subsequent

formation of stable ring canals that interconnect cells and allow transport of cytoplasm and

organelles (de Cuevas et al., 1997; Guertin et al., 2002; Haglund et al., 2011; Pepling and Lei,

2018). Cyst formation has been postulated to increase cytoplasmic volume, enhance sensitivity to

DNA damage, and ensure robust oocyte development (Chaigne et al., 2020; de Cuevas et al., 1997;

Pepling and Lei, 2018; Pepling et al., 1999; Yamashita, 2018). The evolutionary conservation of

cyst formation suggests that germ cell interconnectivity is biologically relevant and in support of

this, stable intercellular bridges are required for optimal fertility in multiple phyla (Haglund et al.,

2011; Lu et al., 2017).

The fruit fly, *Drosophila melanogaster*, is an elegant model of germline cyst formation. In

adult females, ovaries consist of 16-20 ovarioles, in which oocytes are produced in a linear

spatiotemporal arrangement fueled by a tissue-resident stem cell population (Hinnant et al., 2020).

Germline stem cells (GSCs) are housed at the anterior of ovarioles within a somatic cell niche

(Figure 1A). GSCs divide asymmetrically, giving rise to one daughter that will remain a GSC and

a cystoblast which initiates differentiation. Delayed abscission between the GSC and nascent

cystoblast keeps the pair connected well into G2 of the subsequent cell cycle (Ables and

Drummond-Barbosa, 2013; Eikenes Å et al., 2015; Hinnant et al., 2017; Mathieu et al., 2013;

Matias et al., 2015). Following abscission, cystoblasts divide exactly four times with incomplete

cytokinesis, creating 16-cell cysts. During and following mitotic division, cleavage furrows

between germ cells are modified into stable ring canals, which facilitate organelle transport to and

selection of one definitive oocyte within the 16-cell cyst (Grieder et al., 2000; Guertin et al., 2002;

Hinnant et al., 2020). The fusome, an unusual cytoplasmic organelle composed of microtubules,

endoplasmic reticulum-derived vesicles, and a meshwork of membrane skeleton proteins, traverses

the intercellular bridges of GSC/cystoblast pairs and cysts (Hinnant et al., 2020; Ong and Tan,

2010; Pepling and Lei, 2018; Röper, 2007; Snapp et al., 2004). The fusome branches as mitotic

divisions progress, providing a synchronization hub for cell cycle regulation and cell polarity

(Grieder et al., 2000; Hinnant et al., 2020; Lighthouse et al., 2008). These specialized cytological

features of germ cells facilitate interconnectivity and are vital to oocyte selection; however, the

molecular mechanisms underlying fusome morphogenesis and incomplete cytokinesis are not well
understood.

In cells that undergo complete cytokinesis, formation of an actomyosin-based contractile
network at the equatorial cleavage plane is spatiotemporally regulated and requires a dynamic
interplay between the cytoskeleton, the mitotic microtubule-based central spindle, and membrane
trafficking (Frémont and Echard, 2018; Haglund et al., 2011; Nguyen and Robinson, 2020; Pollard
and O'Shaughnessy, 2019). Central to this process is Ras-related nuclear protein (Ran), a small
guanosine triphosphate (GTP)-binding protein implicated in protein transport during mitosis and
interphase (Ozugergin and Piekny, 2020; Sazer and Dasso, 2000). Ran partners with α- and β-
importins to move protein cargo into the nucleus and promote mitotic spindle assembly (Cautain
et al., 2015; Okada et al., 2008). Ran also regulates cell polarization and contractile protein
localization during cleavage furrow ingression (Beaudet et al., 2017; Silverman-Gavrila et al.,
2008). Chromatin-associated Ran-GTP may help recruit and organize essential proteins at the
cortex, thereby ensuring the robustness of cytokinesis (Nguyen and Robinson, 2020). The
biological relevance of importins or Ran to germline cyst formation has not been established.

Here, we identify Ran and β-importins as novel regulators of GSC maintenance and germ
cell cyst formation in the *Drosophila* ovary. We show that Ran and β-importins Tnpo-SR and
Cadmus (Cdm) control cyst formation by regulating key cellular processes, including cell cycle
dynamics, fusome biogenesis, and ring canal stability. Ran function depends on its active, GTP-
bound state, as expression of a dominant-negative *Ran* mutant recapitulates *Ran* loss-of-function.
Ran, *Tnpo-SR*, and *cdm* are not essential for mitotic spindle formation in undifferentiated germ
cells. Moreover, knock-down of *Tnpo-SR* or *cdm* does not completely phenocopy loss of *Ran*,
suggesting that β-importins control cell cycle progression and suppress cytokinesis independent of
Ran activity. Given the conservation of germline cyst formation across phyla, our data suggest that
Ran and specific β-importins may play similar roles in other organisms.

RESULTS
RNAi screen reveals that β-importins and Ran are essential for germ cell development.
We previously identified *Transportin-Serine/Arginine rich* (*Tnpo-SR*) in a genetic mosaic
screen for genes controlling GSC self-renewal (Ables et al., 2016). *Tnpo-SR*, homologous to
mammalian *TNPO3*, encodes a β-importin (Allemand et al., 2002; Lai et al., 2001). To determine

whether *Tnpo-SR* loss-of-function phenotypes reflect a general role for importins in germ cell

mitoses, we performed a secondary reverse genetic screen using germline-enhanced RNA

interference (RNAi) to knock-down expression of nine *Drosophila* importins and the small

GTPase *Ran* specifically in germ cells (Figure 1B-E) (Ni et al., 2011; Rørth, 1998; Van Doren et

al., 1998). Ovarioles were quantified for the presence of germ cells based on the unique nuclear

morphology of nurse cells and oocytes (Figure 1B). Most of the RNAi lines produced ovarioles

with seven to eight egg chambers, similar to driver-only controls (Figure 1B,E). In contrast, three

RNAi lines produced completely agametic ovaries, defined by the absence of egg chambers

(Figure 1C,E). RNAi line *Tnpo/CG8219HMJ23009* carries a short hairpin that recognizes both *Tnpo*

and the related paralog *CG8219*. Since two other RNAi lines specific for *Tnpo* did not display

germ cell phenotypes, we speculate that Tnpo and CG8219 may compensate for each other, as they

share 80% amino acid similarity. Only one RNAi line for the β-importin *moleskin* (*mskHMS00020*)

displayed an agametic phenotype, making it difficult to conclude whether the RNAi accurately

reflects a role in germ cell development. More intriguingly, four RNAi lines corresponding to *Ran*

and the β-importins *Tnpo-SR*, *cadmus* (*cdm*), and *Fs(2)Ket* (also called *Ketel*) exhibited a partial

agametic phenotype (Figure 1D-E). Some ovarioles lacked germ cells, while other ovarioles

contained germ cells in unusual egg chamber morphologies (Figure 1D). Cdm and Tnpo-SR are

paralogs, sharing 42% amino acid similarity, while Ketel is less conserved. These results

demonstrate that Ran and β-importins are necessary for proper oocyte development and may have

distinct yet overlapping functions in the germline.

Ran, Tnpo-SR, Ketel, and cdm are necessary to establish GSCs.

Since GSCs give rise to all germ cells, we hypothesized that agametic ovaries resulting

from *Ran* and *β-importin* knock-down could arise from a failure to establish or maintain GSCs. To

test this, we collected ovaries from control and germ cell *Ran* or *β-importin* RNAi knock-down

females at successive days after eclosion and quantified the number of GSCs per germarium. GSCs

and their progeny can be easily identified by immunolabeling of Hts, a fusome-resident protein in

germ cells, and LaminC (LamC), a nuclear envelope protein highly expressed in cap cells (the

major cellular component of the GSC niche) (Ables and Drummond-Barbosa, 2013). While

fusomes are present in GSCs and their dividing daughters, only GSCs possess an anteriorly

localized fusome juxtaposed to the cap cells (de Cuevas and Spradling, 1998; Lin et al., 1994).

Driver control ovaries averaged 2.8 GSCs per germarium just after eclosion, and consistent with

previous studies, the average decreased slightly over time as females aged (Supp. Figure 1A,C)

(Pan et al., 2007). All four RNAi lines giving a partial agametic phenotype upon germline-specific

knock-down had significantly less GSCs per germarium at three days after eclosion (Supp. Figure

1B-C). Fewer GSCs in early adulthood suggests that *Ran*, *Tnpo-SR*, and *cdm* are required for GSC

establishment during larval/pupal stages or during the proliferative surge in GSCs following the

female's first mating (Ameku and Niwa, 2016; Van Doren et al., 1998). The number of GSCs per

germarium continued to decline with age following knock-down of *Ran*, *Tnpo-SR*, and *cdm* (Supp.

Figure 1C).

150   To test whether Ran or β-importins are also required for GSC maintenance, we used a

conditional knock-out approach to specifically abrogate *Tnpo-SR* in adult GSCs and quantified the

number of GSCs remaining in the tissue over time. The *Flippase* (*FLP*)/*FLP Recognition Target*

(*FRT*) genetic mosaic lineage-tracing system allows for the generation of clonal populations of

homozygous mutant cells in otherwise heterozygous animals (Laws and Drummond-Barbosa,

2015; Xu and Rubin, 1993). We used two strong loss-of-function transposon insertion alleles of

Tnpo-SR (*Tnpo-SR^{KG04870}* and *Tnpo-SR^{LL05552}*) carrying *FRT* sites to generate germline mutant

mosaic germaria (Supp. Figure 1D-F). To quantify GSC loss, we measured the percentage of total

germaria harboring a mutant (GFP-negative) GSC. Mock mosaic germaria were used as controls,

as GFP-negative GSCs can be produced even though all cells are genotypically wild-type (Supp.

Figure 1D). GFP-negative GSC clones were generated at equivalent rates in control, *Tnpo-*

SR^{KG04870}, and *Tnpo-SR^{LL05552}* mosaic germaria at four days after clone induction (Supp. Figure

1F). In contrast to mock controls, *Tnpo-SR^{KG04870}* and *Tnpo-SR^{LL05552}* germline mutant mosaic

germaria rapidly lost marked GSCs over time (Supp. Figure 1E,F), indicating that *Tnpo-SR* is cell-

autonomously required for GSC self-renewal. *Tnpo-SR* mutant GSCs likely adhere to cap cells as

well as adjacent wild-type GSCs, as they express equivalent levels of the adhesion protein E-

cadherin (Supp. Figure 1G,G'). Loss of *Tnpo-SR* appears to uncouple cell growth and cell division,

as *Tnpo-SR* mutant GSCs grow larger than adjacent wild-type GSCs (Supp. Figure 1G,G'),

reminiscent of *Cdk2* or *CycE* null GSCs (Ables and Drummond-Barbosa, 2013). Taken together

with the RNAi data, these results suggest prominent roles for *Ran*, *Tnpo-SR,* and *cdm* in GSC

maintenance.

Loss-of-function mutations in Ran and β-importins alter the number of germ cells per egg
chamber.

The intriguing agametic phenotype resulting from knock-down of *Ran*, *cdm*, *Tnpo-SR*, and
Ketel prompted us to take a closer look at egg chamber morphology. Egg chambers develop in a
stereotypical fashion, where germline cysts are encapsulated by a monolayer of follicle cells as
cysts move posteriorly out of the germarium (Figure 1A). To visualize germline and follicle cells,
we combined anti-Hts antibodies (which detect Hts at the plasma membrane in follicle cells) with
anti-Hts-RC antibodies, which are specific for a cleavage product of the Hts polyprotein that
localizes to ring canals (Petrella et al., 2007; Robinson et al., 1994). Driver control egg chambers
had 16 germ cells connected by 15 ring canals (Figure 2A,D). Knock-down of *Ran*, *cdm*, or *Tnpo-*
SR, however, resulted in significantly less egg chambers with 16 germ cells (Figure 2B-D).
Depletion of *Ketel* did not significantly impact egg chamber morphology (Figure 2D).

Germ cell cysts normally divide synchronously four times, following a 2^n pattern, to
produce cysts with 16 cells (King, 1970). To determine whether extra germ cells in *Ran*, *cdm*, and
Tnpo-SR depleted ovarioles arose by alterations in the number of mitotic divisions, we quantified
the number of germ cells per cyst (Figure 2E). In both control and *Ketel* RNAi ovaries, egg
chambers rarely contained more than 16 cells (Figure 2D,F); in these rare instances, egg chambers
invariably contained 32 cells (Figure 2E). In contrast, knock-down of *Ran*, *cdm,* or *Tnpo-SR*
produced egg chambers with variable numbers of germ cells, ranging anywhere from one to 49
cells per cyst (Figure 2E). While some egg chambers housed germ cells on the order of 2^n, a
substantial number deviated from this pattern (Table 1). Ggerm cell numbers in adjacent chambers
in RNAi-depleted egg chambers did not add up to 8, 16 or 32 cells, arguing that encapsulation
defects alone are not the source of the unusual germ cell number phenotype. Some egg chambers
with 32 or more cells had oocytes with five ring canals, suggesting that some abnormal egg
chambers arose by an extra mitotic division. Other egg chambers, however, likely arose because
of other cellular defects. Moreover, a small percentage of *Tnpo-SR* and *cdm* RNAi knock-down
ovarioles (13.3% and 17.4% of the total number of ovarioles, respectively) contained germline
tumors, in which a few abnormally-shaped egg chambers with a seemingly normal follicle cell
monolayer surrounded mitotically-active germ cells with underdeveloped fusomes (Figure 2G-H;
Supp. Figure 2A-C'). In *Tnpo-SR* RNAi ovarioles, undifferentiated germ cells were found
encapsulated with more-developed germ cells (Supp. Figure 2A). Germline tumors contained cells

with short or unbranched fusomes, resembling those of 2-cell cysts, and immature ring canals that

frequently appeared as dot-like structures rather than rings (Supp. Figure 2A-B). These structures

are reminiscent of the abscission site in the cytokinetic furrow separating GSC/cystoblast pairs

(Supp. Figure 1I-K), suggesting that some tumors arose because of improper abscission events in

2-cell and/or 4-cell cysts. Immunostaining for Sex-lethal (Sxl), which is enriched in GSCs and

cystoblasts in the anterior-most region of the germarium (Supp. Figure 2D-D') (Chau et al., 2009),

confirmed that *Tnpo-SR* RNAi germ cells have not completed differentiation, as Sxl was expressed

at similar levels in cells throughout the germarium (Supp. Figure 2E-E'). We conclude that Ran,

Tnpo-SR, and Cdm impact the synchrony and number of cyst divisions.

Ran, Tnpo-SR, and cdm are required for oocyte specification.

During germ cell mitotic divisions, three to four cyst cells initiate meiosis by assembling

the synaptonemal complex; however, only one of these cells is maintained as the oocyte (Hughes

et al., 2018). Since depletion of *Ran*, *Tnpo-SR*, or *cdm* altered the number of germ cells per cyst,

we hypothesized that oocyte selection was also affected. We analyzed the first three egg chambers

outside of the germarium, where the oocyte-specific proteins Orb (Figure 3A) and Bicaudal D

(BicD) (Figure 3D) had clearly accumulated in one posterior cyst cell in driver control egg

chambers. In contrast, knock-down of *Ran*, *Tnpo-SR*, or *cdm* resulted in egg chambers lacking

either Orb (*Ran^{GL01341}, cdm^{HMS02846}, and Tnpo-SR^{V22}*) or BicD expression (*Tnpo-SR^{V22}*) (Figure 3B-

C, E-F). Moreover, many *Ran*, *Tnpo-SR*, and *cdm* loss-of-function egg chambers lacked a cell with

a condensed nucleus or karyosome, a clear indicator of oocyte specification (Figure 3B,E,F). Since

knock-down of *Ran*, *Tnpo-SR*, and *cdm* can alter the number of cells per cyst, we quantified only

egg chambers with 16 cyst cells (Figure 3G). Although the majority of *nos-Gal4>Ran^{GL01341} and*

nos-Gal4>cdm^{HMS02846} successfully specified a single oocyte, 34% of *nos-Gal4>Tnpo-SR^{V22}* egg

chambers contained only nurse cells (Figure 3G). These data suggest that loss of oocyte identity

is a secondary effect of incorrect mitotic divisions in the absence of *Ran* and *cdm*, but that *Tnpo-*

SR may play independent roles in oocyte specification. We therefore immunostained for Orb in

Tnpo-SR mutant mosaic germaria and counted the number of 16-cell cysts in the germarium with

at least one Orb-positive cell. *Tnpo-SR^{KG04870}* mutant cysts located posterior to wild-type cysts (and

thus at a more advanced stage of development) failed to up-regulate Orb expression to similar

levels as adjacent wild-type cysts (Figure 3H-I). Moreover, a core component of the synaptonemal

complex, C(3)G, which is normally present as early as the 4-cell cyst stage, did not accumulate in

Tnpo-SR^{KG04870} mutant 8-cell or 16-cell cysts to equivalent levels as adjacent wild-type cysts

(Figure 3J-K). Together with the RNAi data, this suggests that *Tnpo-SR* is specifically required for

oocyte specification.

GSC establishment, cyst formation, and oocyte selection require active Ran-GTP.

Since Ran and β-importins work together to drive nucleocytoplasmic transport in other

systems (Chang and Hsia, 2020; Nguyen and Robinson, 2020; Ozugergin and Piekny, 2020), we

expected that depletion of *Ran* would phenocopy loss of either *Tnpo-SR* or *cdm*. Our data, however,

revealed key differences between the loss-of-function phenotypes, suggesting that β-importins

might have roles in germ cell cyst formation independent of Ran, or cellular functions other than

cytoplasmic-to-nuclear trafficking.

To test these possibilities, we first analyzed the subcellular distribution of Ran in

mitotically dividing germ cells. We took advantage of a previously characterized transgene in

which the germline-specific *UASp* promoter drives an N-terminal hemagglutinin (HA)-tagged Ran

coding region (*UASp-Ran^{HA}*) (Cesario and McKim, 2011). Using *nos-Gal4* to drive *UASp-Ran^{HA}*

in germ cells, we then localized Ran using anti-HA antibodies in comparison to mitotic spindles

(labeled with α-tubulin), fusomes, and chromosomes (Figure 4A-A'). Unlike neuroblasts, in which

Ran-HA is nuclearly localized during interphase and overlaps with the mitotic spindle during

mitosis (Cesario and McKim, 2011), we did not observe strong nuclear localization of Ran-HA in

GSCs or dividing germ cells (Figure 4A-A'). Ran-HA was evenly distributed, or slightly higher in

the cytoplasm than in nuclei, of interphase and mitotically-dividing cells. Consistent with

neuroblasts and embryonic cells (Cesario and McKim, 2011; Trieselmann and Wilde, 2002), we

did not observe Ran on chromosomes (Figure 4A'). Ran-HA expression was higher in some cysts

than others, likely due to minor variations in driver strength. Moreover, and in contrast to the

meiotic spindles in late stage egg chambers (Cesario and McKim, 2011), we did not observe strong

accumulation of Ran-HA surrounding mitotic spindles (Figure 4A-A'). Lack of strong nuclear

localization or accumulation at spindles argues against roles for Ran in cytoplasmic-to-nuclear

trafficking or mitotic spindle formation in dividing germ cells.

We next asked whether active Ran-GTP is required for GSC establishment and cyst

formation. Like other small GTP-binding proteins, Ran cycles between active (GTP-bound) and

inactive (GDP-bound) states (Ozugergin and Piekny, 2020). Ran-GTP dissociates importin-bound

cargo, increasing the amount of free importin available to bind cargo. We over-expressed two

transgenes carrying mutated forms of Ran in germ cells under the control of *nos-Gal4*. The first

transgene, *UASp-Ran^{T24N}*, carries an amino acid substitution that locks Ran into a GDP-bound state

and inhibits cargo loading (Cesario and McKim, 2011; Drutovic et al., 2020). Mutant flies were

compared to flies expressing wild-type *UASp-Ran^{HA}*. We expected that constitutively expressing

Ran-GDP would titrate out the amount of active Ran, thus mimicking *Ran* loss-of-function.

Indeed, over-expression of *UASp-Ran^{T24N}* resulted in a partial agametic phenotype (Figure 4B,F),

and the average number of GSCs per germarium was substantially reduced as compared to age-

matched *nos-Gal4>UASp-Ran^{HA}* controls (Figure 4G). Over-expression of dominant negative

Ran^{T24N} also altered the number of germ cells per cyst (Figure 4C,J) and, similar to knock-down

of *Ran* via RNAi (Figure 2E), the number of germ cells did not always follow a 2^n pattern (Figure

4I, Table 1). Over-expression of *Ran^{T24N}* also impaired oocyte selection, as 12-24% of egg

chambers with 16 germ cells lacked an Orb-positive cell (Figure 4J). These phenotypes are

consistent with depletion of *Ran* via RNAi and likely contribute to the female sterility phenotype

noted in prior studies (Cesario and McKim, 2011).

We then used a second transgene, *UASp-Ran^{Q69L}*, to over-express a constitutively active

form of Ran locked in a hydrolysis-resistant GTP-bound state (Cesario and McKim, 2011).

Intriguingly, excess Ran-GTP in germ cells did not result in severe germline defects (Figure 4E-

F). The average number of GSCs per germarium was slightly reduced in *nos-Gal4>UASp-Ran^{Q69L}*

ovarioles (Figure 4G) and a few egg chambers had an incorrect number of cells (Figure 4H-I), but

defects were not as severe or penetrant as either the *Ran* RNAi or Ran-GDP-locked mutants. Since

we did not observe a clear Ran intracellular concentration gradient or accumulation of Ran at

mitotic spindles, these results suggest that Ran-GTP levels are already in excess in mitotically-

active germ cells, perhaps to facilitate timely mitotic divisions.

Ran-GTP, Tnpo-SR, and cdm are dispensable for mitotic spindle formation in dividing cysts.

One well-characterized function of Ran and β-importins is the regulation of spindle

assembly (Forbes et al., 2015; Kosyna and Depping, 2018). Spindle orientation is critical for GSC

maintenance, and mitotic spindle remnants accumulate in fusomes as cysts divide (Cheng et al.,

2008; Deng and Lin, 1997; Grieder et al., 2000; Máthé et al., 2003). We hypothesized that primary

defects in spindle formation could underlie the defective cyst division in our Ran-GTP and
importin-depleted models. Using immunostaining for α-Tubulin, we confirmed that mitotic
spindles in dividing cysts are anchored to the fusome, oriented perpendicularly (Figure 5A)
(Megraw and Kaufman, 2000; Storto and King, 1989). Surprisingly, in *cdm*-depleted (Figure 5B)
or *Tnpo-SR*-depleted (Figure 5C) germ cells, similar-sized spindles formed and were appropriately
anchored and oriented with respect to fusomes. Similarly, we were unable to find defects in spindle
formation or orientation in *nosGal4>UAS-Ran^{T24N}* germ cells (Figure 5D). These data demonstrate
that mitotic spindles can form even when Ran-GTP is depleted, suggesting that very low levels of
active Ran are sufficient to maintain germ cell mitoses.
305
Tnpo-SR and cdm, but not Ran-GTP, are required for fusome biogenesis and maturation.
The fusome extends through ring canals during incomplete cytokinesis, forming an
extensive tubular network (Lighthouse et al., 2008; Röper, 2007; Snapp et al., 2004). The balance
of cytoskeleton to vesicular-like structures in the fusome changes over the course of the mitotic
divisions, as the cytoskeletal core (which can be visualized with anti-Hts antibodies; Figure 5E) is
prominent in GSC and cystoblast fusomes, but eventually degrades as cysts exit the germarium
(Röper, 2007). Loss of *Hts* alters the number of germ cells per cyst (Yue and Spradling, 1992).
Since knock-down of *Ran* or the β-importins also altered the number of germ cells per cyst, we
hypothesized that abnormal fusome morphogenesis might abrogate cyst formation. Using Hts
immunostaining as an indicator of fusome morphology, we observed that when *cdm* (Figure 5F)
or *Tnpo-SR* (Figure 5G and Supp. Figure S2) were depleted, germ cells formed long, thin fusomes
with irregular branching, particularly in cysts in the middle to posterior third of germaria. Fusomes
in *Tnpo-SR*- and *cdm*-depleted germ cells often had thread-like sections separated by thicker
sections, consistent with weak points in the cytoskeletal structure (see arrowheads in Figure 5F-
G). Irregular fusomes were also observed in *Tnpo-SR*-depleted GSC/cystoblast pairs, where new
fusome material builds from the cystoblast to the GSC (Supp. Figure 1 I-J) (de Cuevas and
Spradling, 1998). Importantly, we did not observe fusomes connecting GSCs to other cyst cells in
stem-cysts, as was reported for germ cells in which the chromosomal passenger complex, which
regulates abscission timing in GSCs, is abrogated (Mathieu and Huynh, 2017). We also did not
observe GSCs or cysts with asynchronous mitoses (Supp. Figure S1L-M), as we would predict if
fusomes did not maintain some degree of connectivity between cells. We conclude that the

irregular fusome branching patterns in *Tnpo-SR-* and *cdm*-depleted germ cells likely result from
impaired fusome maintenance or remodeling.

To further investigate how the importins regulate fusome morphogenesis, we recombined
the *Tnpo-SR^{V22}* RNAi transgene with two reporter transgenes in which the coding regions of
fusome-resident proteins were fluorescently tagged. *Scribble* (*scrib*) encodes a scaffolding protein
that regulates vesicle trafficking and is expressed at the cell cortex and in immature fusomes of
GSCs and dividing cysts (Figure 5I) (Lighthouse et al., 2008). By the third mitotic division, *Scrib-*
GFP is largely absent from wild-type fusomes and does not co-localize with Hts in 16-cell cysts.
In *Tnpo-SR*-depleted germ cells, however, *Scrib-GFP* was prominently expressed in the long, thin
fusomes of germ cells in the posterior of germaria (Figure 5J). Conversely, Reticulon-like1 (Rtnl1)
is an endoplasmic reticulum-resident protein that is expressed in low levels in immature fusomes
(Figure 5K) (Röper, 2007). *Rtnl1-GFP* becomes enriched in mature fusomes coincident with cyst
division and inversely correlated with Hts expression, reaching peak concentration in fusomes
after the final mitotic division (Figure 5K). When *Tnpo-SR* is knocked down in germ cells, *Rtnl1-*
GFP and Hts co-localize in fusomes throughout the posterior end of the germarium (Figure 5L).
These data suggest that in the absence of β-importins, cytoskeletal components of the fusome fail
to degrade properly, maintaining the fusome in an immature state that is weakly connected between
cyst cells. Intriguingly, fusome shape and branching in cysts over-expressing the dominant
negative *Ran^{T24N}* mutant (Figure 5D) were indistinguishable from wild-type cysts, suggesting that
Cdm and Tnpo-SR regulate fusome morphogenesis independently of Ran-GTP.

Tnpo-SR and cdm promote cyst formation by suppressing cytokinesis.

Knock-down of *Tnpo-SR* or *cdm* produced egg chambers variable numbers of germ cells;
however, it remained unclear to us exactly how such egg chambers could arise. Key observations
came by closely examining the morphology of *Tnpo-SR* mutant cysts surrounded by wild-type
cysts in mutant mosaic germaria. While knock-down of *Tnpo-SR* by RNAi led to >16 germ cells
per egg chamber, we only rarely found examples of *Tnpo-SR^{KG04870}* or *Tnpo-SR^{LL05552}* mutant egg
chambers with more than 16 mutant germ cells. We did find, however, compound egg chambers
in which a *Tnpo-SR* mutant cyst was encapsulated together with wild-type cysts (Supp. Figure
S3A). We occasionally identified mitotically-dividing *Tnpo-SR^{KG04870}* and *Tnpo-SR^{LL05552}* mutant
germ cells in mosaic egg chambers outside of the germarium, which never occurs in wild-type

germ cells (Supp. Figure S3B). We also frequently found clusters of single *Tnpo-SR* mutant germ cells at the posterior of germaria (Supp. Figure S3C), suggesting that some cyst defects in *Tnpo-SR* mutant and RNAi-knock-down ovarioles could arise by fragmentation of cysts into single germ cells. In the *Tnpo-SR* RNAi model, where all germ cells lack appropriate Tnpo-SR levels, fragmented cysts would likely be encapsulated along with adjacent cysts, thus yielding irregular (i.e., not 2^n) numbers of germ cells per cyst. In support of this idea, we observed many examples of mutant mosaic germaria in which *Tnpo-SR^{KG04870}* mutant cells with dot-like fusomes were adjacent to one another with fusomes aligned along the same plane, as if they were connected (Supp. Figure S3D). Immunostaining with anti-pTyr antibodies to visualize ring canals highlighted *Tnpo-SR^{KG04870}* mutant cysts where ring canal components accumulated on adjacent cyst cells, separated by a very thin line of GFP-positive cytoplasm (Figure 6A and Supp. Figure S3E), suggesting that cells have physically separated from each other after clone induction. Moreover, *Tnpo-SR^{KG04870}* mutant germ cells failed to accumulate proper levels of the cell adhesion protein E-cadherin between germ cells, as occurs in adjacent wild-type 16-cell cysts following the final mitotic division and prior to follicle cell encapsulation (Figure 6B) (Fichelson et al., 2010).

These data raised the possibility that in the absence of *Tnpo-SR*, cysts which normally form via incomplete cytokinesis fail to stay connected by intercellular bridges, and instead break apart at unstable ring canals. Initially, this seemed unlikely, as our first analyses of *Ran*, *cdm*, and *Tnpo-SR* RNAi and *Ran* dominant-negative germ cells focused on egg chambers outside of the germarium, where ring canals containing Hts-RC (a constituent protein of mature ring canals) were easily detected (Figure 2). Moreover, the number of ring canals scaled to the number of germ cells, suggesting that ring canals formed at successive divisions. Spatiotemporal comparison of ring canal formation, however, demonstrated that while Hts-RC is expressed in germ cells from *nos-Gal4>Tnpo-SR^{V22}* ovarioles, its accumulation at ring canals is drastically delayed, such that protein levels are much lower than in age-matched wild-type cysts (compare Figure 6C and D). Similarly, anti-pTyr antibodies are typically the first molecular indicator of mature ring canals (Figure 6E). In the germ cells depleted of *cdm* (Figure 6F) or *Tnpo-SR* (Figure 6G), pTyr antibodies failed to accumulate to wild-type levels in 16-cell cysts. In contrast, both Hts-RC and pTyr accumulated at well-defined ring canals when the dominant negative *Ran^{T24N}* mutant was expressed in germ cells (Figure 6H). These data support the model that Tnpo-SR and Cdm, but not Ran-GTP, function to stabilize germ cell intercellular bridges, maintaining proper cyst formation (Figure 6M).

We then performed immunostaining for Anillin, an actin/myosin-binding protein
previously demonstrated as an importin cargo (Beaudet et al., 2017; Beaudet et al., 2020; Chen et
al., 2015; Piekny and Maddox, 2010; Silverman-Gavrila et al., 2008). Anillin localization in ring
canals begins at the third mitotic division (Figure 6I) (Goldbach et al., 2010). In *Tnpo-SR*-depleted
germ cells, we observed Anillin deposition within the intercellular bridge, rather than on
surrounding rings (Figure 6J). In contrast, Anillin localizes properly to the ring canals in *Ran^{T24N}*
mutants (Figure 6K) and occasionally co-localized with the fusome (Figure 6L), supporting the
model that Tnpo-SR suppresses cytokinesis in dividing cysts independently of Ran-GTP.
Intriguingly, over-expression of a GFP-tagged Anillin in which a nuclear localization signal has
been mutated did not alter Anillin localization in abscising GSC/CB pairs (Supp. Figure 4A-B) or
at ring canals (Supp. Figure 4C-D), and did not change the number of germ cells per cyst. Although
we cannot exclude the possibility that Tnpo-SR may bind Anillin at another protein domain, we
speculate that Tnpo-SR indirectly regulates Anillin by binding other cargo proteins that impact
Anillin localization.

Ectopic CycA accumulates in Tnpo-SR- and cdm-depleted germ cells.
The cell cycle is tightly regulated to coordinate cyst formation. For example, loss of
function of the mitotic cyclin CycB produces cysts with eight cells, while over-expression of CycA
or CycE produces cysts with 32 cells (Lilly et al., 2000; Mathieu et al., 2013; Morris et al., 2005;
Ohlmeyer and Schüpbach, 2003). CycA and CycB accumulate at the fusome and phosphorylation
of CycB by AuroraB Kinase promotes abscission between GSC/cystoblast pairs (Lilly et al., 2000;
Mathieu et al., 2013). Given the phenotypic parallels, we hypothesized that Tnpo-SR and Cdm
might regulate cyclin dynamics in dividing germ cells. We confirmed that CycA accumulates at
the fusome in ~10% of mitotically dividing cysts, but never associates with the fusome of 16-cell
cysts in wild-type tissue (Figure 7A,F). *Ran* depletion in *nos-Gal4>Ran^{GL01341}* germaria did not
alter CycA kinetics (Figure 7B,F). In contrast, nearly 40% of mitotically-dividing cysts in *Tnpo-*
SR and *cdm*-depleted germ cells expressed CycA; moreover, CycA expression was observed in the
fusomes of 16-cell cysts at an equivalent rate as dividing cysts, suggesting that germ cells remain
mitotic (Figure 7C-D,F).
As the highest peak of CycB expression occurs following CycA degradation (Morris et al.,
2005), we then asked whether CycB regulation was likewise impacted by loss of the β-importins.

We utilized the *Drosophila* Fluorescence Ubiquitin-based Cell Cycle Indicator system (Fly-

FUCCI) (Zielke et al., 2014) to ask whether CycB expression was altered in the absence of *Tnpo-*

SR. Importantly, we used a version of the *UASp-CycB-RFP* transgene that includes a canonical

nuclear localization sequence (Hinnant et al., 2017; Zielke et al., 2014). When *Tnpo-SR* was

depleted, the *UASp-CycB-RFP* localized to germ cell nuclei; however, significantly fewer

mitotically dividing cysts expressed CycB as compared to wild-type controls (Figure 7E,G). Taken

together, these data suggest that *Tnpo-SR*, and likely *cdm*, promote proper cyclin dynamics in germ

cells, maintaining cysts in the mitotic expansion phase and facilitating coordination between cyst

division and fusome morphogenesis.

DISCUSSION

During germline cyst formation, cell cycle transitions are precisely controlled to promote

mitotic divisions and block cellular abscission. Our study adds new details to our understanding

of the molecular regulation of germ cell mitotic expansion, identifying previously undescribed

roles for β-importins and the small GTPase Ran in mitotic cell cycle exit during cyst formation.

Remarkably, we do not detect a clear cytoplasmic-to-nuclear gradient of Ran protein localization

in germ cell nuclei. Nor do we find defects in mitotic spindle assembly, cell size, or DNA

condensation in the absence of β-importins or activated Ran. Instead, our data suggest that

importins and Ran-GTP promote mitotic exit by distinct, yet complimentary mechanisms. We

propose the model that Ran-GTP drives germ cell mitoses by mobilizing cell cycle factors,

functioning together with the β-importins Tnpo-SR and Cdm to facilitate transport of cell cycle

factors from the fusome to other parts of the cell. We propose that independent of Ran, Tnpo-SR

and Cdm sequester cell cycle regulators like CycA and Anillin, titrating the levels of protein

accessible to Ran-GTP, counterbalancing mitotic progression. Thus, in the absence of Tnpo-SR or

Cdm, CycA levels accumulate, causing germ cells to continue in a proliferative state and impairing

fusome stability, ring canal maturation, and oocyte terminal differentiation.

Importins may exhibit non-redundant functions in the Drosophila germline.

The *Drosophila* genome encodes 15 importin proteins, including four α-importins, which

function as adaptor proteins linking cargo with nuclear localization signals to β-importins (Kimura

and Imamoto, 2014). Although several α-importins have been implicated in oogenesis prior to our

study (Geles and Adam, 2001; Mason et al., 2002; Máthé et al., 2000), our study is the first to

identify specific roles for β-importins in mitotic expansion. Our data generally support the

hypothesis that specific importins regulate distinct aspects of germ cell development (Hogarth et

al., 2005; Loveland et al., 2015; Mihalas et al., 2015). We cannot exclude the possibility, however,

that RNAi was not sufficient to completely inhibit importin function. For example, Ketel localizes

at the nuclear envelope in early embryos and dividing germ cells (data not shown) (Lippai et al.,

2000; Tirián et al., 2000; Trieselmann and Wilde, 2002; Villányi et al., 2008), yet we did not

observe germ cell mitotic defects in *Ketel*-depleted germ cells. Future studies will need to employ

other loss-of-function approaches to test the roles of Ketel and other β-importins more thoroughly.

The repertoire of cargo carried by a β-importin determines its role in biological processes

and its redundancy with other importins (Kimura and Imamoto, 2014). Although the full extent of

protein-protein interactions mediated by Tnpo-SR remains unknown, structural studies indicate

that TNPO3 binds proteins involved in chromatin organization, mRNA and protein modification,

cell cycle control, and cell-cell communication (Kimura et al., 2017; Maertens et al., 2014). Tnpo-

SR is also likely to bind a broad range of cargo proteins, including SR-rich RNA binding proteins.

Indeed, Tnpo-SR physically interacts with several proteins essential for mRNA splicing and

transport (Allemand et al., 2002). Given that SR-rich proteins facilitate diverse RNA processing

events, there are many potential Tnpo-SR cargo proteins that could influence cyst formation and

GSC self-renewal, which are heavily influenced by translational control (Bradley et al., 2015;

Teixeira and Lehmann, 2019). Our future studies will seek to identify Tnpo-SR and Cdm-

interacting proteins, which should provide additional mechanistic details regarding how importins

regulate germ cell mitotic expansion.

The number of cyst divisions is regulated by Ran-GTP and β-importins.

Two major cell fate transitions accompany the development of *Drosophila* oocytes: stem

cell to differentiating progenitor (the cystoblast), and progenitor to differentiated fate. These

transitions are accompanied by changes in cell cycle control, including a dramatic switch from

mitotic expansion to meiosis as terminal differentiation proceeds. How such cell cycle transitions

are accomplished is of broad scientific interest, both in the germline and in somatic cells, where

exit from the cell cycle accompanies differentiation (Liu et al., 2019). Intriguingly, both cell fate

transitions employ common cell cycle regulatory factors, such as cyclins, Cdks, and ubiquitin

ligases, suggesting that these are largely repurposed or differentially regulated from GSCs to
differentiating cyst cells. Precise spatiotemporal control over cell cycle regulators must therefore
underlie the transition to the oocyte fate. In support of this, the mitotic cyclin CycA is clearly
critical to support differentiation. Failure to degrade CycA leads to self-renewal failure in GSCs,
and extra mitotic divisions and loss of oocyte identity in cyst cells (Chen et al., 2009; Ji et al.,
2017; Lilly et al., 2000; Morris et al., 2005; Ohlmeyer and Schüpbach, 2003; Sugimura and Lilly,
2006). Moreover, the deubiquitinase encoded by *bag of marbles* (*bam*), one of the few known
differentiation factors in *Drosophila* germ cells, is necessary for the transition from GSC to
cystoblast and for cystoblast divisions, at least in part because it stabilizes CycA expression in
dividing cysts (Ji et al., 2017). CycA/Cdk1 activity is further controlled at the level of translation
and ubiquitin-mediated proteolysis. Cyclin/Cdk protein levels and activity are very high in early
cyst divisions but decrease as cysts approach the terminal mitotic division (Hinnant et al., 2017;
Lilly et al., 2000). Since cystoblasts divide exactly four times, several prior studies posited the
model that cystoblasts autonomously limit the number of divisions through a molecular counting
mechanism that involves Bam, CycA, and the fusome (Hinnant et al., 2020; Huynh, 2006; Ji et al.,
2017; King, 1970; Lilly et al., 2000; McKearin, 1997). While a counting mechanism would limit
the number of mitotic divisions, it must also promote modification of contractile ring proteins to
block cytokinesis, stabilize intercellular bridges into stable ring canals, and maintain the
biosynthesis of the fusome (McKearin, 1997).

Our data suggest that the Ran/β-importin intracellular transport system is integrated into
the CycA-dependent cell division counting machinery. Depletion of *Ran*, *cdm*, or *Tnpo-SR*
generated egg chambers with abnormal numbers of cells (Table 1), some on the order of 2^n
reminiscent of cyclin mutants (Lilly et al., 2000; Mathieu et al., 2013; Morris et al., 2005;
Ohlmeyer and Schüpbach, 2003), and others deviating from this pattern, more similar to mutants
which disrupt intercellular bridge formation and fusome biogenesis (de Cuevas et al., 1996; Lin et
al., 1994; Yue and Spradling, 1992). Although we cannot establish a clear temporal sequence of
events mediated by Ran from our current data, we envision two possible mechanisms of action.
We favor the hypothesis that Ran mediates intracellular movement of CycA itself or a factor that
promotes CycA degradation. Binding by Tnpo-SR or Cdm could sequester the cell cycle regulator,
preventing activation by sterically inhibiting interaction with a kinase or by blocking protein
accumulation at the fusome. High levels of Ran-GTP levels would promote release of the Tnpo-

SR cargo, freeing it for activation. In the absence of Tnpo-SR or Cdm, where mitotic regulators
cannot be delivered to Ran, CycA levels instead accumulate at the fusome and in the cytoplasm,
promoting continued mitotic division. Alternatively, Ran-GTP could regulate contractile protein
localization and cell polarization, as has been demonstrated for human Ran in mammalian somatic
cells (Beaudet et al., 2017; Silverman-Gavrila et al., 2008). Although we do not detect clear
evidence of altered cytokinesis or failure to properly localize Anillin to ring canals in the dominant
negative Ran^{T24N} mutant, it is possible that our immunofluorescence assay is not sensitive enough
to detect such defects. Indeed, we have been unable to visualize the cortical accumulation of
Anillin in dividing germ cells described by previous studies (de Cuevas and Spradling, 1998) using
the anti-Anillin or *UASp-Anillin-GFP* reagents available to us.

Tnpo-SR and Cdm may have Ran-independent roles in promoting cyst development.

During nucleocytoplasmic transport, an α-importin/β-importin complex or individual β-
importin binds to a cargo in the cytoplasm and transports it across a nuclear pore (Barbato et al.,
2020; Görlich et al., 1997). Inside the nucleus, Ran-GTP binds to the complex, dissociating the
cargo from the importin. Given the close association between Ran and β-importins, we
hypothesized that depletion of *Ran* by RNAi, or over-expression of dominant negative Ran^{T24N}
mutant would phenocopy β-importin depletion. While this held true for GSC maintenance, cyst
division, and oocyte specification, we were surprised to find that fusome biogenesis defects, cyst
fragmentation, and delayed ring canal maturation phenotypes were unique to loss of *Tnpo-SR* and
cdm. Moreover, the constitutively active Ran^{Q69L} mutant, in which Ran is locked in a GTP-bound
configuration and should deplete the amount of free β-importin in cells, did not have severe defects
in germ cell mitoses. These results could indicate Ran-independent functions of Tnpo-SR and Cdm
in mitotically dividing germ cells. Based on molecular markers of the fusome and ring canals, we
conclude that in *Tnpo-SR-* and *Cdm-* depleted germ cells, the transition from mitosis to meiosis is
abrogated or delayed, such that cell-to-cell connections are not maintained. Delayed maturation of
the fusomes and ring canals appears to create weak connections between cells, resulting in
premature abscission or collapsed intercellular bridges (Figure 6M). Interestingly, we see fusomes
that are fragile and unbranched in structure (Figure 5E-H), but also germline tumors containing
cells with immature fusome morphology resembling that of 2-cell cysts (Figure 2G-H). This may
be the result of altered timing or different concentrations of the β-importins in the first division of

the cystoblast versus the remaining three, perhaps reflecting differing sensitivity to abscission

across the mitotic divisions (Mathieu et al., 2013; McKearin and Ohlstein, 1995). Cyst cell

fusomes are composed of endoplasmic reticulum-like vesicles and cytoskeletal elements that

originate from remnants of the mitotic spindle midbody (de Cuevas and Spradling, 1998; Koch

and King, 1966; Koch et al., 1967; Lin and Spradling, 1995; Mahowald, 1971). We speculate that

Tnpo-SR and Cdm promote reorganization of the germ cell cytoskeleton after mitotic division,

perhaps by shuttling cytoskeletal proteins back to the fusome. Although fusome biogenesis has not

been well-studied, it is clearly integral to cyst formation. Our study provides novel reagents to

further interrogate the molecular mechanisms promoting cyst interconnectivity.

Ran-GTP and β-importins may function as a conserved molecular timer for the mitotic-to-

meiotic transition.

The basic steps in cyst formation have been well described at the morphological level,

particularly in insects (de Cuevas et al., 1997; Lu et al., 2017; Pepling and Lei, 2018; Yamashita,

2018). Yet how cyst formation is guided molecularly remains an open area for study. Insects

provide excellent models, as many exhibit unique mechanisms for achieving cyst formation

(Büning, 1994; de Cuevas et al., 1997; Eastin et al., 2020; Kubrakiewicz, 1997). Given the level

of conservation in cell cycle control across species, we propose that Ran and β-importins may also

play important roles in higher eukaryotes, connecting the intricate programs of cell division, oocyte

differentiation, and cell polarization. This may be of particular consequence for mouse and human

oocytes, in which cyst breakdown occurs prior to meiotic onset (Pepling and Lei, 2018).

Understanding how Ran and importins control cyst formation and breakdown may help elucidate

how the oocyte reserve is established in humans, offering new potential avenues for treatment of

infertility.

MATERIALS AND METHODS

Drosophila strains and husbandry

Flies were maintained at 22°-25°C in standard medium (cornmeal/molasses/yeast/agar)

(NutriFly MF; Genesee Scientific). Female progeny were collected one to two days after eclosion

and maintained on standard medium for 3, 6, and 10 days at 25°C; flies were fed wet yeast paste

for 2-3 days (changed daily) prior to ovary dissection. A complete list of fly lines used in this study

are provided in Supp. Table S1.

Tissue-specific RNA interference (RNAi) generation

All RNAi lines used in this study were carried in *pVALIUM20* or *pVALIUM22* transposons

for maximum germline efficiency (Supp. Table S1). *UAS-Tnpo-SR^{V22}* was generated as described

(www.flyrnai.org/TRiP-HOME.html) (Ni et al., 2011). Primers were designed against the full-

length *Tnpo-SR* RNA via the Designer of Small Interfering RNA tool

(http://biodev.extra.cea.fr/DSIR/DSIRhtml) using default settings (21 nt siRNA; score threshold

90). A sequence with a score of 94.6 and 0 predicted off-targets was selected for hairpin generation

(passenger strand, CGATCCCGTTTACTGGATAGA; guide strand,

TCTATCCAGTAAACGGGATC). Primers were designed according to TRiP recommendations,

annealed to form double-stranded oligos, and ligated into EcoRI-HF/NheI-HF-digested

pVALIUM22. Transgenic flies (*UAS-Tnpo-SR^{V22}*) were generated by phiC31 site-specific integrase

into the *attP2* site on the third chromosome.

For RNAi experiments, germline knock-down was facilitated by expressing the germline-

specific *nos-GAL4::VP16-nos.UTR* (referred to throughout as *nos-Gal4*) (Rørth, 1998; Van Doren

et al., 1998). Driver expression was confirmed using *w*; nos-Gal4, UASp-tubGFP* (BDSC #7253)

(Grieder et al., 2000). Females carrying *nos-Gal4* alone were used as controls.

Drosophila genetic mosaic generation

For genetic mosaic analyses using *flippase (FLP)/FLP recognition target (FRT)*, we

obtained the following mutant alleles of *FRT*-containing chromosome arms: *Tnpo-SR^{KG04870}*

[Kyoto *Drosophila* Stock Center (KDSC) #111581] (Bellen et al., 2004) and *Tnpo-SR^{LL05552}*

(KDSC #141628) (Schuldiner et al., 2008). Other genetic tools are described in FlyBase version

FB2020_05 (Thurmond et al., 2019). Genetic mosaics were generated by *FLP/FRT*-mediated

recombination in 2-3-day old females carrying a mutant allele in trans to a wildtype allele (linked

to a *Ubi-GFP* marker) on homologous *FRT* arms, and a *hs-FLP* transgene, as described (Laws and

Drummond-Barbosa, 2015). Flies were heat shocked at 37°C two times per day for 3 days and

incubated at 25°C for 4, 8, or 12 days and supplemented with wet yeast paste on the last 2 days

prior to dissection. Wild-type alleles were used for the generation of control (mock) mosaics.

Germ cell analyses

GSCs were identified based on the juxtaposition of their fusomes to the junction with
adjacent cap cells (de Cuevas and Spradling, 1998). In *FLP/FRT* mutants, GSC loss was measured
two ways. First, as the percentage of total mosaic germaria showing evidence of recent stem cell
loss; namely, the presence of GFP-negative daughters (cystoblasts/cysts generated from an original
GFP-negative stem cell) in the absence of the GFP-negative mother stem cell (Laws and
Drummond-Barbosa, 2015). Second, as the percentage of total germaria analyzed that had at least
one GFP-negative GSC. All results were subjected to Chi-Square analyses using Microsoft Excel.
In RNAi progeny, GSC loss was measured as the average number of GSCs present over three
timepoints. Results were subjected to Student's two-tailed T-test comparing driver controls versus
driver>UAS-RNAi experimental groups at each timepoint individually using Microsoft Excel.

For analyses involving the egg chambers outside of the germarium, data was collected from
the first three egg chambers posteriorly located from the germarium. Oocytes were identified by
the presence of anti-Orb antibodies. The number of cells and ring canals per egg chamber were
calculated based on visualization of ring canals with anti-pTyr and/or anti-Hts-RC antibodies and
cell nuclei with DAPI. Results were subjected to Student's two-tailed T-test comparing driver
controls versus *driver>UAS-RNAi* experimental groups at each timepoint individually using
Microsoft Excel and Prism (GraphPad).

Immunofluorescence and microscopy

Ovaries were prepared for immunofluorescence microscopy as described (Grieder et al.,
2000; Hinnant et al., 2017). In the standard protocol, ovaries were dissected and teased apart in
Grace's medium without additives (Caisson Labs) and fixed in 5.3% formaldehyde in Grace's
medium for 13 minutes at room temperature. They were then washed extensively in phosphate-
buffered saline (PBS, pH 7.4; Fisher) with 0.1% Triton X-100, and blocked for three hours in
blocking solution [5% bovine serum albumin (Sigma), 5% normal goat serum (MP Biomedicals),
and 0.1% Triton X-100 in PBS] at room temperature. To detect cells in S phase, dissected ovaries
were incubated for one hour at room temperature in Grace's media containing 10 µM 5-ethynyl-
2′-deoxyuridine (EdU; Life Technologies). The following primary antibodies were diluted in
blocking solution and used overnight at 4°C: chicken anti-GFP (#13970, Abcam; 1:2000), rabbit

anti-dsRed (#632496, Clontech; 1:500), rabbit anti-phosphoHistone H3 (pHH3; #06-570,

Millipore; 1:200), rat anti-E-Cadherin (DCAD2, Developmental Studies Hybridoma Bank

(DSHB); 1:20), mouse anti-Orb (4H8 and 6H4; DSHB; 1:500), mouse anti-BicD (1B11 and 4C2;

DSHB; 1:10), mouse anti-p-Tyrosine (Millipore; 1:100), mouse anti-Hts-RC (DSHB; 1:10),

mouse anti-CycA (A12; DSHB; 1:50), rabbit anti-C(3)G (a gift from M. Lilly; 1:3000), rat anti-

HA (3F10; Sigma; 1:25), mouse anti-α-Tubulin-AF555 (DM1A-AF555; Millipore; 1:100), and

rabbit anti-Anillin (a gift from J. Brill; 1:50). Mouse anti-Sxl (m18; DSHB; 1:500) was also used

but fixed in a 4% formaldehyde in Grace's medium. Primary antibodies mouse anti-Hts (1B1,

DSHB; 1:10), and mouse anti-Lamin C (LC28.26, DSHB; 1:100) were incubated over two nights

at 4°C. All primary antibodies are followed by a two hour incubation at room temperature with

AlexaFluor 488-, 568- or 633-conjugated goat species-specific secondary antibodies (Life

Technologies; 1:200). EdU was detected (if necessary) using AlexaFluor-594 or -647 via Click-

It chemistry, following the manufacturer's recommendations (Life Technologies).

For stabilizing structures such as mitotic spindles, ovaries were dissected and teased apart

in room temperature equilibrated Grace's medium and fixed for 10 minutes at room temperature

using 350µl Grace's Medium and 200µl of 16% formaldehyde. They were then extensively washed

in PBS with 0.1% Triton X-100 and blocked at room temperature for at least 30 minutes in

blocking solution (2.5mL NGS, 5mL Triton, 42.5mL 1xPBS). Primary antibody incubations took

place overnight at 4°C followed by extensive washing in PBS with 0.1% Triton X-100 and another

blocking period of at least 30 minutes at room temperature. Detection of secondary antibodies

proceeded as described above.

All ovary samples were stained with 0.5 µg/ml 4'-6-diamidino-2-phenylindole (DAPI;

Sigma) in 0.1% Triton X-100 in PBS, and mounted in 90% glycerol mixed with 20% n-propyl

gallate (Sigma). Confocal z-stacks (1 µm optical sections) were collected with either the Zeiss

LSM700 laser scanning microscope using ZEN Black 2012 software or the Zeiss LSM800 laser

scanning microscope using ZEN Blue 2012. Images were analyzed using Zeiss ZEN software, and

minimally and equally enhanced via histogram using ZEN and Adobe Photoshop Creative Suite.

ACKNOWLEDGMENTS

Many thanks to K. McKim (Rutgers University), B. Riggs (San Francisco State University), K.

Roper (University of Cambridge), J.-Q. Ni (Tsinghua University), M. Lilly (NICHD), D.

Drummond-Barbosa (Johns Hopkins University), M. Buszczak (University of Texas Southwestern
Medical Center), L. Cooley (Yale University), and J. Brill (University of Toronto) for their kind
gifts of fly stocks and antibodies. We would also like to thank the East Carolina University School
of Dental Medicine for use of their Zeiss LSM800 microscope. We are grateful to the Bloomington
Drosophila Stock Center and Genomics Resource Center, the Vienna *Drosophila* Resource Center,
the Kyoto *Drosophila* Genomics and Genetic Resources Center, the Transgenic RNAi Project at
Harvard Medical School, the University of Iowa Developmental Studies Hybridoma Bank for
making other fly lines, antibodies, and plasmids available, and BestGene for *Drosophila*
transgenesis and screening. Many thanks to Tianna Van Cura and members of the Ables laboratory
for assistance with building tools, providing helpful discussions, and critical reading of this
manuscript.
678

679 **COMPETING INTERESTS**
680 The authors declare no competing conflicts of interest.
681

682 **FUNDING**
683 This work was supported by National Institutes of Health R15-GM117502 (E. T. A.), the East
684 Carolina University (ECU) Division of Research and Graduate Studies (A. N. B. and T. D. H.),
685 the ECU Office of Undergraduate Research (A. E. W.), and the ECU Honors College (A. M. P.).
686

687 **DATA AVAILABILITY**
688 Fly strains and plasmids are available upon request. The authors affirm that all data necessary for
689 confirming the conclusions of the article are present within the article and figures.
690

REFERENCES
Ables, E. T. and Drummond-Barbosa, D. (2013). Cyclin E controls Drosophila female germline stem cell
maintenance independently of its role in proliferation by modulating responsiveness to niche
signals. *Development* **140**, 530-540.
Ables, E. T., Hwang, G. H., Finger, D. S., Hinnant, T. D. and Drummond-Barbosa, D. (2016). A Genetic
Mosaic Screen Reveals Ecdysone-Responsive Genes Regulating Drosophila Oogenesis. *G3*
(Bethesda, Md.) **6**, 2629-2642.
Allemand, E., Dokudovskaya, S., Bordonné, R. and Tazi, J. (2002). A conserved Drosophila transportin-
serine/arginine-rich (SR) protein permits nuclear import of Drosophila SR protein splicing factors
and their antagonist repressor splicing factor 1. *Molecular biology of the cell* **13**, 2436-2447.

Ameku, T. and Niwa, R. (2016). Mating-Induced Increase in Germline Stem Cells via the Neuroendocrine
System in Female Drosophila. *PLoS genetics* **12**, e1006123.
Barbato, S., Kapinos, L. E., Rencurel, C. and Lim, R. Y. H. (2020). Karyopherin enrichment at the nuclear
pore complex attenuates Ran permeability. *J Cell Sci* **133**.
Beaudet, D., Akhshi, T., Phillipp, J., Law, C. and Piekny, A. (2017). Active Ran regulates anillin function
during cytokinesis. *Molecular biology of the cell* **28**, 3517-3531.
Beaudet, D., Pham, N., Skaik, N. and Piekny, A. (2020). Importin binding mediates the intramolecular
regulation of anillin during cytokinesis. *Molecular biology of the cell* **31**, 1124-1139.
Bellen, H. J., Levis, R. W., Liao, G., He, Y., Carlson, J. W., Tsang, G., Evans-Holm, M., Hiesinger, P. R.,
Schulze, K. L., Rubin, G. M., et al. (2004). The BDGP gene disruption project: single transposon
insertions associated with 40% of Drosophila genes. *Genetics* **167**, 761-781.
Bradley, T., Cook, M. E. and Blanchette, M. (2015). SR proteins control a complex network of RNA-
processing events. *Rna* **21**, 75-92.
Büning, J. (1994). *The Insect Ovary: Ultrastructure, previtellogenic growth and evolution*: Springer,
Dordrecht.
Buszczak, M., Paterno, S., Lighthouse, D., Bachman, J., Planck, J., Owen, S., Skora, A. D., Nystul, T. G.,
Ohlstein, B., Allen, A., et al. (2007). The carnegie protein trap library: a versatile tool for
Drosophila developmental studies. *Genetics* **175**, 1505-1531.
Cautain, B., Hill, R., de Pedro, N. and Link, W. (2015). Components and regulation of nuclear transport
processes. *The FEBS journal* **282**, 445-462.
Cesario, J. and McKim, K. S. (2011). RanGTP is required for meiotic spindle organization and the
initiation of embryonic development in Drosophila. *J Cell Sci* **124**, 3797-3810.
Chaigne, A., Labouesse, C., White, I. J., Agnew, M., Hannezo, E., Chalut, K. J. and Paluch, E. K. (2020).
Abscission Couples Cell Division to Embryonic Stem Cell Fate. *Developmental cell* **55**, 195-
208.e195.
Chang, C. C. and Hsia, K. C. (2020). More than a zip code: global modulation of cellular function by
nuclear localization signals. *The FEBS journal*.
Chau, J., Kulnane, L. S. and Salz, H. K. (2009). Sex-lethal facilitates the transition from germline stem cell
to committed daughter cell in the Drosophila ovary. *Genetics* **182**, 121-132.
Chen, A., Akhshi, T. K., Lavoie, B. D. and Wilde, A. (2015). Importin β2 Mediates the Spatio-temporal
Regulation of Anillin through a Noncanonical Nuclear Localization Signal. *The Journal of*
biological chemistry **290**, 13500-13509.
Chen, D., Wang, Q., Huang, H., Xia, L., Jiang, X., Kan, L., Sun, Q. and Chen, D. (2009). Effete-mediated
degradation of Cyclin A is essential for the maintenance of germline stem cells in Drosophila.
Development **136**, 4133-4142.
Chen, J., Call, G. B., Beyer, E., Bui, C., Cespedes, A., Chan, A., Chan, J., Chan, S., Chhabra, A., Dang, P.,
et al. (2005). Discovery-based science education: functional genomic dissection in Drosophila by
undergraduate researchers. *PLoS Biol* **3**, e59.
Cheng, J., Türkel, N., Hemati, N., Fuller, M. T., Hunt, A. J. and Yamashita, Y. M. (2008). Centrosome
misorientation reduces stem cell division during ageing. *Nature* **456**, 599-604.
de Cuevas, M., Lee, J. K. and Spradling, A. C. (1996). alpha-spectrin is required for germline cell division
and differentiation in the Drosophila ovary. *Development* **122**, 3959-3968.
de Cuevas, M., Lilly, M. A. and Spradling, A. C. (1997). Germline cyst formation in Drosophila. *Annual*
review of genetics **31**, 405-428.
de Cuevas, M. and Spradling, A. C. (1998). Morphogenesis of the Drosophila fusome and its implications
for oocyte specification. *Development* **125**, 2781-2789.

Deng, W. and Lin, H. (1997). Spectrosomes and fusomes anchor mitotic spindles during asymmetric
germ cell divisions and facilitate the formation of a polarized microtubule array for oocyte
specification in Drosophila. *Developmental biology* **189**, 79-94.
Drutovic, D., Duan, X., Li, R., Kalab, P. and Solc, P. (2020). RanGTP and importin β regulate meiosis I
spindle assembly and function in mouse oocytes. *The EMBO journal* **39**, e101689.
Eastin, K. J., Huang, A. P. and Ferree, P. M. (2020). A novel pattern of germ cell divisions in the
production of hymenopteran insect eggs. *Biol Lett* **16**, 20200137.
**Eikenes Å, H., Malerød, L., Lie-Jensen, A., Sem Wegner, C., Brech, A., Liestøl, K., Stenmark, H. and
Haglund, K.** (2015). Src64 controls a novel actin network required for proper ring canal
formation in the Drosophila male germline. *Development* **142**, 4107-4118.
Fichelson, P., Jagut, M., Lepanse, S., Lepesant, J. A. and Huynh, J. R. (2010). lethal giant larvae is
required with the par genes for the early polarization of the Drosophila oocyte. *Development*
137, 815-824.
Forbes, D. J., Travesa, A., Nord, M. S. and Bernis, C. (2015). Nuclear transport factors: global regulation
of mitosis. *Current opinion in cell biology* **35**, 78-90.
Frémont, S. and Echard, A. (2018). Membrane Traffic in the Late Steps of Cytokinesis. *Current biology :
CB* **28**, R458-R470.
Geles, K. G. and Adam, S. A. (2001). Germline and developmental roles of the nuclear transport factor
importin alpha3 in C. elegans. *Development* **128**, 1817-1830.
Goldbach, P., Wong, R., Beise, N., Sarpal, R., Trimble, W. S. and Brill, J. A. (2010). Stabilization of the
actomyosin ring enables spermatocyte cytokinesis in Drosophila. *Molecular biology of the cell*
21, 1482-1493.
Görlich, D., Dabrowski, M., Bischoff, F. R., Kutay, U., Bork, P., Hartmann, E., Prehn, S. and Izaurralde, E.
(1997). A novel class of RanGTP binding proteins. *J Cell Biol* **138**, 65-80.
Grieder, N. C., de Cuevas, M. and Spradling, A. C. (2000). The fusome organizes the microtubule
network during oocyte differentiation in Drosophila. *Development* **127**, 4253-4264.
Guertin, D. A., Trautmann, S. and McCollum, D. (2002). Cytokinesis in eukaryotes. *Microbiology and
molecular biology reviews : MMBR* **66**, 155-178.
Haglund, K., Nezis, I. P. and Stenmark, H. (2011). Structure and functions of stable intercellular bridges
formed by incomplete cytokinesis during development. *Communicative & integrative biology* **4**,
1-9.
Hinnant, T. D., Alvarez, A. A. and Ables, E. T. (2017). Temporal remodeling of the cell cycle accompanies
differentiation in the Drosophila germline. *Developmental biology* **429**, 118-131.
Hinnant, T. D., Merkle, J. A. and Ables, E. T. (2020). Coordinating Proliferation, Polarity, and Cell Fate in
the Drosophila Female Germline. *Frontiers in cell and developmental biology* **8**, 19.
Hogarth, C., Itman, C., Jans, D. A. and Loveland, K. L. (2005). Regulated nucleocytoplasmic transport in
spermatogenesis: a driver of cellular differentiation? *BioEssays : news and reviews in molecular,
cellular and developmental biology* **27**, 1011-1025.
Hughes, S. E., Miller, D. E., Miller, A. L. and Hawley, R. S. (2018). Female Meiosis: Synapsis,
Recombination, and Segregation in Drosophila melanogaster. *Genetics* **208**, 875-908.
Huynh, J.-R. (2006). Fusome as a Cell-Cell Communication Channel of Drosophila Ovarian Cyst. In *Cell-
Cell Channels* (ed. F. Baluska, D. Volkmann & P. W. Barlow), pp. 217-235. New York, NY: Springer
New York.
Ji, S., Li, C., Hu, L., Liu, K., Mei, J., Luo, Y., Tao, Y., Xia, Z., Sun, Q. and Chen, D. (2017). Bam-dependent
deubiquitinase complex can disrupt germ-line stem cell maintenance by targeting cyclin A.
Proceedings of the National Academy of Sciences of the United States of America **114**, 6316-
6321.

Kimura, M. and Imamoto, N. (2014). Biological significance of the importin-β family-dependent
nucleocytoplasmic transport pathways. *Traffic (Copenhagen, Denmark)* **15**, 727-748.
Kimura, M., Morinaka, Y., Imai, K., Kose, S., Horton, P. and Imamoto, N. (2017). Extensive cargo
identification reveals distinct biological roles of the 12 importin pathways. *eLife* **6**.
King, R. C. R. C. (1970). Ovarian development in Drosophila melanogaster. New York: Academic Press.
Koch, E. A. and King, R. C. (1966). The origin and early differentiation of the egg chamber of Drosophila
melanogaster. *J Morphol* **119**, 283-303.
Koch, E. A., Smith, P. A. and King, R. C. (1967). The division and differentiation of Drosophila cystocytes.
J Morphol **121**, 55-70.
Kosyna, F. K. and Depping, R. (2018). Controlling the Gatekeeper: Therapeutic Targeting of Nuclear
Transport. *Cells* **7**.
Kubrakiewicz, J. (1997). Germ cells cluster organization in polytrophic ovaries of neuroptera. *Tissue and*
Cell **29**, 221-228.
Lai, M. C., Lin, R. I. and Tarn, W. Y. (2001). Transportin-SR2 mediates nuclear import of phosphorylated
SR proteins. *Proc Natl Acad Sci U S A* **98**, 10154-10159.
Laws, K. M. and Drummond-Barbosa, D. (2015). Genetic Mosaic Analysis of Stem Cell Lineages in the
Drosophila Ovary. *Methods in molecular biology (Clifton, N.J.)* **1328**, 57-72.
Lighthouse, D. V., Buszczak, M. and Spradling, A. C. (2008). New components of the Drosophila fusome
suggest it plays novel roles in signaling and transport. *Developmental biology* **317**, 59-71.
Lilly, M. A., de Cuevas, M. and Spradling, A. C. (2000). Cyclin A associates with the fusome during
germline cyst formation in the Drosophila ovary. *Developmental biology* **218**, 53-63.
Lin, H. and Spradling, A. C. (1995). Fusome asymmetry and oocyte determination in Drosophila. *Dev*
Genet **16**, 6-12.
Lin, H., Yue, L. and Spradling, A. C. (1994). The Drosophila fusome, a germline-specific organelle,
contains membrane skeletal proteins and functions in cyst formation. *Development* **120**, 947-
956.
Lippai, M., Tirián, L., Boros, I., Mihály, J., Erdélyi, M., Belecz, I., Máthé, E., Pósfai, J., Nagy, A., Udvardy,
A., et al. (2000). The Ketel gene encodes a Drosophila homologue of importin-beta. *Genetics*
156, 1889-1900.
Liu, L., Michowski, W., Kolodziejczyk, A. and Sicinski, P. (2019). The cell cycle in stem cell proliferation,
pluripotency and differentiation. *Nature cell biology* **21**, 1060-1067.
Loveland, K. L., Major, A. T., Butler, R., Young, J. C., Jans, D. A. and Miyamoto, Y. (2015). Putting things
in place for fertilization: discovering roles for importin proteins in cell fate and spermatogenesis.
Asian journal of andrology **17**, 537-544.
Lu, K., Jensen, L., Lei, L. and Yamashita, Y. M. (2017). Stay Connected: A Germ Cell Strategy. *Trends in*
genetics : TIG **33**, 971-978.
Maertens, G. N., Cook, N. J., Wang, W., Hare, S., Gupta, S. S., Öztop, I., Lee, K., Pye, V. E., Cosnefroy,
O., Snijders, A. P., et al. (2014). Structural basis for nuclear import of splicing factors by human
Transportin 3. *Proc Natl Acad Sci U S A* **111**, 2728-2733.
Mahowald, A. P. (1971). The formation of ring canals by cell furrows in Drosophila. *Z Zellforsch Mikrosk*
Anat **118**, 162-167.
Mason, D. A., Fleming, R. J. and Goldfarb, D. S. (2002). Drosophila melanogaster importin alpha1 and
alpha3 can replace importin alpha2 during spermatogenesis but not oogenesis. *Genetics* **161**,
157-170.
Máthé, E., Bates, H., Huikeshoven, H., Deák, P., Glover, D. M. and Cotterill, S. (2000). Importin-alpha3
is required at multiple stages of Drosophila development and has a role in the completion of
oogenesis. *Developmental biology* **223**, 307-322.

Máthé, E., Inoue, Y. H., Palframan, W., Brown, G. and Glover, D. M. (2003). Orbit/Mast, the CLASP orthologue of Drosophila, is required for asymmetric stem cell and cystocyte divisions and development of the polarised microtubule network that interconnects oocyte and nurse cells during oogenesis. *Development* **130**, 901-915.

Mathieu, J., Cauvin, C., Moch, C., Radford, S. J., Sampaio, P., Perdigoto, C. N., Schweisguth, F., Bardin, A. J., Sunkel, C. E., McKim, K., et al. (2013). Aurora B and cyclin B have opposite effects on the timing of cytokinesis abscission in Drosophila germ cells and in vertebrate somatic cells. *Developmental cell* **26**, 250-265.

Mathieu, J. and Huynh, J. R. (2017). Monitoring complete and incomplete abscission in the germ line stem cell lineage of Drosophila ovaries. *Methods in cell biology* **137**, 105-118.

Matias, N. R., Mathieu, J. and Huynh, J. R. (2015). Abscission is regulated by the ESCRT-III protein shrub in Drosophila germline stem cells. *PLoS genetics* **11**, e1004653.

Matova, N. and Cooley, L. (2001). Comparative aspects of animal oogenesis. *Developmental biology* **231**, 291-320.

McKearin, D. (1997). The Drosophila fusome, organelle biogenesis and germ cell differentiation: if you build it. *BioEssays : news and reviews in molecular, cellular and developmental biology* **19**, 147-152.

McKearin, D. and Ohlstein, B. (1995). A role for the Drosophila bag-of-marbles protein in the differentiation of cystoblasts from germline stem cells. *Development* **121**, 2937-2947.

Megraw, T. L. and Kaufman, T. C. (2000). The centrosome in Drosophila oocyte development. *Current topics in developmental biology* **49**, 385-407.

Mihalas, B. P., Western, P. S., Loveland, K. L., McLaughlin, E. A. and Holt, J. E. (2015). Changing expression and subcellular distribution of karyopherins during murine oogenesis. *Reproduction (Cambridge, England)* **150**, 485-496.

Morris, J. Z., Hong, A., Lilly, M. A. and Lehmann, R. (2005). twin, a CCR4 homolog, regulates cyclin poly(A) tail length to permit Drosophila oogenesis. *Development* **132**, 1165-1174.

Nguyen, L. T. S. and Robinson, D. N. (2020). The Unusual Suspects in Cytokinesis: Fitting the Pieces Together. *Frontiers in cell and developmental biology* **8**, 441.

Ni, J. Q., Zhou, R., Czech, B., Liu, L. P., Holderbaum, L., Yang-Zhou, D., Shim, H. S., Tao, R., Handler, D., Karpowicz, P., et al. (2011). A genome-scale shRNA resource for transgenic RNAi in Drosophila. *Nat Methods* **8**, 405-407.

Ohlmeyer, J. T. and Schüpbach, T. (2003). Encore facilitates SCF-Ubiquitin-proteasome-dependent proteolysis during Drosophila oogenesis. *Development* **130**, 6339-6349.

Okada, N., Ishigami, Y., Suzuki, T., Kaneko, A., Yasui, K., Fukutomi, R. and Isemura, M. (2008). Importins and exportins in cellular differentiation. *J Cell Mol Med* **12**, 1863-1871.

Ong, S. and Tan, C. (2010). Germline cyst formation and incomplete cytokinesis during Drosophila melanogaster oogenesis. *Developmental biology* **337**, 84-98.

Ozugergin, I. and Piekny, A. (2020). Complementary functions for the Ran gradient during division. *Small GTPases*, 1-11.

Pan, L., Chen, S., Weng, C., Call, G., Zhu, D., Tang, H., Zhang, N. and Xie, T. (2007). Stem cell aging is controlled both intrinsically and extrinsically in the Drosophila ovary. *Cell Stem Cell* **1**, 458-469.

Pepling, M. and Lei, L. (2018). Germ Cell Nests and Germline Cysts. In *Encyclopedia of Reproduction (Second Edition)* (ed. M. K. Skinner), pp. 159-166. Oxford: Academic Press.

Pepling, M. E., de Cuevas, M. and Spradling, A. C. (1999). Germline cysts: a conserved phase of germ cell development? *Trends in Cell Biology* **9**, 257-262.

Perkins, L. A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., Yang-Zhou, D., Flockhart, I., Binari, R., Shim, H. S., et al. (2015). The Transgenic RNAi Project at Harvard Medical School: Resources and Validation. *Genetics* **201**, 843-852.

Petrella, L. N., Smith-Leiker, T. and Cooley, L. (2007). The Ovhts polyprotein is cleaved to produce
fusome and ring canal proteins required for Drosophila oogenesis. *Development* **134**, 703-712.
Piekny, A. J. and Maddox, A. S. (2010). The myriad roles of Anillin during cytokinesis. *Seminars in cell &*
developmental biology **21**, 881-891.
Pollard, T. D. and O'Shaughnessy, B. (2019). Molecular Mechanism of Cytokinesis. *Annu Rev Biochem*
88, 661-689.
Robinson, D. N., Cant, K. and Cooley, L. (1994). Morphogenesis of Drosophila ovarian ring canals.
Development **120**, 2015-2025.
Röper, K. (2007). Rtnl1 is enriched in a specialized germline ER that associates with ribonucleoprotein
granule components. *J Cell Sci* **120**, 1081-1092.
Rørth, P. (1998). Gal4 in the Drosophila female germline. *Mech Dev* **78**, 113-118.
Sazer, S. and Dasso, M. (2000). The ran decathlon: multiple roles of Ran. *J Cell Sci* **113 (Pt 7)**, 1111-1118.
Schuldiner, O., Berdnik, D., Levy, J. M., Wu, J. S., Luginbuhl, D., Gontang, A. C. and Luo, L. (2008).
piggyBac-based mosaic screen identifies a postmitotic function for cohesin in regulating
developmental axon pruning. *Developmental cell* **14**, 227-238.
Silverman-Gavrila, R. V., Hales, K. G. and Wilde, A. (2008). Anillin-mediated targeting of peanut to
pseudocleavage furrows is regulated by the GTPase Ran. *Molecular biology of the cell* **19**, 3735-
3744.
Snapp, E. L., Iida, T., Frescas, D., Lippincott-Schwartz, J. and Lilly, M. A. (2004). The fusome mediates
intercellular endoplasmic reticulum connectivity in Drosophila ovarian cysts. *Molecular biology*
of the cell **15**, 4512-4521.
Storto, P. D. and King, R. C. (1989). The role of polyfusomes in generating branched chains of cystocytes
during Drosophila oogenesis. *Dev Genet* **10**, 70-86.
Sugimura, I. and Lilly, M. A. (2006). Bruno inhibits the expression of mitotic cyclins during the prophase
I meiotic arrest of Drosophila oocytes. *Developmental cell* **10**, 127-135.
Teixeira, F. K. and Lehmann, R. (2019). Translational Control during Developmental Transitions. *Cold*
Spring Harb Perspect Biol **11**.
Thurmond, J., Goodman, J. L., Strelets, V. B., Attrill, H., Gramates, L. S., Marygold, S. J., Matthews, B.
B., Millburn, G., Antonazzo, G., Trovisco, V., et al. (2019). FlyBase 2.0: the next generation.
Nucleic Acids Res **47**, D759-d765.
Tirián, L., Puro, J., Erdélyi, M., Boros, I., Papp, B., Lippai, M. and Szabad, J. (2000). The Ketel(D)
dominant-negative mutations identify maternal function of the Drosophila importin-beta gene
required for cleavage nuclei formation. *Genetics* **156**, 1901-1912.
Trieselmann, N. and Wilde, A. (2002). Ran localizes around the microtubule spindle in vivo during
mitosis in Drosophila embryos. *Current biology : CB* **12**, 1124-1129.
Van Doren, M., Williamson, A. L. and Lehmann, R. (1998). Regulation of zygotic gene expression in
Drosophila primordial germ cells. *Current biology : CB* **8**, 243-246.
Villányi, Z., Debec, A., Timinszky, G., Tirián, L. and Szabad, J. (2008). Long persistence of importin-beta
explains extended survival of cells and zygotes that lack the encoding gene. *Mech Dev* **125**, 196-
206.
Xu, T. and Rubin, G. M. (1993). Analysis of genetic mosaics in developing and adult Drosophila tissues.
Development **117**, 1223-1237.
Yamashita, Y. M. (2018). Subcellular Specialization and Organelle Behavior in Germ Cells. *Genetics* **208**,
19-51.
Yue, L. and Spradling, A. C. (1992). hu-li tai shao, a gene required for ring canal formation during
Drosophila oogenesis, encodes a homolog of adducin. *Genes & development* **6**, 2443-2454.

Zielke, N., Korzelius, J., van Straaten, M., Bender, K., Schuhknecht, G. F. P., Dutta, D., Xiang, J. and
Edgar, B. A. (2014). Fly-FUCCI: A versatile tool for studying cell proliferation in complex tissues.
Cell Rep **7**, 588-598.

TABLES

Table 1. Frequency of egg chambers that do not follow the power of two rule in β-importin- and
Ran-depleted ovaries.

Genotype	% of egg chambers that do not follow 2^n germ cells
Driver control	0
$Fs(2)Ket^{GL01273}$	1.3
$Ran^{GL01341}$	2.6
$cdm^{HMS02846}$	55.0
$Tnpo\text{-}SR^{V22}$	29.3
$Ran^{HA\ (ins\ 1)}$	0
$Ran^{HA\ (ins\ 2)}$	0
$Ran^{T24N\ (ins\ 1)}$	12.5
$Ran^{T24N\ (ins\ 2)}$	9.7
$Ran^{Q69L\ (ins\ 1)}$	0
$Ran^{Q69L\ (ins\ 2)}$	2.2

FIGURES

Figure 1. An RNAi screen identified importins essential for germ cell development.

(A) The *Drosophila* ovary is composed of 14-16 ovarioles, each harboring a germarium (top) and

older follicle cells that progress through 14 distinct stages of development (bottom). Within the

germarium, germline stem cells (GSCs; pink) are adjacent to cap cells (yellow) and a population

of escort cells (green). GSCs divide to form daughter cells (cystoblasts; blue) that go on to divide

four additional times to form 16-celled germline cysts (blue). Each cyst is composed of 15 nurse cells and one oocyte (dark blue), connected by 15 intercellular bridges that can be visualized by ring canals and the endoplasmic reticulum-like fusome (red) that transects each bridge. (B-D) Maximum intensity projections of *nos-Gal4* control (B), *nos-Gal4>UAS-* $msk^{HMS00020}$ (C), or *nos-Gal4>UAS-cdm^{HMS02846}* (D) knockdown ovaries labeled with DAPI (white; nuclei). Scale bar, 100 μm. In control ovaries, each ovariole produces germline cysts that are packaged into individual egg chambers. As egg chambers develop, oocytes grow and accumulate yolk, while nurse cells become polyploid, adopting a stereotypical nuclear morphology. In *nos-Gal4>UAS-msk^{HMS00020}* ovaries, ovarioles are present but completely lack germ cells. In *nos-Gal4>UAS-cdm^{HMS02846}* ovaries, ovarioles have a partial agametic phenotype, where germ cells are present in some ovarioles but ovaries are dramatically smaller than wild-type. Arrows indicate oviducts (ovi). (E) Frequencies of ovarioles containing agametic (red) ovarioles. The number of ovarioles analyzed is shown inside bars.

Figure 2. Ran and β-importins regulate the number of germ cells per egg chamber.

(A-C) Egg chambers of driver control (A) or *nos-Gal4* driving RNAi against *Tnpo-SR* (B) and *cdm* (C) ovarioles labeled with anti-Hts-RC (yellow; ring canals), anti-Hts (red; fusomes and follicle cell membranes), anti-LamC (red; nuclear envelopes), and DAPI (blue; nuclei). (D) Frequencies of egg chambers containing less than 16 (grey), 16 (white), or more than 16 (black) germ cells per egg chamber in *nos-Gal4* control or *nos-Gal4* driving *Ran*, *cdm*, *Tnpo-SR*, or *Ketel* RNAi lines at 6 dae. (E) Variation of the number of germ cells per egg chambers. The numbers in/above the bars represent the total number of egg chambers analyzed. (F-H) Germaria from control (F), *nos-Gal4>Tnpo-SR^{V22}* (G), and *nos-Gal4>cdm^{HMS02846}* RNAi (H) females labeled with anti-Orb (yellow; oocyte), anti-Hts and anti-LamC (red), and DAPI (blue). Values in bottom left corner of G-H represent the number of ovarioles containing germline tumors over the total number of ovarioles analyzed. A-C and G are 10 μm subset maximum intensity projections. Scale bar, 20 μm. Error bars, mean ± SEM. ***$p < 0.001$; Chi-squares test.

Figure 3. *Ran* and *Tnpo-SR* are required for proper oocyte selection.

(A-C) Driver control (A), *nosGal4>Tnpo-SR^{V22}* (B), and *nosGal4>Ran^{GL01341}* egg chambers labeled with anti-Orb (yellow; oocyte), anti-Hts (red; fusomes and follicle cell membranes), anti-LamC (red; nuclear envelopes), and DAPI (blue; nuclei). (D-F) Driver control (D) and *nosGal4>Tnpo-SR^{V22}* (E-F) egg chambers labeled with anti-BicD (yellow; oocyte), anti-Hts and anti-LamC (red), and DAPI (blue). Arrowheads indicate oocytes, recognized by their condensed nuclei. (G) Frequencies of egg chambers containing 0 (grey), 1 (white), or 2 (black) oocytes. The numbers in the bars represent the total number of egg chambers analyzed. (H-J) *Tnpo-SR^{KG04870}* mutant mosaic germarium at 8 days after clone induction labeled with anti-GFP (green; wild-type cells), anti-Hts and anti-LamC (blue), and anti-Orb (red) (H-H') or anti-C(3)G (red; synaptonemal complexes) (J-J'). White lines demarcate wild-type (solid) or mutant (dashed) cysts. (I) Percentage of GFP-positive (wild-type) or GFP-negative (*Tnpo-SR^{KG04870}*) 16-cell cysts with an Orb-positive

995   oocyte at 8 days after clone induction. (K) Average number of C(3)G-positive germ cells in GFP-

996   positive (wild-type) or GFP-negative (*Tnpo-SR*^{KG04870}) 8- or 16-cell cysts at 8 days after clone

997   induction. Numbers in the bars represent the total number of cysts analyzed. Scale bar, 20 µm (A-

998   F) and 5 µm (H-J). Error bars, mean ± SEM. ***$p < 0.001$; Student's two-tailed T-test.

Figure 4. Ran function in dividing germ cells requires Ran-GTP activity.

(A-A') Single z-plane image of a *nos-Gal4>Ran^{HA}* germarium labeled with anti-HA (green; Ran protein), anti-Tubulin (red; mitotic spindles), anti-Hts (white; fusomes and follicle cell membranes), anti-LamC (white; nuclear envelopes), and DAPI (blue; nuclei). The panel in A' is the HA-only image of the germarium at left. (B) Maximum intensity projection (10 μm subset) of a germarium from a *nos-Gal4>Ran^{T24N}* ovary immunostained with anti-Hts and anti-LamC (red), and DAPI (blue). (C) Maximum intensity projection (5 μm subset) representative *Ran^{T24N}* germarium stained with anti-Hts-RC (red; ring canals), anti-Hts and anti-LamC (green), and DAPI (blue). (D-E) Representative *Ran^{T24N}* (D, GDP-locked mutant) and *Ran^{Q69L}* (GTP-locked mutant) labeled with anti-Orb (yellow; oocyte), anti-Hts and anti-LamC (red), and DAPI (blue). Arrowheads indicate oocytes, as recognized by their condensed nuclei. (F) Frequencies of agametic (red) or non-agametic (grey) ovarioles in *nos-Gal4* control, *Ran* wild-type, and *Ran* dominant negative lines. Numbers in/above bars represent the total number of ovarioles analyzed. (G) Average number of GSCs at 3, 6, and 10 days after eclosion (dae). Dots indicate average number of GSCs at the point analyzed. More than 15 germaria per genotype at each timepoint were analyzed. (H) Frequencies of egg chambers containing less than 16 (grey), 16 (white), or more than 16 (black) germ cells per egg chamber in *nos-Gal4* control or *nos-Gal4* driving *Ran* wild-type and dominant negative transgenes at 6 dae. (I) Variation of the number of germ cells present per egg chamber. Numbers in/above bars represent the total number of egg chambers analyzed. (J) Frequencies of 16-cell egg chambers containing 0 (grey), 1 (white), or 2 (black) oocytes. In panels F, H, and J, the y-axis labels are identical. In panels H-J, numbers in/above bars represent the total number of egg chambers analyzed. Scale bars, 10 μm (A-A') and 20 μm (B-E). Error bars, mean ± SEM. **$p < 0.01$; ***$p < 0.001$; Student's two-tailed T-test and Chi-squared tests.

Figure 5. *Tnpo-SR* and *cdm* are required for fusome integrity and maturation.

(A-D) Maximum intensity projections of representative driver control (A; 3 µm), *cdm*^{HMS02846} (B; 3 µm), *Tnpo-SR*^{V22} (C; 7 µm), and *Ran*^{T24N} (D; 7 µm) mutant cysts labeled with anti-tubulin (red; mitotic spindles) anti-Hts (green; fusomes and follicle cell membranes, anti-LamC (green; nuclear envelope of cap cells), and DAPI (blue; nuclei). (E-H) Maximum intensity projections of representative driver control (E; 10 µm), *cdm*^{HMS02846} (F; 5 µm), *Tnpo-SR*^{V22} (G; 10 µm), and *Ran*^{T24N} (H; 5 µm) mutant germaria labeled with anti-Hts (red; fusomes and follicle cell membranes, anti-LamC (red; nuclear envelope of cap cells), and DAPI (blue; nuclei). Arrowheads indicate points of the fusome that are thinner than those seen in the control cysts. (I-J) Maximum intensity projections of representative *Scrib-GFP* control (I; 20 µm) and *Tnpo-SR*^{V22}*;Scrib-GFP* mutant (J; 20 µm) germaria labeled with anti-GFP (green), anti-Hts (red; fusomes and follicle cell membranes, and anti-LamC (red; nuclear envelope of cap cells). (K-L) Maximum intensity projections of representative *Rtnl1-GFP* control (K; 10 µm) and *Tnpo-SR*^{V22}*,Rtnl1-GFP* mutant

1039   (L; 10 µm) germaria labeled with anti-GFP (green), anti-Hts (red; fusomes and follicle cell

1040   membranes, and anti-LamC (red; nuclear envelope of cap cells). Arrowheads indicate highest level

1041   of expression of the Scrib (I-J) and Rtnl1 (K-L). Scale bar, 10 µm.

1042

Figure 6. Depletion of *Tnpo-SR* and *cdm*, but not Ran-GTP, delays ring canal protein deposition and fails to suppress cytokinesis in dividing cysts.

(A) *Tnpo-SR^{KG04870}* mutant mosaic germarium labeled with anti-GFP (green; wild-type cells), anti-pTyrosine (pTyr, red; ring canals), anti-Hts (blue; fusomes and follicle cell membranes, and anti-LamC (blue; nuclear envelope of cap cells). (B) Maximum intensity projection of a *Tnpo-SR^{KG04870}* (5 µm subset) mutant mosaic germarium labeled with anti-GFP (green; wild-type cells) and anti-E-cadherin (red). Arrows indicate E-cadherin connections. (C-D) Representative control (C) and

Tnpo-SR^{V22} (D) mutant germaria labeled with Hts-RC (red; ring canals), anti-Hts (green; fusomes

and follicle cell membranes, anti-LamC (green; nuclear envelope of cap cells), and DAPI (blue;

nuclei). (E-H) Maximum intensity projections (5 µm subsets) of representative driver control (E),

cdm^{HMS02846} (F), *Tnpo-SR^{V22}* (G), and *Ran^{T24N}* (H) mutant cysts labeled with anti-pTyrosine (red;

ring canals), anti-Hts (green; fusomes and follicle cell membranes), and anti-LamC (green; nuclear

envelope of cap cells). Closed arrowheads indicate ring canals and open ring canals indicate a cyst

that is lacking any remnants of ring canals. (I-L) Representative control (I), *Tnpo-SR^{V22}* mutant (J)

and *Ran^{T24N}* mutant (K-L) germarium labeled with anti-anillin (green) anti-Hts (red; fusomes and

follicle cell membranes), and anti-LamC (red; nuclear envelope of cap cells). Arrowheads indicate

closed ring canals. Scale bar, 10 µm. (M) Summary of fusome and ring canal phenotypes in *Tnpo-*

SR- and *cdm*-depleted germ cell cysts.

1062

1063

1064

Figure 7. CycA is ectopically expressed in c*dm* and *Tnpo-SR* cysts.

(A-D) Maximum intensity projections (10 µm subsets) of representative driver control (A), *Ran^GL01341* (B), *cdm^HMS02846* (C), and *Tnpo-SR^V22* (D) mutant germaria labeled with anti-CycA (red), anti-Hts (green; fusomes and follicle cell membranes, and anti-LamC (green; nuclear envelope of cap cells). (E) Maximum intensity projection of *nos-Gal4>FUCCI;Tnpo-SR^V22* labeled with anti-GFP (green; E2f1), anti-dsRed (red; CycB), and anti-Hts (white; fusomes and follicle cell membranes, and anti-LamC (white; nuclear envelope of cap cells). Scale bar, 10 µm. (F) Percentage of cells/cysts containing fusome-localized CycA. (G) Percentage of cells/cysts containing *RFP::CycB* in *nos-Gal4* controls and *Tnpo-SR^V22* knockdown germaria. The number of cells/cysts analyzed is shown above bars. ***$p < 0.001$, Student's Chi-squared test.

1077 **SUPPLEMENTAL DATA**

1078

Supp. Figure 1. Ran and β-importins are necessary for GSC maintenance.

(A-B) Max intensity projections (10 μm subset) of *nos-Gal4* control (A) and *nos-Gal4>UAS-*

Ran^{GL01341} (B) germaria labeled with anti-orb (yellow; oocyte), anti-Hts (red; fusomes and follicle

cell membranes, anti-LamC (red; nuclear envelope of cap cells) and DAPI (blue; nuclei). Solid

white lines demarcate GSCs present in the germarium, asterisks indicate cap cells. (C) Average

number of GSCs per germarium in *nos-Gal4* control or *nos-Gal4>UAS-Ran/importin RNAi* ovaries

at 3, 6, and 10 days after eclosion (dae). At least 30 germaria were analyzed at each timepoint. (D-

E) Single z-plane confocal micrographs of *FRT40A* control (D) and *Tnpo-SR^{KG04870} FRT40A* mutant (E) mosaic germaria at 8 days after clone induction labeled with anti-GFP (green; wild-type cells), anti-Hts (red; fusomes and follicle cell membranes), anti-LamC (red; nuclear envelopes) and DAPI (blue; nuclei). Lines demarcate wild-type (solid) or mutant (dashed) GSCs (white) and cystoblast/cysts (yellow). (F) Percentage of total germaria analyzed with a GFP-negative GSC at 4, 8, and 12 days after clone induction. At least 145 germaria were analyzed. (G-G') *Tnpo-SR^{KG04870}* mutant mosaic germaria at 8 days after clone induction labeled with anti-GFP (green; wild-type cells), anti-Hts and anti-LamC (red), and anti-E-cadherin (blue). Lines demarcate wild-type (solid) or mutant (dashed) GSCs. Image in G' is the E-cadherin channel only of the image in G. (H) Average intensity mean value (IMV) of E-cadherin at the GSC-cap cell junction of GFP-positive (wild-type) or GFP-negative (*Tnpo-SR^{KG04870}* mutant) GSCs at 8 days after clone induction. Numbers in the bars represent the total number of GSCs analyzed. (I-J) Max intensity projections (2 μm subset) of *nos-Gal4* control (I) and *nos-Gal4>Tnpo-SR^{V22}* (J) GSC/CB pairs labeled with anti-pTyrosine (red; ring canals), anti-Hts and anti-LamC (green), and DAPI (blue). Arrowheads indicate pTyr+ plug that forms between the GSC/CB pair prior to abscission. (K-L) Max intensity projections (10 μm subset) of *nos-Gal4* control (K) and *nos-Gal4>Tnpo-SR^{V22}* (L) GSC/CB pairs labeled with anti-Edu (magenta; S-phase marker), anti-Hts and anti-LamC (green). Scale bars, 10 μm. Error bars, mean ± SEM. ***$p < 0.001$; Student's two-tailed T-test.

1104

Supp. Figure 2. *Tnpo-SR* germline tumors are primarily composed of mitotically-dividing

undifferentiated germ cells.

(A) Maximum intensity projection (20 μm subset) of a *nos-Gal4>Tnpo-SR^{V22}* egg chamber labeled

with anti-Hts-RC (red; ring canals), anti-Hts (green; fusomes and follicle cell membranes), anti-

LamC (green; nuclear envelopes), and DAPI (blue; nuclei). (B) Maximum intensity projection (20

μm subset) of a *nos-Gal4>Tnpo-SR^{V22}* egg chamber labeled with anti-pTyr (red; ring canals), anti-

Hts and anti-LamC (green), and DAPI (blue). (C) Maximum intensity projection (10 μm subset)

of a *nos-Gal4>Tnpo-SR^{V22}* egg chamber labeled with anti-pHH3 (green; M-phase marker), anti-

Hts and anti-LamC (red), and DAPI (blue). Panel in C' depicts pHH3 channel only. Arrowheads

indicate pHH3-positive cells within the germline tumor. (D-E) Representative control (D) and *nos-*

1115   *Gal4>Tnpo-SR^{V22}* (E) germaria labeled with anti-Sxl (red), anti-Hts and anti-LamC (green), and

1116   DAPI (blue). Panels D' and E' show heat maps corresponding to panels D and E, where cool colors

1117   are low expression and warm colors are high expression. Scale bars, 20 µm.

1118

Supp. Figure 3. *Tnpo-SR* mutants fail to maintain cyst connectivity and some cells are ectopically mitotically active.

(A) *Tnpo-SR^{KG04870}* mutant egg chamber labeled with anti-GFP (green; wild-type cells), anti-Hts (red; fusomes and follicle cell membranes), anti-LamC (red; nuclear envelopes), and DAPI (blue; nuclei). White lines indicate wild-type cysts (solid) and mutant cysts (dashed). (B) *Tnpo-SR^{KG04870}* mutant egg chamber labeled with anti-GFP (green; wild-type cells), anti-pHH3 (white; M-phase), anti-Hts (red; fusomes and follicle cell membranes), anti-LamC (red; nuclear envelopes), and DAPI (blue; nuclei). (C-E''') *Tnpo-SR^{LL05552}* (C and E-E''') and *Tnpo-SR^{KG04870}* (D) mutant germaria labeled with anti-GFP (green; wild-type cells), anti-Hts (red; fusomes and follicle cell membranes), anti-LamC (red; nuclear envelopes), and DAPI (blue; nuclei). Closed arrowhead represents thin thread of fusome material and open arrowheads follow the GFP+ follicle cell projections invaginating the mutant cysts. (D) is a 2 μm subset maximum intensity projection. Scale bar, 10 μm.

1134

Supp. Figure 4. An Anillin-NLS mutant is not efficient in disrupting anillin localization to nuclei and ring canals.

(A-D) Representative controls of *UAS-Scra-GFP* (A and C) and NLS mutated *UAS-Scra-3A-GFP* mutant (B and D) germaria labeled with anti-GFP (green), anti-Hts (red; fusomes and follicle cell membranes), and anti-LamC (red; nuclear envelope of cap cells), and DAPI (blue nuclei) (A-B) or anti-pTyrosine (red; ring canals) (C-D). Arrowheads indicate closed ring canals. Scale bar, 10 µm.

1141

1142 **Supp. Table S1.** Fly stocks and immunofluorescence reagents used in this study.

Reagent	Source	Reference
nos-GAL4::VP16-nos.UTR	D. Drummond-Barbosa	(Rørth, 1998; Van Doren et al., 1998)
w*; nosGal4::VP16, UASp-tubGFP	BLM #7253	(Grieder et al., 2000)
Tnpo-SR^{V22}	Ables Lab	this study
y^{1}v^1; P{y[+t7.7] v[+t1.8]=TRiP.GL01483}attP2	BLM #43142	(Perkins et al., 2015)
y^{1}sc*v^{1}sev^{21}; P{y[+t7.7] v[+t1.8]=TRiP.GL01341}attP40	BLM #42482	(Perkins et al., 2015)
y^{1}sc*v^{1}sev^{21}; P{y[+t7.7] v[+t1.8]=TRiP.HMS02846}attP2	BLM #44551	(Perkins et al., 2015)
y^{1}v^1; P{y[+t7.7] v[+t1.8]=TRiP.HMS00020}attP2	BLM #33626	(Perkins et al., 2015)
y^{1}sc*v^{1}sev^{21}; P{y[+t7.7] v[+t1.8]=TRiP.HMS01408}attP2	BLM #34998	(Perkins et al., 2015)
y^{1}sc*v^{1}sev^{21}; P{y[+t7.7] v[+t1.8]=TRiP.HMS00991}attP2	BLM #34021	(Perkins et al., 2015)
y^{1}v^1; P{y[+t7.7] v[+t1.8]=TRiP.GL01273}attP2	BLM #41845	(Perkins et al., 2015)
y^{1}sc*v1; P{y[+t7.7] v[+t1.8]=TRiP.HMS02968}attP2	BLM #50732	(Perkins et al., 2015)
y^{1}sc*v^{1}sev^{21}; P{y[+t7.7] v[+t1.8]=TRiP.GL00435}attP40	BLM #35598	(Perkins et al., 2015)
y^{1}sc*v^{1}sev^{21}; P{y[+t7.7] v[+t1.8]=TRiP.HMC03738}attP40	BLM #55142	(Perkins et al., 2015)
y^{1}v^1; P{y[+t7.7] v[+t1.8]=TRiP.HMJ22418}attP40/CyO	BLM #58311	(Perkins et al., 2015)

$y^1 sc^* v^1 sev^{21}$; $P\{y[+t7.7]$ $v[+t1.8]=TRiP.HMC04414\}attP40$	BLM #56974	(Perkins et al., 2015)
$y^1 v^1$; $P\{y[+t7.7]$ $v[+t1.8]=TRiP.HMJ23009\}attP40$	BLM #61230	(Perkins et al., 2015)
hsFLP; FRT40A, UbiGFP / CyO	D. Drummond-Barbosa	(Ables et al., 2016)
FRT40A / CyO	D. Drummond-Barbosa	(Ables et al., 2016)
$y^{d2} w^{1118} P\{ry[+t7.2]=ey$-FLP.N$\}2$ $P\{ry[+t7.2]=GMR$-lacZ.C(38.1)$\}TPN1$; $P\{y[+mDint2] w[BR.E.BR]=SUPor$-$P\}Trn$-SR[KG04870] $P\{ry[+t7.2]=neoFRT\}40A$ / CyO, $y[+]$	Kyoto #111581	(Bellen et al., 2004; Chen et al., 2005)
$y^* w^*$; PBac$\{SAstopDsRed\}LL05552$ $P\{ry[+t7.2]=neoFRT\}40A$ $P\{w[+mW.hs]=FRT(w[hs])\}G13 cn^1 bw^1$ / CyO, $S^* bw^1$	Kyoto #141628	(Schuldiner et al., 2008)
w;$P\{UASp:Ran^{HA}\}1.19/SM6$	K. McKim	(Cesario and McKim, 2011)
w;$P\{UASp:Ran^{HA}\}10$-A/TM3	K. McKim	(Cesario and McKim, 2011)
w;$P\{UASp:Ran^{T24NHA}\}18.3/TM3$	K. McKim	(Cesario and McKim, 2011)
w;$P\{UASp:Ran^{T24NHA}\}25.1/TM3$	K. McKim	(Cesario and McKim, 2011)
w;$P\{UASp:Ran^{Q69LHA}\}7.1/TM3$	K. McKim	(Cesario and McKim, 2011)
w;$P\{UASp:ran^{Q69LHA}\}61.2/TM3$	K. McKim	(Cesario and McKim, 2011)
w[1118]; $P\{UASp$-GFP.E2f.1-230$\}26$ $P\{UASp$-mRFP1.NLS.CycB.1-266$\}4/CyO,P\{en1\}wg[en11]$;MKRS/TM6B, Tb1	BLM #55110	(Zielke et al., 2014)
Rtnl1-GFP	B. Riggs, K. Röper	(Röper, 2007)
scrib^{CA07683}	L. Cooley, M. Buszczak	(Buszczak et al., 2007)

w*; P{w[+mC]=UASp-GFP-scra}3	BLM #51348	(Silverman-Gavrila et al., 2008)
w*; wg^{Sp-1}/CyO; P{w[+mC]=UASp-GFP-3A-scra}3	BLM #51347	(Silverman-Gavrila et al., 2008)

1143

1144

1145