β-importins Tnpo-SR and Cadmus and the small GTPase Ran promote ovarian cyst formation in *Drosophila*

Allison N. Beachum, Taylor D. Hinnant, Anna E. Williams, Amanda M. Powell, and Elizabeth T. Ables*

Department of Biology, East Carolina University, Greenville, NC 27858.

*Correspondence: ablese@ecu.edu; 1001 E. 10th St. Mailstop 551, Greenville, NC 27858; Phone: 252-328-9770; Fax: 252-328-4178

RUNNING TITLE

importins promote cyst formation

KEYWORDS

Oocyte, oogenesis, germline, cyst, germline stem cell

CORRESPONDING AUTHOR:

Elizabeth T. Ables

East Carolina University

Department of Biology

1001 E. 10th St., Mailstop 551

553 Science & Technology Building

Greenville, NC 27858

Office phone: 252-328-9770

e-mail: ablese@ecu.edu

SUMMARY STATEMENT

Ran and two β-importins function coordinately to promote oocyte selection and cyst development in the *Drosophila* ovary.

ABSTRACT

Germ cells undergo mitotic expansion via incomplete cytokinesis, forming cysts of undifferentiated cells that remain interconnected prior to meiotic initiation, through mechanisms that are not well-defined. In somatic cells, Ras-related nuclear protein (Ran) spatiotemporally regulates mitotic spindle assembly, cleavage furrow formation and abscission. Here, we identify Ran and β-importins as critical regulators of cyst development in the *Drosophila* ovary. Depletion of *Ran* or the β-importins *Tnpo-SR* and *cadmus* disrupts oocyte selection and results in egg chambers with variable numbers of germ cells, suggesting abnormal cyst development and cyst fragmentation. We demonstrate that Ran, Tnpo-SR, and Cadmus regulate key cellular processes during cyst formation, including cell cycle dynamics, fusome biogenesis, and ring canal stability, yet do so independently of mitotic spindle assembly. Further, Tnpo-SR and Cadmus control cyclin accumulation and suppress cytokinesis independent of Ran-GTP, suggesting that β-importins sequester protein cargos that normally promote the mitotic-to-meiotic transition. Our data demonstrates that Ran and β-importins are critical for germ cell cyst formation, a role that is likely conserved in other organisms.

INTRODUCTION

Germ cells develop as cysts of interconnected undifferentiated cells (Lu et al., 2017; Matova and Cooley, 2001; Pepling and Lei, 2018). Cysts arise via incomplete cytokinesis, manifested by cleavage furrow arrest and the formation of stable ring canals that permit transport of cytoplasm and organelles (de Cuevas et al., 1997; Guertin et al., 2002; Haglund et al., 2011; Pepling and Lei, 2018). Cyst formation increases cytoplasmic volume, enhances sensitivity to DNA damage, and ensures robust oocyte development (Chaigne et al., 2020; de Cuevas et al., 1997; Pepling and Lei, 2018; Pepling et al., 1999; Yamashita, 2018). Cyst formation is evolutionarily conserved and essential for optimal fertility in multiple phyla (Haglund et al., 2011; Lu et al., 2017).

The fruit fly, *Drosophila melanogaster*, is an elegant model of germline cyst formation. In adult females, oocytes are produced in a linear spatiotemporal arrangement fueled by the activity of germline stem cells (GSCs; Figure 1A) (Hinnant et al., 2020). GSCs divide asymmetrically, giving rise to one daughter that remains a GSC and a cystoblast that differentiates. Delayed abscission between the GSC and cystoblast keeps the pair connected well into G2 of the subsequent cell cycle (Ables and Drummond-Barbosa, 2013; Eikenes Å et al., 2015; Hinnant et al., 2017; Mathieu et al., 2013; Matias et al., 2015). Cystoblasts divide four times with incomplete cytokinesis, creating 16-cell cysts. At the completion of the mitotic divisions, cleavage furrows between germ cells are modified into stable ring canals (Grieder et al., 2000; Guertin et al., 2002; Hinnant et al., 2020). The fusome, a cytoplasmic organelle composed of microtubules, endoplasmic reticulum-derived vesicles, and membrane skeleton proteins, traverses the intercellular bridges (Hinnant et al., 2020; Ong and Tan, 2010; Pepling and Lei, 2018; Röper, 2007; Snapp et al., 2004). The fusome branches as mitotic divisions progress, providing a synchronization hub for cell cycle regulation and cell polarity (Grieder et al., 2000; Hinnant et al., 2020; Lighthouse et al., 2008). These specialized cytological features of germ cells facilitate interconnectivity and are vital to oocyte selection; however, the molecular mechanisms underlying fusome morphogenesis and incomplete cytokinesis are not well understood.

Formation of an actomyosin-based contractile network at the equatorial cleavage plane during cytokinesis is spatiotemporally regulated and requires a dynamic interplay between the cytoskeleton, the mitotic microtubule-based central spindle, and membrane trafficking (Frémont and Echard, 2018; Haglund et al., 2011; Nguyen and Robinson, 2020; Pollard and O'Shaughnessy,

2019). Central to this process is Ras-related nuclear protein (Ran), a small guanosine triphosphate

(GTP)-binding protein implicated in protein transport during mitosis and interphase (Ozugergin

and Piekny, 2020; Sazer and Dasso, 2000). Ran partners with α- and β-importins to move protein

cargo into the nucleus and promote mitotic spindle assembly (Cautain et al., 2015; Okada et al.,

2008). Ran also regulates cell polarization and contractile protein localization during cleavage

furrow ingression (Beaudet et al., 2017; Silverman-Gavrila et al., 2008). Chromatin-associated

Ran-GTP may help recruit and organize essential proteins at the cortex, thereby ensuring the

robustness of cytokinesis (Nguyen and Robinson, 2020).

Here, we identify Ran and β-importins as novel regulators of germline cyst formation in

the *Drosophila* ovary. We show that Ran and β-importins Tnpo-SR and Cadmus (Cdm) control

cyst formation by regulating cell cycle dynamics, fusome biogenesis, and ring canal stability. Ran

function depends on its active, GTP-bound state, as expression of a dominant-negative *Ran* mutant

recapitulates *Ran* loss-of-function. *Ran*, *Tnpo-SR*, and *cdm* are not essential for mitotic spindle

assembly in undifferentiated germ cells. Moreover, knock-down of *Tnpo-SR* or *cdm* does not

completely phenocopy loss of *Ran*, suggesting that β-importins control cell cycle progression and

suppress cytokinesis independent of Ran activity. Given the conservation of germline cyst

formation across phyla, our data suggest that Ran and specific β-importins may play similar roles

in other organisms.

RESULTS

RNAi screen reveals that β-importins and Ran are essential for germ cell development.

We previously identified *Transportin-Serine/Arginine rich* (*Tnpo-SR*) in a genetic mosaic

screen for genes controlling GSC self-renewal (Ables et al., 2016). *Tnpo-SR*, homologous to

mammalian *TNPO3*, encodes a β-importin (Allemand et al., 2002; Lai et al., 2001). To determine

whether *Tnpo-SR* loss-of-function phenotypes reflect a general role for importins in germ cell

mitoses, we performed a secondary reverse genetic screen using germline-enhanced RNA

interference (RNAi) to knock-down expression of nine *Drosophila* importins and the small

GTPase *Ran* specifically in germ cells (Figure 1B-D, Table 1) (Ni et al., 2011; Rørth, 1998; Van

Doren et al., 1998). In contrast to driver-only controls (Figure 1B, Table 1), three RNAi lines

produced completely agametic ovaries, defined by the absence of egg chambers (Figure 1C). More

intriguingly, four RNAi lines corresponding to *Ran* and the β-importins *Tnpo-SR*, *cadmus* (*cdm*),

and *Fs(2)Ket* (also called *Ketel*) exhibited a partial agametic phenotype, in which germ cells were

present in some ovarioles, but in unusually shaped egg chambers (Figure 1D; Table 1). Cdm and

Tnpo-SR are paralogs, sharing 42% amino acid similarity, while Ketel is less conserved. These

results demonstrate that Ran and some β-importins are necessary for proper oocyte development

and may have distinct functions.

Since GSCs give rise to all germ cells, we hypothesized that agametic ovaries resulting

from *Ran* and *β-importin* knock-down could arise from a failure to establish or maintain GSCs. To

test this, we quantified the number of GSCs per germarium. GSCs and their progeny can be easily

identified by immunolabeling of Hts, a fusome-resident protein in germ cells, and LaminC

(LamC), a nuclear envelope protein highly expressed in cap cells (the major cellular component

of the GSC niche) (Ables and Drummond-Barbosa, 2013). Unlike cytoblasts, GSCs possess an

anteriorly localized fusome juxtaposed to the cap cells (de Cuevas and Spradling, 1998; Lin et al.,

1994). All four partially agametic RNAi lines had significantly less GSCs per germarium at three

days after eclosion than age-matched driver-only controls (Table 1; Supp. Figure 1A-C). Fewer

GSCs in early adulthood suggests that *Ran*, *Tnpo-SR*, *Ketel*, and *cdm* are required for GSC

establishment during larval/pupal development or during the proliferative surge in GSCs following

the female's first mating (Ameku and Niwa, 2016; Van Doren et al., 1998). Conditional knock-

out of *Tnpo-SR* further demonstrated the requirement for β-importin function in GSC maintenance.

We used two strong loss-of-function transposon insertion alleles of *Tnpo-SR* (*Tnpo-SR^{KG04870}* and

Tnpo-SR^{LL05552}) carrying *FRT* sites to generate germline mutant mosaic germaria under the

Flippase (*FLP*)/*FLP Recognition Target* (*FRT*) genetic mosaic lineage-tracing system (Laws and

Drummond-Barbosa, 2015; Xu and Rubin, 1993). GFP-negative GSC clones were generated at

equivalent rates in control, *Tnpo-SR^{KG04870}*, and *Tnpo-SR^{LL05552}* mosaic germaria at four days after

clone induction (Supp. Figure 1F). In contrast to mock controls, *Tnpo-SR^{KG04870}* and *Tnpo-SR^{LL05552}*

germline mutant mosaic germaria rapidly lost marked GSCs over time (Supp. Figure 1D-F),

indicating that *Tnpo-SR* is cell-autonomously required for GSC self-renewal.

Loss-of-function mutations in Ran and β-importins alter the number of germ cells per egg

chamber.

The intriguing phenotype resulting from knock-down of *Ran* and β-importins prompted us

to examine egg chamber morphology more closely. To visualize germline and follicle cells, we

combined anti-Hts antibodies with anti-Hts-RC antibodies, which are specific for a Hts cleavage

product that localizes to ring canals (Petrella et al., 2007; Robinson et al., 1994). Unlike driver

control egg chambers (Figure 1E; Table 1), knock-down of *Ran*, *cdm*, or *Tnpo-SR* significantly

decreased the number of egg chambers with 16 germ cells (Figure 1F-G; Table 1). Depletion of

Ketel did not significantly impact egg chamber formation (Table 1).

Cystoblasts normally divide synchronously four times, following a 2^n pattern, to produce

cysts with 16 cells (King, 1970). To determine whether extra germ cells in *Ran*, *cdm*, and *Tnpo-*

SR depleted ovarioles arose by alterations in the number of mitotic divisions, we quantified the

number of germ cells per cyst (Figure 1J). In the rare instances where control or *Ketel* RNAi egg

chambers had an abnormal number of germ cells, they invariably contained 32 cells (Figure 1J).

In contrast, knock-down of *Ran*, *cdm,* or *Tnpo-SR* produced egg chambers with variable numbers

of germ cells, ranging anywhere from one to 49 cells per cyst (Figure 1J). While some egg

chambers housed germ cells on the order of 2^n, a substantial number deviated from this pattern

(Table 1). Germ cell numbers in adjacent chambers in RNAi-depleted egg chambers did not add

up to 8, 16, or 32 cells, arguing that encapsulation defects alone are not the source of the unusual

germ cell number phenotype. Some egg chambers with 32 or more cells had oocytes with five ring

canals, suggesting that some abnormal egg chambers arose by an extra mitotic division. Other egg

chambers, however, likely arose because of other cellular defects. Moreover, a small percentage

of *cdm* and *Tnpo-SR* RNAi knock-down ovarioles (17.4% and 13.3% of all ovarioles, respectively)

contained germline tumors, in which a normal follicle cell monolayer surrounded mitotically-

active germ cells with underdeveloped fusomes (Figure 1H-I; Supp. Figure 2A-C'). In *Tnpo-SR*

RNAi ovarioles, undifferentiated germ cells were encapsulated with more-developed germ cells

(Supp. Figure 2A). Germline tumors contained mitotically-active cells with short or unbranched

fusomes and immature ring canals that frequently appeared as dot-like structures, reminiscent of

the abscission site in GSC/cystoblast pairs (Supp. Figure 2B-C'). These structures suggest tumors

arose because of improper abscission events in 2-cell and/or 4-cell cysts. Immunostaining for Sex-

lethal (Sxl), which is enriched in GSCs and cystoblasts in the anterior-most region of the

germarium (Supp. Figure 2D-D') (Chau et al., 2009), confirmed that *Tnpo-SR* RNAi germ cells

have not completed differentiation (Supp. Figure 2E-E'). We conclude that Ran, Tnpo-SR, and

Cdm impact the synchrony and number of cyst divisions.

Ran, Tnpo-SR, and cdm are required for oocyte selection.

During germ cell mitotic divisions, three to four cyst cells initiate meiosis by assembling

the synaptonemal complex; however, only one of these cells is maintained as the oocyte (Hughes

et al., 2018). Since depletion of *Ran*, *Tnpo-SR*, or *cdm* altered the number of germ cells per cyst,

we hypothesized that oocyte selection was also affected. In wild-type egg chambers just posterior

to the germarium, the oocyte-specific proteins Orb (Figure 2A) and Bicaudal D (BicD) (Figure

2D) had clearly accumulated in one posterior cyst cell. In contrast, egg chambers from germline-

depleted *Ran*, *Tnpo-SR*, or *cdm* females did not properly express Orb ($Ran^{GL01341}$, $cdm^{HMS02846}$, *and*

$Tnpo\text{-}SR^{V22}$) or BicD ($Tnpo\text{-}SR^{V22}$) (Figure 2B-C,E-F; Table 1). Many loss-of-function egg

chambers also lacked a cell with a karyosome, a clear indicator of oocyte fate (Figure 3B-C,E-F).

Importantly, most of the *nos-Gal4>$Ran^{GL01341}$* and *nos-Gal4>$cdm^{HMS02846}$* egg chambers with 16

germ cells contained an oocyte (Table 1), suggesting that loss of oocyte identity is a secondary

effect of incorrect mitotic divisions in the absence of *Ran* and *cdm*. This was not the case, however,

for *Tnpo-SR*, as 34% of *nos-Gal4>$Tnpo\text{-}SR^{V22}$* egg chambers with 16 germ cells contained only

nurse cells (Table 1). Similarly, in *Tnpo-SR* germline mutant mosaic germaria, mutant cysts

located posterior to wild-type cysts (thus at a more advanced stage of development) failed to up-

regulate Orb expression to similar levels as adjacent wild-type cysts (Figure 2G-H). Further, a core

component of the synaptonemal complex, C(3)G, normally present as early as the 4-cell cyst stage,

did not accumulate in *Tnpo-SR^{KG04870}* mutant cysts to equivalent levels as adjacent wild-type cysts

(Figure 2I-J). This suggests that *Tnpo-SR* is specifically required for oocyte specification

independent of its role in cyst divisions.

Ran is dynamically localized in cyst cells and dispensable for spindle assembly.

One well-characterized function of Ran and β-importins is the regulation of spindle

assembly (Cesario and McKim, 2011; Forbes et al., 2015; Nguyen and Robinson, 2020; Ozugergin

and Piekny, 2020). Spindle orientation is critical for GSC maintenance, and mitotic spindle

remnants accumulate in fusomes as cysts divide (Cheng et al., 2008; Deng and Lin, 1997; Grieder

et al., 2000; Máthé et al., 2003). Like other small GTP-binding proteins, Ran cycles between active

(GTP-bound) and inactive (GDP-bound) states (Ozugergin and Piekny, 2020). Ran-GTP

dissociates importin-bound cargo in the nucleus and is converted to Ran-GDP in the cytoplasm. In

some cell types, this establishes a gradient of active Ran in the cell that promotes spindle formation.

To determine whether such a gradient may exist in cyst cells, we first analyzed the

subcellular distribution of Ran. We took advantage of a previously characterized transgene in

which the germline-specific *UASp* promoter drives an N-terminal hemagglutinin (HA)-tagged Ran

(*Ran^{HA}*) (Cesario and McKim, 2011). Using *nos-Gal4* to drive *Ran^{HA}* in germ cells, we then

localized Ran using anti-HA antibodies in comparison to mitotic spindles (labeled with α-tubulin),

fusomes, and chromosomes (Figure 4A-A'). Unlike neuroblasts, in which Ran-HA is nuclearly

localized during interphase and overlaps with the mitotic spindle during mitosis (Cesario and

McKim, 2011), we did not observe strong nuclear localization of Ran-HA in GSCs or cyst cells

(Figure 3A-B''). Ran-HA was higher in the cytoplasm than in nuclei of interphase and mitotically-

active cells and, consistent with neuroblasts and embryonic cells (Cesario and McKim, 2011;

Trieselmann and Wilde, 2002), was absent from chromosomes (Figure 3B-B''). Ran-HA

expression was higher in some cysts than others, likely due to minor variations in driver strength.

In contrast to meiotic spindles (Cesario and McKim, 2011), we did not observe strong

accumulation of Ran-HA surrounding mitotic spindles (Figure 3B-B'').

We then asked whether Ran-GDP accumulated in specific subcellular locations in cyst

cells. We over-expressed an HA-tagged *Ran^{T24N}*, which carries an amino acid substitution that

locks Ran into a GDP-bound state and inhibits cargo loading (Cesario and McKim, 2011; Drutovic

et al., 2020). As expected, Ran-GDP was primarily cytoplasmic in interphase cells (Figure 3C).

We also observed that Ran-GDP accumulated in the nucleus of mitotically-dividing cells (Figure

3C), overlapping with chromatin at metaphase (Figure 3D-D''). Mitotic spindles appeared to

assemble correctly (see below), but chromatin was dispersed in interphase cells.

The subcellular localization of Ran-GTP was markedly different than that of Ran-GDP.

We used a second HA-tagged transgene, *Ran^{Q69L}*, to over-express a constitutively active form of

Ran locked in a hydrolysis-resistant GTP-bound state (Cesario and McKim, 2011). Ran-GTP was

excluded from cyst cell nuclei during interphase (Figure 3C) and accumulated at the presumptive

nuclear membrane during mitosis (Figure 3D-D''). Taken together, these experiments suggest that

Ran aids in chromatin condensation or cytokinesis following the completion of mitosis.

Finally, we asked whether Ran and β-importins are necessary for spindle formation. Using

immunostaining for α-Tubulin, we confirmed that mitotic spindles in dividing cysts are anchored

to the fusome, oriented perpendicularly (Figure 3G) (Megraw and Kaufman, 2000; Storto and

King, 1989). Surprisingly, in *cdm*-depleted (Figure 3H) or *Tnpo-SR*-depleted (Figure 3I) germ

cells, similar-sized spindles formed and were appropriately anchored and oriented with respect to

fusomes. Similarly, we were unable to find defects in spindle formation or orientation in

nosGal4>Ran^{T24N} germ cells (Figure 3D,J). We conclude that mitotic spindles can form even when

Ran-GTP is depleted, suggesting that low levels of active Ran are sufficient to maintain germ cell

mitoses.

GSC establishment, cyst formation, and oocyte selection require active Ran-GTP.

Given the intriguing localization of Ran-GTP, we then asked whether active Ran-GTP is

required for GSC establishment and cyst formation. Flies expressing the *Ran^{T24N}* and *Ran^{Q69L}*

mutations in germ cells were compared to flies expressing wild-type *Ran^{HA}*. We expected that

constitutively expressing Ran-GDP would titrate out the amount of active Ran, thus mimicking

Ran loss-of-function. Indeed, over-expression of *Ran^{T24N}* resulted in a partial agametic phenotype

and the average number of GSCs per germarium was substantially reduced as compared to age-

matched controls (Figure 4A; Table 1). Over-expression of dominant negative *Ran^{T24N}* also altered

the number of germ cells per cyst (Figure 4B-C,E) and, similar to knock-down of *Ran* via RNAi,

the number of germ cells did not always follow a 2^n pattern (Figure 4E, Table 1). Over-expression

of *Ran^{T24N}* also impaired oocyte selection, as 12-24% of egg chambers with 16 germ cells lacked

an Orb-positive cell (Table 1). These phenotypes are consistent with depletion of *Ran* via RNAi

and likely contribute to the female sterility phenotype noted in prior studies (Cesario and McKim,

2011).

Intriguingly, excess Ran-GTP in germ cells did not result in severe germline defects

(Figure 4D-E, Table 1). The average number of GSCs per germarium was slightly reduced in *nos-*

Gal4>Ran^{Q69L} ovarioles (Table 1) and a few egg chambers had an incorrect number of cells (Figure

4E), but defects were not as severe or penetrant as either the *Ran* RNAi or Ran-GDP-locked

mutants. Since we did not observe a clear Ran intracellular concentration gradient or accumulation

of Ran at mitotic spindles, these results suggest that Ran-GTP levels are already in excess in

mitotically-active germ cells, perhaps to facilitate timely mitotic divisions.

Tnpo-SR and cdm, but not Ran-GTP, are required for fusome biogenesis and maturation.

The fusome extends through ring canals during incomplete cytokinesis, forming an

extensive tubular network whose structure changes over the course of the mitotic divisions

(Lighthouse et al., 2008; Röper, 2007; Snapp et al., 2004). The cytoskeletal core is prominent in

GSC and cystoblast fusomes, but is replaced with endoplasmic reticulum-like vesicles as cysts exit

the germarium (Figure 5A) (Röper, 2007). Loss of *Hts*, a core fusome cytoskeletal protein, alters

the number of germ cells per cyst (Yue and Spradling, 1992). Since knock-down of *Ran* or the β-

importins produced similar phenotypes, we hypothesized that abnormal fusome morphogenesis

might abrogate cyst formation. Using Hts immunostaining as an indicator of fusome morphology,

we observed that when *cdm* (Figure 5B) or *Tnpo-SR* (Figure 5C and Supp. Figure S3) were

depleted, germ cells formed long, thin fusomes with irregular branching, particularly in cysts in

the middle to posterior third of germaria. Fusomes in *Tnpo-SR*- and *cdm*-depleted germ cells often

had thread-like sections separated by thicker sections, consistent with weak points in the

cytoskeletal structure (Figure 5B-C). Irregular fusomes were also observed in *Tnpo-SR*-depleted

GSC/cystoblast pairs, where new fusome material builds from the cystoblast to the GSC (compare

Figure 5E-F) (de Cuevas and Spradling, 1998). Importantly, we did not observe fusomes

connecting GSCs to other cyst cells in stem-cysts, as was reported for germ cells in which the

chromosomal passenger complex, which regulates abscission timing in GSCs, is abrogated

(Mathieu and Huynh, 2017). We also did not observe GSCs or cysts with asynchronous mitoses

(compare Figure 5G-H), as we would predict if fusomes lost complete connectivity between cells.

We conclude that *Tnpo-SR* and *cdm* promote fusome maintenance or remodeling.

To further investigate how the importins regulate fusome morphogenesis, we recombined

the *Tnpo-SR^{V22}* RNAi transgene with two fluorescently tagged fusome-resident protein reporters.

Scribble (*scrib*) encodes a scaffolding protein that regulates vesicle trafficking and is expressed at

the cell cortex and in immature fusomes of GSCs and dividing cysts (Figure 5I) (Lighthouse et al.,

2008). By the third mitotic division, *Scrib-GFP* is largely absent from wild-type fusomes and does

not co-localize with Hts in 16-cell cysts. In *Tnpo-SR*-depleted germ cells, *Scrib-GFP* was

prominently expressed in the long, thin fusomes of germ cells in the posterior of germaria (Figure

5J). Conversely, Reticulon-like1 (Rtnl1) is an endoplasmic reticulum-resident protein that is

expressed in low levels in immature fusomes (Figure 5K) (Röper, 2007). *Rtnl1-GFP* becomes

enriched in mature fusomes coincident with cyst division and inversely correlated with Hts

expression, reaching peak concentration in fusomes after the final mitotic division (Figure 5K).

When *Tnpo-SR* is depleted, *Rtnl1-GFP* and Hts co-localize in fusomes throughout the posterior of

the germarium (Figure 5L). These data suggest that in the absence of β-importins, cytoskeletal

components of the fusome fail to degrade properly, maintaining the fusome in an immature state

that is weakly connected between cyst cells. Intriguingly, fusome shape and branching in cysts

over-expressing the dominant negative Ran^{T24N} mutant (Figure 5D) were indistinguishable from

wild-type cysts, suggesting that Cdm and Tnpo-SR regulate fusome morphogenesis independently

of Ran-GTP.

Tnpo-SR and cdm promote cyst formation by suppressing cytokinesis.

Knock-down of *Tnpo-SR* or *cdm* produced egg chambers variable numbers of germ cells;

however, it remained unclear to us exactly how such egg chambers could arise. Key observations

came by closely examining the morphology of *Tnpo-SR* mutant cysts surrounded by wild-type

cysts in mutant mosaic germaria. While knock-down of *Tnpo-SR* by RNAi led to >16 germ cells

per egg chamber, we only rarely found examples of *Tnpo-SR^{KG04870}* or *Tnpo-SR^{LL05552}* mutant egg

chambers with more than 16 mutant germ cells. We did find, however, compound egg chambers

in which a *Tnpo-SR* mutant cyst was encapsulated together with wild-type cysts (Supp. Figure

S3A). We occasionally identified mitotically-dividing *Tnpo-SR^{KG04870}* and *Tnpo-SR^{LL05552}* mutant

germ cells in mosaic egg chambers outside of the germarium, which never occurs in wild-type

germ cells (Supp. Figure S3B). We also frequently found clusters of single *Tnpo-SR* mutant germ

cells at the posterior of germaria (Supp. Figure S3C,E-E'''), suggesting that some cyst defects in

Tnpo-SR mutant and RNAi-knock-down ovarioles could arise by cyst fragmentation. In the *Tnpo-*

SR RNAi model, where all germ cells lack appropriate Tnpo-SR levels, fragmented cysts would

likely be encapsulated along with adjacent cysts, thus yielding irregular (i.e., not 2^n) numbers of

germ cells per cyst. In support of this idea, we observed examples of mutant mosaic germaria in

which *Tnpo-SR^{KG04870}* mutant cells with dot-like fusomes were adjacent to one another with

fusomes aligned along the same plane, as if they were connected (Supp. Figure S3D).

Immunostaining with anti-pTyr antibodies to visualize ring canals highlighted *Tnpo-SR^{KG04870}*

mutant cysts where ring canal components accumulated on adjacent cyst cells, separated by a very

thin line of GFP-positive cytoplasm (Figure 6A and Supp. Figure S3E), suggesting that cells have

physically separated from each other after clone induction. Moreover, *Tnpo-SR^{KG04870}* mutant germ

cells failed to accumulate proper levels of the cell adhesion protein E-cadherin between germ cells,

as occurs in adjacent wild-type 16-cell cysts following the final mitotic division and prior to follicle

cell encapsulation (Figure 6B) (Fichelson et al., 2010).

These data raised the possibility that in the absence of *Tnpo-SR*, cysts fail to stay connected

by intercellular bridges, and instead break apart at unstable ring canals. Initially, this seemed

unlikely, as our analyses of *Ran*, *cdm*, and *Tnpo-SR* RNAi and *Ran* dominant-negative germ cells

focused on egg chambers outside of the germarium, where ring canals containing Hts-RC (a

constituent protein of mature ring canals) were easily detected (Figure 1E-G). Moreover, the

number of ring canals scaled to the number of germ cells, suggesting that ring canals formed at

successive divisions. Spatiotemporal comparison of ring canal formation, however, demonstrated

that while Hts-RC is expressed in germ cells from *nos-Gal4>Tnpo-SR^{V22}* ovarioles, its

accumulation at ring canals is drastically delayed, such that protein levels are much lower than in

age-matched wild-type cysts (compare Figure 6C and D). Similarly, germ cells depleted of *cdm*

(Figure 6F) or *Tnpo-SR* (Figure 6G), failed to accumulate pTyr-reactive material (one of the first

indicators of ring canal maturation) to wild-type levels in 16-cell cysts. In contrast, ring canal

maturation proceeded normally in *Ran^{T24N}* mutant cysts (Figure 6H), supporting the model that

Tnpo-SR and Cdm, but not Ran-GTP, stabilize germ cell intercellular bridges, maintaining proper

cyst formation (Figure 6M).

We then performed immunostaining for Anillin, an actin/myosin-binding protein known to

be a β-importin cargo (Beaudet et al., 2017; Beaudet et al., 2020; Chen et al., 2015; Piekny and

Maddox, 2010; Silverman-Gavrila et al., 2008). Anillin localization in ring canals begins at the

third mitotic division (Figure 6I) (Goldbach et al., 2010; Haglund et al., 2010). In *Tnpo-SR*-

depleted germ cells, we observed Anillin deposition within the intercellular bridge, rather than on

surrounding rings (Figure 6J). In contrast, Anillin localizes properly to the ring canals in *Ran^{T24N}*

mutants (Figure 6K) and occasionally co-localized with the fusome (Figure 6L), supporting the

model that Tnpo-SR suppresses cytokinesis in dividing cysts independently of Ran-GTP.

Intriguingly, over-expression of a GFP-tagged Anillin in which a nuclear localization signal has

been mutated did not alter Anillin localization in abscising GSC/CB pairs (Supp. Figure 4A-B) or

at ring canals (Supp. Figure 4C-D), and did not change the number of germ cells per cyst. Although

we cannot exclude the possibility that Tnpo-SR may bind Anillin at another protein domain, we

speculate that Tnpo-SR indirectly regulates Anillin by binding other cargo proteins that impact

Anillin localization.

Ectopic CycA accumulates in Tnpo-SR- and cdm-depleted germ cells.

The cell cycle is tightly regulated to coordinate cyst formation. For example, loss of function of the mitotic cyclin CycB produces cysts with eight cells, while over-expression of CycA or CycE produces cysts with 32 cells (Lilly et al., 2000; Mathieu et al., 2013; Morris et al., 2005; Ohlmeyer and Schüpbach, 2003). CycA and CycB accumulate at the fusome and phosphorylation of CycB by AuroraB Kinase promotes abscission between GSC/cystoblast pairs (Lilly et al., 2000; Mathieu et al., 2013). Given the phenotypic parallels, we hypothesized that Tnpo-SR and Cdm might regulate cyclin dynamics in dividing germ cells. We confirmed that CycA accumulates at the fusome in ~10% of mitotically dividing cysts, but never associates with the fusome of 16-cell cysts in wild-type tissue (Figure 7A,F). *Ran* depletion in *nos-Gal4>Ran^{GL01341}* germaria did not alter CycA kinetics (Figure 7B,F). In contrast, nearly 40% of mitotically-dividing cysts in *Tnpo-SR* and *cdm*-depleted germ cells expressed CycA; moreover, CycA expression was observed in the fusomes of 16-cell cysts at an equivalent rate as dividing cysts, suggesting that germ cells remain mitotic (Figure 7C-D,F).

As the highest peak of CycB expression occurs following CycA degradation (Morris et al., 2005), we then asked whether CycB regulation was likewise impacted by loss of the β-importins. We utilized the *Drosophila* Fluorescence Ubiquitin-based Cell Cycle Indicator system (Fly-FUCCI) (Zielke et al., 2014) to ask whether CycB expression was altered in the absence of *Tnpo-SR*. Importantly, we used a version of the *UASp-CycB-RFP* transgene that includes a canonical nuclear localization sequence (Hinnant et al., 2017; Zielke et al., 2014). When *Tnpo-SR* was depleted, the *UASp-CycB-RFP* localized to germ cell nuclei; however, significantly fewer mitotically dividing cysts expressed CycB as compared to wild-type controls (Figure 7E,G). Taken together, these data suggest that *Tnpo-SR*, and likely *cdm*, promote proper cyclin dynamics in germ cells, maintaining cysts in the mitotic expansion phase and facilitating coordination between cyst division and fusome morphogenesis.

DISCUSSION

During germline cyst formation, cell cycle transitions are precisely controlled to promote mitotic divisions and block cellular abscission. Our study adds new details to our understanding of the molecular regulation of germ cell mitotic expansion, identifying previously undescribed roles for β-importins and the small GTPase Ran in mitotic cell cycle exit during cyst formation. Remarkably, we do not detect a clear cytoplasmic-to-nuclear gradient of Ran protein localization

in germ cell nuclei. Nor do we find defects in mitotic spindle assembly, cell size, or DNA

condensation in the absence of β-importins or activated Ran. Instead, our data suggest that

importins and Ran-GTP promote mitotic exit by distinct, yet complimentary mechanisms. We

propose the model that Ran-GTP drives germ cell mitoses by mobilizing cell cycle factors,

functioning together with the β-importins Tnpo-SR and Cdm to facilitate transport of cell cycle

factors from the fusome to other parts of the cell. We propose that independent of Ran, Tnpo-SR

and Cdm sequester cell cycle regulators like CycA and Anillin, titrating the levels of protein

accessible to Ran-GTP, counterbalancing mitotic progression. Thus, in the absence of Tnpo-SR or

Cdm, CycA levels accumulate, causing germ cells to continue in a proliferative state and impairing

fusome stability, ring canal maturation, and oocyte terminal differentiation.

Importins may exhibit non-redundant functions in the Drosophila germline.

Although several α-importins have been implicated in oogenesis prior to our study (Geles

and Adam, 2001; Mason et al., 2002; Máthé et al., 2000), our study is the first to identify specific

roles for β-importins in mitotic expansion. Our data generally support the hypothesis that specific

importins regulate distinct aspects of germ cell development (Hogarth et al., 2005; Loveland et al.,

2015; Mihalas et al., 2015). We cannot exclude the possibility, however, that RNAi was not

sufficient to completely inhibit importin function. For example, Ketel localizes at the nuclear

envelope in early embryos and dividing germ cells (data not shown) (Lippai et al., 2000; Tirián et

al., 2000; Trieselmann and Wilde, 2002; Villányi et al., 2008), yet we did not observe germ cell

mitotic defects in *Ketel*-depleted germ cells. Future studies will need to employ other loss-of-

function approaches to test the roles of Ketel and other β-importins more thoroughly.

The repertoire of cargo carried by a β-importin determines its role in biological processes

and its redundancy with other importins (Kimura and Imamoto, 2014). Although the full extent of

protein-protein interactions mediated by Tnpo-SR remains unknown, structural studies indicate

that TNPO3 binds proteins involved in chromatin organization, mRNA and protein modification,

cell cycle control, and cell-cell communication (Kimura et al., 2017; Maertens et al., 2014). Tnpo-

SR is also likely to bind a broad range of cargo proteins, including SR-rich RNA binding proteins.

Indeed, Tnpo-SR physically interacts with several proteins essential for mRNA splicing and

transport (Allemand et al., 2002). Given that SR-rich proteins facilitate diverse RNA processing

events, there are many potential Tnpo-SR cargo proteins that could influence cyst formation and

GSC self-renewal, which are heavily influenced by translational control (Bradley et al., 2015;
Teixeira and Lehmann, 2019). Our future studies will seek to identify Tnpo-SR and Cdm-
interacting proteins, which should provide additional mechanistic details regarding how importins
regulate germ cell mitotic expansion.

The number of cyst divisions is regulated by Ran-GTP and β-importins.

Precise spatiotemporal control over cell cycle regulators underlie the transition to the
oocyte fate. In support of this, the mitotic cyclin CycA is clearly critical to support differentiation.
Failure to degrade CycA leads to self-renewal failure in GSCs, and extra mitotic divisions and loss
of oocyte identity in cyst cells (Chen et al., 2009; Ji et al., 2017; Lilly et al., 2000; Morris et al.,
2005; Ohlmeyer and Schüpbach, 2003; Sugimura and Lilly, 2006). Moreover, the deubiquitinase
encoded by *bag of marbles* (*bam*), one of the few known differentiation factors in *Drosophila* germ
cells, is necessary for the transition from GSC to cystoblast and for cystoblast divisions, at least in
part because it stabilizes CycA expression in dividing cysts (Ji et al., 2017). CycA/Cdk1 activity
is very high in early cyst divisions but decrease as cysts approach the terminal mitotic division
(Hinnant et al., 2017; Lilly et al., 2000). Since cystoblasts divide exactly four times, prior studies
posited the model that cystoblasts autonomously limit the number of divisions through a molecular
counting mechanism that involves Bam, CycA, and the fusome (Hinnant et al., 2020; Huynh, 2006;
Ji et al., 2017; King, 1970; Lilly et al., 2000; McKearin, 1997). While a counting mechanism would
limit the number of mitotic divisions, it must also promote modification of contractile ring proteins
to block cytokinesis, stabilize intercellular bridges into stable ring canals, and maintain the
biosynthesis of the fusome (McKearin, 1997).

We suggest that the Ran/β-importin intracellular transport system is integrated into the
CycA-dependent cell division counting machinery. Depletion of *Ran*, *cdm*, or *Tnpo-SR* generated
egg chambers with abnormal numbers of cells (Table 1), some on the order of 2^n reminiscent of
cyclin mutants (Lilly et al., 2000; Mathieu et al., 2013; Morris et al., 2005; Ohlmeyer and
Schüpbach, 2003), and others deviating from this pattern, more similar to mutants which disrupt
intercellular bridge formation and fusome biogenesis (de Cuevas et al., 1996; Lin et al., 1994; Yue
and Spradling, 1992). Although we cannot establish a clear temporal sequence of events mediated
by Ran from our current data, we envision two possible mechanisms of action. We favor the
hypothesis that Ran mediates intracellular movement of CycA itself or a factor that promotes CycA

degradation. Binding by Tnpo-SR or Cdm could sequester the cell cycle regulator, preventing activation by sterically inhibiting interaction with a kinase or by blocking protein accumulation at the fusome. High levels of Ran-GTP levels would promote release of the Tnpo-SR cargo, freeing it for activation. In the absence of Tnpo-SR or Cdm, where mitotic regulators cannot be delivered to Ran, CycA levels instead accumulate at the fusome and in the cytoplasm, promoting continued mitotic division. Alternatively, Ran-GTP could regulate contractile protein localization and cell polarization, as has been demonstrated for human Ran in mammalian somatic cells (Beaudet et al., 2017; Silverman-Gavrila et al., 2008).

Tnpo-SR and Cdm may have Ran-independent roles in promoting cyst development.

Given the close association between Ran and β-importins, we hypothesized that depletion of *Ran* by RNAi, or over-expression of dominant negative Ran^{T24N} mutant would phenocopy β-importin depletion. Yet we observed phenotypes unique to loss of *Tnpo-SR* and *cdm*. In *Tnpo-SR*- and *Cdm*- depleted germ cells, delayed maturation of the fusomes and ring canals appears to create weak connections between cells, resulting in premature abscission or collapsed intercellular bridges (Figure 6M). Interestingly, we see fusomes that are fragile and unbranched (Figure 5E-H), but also germline tumors containing cells with immature fusome morphology resembling that of 2-cell cysts (Figure 2G-H). This may be the result of altered timing or different concentrations of the β-importins in the first division of the cystoblast versus the remaining three, perhaps reflecting differing sensitivity to abscission across the mitotic divisions (Mathieu et al., 2013; McKearin and Ohlstein, 1995). Cyst cell fusomes are composed of endoplasmic reticulum-like vesicles and cytoskeletal elements that originate from remnants of the mitotic spindle midbody (de Cuevas and Spradling, 1998; Koch and King, 1966; Koch et al., 1967; Lin and Spradling, 1995; Mahowald, 1971). This suggests Ran-independent functions of Tnpo-SR and Cdm in mitotically dividing germ cells, perhaps by sequestering negative regulators of fusome maturation or cytokinesis. We speculate that Tnpo-SR and Cdm promote reorganization of the germ cell cytoskeleton after mitotic division, perhaps by shuttling cytoskeletal proteins back to the fusome.

Ran-GTP and β-importins may function as a conserved molecular timer for the mitotic-to-meiotic transition.

The basic steps in cyst formation have been well described at the morphological level,

particularly in insects (de Cuevas et al., 1997; Lu et al., 2017; Pepling and Lei, 2018; Yamashita,

2018). Yet how cyst formation is guided molecularly remains an open area for study. Insects

provide excellent models, as many exhibit unique mechanisms for achieving cyst formation

(Büning, 1994; de Cuevas et al., 1997; Eastin et al., 2020; Kubrakiewicz, 1997). Given the level

of conservation in cell cycle control across species, we propose that Ran and β-importins may also

play important roles in higher eukaryotes, connecting the intricate programs of cell division, oocyte

differentiation, and cell polarization. This may be of particular consequence for mouse and human

oocytes, in which cyst breakdown occurs prior to meiotic onset (Pepling and Lei, 2018).

Understanding how Ran and importins control cyst formation and breakdown may help elucidate

how the oocyte reserve is established in humans, offering new potential avenues for treatment of

infertility.

MATERIALS AND METHODS

Drosophila strains and husbandry

Flies were maintained at 22°-25°C in standard medium (cornmeal/molasses/yeast/agar)

(NutriFly MF; Genesee Scientific). Female progeny were collected one to two days after eclosion

and maintained on standard medium for 3, 6, and 10 days at 25°C; flies were fed wet yeast paste

for 2-3 days (changed daily) prior to ovary dissection. A complete list of fly lines used in this study

are provided in Supp. Table S1.

Tissue-specific RNA interference (RNAi) generation

All RNAi lines used in this study were carried in *pVALIUM20* or *pVALIUM22* transposons

for maximum germline efficiency (Supp. Table S1). *UAS-Tnpo-SR^{V22}* was generated as described

(www.flyrnai.org/TRiP-HOME.html) (Ni et al., 2011). Primers were designed against the full-

length *Tnpo-SR* RNA via the Designer of Small Interfering RNA tool

(http://biodev.extra.cea.fr/DSIR/DSIRhtml) using default settings (21 nt siRNA; score threshold

90). A sequence with a score of 94.6 and 0 predicted off-targets was selected for hairpin generation

(passenger strand, CGATCCCGTTTACTGGATAGA; guide strand,

TCTATCCAGTAAACGGGATC). Primers were designed according to TRiP recommendations,

annealed to form double-stranded oligos, and ligated into EcoRI-HF/NheI-HF-digested

pVALIUM22. Transgenic flies (*UAS-Tnpo-SR^{V22}*) were generated by phiC31 site-specific integrase

into the *attP2* site on the third chromosome.

For RNAi experiments, germline knock-down was facilitated by expressing the germline-

specific *nos-GAL4::VP16-nos.UTR* (referred to throughout as *nos-Gal4*) (Rørth, 1998; Van Doren

et al., 1998). Driver expression was confirmed using *w*; nos-Gal4, UASp-tubGFP* (BDSC #7253)

(Grieder et al., 2000). Females carrying *nos-Gal4* alone were used as controls.

Drosophila genetic mosaic generation

For genetic mosaic analyses using *flippase (FLP)/FLP recognition target (FRT)*, we

obtained the following mutant alleles of *FRT*-containing chromosome arms: *Tnpo-SR^{KG04870}*

[Kyoto *Drosophila* Stock Center (KDSC) #111581] (Bellen et al., 2004) and *Tnpo-SR^{LL05552}*

(KDSC #141628) (Schuldiner et al., 2008). Other genetic tools are described in FlyBase version

FB2020_05 (Thurmond et al., 2019). Genetic mosaics were generated by *FLP/FRT*-mediated

recombination in 2-3-day old females carrying a mutant allele in trans to a wildtype allele (linked

to a *Ubi-GFP* marker) on homologous *FRT* arms, and a *hs-FLP* transgene, as described (Laws and

Drummond-Barbosa, 2015). Flies were heat shocked at 37°C two times per day for 3 days and

incubated at 25°C for 4, 8, or 12 days and supplemented with wet yeast paste on the last 2 days

prior to dissection. Wild-type alleles were used for the generation of control (mock) mosaics.

Germ cell analyses

GSCs were identified based on the juxtaposition of their fusomes to the junction with

adjacent cap cells (de Cuevas and Spradling, 1998). In *FLP/FRT* mutants, GSC loss was measured

two ways. First, as the percentage of total mosaic germaria showing evidence of recent stem cell

loss; namely, the presence of GFP-negative daughters (cystoblasts/cysts generated from an original

GFP-negative stem cell) in the absence of the GFP-negative mother stem cell (Laws and

Drummond-Barbosa, 2015). Second, as the percentage of total germaria analyzed that had at least

one GFP-negative GSC. All results were subjected to Chi-Square analyses using Microsoft Excel.

In RNAi progeny, GSC loss was measured as the average number of GSCs present over three

timepoints. Results were subjected to Student's two-tailed T-test comparing driver controls versus

driver>UAS-RNAi experimental groups at each timepoint individually using Microsoft Excel.

For analyses involving the egg chambers outside of the germarium, data was collected from

the first three egg chambers posteriorly located from the germarium. Oocytes were identified by

the presence of anti-Orb antibodies. The number of cells and ring canals per egg chamber were

calculated based on visualization of ring canals with anti-pTyr and/or anti-Hts-RC antibodies and

cell nuclei with DAPI. Results were subjected to Student's two-tailed T-test comparing driver

controls versus *driver>UAS-RNAi* experimental groups at each timepoint individually using

Microsoft Excel and Prism (GraphPad).

Immunofluorescence and microscopy

Ovaries were prepared for immunofluorescence microscopy as described (Grieder et al.,

2000; Hinnant et al., 2017). In the standard protocol, ovaries were dissected and teased apart in

Grace's medium without additives (Caisson Labs) and fixed in 5.3% formaldehyde in Grace's

medium for 13 minutes at room temperature. They were then washed extensively in phosphate-

buffered saline (PBS, pH 7.4; Fisher) with 0.1% Triton X-100, and blocked for three hours in

blocking solution [5% bovine serum albumin (Sigma), 5% normal goat serum (MP Biomedicals),

and 0.1% Triton X-100 in PBS] at room temperature. To detect cells in S phase, dissected ovaries

were incubated for one hour at room temperature in Grace's media containing 10 μM 5-ethynyl-

2′-deoxyuridine (EdU; Life Technologies). The following primary antibodies were diluted in

blocking solution and used overnight at 4°C: chicken anti-GFP (#13970, Abcam; 1:2000), rabbit

anti-dsRed (#632496, Clontech; 1:500), rabbit anti-phosphoHistone H3 (pHH3; #06-570,

Millipore; 1:200), rat anti-E-Cadherin (DCAD2, Developmental Studies Hybridoma Bank

(DSHB); 1:20), mouse anti-Orb (4H8 and 6H4; DSHB; 1:500), mouse anti-BicD (1B11 and 4C2;

DSHB; 1:10), mouse anti-p-Tyrosine (Millipore; 1:100), mouse anti-Hts-RC (DSHB; 1:10),

mouse anti-CycA (A12; DSHB; 1:50), rabbit anti-C(3)G (a gift from M. Lilly; 1:3000), rat anti-

HA (3F10; Sigma; 1:25), mouse anti-α-Tubulin-AF555 (DM1A-AF555; Millipore; 1:100), and

rabbit anti-Anillin (a gift from J. Brill; 1:50). Mouse anti-Sxl (m18; DSHB; 1:500) was also used

but fixed in a 4% formaldehyde in Grace's medium. Primary antibodies mouse anti-Hts (1B1,

DSHB; 1:10), and mouse anti-Lamin C (LC28.26, DSHB; 1:100) were incubated over two nights

at 4°C. All primary antibodies are followed by a two hour incubation at room temperature with

AlexaFluor 488-, 568- or 633-conjugated goat species-specific secondary antibodies (Life

Technologies; 1:200). EdU was detected (if necessary) using AlexaFluor-594 or -647 via Click-It chemistry, following the manufacturer's recommendations (Life Technologies).

For stabilizing structures such as mitotic spindles, ovaries were dissected and teased apart in room temperature equilibrated Grace's medium and fixed for 10 minutes at room temperature using 350µl Grace's Medium and 200µl of 16% formaldehyde. They were then extensively washed in PBS with 0.1% Triton X-100 and blocked at room temperature for at least 30 minutes in blocking solution (2.5mL NGS, 5mL Triton, 42.5mL 1xPBS). Primary antibody incubations took place overnight at 4°C followed by extensive washing in PBS with 0.1% Triton X-100 and another blocking period of at least 30 minutes at room temperature. Detection of secondary antibodies proceeded as described above.

All ovary samples were stained with 0.5 µg/ml 4'-6-diamidino-2-phenylindole (DAPI; Sigma) in 0.1% Triton X-100 in PBS, and mounted in 90% glycerol mixed with 20% n-propyl gallate (Sigma). Confocal z-stacks (1 µm optical sections) were collected with either the Zeiss LSM700 laser scanning microscope using ZEN Black 2012 software or the Zeiss LSM800 laser scanning microscope using ZEN Blue 2012. Images were analyzed using Zeiss ZEN software, and minimally and equally enhanced via histogram using ZEN and Adobe Photoshop Creative Suite.

ACKNOWLEDGMENTS

Many thanks to K. McKim, B. Riggs, K. Roper, J.-Q. Ni, M. Lilly, D. Drummond-Barbosa, M. Buszczak, L. Cooley, and J. Brill for their kind gifts of fly stocks and antibodies. We are grateful to the Bloomington, Vienna, and Kyoto *Drosophila* Stock Centers, the Drosophila Genomics Resource Center, the Harvard Transgenic RNAi Project, and the University of Iowa Developmental Studies Hybridoma Bank for making fly lines, antibodies, and plasmids available, and BestGene for *Drosophila* transgenesis and screening. Many thanks to Tianna Van Cura and members of the Ables laboratory for assistance with building tools, helpful discussions, and critical reading of this manuscript.

COMPETING INTERESTS

The authors declare no competing conflicts of interest.

FUNDING

602   This work was supported by National Institutes of Health R15-GM117502 (E. T. A.).

DATA AVAILABILITY

605   Fly strains and plasmids are available upon request. The authors affirm that all data necessary for

606   confirming the conclusions of the article are present within the article and figures.

REFERENCES

Ables, E. T. and Drummond-Barbosa, D. (2013). Cyclin E controls Drosophila female germline stem cell
maintenance independently of its role in proliferation by modulating responsiveness to niche
signals. *Development* **140**, 530-540.
Ables, E. T., Hwang, G. H., Finger, D. S., Hinnant, T. D. and Drummond-Barbosa, D. (2016). A Genetic
Mosaic Screen Reveals Ecdysone-Responsive Genes Regulating Drosophila Oogenesis. *G3*
(Bethesda, Md.) **6**, 2629-2642.
Allemand, E., Dokudovskaya, S., Bordonné, R. and Tazi, J. (2002). A conserved Drosophila transportin-
serine/arginine-rich (SR) protein permits nuclear import of Drosophila SR protein splicing factors
and their antagonist repressor splicing factor 1. *Molecular biology of the cell* **13**, 2436-2447.
Ameku, T. and Niwa, R. (2016). Mating-Induced Increase in Germline Stem Cells via the Neuroendocrine
System in Female Drosophila. *PLoS genetics* **12**, e1006123.
Beaudet, D., Akhshi, T., Phillipp, J., Law, C. and Piekny, A. (2017). Active Ran regulates anillin function
during cytokinesis. *Molecular biology of the cell* **28**, 3517-3531.
Beaudet, D., Pham, N., Skaik, N. and Piekny, A. (2020). Importin binding mediates the intramolecular
regulation of anillin during cytokinesis. *Molecular biology of the cell* **31**, 1124-1139.
Bellen, H. J., Levis, R. W., Liao, G., He, Y., Carlson, J. W., Tsang, G., Evans-Holm, M., Hiesinger, P. R.,
Schulze, K. L., Rubin, G. M., et al. (2004). The BDGP gene disruption project: single transposon
insertions associated with 40% of Drosophila genes. *Genetics* **167**, 761-781.
Bradley, T., Cook, M. E. and Blanchette, M. (2015). SR proteins control a complex network of RNA-
processing events. *Rna* **21**, 75-92.
Büning, J. (1994). *The Insect Ovary: Ultrastructure, previtellogenic growth and evolution*: Springer,
Dordrecht.
Buszczak, M., Paterno, S., Lighthouse, D., Bachman, J., Planck, J., Owen, S., Skora, A. D., Nystul, T. G.,
Ohlstein, B., Allen, A., et al. (2007). The carnegie protein trap library: a versatile tool for
Drosophila developmental studies. *Genetics* **175**, 1505-1531.
Cautain, B., Hill, R., de Pedro, N. and Link, W. (2015). Components and regulation of nuclear transport
processes. *The FEBS journal* **282**, 445-462.
Cesario, J. and McKim, K. S. (2011). RanGTP is required for meiotic spindle organization and the
initiation of embryonic development in Drosophila. *J Cell Sci* **124**, 3797-3810.
Chaigne, A., Labouesse, C., White, I. J., Agnew, M., Hannezo, E., Chalut, K. J. and Paluch, E. K. (2020).
Abscission Couples Cell Division to Embryonic Stem Cell Fate. *Developmental cell* **55**, 195-
208.e195.
Chau, J., Kulnane, L. S. and Salz, H. K. (2009). Sex-lethal facilitates the transition from germline stem cell
to committed daughter cell in the Drosophila ovary. *Genetics* **182**, 121-132.
Chen, A., Akhshi, T. K., Lavoie, B. D. and Wilde, A. (2015). Importin β2 Mediates the Spatio-temporal
Regulation of Anillin through a Noncanonical Nuclear Localization Signal. *The Journal of*
biological chemistry **290**, 13500-13509.

Chen, D., Wang, Q., Huang, H., Xia, L., Jiang, X., Kan, L., Sun, Q. and Chen, D. (2009). Effete-mediated degradation of Cyclin A is essential for the maintenance of germline stem cells in Drosophila. *Development* **136**, 4133-4142.

Chen, J., Call, G. B., Beyer, E., Bui, C., Cespedes, A., Chan, A., Chan, J., Chan, S., Chhabra, A., Dang, P., et al. (2005). Discovery-based science education: functional genomic dissection in Drosophila by undergraduate researchers. *PLoS Biol* **3**, e59.

Cheng, J., Türkel, N., Hemati, N., Fuller, M. T., Hunt, A. J. and Yamashita, Y. M. (2008). Centrosome misorientation reduces stem cell division during ageing. *Nature* **456**, 599-604.

de Cuevas, M., Lee, J. K. and Spradling, A. C. (1996). alpha-spectrin is required for germline cell division and differentiation in the Drosophila ovary. *Development* **122**, 3959-3968.

de Cuevas, M., Lilly, M. A. and Spradling, A. C. (1997). Germline cyst formation in Drosophila. *Annual review of genetics* **31**, 405-428.

de Cuevas, M. and Spradling, A. C. (1998). Morphogenesis of the Drosophila fusome and its implications for oocyte specification. *Development* **125**, 2781-2789.

Deng, W. and Lin, H. (1997). Spectrosomes and fusomes anchor mitotic spindles during asymmetric germ cell divisions and facilitate the formation of a polarized microtubule array for oocyte specification in Drosophila. *Developmental biology* **189**, 79-94.

Drutovic, D., Duan, X., Li, R., Kalab, P. and Solc, P. (2020). RanGTP and importin β regulate meiosis I spindle assembly and function in mouse oocytes. *The EMBO journal* **39**, e101689.

Eastin, K. J., Huang, A. P. and Ferree, P. M. (2020). A novel pattern of germ cell divisions in the production of hymenopteran insect eggs. *Biol Lett* **16**, 20200137.

Eikenes Å, H., Malerød, L., Lie-Jensen, A., Sem Wegner, C., Brech, A., Liestøl, K., Stenmark, H. and Haglund, K. (2015). Src64 controls a novel actin network required for proper ring canal formation in the Drosophila male germline. *Development* **142**, 4107-4118.

Fichelson, P., Jagut, M., Lepanse, S., Lepesant, J. A. and Huynh, J. R. (2010). lethal giant larvae is required with the par genes for the early polarization of the Drosophila oocyte. *Development* **137**, 815-824.

Forbes, D. J., Travesa, A., Nord, M. S. and Bernis, C. (2015). Nuclear transport factors: global regulation of mitosis. *Current opinion in cell biology* **35**, 78-90.

Frémont, S. and Echard, A. (2018). Membrane Traffic in the Late Steps of Cytokinesis. *Current biology : CB* **28**, R458-R470.

Geles, K. G. and Adam, S. A. (2001). Germline and developmental roles of the nuclear transport factor importin alpha3 in C. elegans. *Development* **128**, 1817-1830.

Goldbach, P., Wong, R., Beise, N., Sarpal, R., Trimble, W. S. and Brill, J. A. (2010). Stabilization of the actomyosin ring enables spermatocyte cytokinesis in Drosophila. *Molecular biology of the cell* **21**, 1482-1493.

Grieder, N. C., de Cuevas, M. and Spradling, A. C. (2000). The fusome organizes the microtubule network during oocyte differentiation in Drosophila. *Development* **127**, 4253-4264.

Guertin, D. A., Trautmann, S. and McCollum, D. (2002). Cytokinesis in eukaryotes. *Microbiology and molecular biology reviews : MMBR* **66**, 155-178.

Haglund, K., Nezis, I. P., Lemus, D., Grabbe, C., Wesche, J., Liestøl, K., Dikic, I., Palmer, R. and Stenmark, H. (2010). Cindr interacts with anillin to control cytokinesis in Drosophila melanogaster. *Current biology : CB* **20**, 944-950.

Haglund, K., Nezis, I. P. and Stenmark, H. (2011). Structure and functions of stable intercellular bridges formed by incomplete cytokinesis during development. *Communicative & integrative biology* **4**, 1-9.

Hinnant, T. D., Alvarez, A. A. and Ables, E. T. (2017). Temporal remodeling of the cell cycle accompanies differentiation in the Drosophila germline. *Developmental biology* **429**, 118-131.

Hinnant, T. D., Merkle, J. A. and Ables, E. T. (2020). Coordinating Proliferation, Polarity, and Cell Fate in
the Drosophila Female Germline. *Frontiers in cell and developmental biology* **8**, 19.
Hogarth, C., Itman, C., Jans, D. A. and Loveland, K. L. (2005). Regulated nucleocytoplasmic transport in
spermatogenesis: a driver of cellular differentiation? *BioEssays : news and reviews in molecular,*
cellular and developmental biology **27**, 1011-1025.
Hughes, S. E., Miller, D. E., Miller, A. L. and Hawley, R. S. (2018). Female Meiosis: Synapsis,
Recombination, and Segregation in Drosophila melanogaster. *Genetics* **208**, 875-908.
Huynh, J.-R. (2006). Fusome as a Cell-Cell Communication Channel of Drosophila Ovarian Cyst. In *Cell-*
Cell Channels (ed. F. Baluska, D. Volkmann & P. W. Barlow), pp. 217-235. New York, NY: Springer
New York.
Ji, S., Li, C., Hu, L., Liu, K., Mei, J., Luo, Y., Tao, Y., Xia, Z., Sun, Q. and Chen, D. (2017). Bam-dependent
deubiquitinase complex can disrupt germ-line stem cell maintenance by targeting cyclin A.
Proceedings of the National Academy of Sciences of the United States of America **114**, 6316-
6321.
Kimura, M. and Imamoto, N. (2014). Biological significance of the importin-β family-dependent
nucleocytoplasmic transport pathways. *Traffic (Copenhagen, Denmark)* **15**, 727-748.
Kimura, M., Morinaka, Y., Imai, K., Kose, S., Horton, P. and Imamoto, N. (2017). Extensive cargo
identification reveals distinct biological roles of the 12 importin pathways. *eLife* **6**.
King, R. C. R. C. (1970). Ovarian development in Drosophila melanogaster. New York: Academic Press.
Koch, E. A. and King, R. C. (1966). The origin and early differentiation of the egg chamber of Drosophila
melanogaster. *J Morphol* **119**, 283-303.
Koch, E. A., Smith, P. A. and King, R. C. (1967). The division and differentiation of Drosophila cystocytes.
J Morphol **121**, 55-70.
Kubrakiewicz, J. (1997). Germ cells cluster organization in polytrophic ovaries of neuroptera. *Tissue and*
Cell **29**, 221-228.
Lai, M. C., Lin, R. I. and Tarn, W. Y. (2001). Transportin-SR2 mediates nuclear import of phosphorylated
SR proteins. *Proc Natl Acad Sci U S A* **98**, 10154-10159.
Laws, K. M. and Drummond-Barbosa, D. (2015). Genetic Mosaic Analysis of Stem Cell Lineages in the
Drosophila Ovary. *Methods in molecular biology (Clifton, N.J.)* **1328**, 57-72.
Lighthouse, D. V., Buszczak, M. and Spradling, A. C. (2008). New components of the Drosophila fusome
suggest it plays novel roles in signaling and transport. *Developmental biology* **317**, 59-71.
Lilly, M. A., de Cuevas, M. and Spradling, A. C. (2000). Cyclin A associates with the fusome during
germline cyst formation in the Drosophila ovary. *Developmental biology* **218**, 53-63.
Lin, H. and Spradling, A. C. (1995). Fusome asymmetry and oocyte determination in Drosophila. *Dev*
Genet **16**, 6-12.
Lin, H., Yue, L. and Spradling, A. C. (1994). The Drosophila fusome, a germline-specific organelle,
contains membrane skeletal proteins and functions in cyst formation. *Development* **120**, 947-
956.
Lippai, M., Tirián, L., Boros, I., Mihály, J., Erdélyi, M., Belecz, I., Máthé, E., Pósfai, J., Nagy, A., Udvardy,
A., et al. (2000). The Ketel gene encodes a Drosophila homologue of importin-beta. *Genetics*
156, 1889-1900.
Loveland, K. L., Major, A. T., Butler, R., Young, J. C., Jans, D. A. and Miyamoto, Y. (2015). Putting things
in place for fertilization: discovering roles for importin proteins in cell fate and spermatogenesis.
Asian journal of andrology **17**, 537-544.
Lu, K., Jensen, L., Lei, L. and Yamashita, Y. M. (2017). Stay Connected: A Germ Cell Strategy. *Trends in*
genetics : TIG **33**, 971-978.

Maertens, G. N., Cook, N. J., Wang, W., Hare, S., Gupta, S. S., Öztop, I., Lee, K., Pye, V. E., Cosnefroy, O., Snijders, A. P., et al. (2014). Structural basis for nuclear import of splicing factors by human Transportin 3. *Proc Natl Acad Sci U S A* **111**, 2728-2733.

Mahowald, A. P. (1971). The formation of ring canals by cell furrows in Drosophila. *Z Zellforsch Mikrosk Anat* **118**, 162-167.

Mason, D. A., Fleming, R. J. and Goldfarb, D. S. (2002). Drosophila melanogaster importin alpha1 and alpha3 can replace importin alpha2 during spermatogenesis but not oogenesis. *Genetics* **161**, 157-170.

Máthé, E., Bates, H., Huikeshoven, H., Deák, P., Glover, D. M. and Cotterill, S. (2000). Importin-alpha3 is required at multiple stages of Drosophila development and has a role in the completion of oogenesis. *Developmental biology* **223**, 307-322.

Máthé, E., Inoue, Y. H., Palframan, W., Brown, G. and Glover, D. M. (2003). Orbit/Mast, the CLASP orthologue of Drosophila, is required for asymmetric stem cell and cystocyte divisions and development of the polarised microtubule network that interconnects oocyte and nurse cells during oogenesis. *Development* **130**, 901-915.

Mathieu, J., Cauvin, C., Moch, C., Radford, S. J., Sampaio, P., Perdigoto, C. N., Schweisguth, F., Bardin, A. J., Sunkel, C. E., McKim, K., et al. (2013). Aurora B and cyclin B have opposite effects on the timing of cytokinesis abscission in Drosophila germ cells and in vertebrate somatic cells. *Developmental cell* **26**, 250-265.

Mathieu, J. and Huynh, J. R. (2017). Monitoring complete and incomplete abscission in the germ line stem cell lineage of Drosophila ovaries. *Methods in cell biology* **137**, 105-118.

Matias, N. R., Mathieu, J. and Huynh, J. R. (2015). Abscission is regulated by the ESCRT-III protein shrub in Drosophila germline stem cells. *PLoS genetics* **11**, e1004653.

Matova, N. and Cooley, L. (2001). Comparative aspects of animal oogenesis. *Developmental biology* **231**, 291-320.

McKearin, D. (1997). The Drosophila fusome, organelle biogenesis and germ cell differentiation: if you build it. *BioEssays : news and reviews in molecular, cellular and developmental biology* **19**, 147-152.

McKearin, D. and Ohlstein, B. (1995). A role for the Drosophila bag-of-marbles protein in the differentiation of cystoblasts from germline stem cells. *Development* **121**, 2937-2947.

Megraw, T. L. and Kaufman, T. C. (2000). The centrosome in Drosophila oocyte development. *Current topics in developmental biology* **49**, 385-407.

Mihalas, B. P., Western, P. S., Loveland, K. L., McLaughlin, E. A. and Holt, J. E. (2015). Changing expression and subcellular distribution of karyopherins during murine oogenesis. *Reproduction (Cambridge, England)* **150**, 485-496.

Morris, J. Z., Hong, A., Lilly, M. A. and Lehmann, R. (2005). twin, a CCR4 homolog, regulates cyclin poly(A) tail length to permit Drosophila oogenesis. *Development* **132**, 1165-1174.

Nguyen, L. T. S. and Robinson, D. N. (2020). The Unusual Suspects in Cytokinesis: Fitting the Pieces Together. *Frontiers in cell and developmental biology* **8**, 441.

Ni, J. Q., Zhou, R., Czech, B., Liu, L. P., Holderbaum, L., Yang-Zhou, D., Shim, H. S., Tao, R., Handler, D., Karpowicz, P., et al. (2011). A genome-scale shRNA resource for transgenic RNAi in Drosophila. *Nat Methods* **8**, 405-407.

Ohlmeyer, J. T. and Schüpbach, T. (2003). Encore facilitates SCF-Ubiquitin-proteasome-dependent proteolysis during Drosophila oogenesis. *Development* **130**, 6339-6349.

Okada, N., Ishigami, Y., Suzuki, T., Kaneko, A., Yasui, K., Fukutomi, R. and Isemura, M. (2008). Importins and exportins in cellular differentiation. *J Cell Mol Med* **12**, 1863-1871.

Ong, S. and Tan, C. (2010). Germline cyst formation and incomplete cytokinesis during Drosophila melanogaster oogenesis. *Developmental biology* **337**, 84-98.

Ozugergin, I. and Piekny, A. (2020). Complementary functions for the Ran gradient during division.
Small GTPases, 1-11.
Pepling, M. and Lei, L. (2018). Germ Cell Nests and Germline Cysts. In *Encyclopedia of Reproduction*
(Second Edition) (ed. M. K. Skinner), pp. 159-166. Oxford: Academic Press.
Pepling, M. E., de Cuevas, M. and Spradling, A. C. (1999). Germline cysts: a conserved phase of germ
cell development? *Trends in Cell Biology* **9**, 257-262.
Perkins, L. A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., Yang-Zhou, D., Flockhart, I., Binari,
R., Shim, H. S., et al. (2015). The Transgenic RNAi Project at Harvard Medical School: Resources
and Validation. *Genetics* **201**, 843-852.
Petrella, L. N., Smith-Leiker, T. and Cooley, L. (2007). The Ovhts polyprotein is cleaved to produce
fusome and ring canal proteins required for Drosophila oogenesis. *Development* **134**, 703-712.
Piekny, A. J. and Maddox, A. S. (2010). The myriad roles of Anillin during cytokinesis. *Seminars in cell &*
developmental biology **21**, 881-891.
Pollard, T. D. and O'Shaughnessy, B. (2019). Molecular Mechanism of Cytokinesis. *Annu Rev Biochem*
88, 661-689.
Robinson, D. N., Cant, K. and Cooley, L. (1994). Morphogenesis of Drosophila ovarian ring canals.
Development **120**, 2015-2025.
Röper, K. (2007). Rtnl1 is enriched in a specialized germline ER that associates with ribonucleoprotein
granule components. *J Cell Sci* **120**, 1081-1092.
Rørth, P. (1998). Gal4 in the Drosophila female germline. *Mech Dev* **78**, 113-118.
Sazer, S. and Dasso, M. (2000). The ran decathlon: multiple roles of Ran. *J Cell Sci* **113 (Pt 7)**, 1111-1118.
Schuldiner, O., Berdnik, D., Levy, J. M., Wu, J. S., Luginbuhl, D., Gontang, A. C. and Luo, L. (2008).
piggyBac-based mosaic screen identifies a postmitotic function for cohesin in regulating
developmental axon pruning. *Developmental cell* **14**, 227-238.
Silverman-Gavrila, R. V., Hales, K. G. and Wilde, A. (2008). Anillin-mediated targeting of peanut to
pseudocleavage furrows is regulated by the GTPase Ran. *Molecular biology of the cell* **19**, 3735-
3744.
Snapp, E. L., Iida, T., Frescas, D., Lippincott-Schwartz, J. and Lilly, M. A. (2004). The fusome mediates
intercellular endoplasmic reticulum connectivity in Drosophila ovarian cysts. *Molecular biology*
of the cell **15**, 4512-4521.
Storto, P. D. and King, R. C. (1989). The role of polyfusomes in generating branched chains of cystocytes
during Drosophila oogenesis. *Dev Genet* **10**, 70-86.
Sugimura, I. and Lilly, M. A. (2006). Bruno inhibits the expression of mitotic cyclins during the prophase
I meiotic arrest of Drosophila oocytes. *Developmental cell* **10**, 127-135.
Teixeira, F. K. and Lehmann, R. (2019). Translational Control during Developmental Transitions. *Cold*
Spring Harb Perspect Biol **11**.
Thurmond, J., Goodman, J. L., Strelets, V. B., Attrill, H., Gramates, L. S., Marygold, S. J., Matthews, B.
B., Millburn, G., Antonazzo, G., Trovisco, V., et al. (2019). FlyBase 2.0: the next generation.
Nucleic Acids Res **47**, D759-d765.
Tirián, L., Puro, J., Erdélyi, M., Boros, I., Papp, B., Lippai, M. and Szabad, J. (2000). The Ketel(D)
dominant-negative mutations identify maternal function of the Drosophila importin-beta gene
required for cleavage nuclei formation. *Genetics* **156**, 1901-1912.
Trieselmann, N. and Wilde, A. (2002). Ran localizes around the microtubule spindle in vivo during
mitosis in Drosophila embryos. *Current biology : CB* **12**, 1124-1129.
Van Doren, M., Williamson, A. L. and Lehmann, R. (1998). Regulation of zygotic gene expression in
Drosophila primordial germ cells. *Current biology : CB* **8**, 243-246.

Villányi, Z., Debec, A., Timinszky, G., Tirián, L. and Szabad, J. (2008). Long persistence of importin-beta
explains extended survival of cells and zygotes that lack the encoding gene. *Mech Dev* **125**, 196-
206.
Xu, T. and Rubin, G. M. (1993). Analysis of genetic mosaics in developing and adult Drosophila tissues.
Development **117**, 1223-1237.
Yamashita, Y. M. (2018). Subcellular Specialization and Organelle Behavior in Germ Cells. *Genetics* **208**,
19-51.
Yue, L. and Spradling, A. C. (1992). hu-li tai shao, a gene required for ring canal formation during
Drosophila oogenesis, encodes a homolog of adducin. *Genes & development* **6**, 2443-2454.
Zielke, N., Korzelius, J., van Straaten, M., Bender, K., Schuhknecht, G. F. P., Dutta, D., Xiang, J. and
Edgar, B. A. (2014). Fly-FUCCI: A versatile tool for studying cell proliferation in complex tissues.
Cell Rep **7**, 588-598.

TABLES

Table 1. Frequency of ovarioles or egg chambers that showcase germline phenotypes in β-
importin- and Ran-depleted ovaries.

genotype	% of agametic ovarioles	average # of GSCs at 3 days after eclosion[a]	% of egg chambers that do not equal 16 cells	% of egg chambers that do not equal 2^n germ cells	% of 16-cell egg chambers without an oocyte
driver control	0	2.8	0.64	0	0
$Fs(2)Ket^{GL01273}$	3.6	1.8 ***	3.9	1.3	0
$Ran^{GL01341}$	1.5	1.1 ***	11.31***	2.6	4.0
$cdm^{HMS02846}$	33.1	0.27 ***	32.0 ***	55.0	0
$Tnpo\text{-}SR^{V22}$	2.5	1.5 ***	46.0 ***	29.3	33.8
$Ran^{HA.1}$	0	2.9	0	0	0
$Ran^{HA.2}$	0	2.4	0	0	0
$Ran^{T24N.1}$	63.8	0.25 ***	15.6 ***	12.5	23.5
$Ran^{T24N.2}$	49.6	0.38 ***	19.5***	9.7	11.5
$Ran^{Q69L.1}$	0	1.9 ***	0	0	0
$Ran^{Q69L.2}$	2.0	2.0 ***	6.4 **	2.2	0

851   [a]See Suppl. Figure S1.

852   **$p < 0.01$, ***$p < 0.001$. Students two-tailed t-test and Chi-squared test.

853   **FIGURES**

855   **Figure 1. An RNAi screen identified Ran and β-importins necessary for germ cell**

856   **development.**

(A) The *Drosophila* ovary is composed of ovarioles harboring a germarium (top) and older egg chambers containing developing oocytes (bottom). In the germarium, germline stem cells (GSCs; pink) are adjacent to cap cells (yellow) and escort cells (green). GSCs divide to form cystoblasts (blue) that divide four times to form 16-cell cysts (blue). Cysts are composed of 15 nurse cells and one oocyte (dark blue), connected by 15 intercellular bridges that can be visualized by ring canals and the fusome (red) that transects each bridge. (B-D) Maximum intensity projections of *nos-Gal4* control (B), *nos-Gal4>Tnpo/CG8219^{HMJ23009}* (C), or *nos-Gal4>cdm^{HMS02846}* (D) knockdown ovaries labeled with DAPI (white; nuclei). Arrows indicate oviducts (ovi). Scale bar, 200 μm. (E-G) Egg chambers of driver control (E), *nos-Gal4>Tnpo-SR^{V22}* (F), and *nos-Gal4>cdm^{HMS02846}* (G) ovarioles labeled with anti-Hts-RC (yellow), anti-Hts (red), anti-LamC (red), and DAPI (blue). (H-I) *nos-Gal4>Tnpo-SR^{V22}* (H) and *nos-Gal4>cdm^{HMS02846}* (I) females labeled with anti-Orb (yellow; oocyte), anti-Hts and anti-LamC (red), and DAPI (blue). Values in bottom left corner of G-H represent the number of ovarioles containing germline tumors over the total number of ovarioles analyzed. Scale bar, 20 μm. (J) Number of germ cells per egg chamber at 6 days after eclosion. Each dot represents one egg chamber; numbers above dots represent the total number of egg chambers analyzed. Quantifications do not include tumorous egg chambers.

Figure 2. *Ran* and *Tnpo-SR* are required for proper oocyte selection.

(A-C) Driver control (A), *nos-Gal4>Tnpo-SR^{V22}* (B), and *nos-Gal4>Ran^{GL01341}* egg chambers labeled with anti-Orb (yellow), anti-Hts (red), anti-LamC (red), and DAPI (blue). (D-F) Driver control (D) and *nos-Gal4>Tnpo-SR^{V22}* (E-F) egg chambers labeled with anti-BicD (yellow), anti-Hts (red), anti-LamC (red), and DAPI (blue). Arrowheads indicate oocytes, recognized by their condensed nuclei. (G) Frequencies of egg chambers containing 0 (grey), 1 (white), or 2 (black) oocytes. The numbers in the bars represent the total number of egg chambers analyzed. (H-J) *Tnpo-SR^{KG04870}* mutant mosaic germarium at 8 days after clone induction labeled with anti-GFP (green;

wild-type cells), anti-Hts and anti-LamC (blue), and anti-Orb (red) (H-H') or anti-C(3)G (red). (J-

J'). White lines demarcate wild-type (solid) or mutant (dashed) cysts. (I) Percentage of GFP-

positive (wild-type) or GFP-negative (*Tnpo-SR^{KG04870}*) 16-cell cysts with an Orb-positive oocyte

8 days after clone induction. (K) Average number of C(3)G-positive germ cells in GFP-positive

(wild-type) or GFP-negative (*Tnpo-SR^{KG04870}*) 8- or 16-cell cysts at 8 days after clone induction.

Numbers in the bars represent the total number of cysts analyzed. Scale bar, 20 µm (A-F) and 5

µm (H-J). Error bars, mean ± SEM. ***$p < 0.001$; Student's two-tailed T-test.

Figure 3. Ran is dynamically localized in germ cells but dispensable for spindle assembly.

(A-F'') Ovarioles from *nos-Gal4>Ran^{HA}* (A-B''), *nos-Gal4>Ran^{T24N}* (C-D''), and *nos-Gal4>Ran^{Q69L}* (E-F'') labeled with anti-HA (green), anti-Tubulin (red) and DAPI (blue). Panels B'-B'', D'-D'', and F'-F'' show the DAPI and HA channels of panels B, D, and F, respectively. (G-J) Maximum intensity projections of driver control (G), RNAi-depleted *cdm^{HMS02846}* (H) and *Tnpo-SR^{V22}* (I), and *Ran^{T24N}* mutant (J) cysts labeled with anti-tubulin (red) anti-Hts (green), anti-LamC (green), and DAPI (blue). Scale bars, 10 μm.

Figure 4. Ran function in dividing germ cells requires Ran-GTP activity.

(A) Maximum intensity projection of a germarium from a *nos-Gal4>Ran^{T24N}* ovary immunostained with anti-Hts (red), anti-LamC (red), and DAPI (blue). (B) Maximum intensity projection of a *Ran^{T24N}* germarium stained with anti-Hts-RC (red), anti-Hts (green), anti-LamC (green), and DAPI (blue). (C-D) *Ran^{T24N}* (C, GDP-locked mutant) and *Ran^{Q69L}* (D, GTP-locked mutant) germaria labeled with anti-Orb (yellow), anti-Hts (red), anti-LamC (red), and DAPI (blue). Arrowheads indicate oocytes, as recognized by their condensed nuclei. Scale bars, 20 μm. (E) Number of germ cells per egg chamber at 6 days after eclosion. Each dot represents one egg chamber; numbers above dots represent the total number of egg chambers analyzed.

Figure 5. *Tnpo-SR* and *cdm* are required for fusome integrity and maturation.

(A-D) Driver control (A), *cdm^HMS02846^* (B), *Tnpo-SR^V22^* (C), and *Ran^T24N^* (D) germaria labeled with anti-Hts (red), anti-LamC (red), and DAPI (blue). Arrowheads indicate points of the fusome that are thinner than normal. (E-F) Driver control (E) and *nos-Gal4>Tnpo-SR^V22^* (F) GSC/CB pairs labeled with anti-pTyrosine (red), anti-Hts (green), anti-LamC (green), and DAPI (blue). Arrowheads indicate pTyr+ plug that forms between the GSC/CB pair prior to abscission. (G-H) Control (G) and *nos-Gal4>Tnpo-SR^V22^* (H) GSC/CB pairs labeled with anti-EdU (magenta), anti-Hts (green) and anti-LamC (green). (I-J) *Scrib-GFP* control (I) and *Tnpo-SR^V22^;Scrib-GFP* (J) germaria labeled with anti-GFP (green), anti-Hts (red), and anti-LamC (red). (K-L) *Rtnl1-GFP* control (K) and *Tnpo-SR^V22^,Rtnl1-GFP* (L) germaria labeled with anti-GFP (green), anti-Hts (red), and anti-LamC (red). Arrowheads indicate highest level of expression of the Scrib (I-J) and Rtnl1 (K-L). All images are maximum intensity projections. Scale bars, 10 μm.

Figure 6. Depletion of *Tnpo-SR* and *cdm*, but not Ran-GTP, delays ring canal protein deposition and fails to suppress cytokinesis in dividing cysts.

(A) *Tnpo-SR^{KG04870}* mutant mosaic germarium labeled with anti-GFP (green; wild-type cells), anti-pTyrosine (pTyr, red), anti-Hts (blue), and anti-LamC (blue). (B) Maximum intensity projection of a *Tnpo-SR^{KG04870}* mutant mosaic germarium labeled with anti-GFP (green; wild-type cells) and anti-E-cadherin (red). Arrows indicate E-cadherin connections. (C-D) Control (C) and *nos-Gal4>Tnpo-SR^{V22}* (D) germaria labeled with Hts-RC (red), anti-Hts (green), anti-LamC (green),

and DAPI (blue). (E-H) Maximum intensity projections of driver control (E) and RNAi-depleted

cdm (F), *Tnpo-SR* (G), and *Ran* (H) cysts labeled with anti-pTyr (red), anti-Hts (green), and anti-

LamC (green). Arrowheads indicate ring canals; open arrowheads indicate a cyst lacking ring

canals. (I-L) Control (I), *nos-Gal4>Tnpo-SR^{V22}* (J) and *nos-Gal4>Ran^{T24N}* mutant (K-L)

germarium labeled with anti-anillin (green) anti-Hts (red), and anti-LamC (red). Arrowheads

indicate closed ring canals. Scale bars, 10 µm. (M) Summary of fusome and ring canal phenotypes

in *Tnpo-SR-* and *cdm*-depleted germ cell cysts.

Figure 7. CycA is ectopically expressed in c*dm* and *Tnpo-SR* cysts.

(A-D) Driver control (A), *Ran^{GL01341}* (B), *cdm^{HMS02846}* (C), and *Tnpo-SR^{V22}* (D) RNAi knock-down germaria labeled with anti-CycA (red), anti-Hts (green), and anti-LamC (green). (E) *nos-Gal4>FUCCI;Tnpo-SR^{V22}* labeled with anti-GFP (green; E2f1), anti-dsRed (red; CycB), anti-Hts (white), and anti-LamC (white). All images are maximum intensity projections. Scale bars, 10 μm. (F) Percentage of cells/cysts containing fusome-localized CycA. (G) Percentage of cells/cysts containing *RFP::CycB* in *nos-Gal4* controls and *Tnpo-SR^{V22}* knock-down germaria. The number of cells/cysts analyzed is shown above bars. ***$p < 0.001$, Student's Chi-squared test.

SUPPLEMENTAL DATA

Supp. Figure 1. Ran and β-importins are necessary for GSC maintenance.

(A-B) *nos-Gal4* control (A) and *nos-Gal4>Ran^{GL01341}* (B) germaria labeled with anti-Orb (yellow),

anti-Hts (red), anti-LamC (red) and DAPI (blue). Solid white lines demarcate GSCs present in the

germarium, asterisks indicate cap cells. (C) Average number of GSCs per germarium in *nos-Gal4*

control or *nos-Gal4>UAS-Ran/importin RNAi* ovaries at 3, 6, and 10 days after eclosion (dae). At

least 30 germaria were analyzed at each timepoint. (D-E) *FRT40A* control (D) and *Tnpo-SR^{KG04870}*

FRT40A mutant (E) mosaic germaria at 8 days after clone induction labeled with anti-GFP (green),

anti-Hts (red), anti-LamC (red) and DAPI (blue). Lines demarcate wild-type (solid) or mutant

(dashed) GSCs (white) and cystoblast/cysts (yellow). (F) Percentage of total germaria analyzed

with a GFP-negative GSC at 4, 8, and 12 days after clone induction. At least 145 germaria were

analyzed for each genotype at each timepoint.

Supp. Figure 2. *Tnpo-SR* **germline tumors are primarily composed of mitotically-dividing undifferentiated germ cells.**

(A) *nos-Gal4>Tnpo-SR^{V22}* egg chamber labeled with anti-Hts-RC (red), anti-Hts (green), anti-LamC (green), and DAPI (blue). (B) *nos-Gal4>Tnpo-SR^{V22}* egg chamber labeled with anti-pTyr (red), anti-Hts (green), anti-LamC (green), and DAPI (blue). (C-C') *nos-Gal4>Tnpo-SR^{V22}* egg chamber labeled with anti-pHH3 (green), anti-Hts (red), anti-LamC (red), and DAPI (blue). Panel C' depicts pHH3 channel only. Arrowheads indicate pHH3-positive cells within the germline tumor. (D-E) Control (D-D') and *nos-Gal4>Tnpo-SR^{V22}* (E-E') germaria labeled with anti-Sxl (red), anti-Hts (green), anti-LamC (green), and DAPI (blue). Panels D' and E' show heat maps corresponding to panels D and E, where cool colors are low expression and warm colors are high expression. Scale bars, 20 μm.

Supp. Figure 3. *Tnpo-SR* mutants fail to maintain cyst connectivity and some cells are ectopically mitotically active.

(A) *Tnpo-SR^{KG04870}* mutant egg chamber labeled with anti-GFP (green), anti-Hts (red), anti-LamC (red), and DAPI (blue). White lines indicate wild-type cysts (solid) and mutant cysts (dashed). (B) *Tnpo-SR^{KG04870}* mutant egg chamber labeled with anti-GFP (green), anti-pHH3 (white), anti-Hts (red), anti-LamC (red), and DAPI (blue). (C-E''') *Tnpo-SR^{LL05552}* (C and E-E''') and *Tnpo-SR^{KG04870}* (D) mutant germaria labeled with anti-GFP (green), anti-Hts (red), anti-LamC (red), and DAPI (blue). Panels E-E''' are adjacent 1 µm optical sections from the same germarium. Arrowhead indicates thin thread of fusome material; open arrowheads follow the GFP+ somatic cell projections invaginating the mutant cysts. (D) is a 2 µm subset maximum intensity projection. Scale bars, 10 µm.

Supp. Figure 4. An Anillin-NLS mutant is not efficient in disrupting Anillin localization to nuclei and ring canals.

(A-D) Representative controls of *UAS-Scra-GFP* (A and C) and NLS mutated *UAS-Scra-3A-GFP* mutant (B and D) germaria labeled with anti-GFP (green), anti-Hts (red), anti-LamC (red), and DAPI (blue) (A-B) or anti-pTyrosine (red) (C-D). Arrowheads indicate closed ring canals. Scale bar, 10 μm.

988   **Supp. Table S1.** Fly stocks and immunofluorescence reagents used in this study.

Reagent	Source	Reference
nos-GAL4::VP16-nos.UTR	D. Drummond-Barbosa	(Rørth, 1998; Van Doren et al., 1998)
w*; nosGal4::VP16, UASp-tubGFP	BLM #7253	(Grieder et al., 2000)
y¹sc*v¹; P{y[+t7.7] v[+t1.8]=TnpoSR-V22}attP2	Ables Lab	this study
y¹v¹; P{y[+t7.7] v[+t1.8]=TRiP.GL01483}attP2	BLM #43142	(Perkins et al., 2015)
y¹sc*v¹sev²¹; P{y[+t7.7] v[+t1.8]=TRiP.GL01341}attP40	BLM #42482	(Perkins et al., 2015)
y¹sc*v¹sev²¹; P{y[+t7.7] v[+t1.8]=TRiP.HMS02846}attP2	BLM #44551	(Perkins et al., 2015)
y¹v¹; P{y[+t7.7] v[+t1.8]=TRiP.HMS00020}attP2	BLM #33626	(Perkins et al., 2015)
y¹sc*v¹sev²¹; P{y[+t7.7] v[+t1.8]=TRiP.HMS01408}attP2	BLM #34998	(Perkins et al., 2015)
y¹sc*v¹sev²¹; P{y[+t7.7] v[+t1.8]=TRiP.HMS00991}attP2	BLM #34021	(Perkins et al., 2015)
y¹v¹; P{y[+t7.7] v[+t1.8]=TRiP.GL01273}attP2	BLM #41845	(Perkins et al., 2015)
y¹sc*v1; P{y[+t7.7] v[+t1.8]=TRiP.HMS02968}attP2	BLM #50732	(Perkins et al., 2015)
y¹sc*v¹sev²¹; P{y[+t7.7] v[+t1.8]=TRiP.GL00435}attP40	BLM #35598	(Perkins et al., 2015)
y¹sc*v¹sev²¹; P{y[+t7.7] v[+t1.8]=TRiP.HMC03738}attP40	BLM #55142	(Perkins et al., 2015)
y¹v¹; P{y[+t7.7] v[+t1.8]=TRiP.HMJ22418}attP40/CyO	BLM #58311	(Perkins et al., 2015)

$y^1 sc^* v^1 sev^{21}$; $P\{y[+t7.7]$ $v[+t1.8]=TRiP.HMC04414\}attP40$	BLM #56974	(Perkins et al., 2015)
$y^1 v^1$; $P\{y[+t7.7]$ $v[+t1.8]=TRiP.HMJ23009\}attP40$	BLM #61230	(Perkins et al., 2015)
$hsFLP$; $FRT40A$, $UbiGFP$ / CyO	D. Drummond-Barbosa	(Ables et al., 2016)
$FRT40A$ / CyO	D. Drummond-Barbosa	(Ables et al., 2016)
$y^{d2} w^{1118}$ $P\{ry[+t7.2]=ey-FLP.N\}2$ $P\{ry[+t7.2]=GMR-lacZ.C(38.1)\}TPN1$; $P\{y[+mDint2]$ $w[BR.E.BR]=SUPor-P\}Trn-SR[KG04870]$ $P\{ry[+t7.2]=neoFRT\}40A$ / CyO, $y[+]$	Kyoto #111581	(Bellen et al., 2004; Chen et al., 2005)
$y^* w^*$; $PBac\{SAstopDsRed\}LL05552$ $P\{ry[+t7.2]=neoFRT\}40A$ $P\{w[+mW.hs]=FRT(w[hs])\}G13$ $cn^1 bw^1$ / CyO, $S^* bw^1$	Kyoto #141628	(Schuldiner et al., 2008)
$w;P\{UASp:Ran^{HA}\}1.19/SM6$	K. McKim	(Cesario and McKim, 2011)
$w;P\{UASp:Ran^{HA}\}10-A/TM3$	K. McKim	(Cesario and McKim, 2011)
$w;P\{UASp:Ran^{T24NHA}\}18.3/TM3$	K. McKim	(Cesario and McKim, 2011)
$w;P\{UASp:Ran^{T24NHA}\}25.1/TM3$	K. McKim	(Cesario and McKim, 2011)
$w;P\{UASp:Ran^{Q69LHA}\}7.1/TM3$	K. McKim	(Cesario and McKim, 2011)
$w;P\{UASp:ran^{Q69LHA}\}61.2/TM3$	K. McKim	(Cesario and McKim, 2011)
$w[1118]$; $P\{UASp-GFP.E2f.1-230\}26$ $P\{UASp-mRFP1.NLS.CycB.1-266\}4/CyO,P\{en1\}wg[en11];MKRS/TM6B,$ $Tb1$	BLM #55110	(Zielke et al., 2014)
$Rtnl1-GFP$	B. Riggs, K. Röper	(Röper, 2007)

scrib^{CA07683}	L. Cooley, M. Buszczak	(Buszczak et al., 2007)
w; P{w[+mC]=UASp-GFP-scra}3*	BLM #51348	(Silverman-Gavrila et al., 2008)
w; wg*^{Sp-1}*/CyO; P{w[+mC]=UASp-GFP-3A-scra}3*	BLM #51347	(Silverman-Gavrila et al., 2008)

989