CSF metabolites associate with CSF tau and improve prediction of
Alzheimer's disease status

Ruocheng Dong[1], Burcu F. Darst[2], Yuetiva Deming[1], Yue Ma[6], Qiongshi Lu[3],

Henrik Zetterberg[7-10], Kaj Blennow[7,8], Cynthia M. Carlsson[4-6], Sterling C.

Johnson[4-6], Sanjay Asthana[5,6], Corinne D. Engelman[1,4,6]*

[1]Department of Population Health Sciences, University of Wisconsin School of

Medicine and Public Health, Madison, WI, 53726, USA

[2]Center for Genetic Epidemiology, Keck School of Medicine, University of

Southern California, Los Angeles, CA, 90033, USA

[3]Department of Biostatistics and Medical Informatics, University of Wisconsin,

Madison, WI, 53792, USA

[4]Wisconsin Alzheimer's Institute, University of Wisconsin School of Medicine

and Public Health, Madison, WI, 53719, USA

[5]Geriatric Research Education and Clinical Center, Wm. S. Middleton Memorial

VA Hospital, Madison, WI, 53705, USA

[6]Alzheimer's Disease Research Center, University of Wisconsin School of

Medicine and Public Health, Madison, WI, 53792, USA

[7]Institute of Neuroscience and Physiology, The Sahlgrenska Academy at

University of Gothenburg, Mölndal, 41390, Sweden

[8]Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital,

Mölndal, 41345, Sweden

[9]UK Dementia Research Institute at UCL, London, WC1E6BT, UK

[10]Department of Neurodegenerative Disease, UCL Institute of Neurology,

London, WC1H0AL, UK

* Correspondence:

Corinne D. Engelman

610 Walnut Street, 1007A WARF Madison, WI, 53726, USA

Phone: 608-265-5491, email: cengelman@wisc.edu

Abstract

INTRODUCTION: Cerebrospinal fluid (CSF) total tau (t-tau) and

phosphorylated tau (p-tau) are biomarkers of Alzheimer's disease

(AD), yet much is unknown about AD-associated changes in tau

metabolism and tau tangle etiology.

METHODS: We assessed the variation of t-tau and p-tau

explained by 38 previously identified CSF metabolites using linear

regression models in middle-age controls from the Wisconsin

Alzheimer's Disease Research Center, and predicted AD/mild

cognitive impairment (MCI) vs. an independent set of older

controls using metabolites selected by the least absolute shrinkage

and selection operator (LASSO).

RESULTS: The 38 CSF metabolites explained 70.3% and 75.7%

of the variance in t-tau and p-tau. Of these, 7 LASSO-selected

metabolites improved the prediction ability of AD/MCI vs. older

controls (AUC score increased from 0.92 to 0.97 and 0.78 to 0.93)

compared to the base model.

DISCUSSION: These tau-correlated CSF metabolites increase

AD/MCI prediction accuracy and may provide insight into tau

tangle etiology.

Keywords: Alzheimer's disease, CSF, metabolite,

metabolomics, t-tau, p-tau

1. Introduction

One of the defining neuropathological changes in Alzheimer's disease (AD) is the intraneuronal aggregates of hyperphosphorylated and misfolded tau that give rise to neurofibrillary tangles and neuropil threads [1]. Their corresponding biomarkers in cerebrospinal fluid (CSF), total tau (t-tau) and phosphorylated tau (p-tau), can predict clinical AD and its progression [2]. Moreover, a new plasma p-tau biomarker (p-tau181) has recently been associated with AD pathology [3]. Research has been done to understand tau changes and how they happen [4,5]. For example, it has been shown that the dysregulation of kinases and phosphatases results in three to four times greater quantities of phosphorylated tau in the brains of AD patients than in normal adult brains [2], but the pathologic processes remain largely unknown.

Recent advancements in metabolomics technologies allow researchers to study multiple small molecules (<1500 Da), such as amino acids, fatty acids, and carbohydrates, simultaneously within a biological system [6]. Metabolites can be influenced by biological changes resulting from upstream molecular processes such as genetic mutations, as well as exogenous changes caused by environmental exposures (*e.g.*, diet, medications, and physical activity). Moreover, compared to RNA transcripts and proteins,

metabolites are more relevant to the current physiological state of a

cell, and their abnormal levels and relative ratios can reflect

disease progression, thus, metabolites serve as appropriate targets

for health outcomes research [7].

To date, there have been numerous targeted or untargeted

human blood metabolomic studies that focus on AD clinical status

or CSF biomarkers [8]. For example, Toledo *et al.* [9] have

conducted a network analysis using serum metabolites in

participants from the Alzheimer's Disease Neuroimaging Initiative

and found that accumulation of acylcarnitine species indicates

malfunction and alterations in tau metabolism. However, few

studies have been conducted to assess the association between CSF

metabolites and CSF tau. CSF communicates freely with the

interstitial fluid that bathes the neurons and other cell types of the

brain, spinal cord and the cranial and spinal nerves [10], which

makes it an ideal source to study the pathological changes

occurring in AD brains. By linking two well-established AD CSF

biomarkers, CSF t-tau and p-tau, which reflect tau secretion and

phosphorylation, and predict neurodegeneration and cortical tangle

formation, respectively [11], with CSF metabolites, additional

mechanistic information behind the development of pathological

alterations related to tau may be revealed. The findings from

studying CSF metabolites could ultimately be translated into

potential AD prevention through modifiable risk factors (*e.g.*,

dietary interventions), better prognostic indicators, or new drug

targets.

Darst *et al*. [12] constructed an inter-omics network

consisting of whole blood gene expression, plasma metabolites,

CSF metabolites, and AD risk factors in 1,111 non-Hispanic white

participants from the Wisconsin Registry for Alzheimer's

Prevention (WRAP). Within this inter-omics network, a cluster of

38 CSF metabolites was identified in the subset of 141 individuals

in which CSF was collected, with each individual metabolite being

significantly correlated (p threshold: $\leq 6.1 \times 10^{-10}$) with CSF t-tau

and p-tau, and these collective metabolites accounting for 60.7%

and 64.0% of the variation of t-tau and p-tau, respectively. In this

study, we aimed to (1) replicate these findings and evaluate the

predictive ability of these CSF metabolites in an independent

sample (the IMPACT cohort) from the Wisconsin Alzheimer's

Disease Research Center (Wisconsin ADRC); (2) examine the

predictive performance of the same metabolites present in plasma

in WRAP; (3) identify the major metabolites driving this cluster in

the IMPACT and WRAP cohorts and, in an independent sample,

evaluate whether they can be used as potential biomarkers to

enhance the prediction of AD or mild cognitive impairment (MCI);

and (5) understand the biological functions of all 38 metabolites

using pathway analyses to provide insight into disease-related

processes. Our results confirm the previous associations between

38 CSF metabolites and CSF tau and provide potential biological

mechanisms for the development of tau tangles and possible

candidates for CSF metabolite biomarkers or drug targets.

2. Methods

2.1 Participants

The Wisconsin ADRC's clinical core cohort started in 2009

and has well-characterized AD and MCI participants, as well as

healthy older controls (HOC), and the IMPACT cohort of initially

cognitively-unimpaired, asymptomatic middle-aged adults [13–

15]. The replication sample for the main analysis included 158

non-Hispanic white individuals from the IMPACT cohort with

cross-sectional CSF samples.

WRAP began recruitment in 2001 as a prospective cohort

study of initially cognitively-unimpaired, asymptomatic, middle-

aged adults enriched for a parental history of clinical AD [16]. The

WRAP cohort included 130 and 123 non-Hispanic white

individuals with longitudinal CSF and plasma samples,

respectively. Both the CSF and plasma cohorts included five

sibling pairs, one sibling trio and three sibling quartets. The WRAP

dataset was utilized to reproduce and refine the results from Darst

et.al. [12] using similar statistical models as those for the IMPACT

cohort.

This study was conducted with the approval of the

University of Wisconsin Institutional Review Board, and all

participants provided signed informed consent before participation.

2.2 CSF and plasma sample collection and CSF biomarkers

quantification

Fasting CSF samples for the Wisconsin ADRC cohorts and

WRAP were collected via lumbar puncture [13] following the

same protocol and by the same group of well-trained individuals

[13]. Samples were sent together in two batches to the lab of Drs.

Blennow and Zetterberg in Sweden, where commercially available

enzyme-linked immunosorbent assay (ELISA) methods were used

to quantify CSF t-tau, p-tau, and amyloid-beta 1-42 ($A\beta_{42}$)

(INNOTEST® assays HTAU AG, PHOSPHO-TAU[181P], and β-

amyloid1-42, respectively; Fujirebio, Ghent, Belgium) [13]. The

batch-adjusted predicted values for CSF biomarkers were used for

all analyses [17].

In WRAP, fasting blood samples were collected in

ethylenediaminetetraacetic acid (EDTA) tubes; the plasma was

pipetted off within one hour of collection and stored at -80°C [12].

A total of 141 longitudinal samples from 123 individuals in WRAP

with plasma metabolites were available for the main analysis. In

the Wisconsin ADRC, blood samples were collected in heparin

tubes, which could influence metabolite values; as such, plasma

metabolomics data have not been generated in Wisconsin ADRC

blood samples. Further details of how plasma and CSF samples

were processed are explained in an earlier study [12].

2.3 CSF metabolomic profiling and quality control

CSF and plasma metabolomic analyses and quantification

were performed in one batch by Metabolon (Durham, NC) using

an untargeted approach, based on Ultrahigh Performance Liquid

Chromatography-Tandem Mass Spectrometry platform (UPLC-

MS/MS) [18]. Details of the metabolomic profiling were described

in an earlier study [12].

Each metabolite value was first scaled so the median was

equal to one across all samples. Missing values were then imputed

to half the lowest level of detection for each biological metabolite

and 0.0001 (the lowest value that could be accepted in the analytic

software) for each xenobiotic metabolite. The missing percentage

for each of the 38 previously identified CSF metabolites prior to

imputation is shown in Supplemental Table 1. Metabolites with

zero variability between individuals, or with an interquartile range

of zero, were excluded (none of the 38 CSF metabolites were

excluded). Log10 transformation was employed to normalize the

data. After quality control, the previously identified 38 metabolites

were selected for this investigation. The distribution of each of the

38 CSF metabolites after imputation and Log 10 transformation is

shown in Supplemental Figure 1.

2.4 Statistical analysis

2.4.1 Prediction performance of the 38 CSF metabolites

To replicate the previously reported results in WRAP [12],

each metabolite's association with t-tau and p-tau was tested in the

IMPACT cohort and the Bonferroni adjustment was applied to

correct for multiple testing. A meta-analysis was also conducted by

using results from IMPACT and WRAP. To replicate the

performance of the cluster of 38 CSF metabolites in explaining

variation in tau pathology, we used linear regression models to

determine the prediction performance (r^2) of CSF t-tau and p-tau in

IMPACT. The base models, which included age, sex, and years of

education, were compared to models that also included the 38 CSF

metabolites. To reproduce the results in WRAP and compare them

to IMPACT using consistent statistical models, we determined the

prediction performance (r^2) of the 38 CSF metabolites using linear

mixed-effects regression with random intercepts to account for

repeated measures and sibling relationships. In both IMPACT and

WRAP, we randomly split the data into a training (70%) and

validation (30%) set and created plots to compare the observed and

predicted values. Finally, we physically combined the WRAP

baseline samples and IMPACT samples and re-conducted the

analysis to evaluate the explained variance. Sex-stratified

prediction differences were assessed in WRAP by fitting the

mentioned models in males and females separately. The number of

male samples in IMPACT was too small to perform sex-stratified

analyses while meeting the degrees of freedom needed by the

model. Of the original 38 CSF metabolites, 34 were also found in

plasma samples from WRAP and were tested together as predictors

for t-tau and p-tau using linear mixed-effects regression models, as

described above. A sensitivity analysis using only the baseline

samples was also conducted in WRAP for both CSF and plasma

metabolites. The statistical analyses here and below were all

conducted in R version 3.6.2. The lme4 package was used.

2.4.2 LASSO selection of important metabolites and their

prediction of AD/MCI vs. HOC

In order to incorporate a practical number of metabolites in

the prediction model of AD/MCI diagnosis vs. HOC instead of

including all 38 metabolites, the least absolute shrinkage and

selection operator (LASSO) [19] was applied to select the most

important metabolites (those with non-zero estimated effects) for

CSF t-tau and p-tau in both IMPACT and WRAP. In WRAP, the

average of longitudinal CSF measures was used in LASSO

regression. The tau variances explained by the selected metabolites

were re-evaluated using similar model from 2.4.1 in both IMPACT

and WRAP. The ability to enhance the prediction of AD/MCI vs.

HOC status by the metabolites selected from LASSO was

evaluated in an independent set of participants from the Wisconsin

ADRC using logistic regression and an area under the curve

(AUC) score. To determine prediction ability of the selected

metabolites beyond demographic factors and established

biomarkers, base models including age, sex, years of education,

APOE ε4 count, t-tau, p-tau, and Aβ$_{42}$ were compared to the base

model replacing t-tau and p-tau with the selected metabolites and

also the base model plus the selected metabolites from LASSO.

The analysis here used the "glmnet" package in R.

2.4.3 Biological relevance of the 38 CSF metabolites

An exploratory factor analysis was conducted to determine

if subsets of metabolites clustered together in latent factors

associated with t-tau and p-tau. The factor analysis was performed

in IMPACT and WRAP using the "psych" package in R.

Metabolites with a loading of ≥0.4 [20] in one particular factor and

lower loadings for the rest of the factors were considered as

members of that particular factor. Potential functional pathways of

the 38 metabolites were identified from the Homo sapiens KEGG

pathway by conducting pathway analyses using the web-based

software, Metabo-analyst [21], inputting the metabolites' human

metabolome database (HMDB) IDs, and using the default

hypergeometric test and the relative-betweenness centrality, which

is a measure of centrality in a graph based on the shortest paths

that pass through the vertex. Pathways were considered as

important if the FDR was ≤0.05 or the impact was ≥0.1.

3. Results

3.1 Participant characteristics

Characteristics of the participants can be found in Table 1.

Among 158 Wisconsin ADRC IMPACT participants and 130

WRAP participants who had CSF metabolite data available,

females comprised 74.7% of IMPACT participants and 65.4% of

participants in WRAP. The mean baseline age was significantly

younger in IMPACT (57.8 years) compared to WRAP (61.5 years).

The mean years of education was similar (16.0 and 16.1 years in

IMPACT and WRAP, respectively). Mean CSF t-tau was

significantly lower in IMPACT (283.1) compared to WRAP

(311.5). The correlation between t-tau and p-tau was

approximately 0.90 in IMPACT and WRAP. The characteristics of

each additional sub cohort of the Wisconsin ADRC and of the 123

WRAP participants in the plasma prediction analysis can also be

found in Table 1.

3.2 Prediction performance

Each of the 38 CSF metabolites was significantly

associated with t-tau and p-tau in IMPACT and the direction of the

effect was the same as in WRAP (Supplemental Table 2). Meta-

analysis results are shown in Supplemental Table 3. All

metabolites were significantly associated with t-tau and p-tau

except erythritol. Base models only explained approximately 10%

of the variance in t-tau and p-tau in both IMPACT and WRAP

(Table 2). In IMPACT, the statistical model including the 38 CSF

metabolites and demographics together explained 70.3% of the

variance in t-tau and 75.7% of the variance in p-tau values. These

results were similar to those calculated in WRAP, where the model

including the 38 CSF metabolites and demographics explained

62.4% and 65.1% of the variance in t-tau and p-tau values,

respectively. Similarly, in the combined dataset, the 38 CSF

metabolites explained 66.1% and 72.3% of the variance in t-tau

and p-tau, respectively. The results of the same analysis but only

using baseline samples in WRAP are shown in Supplemental Table

4. Supplemental Figure 2 shows plots comparing the observed and

predicted values for t-tau and p-tau in both IMPACT and WRAP.

In WRAP, these metabolites explained more of the variance in the

t-tau and p-tau in males (r^2=0.749 and 0.804) than in females

(r^2=0.591 and 0.640; Table 2). We did not have enough male

participants to fit the sex-stratified model in IMPACT; however,

while the female only r^2 was lower than the overall r^2 in WRAP,

this trend was not seen in IMPACT. In WRAP, the 34 of 38

metabolites present in plasma explained 26.9% and 30.1% of the

variance in CSF t-tau and p-tau, respectively (Table 2), which is

relatively low compared to CSF metabolites. We also examined the

same 34 CSF metabolites' prediction ability and confirmed that the

lower r^2 values for the 34 plasma metabolites were not due to the

absence of the four metabolites (Supplemental Table 4).

3.3 LASSO results

LASSO results for t-tau and p-tau in both IMPACT and

WRAP are shown in Table 3. Eight metabolites with non-zero

coefficients (ranging from 33.25 to 202.10) were chosen in

IMPACT, and twelve metabolites (coefficients ranging from -

112.48 to 333.57) were selected for t-tau in WRAP. Among the

selected metabolites, five were consistent across IMPACT and

WRAP (N-acetylneuraminate, C-glycosyl tryptophan, X-10457, X-

24228, and 1-palmitoyl-GPC(16:0)). Eleven metabolites in

IMPACT and twelve metabolites in WRAP with non-zero

coefficients (ranging from 1.07 to 28.80 in IMPACT and -4.06 to

30.80 in WRAP) were selected for p-tau, with seven metabolites

overlapping (N-acetylneuraminate, C-glycosyl tryptophan, X-

10457, X-24228, 1-oleoyl-GPC(18:1), 1-palmitoyl-GPC(16:0), and

1-myristoyl-2-palmitoyl-GPC(14:0/16:0)), which included the five

metabolites overlapping in the two t-tau models. These seven

metabolites along with demographics explained about 59% and

69% of the variance in t-tau and p-tau, respectively in IMPACT

and 59% and 62%, respectively in WRAP (Table 2).

When predicting AD vs. HOC and MCI vs. HOC, the base

models, including age, sex, years of education, *APOE* ε4 count, t-

tau, p-tau, and Aβ$_{42}$, achieved AUC scores of 0.92 and 0.78,

respectively. Replacing t-tau and p-tau with the seven metabolites

selected by LASSO, that overlapped across IMPACT and WRAP

for t-tau and/or p-tau, achieved AUC scores of 0.94 and 0.82. The

base model plus the seven metabolites collectively improved the

prediction ability of AD vs. HOC (AUC score increased from 0.92

to 0.97) and of MCI vs. HOC (AUC score increased from 0.78 to

0.93; Figure 1). The comparisons of results from the base model

plus seven LASSO selected metabolites to the base model with

seven randomly selected metabolites from 38 metabolites and

seven randomly selected metabolites from all CSF metabolites

with tau outcomes are shown in Supplemental Figure 3.

3.4 Biological relevance of the 38 metabolites

The biochemical names, sub-pathways, and super pathways

of the 38 metabolites can be found in Supplemental Table 5, which

also shows the loadings of each metabolite for three latent factors

produced through exploratory factor analysis. These factors

included the exact same metabolites and similar loadings for each

in both IMPACT and WRAP and explained about 60% of the

variance in the 38 metabolites. Factor 1 included 25 metabolites in

the following pathways: amino acids, nucleotides, carbohydrates,

cofactors and vitamins, energy, xenobiotics, and unknowns (no

confirmed biochemical names). Factor 2 was composed of eleven

lipids. Two lysophospholipids contributed to factor 3 and they

were selected by LASSO for p-tau in both IMPACT and WRAP.

Among the 29 known metabolites, 26 had HMDB IDs and

23 of these were present in the MetaboAnalyst database. In

pathway analyses, these 23 metabolites were enriched in two

KEGG pathways (Figure 2 and Table 4): (1) pentose and

glucuronate interconversions and (2) glycerophospholipid (GP)

metabolism. Three metabolites from Factor 1,

arabinose, arabitol/xylitol, and gulonate, were enriched in pentose

and glucuronate interconversions. Two metabolites, 1-palmitoyl-2-

palmitoleoyl-GPC(16:0/16:1) and 1-oleoyl-GPC(18:1) from

Factors 2 and 3, respectively, were enriched in

glycerophospholipid metabolism.

4. Discussion

Using a cross-sectional sample from the Wisconsin ADRC

IMPACT cohort, we replicated previous findings of 38 CSF

metabolites associated with t-tau and p-tau in WRAP [12]. Not

only was each of the 38 CSF metabolites significantly associated

with both tau outcomes after Bonferroni correction, but the high

amount of variance in tau explained by this cluster of 38 CSF

metabolites was confirmed in IMPACT.

Among these metabolites, there are 13 lipids, 7 amino

acids, 5 carbohydrates, 1 nucleotide, 1 energy metabolite, 1

cofactor and vitamin metabolite, 1 xenobiotic, and 9 unknown

metabolites. Some of these metabolites, such as 1,2-dipalmitoyl-

GPC(16:0/16:0) and stearoyl sphingomyelin(d18:1/18:0), were

previously reported to be associated with AD diagnosis or AD

pathogenesis [22,23]. Orešič et al. (2011) found that serum 1,2-

dipalmitoyl-GPC(16:0/16:0), also called PC(16:0/16:0), was one of

3 metabolites considered to be predictive markers of AD

progression in individuals with MCI [22]. CSF stearoyl

sphingomyelin(d18:1/18:0) , also called SM(d18:1/18:0),

distinguished clinical AD from controls, with an accuracy of 70%

and was significantly increased in patients displaying pathological

levels of $A\beta_{42}$, t-tau and p-tau [23], supporting that this molecule

changes in patients with A/T/N pathology. Additionally, the N-

acetylamino acids, N-acetylvaline, N-acetylthreonine, N-

acetylserine, and N-acetyl-isoputreanine, were identified in our

study. N-acetylthreonine and N-acetylserine are the downstream

metabolites of the cleavage process initiated by lysosomal protease

tripeptidyl peptidase 1 (TPP1) [24], and previous studies [25,26]

suggested that increased levels of TPP1 enhance fibrillar β-

amyloid degradation. In support of this, a secondary analysis in our

study found that N-acetylserine was significantly associated with

$A\beta_{42}$ (beta=480.38, p=0.002), providing evidence that this CSF

metabolite may be involved in brain amyloid pathology.

From the 38 metabolites, 7 were selected by LASSO in

both IMPACT and WRAP: N-acetylneuraminate, C-glycosyl

tryptophan, 1-palmitoyl-GPC(16:0), 1-oleoyl-GPC(18:1), 1-

myristoyl-2-palmitoyl-GPC(14:0/16:0), and two unknown

metabolites (X-10457 and X-24228). These improved the

prediction of AD vs. HOC by approximately 5% and MCI vs

HOC by 15% compared to a model that included the well-

established AD risk factors of age, sex, years of education, *APOE*

ε4 count, t-tau, p-tau, and $A\beta_{42}$. A recent study in a Japanese

cohort found that CSF N-acetylneuraminate was significantly

higher in AD patients, when compared to the idiopathic normal

pressure hydrocephalus, and had a positive correlation with CSF p-

tau (r=0.55) [27]. In our study, CSF N-acetylneuraminate was

positively associated with both t-tau and p-tau. C-glycosyl

tryptophan, a sugar-loaded amino acid, has been reported to be

strongly associated with aging, defined by chronological age

(beta=2.47, p=1.3×10^{-23}), in a human blood metabolome-wide

association study [28]. In our study, CSF C-glycosyl tryptophan

was positively associated with t-tau and p-tau. Two lipids 1-

palmitoyl-GPC(16:0) (also called LysoPC(16:0/0:0)) and 1-

myristoyl-2-palmitoyl-GPC(14:0/16:0) (also called PC(14:0/16:0)),

belong to the class of lysophospholipid (LysoPCs) and

phosphatidylcholines (PCs), respectively. Previous studies have

shown that numerous plasma/serum metabolites from the LysoPC

and PC classes were significantly associated with MCI and AD

dementia or able to discriminate MCI and AD dementia cases from

controls [9,29–34]. In a randomized crossover trial that treated

mild to moderate AD patients with medium-chain triglycerides, 1-

palmitoyl-GPC(16:0) levels increased along with an improvement

in cognition [34]. These seven LASSO-selected metabolites

improved the prediction of AD and MCI status, suggesting they

may be useful biomarkers for clinical AD and MCI diagnosis.

Another interesting discovery from the LASSO results is

that, while five metabolites were overlapping in IMPACT and

WRAP for both t-tau and p-tau, two metabolites,1-oleoyl-

GPC(18:1) (also called LysoPC(18:1(9Z)/0:0)) and 1-myristoyl-2-

palmitoyl-GPC(14:0/16:0) (also called PC(14:0/16:0)) were

selected only for p-tau, not t-tau. Since p-tau is more specific to

AD-related tau pathology than t-tau, these metabolites might

provide insight into the pathological processes involved in tau

tangle formation in AD.

When using the seven metabolites to predict AD/MCI vs.

HOC, the AUC scores from both the base model and base

model+metabolites were higher for AD than for MCI. However, a

greater improvement in prediction accuracy for MCI vs. HOC

(15%) was achieved than AD vs. HOC (5%). One possible reason

could be that the 38 CSF metabolites were originally identified

from the WRAP cohort, whose participants were relatively young

and have not been diagnosed with AD yet. Another explanation

could be that the base model, which included demographics,

APOE ε4 count, and three core AD CSF biomarkers, already

achieved a very high accuracy for predicting AD vs. HOC and had

little room for improvement.

In WRAP, we were able to test the prediction of 34 of the

38 metabolites that were found in plasma. We found that these 34

metabolites collectively did not explain much variation in CSF

concentrations of t-tau and p-tau (r^2 between 0.286 and 0.303).

This was not due to the absence of the 4 metabolites, since the r^2 of

the 34 metabolites in the CSF (0.621 to 0.641) was close to that

with all 38 metabolites (0.624 to 0.651) in WRAP. Moreover, the

correlations between the same 34 metabolites measured in both

CSF and plasma are relatively low (-0.13 to 0.30) [12]

(Supplemental Figure 4). We previously proposed that this low

correlation could be attributed to these metabolites not being able

to cross the blood brain barrier (BBB) [35]. For example,

cholesterol metabolism in the brain relies on its own cells to

produce cholesterol, and the transport of cholesterol from

peripheral circulation into the brain is prevented by the BBB

[36,37]. In this situation, the concentrations and functions of

metabolites like cholesterol are different across the BBB. Thus,

testing for these metabolites in a more readily available body fluid,

like blood, does not appear to be a viable option.

The factor analysis results suggest that the 38 metabolites

are associated with tau through three main clusters (1) the

combination of select amino acids, nucleotides, carbohydrates,

cofactors and vitamins, energy, xenobiotics, and unknown

metabolites; (2) phosphatidylcholines and sphingolipid

metabolism, and (3) lysophospholipids. Five metabolites from

these factors were enriched in (1) pentose and glucuronate

interconversions and (2) glycerophospholipid metabolism from the

pathway analysis. The pentose and glucuronate interconversion

pathway was suggested from genomics and metabolomics studies

to be involved in AD [38–40]. A urinary metabolomics study of

APP/PS1 transgenic mice of AD and a hippocampal metabolomics

study of CRND8 mice also identified this pathway [41,42]. Other

studies have shown that brain glucose dysregulation and pentose

related activities are associated with AD pathology [43–46]. Thus,

our results provide further potential links between molecules in

pentose and glucuronate metabolism, especially the three CSF

metabolites, arabinose, xylitol, and gulonate, and the tau

pathological process of AD.

The brain is the most cholesterol-rich organ, containing

glycerophospholipids, cholesterol, sphingolipids, etc. [47]. The

neural membranes are also composed of these lipids and the

evidence suggests that glycerophospholipids and

glycerophospholipid metabolism may associate with neural

membrane composition alterations, glycerolipid-derived lipid-

mediated oxidative stress, and neuroinflammation [9,48]. For

example, levels of glycerophospholipids were decreased in brain

autopsy samples from AD patients compared to age-matched

controls [49]. In another study, increased glycerophospholipid

levels were associated with increased activities of lipolytic

enzymes and elevated concentrations of phospholipid degradation

metabolites [50]. In our analysis, the two metabolites, 1-palmitoyl-

2-palmitoleoyl-GPC(16:0/16:1) and 1-oleoyl-GPC(18:1), from

Factors 2 and 3 were in a feedback loop and their levels were

influenced by the genes *LCAT*, *PLA2G4B*, and *LPCAT*

(Supplemental Figure 5.) Previous studies have suggested that

LCAT and *LPCAT* are related to AD [51,52]. Thus, by connecting

glycerophospholipids, especially these two metabolites with t-tau

and p-tau, we provide further evidence for their connections with

AD pathogenesis.

Our sample sizes were relatively small for both IMPACT

and WRAP; however, the 38 CSF metabolites' associations with

CSF t-tau and p-tau levels identified before [12] were replicated in

the independent IMPACT data, strengthening our confidence that

these 38 metabolites are important for tau pathology. However,

further research is necessary to understand whether a causal

relationship exists between these CSF metabolites and tau

pathology. One limitation of this study is that both IMPACT and

WRAP are predominantly non-Hispanic white/Caucasian, so the

findings of this study may not be generalizable to other

races/ethnicities. Another limitation is that most of the 38

metabolites are highly correlated with each other. LASSO selected

seven metabolites that have non-zero effects on tau, but the

resulting metabolites are still correlated with each other

(Supplemental Figure 6; range of 0.40 to 0.96). A more

sophisticated approach that can further remove non-independent

metabolites is needed for clinical application. A third limitation is

that in our pathway analysis, only three or two metabolites were

included in the enriched pathways (pentose and glucuronate

interconversions and glycerophospholipid metabolism,

respectively). Future research will be necessary to confirm these

results.

In summary, we aimed to replicate earlier findings of 38

CSF metabolites' correlation with tau and expand the biological

knowledge of them to better understand their roles in AD

pathogenesis. 38 CSF metabolites individually associated with two

tau outcomes significantly and, together, explained a large amount

of variance in tau. A subset of these metabolites, selected by

LASSO, improved the prediction accuracy of AD/MCI vs. HOC

over a model that included established predictors of AD. Two

promising metabolic pathways, pentose and glucuronate

interconversions metabolism and glycerophospholipid metabolism,

were identified in this study and have been shown to be related to

AD in previous literature. IMPACT and WRAP are ongoing

longitudinal studies that are continuing to collect plasma and CSF

from study participants, and additional data will be generated in

the future. These data may help fill in gaps regarding the

mechanisms linking metabolites and AD, improve the

establishment of CSF-based metabolite biomarkers, and identify

novel drug targets.

Acknowledgements

The authors especially thank the WRAP and Wisconsin ADRC

participants and staff for their contributions to the studies. Without

their efforts this research would not be possible. This study was

supported by the National Institutes of Health (NIH) grants

[R01AG27161 (Wisconsin Registry for Alzheimer Prevention:

Biomarkers of Preclinical AD), R01AG054047 (Genomic and

Metabolomic Data Integration in a Longitudinal Cohort at Risk for

Alzheimer's Disease), and P30AG062715 (Wisconsin Alzheimer's

Disease Research Center Grant)], the Helen Bader Foundation,

Northwestern Mutual Foundation, Extendicare Foundation, State

of Wisconsin, the Clinical and Translational Science Award

(CTSA) program through the NIH National Center for Advancing

Translational Sciences (NCATS) grant [UL1TR000427], and the

University of Wisconsin-Madison Office of the Vice Chancellor

for Research and Graduate Education with funding from the

Wisconsin Alumni Research Foundation. This research was

supported in part by the Intramural Research Program of the

National Institute on Aging. Computational resources were

supported by core grants to the Center for Demography and

Ecology [P2CHD047873] and the Center for Demography of

Health and Aging [P30AG017266]. Author Y Deming was

supported by a training grant from the National Institute on Aging

[T32AG000213]. HZ is a Wallenberg Scholar supported by grants

from the Swedish Research Council [#2018-02532], the European

Research Council [#681712], the Swedish state under the

agreement between the Swedish government and the County

Councils, the ALF-agreement [#ALFGBG-720931], the Alzheimer

Drug Discovery Foundation (ADDF), USA [#201809-2016862],

and the UK Dementia Research Institute at UCL.

KB is supported by the Swedish Research Council (#2017-

00915), ADDF, USA [#RDAPB-201809-2016615], the Swedish

Alzheimer Foundation [#AF-742881], Hjärnfonden, Sweden

[#FO2017-0243], the Swedish state under the agreement between

the Swedish government and the County Councils, the ALF-

agreement [#ALFGBG-715986], and European Union Joint

Program for Neurodegenerative Disorders [JPND2019-466-236].

Conflicts of interest

HZ has served at scientific advisory boards for Denali, Roche

Diagnostics, Wave, Samumed and CogRx, has given lectures in

symposia sponsored by Fujirebio, Alzecure and Biogen, and is a

co-founder of Brain Biomarker Solutions in Gothenburg AB

(BBS), which is a part of the GU Ventures Incubator Program. KB

has served as a consultant or at advisory boards for Abcam, Axon,

Biogen, Lilly, MagQu, Novartis and Roche Diagnostics, and is a

co-founder of Brain Biomarker Solutions in Gothenburg AB

(BBS), which is a part of the GU Ventures Incubator Program.

Reference

[1] Serrano-Pozo A, Frosch MP, Masliah E, Hyman BT. Neuropathological Alterations in Alzheimer Disease. Cold Spring Harb Perspect Med 2011;1. https://doi.org/10.1101/cshperspect.a006189.

[2] Zetterberg H a. Cerebrospinal fluid biomarkers for Alzheimer's disease: current limitations and recent developments. Current Opinion in Psychiatry 2015;28:402–9. https://doi.org/10.1097/YCO.0000000000000179.

[3] Karikari TK, Pascoal TA, Ashton NJ, Janelidze S, Benedet AL, Rodriguez JL, et al. Blood phosphorylated tau 181 as a biomarker for Alzheimer's disease: a diagnostic performance and prediction modelling study using data from four prospective cohorts. The Lancet Neurology 2020;19:422–33. https://doi.org/10.1016/S1474-4422(20)30071-5.

[4] Barghorn S, Mandelkow E. Toward a Unified Scheme for the Aggregation of Tau into Alzheimer Paired Helical Filaments. Biochemistry 2002;41:14885–96. https://doi.org/10.1021/bi026469j.

[5] Jouanne M, Rault S, Voisin-Chiret A-S. Tau protein aggregation in Alzheimer's disease: An attractive target for the development of novel therapeutic agents. European Journal of Medicinal Chemistry 2017;139:153–67. https://doi.org/10.1016/j.ejmech.2017.07.070.

[6] Hasin Y, Seldin M, Lusis A. Multi-omics approaches to disease. Genome Biology 2017;18:83. https://doi.org/10.1186/s13059-017-1215-1.

[7] Enche Ady CNA, Lim SM, Teh LK, Salleh MZ, Chin A-V, Tan MP, et al. Metabolomic-guided discovery of Alzheimer's disease biomarkers from body fluid: Metabolomic-based Alzheimer's Disease Biomarkers. J Neuro Res 2017;95:2005–24. https://doi.org/10.1002/jnr.24048.

[8] Wilkins JM, Trushina E. Application of Metabolomics in Alzheimer's Disease. Front Neurol 2018;8. https://doi.org/10.3389/fneur.2017.00719.

[9] Toledo JB, Arnold M, Kastenmüller G, Chang R, Baillie RA, Han X, et al. Metabolic network failures in Alzheimer's disease: A biochemical road map. Alzheimer's & Dementia 2017;13:965–84. https://doi.org/10.1016/j.jalz.2017.01.020.

[10] Fiandaca MS, Mapstone ME, Cheema AK, Federoff HJ. The critical need for defining preclinical biomarkers in Alzheimer's disease. Alzheimer's & Dementia 2014;10:S196–212. https://doi.org/10.1016/j.jalz.2014.04.015.

[11] Sato C, Barthélemy NR, Mawuenyega KG, Patterson BW, Gordon BA, Jockel-Balsarotti J, et al. Tau Kinetics in

Neurons and the Human Central Nervous System. Neuron
2018;98:861–4. https://doi.org/10.1016/j.neuron.2018.04.035.

[12] Darst BF, Lu Q, Johnson SC, Engelman CD. Integrated
analysis of genomics, longitudinal metabolomics, and
Alzheimer's risk factors among 1,111 cohort participants.
Genetic Epidemiology 2019;43:657–74.
https://doi.org/10.1002/gepi.22211.

[13] Darst BF, Koscik RL, Racine AM, Oh JM, Krause RA,
Carlsson CM, et al. Pathway-Specific Polygenic Risk Scores
as Predictors of Amyloid-β Deposition and Cognitive
Function in a Sample at Increased Risk for Alzheimer's
Disease. Journal of Alzheimer's Disease 2016;55:473–84.
https://doi.org/10.3233/JAD-160195.

[14] Racine AM, Koscik RL, Berman SE, Nicholas CR, Clark LR,
Okonkwo OC, et al. Biomarker clusters are differentially
associated with longitudinal cognitive decline in late midlife.
Brain 2016;139:2261–74.
https://doi.org/10.1093/brain/aww142.

[15] Vogt NM, Romano KA, Darst BF, Engelman CD, Johnson
SC, Carlsson CM, et al. The gut microbiota-derived
metabolite trimethylamine N-oxide is elevated in Alzheimer's
disease. Alzheimer's Research & Therapy 2018;10:124.
https://doi.org/10.1186/s13195-018-0451-2.

[16] Johnson SC, Koscik RL, Jonaitis EM, Clark LR, Mueller KD,
Berman SE, et al. The Wisconsin Registry for Alzheimer's
Prevention: A review of findings and current directions.
Alzheimer's & Dementia: Diagnosis, Assessment & Disease
Monitoring 2018;10:130–42.
https://doi.org/10.1016/j.dadm.2017.11.007.

[17] Ma Y, Hulle CAV, Norton DL, Lazar KK, Chappell RJ,
Clark LR, et al. P1-286: TRANSFORMATION OF CSF
BIOMARKER VALUES BETWEEN MEASUREMENT
BATCHES. Alzheimer's & Dementia 2018;14:P393–5.
https://doi.org/10.1016/j.jalz.2018.06.292.

[18] Bridgewater BR EA. High Resolution Mass Spectrometry
Improves Data Quantity and Quality as Compared to Unit
Mass Resolution Mass Spectrometry in High-Throughput
Profiling Metabolomics. Metabolomics 2014;04.
https://doi.org/10.4172/2153-0769.1000132.

[19] Tibshirani R. Regression Shrinkage and Selection Via the
Lasso. Journal of the Royal Statistical Society: Series B
(Methodological) 1996;58:267–88.
https://doi.org/10.1111/j.2517-6161.1996.tb02080.x.

[20] Archer N, Brown RG, Reeves SJ, Boothby H, Nicholas H,
Foy C, et al. Premorbid Personality and Behavioral and
Psychological Symptoms in Probable Alzheimer Disease. The

American Journal of Geriatric Psychiatry 2007;15:202–13. https://doi.org/10.1097/01.JGP.0000232510.77213.10.

[21] Chong J, Soufan O, Li C, Caraus I, Li S, Bourque G, et al. MetaboAnalyst 4.0: towards more transparent and integrative metabolomics analysis. Nucleic Acids Res 2018;46:W486–94. https://doi.org/10.1093/nar/gky310.

[22] Orešič M, Hyötyläinen T, Herukka S-K, Sysi-Aho M, Mattila I, Seppänan-Laakso T, et al. Metabolome in progression to Alzheimer's disease. Translational Psychiatry 2011;1:e57–e57. https://doi.org/10.1038/tp.2011.55.

[23] Koal T, Klavins K, Seppi D, Kemmler G, Humpel C. Sphingomyelin SM(d18:1/18:0) is Significantly Enhanced in Cerebrospinal Fluid Samples Dichotomized by Pathological Amyloid-β42, Tau, and Phospho-Tau-181 Levels. JAD 2015;44:1193–201. https://doi.org/10.3233/JAD-142319.

[24] Sindelar M, Dyke JP, Deeb RS, Sondhi D, Kaminsky SM, Kosofsky BE, et al. Untargeted Metabolite Profiling of Cerebrospinal Fluid Uncovers Biomarkers for Severity of Late Infantile Neuronal Ceroid Lipofuscinosis (CLN2 , Batten Disease). Sci Rep 2018;8:1–12. https://doi.org/10.1038/s41598-018-33449-0.

[25] Solé-Domènech S, Rojas AV, Maisuradze GG, Scheraga HA, Lobel P, Maxfield FR. Lysosomal enzyme tripeptidyl peptidase 1 destabilizes fibrillar Aβ by multiple endoproteolytic cleavages within the β-sheet domain. PNAS 2018;115:1493–8. https://doi.org/10.1073/pnas.1719808115.

[26] Cruz D, El-Banna M, Majumdar A, Sleat DE, Muldowney M, Lobel P, et al. Lysosomal enzyme tripeptidyl peptidase 1 plays a role in degradation of beta amyloid fibrils. BioRxiv 2019:639682. https://doi.org/10.1101/639682.

[27] Nagata Y, Hirayama A, Ikeda S, Shirahata A, Shoji F, Maruyama M, et al. Comparative analysis of cerebrospinal fluid metabolites in Alzheimer's disease and idiopathic normal pressure hydrocephalus in a Japanese cohort. Biomark Res 2018;6:1–11. https://doi.org/10.1186/s40364-018-0119-x.

[28] Menni C, Kastenmüller G, Petersen AK, Bell JT, Psatha M, Tsai P-C, et al. Metabolomic markers reveal novel pathways of ageing and early development in human populations. Int J Epidemiol 2013;42:1111–9. https://doi.org/10.1093/ije/dyt094.

[29] Mapstone M, Cheema AK, Fiandaca MS, Zhong X, Mhyre TR, MacArthur LH, et al. Plasma phospholipids identify antecedent memory impairment in older adults. Nature Medicine 2014;20:415–8. https://doi.org/10.1038/nm.3466.

[30] Klavins K, Koal T, Dallmann G, Marksteiner J, Kemmler G, Humpel C. The ratio of phosphatidylcholines to

lysophosphatidylcholines in plasma differentiates healthy
controls from patients with Alzheimer's disease and mild
cognitive impairment. Alzheimer's & Dementia: Diagnosis,
Assessment & Disease Monitoring 2015;1:295–302.
https://doi.org/10.1016/j.dadm.2015.05.003.
[31] Casanova R, Varma S, Simpson B, Kim M, An Y, Saldana S,
et al. Blood metabolite markers of preclinical Alzheimer's
disease in two longitudinally followed cohorts of older
individuals. Alzheimer's & Dementia 2016;12:815–22.
https://doi.org/10.1016/j.jalz.2015.12.008.
[32] Li D, Misialek JR, Boerwinkle E, Gottesman RF, Sharrett
AR, Mosley TH, et al. Plasma phospholipids and prevalence
of mild cognitive impairment and/or dementia in the ARIC
Neurocognitive Study (ARIC-NCS). Alzheimer's &
Dementia: Diagnosis, Assessment & Disease Monitoring
2016;3:73–82. https://doi.org/10.1016/j.dadm.2016.02.008.
[33] Varma VR, Oommen AM, Varma S, Casanova R, An Y,
Andrews RM, et al. Brain and blood metabolite signatures of
pathology and progression in Alzheimer disease: A targeted
metabolomics study. PLOS Medicine 2018;15:e1002482.
https://doi.org/10.1371/journal.pmed.1002482.
[34] Xu Q, Zhang Y, Zhang X, Liu L, Zhou B, Mo R, et al.
Medium-chain triglycerides improved cognition and lipid
metabolomics in mild to moderate Alzheimer's disease
patients with APOE4-/-: A double-blind, randomized,
placebo-controlled crossover trial. Clin Nutr 2019.
https://doi.org/10.1016/j.clnu.2019.10.017.
[35] Pardridge WM. The blood-brain barrier: Bottleneck in brain
drug development. Neurotherapeutics 2005;2:3–14.
https://doi.org/10.1602/neurorx.2.1.3.
[36] Björkhem Ingemar, Meaney Steve. Brain Cholesterol: Long
Secret Life Behind a Barrier. Arteriosclerosis, Thrombosis,
and Vascular Biology 2004;24:806–15.
https://doi.org/10.1161/01.ATV.0000120374.59826.1b.
[37] Zhang J, Liu Q. Cholesterol metabolism and homeostasis in
the brain. Protein Cell 2015;6:254–64.
https://doi.org/10.1007/s13238-014-0131-3.
[38] Liu G, Yao L, Liu J, Jiang Y, Ma G, Chen Z, et al.
Cardiovascular disease contributes to Alzheimer's disease:
evidence from large-scale genome-wide association studies.
Neurobiology of Aging 2014;35:786–92.
https://doi.org/10.1016/j.neurobiolaging.2013.10.084.
[39] Shang H, Liu G, Jiang Y, Fu J, Zhang B, Song R, et al.
Pathway Analysis of Two Amyotrophic Lateral Sclerosis
GWAS Highlights Shared Genetic Signals with Alzheimer's

Disease and Parkinson's Disease. Mol Neurobiol
2015;51:361–9. https://doi.org/10.1007/s12035-014-8673-1.
[40] Chen J, Xie C, Zhao Y, Li Z, Xu P, Yao L. Gene expression
analysis reveals the dysregulation of immune and metabolic
pathways in Alzheimer's disease. Oncotarget 2016;7.
https://doi.org/10.18632/oncotarget.12505.
[41] Lin S, Liu H, Kanawati B, Liu L, Dong J, Li M, et al.
Hippocampal metabolomics using ultrahigh-resolution mass
spectrometry reveals neuroinflammation from Alzheimer's
disease in CRND8 mice. Anal Bioanal Chem 2013;405:5105–
17. https://doi.org/10.1007/s00216-013-6825-1.
[42] Yu J, Kong L, Zhang A, Han Y, Liu Z, Sun H, et al. High-
Throughput Metabolomics for Discovering Potential
Metabolite Biomarkers and Metabolic Mechanism from the
APPswe/PS1dE9 Transgenic Model of Alzheimer's Disease.
J Proteome Res 2017;16:3219–28.
https://doi.org/10.1021/acs.jproteome.7b00206.
[43] Schubert D. Glucose metabolism and Alzheimer's disease.
Ageing Research Reviews 2005;4:240–57.
https://doi.org/10.1016/j.arr.2005.02.003.
[44] Kandimalla R, Thirumala V, Reddy PH. Is Alzheimer's
disease a Type 3 Diabetes? A critical appraisal. Biochimica et
Biophysica Acta (BBA) - Molecular Basis of Disease
2017;1863:1078–89.
https://doi.org/10.1016/j.bbadis.2016.08.018.
[45] An Y, Varma VR, Varma S, Casanova R, Dammer E,
Pletnikova O, et al. Evidence for brain glucose dysregulation
in Alzheimer's disease. Alzheimer's & Dementia
2018;14:318–29. https://doi.org/10.1016/j.jalz.2017.09.011.
[46] Nho K, Kueider-Paisley A, Ahmad S, MahmoudianDehkordi
S, Arnold M, Risacher SL, et al. Association of Altered Liver
Enzymes With Alzheimer Disease Diagnosis, Cognition,
Neuroimaging Measures, and Cerebrospinal Fluid
Biomarkers. JAMA Netw Open 2019;2:e197978.
https://doi.org/10.1001/jamanetworkopen.2019.7978.
[47] Dietschy JM, Turley SD. Cholesterol metabolism in the brain.
Current Opinion in Lipidology 2001;12:105.
[48] Frisardi V, Panza F, Seripa D, Farooqui T, Farooqui AA.
Glycerophospholipids and glycerophospholipid-derived lipid
mediators: A complex meshwork in Alzheimer's disease
pathology. Progress in Lipid Research 2011;50:313–30.
https://doi.org/10.1016/j.plipres.2011.06.001.
[49] Nitsch R, Pittas A, Blusztajn JK, Slack BE, Growdon JH,
Wurtman RJ. Alterations of phospholipid metabolites in
postmortem brain from patients with Alzheimer's disease.

Ann N Y Acad Sci 1991;640:110–3.

https://doi.org/10.1111/j.1749-6632.1991.tb00200.x.

[50] Mulder C, Wahlund L-O, Teerlink T, Blomberg M, Veerhuis

R, van Kamp GJ, et al. Decreased

lysophosphatidylcholine/phosphatidylcholine ratio in

cerebrospinal fluid in Alzheimer's disease. J Neural Transm

2003;110:949–55. https://doi.org/10.1007/s00702-003-0007-

9.

[51] Ross BM, Moszczynska A, Erlich J, Kish SJ. Phospholipid-

Metabolizing Enzymes in Alzheimer's Disease: Increased

Lysophospholipid Acyltransferase Activity and Decreased

Phospholipase A2 Activity. Journal of Neurochemistry

1998;70:786–93. https://doi.org/10.1046/j.1471-

4159.1998.70020786.x.

[52] Zannis VI, Koukos G, Drosatos K, Vezeridis A, Zanni EE,

Kypreos KE, et al. Discrete roles of apoA-I and apoE in the

biogenesis of HDL species: Lessons learned from gene

transfer studies in different mouse models. Annals of

Medicine 2008;40:14–28.

https://doi.org/10.1080/07853890701687219.

Table 1. Wisconsin ADRC and WRAP participant characteristics.

Participant Characteristics	CSF						Plasma
	Wisconsin ADRC						
	AD (n = 38)	MCI (n = 29)	HOC (n = 40)	IMPACT (n = 158)	WRAP (n = 130†)	P values*	WRAP (n = 123)
Female: n (%)	12 (31.6)	8 (27.6)	22 (55.0)	118 (74.7)	85 (65.4)	0.11	80 (65.0)
Age in years: mean (SD)	71.4 (8.9)	74.2 (8.3)	74.1 (4.8)	57.8 (5.3)	61.5 (6.6)	<0.001	62.6 (6.5)
Education in years: mean (SD)	14.8 (2.9)	16.3 (2.8)	16.4 (2.9)	16.0 (2.3)	16.1 (2.2)	0.50	16.1 (2.2)
CSF t-tau: mean (SD)	775.5 (332.9)	573.0 (346.9)	436.8(194.5)	283.1 (144.2)	311.5 (117.7)	0.03	318.9 (122.7)
CSF p-tau: mean (SD)	74.8 (27.3)	62.4(36.2)	52.9 (19.4)	39.8 (14.5)	41.5 (13.4)	0.10	42.6 (13.5)

* The p values are based on a two-sample t test conducted between the Wisconsin ADRC IMPACT cohort and WRAP (shaded columns). †The 130 individuals had 210 longitudinal samples.

866 **Table 2. Prediction performance (r²) of each model in IMPACT and WRAP.**

	IMPACT		WRAP	
	T-tau	**P-tau**	**T-tau**	**P-tau**
Null models: demographics* only	0.083	0.106	0.085	0.087
Enhanced models: demographics and 38 CSF metabolites				
Overall	0.703	0.757	0.624	0.651
Female	0.787	0.791	0.591	0.640
Male	NA†	NA†	0.794	0.804
Demographics and 7 LASSO selected CSF metabolites	0.594	0.692	0.585	0.615
Demographics and 34 Plasma metabolites	NA†	NA†	0.269	0.301

867 * Demographics are age, sex, and years of education.
868 † The sample size for males was too small to perform the analysis in IMPACT.

886　**Table 3. LASSO results for CSF t-tau and p-tau in IMPACT and WRAP.**

IMPACT		WRAP	
T-tau			
Biochemical Name	**Coefficient**	**Biochemical Name**	**Coefficient**
X - 24228	202.10	**N-acetylneuraminate**	333.57
N-acetylneuraminate	188.68	**C-glycosyl tryptophan**	264.20
Beta-citrylglutamate	151.01	X - 12906	97.42
C-glycosyl tryptophan	92.25	1-oleoyl-GPC (18:1)	69.00
N-acetylthreonine	78.41	**X - 10457**	62.67
1-palmitoyl-GPC (16:0)	72.78	N6-succinyladenosine	50.65
X - 10457	42.70	**X - 24228**	36.65
X - 24699	33.25	X - 23739	28.95
		1-palmitoyl-GPC (16:0)	1.38
		1-palmitoyl-2-palmitoleoyl-GPC (16:0/16:1)	0.21
		X - 18887	-3.56
		Ribonate	-112.48
P-tau			
C-glycosyl tryptophan	28.80	**N-acetylneuraminate**	30.80
N-acetylneuraminate	24.98	**C-glycosyl tryptophan**	30.55
Beta-citrylglutamate	15.71	**X - 10457**	9.80
X - 24228	7.37	N6-succinyladenosine	7.41
X - 10457	5.47	**X - 24228**	5.59
1-palmitoyl-GPC (16:0)	4.26	X - 12906	3.66
Cholesterol	2.52	Sphingomyelin (d18:1/14:0, d16:1/16:0)	3.05
1-oleoyl-GPC (18:1)	2.44	X - 24699	3.04
X - 24329	1.25	**1-oleoyl-GPC (18:1)**	2.33
Gulonate	1.13	**1-myristoyl-2-palmitoyl-GPC (14:0/16:0)**	0.88
1-myristoyl-2-palmitoyl-GPC (14:0/16:0)	1.07	X - 18887	-2.50
		Ribonate	-4.06

887　Metabolites shaded in light grey with bold font are consistent across IMPACT and WRAP.

Table 4. Pathway analysis results for the 38 CSF metabolites.

KEGG Pathway	Pathway Enrichment					Pathway Impact
	Total # Metabolites in KEGG Pathway	# Metabolites Identified in Present Study	Raw p	-Log(p)	FDR	Impact
Pentose and glucuronate interconversions	18	3	2E-04	8.80	0.01	0.25
Glycerophospholipid metabolism	36	2	0.02	3.86	0.88	0.11
Linoleic acid metabolism	5	1	0.03	3.45	0.89	0
Ascorbate and aldarate metabolism	8	1	0.05	2.98	1	0
alpha-Linolenic acid metabolism	13	1	0.08	2.51	1	0
Sphingolipid metabolism	21	1	0.13	2.06	1	0
Arachidonic acid metabolism	36	1	0.21	1.56	1	0
Steroid biosynthesis	42	1	0.24	1.42	1	0.03
Primary bile acid biosynthesis	46	1	0.26	1.34	1	0.05
Steroid hormone biosynthesis	85	1	0.43	0.84	1	0.005

Raw p is the original p value calculated from the enrichment analysis; FDR p is the p value adjusted using False Discovery Rate; Impact is the pathway impact value calculated from the pathway topology analysis. The shaded two pathways are considered important.

904
905
(a)

906
907
(b)

908 **Figure 1. ROC curves and AUC scores of predictions by 6 models in the Wisconsin ADRC (a) AD vs. HOC (b)**
909 **MCI vs. HOC.** Base model: age, sex, years of education, *APOE* ε4 count, t-tau, p-tau, and Aβ42; base model replacing
910 t-tau and p-tau with the seven selected metabolites from LASSO; and base model plus the seven selected
911 metabolites from LASSO.

912
Figure 2. Pathway analysis results for 23 CSF metabolites. The x-axis represents the pathway impact, and y-axis
represents the pathway enrichment. Larger sizes and darker colors represent higher pathway impact and enrichment,
respectively.
916