Title: **Extracellular matrix stiffness regulates degradation of the Hippo kinase MST2 via SCF βTrCP**

Authors: Ana Paula Zen Petisco Fiore[1], Ana Maria Rodrigues da Silva[1], Helder Veras Ribeiro Filho[2], Antonio Carlos Manucci[1], Pedro de Freitas Ribeiro[1], Mayara Carolinne Silva Botelho[1], Paulo Sergio Lopes de Oliveira[2], Michele Pagano[3] & Alexandre Bruni-Cardoso[1*].

[1]Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, São Paulo 05508-000, Brazil.
[2]Laboratório Nacional de Biociências, Centro Nacional de Pesquisa em Energia e Materiais, Campinas 13083-970, Brazil.
[3]Department of Biochemistry and Molecular Pharmacology, and Howard Hughes Medical Institute, New York University Grossman School of Medicine, New York, NY 10016, USA.

[*]corresponding author (brunicar@iq.usp.br)

Abstract

Tumor microenvironments display disrupted mechanical properties, including altered extracellular matrix (ECM) rigidity. ECM stiffening perturbs cell tensional homeostasis resulting in activation of mechanosensing transcriptional co-activators, such as the Hippo pathway effectors YAP and TAZ. The Hippo pathway plays central roles in development and tumorigenesis, but how the proteostasis of the Hippo kinase MST2 is regulated remains unknown. We show that MST2 levels decrease upon changes in ECM rigidity via proteasome degradation. MST2 degradation is enhanced in human breast epithelial cells that are cultured in stiffer microenvironments due to integrin and integrin-linked kinase activation. MST2 knockdown resulted in increased nucleus-to-cytoplasm ratio of YAP, increased proliferation rates in a soft microenvironment and F-actin alignment in cells cultured in intermediate stiffness. We found that MST2 is ubiquitinated by the SCF βTrCP ubiquitin ligase, and site-directed mutagenesis combined with computational molecular dynamics studies revealed that βTrCP binds MST2 via a noncanonical degradation motif. Our study uncovers the underlying biochemical mechanisms controlling MST2 degradation and demonstrates how alterations in the microenvironment rigidity regulate the proteostasis of a central Hippo pathway component.

Introduction

The tissue microenvironment is a source of mechanical signals, such as those provided by stiffness of the extracellular matrix (ECM), that regulate cell shape and several cellular processes (*1, 2*). Cells sense composition and mechanical properties of the ECM eliciting adjustment in tension and arrangement of the actomyosin cytoskeleton (*3, 4*). Altered ECM mechanical properties are found in many diseases, including cancers (*5-7*). As cancer progresses, malignant cells disturb the surrounding tissue physical properties (*5*). Reciprocally, mounting evidence has shown that tissue stiffening can induce tumorigenesis (*4, 5, 8, 9*). Commonly used as diagnosis and prognosis of several types of cancer, increased tissue stiffness is caused by ECM deposition and remodeling. And, a stiff ECM activates nuclear mechanosensing relays, such as the co-activator factors YAP (Yes-associated protein)/TAZ (transcriptional co-activator with PDZ-binding motif, also known as WWTR1) (*10*), that promote proliferation and invasion (*5, 8, 9*).

In epithelial cells exposed to physiological rigidities, YAP/TAZ are predominantly cytoplasmic but translocate to the nucleus in stiff microenvironments (*11*). YAP/TAZ subcellular localization and transcriptional activity are modulated by Hippo, a pathway cored by the kinases MST1/2 and LATS1/2. On the other hand, YAP/TAZ mechanosignaling promotes fibrosis, resulting in microenvironment stiffening (*12-24*). The transduction of mechanical signals that results in cell shape remodeling and in changes in gene expression are initially sensed by the ECM receptors, integrins, which in turn trigger assembly of focal adhesions (FAs), multiprotein complexes linked to filamentous actin (*2, 25*). FAs are composed of Focal Adhesion Kinase (FAK) and/or Integrin-Linked Kinase (ILK), integrin clusters and scaffold proteins, and are modulated by substrate rigidity in response to intracellular tension built in the actomyosin cytoskeleton (*26, 27*). FAK and ILK phosphorylate multiple substrates and help converge integrin and growth factor-receptor triggered signaling pathways (*28*).

Although a great deal is known regarding regulation of YAP/TAZ levels and activities by the ECM stiffness, little is known about the mechanical regulation of the Hippo pathway components. We sought at investigating whether MST2, a core Hippo serine threonine kinase, was modulated by microenvironment rigidity and found that a stiff microenvironment induces MST2 proteolysis in human breast epithelial cells. Silencing of *STK3*, the gene encoding MST2, resulted in increased nucleus/cytoplasm ratio of YAP, increased proliferation rates in a soft microenvironment and F-actin alignment in cells cultured in intermediate stiffness. We found that MST2 is ubiquitinated by the E3-ligase

complex Skp1/Cul1/F-box-protein/β-transducing-repeat-containing protein (SCF$^{\beta TrCP}$) and degraded via the proteasome. The E3-ligase substrate-recognition subunit βTrCP binds to MST2 via a noncanonical degradation motif (degron). We also found that signaling from the integrin-ILK (integrin-linked kinase) axis results in increased MST2 and βTrCP binding, leading to MST2 degradation. Our findings shed light on the mechanisms controlling Hippo activity by targeted proteolysis and that degradation of MST2 is a crucial part in the mechanoregulatory circuitry that senses and transduces physical cues from the cell microenvironment.

Results

Stiff microenvironments reduce MST2 levels

We asked whether the level of MST2 was regulated by ECM stiffness. For this, we cultured MCF10A cells, a non-tumoral human mammary cell line highly sensitive to modulation of substrate stiffness (*8*), on fibronectin-coated polyacrylamide hydrogels that can be tuned in rigidity by changing the concentration of polyacrylamide (*29*). Three degrees of elastic modulus were used: 0.48 KPa (Low; a physiological rigidity), 4.47 KPa (Mid; average rigidity of the mammary tumor stroma) and 40.40 KPa (Hi; a supraphysiological stiffness).

Immunoblotting of MCF10A-cell lysates revealed that levels of MST2 decrease in response to a stiffer ECM (Fig. 1A). Phosphorylation of YAP at Serine 127, a mark for cytoplasmic retention of YAP due to Hippo signaling activity, was decreased in stiffer ECM substrates (Fig. 1A). Intriguingly, *STK3* (the gene encoding MST2) mRNA levels did not change is response to different rigidities (Fig. 1B), suggesting that regulation of MST2 protein levels could be via proteolytic degradation in different mechanical contexts.

We silenced *STK3* using siRNA to evaluate whether modulation of MST2 levels would reflect in alterations in the nuclear to cytoplasm ratio of YAP and also phenotypical changes in MCF10A cells (Fig. 1C, Supp. Fig. 1A and Fig. 1D). In unperturbed MCF10A cells grown in Low stiffness,

Figure 1: The extracellular matrix stiffness regulates the level of MST2, and *STK3* knockdown alter cellular mechanoregulatory outputs.

A) Immunoblots for MST2, phosphorylated YAP^{Ser127} (pYAP^{Ser127}), total YAP and α-tubulin (used as a loading control) of MCF10A cells cultured on fibronectin-1 (FN1) coated hydrogels with different rigidities: 0.48 KPa (Low), 4.47 KPa (Mid) and 40.40 (Hi). Fold changes of MST2 levels are shown under MST2 immunoblots.

B) Quantitative RT-PCR for *STK3* (gene that encodes MST2) of MCF10A cells cultured on hydrogels with different degrees of rigidity. GAPDH was used as an endogenous control to normalize gene expression. Data are expressed as mean ± SEM of e$^{-\Delta\Delta ct}$ of 3 replicates.

C) Quantitative RT-PCR for *STK3* of MCF10A cells silenced for *STK3* cultured on hydrogels. GAPDH was used as an endogenous control to normalize gene expression. MCF10A cells were transfected with an siRNA for *STK3* (siSTK3) or nontargeting siRNA (siControl). Data are expressed as mean ± SEM of $e^{-\Delta\Delta ct}$. See also Figure S1A for MST2 protein expression after siRNA treatments.

D) Representative microscope images of immunofluorescence (IF) for YAP (red) of MCF10A cells silenced for *STK3* cultured on hydrogels. See Figure 1S2 for images of YAP staining paired with DNA labeling with DAPI.

E) Quantification of nuclear-to-cytoplasmic ratio of YAP staining of MCF10A cells silenced for *STK3*.

F) Quantification of EdU-positivity reflecting proliferation of MCF10A cells silenced for *STK3*.

G) Representative low magnification Airy-Scan superresolution microscope images of MCF10 cells silenced for *STK3* and stained with fluorescent phalloidin for F-actin visualization (magenta).

H) Aspect ratio changes of MCF10A cells silenced for *STK3*. Top: schematic depicting a rounded cell and an elliptical cell with highlighting its major and minor axes. Bottom: Quantification of cell aspect ratio.

I) Quantification of the area of MCF10A cells silenced for *STK3*.

J) Representative high magnification Airy-Scan superresolution microscope images of MCF10 cells silenced for *STK3* and stained with phalloidin (magenta) and vinculin (green).

K) Quantification of F-actin alignment in MCF10A cells silenced for *STK3*. Data are plotted for the three degrees of stiffness as frequency of fibers as a function of their directionality in degrees (°) with respect to the major cell axis (a directionality of 0° means perfect alignment with the major cell axis).

Statistical significance = *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$ and n.s denotes no statistical significance. Scale bars, D =10 µm, G = 40 µm, J = 5 µm.

YAP is predominantly cytoplasmic, as determined by measuring nuclear to cytoplasmic YAP ratio (Fig. 1D-E and Supp. Fig. 1B). YAP preferentially localizes in the nucleus in response to stiffer substrates (*10*). Silencing *STK3* results in increased nucleus to cytoplasm YAP ratios in Low and Mid stiffness in comparison to cells treated with a control siRNA (siControl). In stiff ECM, knockdown of *STK3* does not change the nucleus-to-cytoplasm YAP ratio. Cell proliferation, as measured by EdU incorporation, increases in Mid and Hi rigidities in comparison to cells cultured in a soft ECM (Fig. 1F and Supp. Fig. 1C). However, silencing *STK3* resulted in higher percentages of EdU-positive cells in all three degrees of rigidity.

Knocking down MST2 resulted in striking morphological changes particularly in Low and Mid stiffnesses (Fig. 1G-J). In low stiffness, siControl MCF10A cells behave as a normal epithelium: cells acquire a rounded morphology, F-actin is largely cortical, vinculin (a focal adhesion protein) staining is not defined in a classical focal adhesion (FA) pattern (Fig. 1G and Fig. 1J). On the other hand, in stiffer hydrogels, MCF10A cells are flatter and more spread - despite not losing cell-to-cell contact – display F-actin bundles and a classical focal FA pattern of vinculin staining. In Hi stiffness, cells were less rounded as evidenced by increased cell aspect ratio. (Fig. 1G and Fig. 1H). MCF10A cells treated with siSTK3 were larger in

Low and Mid rigidities and displayed a less rounded morphology in Mid rigidity in comparison to cells treated with control siRNA (Fig. 1I). In correspondence to the morphological changes, siSTK3-treated cells displayed in Mid and Hi rigidities pronounced stress fibers that were more aligned (in relation to the cell's major axes) and colocalized with typical FAs, as revealed by Airy-Scan superresolution microscopy (Fig. 1J-K). Altogether, these data revealed ECM rigidity modulates MST2 levels and that this can affect changes in YAP subcellular localization and result in actin cytoskeleton and morphological changes.

MST2 is degraded via the proteasome after ubiquitination by SCF$^{\beta TrCP}$

Because MST2 protein levels decreased in response to stiffer ECMs - albeit its coding mRNA remained unaltered in the same conditions- we decided to investigate whether inhibition of the proteasome with MG132 or cullin-1 neddylation (covalent conjugation of the ubiquitin-like protein NEDD8 required for activation of SCF E3-ligase complexes) with MLN4924 would result in accumulation of MST2 in a panel of non-malignant and malignant human breast epithelial cell lines. Accordingly, treatment with MG132 or MLN4924 resulted in MST2 accumulation in 4 of 7 human breast cell lines grown on regular cell culture plastic plates (Fig. 2A).

To identify which specific E3-ligase complex would be responsible for ubiquitinating MST2, we assessed the literature and the available MST2 interactomes. Mass spectrometry analysis performed in our laboratory and by others independently identified three E3-ligase substrate-recognition subunits in the MST2 interactomes: FBXO3, SKP2 and βTrCP (*30-32*). Furthermore, a recent report demonstrated that βTrCP binds MST2 in an precipitation assay of ectopically expressed proteins and that inhibition of the proteasome resulted in MST2 accumulation in HEK293T cells (*33*). We transfected HEK293T cells with flag-tagged βTrCP, FBXO3, or SKP2 constructs and performed immunoprecipitation assays followed by immunoblotting (Fig. 2B). Endogenous MST2 was detected only in the flag-tagged βTrCP immunoprecipitants. βTrCP is part of the SCF$^{\beta TrCP}$ E3-ligase complex which tags key regulators of proliferation (including YAP/TAZ (*34*)), DNA repair and apoptosis for degradation (*35-37*). As other SCF complexes, SCF$^{\beta TrCP}$ consists of 4 main subunits: SKP1,

A

B

IP Flag

	FBXO3	βTrCP	SKP2

CUL1 ←Nedd

MST2

SKP1

Flag

C

substrate

βTrCP

SKP1

CUL1 N

RBX1

E2 U

E1 U

proteassome

D

IP Flag

MG − + − + − +

empty empty βΔFbox βΔFbox βTRCP βTRCP

MST2

CUL 1 ←Nedd

SKP1

Flag

E

βΔFbox-Flag	−	−	+	−
βTrCP-Flag	−	−	−	+
MST2-FFSS	−	+	+	+
Ub-Myc	+	+	+	+

Myc-Tag (Ub)

MST2

F

MCF10A

	Si Control	Si βTRCP 24h

MST2
1.0   2.2

βTrCP1

p21

Cul1

SKP1

β-actin

G

MCF10A

IP   IgG   βTrCP

MG   −   +   −   +

MST2 ← ← IgG heavy chain

βTrCP ← ← IgG heavy chain

CUL1

SKP1 ← ← IgG light chain

Figure 2: MST2 is ubiquitinated by the SCF$^{\beta TrCP}$ E3 ligase complex and degraded via proteasome:
A) Immunoblots for MST2 in protein extract from breast epithelial cell lines cultured on plastic plates in the absence (-) or presence (+) of the proteasome inhibitor MG132 (MG) or the CUL1 neddylation inhibitor MLN4924 (MN) for 3 hours. Cell lines in which MST2 was accumulated upon treatments are highlighted (red outline). β-actin was used as a loading control. Fold changes of MST2 levels are shown under immunoblots for MST2.
B) Co-immunoprecipitation (co-IP) FBXO3, βTrCP and SKP2 followed by immunoblotting for CUL1, MST2, SKP1 and Flag. HEK293T cells were transfected with Flag-tagged constructs FBXO3, βTrCP and SKP2. Transfected cells were treated for 1.5 hour with MG132, lysed and the immunoprecipitation was performed using M2 antibody coated agarose beads. Both CUL1 and SKP1 are proteins that interact with F-box proteins and were used as Co-IP controls. Flag was detected to assess the success of precipitation of the constructs.
C) Schematic of the E3 ligase complex SCF$^{\beta TrCP}$. As other SKP1-CUL1-Fbox (SCF) complexes, SCF$^{\beta TrCP}$ consists of 4 main subunits: SKP1, CUL1, RBX1 and the Fbox protein βTrCP that functions as a receptor for target proteins, giving specificity to the complex. Protein degradation by the 26S proteasome depends on a cascade of enzymes. The E1 enzyme binds and activate ubiquitin (U), followed by the recruitment of an E2 ubiquitin-conjugating enzyme that is then loaded with activated ubiquitin. E2 interacts with the E3 complex via the adaptor protein (RBX). After a sequence of ubiquitin conjugation cycles the target protein is degraded by the proteasome. CUL neddylation (N) is required for SCF activation.
D) Co-IP confirming the direct binding of MST2 and βTrCP. HEK293T cells were transfected with Flag-tag constructs fused to full length βTrCP or the dominant negative βTrCP (βΔFbox), were treated or not with MG132 for 1.5 hour, lysed and Flag immunoprecipitation was performed using M2 antibody coated beads. Endogenous MST2, CUL1 and SKP1 (used as controls for the co-IP) and flag were detected by immunoblotting. Flag was detected to assess the success of the assay.
E) Detection of MST2 ubiquitination. HEK293T cells were transfected with Flag-Flag-Strep-Step-tagged MST2 (FFSS-MST2) Flag-tagged β-TrCP or Flag-tagged βΔFbox and Myc-tagged Ubiquitin (Ub-Myc). After a stringent extraction, the FFSS_MST2 was precipitated with tactin-coated beads. To evaluate ubiquitination of MST2, Ub-Myc was detected with immunoblotting using an antibody against Myc-tag. MST2 was also detected by immunoblotting.
F) Silencing of βTRCP in MCF10A cells results in MST2 accumulation. Cells were cultured in the presence of siRNA for βTRCP or a non-targeting siRNA (siControl). Immunoblots confirming that silencing βTRCP was efficient and resulted in increased MST2 levels. Immunoblot of p21 was used as a negative control, since it is not a βTRCP substrate. SKP1 and CUL1 did not vary with βTRCP silencing. β-actin was used as loading control. Fold changes of MST2 levels are shown under MST2 immunoblots.
G) Co-IP for βTRCP confirming the interaction of endogenous MST2 and βTrCP in MCF10A cells. Cells were culture for 1.5 hour in the presence of MG132, lysed and immunoprecipitation was performed with a βTrCP antibody or a control IgG, followed by incubation with protein A/G-coated magnetic beads. MST2, CUL1 and SKP1 were only detected by immunoblot in βTrCP immunoprecipitants. Arrows indicate bands corresponding to IgG that were detected by the secondary antibody.

CUL1, RBX1 and the Fbox protein βTrCP, the latter acts as a substrate receptor, providing specificity to the complex (*38*) (Fig. 2C).

To confirm the direct interaction of βTrCP and MST2, we transfected HEK293T cells with a full length βTrCP or a βTrCP lacking the whole Fbox domain (βΔFbox), a sequence of ~40 amino acids that is required for Fbox proteins to bind SKP1 and, hence, to assemble the SCF complex (*39*). Co-IP followed by immunoblotting revealed that MST2 binds to βΔFbox showing a more intense band than the βTrCP lane (Fig. 2D). Proteasome inhibition increased the interaction between βTrCP and MST2, but resulted in no increment of βΔFbox and MST2 binding. This occurred, probably, because substrates bound to βΔFbox are not ubiquitinated. These co-IP experiments confirmed that MST2 interacts with βTrCP even in the absence of the intact SCF^{βTrCP} complex, indicating that MST2 binds directly βTrCP and not via other SCF subunits. A ubiquitination assay using Myc-tagged ubiquitin confirmed that MST2 is, indeed, ubiquitinated only in the presence of βTrCP but not βΔFbox (Fig. 2E).

Next, we assessed if our findings of βTrCP binding and regulation of MST2 levels in HEK293T cells would be consistent in a breast epithelial cell line without ectopic overexpression. Silencing βTrCP in MCF10A cells resulted in increased levels of MST2 (Fig. 2F). The level of p21, which is not a βTrCP substrate, CUL1 and SKP1 did not change upon βTrCP knockdown. Crucially, a co-IP for MCF10A cell lysates confirmed the interaction of endogenous βTrCP and MST2 proteins, reinforcing the robustness of our findings and corroborating that the results from assays with HEK293T cells are not specific to this cell line neither an artifact resulting from nonphysiological overexpression of protein constructs (Fig. 2G).

Identification of a βTrCP binding site in MST2

The F-box protein subunits of SCFs, such as βTrCP, recognize their substrates via degradation motifs termed degrons (*40*). Most known SCF^{βTrCP} substrates contain the consensus sequence DpSGXX(X)pS (Fig. 3A), a biphosphorylated motif, usually localized in structurally disordered regions (*41*), that mediates the interaction of the substrate with the WD40 repeats of βTrCP. Although, MST2 does not contain a canonical βTrCP degron, we searched for sequences chemically similar to the highly conserved three amino-acid residues of the canonical degron (DpSG). Four putative tripeptide degrons are present in the human MST2 protein sequence: EDS (16-18), ESG (48-50), ESD (359-361), and EDG (375-377) (Fig. 3B). Noteworthy, similar sequences are found in other noncanonical degrons of experimentally validated SCF^{βTrCP} targets (*42-46*). We generated four MST2 constructs each containing a deletion for one of the four tripeptides and transfected them into HEK293T cells for co-IP experiments using an antibody against

βTrCP. All 4 deletions affected MST2 binding to βTrCP, however deletion of the tripeptide EDG (375-377) almost abolished the interaction (Fig. 3C).

Because deletion of the MST2 EDG tripeptide produced the most prominent impairment in βTrCP interaction, we decided to model the MST2 segment that includes the EDG tripeptide in a complex with the βTrCP WD40 domain to investigate the structural basis that guides their recognition and interaction. The EDG tripeptide and most of its flanking residues are localized, as most degron motifs of E3-ligase targets, in a disordered region of MST2 (Supplementary Figure 2A). In comparison to the canonical βTrCP degron motif (Figure 3A), the first phosphorylated serine is substituted by an aspartic acid residue (D376) in MST2 EDG tripeptide, maintaining its negative charge feature (Figure 3A). Additional glutamic acid residues N-terminal to EDG could also help provide electrostatic forces that compensate the lack of phosphorylation. The second phosphorylated serine from the canonical degron (position +4 to +5 in relation to the first phosphorylated serine) is not present in the linear sequence of MST2, and two positively charged residues occupy this position instead.

Molecular dynamics (MD) simulation of an MST2 11-mer peptide (EE**EDG**TMKRNA), starting from the position of β-catenin 11-mer peptide crystallographic structure in complex with βTrCP (see methods), shows a small deviation at the EDG tripeptide region relative to its initial position (Supplementary Figure 2B-D). The stability in this region is a consequence of putative hydrogen bonds between D376 in MST2 (structural position similar to pS33 from β-catenin peptide) and R285 in βTrCP as well as E375 in MST2 (structural position similar to D32 from β-catenin peptide) and R474 in β-TrCP (Figure 3D and Supplementary Figure 2E). In MST2, E373 also contacts R285 through hydrogen bonds. Another similarity between MST2 and β-catenin peptide binding is the presence of a hydrophobic residue (M379 of MST2 and I35 in the β-catenin peptide) pointing towards the βTrCP β-propeller channel (Figure 3D and Supplementary Figure 2E).

During the simulation, K380 in MST2 is displaced from its initial position to avoid repulsive contact with R431 in βTrCP (Supplementary Figure 2B), which anchors the second phosphorylated serine in canonical degrons (Supplementary Figure 2E). The presence of βTrCP positive charges favors K380 in MST2 to be exposed to solvent. This is an interesting finding, since an *in silico* analysis predicted K380 could be ubiquitinated (Supplementary Figure 3A).

A

B

Human MST2:

```
01 MEQPPAPKSKLKKLSEDSLTKQPEEVFDVLEKLGEGSYGSVFKAIHKESG 50
51 QVVAIKQVPVESDLQEIIKEISIMQQCDSPYVVKYYGSYFKNTDLWIVME 100
101 YCGAGSVSDIIRLRNKTLIEDEIATILKSTLKGLEYLHFMRKIHRDIKAG 150
151 NILLNTEGHAKLADFGVAGQLTDTMAKRNTVIGTPFWMAPEVIQEIGYNC 200
201 VADIWSLGITSIEMAEGKPPYADIHPMRAIFMIPTNPPPTFRKPELWSDD 250
251 FTDFVKKCLVKNPEQRATATQLLQHPFIKNAKPVSILRDLITEAMEIKAK 300
301 RHEEQQRELEEEEENSDEDELDSHTMVKTSVESVGTMRATSTMSEGAQTM 350
351 IEHNSTMLESDLGTMVINSEDEEFEDGTMKRNATSPQVQRPSFMDYFDKQ 400
401 DFKNKSHENCNQNMHEPFPMSKNVFPDNWKVPQDGDFDFLKNLSLEELQM 450
451 RLKALDPMMEREIEELRQRYTAKRQPILDAMDAKKRRQQNF 491
```

C

cell extract   IP. βTrCP

FFSS_MST2
(motif deletion)

wt 16-18 (EDS) 48-50 (ESG) 359-361 (ESD) 375-377 (EDG)

MST2 (Flag)

βTrCP

D

β-TrCP WD40 domain

MST2 segment

90°

β-TrCP WD40 domain

MST2 segment

E

```
STK3_HUMAN       346  GAQ-----TMIEHNSTMLESDLGTMVINSEDE------EEEDGTMKRNATSPQVQRPSFM  394
STK3_MOUSE       346  GAQ-----TMIEHNSTMLESDLGTMVINSEEEEEEEEEEEEEEDGTMKRNATSPQVQRPSFM  400
STK3_RAT         346  GAQ-----TMIEHNSTMLESDLGTMVINSEDE------EEEDGTMKRNATSPQVQRPSFM  394
STK3_DANRE       346  GAQ-----TMIEHGSTMLESNLGTMVINSDDEEE----EEDLGSMRRNPTSQQIQRPSFM  396
STK3_XENLA       345  GAQ-----TMIEHNSTMLESDLGTMVINSDDEEE-----EEEDGTMKRNATSPQGPRPSFM  396
A0A2I3SVA8_PANTR 374  GAQ-----TMIEHNSTMLESDLGTMVINSEDE------EEEDGTMKRNATSPQVQRPSFM  422
HIPPO_DROME      408  AASPLDMPAVDSGTMVELESNLGTMVINSDSDDSTTAKNNDD-----QKPRNRYRPQFL  461
```

F

MST2wt   MST2delEDG

DOX washout (hours) 3 6 9 12   3 6 9 12

FFSS_MST2 →

endogenous MST2 →

G

MST2 wt
MST2 delEDG

FFSS_MST2/MST2

washout (hours)

Figure 3: βTrCP binds to MST2 via a noncanonical degron.

A) βTrCP consensus degron motif. The motif logo was generated with using MEME (http://meme-suite.org/tools/meme) according to Low et al (*5*).

B) MST2 protein sequence. Four tripeptides chemically similar to DSG, the most conserved amino acid residues of the βTrCP canonical degron (Fig. 3A), are present (highlighted) in MST2 sequence: EDS (16-18), ESG (48-50), ESD (359-361) and EDG (375-377).

C) Deletion of EDG (375-377) strongly impacts MST2 interaction with βTrCP. HEK293T cells were transfected with FFSS-tagged MST2 constructs which the tripeptides identified in (B) were deleted or a wild type (wt) construct. Cells were treated with MG132 (5 µM) for 1.5 hour and lysed. Endogenous βTrCP was immunoprecipitated, using a specific antibody and protein A/G-coated beads. Immunoblot for Flag was used to detect FFSS-MST2. βTrCP was detected with the same antibody used for the immunoprecipitation. Total cell extracts were also probed to assess the levels of the different constructs.

D) Structural model of the MST2 EE**EDG**TMKRNA peptide segment bound to βTrCP WD40 domain obtained from 200-ns MD simulation. βTrCP WD40 domain surface is colored by element. MST2 peptide is colored in magenta. MST2 residues are indicated by dashed lines and β-TrCP residues are indicated in boxes. The right image shows a sliced view of the structure.

E) Interspecies alignment of MST2 protein sequence showing that the degron sequence EDGTMKRNA is conserved from Zebra Fish to Humans. DANRE = *Danio rerio* (zebrafish); XENLA = *Xenopus laevis* (African clawed frog); PANTR = *Pan troglodytes* (chimpanzee); and DROME = *Drosophila melanogaster* (fruit fly).

F-G) MST2 with deleted EDG (375-377), MST2- delEDG, is more resistant to degradation than the wild type MST2 (MST2wt).

F) HEK293T cells were transfected with FFSS-tagged MST2wt or MST2-delEDG constructs and treated for 24 h with doxycycline (Dox). After Dox washout, the cells were collected for protein extraction every 3 hours for 12 hours and submitted to immunoblotting with an antibody for MST2. Both ectopic and endogenous MST2 were detected as indicated in the figure.

G) Quantification of MST2wt and MST2-delEDG after Dox washout. The intensity of FFSS-tagged MST2wt and MST2-delEDG bands was normalized to the intensity of the endogenous MST2. Data are show as mean ± SEM.

Protein degradation motifs are commonly conserved across species (*42*). We performed interspecies alignment of MST2 and found that the MST2 EE**EDG**TMKRNA peptide is conserved from zebra fish to Mammalians, but it is not present in the ancestor orthologue Hippo in Drosophila (Fig. 3E).

We then assessed whether deleting the tripeptide EDG would alter MST2 susceptibility to degradation. For this, we cloned MST2-wild type (wt) or MST2- delEDG in a tetracycline inducible expression vector transfected them into HEK293T cells. After Doxycycline (Dox) treatment, both vectors showed robust protein expression (Supplementary Figure 4). A twelve-hour pulse chase-like experiment - stimulation with Dox followed by its removal - confirmed that MST2- delEDG was indeed more resistant to degradation than the wild type protein (Fig. 3F-G). Altogether, our experimental and computational

analyses revealed that MST2 can bind to βTrCP WD40 domain through the noncanonical EDG site, but maintaining structural and chemical characteristics of canonical degrons, such as the β-catenin degron.

Stiff ECM substrates and activation of integrin signaling triggers MST2 degradation

We sought at investigating the signaling pathways activating MST2 degradation in stiffer substrates. First, we confirmed that inhibition of CUL1 neddylation resulted in accumulation of MST2 in MCF10A cells cultured on stiff ECM (Mid and Hi hydrogels) (Fig. 4A). Because stiff microenvironments trigger integrin signaling (*4, 9, 47, 48*), we asked whether modulation of integrin signaling would result in changes of MST2 levels. We mimicked integrin hyperactivation by treating MCF10A cells with $MnCl_2$. Integrins drastically increase affinity to their ligands when bound to Mn^{2+} (*49, 50*). A time-course experiment showed that MST2 decreases in response to Mn^{2+} (Fig. B). Importantly, blocking integrin binding to fibronectin with the RGDS peptide (an integrin binding motif) resulted in increased levels of MST2 (Fig. 4C).

Inhibition of CUL1 neddylation prevented the effects of integrin hyperactivation with Mn^{2+} (Fig. 4D). A 3-hour treatment with Mn^{2+} (Fig. 4E) resulted in 20% reduction of MST2 levels, but a co-treatment with cycloheximide (CHX), a protein synthesis inhibitor, resulted in robust MST2 decrease, similar to the response shown by p21, a well-established Ubiquitin-proteasome system target. Inhibition of CUL1 neddylation or proteasome activity reverted the effects on MST2 content caused by the co-treatment (Fig. 4E). To assess the direct association between integrin activation and MST2 ubiquitination by $SCF^{\beta TrCP}$, we silenced βTrCP in combination with integrin hyperactivation by Mn^{+2}. In fact, forcing decreased levels of βTrCP abrogated the effects of integrin signaling on MST2 (Fig. 4F).

FAs are large protein complexes composed of kinases, FAK and/or ILK, integrins and scaffold proteins that mechanically and biochemically link the ECM to non-cytoskeleton and cytoskeleton

Figure 4: ECM stiffness and integrin activation triggers MST2 degradation

A) Immunoblots of MCF10A cells cultured on Low and Hi rigidities and treated with CUL1 neddylation inhibitor (MLN4924). After MLN4924 (MLN) treatment, MST2 increases in both degrees of rigidity. Note that the band referring to CUL1 neddylation (indicated by an arrow) disappears in the presence of MLN4924 indicating that the inhibition was efficient. Both βTrCP and SKP1 increase after the treatment. Because β-actin varies as a function of stiffness and α-tubulin increases when CUL1 neddylation is inhibited (*51*), Ponceau staining was used to show equal gel loading. Fold changes of MST2 levels are shown under MST2 immunoblots.

B) Immunoblots of MCF10A cells treated with $MnCl_2$ (1 mM) for 1, 3 and 6 hours. Src phosphorylation was detected to assess integrin signaling activation. β-actin was use as loading control. Fold changes of MST2 levels and phosphorylation (pSRC/total SRC) are shown under their respective immunoblots.

C) Immunoblots of MCF10A cells cultured on fibronectin (FN1)-coated plastic plates with or without RGDS, a peptide that blocks integrin binding to FN1. β-actin was used a loading control. Fold changes of MST2 levels are shown under MST2 immunoblots.

D) Immunoblots of MCF10A cells pretreated with MLN for 1.5 hours and then treated or not with $MnCl_2$ (1 mM) for 6 hours. β-actin was used as loading control. Note that the band corresponding to neddylated

CUL1 (indicated by an arrow) disappears in the presence of MLN4924 indicating efficient inhibition. Fold changes of MST2 levels are shown under MST2 immunoblots.

E) Immunoblots of MCF10A cells treated with $MnCl_2$ for 3 hours, with the protein synthesis inhibitor cycloheximide (CHX) for 4 hours, with MLN4294 and MG132 for 1.5 hours. The different combinations of treatments are indicated in the figure panel. Note that the band referring to CUL1 neddylation (indicated by an arrow) disappears in the presence of MLN4924 indicating that the inhibition was efficient. MST2 levels robustly decrease in the presence of $MnCl_2$ and CHX, but inhibition of the proteasome or neddylation reverses this effect. Immunoblotting for p21 was used as a positive control to assess efficiency of CHX, MLN4294 and MG132 treatments. β-actin was used as loading control. Fold changes of MST2 levels are shown under MST2 immunoblots.

F) Silencing of βTrCP in MCF10A cells results in MST2 accumulation, despite integrin signaling hyperactivation. Cells were cultured in the presence of siRNA for βTrCP or a non-targeting siRNA (siControl) and treated or not with $MnCl_2$ for 6 hours. Immunoblots confirming that silencing of βTRCP was efficient and resulted in increased MST2 levels even in the presence of $MnCl_2$. β-actin was used as loading control. Fold changes of MST2 levels are shown under MST2 immunoblots.

proteins resulting in a multitude of cellular outputs (*52, 53*). Our hypothesis was that pharmacological perturbation of the main FA kinases (FAK and ILK) in MCF10A cells would impact MST2 proteostasis. Immunoblots for phospho-AKT and phospho-FAK confirmed that ILK and FAK were inhibited. Surprisingly, kinetical studies showed that only ILK pharmacological inhibition resulted in higher amounts of MST2, whereas FAK inhibition did not alter the level of MST2 (Fig. 5A-B). To further verify the involvement of ILK in MST2 proteostasis, we inhibited ILK in conjunction with integrin hyperactivation by Mn^{2+}. ILK inhibition resulted in increased levels of MST2 (Fig. 5C), and decreased MST2- βTrCP binding in MCF10A cells, as assessed by Co-IP of endogenous MST2 (Fig. 5D).

Stiff substrates elevate the activity of growth factor activated pathways in cancer, fibrosis and other diseases (*54, 55*). For example, PIK3-AKT is a major growth promoting pathway that can be amplified by integrin activation. However, pharmacological inhibition PI3K or AKT did not affect MST2 concentration (Suppl. Fig 5A-B).

Because stiff microenvironments activate actomyosin contraction, we pharmacologically inhibited myosin-II with ML-7, a myosin light chain kinase (MLCK) inhibitor, or blebbistatin, an inhibitor of myosin ATPase activity, and measured the level of MST2. We could observe a relatively transient increase in MST2 levels after myosin contraction inhibition by both drugs (Fig. 5E-F), indicating that intracellular tension stored in the actomyosin cytoskeleton plays a role in MST2 proteostasis. This set of experiments revealed that ILK directly links integrin activation to MST2 degradation control in an actomyosin-cytoskeleton manner (Fig. 5G).

Figure 5: Perturbation of ILK and the actomyosin cytoskeleton regulates MST2 levels:

A) Immunoblots of MCF10A cells treated with an Integrin-Linked Kinase (ILK) inhibitor (CPD022) for 1, 3 and 6 hours. AKT phosphorylation was detected to assess the efficiency of ILK inhibition. β-actin was used as loading control. Fold changes of MST2 levels and AKT activation (pAKT/AKT) are shown, respectively, under MST2 immunoblots and pAKT immunoblots.

B) Immunoblot for MST2 of MCF10A cells treated with FAK inhibitor (FAKi 14) for 1, 3 and 6 hours. FAK phosphorylation was detected to monitor its inhibition. β-actin was used as loading control. Fold changes of MST2 levels and FAK phosphorylation (pFAK/FAK) are shown, respectively, under MST2 immunoblots and pFAK immunoblots.

C) Immunoblots of MCF10A cells treated with CPD022 or MLN4294 in combination with $MnCl_2$ for 6 hours. Both inhibition of CUL1 neddylation and ILK reverses the effect of integrin hyperactivation. β-actin was used as loading controls. Note that the band corresponding to CUL1 neddylation (indicated by an arrow) disappears in the presence of MLN4924 indicating that the inhibition was efficient. Fold changes of MST2 levels are shown under MST2 immunoblots.

D) Co-IP revealed that the interaction of endogenous MST2 and βTrCP decreases after ILK inhibition, despite integrin hyperactivation. MCF10A cells treated with DMSO (ct), CPD022, $MnCl_2$ or CPD022 concomitantly with $MnCl_2$ for 6 hours. Co-IP was performed with an MST2 antibody or a control IgG, followed by incubation with protein A/G-coated beads. βTrCP and MST2 were only detected in the MST2 immunoprecipitants. MST2 was detected to confirm the Co-IP. Arrows indicate bands corresponding to IgG that were detected by the secondary antibody. Fold changes of MST2 levels are shown under MST2 immunoblots.

E) Immunoblot for MST2 of MCF10A cells treated with MLCK inhibitor (ML7) for 1, 3 and 6 hours. β-actin was used as loading controls.

F) Immunoblot for MST2 of MCF10A cells treated with a myosin ATPase activity inhibitor (Blebbistatin, 2.5 µM) for 1, 3 and 6 hours. β-actin was used as loading control.

G) Schematic depicting the pathway of MST2 degradation induced by ECM stiffness. In a stiff ECM, hyperactive integrin signaling results in ILK activation and actomyosin contraction leading to ubiquitination of MST2 by $SCF^{βTrCP}$ and consequent MST2 degradation via proteasome 26S. This mechanism influences the level of YAP in the nucleus.

Discussion

Besides playing crucial roles in tissue development and regeneration, Hippo kinases can be considered bona fide tumor suppressors(*56*). Here we uncovered the molecular mechanism by which MST2, one of the core Hippo kinases, degradation is regulated (Fig. 5G). We not only identified that $SCF^{βTrCP}$ ubiquitinates MST2 for proteasomal proteolysis and mapped the βTrCP binding motif (degron) in MST2, but also described a pathway based on integrin and ILK activation that triggers MST2 degradation in breast epithelial cells.

The cell's microenvironment provide active and passive mechanical cues that elicit biochemical signaling through the cytoskeleton and kinase cascades regulating a milieu of cellular processes such as cell proliferation (*57*), invasion (*4*), metabolism (*55*) and differentiation (*58*). A crucial signal is rigidity of the ECM that is initially sensed by integrins that are intracellularly linked to the actomyosin cytoskeleton that in response to stiffness increase cell contractility. Tension built in contractile actomyosin reciprocally activates the assembly of FA (*26, 57*). YAP/TAZ accumulate in the nuclei of epithelial cells, but compelling evidence has shown YAP/TAZ exerts at least part of their mechanoregulatory activity by directly inducing the expression of genes encoding for FA proteins (*59*). Thus, reinforcing a feed-forward activation cycle where a stiff ECM increases YAP/TAZ nuclear activity and YAP/TAZ transcriptional

control activity results in increased FA assembly. We indeed observed that silencing MST2 results in increased nuclear YAP and that is accompanied by increased cell proliferation and spreading, and F-actin alignment to the major cell axis, suggesting that in our experiments YAP is regulating the actin cytoskeleton and FA assembly.

Herewith, a series of biochemical assays, using ectopic protein expression in HEK293T cells, unveiled that SCF$^{\beta TrCP}$ tags MST2 for degradation after binding to a non-canonical degron in MST2. Importantly, by using a human breast epithelial cell line and probing for endogenous proteins, we not only confirmed these findings but also showed that integrin hyperactivation increases SCF$^{\beta TrCP}$ and MST2 binding. We also observed that inhibition of ILK and dissipation of tension built in actomyosin stress fibers caused MST2 accumulation in breast epithelial cells and inhibited SCF$^{\beta TrCP}$ and MST2 binding, indicating that MST2 degradation is directly dependent upon the integrin-ILK axis activation.

There are conflicting data in the literature as to whether the mechanical cues that influence YAP/TAZ localization and activity are mediated by Hippo kinases (*10, 60, 61*). This might be attributed to the use of different cell lines, type of ECM molecule used for coating, hydrogel preparation and degrees of stiffness. Nevertheless, in the present work the outputs in our MST2 silencing assays were different depending on the cellular process assessed: MST2 knock down resulted in increased proliferation in all three degrees of rigidity while cell shape and F-actin arrangement were only perturbed in stiffer substrates indicating that regulation of cell proliferation and cytoskeleton/cell shape might be independently regulated in physiological stiffness and also that YAP/TAZ regulation of FA assembly might occur only in stiffer ECMs. Although our data clearly shows that MST2 level is mechanoresponsive, other complementary factors could act on YAP/TAZ as previously reported (*10, 62*). Also, MST2 can phosphorylate a vast number of substrates (*63*) and play roles nonrelated to its kinase activity ((*64*). Therefore, part of the phenotypic changes due to MST2 modulation might not relate to regulation of YAP/TAZ but by action of other effectors. In this context, there is evidence showing that cytoskeletal disruption activated MST as well as JNK1 in an MST-dependent manner (*65*). Inhibition of actin cytoskeleton with tat-C3 resulted in higher MST activity (*65*).

The βTrCP degron in MST2 does not contain serine amino-acid residues as the canonical degron, raising the question of how the MST2 and βTrCP interaction is molecularly regulated. Nevertheless, the chemical properties of the MST2 tripeptide EDG are similar to the conserved DpSG in the canonical degrons and

the *in silico* studies showed that an 11-mer MST2 peptide that includes the EDG tripeptide stably binds to the βTrCP, despite the fact that the segment does not present a C-terminus negative amino acid that would substitute for the second phosphorylated serine in the degron. It remains, thus, to be elucidated whether phosphorylation in other regions of MST2 or other posttranslational modifications would influence MST2 and βTrCP binding. This will be important to pinpoint which molecule is linking integrin-ILK activation to MST2 and βTrCP. There are, indeed, other instances in the literature of degrons that do not follow the DpSGXX(X)pS rule. For example, one of the WEE1 degrons (SWEEEGFGSS) (*43*) and the CDH1 degron (SSPDDGNDVS) (*44*) do not bear the first serine found in the canonical degron. And, there is at least one report of a degron (the CDC25B degron; TEEDDGFVDI) analogous to the MST2 degron we identified, that does not contain any serine (*45*).

Frequently, tumors are stiffer than normal tissues eliciting integrin hyperactivation (*4, 8, 48*). We propose that engagement of integrin-ILK in focal adhesion arrays in stiffer ECM, could result in decreased levels of MST2 due to enhanced protein degradation and this process could cooperate with other aberrant signals in reinforcing malignant phenotypes in tumor microenvironments. We also suggest that the pathway, integrin-ILK-MST2, might consist in a novel molecular target for treating breast tumors, and other types of cancers, that display increased density mammograms and low levels of Hippo activity. Based on the degron we identified, a possible strategy would be developing peptides or small molecules that would block MST2 degradation by blunting the anchoring of βTrCP to MST2. MST2 bears a noncanonical βTrCP degron, hence precisely targeting the βTrCP-MST2 interaction interface could reduce the chance of undesirably off-targeting SCF^{βTrCP} substrates with opposing biological functions. For instance, although YAP and TAZ are substrates of SCF^{βTrCP} their degrons differ considerably from the one we described for MST2. Finally, we believe that our work unveiled a detailed mechanism regulating MST2 levels and this is expected to have wide ramifications for cell tensional homeostasis determination and provide opportunity for therapeutic strategies.

Materials and Methods

Cell lines and cell culture media

Non tumoral breast epithelial cell lines MCF10A (Gently donated by Mina J. Bissell -Lawrence Berkeley National Laboratory) and MCF12A (Gently donated by Carlos Frederico Martins Menck, from Instituto de Ciências Biomédicas-USP) were cultured in Dulbecco Modified Eagle's/ HAM F-12 medium (1:1,

DMEM/F-12, Gibco, #12-500-062), supplemented with 5% horse serum (Gibco # 16050-122), Insulin (10 µg/mL, Sigma- Aldrich, #I5500), Hydrocortisone (10 µg/mL, Sigma-Aldrich # H4001), Choleric toxin (1 µg/mL, Sigma-Aldrich, #C8052) and Epidermal Growth Factor (20 ng/mL, Sigma-Aldrich, #E4127).

Breast cancer epithelial cell lines MCF7 and MDA-MB231 (both gently donated by Mina J. Bissell - Lawrence Berkeley National Laboratory), and MDA-MB468 (purchased from American Type Culture Collection, ATCC, #HTB-132) cell lines were cultured in DMEM high glucose (Gibco, #12100046) supplemented with 10% of fetal bovine serum (FBS, Gibco #12657-029); BT474 and SKBR3 (both purchased from ATCC #HTB-20 and #HTB-30, respectively) were cultured in RPMI (Gibco #11875093) supplemented with 10% FBS, Hyclone non-essential amino acids (0.1 mM, Thermo #SH3023801), L-glutamine (2 mM, Gibco, #25030081) and sodium pyruvate (1 mM, Gibco, #11360070).

Human Embryonic Kidney (HEK) 293T cells (purchased from ATCC, #CRL-11268) were cultured in Dulbecco's Modified Eagle's Medium (DMEM High Glucose, Thermo Fisher) supplemented with 10% Fetal Bovine Serum (Gibco), Hyclone non-essential amino acids (0.1 mM, Thermo), L-Glutamine (2 mM, Gibco) and sodium pyruvate (1 mM, Gibco).

Cell culture on polyacrylamide hydrogels

The polyacrylamide hydrogels with tunable mechanical properties were made in three elastic modulus levels: 0.48 kPa (3% acrylamide, 0.06% bis-acrylamide), 4.47 kPa (5% acrylamide, 0.15% bis-acrylamide) and 40.40 kPa (8% acrylamide, 0.48% bis acrylamide), as previously described (*29*). The hydrogels were coated with fibronectin-1 (10 µg/ml) by using the crosslinker Sulfo-SANPAH (Pierce Biotechnology, # A35395). MCF10A cells were seeded at a density of 3×10^4 cells per cm^2 in complete medium and kept in culture for 72 hours.

cDNA constructs, plasmids and transfection

All plasmids for protein expression used in this study are listed in Table S1. MST2 ORF (*STK3,* accession number NM_006281.4) was amplified from MCF10A human breast cells using specific primers (listed in Table S2) with Q5 High-Fidelity 2X Master Mix (New England BioLabs, # M0492), following the manufacturer's instructions. The MST2 ORF was inserted in the pCDNA3.1_Flag2x _Strep2x vector using StickyEnd ligase (New England Biolabs, # M0370L). The plasmids pCDNA3.1_Flag_ β-TrCP, pCDNA3.1_Flag_ βΔFbox and pCDNA3.1_myc_ubiquitin were described previously (*66, 67*). To obtain the MST2 constructs with the tripeptide deletions, we used the pCDNA3.1_Flag2x _Strep2x _MST2

plasmid as a template, specific primers (MST2_BAMHI_Fw and MST2_XHOI_Rv, listed in Table S2) and the site directed deletions and amplification were performed using Pfu turbo DNA polymerase (Agilent, #600250) following manufacturers protocol. All constructs sequences were confirmed by Sanger sequencing (GeneWitz, NJ) and were transfected in HEK293T cells using the Lipofectamine 3000 reagent (Thermo Fischer # L3000008).

To generate inducible vectors for MST2 wild type and del EDG we amplified the Flag2x _Strep2x fused to the MST2 constructs (wild type and del_EDG) with Q5 High-Fidelity Polymerase (New England BioLabs, # M0492) using specific primers (EcoRI_Flag_Fw and MST2_AgeI_Rv, listed in table S2) and inserted the 2xFlag_2xStrep_MST2 constructs (wild type or del_EDG) in the inducible vector pLVX (Clontech, #631847).

Gene silencing by siRNA

The following pre-designed and validated duplexes siRNA oligos (Ambion # 4392420): *STK3* (s13567), *βTrCP* (s17109), and non-targeting (siRNA #1, AM4635) were used. Oligo duplexes were transfected using Lipofectamine RNAiMax (ThermoFisher Scientific # 13778030) according to the manufacturer's instructions, using 200 nM of each oligo. The durations of transfection are indicated in the figures and/or figure legends.

Treatment with inhibitors and blocking peptides

Inhibitions of Cullin (CUL1) neddylation or proteasome activity were achieved by treating cells with 2.5 µM MLN4924 (Cayman Biochemicals # 15217) or 5 µM MG132 (Peptides International # IZL-3175-v) for the durations indicated in the figure legends. To assess the protein stability of MST2, we inhibited global protein synthesis with cycloheximide (CHX, 100 µg/mL, Sigma Aldrich # C4859).

Integrin activation was induced in MCF10A cells with 1 mM $MnCl_2$ (Sigma Aldrich # M1787) treatment for the durations indicated in the figure legends (*68*). To screen the pathways mediating MST2 degradation, MCF10A cells were treated for 1, 3 and 6 hours with the inhibitors for ILK (1 µM CPD022, Calbiochem, #407331), FAK (5 µM FAK inhibitor 14, Tocris # 3414), PI3K (30 µM LY294002, Gibco # PHZ1144), AKT (20 µM MK2206, Cayman Biochemicals #1032350-13-2), myosin ATPase activity inhibitor (5 µM Blebbistatin, Tocris #1852) and myosin light chain kinase (MLCK) inhibitor (5 µM ML-7, Tocris #4310).

For blocking integrin binding to the ECM, cell-culture plates were coated with fibronectin-1 (FN1) (10 µg/mL, Corning # 356008), following the manufacturer instructions. MCF10A cells were seeded on cell-culture plates coated with FN1 with concomitant addition of RGDS peptide (250 µM, Chinese peptides # RGDP-010) for 13 hours.

MST2 stability assay using an inducible protein expression system

HEK293T cells were transiently transfected using Lipofectamine 3000 (ThermoFisher Scientific) based on the manufacturer's recommendation with wild type MST2 or MST2_delEDG subcloned in pLVX_TetOne-Puro (Clontech # 631849). Cells were cultured in presence of Doxycycline (Dox, 500 ng/mL Sigma Aldrich, #D9891) to induce MST2 and MST2_delEDG expression. After 24 hours, Dox was washed with PBS 3 times for 30 minutes and complete medium was added. Cells were collected for protein extraction 3, 6, 9 and 12 hours after Dox removal.

Protein extraction, immunoprecipitation, SDS-PAGE and Immunoblot

For total cell lysate protein extraction, samples were lysed using modified RIPA buffer (Pierce) supplemented with protease (Sigma-Aldrich) and phosphatase inhibitors (Sigma-Aldrich). Protein concentration was determined using the DC Protein Assay kit (BioRad, #5000111). Protein lysate was resuspended at a 0.5 µg/µL concentration in Laemmli sample buffer (0.1% 2-Mercaptoethanol, 0.005% Bromophenol blue, 10% Glycerol, 2% SDS, 63 mM Tris-HCl), heated at 95°C for 5 minutes and kept at -20 °C.

For the Co-immunoprecipitation (Co-IP) cells were lysed in immunoprecitation lysis buffer (25 mM Tris pH 8.0, 150 mM NaCl, 10% glycerol, 1 mM EDTA, 1 mM EGTA, 1 mM 1,4-Dithiothreitol (DTT) and 0.1% NP-40) containing protease (Sigma-Aldrich) and phosphatase inhibitors (Sigma-Aldrich). The insoluble fraction was removed by centrifugation (20,000 x g x 15 min at 4°C). Protein concentration was determined using the DC Protein Assay kit (BioRad #5000111). For both MST2 and βTrCP immunoprecipitation, we incubated 2 mg of protein extracts with specific antibodies (listed in Table S3) overnight at 4°C. Subsequently, we added Protein A/G magnetic beads (Pierce, # 88802), incubated at 4°C for 2 hours, and proceeded with extensive washing using the lysis buffer. After the final wash, the beads with the immunoprecipitant were resuspended with 1x Laemmli sample buffer, heated at 95°C for 5 minutes and kept at -20 °C.

The flag-tag immunoprecipitations were carried out using FLAG-M2 coated agarose beads (Sigma-Aldrich #F2426) for 2 hours at 4 °C. The beads were then extensively washed in lysis buffer and elution was carried out with 50 µL of 3×FLAG peptide (Sigma-Aldrich #F4799), the supernatant was separated from the beads by centrifugation, resuspended in the same volume of 2x Laemmli sample buffer, heated at 95°C for 5 minutes and kept at -20 °C.

Total protein extracts or immunoprecipitants were resolved in SDS-PAGE followed by immunoblot. In brief, 10 µL (5 µg) of protein sample and 20 µL of IP samples were heated at 95°C for 5 minutes and loaded into 10% tris-glycine polyacrylamide gels. Resolved proteins were transferred to PVDF membrane (0.45 µm, Millipore, #05317) followed by 1-hour incubation in a blocking buffer (TBS, 0.5% Tween-20 with 1% BSA). Membranes were incubated in buffer (TBS, 0.5% Tween-20) containing primary antibodies (listed in Table S3) overnight., and finally incubated with horseradish peroxidase (HRP) for 1 hour at room temperature in the blocking buffer. HRP was detected by Pierce SuperSignal detection kit (Thermo-Fisher Scientific, #A45916) and the chemiluminescence signal was captured with a ChemiDoc MP Imaging System (BioRad).

RNA extraction, reverse transcription, and quantitative polymerase chain reaction (RT-qPCR)

Total RNA was extracted using RNeasy kit (Qiagen #74106) following the manufacturer's guidelines. The RNA 260/280 and 260/230 ratio and the concentration were measured with a Nanodrop (Thermo) UV-spectrophotometer. The cDNA was generated using 1 µg of total RNA using the kit First-Strand cDNA Synthesis (Thermo Scientific # 18080051) using oligoDT and random hexamer primers, following the manufacturer's instructions. For the qPCR, 10-µL reactions were performed in triplicates with 25 ng of cDNA using 2x SYBR Green PCR Master Mix (Applied Biosystems # 4364346) and specific primers (listed in table 2) in MicroAmp Optical 96-Well Reaction plates (Applied Biossystems #N8010560). The reactions were performed in the AB-9500 system (Applied Biosystems) real-time thermocycler. GAPDH was used as an endogenous control and mRNA fold-changes were calculated according to the Pfaffl method (*69*).

5-ethynyl-2'-deoxyuridine (EdU) incorporation assay.

The Click-iT® EdU kit (Life Technologies, # C10639) was used for measuring the percentage of cells in S-phase in different conditions cultured on the hydrogels. Only DNA-replicating cells incorporate the base analog EdU. Cells were incubated with 20 µM EdU for 1 hour, and fixed with 4% PFA (Electron Microscopy Sciences, #15714) in PBS. EdU incorporation was detected following the manufacturer's

instructions as previously described (*70*). The cells were then counterstained with 4'6-diamidino-2-phenylindole (DAPI, 0.5ug/mL, Sigma-Aldrich #D9542) for 10 minutes, the slides/ coverslips were mounted with Prolong® Diamond (Thermo Scientific # P36965) and analyzed with a Leica DMi8 wide-field microscope (Leica Microsystems) equipped with a 40x objective.

Immunofluorescence and phalloidin staining

The cells were fixed with 4% paraformaldehyde (PFA) (Electron Microscopy Sciences) in PBS. Residual PFA reactivity was quenched with PBS–25 mM glycine and cells were permeabilized with 0.5% Triton X-100 in PBS for 30 minutes. The cells were then incubated in blocking buffer 1% Bovine Serum Albumin (MP Biomedicals #160069) and 5% goat-serum in PBS (Gibco # 6210-064) for 1 hour at room temperature, followed by overnight incubation with primary antibody (listed in table S3) in blocking buffer. Cells were then incubated in blocking buffer with Alexa 488 or Alexa 555 conjugated secondary antibodies (listed in table S3) for 45 minutes, counterstained with DAPI for 10 minutes and mounted with Prolong® Diamond (Thermo Scientific #, P36970). and analyzed with a Leica DMi8 wide field microscope. Image stacks were captured with a 63X/1.4 NA oil objective (Leica) for YAP detection. We used the optimized step intervals along the Z-axis to capture 3D images and the deconvolution of the acquired images was performed in the Leica Application Suite-X software using the tool blind deblur with 10 iterations (Leica Microsystems).

Superresolution fluorescence microscopy images of cells stained with phalloidin-Alexa 488 (1:50, Thermo Scientific #A12379) and vinculin antibody were obtained of with an LSM-880 confocal microscope equipped with the AiryScan system (Carl Zeiss) using a 63x/1.4 NA oil objective lens and optimized acquisition settings.

In vitro **ubiquitination assay**

We co-transfected HEK293T cells with plasmids containing myc-tagged ubiquitin, 2x Flag 2x Strep MST2 and Flag βTrCP (listed in table1) for 24 hours. Three hours prior to collection, we added 5 μM MG132, a proteasome inhibitor. Protein extraction was performed under denaturing conditions (50 mM TRIS, 150 mM NaCl, 1 mM EDTA, 1% SDS) containing protease (Sigma-Aldrich #11836145001) and phosphatase inhibitors (Sigma-Aldrich #P5726-5ML) and the deubiquitinase inhibitor N-ethylmaleimide (1 μM, Sigma-Aldrich, #E3876). Samples were boiled at 98 °C for 10 minutes and sonicated 3 times for 10 seconds on ice and the insoluble fraction was removed by centrifugation. Next, 2xFlag 2xStrep MST2

was purified by affinity, using Strep-Tactin Superflow beads (IBA # 2-1206-025) in immunoprecipitation lysis buffer (50 mM TRIS, 150 mM NaCl, 2.5 mM EDTA, 1% Glycerol, 0,1% NP40) for 2 hours at 4°C, beads were extensively washed in lysis buffer and elution was carried out with Strep-Tactin Elution Buffer containing Desthiobiotin (IBA # 2-1000-025).

Image analysis

Fluorescence microscope and gel image quantification were performed using the ImageJ software (2.0.0-rc-68/1.52e; Java 1.8.0_66 64-bit).

For the relative quantification of the protein levels of the immunoblot experiments, and each band integrated optical density was determined using the gel analyzer application in Image. The endogenous control (β-actin, α-tubulin or ponceau staining, depending on the experiment) was also determined and the fold change was determined as the ratio between target protein level and endogenous control. For phosphorylated proteins, we used the ratio phosphorylated protein/ total protein/endogenous control.

For quantification of YAP staining, we used a customized ImageJ macro (available upon request). In brief, the DAPI channel was converted to a binary mask to delimit the nuclei and the analyze particles tool in ImageJ tool was used to generate selections (outlines) of individual nuclei. The imageJ tool *enlarge* was used to produce an outer and an inner ring by concentrically expanding or shrinking the nuclear. To determine the intensity of YAP level in the cytoplasm, the YAP intensity inside the nuclear outline was subtracted from the YAP staining in the outer ring. Finally, to obtain the YAP cytoplasm to nucleus ratio for each cell, the intensity of YAP signal inside the inner ring was divided by the intensity of YAP in the cytoplasm of the same cell.

To quantify the EdU assay, the number of total and Edu-stained cells was obtained by using the *cell counter* application in ImageJ and the percentages of EdU-positive cells were determined.

For quantification of F-actin directionality, we used the *directionality* plugin (https://imagej.net/Directionality) in ImageJ was used. For this, superresolution images of individual cells stained with phalloidin were aligned in respect to the cell's major axis.

All imaging acquisition parameters were kept constant for each experiment.

Statistics

Statistical analyses were performed using GraphPad Prism 7.0. The number of samples analyzed and the number of times that the experiments were performed are indicated in the respective figure legends. All the data were submitted to a normality test. Nonparametric data were submitted to Kruskal-Wallis test, while parametric data were submitted to ANOVA followed by Dunnett test. Differences were considered significant when the p-value was less than 0.05. All graphs were generated with GraphPad and the data are presented as mean $\pm$ SEM.

Molecular modeling and dynamic simulation and protein sequence alignment

Human MST2 secondary structure was predicted using the PsiPred server (http://bioinf.cs.ucl.ac.uk/psipred/) (*71*). An initial model of the MST2 segment that comprises EDG tripeptide (EEEDGTMKRNA) bound to β-TrCP WD40 domain was built using the main chain atoms coordinates of β-catenin peptide in a crystallographic complex with β-TrCP (PDB ID: 1P22) and changing the sidechains to match MST2 sequence. This process was performed using YASARA software (http://www.yasara.org/). Only β-TrCP WD40 domain was used in modeling and simulations.

The molecular dynamic (MD) simulations were performed using YASARA with AMBER14 forcefield. For all simulations, the structures were solvated in a 10 Å cubic cell around all atoms, neutralized and submitted to energy minimization. The production runs were performed at 298 K and pH 7.4 with 2*2.5 fs timestep. All bonds and angles involving hydrogens were constrained using LINCS algorithm and an 8 Å cutoff was used for Van der Waals and Coulomb force calculation and long-range Coulomb interaction were calculated using the Particle Mesh Ewald algorithm. Simulation snapshots were saved each 250 ps. The production run of MST2 11-mer peptide EEEDGTMKRNA was performed for 200 ns and the last frame was submitted to energy minimization in YASARA and evaluated.

For MD simulations analysis, simulation snapshots were aligned using the βTrCP WD40 domain first snapshot as reference in VMD (*72*). RMSD (root-mean-square deviation) and RMSF (root-mean-square fluctuation) were calculated for the MST2 peptide using the R Bio3D package (*73*).

For alignment of MST2 sequence from different species, we used the multi sequence alignment tool from the software Jalview 2.11.1.3 (https://www.jalview.org/).

Acknowledgements: This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) Young Investigator Award (14/10492-0), by a FAPESP grant (2019/26767-2) by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq- Universal 444597/2014-0) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Finance Code 001). MP is an investigator with the Howard Hughes Medical Institute and his lab is partially funded by grants from the National Institute of Health (R01-CA76584 and R35-GM136250). APZPF is a FAPESP Postdoctoral fellowship awardee (14/25832-1 and 17/18641-3). AMRS is funded by CAPES PhD-scholarship (PROEX), PFR is a FAPESP PhD-scholarship recipient (17/18067-5), MCSB is a FAPESP PhD-scholarship recipient (2017/25437-3) and ACM is a CNPq PhD-scholarship recipient (14668/2019-9). The authors would like to thank Professor Ricardo Giordano [Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo (DBQ-IQUSP)] for the RGDS peptide donation and Dr Gergely Róna (Department of Biochemistry and Molecular Pharmacology, New York University School of Medicine), Dr Rebeka Tomasin (DBQ-IQUSP), Dr. Deborah Schechtman (DBQ-IQUSP), Dr. Marcelo Damário Gomes (Faculdade de Medicina de Ribeirão Preto – Universidade de São Paulo) and Dr. Hernandes F. Carvalho (UNICAMP) for all the fruitful discussions and valuable advices during the progress of this project. The authors would also like to thank Jeffrey Estrada, Celia Ludio Braga, Maria Luiza Baldini and Izaura Nobuko Toma for technical assistance.

Author contribution: APZPF planned, performed, analyzed most experiments and co-wrote the manuscript. AMRS performed all the RT-qPCRs and immunoblottings in figure 5. PRF manufactured all hydrogels. ACM generated cDNA constructs and performed the protein stability assay. MCSB cultured the breast cell lines used in the experiments shown in figure 2A. HVRF and PSLO performed the molecular dynamics studies. MP co-supervised part of the study. AB-C planned, analyzed data, performed fluorescence microscopy, supervised the study and wrote the manuscript.

Competing financial interests

M.P. is a consultant for and has financial interests in Coho Therapeutics, CullGen, Kymera Therapeutics, and SEED Therapeutics. M.P. is a cofounder of Coho Therapeutics, is on the SAB of CullGen and Kymera Therapeutics, and is a consultant for Santi Therapeutics. The other authors declare no competing interests.

Supplementary Figures

Supplementary Figure 1: Validation of STK3 knock down in MCF10A cells and fluorescence microscopy images accompanying figure 1:
A) Immunoblot for MST2 confirming STK3 silencing in MCF10A cells. MCF10A cells were cultured on regular plastic plates and treated with a control siRNA or an siRNA for STK3 for 24 hours.
B) This panel refers to Figure 1B and displays images of YAP immunofluorescence (red) along with their corresponding nuclear staining (DAPI = blue).
C) This panel refers to the quantification shown in the bar graph in Figure 1F and displays fluorescence microscopy images of EdU staining (red) along with their corresponding nuclear staining (DAPI = blue). Scale bars = B, 10 µm and C, 20 µm.

Supplementary Figure 2: Root-mean-square deviation (RMSD) plot:

A) MST2 secondary <u>structure</u> prediction. The prediction was performed in the PsiPred server. The MST2 11-mer peptide EEEDGTMKRNA is localized within a structurally disordered – protein binding region.

B) 200 ns MD simulation of MST2 11-mer peptide EEEDGTMKRNA in complex with β-TrCP WD40 domain. MST2 peptide snapshots of each 10 ns are shown as cartoon representation and colored by timestep. MST2 K380 is shown in sticks representing the first (red) and the last (blue) simulation frame.

C) Backbone RMSD (in Å) of MST2 peptide during the MD simulation.

D) RMSF calculated from all atoms of MST2 segment during the MD simulation.

E) Structural alignment between β-catenin peptide crystallographic structure (PDB ID: 1P22) and MST2 peptide model bound to βTrCP. β-catenin peptide is presented in yellow and MST2 in magenta. Important residue for MST2 and β-TrCP interaction are presented in sticks.

A

```
MEQPPAPKSKLKKLSEDSLTKQPEEVFDVLEKLGEGSYGSVFKAIHKESGQVVAIKQVPVESDLQEIIKEISIMQQCDSP
YVVKYYGSYFKNTDLWIVMEYCGAGSVSDIIRLRNKTLIEDEIATILKSTLKGLEYLHFMRKIHRDIKAGNILLNTEGHA
KLADFGVAGQLTDTMAKRNTVIGTPFWMAPEVIQEIGYNCVADIWSLGITSIEMAEGKPPYADIHPMRAIFMIPTNPPPT
FRKPELWSDDFTDFVKKCLVKNPEQRATATQLLQHPFIKNAKPVSILRDLITEAMEIKAKRHEEQQRELEEEENSDEDE
LDSHTMVKTSVESVGTMRATSTMSEGAQTMIEHNSTMLESDLGTMKRNATSPQVQRPSFMDYFDKQ
DFKNKSHENCNQNMHEPFPMSKNVFPDNWKVPQDGDFDFLKNLSLEELQMRLKALDPMMEREIEELRQRYTAKRQPILDA
MDAKKRRQQNF
```

Supplementary Figure 3: The MST2 K380 residue is predicted to be ubiquitinated:

A) MST2 ubiquitination prediction. The prediction was performed using the UbPred server (http://www.ubpred.org/index.html). The colored boxes indicate the result confidence. Green, blue and red represents low, medium and high confidence, respectively. Among the MST2 lysine residues, K380 (black arrow) has the highest confidence value (0.96).

Supplementary Figure 4. Validation of doxycycline-inducible expression of MST2wt and MST2delEDG:

A) Immunoblot using an antibody for MST2 of HEK293T cells transfected with FFSS-tagged MST2wt or MST2-delEDG constructs and treated for 24 h or 36 hours with dox. Both constructs showed robust and comparable protein expression after induction.

Supplementary Figure 5. Inhibition of PI3K and AKT does not alter MST2 levels

A) Immunoblots of MCF10A cells treated with PI3 Kinase (PI3K) inhibitor (LY294002) for 1, 3 and 6 hours. AKT phosphorylation was detected to assess PI3K inactivation. β-actin was used as loading control. Fold changes of MST2 levels are shown under immunoblots for MST2

B) Immunoblot for MST2 of lysates from MCF10A cells treated with AKT inhibitor (MK2206) for 1, 3 and 6 hours. AKT phosphorylation was detected to assess its inhibition. β-actin was used as loading control. Fold changes of MST2 levels are shown under immunoblots for MST2.

Supplementary tables:

Table1. List of Plasmids used in this study

Plasmid	Backbone	cDNA cloned	Reference
pCDNA3.1_Flag_β-TrCP	pCDNA3.1_Flag	βTrCP	Guardavaccaro et al 2003, (*74*)
pCDNA3.1_Flag_βΔFbox	pCDNA3.1_Flag	βTrCP with deletion of Fbox binding motif	Guardavaccaro et al 2003, (*74*)
pCDNA3.1_FFSS-MST2	pCDNA3.1_Flag2xStrep2x	MST2	This article
pCDNA3.1_FFSS-MST2[del16-18]	pCDNA3.1_Flag2xStrep2x	MST2 with deletion of EDS (amino acids 16-18)	This article

pCDNA3.1_FFSS-MST2del48-50	pCDNA3.1_Flag2xStrep2x	MST2 with deletion of ESG (amino acids 48-50)	This article
pCDNA3.1_FFSS-MST2del359-361	pCDNA3.1_Flag2xStrep2x	MST2 with deletion of ESD (amino acids 359-361)	This article
pCDNA3.1_FFSS-MST2del375-377	pCDNA3.1_Flag2xStrep2x	MST2 with deletion of EDG (amino acids 375-377)	This article
pCDNA3.1_myc_Ub	pCDNA3.1_myc	Ubiquitin	Marzio et al 2019, (*66*)
pLVX_FFSS_MST2	pLVX-TetOne-Puro	MST2	This article
pLVX_FFSS_MST2delEDG	pLVX-TetOne-Puro	MST2 with the deletion of EDG (amino acids 375-377)	This article

Table 2. List of primers used in this study

Primer name		type of use	sequence
MST2_BAMHI	Fw	Cloning in pCDNA3.1	CGCGGATCCGAGCAGCCGCCGGCGCCT
MST2_XHOI	Rv	Cloning in pCDNA3.1	CCGCTCGAGTCAAAAGTTTTGCTGCCTTCTTTTCTTTGC
MST2_del16-18	Fw	Deletion	GAGTAAACTAAAAAAGCTGAGTTTGACTAAGCAGCCTGAAGAAG
MST2_del16-18	Rv	Deletion	CTTCTTCAGGCTGCTTAGTCAAACTCAGCTTTTTTAGTTTACTC
MST2_del48-50	Fw	Deletion	AAGTGTATTTAAAGCAATACACAAGCAAGTTGTCGCAATTAAACAAGTAC

MST2_del48-50	Rv	Deletion	GTACTTGTTTAATTGCGACAACTTGCTTGTGTATTGCTTTAAATACACTT
MST2_del359-361	Fw	Deletion	GAACATAATAGCACGATGTTGTTGGGGACCATGGTGATAAAC
MST2_del359-361	Rv	Deletion	GTTTATCACCATGGTCCCCAACAACATCGTGCTATTATGTTC
MST2_del375-377	Fw	Deletion	GATAAACAGTGAGGATGAGGAAGAAACTATGAAAAGAAATGCAACC
MST2_del375-377	Rv	Deletion	GGTTGCATTTCTTTTCATAGTTTCTTCCTCATCCTCACTGTTTATC
ECORI_Flag	Fw	Cloning	TCCATGAATTCATGGACTACAAGGACGACGAT
MST2_AgeI	Rv	Cloning	TCCATACCGGTTCAAAAGTTTTGCTGCCTT
MST2	Fw	qPCR	GTTGGTCCGAGTTCCCAGAG
MST2	Rv	qPCR	ACAGAGAGAGGGACCTGGTC
GAPDH	Fw	qPCR	AGGGCTGCTTTTAACTCTGGT
GAPDH	Rv	qPCR	CCCCACTTGATTTTGGAGGGA

Table 3. List of antibodies

Target	Host	Dilution	Company (catalog)
MST2	Rabbit	1:1000 (IB) 1:400 (IF) 1.5 µg (IP)	Cell Signaling Technology (3952S)
MST2	Mouse	1:1000 (IB)	Proteintech (66637-1-Ig)
β-TrCP	Rabbit	1:1000 (IB) 1.5 µg (IP)	Cell Signaling Technology (4394S)
YAP	Rabbit	1:1000 (IB) 1:400 (IF)	Cell Signaling Technology (4912S)
pYAP^{S127}	Rabbit	1:1000 (IB)	Cell Signaling Technology (13008s)
LATS1	Rabbit	1:1000 (IB)	Cell Signaling Technology (9153S)

pLATS1/2	Rabbit	1:1000 (IB)	Cell Signaling Technology (8654S)
SRC	Rabbit	1:1000 (IB)	Cell Signaling Technology (2109S)
pSRC	Rabbit	1:1000 (IB)	Cell Signaling Technology (6943S)
FAK	Rabbit	1:1000 (IB)	Cell Signaling Technology (3285S)
pFAK^{Y861}	Rabbit	1:1000 (IB)	Thermo Scientific (44-626G)
CUL1	Rabbit	1:1000 (IB)	Thermo Scientific (71-8700)
SKP1	Rabbit	1:5000 (IB)	Pagano lab (75) (N/A)
Flag	Mouse	1:4000 (IB)	Sigma-Aldrich (4503365)
Myc-Tag	Mouse	1:1000 (IB	Cell Signaling Technology (2276S)
PI3K	Rabbit	1:1000 (IB)	Cell Signaling Technology (4292S)
pPI3K	Rabbit	1:500 (IB)	Cell Signaling Technology (4228S)
AKT	Rabbit	1:1000 (IB)	Cell Signaling Technology (9272S)
pAKT	Rabbit	1:1000 (IB)	Cell Signaling Technology (4060S)
β-actin	Mouse	1:10000 (IB)	Sigma-Aldrich (A5441)
α-tubulin	Mouse	1:10000 (IB)	Thermo Scientific (2036)
Vinculin	Mouse	1:500 (IF)	Thermo Scientific (1209)
Mouse (secondary)	Goat (Alexa 555)	1:500 (IF)	Thermo Scientific (A28180)
Rabbit (secondary)	Goat (Alexa 555)	1:500 (IF)	Thermo Scientific (4060S)
Mouse (secondary)	Goat (HRP)	1:10000 (IB)	Thermo Scientific (31430)
Rabbit (secondary)	Goat (HRP)	1:10000 (IB)	Thermo Scientfic (32460)
Mouse light-chain (secondary)	Goat (HRP)	1:10000 (IB)	Jackson ImmunoResearch (115-035-174)
Rabbit light-chain (secondary)	Goat (HRP)	1:10000 (IB)	Jackson ImmunoResearch (211-032-171)

(IB) Immunoblot, (IF) Immunofluorescence, (IP) Immunoprecipitation.

References

1.   A. P. Z. P. Fiore, P. F. Ribeiro, A. Bruni-Cardoso, Sleeping Beauty and the Microenvironment Enchantment: Microenvironmental Regulation of the Proliferation-Quiescence Decision in Normal Tissues and in Cancer Development. *Front Cell Dev Biol* **6**, 59 (2018).
2.   J. D. Humphrey, E. R. Dufresne, M. A. Schwartz, Mechanotransduction and extracellular matrix homeostasis. *Nat Rev Mol Cell Biol* **15**, 802-812 (2014).
3.   S. Sengupta, K. E. Rothenberg, H. Li, B. D. Hoffman, N. Bursac, Altering integrin engagement regulates membrane localization of K. *J Cell Sci* **132**, (2019).
4.   K. R. Levental *et al.*, Matrix crosslinking forces tumor progression by enhancing integrin signaling. *Cell* **139**, 891-906 (2009).
5.   H. T. Nia, L. L. Munn, R. K. Jain, Physical traits of cancer. *Science* **370**, (2020).
6.   P. Lu, V. M. Weaver, Z. Werb, The extracellular matrix: a dynamic niche in cancer progression. *J Cell Biol* **196**, 395-406 (2012).
7.   G. Charras, E. Sahai, Physical influences of the extracellular environment on cell migration. *Nat Rev Mol Cell Biol* **15**, 813-824 (2014).
8.   M. J. Paszek *et al.*, Tensional homeostasis and the malignant phenotype. *Cancer Cell* **8**, 241-254 (2005).
9.   O. Chaudhuri *et al.*, Extracellular matrix stiffness and composition jointly regulate the induction of malignant phenotypes in mammary epithelium. *Nat Mater* **13**, 970-978 (2014).
10.   S. Dupont *et al.*, Role of YAP/TAZ in mechanotransduction. *Nature* **474**, 179-183 (2011).
11.   A. Das, R. S. Fischer, D. Pan, C. M. Waterman, YAP Nuclear Localization in the Absence of Cell-Cell Contact Is Mediated by a Filamentous Actin-dependent, Myosin II- and Phospho-YAP-independent Pathway during Extracellular Matrix Mechanosensing. *J Biol Chem* **291**, 6096-6110 (2016).
12.   F. Liu *et al.*, Mechanosignaling through YAP and TAZ drives fibroblast activation and fibrosis. *Am J Physiol Lung Cell Mol Physiol* **308**, L344-357 (2015).
13.   S. G. Szeto *et al.*, YAP/TAZ Are Mechanoregulators of TGF-. *J Am Soc Nephrol* **27**, 3117-3128 (2016).
14.   K. Martin *et al.*, PAK proteins and YAP-1 signalling downstream of integrin beta-1 in myofibroblasts promote liver fibrosis. *Nat Commun* **7**, 12502 (2016).
15.   I. Mannaerts *et al.*, The Hippo pathway effector YAP controls mouse hepatic stellate cell activation. *J Hepatol* **63**, 679-688 (2015).
16.   K. Zhang *et al.*, ω-3 PUFAs ameliorate liver fibrosis and inhibit hepatic stellate cells proliferation and activation by promoting YAP/TAZ degradation. *Sci Rep* **6**, 30029 (2016).
17.   M. Swiderska-Syn *et al.*, Hedgehog regulates yes-associated protein 1 in regenerating mouse liver. *Hepatology* **64**, 232-244 (2016).
18.   E. Seo *et al.*, The Hippo-Salvador signaling pathway regulates renal tubulointerstitial fibrosis. *Sci Rep* **6**, 31931 (2016).
19.   J. Xu *et al.*, Involvement of the Hippo pathway in regeneration and fibrogenesis after ischaemic acute kidney injury: YAP is the key effector. *Clin Sci (Lond)* **130**, 349-363 (2016).
20.   S. Anorga *et al.*, Deregulation of Hippo-TAZ pathway during renal injury confers a fibrotic maladaptive phenotype. *FASEB J* **32**, 2644-2657 (2018).
21.   K. Schlegelmilch *et al.*, Yap1 acts downstream of α-catenin to control epidermal proliferation. *Cell* **144**, 782-795 (2011).
22.   M. J. Lee, M. R. Byun, M. Furutani-Seiki, J. H. Hong, H. S. Jung, YAP and TAZ regulate skin wound healing. *J Invest Dermatol* **134**, 518-525 (2014).
23.   S. Noguchi *et al.*, TAZ contributes to pulmonary fibrosis by activating profibrotic functions of lung fibroblasts. *Sci Rep* **7**, 42595 (2017).
24.   T. Bertero *et al.*, A YAP/TAZ-miR-130/301 molecular circuit exerts systems-level control of fibrosis in a network of human diseases and physiologic conditions. *Sci Rep* **5**, 18277 (2015).
25.   D. J. Klionsky *et al.*, Guidelines for the use and interpretation of assays for monitoring autophagy. *Autophagy* **8**, 445-544 (2012).
26.   M. A. Wozniak, K. Modzelewska, L. Kwong, P. J. Keely, Focal adhesion regulation of cell behavior.

Biochim Biophys Acta **1692**, 103-119 (2004).

27.   G. Halder, S. Dupont, S. Piccolo, Transduction of mechanical and cytoskeletal cues by YAP and TAZ. *Nat Rev Mol Cell Biol* **13**, 591-600 (2012).

28.   S. Cabodi *et al.*, Integrins and signal transduction. *Adv Exp Med Biol* **674**, 43-54 (2010).

29.   J. R. Tse, A. J. Engler, Preparation of hydrogel substrates with tunable mechanical properties. *Curr Protoc Cell Biol* **Chapter 10**, Unit 10.16 (2010).

30.   T. Y. Kim *et al.*, Substrate trapping proteomics reveals targets of the βTrCP2/FBXW11 ubiquitin ligase. *Mol Cell Biol* **35**, 167-181 (2015).

31.   E. Coyaud *et al.*, BioID-based Identification of Skp Cullin F-box (SCF)β-TrCP1/2 E3 Ligase Substrates. *Mol Cell Proteomics* **14**, 1781-1795 (2015).

32.   E. L. Huttlin *et al.*, Architecture of the human interactome defines protein communities and disease networks. *Nature* **545**, 505-509 (2017).

33.   G. W. Won, S. H. Park, J. Park, Y. Lee, Y. H. Lee, Mammalian Hippo kinase pathway is downregulated by BCL-2 via protein degradation. *Biochem Biophys Res Commun* **512**, 87-92 (2019).

34.   B. Zhao, L. Li, K. Tumaneng, C. Y. Wang, K. L. Guan, A coordinated phosphorylation by Lats and CK1 regulates YAP stability through SCF(beta-TRCP). *Genes Dev* **24**, 72-85 (2010).

35.   J. R. Skaar, J. K. Pagan, M. Pagano, SCF ubiquitin ligase-targeted therapies. *Nat Rev Drug Discov* **13**, 889-903 (2014).

36.   Y. Cheng *et al.*, β-TrCP-mediated ubiquitination and degradation of liver-enriched transcription factor CREB-H. *Sci Rep* **6**, 23938 (2016).

37.   L. Busino *et al.*, Degradation of Cdc25A by beta-TrCP during S phase and in response to DNA damage. *Nature* **426**, 87-91 (2003).

38.   T. Cardozo, M. Pagano, The SCF ubiquitin ligase: insights into a molecular machine. *Nat Rev Mol Cell Biol* **5**, 739-751 (2004).

39.   N. Belaïdouni *et al.*, Overexpression of human beta TrCP1 deleted of its F box induces tumorigenesis in transgenic mice. *Oncogene* **24**, 2271-2276 (2005).

40.   , (!!! INVALID CITATION !!! {}).

41.   M. Guharoy, P. Bhowmick, P. Tompa, Design Principles Involving Protein Disorder Facilitate Specific Substrate Selection and Degradation by the Ubiquitin-Proteasome System. *J Biol Chem* **291**, 6723-6731 (2016).

42.   T. Y. Low *et al.*, A systems-wide screen identifies substrates of the SCFβTrCP ubiquitin ligase. *Sci Signal* **7**, rs8 (2014).

43.   N. Watanabe *et al.*, M-phase kinases induce phospho-dependent ubiquitination of somatic Wee1 by SCFbeta-TrCP. *Proc Natl Acad Sci U S A* **101**, 4419-4424 (2004).

44.   H. Fukushima *et al.*, SCF-mediated Cdh1 degradation defines a negative feedback system that coordinates cell-cycle progression. *Cell Rep* **4**, 803-816 (2013).

45.   Y. Kanemori, K. Uto, N. Sagata, Beta-TrCP recognizes a previously undescribed nonphosphorylated destruction motif in Cdc25A and Cdc25B phosphatases. *Proc Natl Acad Sci U S A* **102**, 6279-6284 (2005).

46.   R. E. Amir, H. Haecker, M. Karin, A. Ciechanover, Mechanism of processing of the NF-kappa B2 p100 precursor: identification of the specific polyubiquitin chain-anchoring lysine residue and analysis of the role of NEDD8-modification on the SCF(beta-TrCP) ubiquitin ligase. *Oncogene* **23**, 2540-2547 (2004).

47.   B. Doornaert *et al.*, Time course of actin cytoskeleton stiffness and matrix adhesion molecules in human bronchial epithelial cell cultures. *Exp Cell Res* **287**, 199-208 (2003).

48.   M. F. Pang *et al.*, Tissue Stiffness and Hypoxia Modulate the Integrin-Linked Kinase ILK to Control Breast Cancer Stem-like Cells. *Cancer Res* **76**, 5277-5287 (2016).

49.   C. Cluzel *et al.*, The mechanisms and dynamics of (alpha)v(beta)3 integrin clustering in living cells. *J Cell Biol* **171**, 383-392 (2005).

50.   G. L. Lin *et al.*, Activation of beta 1 but not beta 3 integrin increases cell traction forces. *FEBS Lett* **587**, 763-769 (2013).

51.   C. Arquint, F. Cubizolles, A. Morand, A. Schmidt, E. A. Nigg, The SKP1-Cullin-F-box E3 ligase βTrCP

and CDK2 cooperate to control STIL abundance and centriole number. *Open Biol* **8**, (2018).

52. V. Rausch, C. G. Hansen, The Hippo Pathway, YAP/TAZ, and the Plasma Membrane. *Trends Cell Biol* **30**, 32-48 (2020).

53. N. Elad *et al.*, The role of integrin-linked kinase in the molecular architecture of focal adhesions. *J Cell Sci* **126**, 4099-4107 (2013).

54. G. Burgstaller *et al.*, The instructive extracellular matrix of the lung: basic composition and alterations in chronic lung disease. *Eur Respir J* **50**, (2017).

55. J. S. Park *et al.*, Mechanical regulation of glycolysis via cytoskeleton architecture. *Nature* **578**, 621-626 (2020).

56. N. G. Kim, E. Koh, X. Chen, B. M. Gumbiner, E-cadherin mediates contact inhibition of proliferation through Hippo signaling-pathway components. *Proc Natl Acad Sci U S A* **108**, 11930-11935 (2011).

57. P. P. Provenzano, D. R. Inman, K. W. Eliceiri, P. J. Keely, Matrix density-induced mechanoregulation of breast cell phenotype, signaling and gene expression through a FAK-ERK linkage. *Oncogene* **28**, 4326-4343 (2009).

58. A. J. Engler, S. Sen, H. L. Sweeney, D. E. Discher, Matrix elasticity directs stem cell lineage specification. *Cell* **126**, 677-689 (2006).

59. G. Nardone *et al.*, YAP regulates cell mechanics by controlling focal adhesion assembly. *Nat Commun* **8**, 15321 (2017).

60. J. Seo, J. Kim, Regulation of Hippo signaling by actin remodeling. *BMB Rep* **51**, 151-156 (2018).

61. S. Alsamman *et al.*, Targeting acid ceramidase inhibits YAP/TAZ signaling to reduce fibrosis in mice. *Sci Transl Med* **12**, (2020).

62. M. Aragona *et al.*, A mechanical checkpoint controls multicellular growth through YAP/TAZ regulation by actin-processing factors. *Cell* **154**, 1047-1059 (2013).

63. J. A. Galan, J. Avruch, MST1/MST2 Protein Kinases: Regulation and Physiologic Roles. *Biochemistry* **55**, 5507-5519 (2016).

64. M. Chatzifrangkeskou *et al.*, RASSF1A is required for the maintenance of nuclear actin levels. *EMBO J* **38**, e101168 (2019).

65. R. M. Densham *et al.*, MST kinases monitor actin cytoskeletal integrity and signal via c-Jun N-terminal kinase stress-activated kinase to regulate p21Waf1/Cip1 stability. *Mol Cell Biol* **29**, 6380-6390 (2009).

66. A. Marzio *et al.*, The F-Box Domain-Dependent Activity of EMI1 Regulates PARPi Sensitivity in Triple-Negative Breast Cancers. *Mol Cell* **73**, 224-237.e226 (2019).

67. D. Guardavaccaro *et al.*, Control of meiotic and mitotic progression by the F box protein beta-Trcp1 in vivo. *Dev Cell* **4**, 799-812 (2003).

68. G. Bazzoni, D. T. Shih, C. A. Buck, M. E. Hemler, Monoclonal antibody 9EG7 defines a novel beta 1 integrin epitope induced by soluble ligand and manganese, but inhibited by calcium. *J Biol Chem* **270**, 25570-25577 (1995).

69. M. W. Pfaffl, A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* **29**, e45 (2001).

70. A. P. Z. P. Fiore *et al.*, Laminin-111 and the Level of Nuclear Actin Regulate Epithelial Quiescence via Exportin-6. *Cell Rep* **19**, 2102-2115 (2017).

71. E. Krieger, G. Vriend, YASARA View - molecular graphics for all devices - from smartphones to workstations. *Bioinformatics* **30**, 2981-2982 (2014).

72. W. Humphrey, A. Dalke, K. Schulten, VMD: visual molecular dynamics. *J Mol Graph* **14**, 33-38, 27-38 (1996).

73. B. J. Grant, A. P. Rodrigues, K. M. ElSawy, J. A. McCammon, L. S. Caves, Bio3d: an R package for the comparative analysis of protein structures. *Bioinformatics* **22**, 2695-2696 (2006).

74. T. Y. Low *et al.*, A systems-wide screen identifies substrates of the SCFbetaTrCP ubiquitin ligase. *Sci Signal* **7**, rs8 (2014).

75. S. Duan *et al.*, FBXO11 targets BCL6 for degradation and is inactivated in diffuse large B-cell lymphomas. *Nature* **481**, 90-93 (2012).