Meta-Analysis of Oxidative Transcriptomes in Insects

Hidemasa Bono [1,2,*]

[1] Program of Biomedical Science, Graduate School of Integrated Sciences for Life, Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-0046, Japan; bonohu@hiroshima-u.ac.jp

[2] Database Center for Life Science (DBCLS), Joint Support-Center for Data Science Research, Research Organization of Information and Systems, 178-4-4 Wakashiba, Kashiwa, Chiba 277-0871, Japan; bono@dbcls.rois.ac.jp

* Correspondence: bonohu@hiroshima-u.ac.jp; Tel.: +81-82-424-4013

Abstract: With the increasing compilation of data, meta-analysis, a statistical analysis combining results from multiple studies, is being widely used. To draw functional insights in a data-driven manner, we aimed to conduct a meta-analysis of oxidative transcriptomes in insect species based on data archived in public databases. We first explored and collected oxidative stress response-related RNA sequencing (RNA-seq) data for a wide variety of insect species, mainly from public gene expression databases by manually curating the results obtained. Only RNA-seq data of *Drosophila melanogaster* were found, which were then systematically analyzed with a newly developed RNA-seq analysis workflow for species without a reference genome sequence. The results were evaluated by two metric methods to construct a reference dataset for oxidative stress response studies. The results showed that many genes were downregulated under oxidative stress, which were found to be related to organ system process (GO:0003008) and adherens junction organization (GO:0034332) according to the gene enrichment analysis. A cross-species analysis was also performed. The RNA-seq data of *Caenorhabditis elegans* were curated since no RNA-seq data of insect species are currently available in public databases. This method, including the workflow developed, represents a powerful tool for deciphering conserved networks in oxidative stress response.

Keywords: insect, meta-analysis; oxidative stress; public database; RNA-seq; transcriptome

---

1. Introduction

With the increasing data collection, a statistical analysis called meta-analysis, which combines the results from multiple studies, is being widely used. By the end of 2020, over 200,000 records could be retrieved from PubMed (https://pubmed.gov/) by searching the keyword 'meta-analysis.' Non-typical data analysis specially designed for the collected data is now more often employed, with scientists becoming more aware of its potential and usefulness, paving the way for new findings that could not be achieved with the standard hypothesis-driven research methods [1]. Despite its useful analytical power, performing a meta-analysis can also be challenging owing to difficulties in collecting data from public databases and analyzing them. Nonetheless, this approach has been often used for investigating human-related subjects, allowing us to overcome major setbacks of human research, such as limited primary sample collection and small patient cohorts.

In life sciences, several datasets are archived in public databases that can be freely accessed and reused; this is particularly true for nucleotide sequence data. The database for nucleotide sequence, the Sequence Read Archive (SRA), now contains over 20 Petabases of open access nucleotide sequences (https://www.ncbi.nlm.nih.gov/sra/docs/sragrowth/). For example, chromatin immunoprecipitation sequencing (ChIP-seq) data in the SRA database were collected, curated, and pre-calculated for reuse in the ChIP-Atlas database (https://chip-atlas.org)[2] . An increasing number of meta-analyses can be found that re-use the public sequence data on a large scale, especially for coronavirus disease 2019 (COVID-19) studies. Gene set enrichment analysis re-using biomedical data for COVID-19, named Coronascape (https://metascape.org/COVID), makes it possible to

analyze the functional features of a gene list, specifically for COVID-19 studies [3]. However, few
studies reuse those resources when considering the amount of available data. In particular,
transcriptomes exposed to stress are a very interesting target for data meta-analysis. For example,
hypoxic transcriptomes could be collected from public gene expression data from human and mouse
cell lines and integrated with ChIP-seq data of transcription factors related to hypoxic stress by
using the ChIP-Atlas database [4].

Oxidative stress is a state of imbalance between the biological action of producing reactive
oxygen species (ROS) and the ability of the biological system to directly detoxify ROS or to repair the
damage caused. When the normal redox state of the biological system is disturbed, peroxides and
free radicals can damage proteins, lipids, and DNA, thereby resulting in the impaired function of
various intercellular mechanisms/organelles. In humans, oxidative stress is believed to be involved
in the development of cancer and Parkinson's disease, among other pathological conditions [5]. We
have explored the potential of silkworm (*Bombyx mori*) as a new model organism for investigating
Parkinson's disease [6] and superoxide dismutases [7,8]. Therefore, we focused on various effect of
oxidative stress to the living organism, instead of using cultured cells.

Supported by the above mentioned information, the present study aimed to apply a
meta-analysis approach to shed light on the oxidative transcriptomes of insects based on publicly
available data. We first pursued and collected oxidative stress response-related RNA sequencing
(RNA-seq) data for a wide variety of insect species. The collected gene expression data were
manually curated and then systematically analyzed with a new RNA-seq workflow designed
specifically for insects. The results were evaluated by two metric methods to construct a reference
dataset for oxidative stress response studies.

2. Materials and Methods

2.1 Curation of public gene expression data

Several public databases for gene expression data are available, whereas only one repository
archives nucleotide sequence data under the International Nucleotide Sequence Database
Collaboration (https://www.insdc.org/). This makes it difficult to re-use gene expression data in
public repositories. Therefore, we initially used the All Of gene Expression (AOE;
https://aoe.dbcls.jp/) online tool [9] to acquire oxidative stress-related gene expression data from the
public database. AOE integrates metadata not only from the Gene Expression Omnibus (GEO)
database from the U.S. National Center for Biotechnology Information (NCBI) [10] but also from the
ArrayExpress from the European Bioinformatics Institute (EBI) [11] and the Genomic Expression
Archive (GEA) from the DNA Data Bank of Japan (DDBJ) [12]. Moreover, AOE covers RNA-seq data
archived only in the SRA [13].

Conventional search using the keywords 'oxidative' and 'insect' yielded no hits because the
metadata for those were not rich enough for the given query. Therefore, extensive searches using
additional keywords (for example, 'paraquat' and 'rotenone' and other) were manually performed
to populate the dataset. Detailed search in the NCBI GEO web interface was also used to assess
records for insect species because some entries were not found by the AOE.

2.2 Retrieval and quality control of sequence data

For data retrieval from the SRA database and file conversion to FASTQ, we used the *prefetch*
and *fasterq-dump* tools from the SRA Toolkit (v2.9.6) (https://github.com/ncbi/sra-tools), respectively.
A parallel implementation of gzip (pigz) was used to compress FASTQ files and reducing the time
consumed in this process. For the trimming and quality control of reads from the SRA database,
Trim Galore! (v0.6.6) (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) with
Cutadapt (v1.18) [14] was applied to filter reads with unsatisfied quality. This step is
time-consuming, but it is indispensable for biologists to re-use reads archived in SRA, as the quality
of reads is not uniform.

2.3 Gene expression quantification

Salmon was used for quantifying the expression of transcripts using RNA-seq data [15]. For the
reproducibility of data analysis, Common Workflow Language (CWL) was introduced
(https://www.commonwl.org) (DOI: 10.6084/m9.figshare.3115156.v2). In the shell scripts to run
Salmon, we deployed the CWL definition files of salmon (indexing and quantification for single-end
and paired-end reads) in Pitagora Workflows (v0.1.0) maintained by Pitagora Network, as known as
Galaxy Community Japan (DOI: 10.5281/zenodo.2583024). Computer programs to run gene
expression quantification, including retrieval and quality control of sequence data, were freely
accessible in the GitHub repository (https://github.com/bonohu/SAQE).
The ratio of all gene pairs (termed ON-ratio) was calculated by pairing oxidative stressed and
normal state (Equation 1). A small number (in this case, 0.01) was added for avoiding the calculation
of logarithm of zero.
108

$$\text{ON-ratio} = \log(\text{gene expression}_{oxidative\ stress} + 0.01) - \log(\text{gene expression}_{normal\ state} + 0.01) \quad (1)$$

ON-ratio values for all paired samples can be classified into three groups: upregulated,
downregulated, or unchanged. When the ON-ratio was over the threshold for upregulation, the
gene was treated as 'upregulated'. Similarly, when the ON-ratio was under the threshold for
downregulation, the gene was treated as 'downregulated'. If the gene was treated as neither
'upregulated' nor 'downregulated', it was classified as 'unchanged'. The numbers of counts for
upregulated, downregulated, and unchanged were calculated for all genes. Several thresholds were
tested to optimize the calibration, and we adopted a 10-fold threshold for upregulation and 0.1-fold
threshold for downregulation.
An oxidative stress–normal state score (ON-score) was then calculated, as shown in Equation 2.
A detailed description of the ON-score is provided in the Results section.
120

$$\text{ON-score} = \text{count number}_{upregulated} - \text{count number}_{downregulated} \quad (2)$$

ON-ratio and ON-score were formally introduced as HN-ratio and HN-score in the
meta-analysis of the hypoxic transcriptomes [4].

2.4 Functional annotation and gene set enrichment analysis

Ensembl Biomart (https://www.ensembl.org/biomart/martview/) was used to extract the list of
transcript stable identifiers (IDs) and gene names, and corresponding gene ontology annotations for
Drosophila melanogaster and *Caenorhabditis elegans* from the Ensembl database (v101) [16]. The
orthologous gene relationship between these two species was also retrieved from the Ensembl
Biomart.
Metascape (https://metascape.org/) was used for the gene set enrichment analysis [3]. In the
enrichment analysis, the functional annotation provided by Metascape was used for the queried
genes.

2.5 Visualization

In addition to join two sets of data column by ID, we used TIBCO Spotfire Desktop (v7.6.0;
TIBCO Software Inc., Palo Alto, CA, USA) for producing scatter plots and histograms.

3. Results

3.1. Data collection

The overall procedure of our study is depicted in Figure 1. We first pursued oxidative
stress-related transcriptome data from insect species in public databases using the AOE tool.
Because the conventional keyword search by 'oxidative' and 'insect' in AOE yielded no hits, we
manually curated the search results for 'oxidative' by limiting the species to insects. Most data only
contained the word 'oxidative' in corresponding metadata without gene expression results by
oxidative stress.

In order to re-use the data for the comparative study in meta-analysis, we limited the
transcriptome data to RNA-seq, whereas several datasets in the public databases were quantified by
microarrays. The probe design for microarrays can be different from that used in RNA-seq; hence,
the comparative analysis of expression by genes can be complicated.

Currently, only RNA-seq data for *Drosophila melanogaster* can be found when using 'insects' as a
keyword in the public databases. Therefore, we also explored potential data based on names of
oxidative stress-related reagents in the NCBI GEO web interface.

Upon curating all candidate data, pair sets (oxidative stressed and control) were defined (Table
1). Paraquat was the most popular reagent to cause oxidative stress (source of stress) in this dataset.
153
154

155   **Figure 1**. Schematic view of insect oxidative transcriptome meta-analysis. Public databases were
156   searched, oxidative stress-related RNA-sequencing data were manually curated, and a meta-analysis
157   was performed.

158   **Table 1.** Dataset generated by the curation

SRA project ID	Oxidative stressed	Control	Source of stress (Reagent)	conc. (mM)	time (h)
SRP005712	SRR1509509	SRR1509506[l]	Rotenone	NA	NA
SRP005712	SRR1509508	SRR1509506[l]	Rotenone	NA	NA
SRP005712	SRR124259	SRR1509506[l]	Paraquat	5	NA

SRP005712	SRR103718	SRR1509506[1]	Paraquat	5	NA
SRP005712	SRR103721	SRR1509506[1]	Paraquat	10	NA
SRP005712	SRR103722	SRR1509506[1]	Paraquat	10	NA
SRP132308	SRR6677984	SRR6677982	Paraquat	50	6
SRP132308	SRR6677985	SRR6677983	Paraquat	50	6
SRP132308	SRR6677986	SRR6677982	Paraquat	50	12
SRP132308	SRR6677987	SRR6677983	Paraquat	50	12
SRP136174	SRR6874835	SRR6874832	Paraquat	10	24
SRP136174	SRR6874836	SRR6874833	Paraquat	10	24
SRP136174	SRR6874837	SRR6874834	Paraquat	10	24
SRP136174	SRR6874841	SRR6874838	Paraquat	10	24
SRP136174	SRR6874842	SRR6874839	Paraquat	10	24
SRP136174	SRR6874843	SRR6874840	Paraquat	10	24
SRP060444	SRR2088914	SRR2088913	Paraquat	NA	NA
SRP044035	SRR1505749	SRR1505771	UVA	NA	0.5
SRP044035	SRR1505750	SRR1505771	UVA	NA	1

[1] SRR1509506, SRR1509507, SRR1509516, SRR1509517. Abbreviations: NA, not available; UVA,

ultraviolet A radiation.

3.2. Quantification of transcripts using RNA-seq

For analyzing human and mouse RNA-seq data, there exists a pipeline centered on Salmon
called ikra (doi: 10.5281/zenodo.3352573). ikra not only automates the RNA-seq data analysis
process but also makes it easy to re-use data from public databases for meta-analysis. Currently, no
such pipeline exists for insect species. Therefore, we created a new generic pipeline for quantifying
RNA-seq data called Systematic Analysis for Quantification of Expression (SAQE), which consists of
the following steps:

1.   Retrieval of RNA-seq reads from the SRA database
2.   Conversion of data format and compression
3.   Trimming and quality control of reads
4.   Transcriptome assembly by Trinity
5.   Expression quantification by salmon
173

Step 4 (transcriptome assembly) was not used in this study because well-curated transcriptome
sequences were available for the target organisms. Nonetheless, this step is important for insects
whose genome sequences are not available for the data analysis.
For reproducibility of the study, docker containers with fixed versions were employed in the
last two steps. Furthermore, a script language, CWL, was partly employed in SAQE to call Docker
containers, which is an open standard for describing analysis workflows and tools in scientific
computing.   All   codes   of   SAQE   are   freely   available   from   GitHub
(https://github.com/bonohu/SAQE/).
As an output, SAQE yields the matrix for quantified expression data in transcript per million
(TPM). This expression matrix was uploaded to figshare and is publicly available (DOI:
10.6084/m9.figshare.13594538.v1).
Using the expression matrix, hierarchical clustering for sample direction was analyzed. A
dendrogram revealed that RNA-seq data from the same project clustered together (Figure S1),
suggesting the need for analysis of differences in the gene expression profile between samples
subjected to oxidative stress and those in the normal state.
189

3.3. Differentially expressed genes under oxidative stress

Thus, we tried to evaluate differentially expressed genes under oxidative stress. By pairing
oxidative stressed and normal state samples, we calculated the ratio of all gene pairs (termed
ON-ratio). The ON-ratio was previously introduced as HN-ratio (Hypoxia vs. Normoxia) in
meta-analysis of hypoxic transcriptomes [4]. After detailed investigation of data, we employed in
this study a smaller value (0.01, previously 1) for genes with weak expression. ON-ratio was
calculated for all sample pairs (Table 1). The complete list of ON-ratio is publicly available (DOI:
10.6084/m9.figshare.13596257.v1).

For meta-analysis of these values, the simplest way is to use 'averaged' ON-ratio for all
samples. The histogram of the averaged ON-ratio showed a skewed distribution curve (Figure 2A),
with few genes having high ON-ratios and more genes with negative ON-ratios.

In order to evaluate the oxidative stress-inducible genes, an oxidative stress-normal state score
(ON-score) was calculated for all genes, as described in the Materials and Methods. The complete list
of ON-score is also publicly available (DOI: 10.6084/m9.figshare.13596203.v2). In contrast to
ON-ratio, ON-score is quantal because it is calculated by subtraction of the number of counts of
upregulated from those of downregulated genes (Figure 2B). ON-score also showed skewed
distribution curve, with more genes having ON-scores below zero. These results showed that several
genes were downregulated under oxidative stress.

(A)

(B)

Figure 2. Distribution of calculated differential values for all genes. (**A**) Differential values averaged
for all experiments. (**B**) Differential values normalized by newly introduced metrics (ON-score).

We then performed gene set enrichment analysis using Metascape [3]. The analysis revealed
that genes upregulated in many samples with high ON-scores were related to cellular component
morphogenesis (GO: 0032989) and blood circulation (GO: 0008015) (Figure 3A). Metascape also
uncovered the functions of downregulated genes with low ON-scores (Figure 3B), being related to
organ system process (GO:0003008) and adherens junction organization (GO:0034332).

(A)

(B)

Figure 3. Histogram of gene set enrichment analysis for **(A)** upregulated (ON-score > 6) and **(B)** downregulated (ON-score < –7) genes under oxidative stress.

We then investigated ON-scores for genes annotated with GO:0006979 (response to oxidative stress), since the initial analyses did not show functional enrichment for oxidative stress-related annotated genes. In addition to GO:0006979, child terms of this GO group were included in the new differentially expressed gene analysis (Table 2).

Table 2. Child terms (direct descendants) of the gene ontology term GO:0006979 (response to oxidative stress)

Child GO ID	Child GO Term	Relationship to GO:0006979
GO:0001306	age-dependent response to oxidative stress	is_a
GO:0033194	response to hydroperoxide	is_a
GO:1902882	regulation of response to oxidative stress	regulates
GO:1902883	negative regulation of response to oxidative stress	negatively_regulates
GO:0070994	detection of oxidative stress	is_a
GO:1902884	positive regulation of response to oxidative stress	positively_regulates
GO:0000302	response to reactive oxygen species	is_a
GO:0080183	response to photooxidative stress	is_a
GO:2000815	regulation of mRNA stability involved in response to oxidative stress	part_of
GO:0034599	cellular response to oxidative stress	is_a

Original source: https://www.ebi.ac.uk/QuickGO/term/GO:0006979

(A)

(B)

Figure 4. ON-scores for **(A)** upregulated and **(B)** downregulated genes annotated with GO:0006979 (response to oxidative stress).

Among the identified oxidative stress-related genes, much more genes had negative ON-scores (Figure 4).

3.4. Cross-species analysis of transcriptomes under oxidative stress

For the cross-species study, we collected RNA-seq data from *C. elegans* because no RNA-seq data of insect species can be currently found in public databases. Similar to the approach used before, transcriptome sequencing data were curated for *C. elegans* and RNA-seq data were manually collected from the public databases (Table 3). In this dataset for *C. elegans*, rotenone was the most dominant reagent to cause oxidative stress, whereas paraquat was the most dominant reagent in the dataset for *D. melanogaster*.

Table 3. *C. elegans* oxidative transcriptome dataset generated by the curation

SRA project ID	Oxidative stressed	Control	Source of stress (Reagent)	conc. (mM)	time (h)
SRP070204	SRR3173735[1]	SRR3173720[2]	Sodium arsenite	10mM	5h
SRP021083	SRR827426	SRR827423	Rotenone	100nM	NA
SRP021083	SRR827427	SRR827424	Rotenone	100nM	NA
SRP021083	SRR827428	SRR827425	Rotenone	100nM	NA
SRP021083	SRR827432	SRR827429	Rotenone	100nM	NA
SRP021083	SRR827433	SRR827430	Rotenone	100nM	NA
SRP021083	SRR827434	SRR827431	Rotenone	100nM	NA
SRP021083	SRR827438	SRR827435	Rotenone	100nM	NA
SRP021083	SRR827439	SRR827436	Rotenone	100nM	NA
SRP021083	SRR827440	SRR827437	Rotenone	100nM	NA
SRP021083	SRR827443	SRR827441	Rotenone	100nM	NA
SRP021083	SRR827444	SRR827442	Rotenone	100nM	NA
ERP117708	ERR3580218	ERR3580215	Gamma radiation	10Gray	

ERP117708	ERR3580219	ERR3580216	Gamma radiation	10Gray
ERP117708	ERR3580220	ERR3580217	Gamma radiation	10Gray
ERP117708	ERR3580221	ERR3580215	Gamma radiation	100Gray
ERP117708	ERR3580222	ERR3580216	Gamma radiation	100Gray
ERP117708	ERR3580223	ERR3580217	Gamma radiation	100Gray
ERP117708	ERR3580224	ERR3580215	Gamma radiation	0.4Gray
ERP117708	ERR3580225	ERR3580216	Gamma radiation	0.4Gray
ERP117708	ERR3580226	ERR3580217	Gamma radiation	0.4Gray

[1] SRR3173735 – SRR3173746. [2] SRR3173720 – SRR3173734.

The same procedure for retrieval and quality control of sequence data and the same gene
expression quantification using SAQE as described above was done for this dataset (DOI:
10.6084/m9.figshare.13594541.v1). Using calculated differential values (ON-ratio; DOI:
10.6084/m9.figshare.13596263.v1) and ON-score (DOI: 10.6084/m9.figshare.13596206.v2),
transcriptomes with oxidative stresses in *D. melanogaster* and *C. elegans* were compared and
visualized in scatter plots (Figure 5). The integrated lists for ON-ratio and ON-score are publicly
available (DOI: 10.6084/m9.figshare.13597421.v1 and DOI: 10.6084/m9.figshare.13597241.v1
respectively). There was no. No striking correlation between these two organisms was observed.
Nonetheless, genes with correlated expression pattern could be useful for the candidate gene
selection.
250

(A)   (B)

Figure 5. Comparison of differentially expressed genes between *D. melanogaster* (X-axis) and *C.*
elegans (Y-axis). Colored dots correspond to genes annotated with GO:0006979 (response to oxidative
stress). **(A)** Differential values averaged for all experiments. **(B)** ON-score values.

We then looked into genes annotated with GO:0006979 (response to oxidative stress).
Upregulated experiments were dominant (ON-score positive) in nine genes (*CanA1, GstS1, Hsp22,*
Pde8, Sod2, foxo, per, ple, and *rl*) in both *D. melanogaster* and *C. elegans*, whereas downregulated
experiments were dominant (ON-score negative) in 13 genes (*CG9314, CG9416, CYLD, Ddc, Itp-r83A,*
ND-B17.2, Trap1, alph, bsk, mtd, park, ple, and *whd*) (Table 4). More genes had distinct ON-scores but
are not shown in Figure 5 because no *C. elegans* orthologs for those genes could be assigned.
The complete list of GO:0006979 (response to oxidative stress) annotated genes with ON-scores
for Table 4 is publicly available (DOI: 10.6084/m9.figshare.13597781.v1).
262

Table 4. *D. melanogaster* oxidative stress-related genes with ON-score feature (positive values or not) and *C. elegans* ortholog

ON-score	*C. elegans* ortholog	Genes
+	+	*CanA1, GstS1, Hsp22, Pde8, Sod2, foxo, per, ple, rl*
+	-	*CG10211, CG30487, CG31659, CG42331, CG6888, Chchd2, Daxx, GLaz, Gyc89Da, Gyc89Db, Jafrac1, MTF-1, Mnn1, Mt2, Nf1, Pi3K92E, SelR, TFAM, Thor, TotX, inaE, p38a, puc, scyl, srl*
-	+	*CG9314, CG9416, CYLD, Ddc, Itp-r83A, ND-B17.2, Trap1, alph, bsk, mtd, park, ple, whd*
-	-	*Akt1, CG31659, CG4009, CG42331, CG5948, Ccs, Daxx, GLaz, Gyc88E, Gyc89Da, InR, Irc, Jafrac1, Jafrac2, Karl, Keap1, MTF-1, Mnn1, Mt2, NLaz, Nf1, Pi3K92E, Prx5, Pxd, Ric, SelR, Sid, TotC, TotZ, cd, cnc, dec-1, fbl, inaE, mth, p38a, rut, spz, srl*

4. Discussion

In the present study, we collected transcriptome data specific to oxidative stress responses based on data stored in public databases, such as the NCBI GEO, EBI ArrayExpress, DDBJ GEA, and the SRA, by using the AOE web tool. Although RNA-seq is now widely used for transcriptomic studies in a wide variety of insect species, RNA-seq data specific from oxidative stress conditions and including a normal control were only found available in *D. melanogaster,* in spite of extensive manual curation. In most cases, the dataset only contains oxidative stressed RNA-seq data for specific knockdown (or knockout) conditions. Hence, a continuous effort to collect data from public databases is needed.

Collecting data is a critical and laborious step because there are many 'oxidative' descriptions in the metadata of public databases. Moreover, many of these are 'noise', which do not contain any transcriptome data for oxidative stress responses. In order to overcome this handicap, we must re-organize public databases for semantic queries. Currently, there is no computational method to filter these 'noise' events and manual curation by experts with domain knowledge is indispensable. Lack of information in the databases can be supplemented by searching published manuscripts (even supplemental materials in some cases). Deciphering metadata to collect the needed information from English texts also requires professional knowledge on sample preparation. Manual curation and creation of datasets are crucial for conducting this type of meta-analysis, which also represent a major hurdle for the meta-analysis.

As a new generic pipeline for quantifying RNA-seq data, SAQE was developed for insect species without reference genome sequences. Herein, SAQE was applied for insect species with reference transcriptomes in the public database. Only the transcriptome sequences of *D. melanogaster* and *C. elegans* were used, since both organisms are typical experimental models and transcriptome sequence sets are well-curated. Nonetheless, as it uses transcriptome assembly obtained from the execution of Trinity, SAQE can also be applied for non-model organisms without reference transcriptomes. Technically, SAQE is not pure 'workflow' in CWL currently. It is a group of shell scripts making full use of CWL command line tools and workflows for practical use in the insect study. Continuous development of SAQE itself is also needed.

The gene set enrichment analysis by Metascape for upregulated and downregulated genes showed interesting functional enrichments but is difficult to be interpreted in the context of oxidative stress. These results contrasted with those from the analysis for hypoxic transcriptomes in human cultured cells, which identified HIF1 TFPATHWAY, GO:0001666 response to hypoxia, and GO:0005996 monosaccharide metabolic process as being enriched. This is due to the various sources of oxidative stress and transcriptomic responses to oxidative stress, which are also not yet well-studied. Additionally, in insect species, functional annotation of genes is still underdeveloped, as compared with that in humans and mice. In other words, there are chances to find important discoveries from meta-analysis of these data.

Herein, the collection of datasets was relatively small and biased. Only 19 pairs of
oxidative-stress and normal condition samples were collected, contrasting to over hundred pairs of
hypoxic and normoxic transcriptomes meta-analyzed in the previous study. In addition, the most
represented stress source was paraquat (Table 1). Hence, more data and more species are needed for
a better meta-analysis. Nevertheless, interesting genes could be found in this meta-analysis.
Knockdown of *Sod2* in *Tribolium castaneum* was reported to impair sensitivity to paraquat [17], and
Sod2 was one of the genes with positive ON-scores in both species (Table 4). Therefore, we believe
that this meta-analysis is a repository of future findings.
Our future work will gather oxidative stress-related transcriptome data for insect species to
continuously update this meta-analysis. This approach will potentially uncover conserved
mechanisms at the molecular level.

5. Conclusions

In this study, we tried to collect oxidative stress-related transcriptomic data in a wide variety of
insect species from public databases. To date, such transcriptomes are only available for *D.*
melanogaster; however, around 20 pairs of oxidative stressed and normal state transcriptomes were
listed.
For analyzing such transcriptomes, an RNA-seq analysis workflow for species without
reference genome sequences was developed. Downregulated genes were dominant in spite of the
use of several oxidative stress reagents at variable conditions.
Currently, no distinct features were retrieved from enrichment analyses of upregulated and
downregulated gene sets, but enrichments for other functional groups provided interesting new
insights. We focused on genes annotated with oxidative stress-related gene ontology terms in the
comparative analyses of *C. elegans* oxidative stressed transcriptomes from the same curation process
to *D. melanogaster*.
This method, including the workflow developed, will represent a powerful tool for deciphering
conserved networks in oxidative stress response.
328

Supplementary Materials: Figure S1: Hierarchical clustering of all data (samples) in transcript per million
(TPM).

Data Availability Statement: The data presented in this study are openly available in FigShare at
https://doi.org/10.6084/m9.figshare.c.5271731). Source codes to replicate the study are also freely available at
GitHub (https://github.com/bonohu/SAQE).

Author Contributions: Conceptualization, H.B.; methodology, H.B.; software, H.B.; validation, H.B.; formal
analysis, H.B.; investigation, H.B.; resources, H.B.; data curation, H.B.; writing—original draft preparation,
H.B.; writing—review and editing, H.B.; visualization, H.B.; supervision, H.B.; project administration, H.B.;
funding acquisition, H.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by ROIS-DS-JOINT (010RP2020).

Acknowledgments: We thank Ms. Yoko Ono for helpful discussion. Computations were performed on the
computers at Hiroshima University Genome Editing Innovation Center.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the
study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to
publish the results.

References

1.   Parkinson, N.; Rodgers, N.; Head Fourman, M.; Wang, B.; Zechner, M.; Swets, M.C.; Millar, J.E.; Law, A.;
Russell, C.D.; Baillie, J.K.; et al. Dynamic Data-Driven Meta-Analysis for Prioritisation of Host Genes
Implicated in COVID-19. *Sci. Rep.* **2020**, *10*, 22303, doi:10.1038/s41598-020-79033-3.

2. Oki, S.; Ohta, T.; Shioi, G.; Hatanaka, H.; Ogasawara, O.; Okuda, Y.; Kawaji, H.; Nakaki, R.; Sese, J.; Meno, C. ChIP-Atlas: A Data-Mining Suite Powered by Full Integration of Public ChIP-Seq Data. *EMBO Rep.* **2018**, *19*, doi:10.15252/embr.201846255.

3. Zhou, Y.; Zhou, B.; Pache, L.; Chang, M.; Khodabakhshi, A.H.; Tanaseichuk, O.; Benner, C.; Chanda, S.K. Metascape Provides a Biologist-Oriented Resource for the Analysis of Systems-Level Datasets. *Nat. Commun.* **2019**, *10*, 1523, doi:10.1038/s41467-019-09234-6.

4. Bono, H.; Hirota, K. Meta-Analysis of Hypoxic Transcriptomes from Public Databases. *Biomedicines* **2020**, *8*, doi:10.3390/biomedicines8010010.

5. Dias, V.; Junn, E.; Mouradian, M.M. The Role of Oxidative Stress in Parkinson's Disease. *J. Park. Dis.* **2013**, *3*, 461–491, doi:10.3233/JPD-130230.

6. Tabunoki, H.; Ono, H.; Ode, H.; Ishikawa, K.; Kawana, N.; Banno, Y.; Shimada, T.; Nakamura, Y.; Yamamoto, K.; Satoh, J.-I.; et al. Identification of Key Uric Acid Synthesis Pathway in a Unique Mutant Silkworm Bombyx Mori Model of Parkinson's Disease. *PloS One* **2013**, *8*, e69130, doi:10.1371/journal.pone.0069130.

7. Nojima, Y.; Ito, K.; Ono, H.; Nakazato, T.; Bono, H.; Yokoyama, T.; Sato, R.; Suetsugu, Y.; Nakamura, Y.; Yamamoto, K.; et al. Superoxide Dismutases, SOD1 and SOD2, Play a Distinct Role in the Fat Body during Pupation in Silkworm Bombyx Mori. *PloS One* **2015**, *10*, e0116007, doi:10.1371/journal.pone.0116007.

8. Kobayashi, Y.; Nojima, Y.; Sakamoto, T.; Iwabuchi, K.; Nakazato, T.; Bono, H.; Toyoda, A.; Fujiyama, A.; Kanost, M.R.; Tabunoki, H. Comparative Analysis of Seven Types of Superoxide Dismutases for Their Ability to Respond to Oxidative Stress in Bombyx Mori. *Sci. Rep.* **2019**, *9*, 2170, doi:10.1038/s41598-018-38384-8.

9. Bono, H. All of Gene Expression (AOE): An Integrated Index for Public Gene Expression Databases. *PloS One* **2020**, *15*, e0227076, doi:10.1371/journal.pone.0227076.

10. Barrett, T.; Wilhite, S.E.; Ledoux, P.; Evangelista, C.; Kim, I.F.; Tomashevsky, M.; Marshall, K.A.; Phillippy, K.H.; Sherman, P.M.; Holko, M.; et al. NCBI GEO: Archive for Functional Genomics Data Sets--Update. *Nucleic Acids Res.* **2013**, *41*, D991-995, doi:10.1093/nar/gks1193.

11. Athar, A.; Füllgrabe, A.; George, N.; Iqbal, H.; Huerta, L.; Ali, A.; Snow, C.; Fonseca, N.A.; Petryszak, R.; Papatheodorou, I.; et al. ArrayExpress Update - from Bulk to Single-Cell Expression Data. *Nucleic Acids Res.* **2019**, *47*, D711–D715, doi:10.1093/nar/gky964.

12. Kodama, Y.; Mashima, J.; Kosuge, T.; Ogasawara, O. DDBJ Update: The Genomic Expression Archive (GEA) for Functional Genomics Data. *Nucleic Acids Res.* **2019**, *47*, D69–D73, doi:10.1093/nar/gky1002.

13. Kodama, Y.; Shumway, M.; Leinonen, R.; International Nucleotide Sequence Database Collaboration The Sequence Read Archive: Explosive Growth of Sequencing Data. *Nucleic Acids Res.* **2012**, *40*, D54-56, doi:10.1093/nar/gkr854.

14. Martin, M. Cutadapt Removes Adapter Sequences from High-Throughput Sequencing Reads. *EMBnet.journal* **2011**, *17*, 10–12, doi:10.14806/ej.17.1.200.

15. Patro, R.; Duggal, G.; Love, M.I.; Irizarry, R.A.; Kingsford, C. Salmon Provides Fast and Bias-Aware Quantification of Transcript Expression. *Nat. Methods* **2017**, *14*, 417–419, doi:10.1038/nmeth.4197.

16. Kinsella, R.J.; Kähäri, A.; Haider, S.; Zamora, J.; Proctor, G.; Spudich, G.; Almeida-King, J.; Staines, D.; Derwent, P.; Kerhornou, A.; et al. Ensembl BioMarts: A Hub for Data Retrieval across Taxonomic Space. *Database J. Biol. Databases Curation* **2011**, *2011*, bar030, doi:10.1093/database/bar030.

389   17.   Tabunoki, H.; Gorman, M.J.; Dittmer, N.T.; Kanost, M.R. Superoxide Dismutase 2 Knockdown Leads to

390   Defects in Locomotor Activity, Sensitivity to Paraquat, and Increased Cuticle Pigmentation in Tribolium

391   Castaneum. *Sci. Rep.* **2016**, *6*, 29583, doi:10.1038/srep29583.

393