1 Outer Hair Cell Function is Normal in βV Spectrin Knockout Mice.
2

3 Michael C. Stankewich[1*], Jun-Ping Bai[2*], Paul R. Stabach[1], Saaim Khan[3], Winston
4 J.T. Tan[6], Alexei Surguchev[6], Lei Song[4], Jon S. Morrow[1,5], Joseph Santos-
5 Sacchi[6,7,8], Dhasakumar S. Navaratnam[2,6,7 ^]
6
7
8
9 1 Department of Pathology, Yale School of Medicine, New Haven, CT, U.S.A
10
11 2 Department of Neurology, Yale School of Medicine, New Haven, CT, U.S.A
12
13 3 Cranbrook Kingswood Upper School, MI, USA
14
15 4 Department of Otolaryngology-Head and Neck Surgery, Shanghai Ninth
16 People's Hospital, Shanghai Jiaotong University School of Medicine, Shanghai,
17 China
18
19 5 Department of Molecular, Cellular, and Developmental Biology, Yale University,
20 New Haven, CT, U.S.A
21
22 6 Department of Surgery, Yale School of Medicine, New Haven, CT, U.S.A
23
24 7 Department of Neuroscience, Yale School of Medicine, New Haven, CT, U.S.A
25
26 8 Department of Cellular and Molecular Physiology, Yale University, New Haven,

27 CT, U.S.A

28 * Co-first authors
29 ^ Corresponding Author
30
31 Address correspondence to Dr. Dhasakumar Navaratnam
32 Yale School of Medicine, Department of Neurology, PO Box 208018,
33 New Haven, CT., 06519
34 email: dhasakumar.navaratnam@yale.edu
35

Keywords: cochlea, auditory brainstem response, electromotility, distortion

product otoacoustic emissions, Spnb5, ribbon synapse, cochlear nerve

ABSTRACT

Reports have proposed a putative role for βV spectrin in outer hair cells

(OHCs) of the cochlea. In an ongoing investigation of the role of the cytoskeleton

in electromotility, we tested mice with a targeted exon deletion of βV spectrin

(Spnb5), and unexpectedly find that Spnb5$^{(-/-)}$ animals' auditory thresholds are

unaffected. Similarly, these mice have normal OHC electromechanical activity

(otoacoustic emissions) and non-linear capacitance. In contrast, magnitudes of

auditory brainstem response (ABR) wave 1-amplitudes are significantly reduced.

Evidence of a synaptopathy was absent with normal hair cell CtBP2 counts.

In Spnb5$^{(-/-)}$ mice, the number of afferent and efferent nerve fibers is decreased.

Consistent with this data, Spnb5 mRNA is present in Type I and II spiral ganglion

neurons, but undetectable in OHCs. Together, these data establish that βV

spectrin is important for hearing, affecting neuronal structure and function.

Significantly, these data exclude βV spectrin as functionally important to OHCs as

has been previously suggested.

53
54

INTRODUCTION

In the early 1990s, several groups made key seminal observations of a spectrin/actin lattice in the cochlear outer hair cells (OHCs) (Holley & Ashmore, 1990) (Holley, Kalinec, & Kachar, 1992). Their electron micrographs propelled an exquisite model of the spectrin/actin lattice assembled just beneath the OHC lateral membrane with longitudinal spectrin molecules linked to circumferential actin filaments that, in turn, were linked to the plasma membrane by electron-dense pillars. Their work also identified a pool of spectrin in the actin-rich cuticular plate. Another study evaluated spectrin proteins in cochlear innervation during rodent development (Hafidi, Despres, & Romand, 1990). They identified differences in developmental expression of the erythroid beta spectrin (Spnb1 spectrin) compared to the most ubiquitous nonerythroid form $\alpha II / \beta II$ spectrin, similar to what is observed in the brain. What is not certain from these earlier studies is the specific spectrins involved, a family now known to consist of two alpha spectrins (Spna1 and 2) and five beta spectrins (Spnb1-5). More recently, it has been shown that Spnb4 is restricted to the nodes of Ranvier in myelinated axons and is also a deafness-related gene (Parkinson et al., 2001) (Lewis et al., 2018). It appears from these studies that the deafness in the quivering mouse is likely not originating in the cochlea but rather is central in origin. Furthermore, Spnb2 spectrin, when selectively deleted from the mouse cochlea sensory epithelia by crossing Spnb2 floxed mice with Atoh-1 Cre-recombinase, causes profound deafness due to disruption of the stereocilia rootlets in the cuticular plate in hair cells (Liu et al., 2019). Thus, the function of specific spectrins, particularly

as it relates to the OHC lateral membrane and cochlear innervation, remains

unclear. In humans, the molecule βV spectrin (SPTBN5) that was first discovered

in the retina, is involved with the retinal disorder, Usher syndrome (Stabach &

Morrow, 2000) (Papal et al., 2013). Further reports have proposed a putative role

for βV spectrin in a second sensorineural function, hearing. One group's

observations demonstrated that antisera raised against the C-terminal peptide of

SPTBN5, labeled structures along the lateral membrane of the OHC (Legendre,

Safieddine, Kussel-Andermann, Petit, & El-Amraoui, 2008) (Cortese et al., 2017).

This observation drew attention because detection of this giant spectrin in tissue

other than retina and testis has been elusive. This finding raised the possibility that

βV spectrin was involved in the electromotile response of OHCs.

Continuing an ongoing effort in our lab to define the cytoskeleton's role in

electromotility (Bai et al., 2010), we sought to determine the role of βV spectrin in

OHC function. In this study, we analyzed βV spectrin expression in the cochlea

and addressed its function in mice with a targeted deletion of Spnb5. We tested

hearing by auditory brainstem responses (ABRs), measured OHC-mediated

otoacoustic emissions and OHC nonlinear capacitance (NLC), and evaluated the

number of inner hair cell (IHC) ribbon synapses and nerve fibers that innervate the

cochlear sensory epithelium. We find that while OHC function is unaffected in the

absence of βV spectrin, nerve counts are reduced at the habenular, in all three

turns of the cochlea, and in efferent fibers in the mid and basal turns, which

collectively account for the reduced wave I amplitudes in ABRs. Using RNAscope

in situ hybridization, we confirm expression of Spnb5 mRNA in Type I and II SGNs

and absent expression in OHCs. Moreover, mRNA transcript was significantly

reduced in the SGN of Spnb5$^{(-/-)}$ mice, confirming knockdown of the gene. We

conclude that βV spectrin is necessary to maintain the integrity or function of

afferent and efferent nerves of the peripheral auditory system, but plays no role in

the OHC itself.

RESULTS
109

Mouse βV spectrin amino acid sequence is divergent from the human

protein.

Surprisingly, a sequence alignment comparing the mouse amino acid sequence to

the human amino acid sequence reveals a significant divergence. In contrast, other

beta spectrins, including beta I-IV, have ~ 90% amino acid conservation between

human and mouse sequences. By aligning human and mouse βV spectrin, we note

a significant divergence at the N- and C-termini of the proteins and an internal

stretch of amino acids from position 817-922 (**Figure 1A**). Of note is that the

initiator methionine residue in the human form is conserved between mouse and

human genes and lies at position 36 and within exon 2 (**Figure 2**). The mouse

sequence incorporates an unusual homorepeat of glutamines and, notably, does

not have a bona fide pleckstrin homology (PH) domain seen in the human form.

Spnb5 mRNA tissue expression.

Spnb5 spectrin is expressed in most mouse tissues but at low levels. In ENCODE,

the level of Spnb5 transcripts is an order of magnitude less abundant than the

ubiquitous form Spnb2 (**Figure 1B**) (Yue et al., 2014). The peak abundance of

Spnb2 approaches 50 RPKM while Spnb5 is 0.5 RPKM in testis (**Figure 1B, C**).

Interestingly, the GTEX dataset shows high expression of Spnb5 in the peripheral

(tibial) nerve (**Figure 1D**). To validate that mRNA is detected in mice, we tested

RNA from the testis with primer sets that bridged exons. By standard RT-PCR,

Spnb5 mRNA fragments encoding both the N-terminus and C-terminus were

amplified from testis and brain RNA, and its identity was confirmed by Sanger

sequencing. In contrast to brain and testicular tissue, we could not detect Spnb5

mRNA in isolated OHCs and whole organ of Corti by both conventional and nested

PCR (Table 1 details the gene-specific primer sets). Using RT-PCR from ~3000

OHCs and organ of Corti, we detect both Spnb1 and Spnb2. Using nested PCR,

amplicons for these spectrins are robust, while we do not visualize any product for

Spnb5 (**Figure 1E**). These data are in accord with data derived from the public

data sets of transcriptome profiling for OHCs in gEAR (https://umgear.org/) (**See**

supplementary Figure1)**.** Furthermore, *in situ* hybridization for Spnb5 using

RNAscope corroborates these findings. We are unable to detect Spnb5 mRNA in

OHCs (**see Supplementary Figure 2, panels C and S**), whereas mRNA for the

potassium channel, Kcnma1, is readily detected in OHCs (green signal in **panel**

B). In contrast, the Spnb5 probe is detected in spiral ganglion neurons (SGNs) (red

signal in both **panels C and P**). To further characterize the expression of Spnb5

in SGNs, we co-labeled Spnb5 (green) with known markers of myelinated Type I

ganglion cells (tyrosine hydroxylase, (TH)) and unmyelinated Type II ganglion cells

(Epha4) in red (see Supplementary Figure 3). We find that Spnb5 signal overlaps

with the signals from both TH (panel A) and Epha4 (panel B). These data

demonstrate that Spnb5 is expressed in both Type I and II SGNs.

A

Figure 1. (A) Spnb5 mouse (ENSMUST00000156159.4) **amino acid sequence differs from humans.** Shown is the comparison of the deduced amino acid sequence of mouse (lower) and human (upper) Spnb5. Conserved amino acids are marked in red. There is divergence at the N and C termini. In addition, a region in the middle of the protein from posiiton 817-922 shows significant divergence. The mouse version of this gene shares only 65.6 percent identity with the human form. **(B, C) Spnb5 is expressed in brain and testis.** The relative abundance of Spnb2 transcripts as compared to Spnb5 in mouse tissues. Shown are the expression of Spnb2 and 5 in RNA Seq. NCBI reference datasets (ENCODE; Bioproject PRJNA66167 (Yue et al., 2014). On average, the abundance of Spnb5 mRNA is an order of magnitude less abundant than Spnb2. Of the tissues examined, adult testis has the most mRNA for Spnb5. **(D) In human, Spnb5 is most expressed in peripheral nerve.** The relative abundance of Spnb5 in different tissues in GTEX database is presented (version 6) (Consortium, 2013). Of all the tissues examined Spnb5 is expressed in highest amounts in the peripheral nerve (tibial nerve). **(E) Spnb5 mRNA is not detectable in the organ of Corti or OHCs.** Shown are PCR amplifications from mouse brain (lanes 2-4), organ of Corti (lanes 6-8) and OHCs (lanes 10-12) of Spnb1 (lanes 2, 6 and 10), Spnb2 (lanes 3, 7 and 11) and Spnb5 (lanes 4, 8 and 12). The upper panel shows amplicons for these three mRNA transcripts using outer primers while the lower panel shows amplicons of primers using a nested set of primers. All three spectrins are detectable in brain. Spnb1 and 2 are detectable in both the organ of Corti and OHCs (lanes 6-7 and 10-11). Spnb5 however, is not detectable using

B

Spnb2 tissue expression: ENCODE

C

Spnb5 tissue expression: ENCODE

D SPTBN5 tissue expression: GTEX

E

Figure 1

154 **Supplementary Figure 1.**

Figure S1. Expression levels for each of the beta spectrins in OHCs.
Mining data from the gene Expression Analysis Resource (gEAR) we graphed the abundance of each of the five beta spectrin genes. This data set was derived from transcriptome analysis using sc-RNAseq of cell populations isolated from mouse cochleae (Li et al., 2018).

155
156
157

Targeted knockout of Spnb5 in mice using homologous recombination.

In order to properly target the βV spectrin gene in mice, we needed to know its

precise gene structure. While the human gene has been well characterized and

indexed since its discovery in 2000 (Stabach & Morrow, 2000), there has been

ambiguity in the intron/exon usage in the mouse gene found on chromosome 2.

The GENBANK entry for Spnb5 has been modified many times in the last several

years, with the last significant annotation in 2007 (Benson et al., 2013). This

annotation was derived by automated computational analysis using gene

prediction algorithms. The exon prediction is made difficult because 1) the gene is

very large, so PCR amplifying a full-length cDNA is not trivial, and 2) the

conservation between human and mouse intron/exon boundaries is weak at the N-

and C-termini. The current and accepted annotated version in both the UCSC

genome browser and in Ensembl shows Spnb5 spectrin mRNA in the mouse to be

14.4 kb with a 10,872 bp ORF that has 66 exons and codes for 3623 amino acids:

NM_001370938: NP_001357867.1. This version has been validated in the

ENCODE project based on the robustness of next-generation sequencing efforts

over the last several years (Yue et al., 2014).

Shown in the schematic in Figure 2A is the strategy for targeted deletion of exon

2 of Spnb5. The knockout strategy was based on the transcript ID:

ENSMUST00000156159.3, and a targeting construct was generated in which

three Lox P sites were inserted to flank the neomycin selection marker and exon

2 containing the start codon, resulting in disruption of the open reading frame upon

Cre-mediated excision. Exon 2 contains the ATG start codon (with a consensus

Kozak sequence) and an additional 37 amino acids at its N-terminus. Notably, the

codon for the methionine residue corresponding to the human start codon is also

contained within the second exon. A genomic bacterial artificial chromosome

containing the Spnb5 locus was obtained, and genomic DNA fragments containing

the first exons of Spnb5 were subcloned in pEasy (a gift from Sankar Ghosh). The

targeting vector comprised a 4.3 kb long arm, a 4.1 kb short arm, and a middle

selection cassette which contains exon two flanked by a PGK–neomycin

resistance (Neo), two FRT sites, and the herpes simplex thymidine kinase gene

(Figure 2B and Primer Sets in Table 2). The targeting plasmid DNA was linearized

with Cla1 and electroporated into mouse embryonic stem cells (ESCs)

(129SV/EV). ESCs were selected for neomycin/G418 resistance. Positive clones

were identified by PCR and confirmed by Southern blot analysis using external

and internal probes. A correctly targeted ESC clone was injected into C57BL/6J

blastocysts to produce germline transmitting chimeric mice. These were then

mated with actin-Cre C57BL/6J transgenic mice for generation of heterozygous

Spnb5 transgenic mice with both the Neo cassette and exon two removed

(Spnb5$^{(+/-)}$). The resulting Spnb5$^{(+/-)}$ offspring were crossed to generate βV

spectrin-deficient mice. The genotypes of these mice were confirmed by PCR

amplification (Figure 2C). Chimeric mice were backcrossed onto the C57BL/6J

strain. Litters from heterozygous breeders Spnb5$^{(+/-)}$ / Spnb5$^{(+/-)}$ were of normal

number, and the mice of all genotypes were phenotypically indistinguishable with

normal lifespans >2 yrs. old. We confirmed decreased expression of Spnb5 in

cDNA from testes of Spnb5$^{(-/-)}$ mice, where we found equivalent expression of

Spnb2 and Spna2 (Figure 2D and Primer Sets in Table 3). The decreased level of

Spnb5 mRNA was corroborated by RNAScope in situ hybridization (ISH)

experiments. In low power images of cochlear tissue (Supplementary Figure 2,

panels A- I), the signal for Spnb5 (red) in the SGN is significantly reduced in KO

mice. Quantitation of the mean fluorescence intensity shown in panel V reveals

that the amount of signal in SGNs is significantly reduced by greater than 60% in

the SGN of Spnb5$^{(-/-)}$ animals. Higher power magnification ISH images, when

labeled with Spnb5 probe alone, reveals a significant reduction in both the number

of fluorescent particles per neuron in KO tissue (panel W). On average, the number

of particles per neuron with positive granules was significantly reduced by greater

than ~50%. In addition, both the total fluorescence intensity and the number of

neurons that contained Spnb5 is dramatically reduced in the SGNs of Spnb5$^{(-/-)}$

animals (panel C vs. F). The binding site for the RNAscope *in situ* probes (at

218 position 7163 – 8000 in XM_006500537; 3669-4506 in ENSMUST00000156159.4)

219 corresponds to ~ 1300 amino acids after the initiator methionine. These data

220 suggest that the transcript up to position 3669-4506 was significantly reduced in

221 the SGN of knockout animals. It should be noted that the actin binding of spectrins

222 occurs through calponin homology (CH1 and CH2) sites present at the N-terminus

223 of the proteins. In Spnb5, these sites are located at aa 20-124 (CH1) and aa 143-

224 247 (CH2) and encoded by the transcript well before the RNAscope *in situ* probe

225 binding sites. In other beta spectrins, loss of function mutations occur within the

226 CH1 and CH2 actin binding domains (Cousin et al., 2020). While it is still possible

227 that there are mRNA transcripts that are shorter with in-frame methionine start

228 sites, these transcripts would not be expected to produce a functional protein in

229 lieu of it lacking the CH1 and CH2 actin binding domains (there are, for instance,

230 43 in-frame methionine codons, 19 of which also contain consensus Kozak

231 sequences, that lie downstream of the *in situ* hybridization probe binding sites).

A.

B.

C.

D.

Figure 2. Exon 2 of the Spnb5 gene was removed using homologous recombination. (A) Schematic of the strategy used to delete exon 2. The construct contained a short arm of exon 1 and a long arm of exons 3, 4, and 5 together with the intervening introns. Both the selection marker Neo and exon two are flanked by Lox P sites. Linearized plasmid was injected into ES cells and clones with homologous recombination and selected by neomycin (Neo)/ G418 resistance. Mice were then crossed with Actin-Cre mice to generate floxed mice lacking the Neo cassette and exon 2. **(B) Vector map of pEASY used to target Spnb5 deletion.** Exon 2 was subcloned between two Lox P sites adjacent to the Neo gene. These two genomic fragments were in turn placed between a 4.1kb short arm containing Exon 1 and a 4.3kb long arm containing exons 3, 4, and 5 and the intervening introns. The vector also contains a thymidine kinase and ampicillin gene allowing for selection. The plasmid was linearized with Cla1. **(C) PCR amplification of genomic DNA confirming deletion of exon two and flanking intronic sequence.** In genomic DNA from Spnb5$^{(-/-)}$ mice**,** primers that amplify a product no longer contain part of intron 1; exon 2; and part of intron 2/3. Lane 1 base pair standards, lane 2 Spnb5$^{(+/-)}$, lane 3 Spnb5$^{(-/-)}$, lane 4 Spnb5$^{(+/+)}$. The larger fragment of 665 bp consists of unexcised exon two while the smaller fragment consists of genomic DNA fragment from which exon two has been excised. **(D) Decreased expression of Spnb5 in Spnb5$^{(-/-)}$ testicular cDNA compared to Spnb2 and Spna2**. Spnb5 exons 58-60 in lanes 2-3, Spnb5 exon 53-55 in lanes 4-5; Spnb2 in lanes 7-8 (exons 28-29) and Spna2 (exons 54-55) in lanes 10-11**.** WT in lanes 2, 4, 7 and 10. Spnb5$^{(-/-)}$ in lanes 3, 5, 8 and 11.

Figure 2

234
235
236

Supplementary Figure 2.

Figure S2. *In situ* hybridization (RNAscope) shows the expression of Spnb5 and Kcnma1 in WT and Spnb5[(-/-)] cochlear tissue. In the top image, panels (A-F) are a low magnification (10x) comparison of Spnb5 mRNA (red, C & F) and Kcnma1 mRNA (green, B & E) between WT and KO from the apical turn of the organ of Corti. Merged images are displayed in panels (A & D) and the DapB negative control images (-ve control) are shown in panels (G-I). Single asterisks indicate spiral ganglion neurons (SGNs) and double asterisks highlight regions with hair cells (HC). Mean SGN Spnb5 fluorescence intensity of WT animals were significantly different from KO animals and negative controls (V) (WT 1.744 ± 0.13 SEM, n = 9, KO 0.80 ± 0.15 SEM, n = 7, -ve control 0.60 ± 0.09 SEM, n = 5 one-way ANOVA, $p<0.001$). Areas of SGNs were demarcated based on staining of nuclei (DAPI). Scale bar, 200 μm. The bottom image (J-U) is a single-color, high magnification (40x) for WT and KO tissue. mRNA for Spnb5 was assessed in the SGNs (P-R) and HCs (S-U). The upper panels (J-O) are the same images with DAPI demarcating cell nuclei. Panels (J,P,K,Q) representing Spnb5 mRNA levels (red signal) in WT and KO animals show reduced expression of Spnb5 mRNA in KO animals. Negative control (DapB) images are shown in panels L & R. Of note, Spnb5 is not expressed in all SGNs. In KO animals, the number of Spnb5 granules per neuron, the overall fluorescence intensity of these granules, and the number of neurons with granules were all reduced compared to WT animals. Mean number of Spnb5 mRNA granules per SGN in WT and KO mice (W) were significantly different (WT 23.8 ± 1.3 SEM, n = 23 and KO 14.4 ± 1.7 SEM, n = 28 neurons, respectively, students t test, P <0.001). Here, only those neurons with Spnb5 signal were included. Panels (M,S,N,T) of Spnb5 in HCs of WT and KO cochlea show no signal above DapB background (O & U). Scale bar, 20 μm.

258 **Supplementary Figure 3.**

259
260 Figure S3. *In situ* hybridization (RNAscope) shows the expression of Spnb5 in SGNs I and
261 II. In the top image, panel (A) is a merged high magnification (40x) comparison of Spnb5 (green)
262 and tyrosine hydroxylase (TH) (red) mRNA. Yellow granules in this image demarcate coincident
RNA labeling in SGN Type I cells. Panel (B) is a merged image of Spnb5 (green) and Epha4 (red)
263 signals. Yellow granules are coincident labeling of Spnb5 and Epha4 mRNA in three SGN Type
II cells. Scale bar, 20 μm.

Auditory thresholds and OHC function are similar between Spnb5$^{(-/-)}$ and

control mice.

To investigate how the loss of Spnb5 impacts auditory information processing, we

compared the *in vivo* ABRs of 2-month-old Spnb5$^{(+/+)}$ and Spnb5$^{(-/-)}$ mice. ABR

thresholds remained unaltered in Spnb5$^{(-/-)}$ mice across all sound frequencies

tested (**Figure 3A**). We also found that distortion product otoacoustic emissions

(DPOAEs), a measure of the robustness of OHC function, were unaffected by the

loss of Spnb5 (**Figure 3B**). These findings with preserved DPOAE thresholds also

extended to suprathreshold responses at sound intensities of 80-90 dB SPL

(**Figure 3C**). Since spectrins have been implicated in membrane integrity and

protein shuttling, we also assayed prestin expression in the lateral wall using its

gating charge movement (NLC) at postnatal day 13 (when prestin expression

increases in OHCs) and at two months (Abe et al., 2007). At both these time points,

we found NLC to be unaffected in Spnb5$^{(-/-)}$ mice (**Figure 4A**). Our measures of

the different parameters of NLC, Q_{sp}, V_h, and z, were similar between WT and

Spnb5$^{(-/-)}$ mice (**Figure 4, A-D**).

Figure 3

Figure 3. ABR analysis suggests a neuronal and not OHC dysfunction in Spnb5$^{(-/-)}$ mice. (A) ABR thresholds at frequencies between 2-32 kHz remain unchanged in Spnb5$^{(-/-)}$ mice. ABR thresholds in Spnb5$^{(-/-)}$ (open circle) and littermate WT (solid circle) mice at 2 months of age are plotted as a function of stimulus frequency. There are no significant differences between WT and Spnb5$^{(-/-)}$ mice at all frequencies tested from 2 to 32 kHz. (B) Similar 2f$_1$-f$_2$ DPOAE thresholds are observed in 2-month-old WT and Spnb5$^{(-/-)}$ mice at all f$_2$ frequencies tested. (C) 2f$_1$-f$_2$ DPOAE amplitudes are unaffected in Spnb5$^{(-/-)}$ mice. Input/output (I/O) function analysis of 2f$_1$-f$_2$ DPOAE responses in WT (solid circle) and Spnb5$^{(-/-)}$ (open circle) mice as a function of sound intensity at the f$_2$ frequency of 17.4 kHz at 2 months of age are graphed. DPOAE amplitudes in Spnb5$^{(-/-)}$ mice show no significant differences across all sound intensities compared to WT mice. (D) ABR wave I amplitudes are significantly altered between Spnb5$^{(-/-)}$ and WT at high stimulus intensities at frequencies between 5.6-32 kHz, but not at frequencies between 2-4 kHz. Shown are wave I amplitudes at 90 dB SPL. There is a consistent reduction in wave I amplitudes at these suprathreshold intensities from 5.6-32 kHz. (E) ABR wave I amplitudes at 16 kHz are significantly reduced in Spnb5$^{(-/-)}$ mice at high sound intensities (75-90 dB SPL). (F) ABR wave I latencies at 16 kHz are significantly delayed in Spnb5$^{(-/-)}$ mice at 90 dB SPL. Latencies at all other frequencies are not significantly delayed (data not shown). Data are means +/- SEM. The number (n) of animals per experiment is shown in each graph. Asterisks indicate p<0.05.

Figure 4

285
286

Figure 4. Spnb5$^{(-/-)}$ mice have similar nonlinear capacitance characteristics as controls at two age groups. NLC was measured at postnatal day 13, a midpoint of NLC development, and at two months when hearing is mature. **(A)** **There are no significant effects of OHC NLC in Spnb5$^{(-/-)}$ mice.** Shown are representative traces of NLC measurement in Spnb5$^{(-/-)}$ and WT mice. There are no differences in NLC between Spnb5$^{(-/-)}$ and WT mice at these two time points. **B, C, and D**. **NLC parameters are unaffected in Spnb5$^{(-/-)}$ mice.** Shown are means (+/- SEM) of NLC parameters in Spnb5$^{(+/+)}$ and Spnb5$^{(-/-)}$ mice at 13 days and two months, with no significant differences ($P > 0.05$) in all of these measures (Q_{sp}, V_h and z). **(B) Q_{sp}, a specific measure of net charge movement per unit of membrane corrected for linear capacitance. (C) V_h, a measure of the voltage dependence of charge movement. (D) z, a measure of the slope in NLC voltage dependence and indicative of the unitary charge movement.**

Loss of βV spectrin affects hearing sensitivity in a manner suggesting a

neural processing disorder.

The ABR is characterized by a series of electrical waves representing the

progressive transfer of the auditory signal from the periphery to the CNS (Song,

McGee, & Walsh, 2006). Wave I represents the summated response from the

spiral ganglion and auditory nerve (Melcher et al., 1996). While we see no

differences in ABR thresholds, we note that wave I amplitudes from 5.6 kHz and

upwards are significantly altered between Spnb5[(-/-)] and WT mice at high stimulus

intensities (suprathreshold) (**Figure 3D**). We also analyzed the latency of wave I

to investigate any potential changes in the rate of propagation of neural

transmission. Wave I latency at 16 kHz was progressively delayed at high sound

intensities reaching significance at the highest sound intensity (90 dB SPL) (**Figure**

3F). Latencies at other frequencies were not significantly different from WT.

Spnb5 plays a critical role in sensorineural processing.

What could be the cause of the differences in amplitudes of wave I between WT

and Spnb5[(-/-)] mice? We sought to exclude synaptopathy as a cause of the

decrease in wave I amplitudes. For instance, spectrins have been shown to be

important in targeting structural elements of the synaptic apparatus as well as

maintaining the integrity and compartmentalization of axons and dendrites

((Hammarlund, Jorgensen, & Bastiani, 2007), (Stankewich et al., 2010), (Efimova

et al., 2017; Stankewich et al., 2011), (Lorenzo et al., 2019). We compared the

number of IHC ribbon synapses in the basal, middle, and apical turn of the organ

of Corti between Spnb5$^{(-/-)}$ and WT mice. We used antibody labeling of the marker

protein CtBP2 (Bramhall, McMillan, Kujawa, & Konrad-Martin, 2018), a major

component of the ribbon synapses, and a well-defined marker of synaptopathy

(Sergeyenko, Lall, Liberman, & Kujawa, 2013). We found no differences in CtBP2-

labeled synaptic boutons of IHCs in the three cochlear turns between WT and

Spnb5$^{(-/-)}$ mice (**Figure 5A**). We then sought to identify morphological differences

in afferent nerve numbers as a possible cause of the reduced wave I amplitudes.

We found significant reductions in neurofilament labeled, nonmyelinated, afferent

fibers in the habenular in all three turns as well as efferent fibers in the basal and

middle turns of the cochlea (**Figure 5H, O**).

Figure 5

Figure 5. Spnb5$^{(-/-)}$ mice show evidence of neuropathy and no evidence of a synaptopathy. Shown are representative z axis projections of confocal micrographs of WT and βV spectrin KO mice in which synaptic boutons are labeled with antibodies to CtBP2 (**B-G**). **Afferent (I-N) and efferent (P-U) nerve fibers are labeled with antibodies to neurofilament.** The scale bar equates to 20 microns. There is a significant reduction in the density of afferent fibers at the habenular in all three turns and efferent fibers in the cochlear canal of Spnb5$^{(-/-)}$ mice. In contrast, there are no differences in CtBP2-positive bouton counts in all three turns of the cochlea. (**A, H, O) Shown are mean (+/- SEM) counts of synapses per IHC (A), afferent fibers (H), and efferent fibers (O) in all three cochlear turns.** The differences in afferent and efferent counts are significant (* P<0.05, n = 4 cochlea) between WT and Spnb5$^{(-/-)}$ mice in all three turns of the cochlea (afferent), and in the middle and basal turns of the cochlea (efferent). CtBP2 counts between WT and Spnb5$^{(-/-)}$ mice are not significantly different (P>0.05).

DISCUSSION

In this paper, we elucidate the function of βV spectrin in hearing using a mouse

gene deletion model. Removal of exon 2 containing the initiator methionine and an

N-terminal 37 amino acids, which, in part, compose its actin-binding domain,

results in a mouse with normal ABR thresholds, and normal measures of OHC

function including DPOAEs and NLC. Typical DPOAEs are a strong indication of

normal OHC electromotility (Shera & Guinan, 1999). The presence of normal NLC

measures further reiterates this interpretation. Moreover, we were unable to detect

Spnb5 mRNA in the organ of Corti or OHCs.

In contrast, we find a reduction in ABR wave I amplitudes that are concordant with

our finding of reduced numbers of nerve fibers. These data point to βV spectrin as

critical for the maintenance and function of afferent and efferent nerve fibers. The

normal ABR thresholds in the presence of reduced wave I amplitudes we observe

in mice have been seen previously in mutations of other proteins and, in particular,

in the recovery phase of noise-induced hearing loss (Maison, Pyott, Meredith, &

Liberman, 2013) (Kujawa & Liberman, 2009).

How do we explain the preserved ABR thresholds in the presence of reduced wave

I amplitudes? Loss of neurons of up to 80% has been shown to be associated with

normal thresholds as long as hair cells were functional (Schuknecht & Woellner,

1953). Although speculative, other authors have reasoned that a small increase in

sound pressure of 5 dB results in wider recruitment of neurons adjacent to the

index sound frequency that could compensate for a 50% loss in afferent neurons

using insensitive measures such as compound action potentials and ABRs

(Taberner and Liberman 2005), (Lin, Furman, Kujawa, & Liberman, 2011). An

important distinction in our data is that, unlike in noise-induced hearing loss

models, we see no differences in CtBP2 counts (Liberman & Kujawa, 2017). Our

CtBP2 counts and preliminary analysis using nerve counts suggest that the

functional effects in these mice lie in the nerve. Ongoing work in our lab includes

detailed analysis of high and low threshold fiber counts in these animals, together

with studies that will better determine the exact molecular mechanisms which

underlie the reduction in dendritic and axonal numbers.

We find absent effects on OHC function in $Spnb5^{(-/-)}$ mice that is consistent with

the undetectable expression of these transcripts in the cochlea and OHC in

particular. We note similar low/absent Spnb5 RNA expression upon interrogation

of both OHC single-cell RNA and OHC RNA expression datasets of adult mice that

have been published (Li et al., 2018) and gEAR portal (Shen, Scheffer, Kwan, &

Corey, 2015), (Liu, Chen et al. 2018), (Hoa et al., 2020). Together, these data

question the validity of βV spectrin antibody staining of OHCs, the principal reason

for ascribing its presence in OHCs (Legendre et al., 2008), (Cortese et al., 2017).

These papers led to the belief that βV spectrin had an integral role in the

cytoskeleton of hair cells. Our assertion that βV spectrin is not important for OHC

function is based, in part, on the absent phenotype in the knockout mice. It is

reasonable to ask if the knockout mouse resulted in an actual loss of function in

the gene. We feel confident that the deletion of the second exon in Spnb5$^{(-/-)}$

resulted in a silencing of the gene. In addition to the neuronal phenotype we

describe herein with the Spnb5$^{(-/-)}$ mice, we were also unable to detect Spnb5

transcript from the testis of knockout mice that may relate to degradation of

nonsense transcripts and a phenomenon observed in deletion of other genes

(Baker & Parker, 2004). RNAscope *in situ* hyridization data also show reduction in

the amount of detected mRNA in the SGN of KO animals. While the phenotype of

mice is consistent with the absent expression of Spnb5 transcript in OHCs, we

have to be mindful that normal OHC function in these mice could be due to other

beta spectrins substituting for Spnb5 spectrin. In this context, the presence of a

clear auditory phenotype in the Spnb2 spectrin knockout mice would suggest that

such an interchange and substitution by other spectrins is, however, not likely.

Moreover, Spnb4$^{(qv/qv)}$ spectrin has a neuronal phenotype, similar to Spnb5$^{(-/-)}$.

In conclusion, we present data in this study that βV spectrin plays an important role

in auditory function. Our data confirm that βV spectrin is necessary to maintain the

integrity or function of afferent and efferent nerves of the peripheral auditory

system. Importantly, our data show no evidence that βV spectrin has a function in

OHCs, as has been previously suggested.

MATERIALS AND METHODS

Mice.

All procedures conducted with mice were approved by the Institutional Animal Care

and Use Committee at the Yale University School of Medicine. C57BL/6J mice

were housed under pathogen-free conditions and maintained as a breeding

colony. In our studies, C57BL/6J mice were engineered to harbor a gene knockout

of Spnb5 (βV spectrin). The floxed exon 2 of Spnb5 was excised with Cre-

recombinase from embryonic stem (ES) cells in culture before injecting them into

mouse blastocysts under the guidance of the Yale Gene Targeting Center. This

construct was designed to create a constitutive loss of βV spectrin. For genotyping

animals, tail snips were collected from mice at their weaning age. Genomic DNA

was prepared by an overnight 56˚C incubation in tail lysis buffer 50mM Tris pH 8.0,

50mM KCL, 2.5mM EDTA, 0.4% Igepal, 0.4% Tween 20 supplemented with

Proteinase K (400 μg/ml). gDNA was assayed by PCR using the following primer

set: Forward 5'-CTGTCCGCCAAAGAGCAGGGGCCATG-3' Reverse 5'-

GATGTGGGGTCAGCCCAGCATCTCC-3' and PCR amplimers were visualized

on a 2% agarose gel.

OHC isolation.

Prestin-YFP mice of 2-3 months of age were euthanized, the cochleae extracted,

and the organ of Corti dissected in ice cold 1x DPBS. OHCs were dissociated by

enzymatic digestion with pre-warmed 0.05% trypsin for 8 minutes at 37˚C followed

by gentle trituration in 10% FBS in DMEM with a 1 mL pipette tip. The cell

suspension was spun at 1500 g for 10 minutes. The cell pellet was resuspended

in DMEM supplemented with 10% FBS and then filtered through a 40-micron cell

strainer. The cells were FACS sorted, and YFP-positive OHCs were collected.

mRNA detection.

Total RNA from mouse testis and the whole brain was either obtained from Takara

or prepared from 100 mg of testis tissue from 3-month old WT and βV KO

C57BL/6J animals with Trizol reagent (Chomczynski & Mackey, 1995). cDNA was

prepared with Superscript IV reverse transcriptase (ThermoFisher Scientific,

Waltham, MA,) and quantified with a Nanodrop spectrophotometer. Standard

procedures followed reagent manufacturer's directions for first-strand synthesis.

The organ of Corti and OHCs (prepared as described above) were isolated and

proteolyzed with 1 unit of proteinase K. After inactivating at 95°C for 5 minutes,

non-purified RNA was treated with Superscript IV to generate cDNA. Gene-specific

PCR primers bridging consecutive exons were designed to ensure the exclusion

of genomic DNA from amplification in **Table 1**.

RNAscope *in situ* Hybridization.

RNAscope *in situ* hybridization was performed on cochlear whole mount tissue

using RNAscope Multiplex Fluorescent v2 Assay (Advanced Cell Diagnostics

(ACD), Inc. Newark, CA, USA) according to the manufacturer's instructions

(Document # 323100-USM). RNAscope Probes and RNAscope Mulitplex

Fluorescent Reagent Kit v2 (Catalog # 323100) required for the assay were

purchased directly from ACD. Briefly, the cochlear apical turn of the organ of Corti

was dissected in 1x DPBS and fixed in 4% PFA. Cochlear tissue was pre-treated

with hydrogen peroxidase for 10 min at room temperature (RT), then with Protease

Plus for 15 min at 40°C in the HybEZ II Oven (ACD). Tissues were washed thrice

with distilled water for 5 min each at RT. The RNAscope Assay used the following

probes: target probe for Spnb5 (Catalog # 548431); positive control probe, Kcnma1

(Catalog #476251-C2); and negative control probe, DapB (Catalog #310043). For

SGNs the two probes used were Epha4 (Catalog #419081) and tyrosine

hydroxylase (Catalog #317612-C3). All probes were pre-warmed (10 min at 40°C),

brought to RT, then hybridized to the tissue with an 8 h incubation at 40°C. After

the tissue was stored in 5x saline sodium citrate (SCC) overnight at RT, hybridized

probe signal was amplified at 40°C with the consecutive addition of three

amplifiers: AMP1 (30 min), AMP2 (30 min), and AMP3 (15 min). HRP signal was

developed with a 15 min incubation in HRP-C1 at 40°C. For a single label

experiment, the TSA plus fluorophore CY3, diluted (1:50), (TSA Fluorescence Kit,

Catalog # NEL760001KT; PerkinElmer, Inc., MA, USA) was added for 30 min at

40°C, followed by a 15 min treatment with HRP blocker at 40°C. For double

fluorescent experiments, probes were developed using both HRP-C1 and HRP-C2

with alternating treatments of each fluorophore (fluorescein and CY3) and the HRP

blocker step. Between steps, tissue was washed twice with 1x Wash Buffer for 2

min each at RT. Finally, tissue was mounted with Prolong Gold antifade reagent

with DAPI (Thermo Fisher Scientific, OR, USA) on a glass bottom dish, and

fluorescent signal was visualized and imaged using a confocal microscope (Zeiss

Observer Z1).

Auditory brainstem responses (ABR).

ABR measurements were conducted in a manner similar to that described by Song

et. al. (El-Hassar et al., 2019). Briefly, animals were anesthetized with 480 mg/kg,

i.p., chloral hydrate, and all recordings were conducted in a sound-attenuating

chamber (Industrial Acoustics Corp., Bronx, NY). A customized TDT System 3

(Tucker-Davis Technologies Inc., Alachua, FL) was used for ABR recordings.

Subdermal needle electrodes (Rochester Electro-Medical, Lutz, FL) were

positioned at the vertex (active, noninverting), the right infra-auricular mastoid

region (reference, inverting), and the left neck region (ground). Differentially

recorded scalp potentials were bandpass filtered between 0.05 and 3 kHz over a

15 ms epoch. A total of 400 responses were averaged for each waveform for each

stimulus condition.

Symmetrically shaped tone bursts were 3 ms long (1 ms raised cosine on/off ramps

and 1 ms plateau). All acoustic stimuli were delivered free field via a speaker (EC1

Electrostatic Speaker, Tucker-Davis Technologies) positioned 10 cm from the

vertex. Stimulus levels were calibrated using a 0.5-inch condenser microphone

(model 4016, ACO Pacific, Belmont, CA) positioned at the approximate location of

the animal's head during recording sessions and are reported in decibels of sound

pressure level (SPL; referenced to 20 μPa). Stimuli of alternating polarity were

delivered at a rate of 21/s. Tone burst responses were collected in half-octave

steps ranging from 32 to 2.0 kHz. Stimuli were presented in 5 dB decrements of

sound intensity from a maximum level of 90 dB SPL. The ABR threshold was

defined as the lowest intensity of sound level capable of evoking a reproducible,

visually detectable response. Latencies and amplitudes of the initial ABR wave

were determined at a frequency of 16 kHz, one of the most sensitive frequency of

hearing in mice. Latencies and amplitudes of wave I at 90 dB SPL were also

measured at all the other frequencies. The analysis was performed in BioSigRP,

a digital signal processing software (TDT), on traces with discernible peaks by

setting cursors at the maxima and minima of the peaks. Latency was defined as

the time from the onset of the stimulus to the peak, while amplitude was measured

by taking the mean of the ΔV of the positive and negative deflections of the ABR

wave.

Distortion product otoacoustic emissions (DPOAEs).

The acoustic stimuli for DPOAEs were produced, and responses were recorded

using a customized TDT System 3 (Tucker-Davis Technologies Inc., Alachua, FL)

controlled by BioSigRP (TDT). To deliver acoustic stimuli and record the

responses, two speakers (MF1 Multi-Field Magnetic Speakers, TDT) and a

microphone probe (ER-10B+, Etymotic Research, Elk Grove Village, IL) were

connected by short PVC tubing, and the microphone probe was inserted into the

external auditory meatus of the anaesthetized mouse via a tapered plastic pipette

tip. Two simultaneous continuous pure tones (f_1 and f_2) that have a frequency ratio

of 1.2 (f_2/f_1) and equal sound level ($L_1 = L_2$) were delivered at center frequencies

of 8, 12, 16, 24 and 32 kHz with sound levels from 80 to 20 dB SPL in 10 dB

decrements. Stimuli duration was 83.88 ms at a repetition rate of 11.92 per second.

The acquired DPOAE responses were averaged 100 times. The DPOAE of interest

was at $2f_1$-f_2, the largest and most prominent DPOAE. The DPOAE threshold was

defined as the lowest sound intensity capable of evoking a visually detectable $2f_1$-

f_2 signal above the noise floor. DPOAE amplitudes were analyzed offline in

BioSigRP by setting cursors at the peak of the $2f_1$-f_2 signals.

Nonlinear capacitance (NLC).

Nonlinear capacitance was conducted in a fashion much like that previously

described (Song & Santos-Sacchi, 2016). C57BL/6J WT and βV spectrin KO

littermates at postnatal day 13 and at 2 months of age were compared. Apical turns

of the organ of Corti were isolated from each and recorded with ionic current

blocking solutions to minimize interference on measures of membrane

capacitance. The extracellular solution contained (in mM): 100 NaCl, 20

tetraethylammonium (TEA)-Cl, 20 CsCl, 2 $CoCl_2$, 1 $MgCl_2$, 1 $CaCl_2$, 10 HEPES,

pH 7.2. The intracellular solution contained (in mM): 140 CsCl, 2 $MgCl_2$, 10

HEPES, and 10 EGTA, pH 7.2. Pipettes had resistances of 3-5 MΩ. Gigohm seals

were made, and stray capacitance was balanced out with amplifier circuitry prior

to establishing whole-cell conditions. A Nikon Eclipse E600-FN microscope with

40× water immersion objective lens was used to observe cells during voltage

clamp. Whole-cell voltage-clamp recordings were performed from all three rows of

OHCs on whole-mount organ of Corti. OHCs were recorded at room temperature

using jClamp software and an Axopatch 200B amplifier (Molecular Devices, LCC,

San Jose, CA). Data were low pass filtered at 10 kHz and digitized at 100 kHz with

a Digidata 1320A (Molecular Devices, LCC, San Jose, CA). In order to extract

Boltzmann parameters, capacitance-voltage data were fit to the first derivative of

a two-state Boltzmann function.

540
541   $$C_m = NLC + C_{sa} + C_{lin} = Q_{max} \frac{ze}{k_B T} \frac{b}{(1+b)^2} + C_{sa} + C_{lin} \qquad \text{(eq.1)}$$
542
543
544   where $b = exp\left(-ze\frac{V_m - V_h}{k_B T}\right)$, $C_{sa} = \frac{\Delta C_{sa}}{(1+b^{-1})}$
545
Q_{max} is the maximum nonlinear charge moved, V_h is the voltage at peak

capacitance or equivalently, at half-maximum charge transfer, V_m is R_s-corrected

membrane potential, z is valence, C_{lin} is linear membrane capacitance, e is the

electron charge, kB is Boltzmann's constant, and T is the absolute temperature.

Immunohistochemistry.

For immunohistochemistry, staining was performed on fixed and decalcified three

turns of the organ of Corti to compare the number of neuronal fibers and synapses

between all three cochlear turns in KO and WT mice. Temporal bones were fixed

in 4% PFA overnight at 4°C and then decalcified in 120 mM

ethylenediaminetetraacetic acid (EDTA) for 2 days. Whole-mount of the organ of

Corti was dissected, washed in PBS 0.05% Tween 20, 0.05% Triton-X 100

(incubation buffer) three times for 3 minutes each, and incubated with primary

antibody in incubation buffer overnight at 4°C with mouse anti-CtBP2 antibody

(1:50, BD Transduction, San Jose, CA, cat# 612044), chicken anti-neurofilament

(NF) antibody (1:500, MilliporeSigma, Billerica, MA, cat# AB5539). After three

washes in incubation buffer, the tissue was incubated with secondary antibody in

incubation buffer for 1 hour at room temperature. The secondary antibodies used

included donkey anti-rabbit Alexa Fluor 568/647 (1:200; Thermo Fisher Scientific,

cat # A-10042/A-31573), goat anti-mouse Alexa Fluor 568/647 (1:200; Thermo

Fisher Scientific, cat # A-21124/A-21235) and goat anti-chicken Alexa Fluor

488/594 (1:200; Thermo Fisher Scientific, cat # A-11039/A-11042). After antibody

treatment, cells were washed in incubation buffer and mounted in Vectashield

(Vector Laboratories, Inc., Burlingame, CA), and viewed using a Zeiss confocal

laser-scanning microscope (Oberkochen, Germany).

Nerve (NF positive) and synaptic vesicle (CtBP2 positive) counts were performed

posthoc in Fiji (ImageJ). Briefly, z stacks of images were separated by probes,

background-subtracted (using a 50-pixel rolling ball radius), and z sections

containing the images of interest identified for further analysis (slice keeper).

CtBP2 labeled IHCs and NF labeled afferent fibers surrounding IHCs in proximity

to the habenula and efferent fibers in the cochlear canal were counted. Z

projections (sum) of the respective sections were obtained and automated

thresholds determined (max entropy). Counts of particles (synaptic vesicles)

immunopositive for CtBP2 larger than 0.1 microns were scored. Neuronal numbers

were established by counting the peaks in a profile plotted perpendicular to neural

fibers in the cochlea canal. For afferent nerve counts the profile was plotted at the

midpoint of nerve densities.

Statistical Analysis.

Data were analyzed using FIJI, Microsoft Excel, SigmaPlot and GraphPad Prism.

The measures of variability are depicted as error bars +/- the SEM and stated

numerically in the figure legends. Unpaired, one or two-tailed Student's t-test were

performed using both Excel and GraphPad Prism to make comparisons between

WT and Spnb5$^{(-/-)}$ data points. P-values are indicated in the figure legends to two

significant figures. The sample size (n) is stated in the figures.

COMPETING INTERESTS

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

MCS, JPB, JSM, DN and JSS conceptualized the project. PRS and JSM

generated the Spnb5 deleted mice. MCS and JPB conducted animal breeding,

tissue collection and genotyping. MCS, JPB, SK, WT, AS and DN performed gene

expression and immunohistochemistry analysis. JPB and LS performed assays to

assess hearing function. MCS and DN wrote the manuscript, and all authors

contributed to its editing and refinement.

ACKNOWLEDGEMENTS

These studies were supported by grants R01 DC 008130 and R01 DC 008130

from the National Institute of Health to JSS and DN and grants P01 NS35476

(project 4) and R01 HL28560 to JSM. Other support includes the Raymond Yesner

Endowment to Yale University (JSM). We also would like to thank Lan Ji for her

technical assistance.

REFERENCES

Abe, T., Kakehata, S., Kitani, R., Maruya, S., Navaratnam, D., Santos-Sacchi, J.,
      & Shinkawa, H. (2007). Developmental expression of the outer hair cell
      motor prestin in the mouse. *J Membr Biol, 215*(1), 49-56.
      doi:10.1007/s00232-007-9004-5

Bai, J. P., Surguchev, A., Ogando, Y., Song, L., Bian, S., Santos-Sacchi, J., &
      Navaratnam, D. (2010). Prestin surface expression and activity are
      augmented by interaction with MAP1S, a microtubule-associated protein.
      J Biol Chem, 285(27), 20834-20843. doi:10.1074/jbc.M110.117853

Baker, K. E., & Parker, R. (2004). Nonsense-mediated mRNA decay: terminating
      erroneous gene expression. *Curr Opin Cell Biol, 16*(3), 293-299.
      doi:10.1016/j.ceb.2004.03.003

Benson, D. A., Cavanaugh, M., Clark, K., Karsch-Mizrachi, I., Lipman, D. J.,
      Ostell, J., & Sayers, E. W. (2013). GenBank. *Nucleic Acids Res,
      41*(Database issue), D36-42. doi:10.1093/nar/gks1195

Bramhall, N. F., McMillan, G. P., Kujawa, S. G., & Konrad-Martin, D. (2018). Use
      of non-invasive measures to predict cochlear synapse counts. *Hear Res,
      370*, 113-119. doi:10.1016/j.heares.2018.10.006

Chomczynski, P., & Mackey, K. (1995). Short technical reports. Modification of
      the TRI reagent procedure for isolation of RNA from polysaccharide- and
      proteoglycan-rich sources. *Biotechniques, 19*(6), 942-945. Retrieved from
      https://www.ncbi.nlm.nih.gov/pubmed/8747660

Consortium, G. T. (2013). The Genotype-Tissue Expression (GTEx) project. *Nat
      Genet, 45*(6), 580-585. doi:10.1038/ng.2653

Cortese, M., Papal, S., Pisciottano, F., Elgoyhen, A. B., Hardelin, J. P., Petit, C., .
      . . El-Amraoui, A. (2017). Spectrin betaV adaptive mutations and changes
      in subcellular location correlate with emergence of hair cell electromotility
      in mammalians. *Proc Natl Acad Sci U S A, 114*(8), 2054-2059.
      doi:10.1073/pnas.1618778114

Cousin, M., Breau, K., Creighton, B.A., Spillmann, R.C., Torti, E., Dontu, S.,
      Tripathi, S., Ajit, D., Harper, K.M., Stankewich, M.C… Lorenzo, D.N.
      (2020). Pathogenic SPTBN1 variants cause a novel autosomal dominant

neurodevelopmental syndrome. medRxiv 2020.08.31.20184481; doi: https://doi.org/10.1101/2020.08.31.20184481

Efimova, N., Korobova, F., Stankewich, M. C., Moberly, A. H., Stolz, D. B., Wang, J., . . . Svitkina, T. (2017). betaIII Spectrin Is Necessary for Formation of the Constricted Neck of Dendritic Spines and Regulation of Synaptic Activity in Neurons. *J Neurosci, 37*(27), 6442-6459. doi:10.1523/JNEUROSCI.3520-16.2017

El-Hassar, L., Song, L., Tan, W. J. T., Large, C. H., Alvaro, G., Santos-Sacchi, J., & Kaczmarek, L. K. (2019). Modulators of Kv3 Potassium Channels Rescue the Auditory Function of Fragile X Mice. *J Neurosci, 39*(24), 4797-4813. doi:10.1523/JNEUROSCI.0839-18.2019

Hafidi, A., Despres, G., & Romand, R. (1990). Cochlear innervation in the developing rat: an immunocytochemical study of neurofilament and spectrin proteins. *J Comp Neurol, 300*(2), 153-161. doi:10.1002/cne.903000202

Hammarlund, M., Jorgensen, E. M., & Bastiani, M. J. (2007). Axons break in animals lacking beta-spectrin. *J Cell Biol, 176*(3), 269-275. doi:10.1083/jcb.200611117

Hoa, M., Olszewski, R., Li, X., Taukulis, I., Gu, S., DeTorres, A., . . . Kelley, M. W. (2020). Characterizing Adult Cochlear Supporting Cell Transcriptional Diversity Using Single-Cell RNA-Seq: Validation in the Adult Mouse and Translational Implications for the Adult Human Cochlea. *Front Mol Neurosci, 13*, 13. doi:10.3389/fnmol.2020.00013

Holley, M. C., & Ashmore, J. F. (1990). Spectrin, actin and the structure of the cortical lattice in mammalian cochlear outer hair cells. *J Cell Sci, 96 (Pt 2)*, 283-291. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/2211870

Holley, M. C., Kalinec, F., & Kachar, B. (1992). Structure of the cortical cytoskeleton in mammalian outer hair cells. *J Cell Sci, 102 (Pt 3)*, 569-580. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/1506434

Kujawa, S. G., & Liberman, M. C. (2009). Adding insult to injury: cochlear nerve degeneration after "temporary" noise-induced hearing loss. *J Neurosci, 29*(45), 14077-14085. doi:10.1523/JNEUROSCI.2845-09.2009

Legendre, K., Safieddine, S., Kussel-Andermann, P., Petit, C., & El-Amraoui, A. (2008). alphaII-betaV spectrin bridges the plasma membrane and cortical lattice in the lateral wall of the auditory outer hair cells. *J Cell Sci, 121*(Pt 20), 3347-3356. doi:10.1242/jcs.028134

Lewis, M. A., Nolan, L. S., Cadge, B. A., Matthews, L. J., Schulte, B. A., Dubno, J. R., . . . Dawson, S. J. (2018). Whole exome sequencing in adult-onset hearing loss reveals a high load of predicted pathogenic variants in known deafness-associated genes and identifies new candidate genes. *BMC Med Genomics, 11*(1), 77. doi:10.1186/s12920-018-0395-1

Li, Y., Liu, H., Giffen, K. P., Chen, L., Beisel, K. W., & He, D. Z. Z. (2018). Transcriptomes of cochlear inner and outer hair cells from adult mice. *Sci Data, 5*, 180199. doi:10.1038/sdata.2018.199

Liberman, M. C., & Kujawa, S. G. (2017). Cochlear synaptopathy in acquired sensorineural hearing loss: Manifestations and mechanisms. *Hear Res, 349*, 138-147. doi:10.1016/j.heares.2017.01.003

Lin, H. W., Furman, A. C., Kujawa, S. G., & Liberman, M. C. (2011). Primary neural degeneration in the Guinea pig cochlea after reversible noise-induced threshold shift. *J Assoc Res Otolaryngol, 12*(5), 605-616. doi:10.1007/s10162-011-0277-0

Liu, Y., Qi, J., Chen, X., Tang, M., Chu, C., Zhu, W., . . . Zhong, G. (2019). Critical role of spectrin in hearing development and deafness. *Sci Adv, 5*(4), eaav7803. doi:10.1126/sciadv.aav7803

Lorenzo, D. N., Badea, A., Zhou, R., Mohler, P. J., Zhuang, X., & Bennett, V. (2019). betaII-spectrin promotes mouse brain connectivity through stabilizing axonal plasma membranes and enabling axonal organelle transport. *Proc Natl Acad Sci U S A, 116*(31), 15686-15695. doi:10.1073/pnas.1820649116

Maison, S. F., Pyott, S. J., Meredith, A. L., & Liberman, M. C. (2013). Olivocochlear suppression of outer hair cells in vivo: evidence for combined action of BK and SK2 channels throughout the cochlea. *J Neurophysiol, 109*(6), 1525-1534. doi:10.1152/jn.00924.2012

Melcher, J. R., Knudson, I. M., Fullerton, B. C., Guinan, J. J., Jr., Norris, B. E., & Kiang, N. Y. (1996). Generators of the brainstem auditory evoked potential

in cat. I. An experimental approach to their identification. *Hear Res, 93*(1-
2), 1-27. doi:10.1016/0378-5955(95)00178-6

Papal, S., Cortese, M., Legendre, K., Sorusch, N., Dragavon, J., Sahly, I., . . . El-
Amraoui, A. (2013). The giant spectrin betaV couples the molecular
motors to phototransduction and Usher syndrome type I proteins along
their trafficking route. *Hum Mol Genet, 22*(18), 3773-3788.
doi:10.1093/hmg/ddt228

Parkinson, N. J., Olsson, C. L., Hallows, J. L., McKee-Johnson, J., Keogh, B. P.,
Noben-Trauth, K., . . . Tempel, B. L. (2001). Mutant beta-spectrin 4 causes
auditory and motor neuropathies in quivering mice. *Nat Genet, 29*(1), 61-
65. doi:10.1038/ng710

Schuknecht, H. F., & Woellner, R. C. (1953). Hearing losses following partial
section of the cochlear nerve. *Trans Am Laryngol Rhinol Otol Soc, 58*(57th
Meeting), 369-393; discussion, 393-364. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/13146770

Sergeyenko, Y., Lall, K., Liberman, M. C., & Kujawa, S. G. (2013). Age-related
cochlear synaptopathy: an early-onset contributor to auditory functional
decline. *J Neurosci, 33*(34), 13686-13694.
doi:10.1523/JNEUROSCI.1783-13.2013

Shen, J., Scheffer, D. I., Kwan, K. Y., & Corey, D. P. (2015). SHIELD: an
integrative gene expression database for inner ear research. *Database
(Oxford), 2015*, bav071. doi:10.1093/database/bav071

Shera, C. A., & Guinan, J. J., Jr. (1999). Evoked otoacoustic emissions arise by
two fundamentally different mechanisms: a taxonomy for mammalian
OAEs. *J Acoust Soc Am, 105*(2 Pt 1), 782-798. doi:10.1121/1.426948

Song, L., McGee, J., & Walsh, E. J. (2006). Frequency- and level-dependent
changes in auditory brainstem responses (ABRS) in developing mice. *J
Acoust Soc Am, 119*(4), 2242-2257. doi:10.1121/1.2180533

Song, L., & Santos-Sacchi, J. (2016). A Walkthrough of Nonlinear Capacitance
Measurement of Outer Hair Cells. *Methods Mol Biol, 1427*, 501-512.
doi:10.1007/978-1-4939-3615-1_28

Stabach, P. R., & Morrow, J. S. (2000). Identification and characterization of beta
742   V spectrin, a mammalian ortholog of Drosophila beta H spectrin. *J Biol*
743   *Chem, 275*(28), 21385-21395. doi:10.1074/jbc.C000159200

744   Stankewich, M. C., Cianci, C. D., Stabach, P. R., Ji, L., Nath, A., & Morrow, J. S.
745   (2011). Cell organization, growth, and neural and cardiac development
746   require alphaII-spectrin. *J Cell Sci, 124*(Pt 23), 3956-3966.
747   doi:10.1242/jcs.080374

748   Stankewich, M. C., Gwynn, B., Ardito, T., Ji, L., Kim, J., Robledo, R. F., . . .
749   Morrow, J. S. (2010). Targeted deletion of betaIII spectrin impairs
750   synaptogenesis and generates ataxic and seizure phenotypes. *Proc Natl*
751   *Acad Sci U S A, 107*(13), 6022-6027. doi:10.1073/pnas.1001522107

752   Yue, F., Cheng, Y., Breschi, A., Vierstra, J., Wu, W., Ryba, T., . . . Mouse, E. C.
753   (2014). A comparative encyclopedia of DNA elements in the mouse
754   genome. *Nature, 515*(7527), 355-364. doi:10.1038/nature13992
755
756

758 **Table1. Beta spectrin gene specific primers.**

Primers					Amplimer (bps)
Spnb1	forward	CAAAGAGCAGGAGGTGTCAGC	reverse	TCTAGGCAGGTGCTGAAGTTC	264
	nested forward	GATGCGTGTGCCGGGCGCC	nested reverse	CTGGTATTGATCTCTGCCT	197
Spnb2	forward	CCGGTGGGAGTGGTTAAGAC	reverse	CATCCCAGGTCGCTGCAGAC	188
	nested forward	GATTTTGGAGGTCCATCAG	nested reverse	TCAAACGCCTCATGGCGCT	144
Spnb5	forward	CTCCGGAGCTTTGAGCAGGCTGC	reverse	CCTGAGCCCACGCTAGCAATTCC	439
	nested forward	GATACAGGAGAAGACCACC	nested reverse	TTCCTGGACTTTGGCCTCC	319
Spnb5	forward	TGGCCTTCCTCAGGGCCAAGGT	reverse	GCACTCATCTGGATGCAGGGCTG	440
	nested forward	CTTGGACAGGGAGGAGTTTGG	nested reverse	GCACTCATCTGGATGCAGGGCTG	313

759
760 Primer sets were used to detect for mRNA of Spnb1, Spnb2, and Spnb5 in the
761 brain, organ of Corti, and isolated OHCs. The PCR amplimer sizes from mRNA
762 are presented in the final column.

764 **Table 2. Primers for the Spnb5 gene used to prepare the three DNA fragments**
765 **that were subcloned into pEasy-Flox.**

Fragment 1	forward	CAGCGGCCGCAGGGGCTGCCTCCTACTGTACC	reverse	GAGCGGCCGCACTTGAATGACCCAGAGCCACCCCT
Fragment 2	forward	GGTCTAGACCTCTTACTCGCTTAGGTTCCAATC	reverse	GCTCTAGATGACTCTACAGCAGACCCCCAGGTC
Fragment 3	forward	GACTCGAGCCTGTAGCTATTTCTGGCCTACCCCTCC	reverse	GACTCGAGGGAGGGAGGGTTACTGTAGTAGAAGGGC

766
The targeting vector for homologous recombination was constructed by cloning
three PCR generated fragments into the pEASY-Flox vector. Fragment 1 is a 4115
bp product, fragment 2 is a 469 bp product that contains Spnb5 exon 2, and
fragment 3 is 4371 bp of sequence that spans up to exon 5. The three fragments
were sequence verified. The oligos contain common restriction sites at their 5' end
for easy cloning into the targeting vector.

Table 3. Primers for rt-PCR in WT and Spnb5$^{(-/-)}$ tissue.

Spnb5	exon 58/60	forward	CTCCGGAGCTTTGAGCAGGCTGC	reverse	CCTGAGCCCACGCTAGCAATTCC	439
Spnb5	exon 53/55	forward	AGGAGAAACACCAGAATTTGGAGAATG	reverse	TGTCCACTCTCCAGAAGGGCAACTG	412
Spnb2	exon 28/29	forward	GATACAGGAGAAGACCACC	reverse	TTCCTGGACTTTGGCCTCC	144
Spna2	exon 54/55	forward	AACAGGAGTGACTGAGGAGG	reverse	CCTTCCTCAACCATTGGCAGA	144

775
776 These primer sets were tested on cDNA from testis isolated from WT and Spnb5$^{(-}$
777 $^{/-)}$ 3-month-old mice. Because the C-terminus of these large genes tends to be
778 more abundant than N-terminus exons, we designed primer sets that will amplify
779 exons near the poly-A tail.

780
781