1 Outer Hair Cell Function is Normal in βV Spectrin Knockout Mice.
2

3 Michael C. Stankewich[1*], Jun-Ping Bai[2*], Paul R. Stabach[1], Saaim Khan[3], Lei
4 Song[4], Jon S. Morrow[1,5], Joseph Santos-Sacchi[6,7,8], Dhasakumar S.
5 Navaratnam[2,6,7 ^]
6
7
8
9 1 Department of Pathology, Yale School of Medicine, New Haven, CT, U.S.A
10
11 2 Department of Neurology, Yale School of Medicine, New Haven, CT, U.S.A
12
13 3 Cranbrook Kingswood Upper School, MI, USA
14
15 4 Department of Otolaryngology-Head and Neck Surgery, Shanghai Ninth
16 People's Hospital, Shanghai Jiaotong University School of Medicine, Shanghai,
17 China
18
19 5 Department of Molecular, Cellular, and Developmental Biology, Yale University,
20 New Haven, CT, U.S.A
21
22 6 Department of Surgery, Yale School of Medicine, New Haven, CT, U.S.A
23
24 7 Department of Neurobiology, Yale School of Medicine, New Haven, CT, U.S.A
25
26 8 Department of Cellular and Molecular Physiology, Yale University, New Haven,

27 CT, U.S.A

28 * Co-first authors
29 ^ Corresponding Author
30
Address correspondence to Dr. Dhasakumar Navaratnam
Yale School of Medicine, Department of Neurology, PO Box 208018,
New Haven, CT., 06519
email: dhasakumar.navaratnam@yale.edu

Keywords: cochlea, auditory brainstem response, electromotility, distortion

product otoacoustic emissions, Spnb5, ribbon synapse, cochlear nerve

ABSTRACT

Reports have proposed a putative role for βV spectrin in outer hair cells (OHCs) of the cochlea. In an ongoing investigation of the role of the cytoskeleton in electromotility, we tested mice with a targeted exon deletion of βV spectrin (Spnb5), and unexpectedly find that Spnb5$^{(-/-)}$ animals' auditory thresholds are unaffected. Similarly, these mice have normal OHC electromechanical activity (otoacoustic emissions) and non-linear capacitance. Moreover, Spnb5 mRNA is undetectable in the organ of Corti or OHCs. In contrast, magnitudes of auditory brainstem response (ABR) peak 1-amplitudes are significantly reduced. Evidence of a synaptopathy was absent with normal hair cell CtBP-2 counts. In Spnb5$^{(-/-)}$ mice, the number of afferent and efferent nerve fibers is decreased. Taken together, these data establish that βV spectrin is important for hearing, affecting neuronal structure and function. Significantly, these data exclude βV spectrin as functionally important to OHCs as has been previously suggested.

INTRODUCTION

In the early 1990s, several groups made key seminal observations of a spectrin/actin lattice in the cochlear outer hair cells (OHCs) (Holley & Ashmore, 1990) (Holley, Kalinec, & Kachar, 1992). Their electron micrographs propelled an exquisite model of the spectrin/actin lattice assembled just beneath the OHC lateral membrane with longitudinal spectrin molecules linked to circumferential actin filaments that, in turn, were linked to the plasma membrane by electron-dense pillars. Their work also identified a pool of spectrin in the actin-rich cuticular plate. Another study evaluated spectrin proteins in cochlear innervation during rodent development (Hafidi, Despres, & Romand, 1990). They identified differences in developmental expression of the erythroid beta spectrin (Spnb1 spectrin) compared to the most ubiquitous nonerythroid form $\alpha II/\beta II$ spectrin, similar to what is observed in the brain. What is not certain from these earlier studies is the specific spectrins involved, a family now known to consist of two alpha spectrins (Spna1 and 2) and five beta spectrins (Spnb1-5). More recently, it has been shown that Spnb4 is restricted to the nodes of Ranvier in myelinated axons and is also a deafness-related gene (Parkinson et al., 2001) (Lewis et al., 2018). It appears from these studies that the deafness in the quivering mouse is likely not originating in the cochlea but rather is central in origin. Furthermore, Spnb2 spectrin, when selectively deleted from the mouse cochlea sensory epithelia by crossing Spnb2 floxed mice with Atoh-1 Cre-recombinase, causes profound deafness due to a disruption of the stereocilia rootlets in the cuticular plate in hair cells (Liu et al., 2019). Thus, the function of specific spectrins,

particularly as it relates to the OHC lateral membrane and cochlear innervation, remains unclear. In humans, the molecule βV spectrin (SPTBN5) that was first discovered in the retina, is involved with the retinal disorder, Usher syndrome (Stabach & Morrow, 2000) (Papal et al., 2013). Further reports have proposed a putative role for βV spectrin in a second sensorineural function, hearing. One group's observations demonstrated that antisera raised against the C-terminal peptide of SPTBN5, the largest form of spectrin, and an orthologue of beta(Heavy) spectrin in Drosophila, labeled structures along the lateral membrane of the OHC (Legendre, Safieddine, Kussel-Andermann, Petit, & El-Amraoui, 2008) (Cortese et al., 2017). This observation drew attention because detection of this giant spectrin in tissue other than retina and testis has been elusive. This finding raised the possibility that βV spectrin was involved in the electromotile properties of OHCs.

Continuing an ongoing effort in our lab to define the cytoskeleton's role in electromotility, we sought to determine the role of βV spectrin in OHC function (Bai et al., 2010). In this study, we analyzed βV spectrin expression in the cochlea and addressed its function in mice with a targeted deletion of Spnb5. We tested hearing by auditory brainstem response (ABR), measured OHC-mediated otoacoustic emissions and OHC nonlinear capacitance (NLC), and evaluated the number of inner hair cell (IHC) ribbon synapses and nerve fibers that innervate the cochlear sensory epithelium. We find that while OHC function is unaffected in the absence of βV spectrin, nerve counts are reduced at the habenular, in all three turns of the cochlea, and in efferent fibers in the mid and basal turns, which collectively account for the reduced wave I amplitudes in ABRs. We conclude that βV spectrin is

100   necessary to maintain the integrity or function of afferent and efferent nerves of the

101   peripheral auditory system, but plays no role in the OHC itself.

 RESULTS
105

Mouse βV spectrin amino acid sequence is divergent from the human

protein.

Surprisingly, a sequence alignment comparing the mouse amino acid sequence to

the human amino acid sequence reveals a significant divergence. In contrast, other

beta spectrins, including beta I-IV, have ~ 90% amino acid conservation between

human and mouse sequences. By aligning human and mouse βV spectrin, we note

a significant divergence at the N- and C-termini of the proteins and an internal

stretch of amino acids from position 817-922 (**Figure 1A**). Of note is that the

initiator methionine residue in the human form is conserved between mouse and

human genes and lies at position 36 and within exon 2 (**Figure 2**). The mouse

sequence incorporates an unusual homorepeat of glutamines and, notably, does

not have a bona fide pleckstrin homology (PH) domain seen in the human form.

Spnb5 mRNA tissue expression.

Spnb5 spectrin is expressed in most mouse tissues but at low levels. In ENCODE,

the level of Spnb5 transcripts is an order of magnitude less abundant than the

ubiquitous form Spnb2 (**Figure 1B**) (Yue et al., 2014). The peak abundance of

Spnb2 approaches 50 RPKM while Spnb5 is 0.5 RPKM in testis (**Figure 1B, C**).

Interestingly, the GTEX dataset shows high expression of Spnb5 in the peripheral

(tibial) nerve (**Figure 1D**). To validate that mRNA is detected in mice, we tested

RNA from the testis with primer sets that bridged exons. By standard rt-PCR,

Spnb5 mRNA fragments encoding both the N-terminus and C-terminus were

amplified from testis and brain RNA, and its identity was confirmed by Sanger

sequencing. In contrast to brain and testicular tissue, we could not detect Spnb5

mRNA in isolated OHCs and whole organ of Corti by both conventional and nested

PCR (Table 1 details the gene-specific primer sets). Using rt-PCR from ~3000

OHCs and organ of Corti, we detect both Spnb1 and Spnb2. Using nested PCR,

amplicons for these spectrins are robust, while we do not visualize any product for

Spnb5 (**Figure 1E**). These data are in accord with data derived from the public

data sets of transcriptome profiling for OHCs in gEAR (https://umgear.org/) (**See**

supplementary Figure1).

A

B

Spnb2 tissue expression: ENCODE

C

Spnb5 tissue expression: ENCODE

D SPTBN5 tissue expression: GTEX

E

Figure 1

Figure 1. (A) Spnb5 amino acid sequence differs from humans. Shown is the comparison of the deduced amino acid sequence of mouse and human Spnb5. Conserved amino acids are marked in red. There is divergence at the N and C termini. In addition, a region in the middle of the protein from posiiton 817-922 shows significant divergence. The mouse version of this gene shares only 65.6 percent identity with the human form. (B, C) Spnb5 is expressed in brain and testis. The relative abundance of Spnb2 transcripts as compared to Spnb5 in mouse tissues. Shown are the expression of Spnb2 and 5 in RNA Seq. NCBI reference datasets (ENCODE; Bioproject PRJNA66167 (Yue et al., 2014). On average, the abundance of Spnb5 mRNA is an order of magnitude less abundant than Spnb2. Of the tissues examined, adult testis has the most mRNA for Spnb5. (D) In human, Spnb5 is most expressed in peripheral nerve. The relative abundance of Spnb5 in different tissues in GTEX database is presented (version 6) (Consortium, 2013). Of all the tissues examined Spnb5 is expressed in highest amounts in the peripheral nerve (tibial nerve). (E) Spnb5 mRNA is not detectable in the organ of Corti or OHCs. Shown are PCR amplifications from mouse brain (lanes 2-4), organ of Corti (lanes 6-8) and OHCs (lanes 10-12) of Spnb1 (lanes 2, 6 and 10), Spnb2 (lanes 3, 7 and 11) and Spnb5 (lanes 4, 8 and 12). The upper panel shows amplicons for these three mRNA transcripts using outer primers while the lower panel shows amplicons of primers using a nested set of primers. All three spectrins are detectable in brain. Spnb1 and 2 are detectable in both the organ of Corti and OHCs (lanes 6-7 and 10-11). Spnb5 however, is not detectable using either primer set (lanes 8 and 12).

139 **Supplementary Figure 1.**

140
141 **Expression levels for each of the beta spectrins in OHCs.**
142 Mining data from the gene Expression Analysis Resource (gEAR) we graphed the
143 abundance of each of the five beta spectrin genes. This data set was derived from
144 transcriptome analysis using sc-RNAseq of cell populations isolated from mouse
145 cochleae (Li et al., 2018).
146
Targeted knockout of Spnb5 in mice using homologous recombination.

In order to properly target the βV spectrin gene in mice, we needed to know its

precise gene structure. While the human gene has been well characterized and

indexed since its discovery in 2000 (Stabach & Morrow, 2000), there has been

ambiguity in the intron/exon usage in the mouse gene found on chromosome 2.

The GENBANK entry for Spnb5 has been modified many times in the last several

years, with the last significant annotation in 2007 (Benson et al., 2013). This

annotation was derived by automated computational analysis using gene

prediction algorithms. The exon prediction is made difficult because 1) the gene is

very large, so PCR amplifying a full-length cDNA is not trivial, and 2) the

conservation between human and mouse intron/exon boundaries is weak at the N-

and C-termini. The current and accepted annotated version in both the UCSC

genome browser and in Ensembl shows Spnb5 spectrin mRNA in the mouse to be

14.4 kb with a 10,872 bp ORF that has 66 exons and codes for 3623 amino acids:

NM_001370938: NP_001357867.1. This version has been validated in the

ENCODE project based on the robustness of next-generation sequencing efforts

over the last several years (Yue et al., 2014).

Shown in the schematic in **Figure 2A** is the strategy for targeted deletion of exon

2 of Spnb5. The knockout strategy was based on the transcript ID:

ENSMUST00000156159.3, and a targeting construct was generated in which

three Lox P sites were inserted to flank the neomycin selection marker and exon

2 containing the start codon, resulting in disruption of the open reading frame upon

Cre-mediated excision. Exon 2 contains the ATG start codon (with a consensus

Kozak sequence) and an additional 37 amino acids at its N-terminus. Notably, the

codon for the methionine residue corresponding to the human start codon is also

contained within the second exon. A genomic bacterial artificial chromosome

containing the Spnb5 locus was obtained, and genomic DNA fragments containing

the first exons of Spnb5 were subcloned in pEasy (a gift from Sankar Ghosh). The

targeting vector comprised a 4.3 kb long arm, a 4.1 kb short arm, and a middle

selection cassette which contains exon two flanked by a PGK–neomycin

resistance (Neo), two FRT sites, and the herpes simplex thymidine kinase gene

(**Figure 2B and Primer Sets in Table 2**). The targeting plasmid DNA was

linearized with Cla1 and electroporated into mouse embryonic stem cells (ESCs)

(129SV/EV). ESCs were selected for neomycin/G418 resistance. Positive clones

were identified by PCR and confirmed by Southern blot analysis using external

and internal probes. A correctly targeted ESC clone was injected into C57BL/6J

blastocysts to produce germline transmitting chimeric mice. These were then

mated with actin-Cre C57BL/6J transgenic mice for generation of heterozygous

Spnb5 transgenic mice with both the Neo cassette and exon two removed

(Spnb5$^{(+/-)}$). The resulting Spnb5$^{(+/-)}$ offspring were crossed to generate βV

spectrin-deficient mice. The genotypes of these mice were confirmed by PCR

amplification (**Figure 2C**). Chimeric mice were backcrossed onto the C57BL/6J

strain. Litters from heterozygous breeders Spnb5$^{(+/-)}$ / Spnb5$^{(+/-)}$ were of normal

number, and the mice of all genotypes were phenotypically indistinguishable with

normal lifespans >2 yrs old. We confirmed decreased expression of Spnb5 in

cDNA from testes of Spnb5$^{(-/-)}$ mice, where we found equivalent expression of

Spnb2 and Spna2 (**Figure 2D and Primer Sets in Table 3**).

A.

B.

Exon1 (4.1kb)

Intron/Exon2/Intron (469bp)

pEASY F1 F2 F3

Intron / Exon 3 / Intron / Exon 4/ Intron / Exon 5 (4.3kb)

C.

D

| Spnb5 | | | | | Spnb2 | | | Spna2 | | |
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 |

Figure 2. Exon 2 of the Spnb5 gene was removed using homologous recombination. (A) Schematic of the strategy used to delete exon 2. The construct contained a short arm of exon 1 and a long arm of exons 3, 4, and 5 together with the intervening introns. Both the selection marker Neo and exon two are flanked by Lox P sites. Linearized plasmid was injected into ES cells and clones with homologous recombination and selected by neomycin (Neo)/ G418 resistance. Mice were then crossed with Actin-Cre mice to generate floxed mice lacking the Neo cassette and exon 2. **(B) Vector map of pEASY used to target Spnb5 deletion.** Exon 2 was subcloned between two Lox P sites adjacent to the Neo gene. These two genomic fragments were in turn placed between a 4.1kb short arm containing Exon 1 and a 4.3kb long arm containing exons 3, 4, and 5 and the intervening introns. The vector also contains a thymidine kinase and ampicillin gene allowing for selection. The plasmid was linearized with Cla1. **(C) PCR amplification of genomic DNA confirming deletion of exon two and flanking intronic sequence.** In genomic DNA from Spnb5$^{(-/-)}$ mice, primers that amplify a product no longer contain part of intron 1; exon 2; and part of intron 2/3. Lane 1 base pair standards, lane 2 Spnb5$^{(+/-)}$, lane 3 Spnb5$^{(-/-)}$, lane 4 Spnb5$^{(+/+)}$. The larger fragment of 665 bp consists of unexcised exon two while the smaller fragment consists of genomic DNA fragment from which exon two has been excised. **(D) Decreased expression of Spnb5 in Spnb5$^{(-/-)}$ testicular cDNA compared to Spnb2 and Spna2.** Spnb5 exons 58-60 in lanes 2-3, Spnb5 exon 53-55 in lanes 4-5; Spnb2 in lanes 7-8 (exons 28-29) and Spna2 (exons 54-55) in lanes 10-11. WT in lanes 2, 4, 7 and 10. Spnb5$^{(-/-)}$ in lanes 3, 5, 8 and 11.

Figure 2

Auditory thresholds and OHC function are similar between Spnb5$^{(-/-)}$ and

control mice.

To investigate how the loss of Spnb5 impacts auditory information processing, we

compared the *in vivo* ABR of 2-month-old Spnb5$^{(+/+)}$ and Spnb5$^{(-/-)}$ mice. ABR

thresholds were not altered in Spnb5$^{(-/-)}$ mice at all sound frequencies tested from

2 to 32 kHz (**Figure 3A**). We also found that distortion product otoacoustic

emissions (DPOAE) measurements, a measure specific to OHC function, are not

affected by the loss of βV spectrin (**Figure 3B**). These findings with preserved

DPOAE thresholds also extended to suprathreshold responses at sound

intensities of 80-90 dB SPL (**Figure 3C**). Since spectrins have been implicated in

membrane integrity and protein shuttling, we also assayed prestin expression in

the lateral wall using its gating charge movement (NLC) at postnatal day 13 (when

prestin expression increases in OHCs) and at two months (Abe et al., 2007). At

both these time points, we found NLC to be unaffected in Spnb5$^{(-/-)}$ mice (**Figure**

4A). Our measures of the different parameters of NLC, Q_{sp}, V_h, and z, were similar

between WT and Spnb5$^{(-/-)}$ mice (**Figure 4, A-D**).

Figure 3

214
215
216
Figure 3. ABR analysis suggests a neuronal and not OHC dysfunction in Spnb5$^{(-/-)}$ mice. (A)
thresholds at frequencies from 2-32 kHz reveal no change in Spnb5$^{(-/-)}$ mice. ABR thresholds
in Spnb5$^{(-/-)}$ (open circle) and littermate WT (solid circle) mice at 2-month-old are plotted over a
range of frequencies. There was not a significant difference between WT and Spnb5$^{(-/-)}$ mice at all
frequencies tested from 2 to 32 kHz. **(B) Similar DPOAE thresholds were observed in 2-month-**
old WT and Spnb5$^{(-/-)}$ mice at all frequencies. **(C) DPOAE input/output function are unaffected**
in Spnb5$^{(-/-)}$ mice. Input/output (I/O) function analysis of DPOAE responses in WT (solid circle) and
Spnb5$^{(-/-)}$ (open circle) mice as a function of sound intensity at 16 kHz at two months of age are
graphed. DPOAE amplitudes in Spnb5$^{(-/-)}$ mice showed no significant difference across all sound
intensities compared to WT mice. **(D) Wave I amplitudes are significantly altered between**
Spnb5$^{(-/-)}$ and WT at high stimulus intensities at frequencies 5.6-32 kHz, but not at
frequencies 2-4 kHz. Shown is the wave I amplitudes at 90 dB SPL. There is a consistent reduction
in the amplitudes of wave I amplitudes at these suprathreshold intensities from 5.6-32 kHz. **(E)**
Wave I amplitudes were significantly reduced in Spnb5$^{(-/-)}$ mice at higher sound intensities
(75-90 dB SPL). (F) Wave I latencies at 16 kHz show significant delays at 90 dB SPL. Latencies
at all other frequencies are not significantly delayed (data not shown). Data are means +/- SEM.
The number (n) of animals per experiment is shown in each graph. Asterisks indicate $P<0.05$.
234

Figure 4

Figure 4. Spnb5$^{(-/-)}$ mice have similar nonlinear capacitance characteristics as controls at two age groups. NLC was measured at postnatal day 13, a midpoint of NLC development, and at two months when hearing is mature. **(A**) **There are no significant effects of OHC NLC in Spnb5$^{(-/-)}$ mice.** Shown are representative traces of NLC measurement in Spnb5$^{(-/-)}$ and WT mice. There are no differences in NLC between Spnb5$^{(-/-)}$ and WT mice at these two time points. **B, C, and D**. **NLC parameters are unaffected in Spnb5$^{(-/-)}$ mice.** Shown are means (+/- SEM) of NLC parameters in Spnb5$^{(+/+)}$ and Spnb5$^{(-/-)}$ mice at 13 days and two months, with no significant differences (P>0.05) in all of these measures (Q$_{sp}$, V$_h$ and z). **(B) Q$_{sp}$, a specific measure of net charge movement per unit of membrane corrected for linear capacitance. (C) V$_h$, a measure of the voltage dependence of charge movement. (D) z, a measure of the slope in NLC voltage dependence and indicative of the unitary charge movement.**

Loss of βV spectrin affects hearing sensitivity in a manner suggesting a neural processing disorder.

The ABR is characterized by a series of electrical waves representing the progressive transfer of the auditory signal from the periphery to the CNS (Song, McGee, & Walsh, 2006). Wave I represents the summated response from the spiral ganglion and auditory nerve (Melcher et al., 1996). While we see no differences in ABR thresholds, we note that Wave I amplitudes from 5.6 kHz and upwards are significantly altered between Spnb5$^{(-/-)}$ and WT at high stimulus intensities (suprathreshold). We also analyzed the latency of wave I to investigate any potential changes in the rate of propagation of neural transmission (**Figure 5**). Wave I latency at 16 kHz was progressively delayed at high sound intensities reaching significance at the highest sound intensity (90 dB SPL). Latencies at other frequencies were not significantly different from WT.

Spnb5 plays a critical role in sensorineural processing.

What could be the cause of the differences in amplitudes of wave I between WT and Spnb5$^{(-/-)}$ mice? We sought to exclude synaptopathy as a cause of the decrease in wave I amplitudes. For instance, spectrins have been shown to be important in targeting structural elements of the synaptic apparatus as well as maintaining the integrity and compartmentalization of axons and dendrites ((Hammarlund, Jorgensen, & Bastiani, 2007), (Stankewich et al., 2010), (Efimova et al., 2017; Stankewich et al., 2011), (Lorenzo et al., 2019). We compared the numbers of ribbon synapses at the base, middle, and apex of the organ of Corti

between Spnb5$^{(-/-)}$ and WT mice. We used antibody labeling of the marker protein

CtBP-2 (Bramhall, McMillan, Kujawa, & Konrad-Martin, 2018), a major component

of the ribbon synapses, and a well-defined marker of synaptopathy (Sergeyenko,

Lall, Liberman, & Kujawa, 2013). We found no differences in CtBP-2 labeled

synaptic boutons of IHC in the three cochlea turns between WT and Spnb5$^{(-/-)}$ mice

(**Figure 5I**). We then sought to identify morphological differences in afferent nerve

numbers as a possible cause of the reduced wave I amplitudes. We found

significant reductions in neurofilament labeled, nonmyelinated, afferent fibers in

the habenular in all three turns as well as efferent fibers in the basal and middle

turns of the cochlea (**Figure 5G, H**).

Figure 5

Figure 5. Spnb5$^{(-/-)}$ mice show evidence of neuropathy and no evidence of a synaptopathy.
Shown are representative z axis projections of confocal micrographs of WT and βV spectrin KO
mice in which synaptic boutons are labeled with antibodies to CtBP-2 (**B-G**). **Afferent (I-N) and**
efferent (P-U) nerve fibers are labeled with antibodies to neurofilament. The scale bar

equates to 20 microns. There is a significant reduction in the density of afferent fibers at the
habenular in all three turns and efferent fibers in the cochlear canal of Spnb5[(-/-)] mice. In contrast,
there were no differences in CtBP-2 positive bouton counts in all three turns of the cochlea. (**A, H,**
O) Shown are the mean (+/- SEM) synaptic (A) counts, afferent (H) and efferent (O) neuronal
and in all three turns of the cochlea. The differences in afferent and efferent counts were
significant (* P<0.05, n = 4 cochlea) between WT and Spnb5[(-/-)] mice in all three turns of the cochlea
(afferent), and in the middle and basal turns of the cochlea (efferent). CtBP-2 counts between WT
and Spnb5[(-/-)] mice were not significantly different (P>0.05).
295

DISCUSSION

In this paper, we elucidate the function of βV spectrin in hearing using a mouse

gene deletion model. Removal of exon 2 containing the initiator methionine and an

N-terminal 37 amino acids, which, in part, compose its actin-binding domain,

results in a mouse with normal ABR thresholds, and normal measures of OHC

function including DPOAE and NLC. Typical DPOAEs are a strong indication of

normal OHC electromotility (Shera & Guinan, 1999). The presence of normal NLC

measures further reiterates this interpretation. Moreover, we were unable to detect

Spnb5 mRNA in the organ of Corti or OHCs.

In contrast, we find a reduction in wave I amplitudes that are concordant with our

finding of reduced numbers of nerve fibers. These data point to βV spectrin as

critical for the maintenance and function of afferent and efferent nerve fibers. The

normal ABR thresholds in the presence of reduced wave I amplitudes we observe

in mice have been seen previously in mutations of other proteins and, in particular,

in the recovery phase of noise-induced hearing loss (Maison, Pyott, Meredith, &

Liberman, 2013) (Kujawa & Liberman, 2009).

How do we explain the preserved ABR thresholds in the presence of reduced wave

I amplitudes? Loss of neurons of up to 80% has been shown to be associated with

normal thresholds as long as hair cells were functional (Schuknecht & Woellner,

1953). Although speculative, other authors have reasoned that a small increase in

sound pressure of 5 dB results in wider recruitment of neurons adjacent to the

index sound frequency that could compensate for a 50% loss in afferent neurons

using insensitive measures such as compound action potentials and ABRs

(Taberner and Liberman 2005), (Lin, Furman, Kujawa, & Liberman, 2011). An

important distinction in our data is that, unlike in noise-induced hearing loss

models, we see no differences in CtBP-2 counts (Liberman & Kujawa, 2017). Our

CtBP-2 counts and preliminary analysis using nerve counts suggest that the

functional effects in these mice lie in the nerve. Ongoing work in our lab includes

detailed analysis of high and low threshold fiber counts in these animals, together

with studies that will better determine the exact molecular mechanisms which

underlie the reduction in dendritic and axonal numbers.

We find absent effects on OHC function in Spnb5[(-/-)] mice that is consistent with

the undetectable expression of these transcripts in the cochlea and OHC in

particular. We note similar low/absent Spnb5 RNA expression upon interrogation

of both OHC single-cell RNA and OHC RNA expression datasets of adult mice that

have been published (Li et al., 2018) and gEAR portal (Shen, Scheffer, Kwan, &

Corey, 2015), (Liu, Chen et al. 2018), (Hoa et al., 2020). Together, these data

question the validity of βV spectrin antibody staining of OHCs, the principal reason

for ascribing its presence in OHCs (Legendre et al., 2008), (Cortese et al., 2017).

These papers led to the belief that βV spectrin had an integral role in the

cytoskeleton of hair cells. Our assertion that βV spectrin is not important for OHC

function is based, in part, on the absent phenotype in the knockout mice. It is

reasonable to ask if the knockout mouse resulted in an actual loss of function in

the gene. We feel confident that the deletion of the second exon in Spnb5$^{(-/-)}$

resulted in a silencing of the gene. In addition to the neuronal phenotype we

describe herein with the Spnb5$^{(-/-)}$ mice, we were also unable to detect Spnb5

transcript from the testis of knockout mice that may relate to degradation of

nonsense transcripts and a phenomenon observed in deletion of other genes

(Baker & Parker, 2004). While the phenotype of mice is consistent with the absent

expression of Spnb5 transcript in OHCs, we have to be mindful that normal OHC

function in these mice could be due to other beta spectrins substituting for Spnb5

spectrin. In this context, the presence of a clear auditory phenotype in the Spnb2

spectrin knockout mice would suggest that such an interchange and substitution

by other spectrins is, however, not likely. Moreover, Spnb4$^{(qv/qv)}$ spectrin has a

neuronal phenotype, similar to Spnb5$^{(-/-)}$.

In conclusion, we present data in this study that βV spectrin plays an important role

in auditory function. Our data confirm that βV spectrin is necessary to maintain the

integrity or function of afferent and efferent nerves of the peripheral auditory

system. Importantly, our data show no evidence that βV spectrin has a function in

OHCs, as has been previously suggested.

MATERIALS AND METHODS

Mice.

All procedures conducted with mice were approved by the Institutional Animal Care and Use Committee at the Yale University School of Medicine. C57BL/6J mice were housed under pathogen-free conditions and maintained as a breeding colony. In our studies, C57BL/6J mice were engineered to harbor a gene knockout of Spnb5 (βV spectrin). The floxed exon 2 of Spnb5 was excised with Cre-recombinase from embryonic stem (ES) cells in culture before injecting them into mouse blastocysts under the guidance of the Yale Gene Targeting Center. This construct was designed to create a constitutive loss of βV spectrin. For genotyping animals, tail snips were collected from mice at their weaning age. Genomic DNA was prepared by an overnight 56°C incubation in tail lysis buffer 50mM Tris pH 8.0, 50mM KCL, 2.5mM EDTA, 0.4% Igepal, 0.4% Tween20 supplemented with Proteinase K (400 μg/ml). gDNA was assayed by PCR using the following primer set: Forward 5'-CTGTCCGCCAAAGAGCAGGGGCCATG-3' Reverse 5'-GATGTGGGGTCAGCCCAGCATCTCC-3' and PCR amplimers were visualized on a 2% agarose gel.

Organ of Corti and OHC isolation.

Prestin-YFP mice that were 2-3 months old were euthanized and the cochleae extracted. The organ of Corti was dissected and treated with pre-warmed 0.05% trypsin for 8 minutes at 37°C. The organ of Corti was dissociated by triturating the tissue 20-30 times in a 1 ml pipet tip. The cell suspension was spun at 1500 g for

10 minutes. The cell pellet was resuspended in DMEM supplemented with 10%

FBS and then filtered through a 40-micron cell strainer. The cells were FACS

sorted, and YFP positive OHCs were collected.

mRNA detection.

Total RNA from mouse testis and the whole brain was either obtained from Takara

or prepared from 100 mg of testis tissue from 3 month WT and βV KO C57BL/6J

animals with Trizol reagent (Chomczynski & Mackey, 1995). cDNA was prepared

with Superscript IV reverse transcriptase (ThermoFisher Scientific, Waltham, MA,)

and quantified with a Nanodrop spectrophotometer. Standard procedures followed

reagent manufactures directions for first-strand synthesis. The organ of Corti and

OHCs (prepared as described above) were isolated and proteolyzed with 1 unit of

proteinase K. After inactivating at 95˚C for 5 minutes, non-purified RNA was

treated with Superscript IV to generate cDNA. Gene-specific PCR primers bridging

consecutive exons were designed to ensure the exclusion of genomic DNA from

amplification in **Table 1**.

Auditory brainstem responses (ABR).

ABR experiments were conducted in a manner similar to that described by Song

et. al. (El-Hassar et al., 2019). Briefly, animals were anesthetized with 480 mg/kg,

i.p., chloral hydrate, and all recordings were conducted in a sound-attenuating

chamber (Industrial Acoustics Corp., Bronx, NY). A customized TDT3 system

(Tucker-Davis Technologies Inc., Alachua, FL) was used for ABR recordings.

Subdermal needle electrodes (Rochester Electro-Medical, Lutz, FL) were

positioned at the vertex (active, noninverting), the infra-auricular mastoid region

(reference, inverting), and the neck region (ground). Differentially recorded scalp

potentials were bandpass filtered between 0.05 and 3 kHz over a 15 ms epoch. A

total of 400 trials was averaged for each waveform for each stimulus condition.

Symmetrically shaped tone bursts were 3 ms long (1 ms raised cosine on/off ramps

and 1 ms plateau). All acoustic stimuli were delivered free field via a speaker (Part

FF1 2021, Tucker-Davis Technologies) positioned 10 cm from the vertex. Stimulus

levels were calibrated using a 0.5-inch condenser microphone (model 4016, ACO

Pacific, Belmont, CA) positioned at the approximate location of the animal's head

during recording sessions and are reported in decibels of sound pressure level

(SPL; referenced to 20 μPa). Stimuli of alternating polarity were delivered at a rate

of ~20/s. Tone burst responses were collected in half-octave steps ranging from

32 to 2.0 kHz. The effects of the level were determined by decreasing stimulus

intensity in 5 dB steps up to a maximum stimulus level of 90 dB SPL. The ABR

threshold was defined as the lowest intensity of sound level capable of evoking a

reproducible, visually detectable response. Latencies and amplitudes of the initial

ABR peaks were measured from sound stimuli delivered at a rate of 20/s and at a

frequency of 16 kHz, one of the most sensitive frequency ranges of hearing in

mice. Latencies and amplitudes of wave I at 90 dB SPL were also measured at all

the other frequencies. The analysis was performed in BioSig software (BioSig

Technologies, Westport, CT) on traces with discernible peaks by setting cursors

429 at the maxima and minima of the peaks. Latency was defined as the time from the

430 onset of the stimulus to the peak, while amplitude was measured by taking the

431 mean of the ΔV of the positive and negative deflections of the ABR wave.

433 **Distortion product otoacoustic emissions (DPOAE).**

434 The acoustic stimuli were produced, and responses were recorded using a

435 customized TDT3 system (Tucker-Davis Technologies Inc., Alachua, FL)

436 controlled by BioSig, digital signal processing software. DPOAE was measured

437 by inserting a probe into anesthetized mouse external ear canal to deliver sound

438 stimuli, and emissions of sound were collected by a microphone and recorded for

439 subsequent analysis. To deliver acoustic stimuli and record the responses, two

440 speakers (f_1 and f_2) (MF1 Multi-Field Magnetic Speakers, TDT) and a microphone

441 (ER-10B, Etymotic Research, Elk Grove Village, IL) were connected to a short

442 PVC tube, and tapered plastic probe that was inserted into the external auditory

443 meatus of the mice. Two simultaneous continuous pure tones (f_1 and f_2) that have

444 a frequency ratio of 1.2 (f_2/f_1) and equal sound level ($L_1 = L_2$) were delivered at

445 center frequencies of 8, 12, 16, 24 and 32 kHz with sound levels from 80 to 20 dB

446 SPL in 10 dB decrements. Stimuli duration was 83.88 ms at a repetition rate of

447 11.92 per second. The acquired DPOAE responses were averaged 100 times. The

448 DPOAE of interest was at $2f_1$-f_2, the largest and most prominent DPOAE. The

449 DPOAE threshold was defined as the lowest sound intensity capable of evoking a

450 visually detectable $2f_1$-f_2 signal above the noise floor. DPOAE amplitudes were

451 analyzed offline in BioSigRP by setting cursors at the peak of the $2f_1$-f_2 signals.

Nonlinear capacitance (NLC).

Nonlinear capacitance was conducted in a fashion much like that previously described in the Methods Mol. Biol. 2016 Song and Santos-Sacchi (Song & Santos-Sacchi, 2016). WT C57BL/6J and βV spectrin KO littermates at postnatal day 13 and at 2-month-old were compared. Apical turns of the organ of Corti were isolated from each and recorded with ionic current blocking solutions to minimize interference on measures of membrane capacitance. The extracellular solution contained (in mM): 100 NaCl, 20 tetraethylammonium (TEA)-Cl, 20 CsCl, 2 CoCl$_2$, 1 MgCl$_2$, 1 CaCl$_2$, 10 HEPES, pH 7.2. The intracellular solution contained (in mM): 140 CsCl, 2 MgCl$_2$, 10 HEPES, and 10 EGTA, pH 7.2. Pipettes had resistances of 3-5 MΩ. Gigohm seals were made, and stray capacitance was balanced out with amplifier circuitry prior to establishing whole-cell conditions. A Nikon Eclipse E600-FN microscope with 40× water immersion objective lens was used to observe cells during voltage clamp. Whole-cell voltage-clamp recordings were performed from all three rows of OHCs on whole-mount organ of Corti. OHCs were recorded at room temperature using jClamp software and an Axopatch 200B amplifier. Data were low pass filtered at 10 kHz and digitized at 100 kHz with a Digidata 1320A. In order to extract Boltzmann parameters, capacitance-voltage data were fit to the first derivative of a two-state Boltzmann function.

$$C_m = NLC + C_{sa} + C_{\text{lin}} = Q_{\max} \frac{ze}{k_B T} \frac{b}{(1+b)^2} + C_{sa} + C_{\text{lin}} \qquad \text{(eq.1)}$$

where $b = exp\left(-ze\frac{V_m - V_h}{k_B T}\right), \; C_{sa} = \frac{\Delta C_{sa}}{(1 + b^{-1})}$

Q_{max} is the maximum nonlinear charge moved, V_h is the voltage at peak

capacitance or equivalently, at half-maximum charge transfer, V_m is Rs-corrected

membrane potential, z is valence, C_{lin} is linear membrane capacitance, e is the

electron charge, kB is Boltzmann's constant, and T is the absolute temperature.

Immunocytochemistry (ICC).

For ICC, staining was performed on fixed and decalcified 3 turns of the organ of

Corti to compare the number of neuronal fibers and synapses between all 3 turns

organ of Corti in KO and WT mice. Temporal bones were fixed overnight in 4°C

and then decalcified in 120 mM ethylenediaminetetraacetic acid (EDTA) for 2 days.

Whole-mount of the organ of Corti was dissected, washed in PBS 0.05% Tween

20, 0.05% Triton-X 100 (incubation buffer) three times for 3 minutes, and incubated

with primary antibody in incubation buffer overnight at 4°C with mouse anti-CtBP-

2 antibody (1:50, BD Transduction, San Jose, CA, cat# 612044), chicken anti-

neurofilament (NF) antibody (1:500, MilliporeSigma, Billerica, MA, cat# AB5539).

After three washes in incubation buffer, the tissue was incubated with secondary

antibody in incubation buffer for 1 hour at room temperature. The secondary

antibodies used included donkey anti-rabbit Alexa Fluor 568/647 (1:200; Thermo

Fisher Scientific, cat # A-10042/A-31573), goat anti-mouse Alexa Fluor

568/647 (1:200; Thermo Fisher Scientific, cat # A-21124/A-21235) and goat anti-

chicken Alexa Fluor 488/594 (1:200; Thermo Fisher Scientific, cat # A-11039/A-

11042). After antibody treatment, cells were washed in incubation buffer and

mounted in Vectashield (Vector Laboratories, Inc., Burlingame, CA), and viewed

using a Zeiss confocal laser-scanning microscope (Oberkochen, Germany).

Nerve (NF positive) and synaptic vesicle (CtBP-2 positive) counts were performed

posthoc in Fiji (ImageJ). Briefly, z stacks of images were separated by probes,

background-subtracted (using a 50-pixel rolling ball radius), and z sections

containing the images of interest identified for further analysis (slice keeper). CtBP-

2 labeled IHCs and NF labeled afferent fibers surrounding IHCs in proximity to the

habenula and efferent fibers in the cochlear canal were counted. Z projections

(sum) of the respective sections were obtained and automated thresholds

determined (max entropy). Counts of particles (synaptic vesicles) immunopositive

for CtBP-2 larger than 0.1 microns were scored. Neuronal numbers were

established by counting the peaks in a profile plotted perpendicular to neural fibers

in the cochlea canal. For afferent nerve counts the profile was plotted at the

midpoint of nerve densities.

Statistical Analysis.

Data were analyzed using FIJI, Microsoft Excel, SigmaPlot and GraphPad Prism.

The measures of variability are depicted as error bars +/- the SEM and stated

numerically in the figure legends. Unpaired, one or two-tailed Student's t-test were

performed using both Excel and GraphPad Prism to make comparisons between

WT and Spnb5$^{(-/-)}$ data points. P-values are indicated in the figure legends to two

significant figures. The sample size (n) is stated in the figures.

COMPETING INTERESTS

526 The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

529 MCS, JPB, JSM, DN and JSS conceptualized the project. PRS and JSM generated the Spnb5 deleted mice. MCS and JPB conducted animal breeding and genotyping. MCS, JPB, SK, and DN performed gene expression and immunohistochemistry analysis. JPB and LS performed assays to assess hearing function. MCS and DN wrote the manuscript, and all authors contributed to its editing and refinement.

ACKNOWLEDGEMENTS

537 These studies were supported by grants R01 DC 008130 and R01 DC 008130 from the National Institute of Health to JSS and DN and grants P01 NS35476 (project 4) and R01 HL28560 to JSM. Other support includes the Raymond Yesner Endowment to Yale University (JSM). We also would like to thank Winston Tan, Alexei Surguchev and Lan Ji for their technical assistance.

REFERENCES

Abe, T., Kakehata, S., Kitani, R., Maruya, S., Navaratnam, D., Santos-Sacchi, J., & Shinkawa, H. (2007). Developmental expression of the outer hair cell motor prestin in the mouse. *J Membr Biol, 215*(1), 49-56. doi:10.1007/s00232-007-9004-5

Bai, J. P., Surguchev, A., Ogando, Y., Song, L., Bian, S., Santos-Sacchi, J., & Navaratnam, D. (2010). Prestin surface expression and activity are augmented by interaction with MAP1S, a microtubule-associated protein. *J Biol Chem, 285*(27), 20834-20843. doi:10.1074/jbc.M110.117853

Baker, K. E., & Parker, R. (2004). Nonsense-mediated mRNA decay: terminating erroneous gene expression. *Curr Opin Cell Biol, 16*(3), 293-299. doi:10.1016/j.ceb.2004.03.003

Benson, D. A., Cavanaugh, M., Clark, K., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., & Sayers, E. W. (2013). GenBank. *Nucleic Acids Res, 41*(Database issue), D36-42. doi:10.1093/nar/gks1195

Bramhall, N. F., McMillan, G. P., Kujawa, S. G., & Konrad-Martin, D. (2018). Use of non-invasive measures to predict cochlear synapse counts. *Hear Res, 370*, 113-119. doi:10.1016/j.heares.2018.10.006

Chomczynski, P., & Mackey, K. (1995). Short technical reports. Modification of the TRI reagent procedure for isolation of RNA from polysaccharide- and proteoglycan-rich sources. *Biotechniques, 19*(6), 942-945. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/8747660

Consortium, G. T. (2013). The Genotype-Tissue Expression (GTEx) project. *Nat Genet, 45*(6), 580-585. doi:10.1038/ng.2653

Cortese, M., Papal, S., Pisciottano, F., Elgoyhen, A. B., Hardelin, J. P., Petit, C., . . . El-Amraoui, A. (2017). Spectrin betaV adaptive mutations and changes in subcellular location correlate with emergence of hair cell electromotility in mammalians. *Proc Natl Acad Sci U S A, 114*(8), 2054-2059. doi:10.1073/pnas.1618778114

Efimova, N., Korobova, F., Stankewich, M. C., Moberly, A. H., Stolz, D. B., Wang, J., . . . Svitkina, T. (2017). betaIII Spectrin Is Necessary for Formation of the Constricted Neck of Dendritic Spines and Regulation of Synaptic

Activity in Neurons. *J Neurosci, 37*(27), 6442-6459.
doi:10.1523/JNEUROSCI.3520-16.2017

El-Hassar, L., Song, L., Tan, W. J. T., Large, C. H., Alvaro, G., Santos-Sacchi, J.,
& Kaczmarek, L. K. (2019). Modulators of Kv3 Potassium Channels
Rescue the Auditory Function of Fragile X Mice. *J Neurosci, 39*(24), 4797-
4813. doi:10.1523/JNEUROSCI.0839-18.2019

Hafidi, A., Despres, G., & Romand, R. (1990). Cochlear innervation in the
developing rat: an immunocytochemical study of neurofilament and
spectrin proteins. *J Comp Neurol, 300*(2), 153-161.
doi:10.1002/cne.903000202

Hammarlund, M., Jorgensen, E. M., & Bastiani, M. J. (2007). Axons break in
animals lacking beta-spectrin. *J Cell Biol, 176*(3), 269-275.
doi:10.1083/jcb.200611117

Hoa, M., Olszewski, R., Li, X., Taukulis, I., Gu, S., DeTorres, A., . . . Kelley, M.
W. (2020). Characterizing Adult Cochlear Supporting Cell Transcriptional
Diversity Using Single-Cell RNA-Seq: Validation in the Adult Mouse and
Translational Implications for the Adult Human Cochlea. *Front Mol
Neurosci, 13*, 13. doi:10.3389/fnmol.2020.00013

Holley, M. C., & Ashmore, J. F. (1990). Spectrin, actin and the structure of the
cortical lattice in mammalian cochlear outer hair cells. *J Cell Sci, 96 (Pt
2)*, 283-291. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/2211870

Holley, M. C., Kalinec, F., & Kachar, B. (1992). Structure of the cortical
cytoskeleton in mammalian outer hair cells. *J Cell Sci, 102 (Pt 3)*, 569-
580. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/1506434

Kujawa, S. G., & Liberman, M. C. (2009). Adding insult to injury: cochlear nerve
degeneration after "temporary" noise-induced hearing loss. *J Neurosci,
29*(45), 14077-14085. doi:10.1523/JNEUROSCI.2845-09.2009

Legendre, K., Safieddine, S., Kussel-Andermann, P., Petit, C., & El-Amraoui, A.
(2008). alphaII-betaV spectrin bridges the plasma membrane and cortical
lattice in the lateral wall of the auditory outer hair cells. *J Cell Sci, 121*(Pt
20), 3347-3356. doi:10.1242/jcs.028134

Lewis, M. A., Nolan, L. S., Cadge, B. A., Matthews, L. J., Schulte, B. A., Dubno,
J. R., . . . Dawson, S. J. (2018). Whole exome sequencing in adult-onset
hearing loss reveals a high load of predicted pathogenic variants in known
deafness-associated genes and identifies new candidate genes. *BMC*
Med Genomics, 11(1), 77. doi:10.1186/s12920-018-0395-1

Li, Y., Liu, H., Giffen, K. P., Chen, L., Beisel, K. W., & He, D. Z. Z. (2018).
Transcriptomes of cochlear inner and outer hair cells from adult mice. *Sci*
Data, 5, 180199. doi:10.1038/sdata.2018.199

Liberman, M. C., & Kujawa, S. G. (2017). Cochlear synaptopathy in acquired
sensorineural hearing loss: Manifestations and mechanisms. *Hear Res,*
349, 138-147. doi:10.1016/j.heares.2017.01.003

Lin, H. W., Furman, A. C., Kujawa, S. G., & Liberman, M. C. (2011). Primary
neural degeneration in the Guinea pig cochlea after reversible noise-
induced threshold shift. *J Assoc Res Otolaryngol, 12*(5), 605-616.
doi:10.1007/s10162-011-0277-0

Liu, Y., Qi, J., Chen, X., Tang, M., Chu, C., Zhu, W., . . . Zhong, G. (2019).
Critical role of spectrin in hearing development and deafness. *Sci Adv,*
5(4), eaav7803. doi:10.1126/sciadv.aav7803

Lorenzo, D. N., Badea, A., Zhou, R., Mohler, P. J., Zhuang, X., & Bennett, V.
(2019). betaII-spectrin promotes mouse brain connectivity through
stabilizing axonal plasma membranes and enabling axonal organelle
transport. *Proc Natl Acad Sci U S A, 116*(31), 15686-15695.
doi:10.1073/pnas.1820649116

Maison, S. F., Pyott, S. J., Meredith, A. L., & Liberman, M. C. (2013).
Olivocochlear suppression of outer hair cells in vivo: evidence for
combined action of BK and SK2 channels throughout the cochlea. *J*
Neurophysiol, 109(6), 1525-1534. doi:10.1152/jn.00924.2012

Melcher, J. R., Knudson, I. M., Fullerton, B. C., Guinan, J. J., Jr., Norris, B. E., &
Kiang, N. Y. (1996). Generators of the brainstem auditory evoked potential
in cat. I. An experimental approach to their identification. *Hear Res, 93*(1-
2), 1-27. doi:10.1016/0378-5955(95)00178-6

Papal, S., Cortese, M., Legendre, K., Sorusch, N., Dragavon, J., Sahly, I., . . . El-
Amraoui, A. (2013). The giant spectrin betaV couples the molecular

motors to phototransduction and Usher syndrome type I proteins along
their trafficking route. *Hum Mol Genet, 22*(18), 3773-3788.
doi:10.1093/hmg/ddt228

Parkinson, N. J., Olsson, C. L., Hallows, J. L., McKee-Johnson, J., Keogh, B. P.,
Noben-Trauth, K., . . . Tempel, B. L. (2001). Mutant beta-spectrin 4 causes
auditory and motor neuropathies in quivering mice. *Nat Genet, 29*(1), 61-
65. doi:10.1038/ng710

Schuknecht, H. F., & Woellner, R. C. (1953). Hearing losses following partial
section of the cochlear nerve. *Trans Am Laryngol Rhinol Otol Soc, 58*(57th
Meeting), 369-393; discussion, 393-364. Retrieved from
https://www.ncbi.nlm.nih.gov/pubmed/13146770

Sergeyenko, Y., Lall, K., Liberman, M. C., & Kujawa, S. G. (2013). Age-related
cochlear synaptopathy: an early-onset contributor to auditory functional
decline. *J Neurosci, 33*(34), 13686-13694.
doi:10.1523/JNEUROSCI.1783-13.2013

Shen, J., Scheffer, D. I., Kwan, K. Y., & Corey, D. P. (2015). SHIELD: an
integrative gene expression database for inner ear research. *Database
(Oxford), 2015*, bav071. doi:10.1093/database/bav071

Shera, C. A., & Guinan, J. J., Jr. (1999). Evoked otoacoustic emissions arise by
two fundamentally different mechanisms: a taxonomy for mammalian
OAEs. *J Acoust Soc Am, 105*(2 Pt 1), 782-798. doi:10.1121/1.426948

Song, L., McGee, J., & Walsh, E. J. (2006). Frequency- and level-dependent
changes in auditory brainstem responses (ABRS) in developing mice. *J
Acoust Soc Am, 119*(4), 2242-2257. doi:10.1121/1.2180533

Song, L., & Santos-Sacchi, J. (2016). A Walkthrough of Nonlinear Capacitance
Measurement of Outer Hair Cells. *Methods Mol Biol, 1427*, 501-512.
doi:10.1007/978-1-4939-3615-1_28

Stabach, P. R., & Morrow, J. S. (2000). Identification and characterization of beta
V spectrin, a mammalian ortholog of Drosophila beta H spectrin. *J Biol
Chem, 275*(28), 21385-21395. doi:10.1074/jbc.C000159200

Stankewich, M. C., Cianci, C. D., Stabach, P. R., Ji, L., Nath, A., & Morrow, J. S.
(2011). Cell organization, growth, and neural and cardiac development
require alphaII-spectrin. *J Cell Sci, 124*(Pt 23), 3956-3966.
doi:10.1242/jcs.080374

Stankewich, M. C., Gwynn, B., Ardito, T., Ji, L., Kim, J., Robledo, R. F., . . .
Morrow, J. S. (2010). Targeted deletion of betaIII spectrin impairs
synaptogenesis and generates ataxic and seizure phenotypes. *Proc Natl*
Acad Sci U S A, 107(13), 6022-6027. doi:10.1073/pnas.1001522107

Yue, F., Cheng, Y., Breschi, A., Vierstra, J., Wu, W., Ryba, T., . . . Mouse, E. C.
(2014). A comparative encyclopedia of DNA elements in the mouse
genome. *Nature, 515*(7527), 355-364. doi:10.1038/nature13992
681
682

684 **Table1. Beta spectrin gene specific primers.**

Primers					Amplimer (bps)
Spnb1	forward	CAAAGAGCAGGAGGTGTCAGC	reverse	TCTAGGCAGGTGCTGAAGTTC	264
	nested forward	GATGCGTGTGCCGGGCGCC	nested reverse	CTGGTATTGATCTCTGCCT	197
Spnb2	forward	CCGGTGGGAGTGGTTAAGAC	reverse	CATCCCAGGTCGCTGCAGAC	188
	nested forward	GATTTTGGAGGTCCATCAG	nested reverse	TCAAACGCCTCATGGCGCT	144
Spnb5	forward	CTCCGGAGCTTTGAGCAGGCTGC	reverse	CCTGAGCCCACGCTAGCAATTCC	439
	nested forward	GATACAGGAGAAGACCACC	nested reverse	TTCCTGGACTTTGGCCTCC	319
Spnb5	forward	TGGCCTTCCTCAGGGCCAAGGT	reverse	GCACTCATCTGGATGCAGGGCTG	440
	nested forward	CTTGGACAGGGAGGAGTTTGG	nested reverse	GCACTCATCTGGATGCAGGGCTG	313

685
686 Primer sets were used to detect for mRNA of Spnb1, Spnb2, and Spnb5 in the
687 brain, organ of Corti, and isolated OHCs. The PCR amplimer sizes from mRNA
688 are presented in the final column.
689

690 **Table 2. Primers for the Spnb5 gene used to prepare the three DNA fragments**
691 **that were subcloned into pEasy-Flox.**

Fragment 1	forward	CAGCGGCCGCAGGGGCTGCCTCCTACTGTACC	reverse	GAGCGGCCGCACTTGAATGACCCAGAGCCACCCCT
Fragment 2	forward	GGTCTAGACCTCTTACTCGCTTAGGTTCCAATC	reverse	GCTCTAGATGACTCTACAGCAGACCCCCAGGTC
Fragment 3	forward	GACTCGAGCCTGTAGCTATTTCTGGCCTACCCCTCC	reverse	GACTCGAGGGAGGGAGGGTTACTGTAGTAGAAGGGC

692
The targeting vector for homologous recombination was constructed by cloning
three PCR generated fragments into the pEASY-Flox vector. Fragment 1 is a 4115
bp product, fragment 2 is a 469 bp product that contains Spnb5 exon 2, and
fragment 3 is 4371 bp of sequence that spans up to exon 5. The three fragments
were sequence verified. The oligos contain common restriction sites at their 5' end
for easy cloning into the targeting vector.
699

700 **Table 3. Primers for rt-PCR in WT and Spnb5[(-/-)] tissue.**

Spnb5	exon 58/60	forward	CTCCGGAGCTTTGAGCAGGCTGC	reverse	CCTGAGCCCACGCTAGCAATTCC	439
Spnb5	exon 53/55	forward	AGGAGAAACACCAGAATTTGGAGAATG	reverse	TGTCCACTCTCCAGAAGGGCAACTG	412
Spnb2	exon 28/29	forward	GATACAGGAGAAGACCACC	reverse	TTCCTGGACTTTGGCCTCC	144
Spna2	exon 54/55	forward	AACAGGAGTGACTGAGGAGG	reverse	CCTTCCTCAACCATTGGCAGA	144

701
702 These primer sets were tested on cDNA from testis isolated from WT and Spnb5[(-
703 /-)] 3-month-old mice. Because the C-terminus of these large genes tends to be
704 more abundant than N-terminus exons, we designed primer sets that will amplify
705 exons near the poly-A tail.
706
707