Drosophila Evolution over Space and Time (DEST) - A New Population Genomics Resource

Martin Kapun[1,2,*,☮,§], Joaquin C. B. Nunez[3,*], María Bogaerts-Márquez[4,*,§], Jesús Murga-Moreno[5,6,*,§], Margot Paris[7,*,§], Joseph Outten[3], Marta Coronado-Zamora[4,§], Courtney Tern[3], Omar Rota-Stabelli[8,§], Maria P. García Guerreiro[5,§], Sònia Casillas[5,6,§], Dorcas J. Orengo[9,10,§], Eva Puerma[9,10,§], Maaria Kankare[11,§], Lino Ometto[12,§], Volker Loeschcke[13,§], Banu S. Onder[14,§], Jessica K. Abbott[15,§], Stephen W. Schaeffer[16,#], Subhash Rajpurohit[17,18,#], Emily L Behrman[17,19,#], Mads F. Schou[13,15,§], Thomas J.S. Merritt[20,#], Brian P Lazzaro[21,#], Amanda Glaser-Schmitt[22,§], Eliza Argyridou[22,§], Fabian Staubach[23,§], Yun Wang[23,§], Eran Tauber[24,§], Svitlana V. Serga[25,26,§], Daniel K. Fabian[27,#], Kelly A. Dyer[28,#], Christopher W. Wheat[29,§], John Parsch[22,§], Sonja Grath[22,§], Marija Savic Veselinovic[30,§], Marina Stamenkovic-Radak[30,§], Mihailo Jelic[30,§], Antonio J. Buendía-Ruíz[31,§], M. Josefa Gómez-Julián[31,§], M. Luisa Espinosa-Jimenez[31,§], Francisco D. Gallardo-Jiménez[32,§], Aleksandra Patenkovic[33,§], Katarina Eric[33,§], Marija Tanaskovic[33,§], Anna Ullastres[4,§], Lain Guio[4,§], Miriam Merenciano[4,§], Sara Guirao-Rico[4,§], Vivien Horváth[4,§], Darren J. Obbard[34,§], Elena Pasyukova[35,§], Vladimir E. Alatortsev[35,§], Cristina P. Vieira[36,37,§], Jorge Vieira[36,37,§], J. Roberto Torres[38,§], Iryna Kozeretska[25,26,§], Oleksandr M. Maistrenko[26,39,§], Catherine Montchamp-Moreau[40,§], Dmitry V. Mukha[41,§], Heather E. Machado[42,43,#], Tânia F. Paulo[44,§], Antonio Barbadilla[5,6,§], Dmitri Petrov[42,☮,#], Paul Schmidt[16,☮,#,§], Josefa Gonzalez[4,☮,#,§], Thomas Flatt[7,☮,#,§], Alan O. Bergland[3,☮,#,§]

* equal contribution

☮ To whom correspondence should be addressed

§ The European *Drosophila* Population Genomics Consortium (DrosEU)

The *Drosophila* Real-Time Evolution Consortium (DrosRTEC)

1   Department of Evolutionary Biology and Environmental Studies, University of Zürich, Switzerland

2   Department of Cell & Developmental Biology, Center of Anatomy and Cell Biology, Medical University of Vienna, Vienna, Austria

3   Department of Biology, University of Virginia, Charlottesville, USA

4   Institute of Evolutionary Biology, CSIC-Universitat Pompeu Fabra, Barcelona,

Spain

Department of Genetics and Microbiology, Universitat Autònoma de Barcelona, Barcelona, Spain

Institute of Biotechnology and Biomedicine, Universitat Autònoma de Barcelona, Barcelona, Spain

Department of Biology, University of Fribourg, Fribourg, Switzerland

Center Agriculture Food Environment, University of Trento, San Michele all' Adige, Italy

Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de Barcelona, Barcelona, Spain

Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona, Barcelona, Spain

Department of Biological and Environmental Science, University of Jyväskylä, Jyväskylä, Finland

Department of Biology and Biotechnology, University of Pavia, Pavia, Italy

Department of Biology, Aarhus University, Aarhus, Denmark

Department of Biology, Hacettepe University, Ankara, Turkey

Department of Biology, Lund University, Lund, Sweden

Department of Biology, The Pennsylvania State University, University Park, USA

Department of Biology, University of Pennsylvania, Philadelphia, USA

Division of Biological and Life Sciences, School of Arts and Sciences, Ahmedabad University, Ahmedabad, India

Janelia Research Campus, Ashburn, USA

Department of Chemistry & Biochemistry, Laurentian University, Sudbury, Canada

Department of Entomology, Cornell University, Ithaca, USA

Division of Evolutionary Biology, Faculty of Biology, Ludwig-Maximilians-Universität, Munich, Germany

Department of Evolution and Ecology, University of Freiburg, Freiburg, Germany

Department of Evolutionary and Environmental Biology, Institute of Evolution, University of Haifa, Haifa, Israel

Department of General and Medical Genetics, Taras Shevchenko National University of Kyiv, Kyiv, Ukraine

State Institution National Antarctic Scientific Center, Ministry of Education and Science of Ukraine, Kyiv, Ukraine

Department of Genetics, University of Cambridge, Cambridge, UK

Department of Genetics, University of Georgia, Athens GA, USA

Department of Zoology, Stockholm University, Stockholm, Sweden

Faculty of Biology, University of Belgrade, Belgrade, Serbia

IES Eladio Cabañero, Tomelloso, Spain

IES Jose de Mora, Baza, Spain

Institute for Biological Research "Siniša Stanković", National Institute of Republic of
        Serbia, University of Belgrade, Belgrade, Serbia

34   Institute of Evolutionary Biology, University of Edinburgh, UK

35   Institute of Molecular Genetics of the National Research Centre "Kurchatov
        Institute", Moscow, Russia

36   Instituto de Biologia Molecular e Celular (IBMC), Porto, Portugal

37   Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Porto,
        Portugal

38   La ciència al teu mòn, Barcelona, Spain

39   Structural and Computational Biology Unit, European Molecular Biology Laboratory,
        Heidelberg, Germany

40   UMR Évolution, Génomes, Comportement et Écologie, Université Paris-Saclay,
        CNRS, Gif-sur-Yvette, France

41   Vavilov Institute of General Genetics, Russian Academy of Sciences, Moscow,
        Russia

42   Department of Biology, Stanford University, Stanford, USA

43   Wellcome Trust Sanger Institute, Hinxton, UK

44   Instituto Gulbenkian de Ciência, Oeiras, Portugal

ABSTRACT

Drosophila melanogaster is a premier model in population genetics and genomics, and a growing number of whole-genome datasets from natural populations of this species have been published over the last 20 years. A major challenge is the integration of these disparate datasets, often generated using different sequencing technologies and bioinformatic pipelines, which hampers our ability to address questions about the evolution and population structure of this species. Here we address these issues by developing a bioinformatics pipeline that maps pooled sequencing (Pool-Seq) reads from *D. melanogaster* to a hologenome consisting of fly and symbiont genomes and estimates allele frequencies using either a heuristic (PoolSNP) or a probabilistic variant caller (SNAPE-pooled). We use this pipeline to generate the largest data repository of genomic data available for *D. melanogaster* to date, encompassing 271 population samples from over 100 locations in >20 countries on four continents. Several of these locations are sampled at different seasons across multiple years. This dataset, which we call *Drosophila Evolution over Space and Time* (DEST), is coupled with sampling and environmental meta-data. A web-based genome browser and web portal provide easy access to the SNP dataset. Our aim is to provide this scalable platform as a community resource which can be easily extended via future efforts for an even more extensive cosmopolitan dataset. Our resource will enable population geneticists to analyze spatio-temporal genetic patterns and evolutionary dynamics of *D. melanogaster* populations in unprecedented detail.

Keywords: *Drosophila melanogaster*, population genomics, SNPs, evolution, adaptation, demography

Introduction

The vinegar fly *Drosophila melanogaster* is one of the oldest and most important genetic model systems and has played a key role in the development of theoretical and empirical

population genetics (Schneider 2000; Hales *et al.* 2015; Haudry *et al.* 2020). Through decades of work, we now have a basic picture of the evolutionary origin (David and Capy

1988; Lachaise *et al.* 1988; Keller 2007; Sprengelmeyer *et al.* 2020), colonization history and demography (Caracristi and Schlötterer 2003; Li and Stephan 2006; Duchen *et al.* 2013;

Grenier *et al.* 2015; Arguello *et al.* 2019; Kapopoulou *et al.* 2020), and spatio-temporal diversification patterns of this species and its close relatives (Kolaczkowski *et al.* 2011;

Fabian *et al.* 2012; Bergland *et al.* 2014; Lack *et al.* 2016; Machado *et al.* 2016; Kapun *et al.* 2016, 2020). The availability of high-quality reference genomes (Adams 2000; Celniker and

Rubin 2003; dos Santos *et al.* 2015) and genetic tools (Schneider 2000; Duffy 2002; Jennings 2011; Hales *et al.* 2015; Haudry *et al.* 2020) efficiently facilitates placing

evolutionary studies of flies in a mechanistic context, allowing for the functional characterization of ecologically relevant polymorphism (e.g., de Jong and Bochdanovits

2003; Paaby *et al.* 2010, 2014; Mateo *et al.* 2014; Kapun *et al.* 2016; Durmaz *et al.* 2018, 2019; Ramaekers *et al.* 2019).

Recently, work on the evolutionary biology of *Drosophila* has been fueled by the growing number of population genomic datasets from field collections across a large portion

of *D. melanogaster*'s range (Grenier *et al.* 2015; Machado *et al.* 2019; Guirao-Rico and González 2019; Arguello *et al.* 2019). These genomic data consist either of re-sequenced

inbred (or haploid) individuals (e.g., Mackay *et al.* 2012; Langley *et al.* 2012; Grenier *et al.* 2015; Lack *et al.* 2015, 2016; Mateo *et al.* 2018; Kapopoulou *et al.* 2020) or pooled

sequencing (Pool-Seq; e.g., Kolaczkowski *et al.* 2011; Fabian *et al.* 2012; Bastide *et al.* 2013; Campo *et al.* 2013; Bergland *et al.* 2014; Machado *et al.* 2016, 2019; Kapun *et al.*

2016, 2020) of outbred population samples. Pooled re-sequencing provides accurate and precise estimates of allele frequencies across most of the allele frequency spectrum (Zhu *et*

al. 2012; Lynch *et al.* 2014; Schlötterer *et al.* 2014) at a fraction of the cost of individual-based sequencing. Although Pool-Seq retains limited information about linkage

disequilibrium (LD) relative to individual sequencing (Feder *et al.* 2012), Pool-Seq data can be used to infer complex demographic histories (e.g., Cheng *et al.* 2012; Bergland *et al.*

2016; Deitz *et al.* 2016; Gould *et al.* 2017; Corbett-Detig and Nielsen 2017; Giesen *et al.* 2020), characterize levels of diversity (Kofler *et al.* 2011a, 2011b; Ferretti *et al.* 2013; Kapun

et al. 2020), and infer genomic loci involved in recent adaptation in nature (Flatt 2016; Kapun *et al.* 2016, 2020; Gould *et al.* 2017; Machado *et al.* 2019; Bogaerts-Márquez *et al.* 2020)

and during experimental evolution (e.g. Turner *et al.* 2011; Orozco-terWengel *et al.* 2012;

Burke 2012; Kofler and Schlötterer 2014). However, the rapidly increasing number of genomic datasets processed with different bioinformatic pipelines makes it difficult to compare results across studies and to jointly analyze multiple datasets. Differences among bioinformatic pipelines include filtering methods for the raw reads, mapping algorithms, the choice of the reference genome or SNP calling approaches, potentially generating biases when combining processed datasets from different sources for joint analyses (e.g., Gautier *et al.* 2013; Hoban *et al.* 2016).

To address these issues, we have developed a modular bioinformatics pipeline to map Pool-Seq reads to a hologenome consisting of fly and microbial genomes, to remove reads from potential *D. simulans* contaminants, and to estimate allele frequencies using two complementary SNP callers. Our pipeline is available as a Docker image (available from https://dest.bio) to standardize versions of software used for filtering and mapping, to make the pipeline available independently of the operating system used and to facilitate future updates and modification of the pipeline. In addition, our pipeline allows using either heuristic or probabilistic methods for SNP calling, based on PoolSNP (Kapun *et al.* 2020) and SNAPE-pooled (Raineri *et al.* 2012). We also provide tools for performing *in-silico* pooling of existing inbred (haploid) lines that exist as part of other *Drosophila* population genomic resources (Pool *et al.* 2012; Langley *et al.* 2012; Grenier *et al.* 2015; Kao *et al.* 2015; Lack *et al.* 2015, 2016). This pipeline is also designed to be flexible, facilitating the streamlined addition of new population samples as they arise.

Using this pipeline, we generated a unified dataset of pooled allele frequency estimates of *D. melanogaster* sampled across large portions of Europe and North America. This dataset is the result of the collaborative efforts of the European DrosEU (Kapun *et al.* 2020) and DrosRTEC (Machado *et al.* 2019) consortia and combines both novel and previously published population genomic data. Our dataset combines samples from 100 localities, 55 of which were sampled at two or more time points across the reproductive season (~10-15 generations/year) for one or more years. Collectively, these samples represent >13,000 individuals, cumulatively sequenced to >16,000x coverage. The cost-effectiveness of Pool-Seq has enabled us to estimate genome-wide allele frequencies over geographic space (continental and sub-continental) and time (seasonal, annual and decadal) scales, thus making our data a unique resource for advancing our understanding of fundamental adaptive and neutral evolutionary processes. We provide data in two file formats (VCF and GDS: (Danecek *et al.* 2011; Zheng *et al.* 2017), thus allowing researchers to utilize a variety of tools for computational analyses. Our dataset also contains sampling and environmental meta-data to enable various downstream analyses of biological interest.

102   **Materials and Methods**

Data sources. The genomic dataset presented here has been assembled from a
104   combination of Pool-Seq libraries and *in-silico* pooled haplotypes. We combined 246 Pool-
Seq libraries of population samples from Europe, North America and the Caribbean that
106   were sampled through space and time by two collaborating consortia in North America
(DrosRTEC: https://web.sas.upenn.edu/paul-schmidt-lab/dros-rtec/) and Europe (DrosEU:
108   http://droseu.net) between 2003 and 2016. In addition, we integrated genomic data from
>900 inbred or haploid genomes from 25 populations in Africa, Europe, Australia, and North
110   America available from the *Drosophila* Genome Nexus dataset (DGN; Lack *et al.* 2015,
2016). We further included the *D. simulans* haplotype, built as part of the DGN dataset, as
112   an outgroup, making this repository of 272 (246 pool-seq + 25 DGN + 1 *D. simulans*) whole-
genome sequenced samples the largest dataset of genome-wide SNPs available for *D.
114   melanogaster* to date.

116   **Metadata.** We assembled uniform meta-data for all samples (Supplemental Material, Table
S1). This information includes collection coordinates, collection date, and the number of flies
118   per sample. Samples are also linked to bioclimatic variables from the nearest WorldClim
(Hijmans *et al.* 2005) raster cell at a resolution of 2.5° and to weather stations from the
120   Global Historical Climatology Network (GHCND; ftp://ftp.ncdc.noaa.gov/pub/data/ghcn/daily/)
for future analysis of the environmental drivers that might underlie genetic change. We also
122   provide summaries of basic attributes of each sample derived from the sequencing data
including average read depth, PCR-duplicate rate, *D. simulans* contamination rate, relative
124   abundances of non-synonymous versus synonymous polymorphisms (p_N/p_S), the number of
private polymorphisms, and diversity statistics (Watterson's θ, π and Tajima's D).

Sample collection. The majority of population samples contributed by the DrosEU and the
128   DrosRTEC consortia was collected in a coordinated fashion to generate a consistent dataset
with minimized sampling bias. In brief, fly collections were performed exclusively in natural
130   or semi-natural habitats, such as orchards, vineyards and compost piles. For most European
collections, flies were collected using mashed banana, or apples with live yeast as bait in
132   traps placed at sampling sites for multiple days to attract flies or by sweep netting (see
Kapun *et al.* 2020 for more details). For North American collections, flies were collected by
134   sweep-net, aspiration, or baiting over natural substrate or using baited traps (see Behrman
et al. 2018; Machado *et al.* 2019 for details). Samples were either field caught flies (n-227),

136   from F1 offspring of wild caught females (n=7), from a mixture of F1 and wild caught flies
       (n=7), or from flies kept as isofemale lines in the lab for 5 generations or less (n=4); see
138   Supplemental Table 1 for more information. To minimize cross-contamination with the
       closely related sympatric sister species *D. simulans*, we only sequenced male *D.*
140   *melanogaster* specimens, allowing for higher confidence discrimination between the two
       species based on the morphology of male genitalia (Capy and Gibert 2004; Markow and
142   O'Grady 2005). Samples were stored in 95% ethanol at -20°C before DNA extraction.

144   **DNA extraction and sequencing.** The DrosEU and DrosRTEC consortia centralized
       extractions from pools of flies. DNA was extracted either using chloroform/phenol-based
146   (DrosEU: Kapun *et al.* 2020) or lithium chloride/potassium acetate extraction protocols
       (DrosRTEC: Bergland *et al.* 2014; Machado *et al.* 2019) after homogenization with bead
148   beating or a motorized pestle. DrosEU samples from the 2014 collection were sequenced on
       an Illumina NextSeq 500 sequencer at the Genomics Core Facility of Pompeu Fabra
150   University in Barcelona, Spain. Libraries of the previously unpublished DrosEU samples
       from 2015 and 2016 were constructed using the Illumina TruSeq PCR Free library
152   preparation kit following the manufacturer's instructions and sequenced on the Illumina
       HiSeq X platform as paired-end fragments with 2 x 150 bp length at NGX Bio (San
154   Francisco, California, USA). The previously published samples of the DrosRTEC consortium
       were prepared and sequenced on GAIIX, HiSeq2000 or HiSeq3000 platforms, as described
156   in Bergland *et al.* (2014) and Machado *et al.* (2019). For information on DNA extraction and
       sequencing methods of the various DGN samples see Lack *et al.* (2016).

       Mapping pipeline. The joint analysis of genomic data from different sources requires the
160   application of uniform quality criteria and a common bioinformatics pipeline. To accomplish
       this, we developed a standardized pipeline that performs filtering, quality control and
162   mapping of any given Pool-Seq sample (see Supplemental Information Figure S1). This
       pipeline performs quality filtering of raw reads, maps reads to a hologenome (see below),
164   performs realignment and filtering around indels, and filters for mapping quality. The output
       of this pipeline includes quality control metrics, bam files, pileup files, and allele frequency
166   estimates for every site in the genome (gSYNC, see below). Our pipeline is provided as a
       Docker image, which automatically installs external software and executes the pipeline
168   across various operating systems. Our pipeline will facilitate the integration of future samples
       to extend the worldwide *D. melanogaster* SNP dataset presented here.
170   The mapping pipeline includes the following major steps. Prior to mapping, we removed
       sequencing adapters and trimmed the 3' ends of all reads using cutadapt (Martin 2011). We
172   enforced a minimum base quality score ≥ 18 (-q flag in cutadapt) and assessed the quality of

raw and trimmed reads with FASTQC (Andrews 2010). Trimmed reads with minimum length
174 < 75 bp were discarded and only intact read pairs were considered for further analyses.
Overlapping paired-end reads were merged using *bbmerge* (v. 35.50; (Bushnell *et al.* 2017).
176 Trimmed reads were mapped against a compound reference genome ("hologenome")
consisting of the genomes of *D. melanogaster* (v.6.12) and *D. simulans* (Hu *et al.* 2013) as
178 well as genomes of common commensals and pathogens, including *Saccharomyces
cerevisiae* (GCF_000146045.2), *Wolbachia pipientis* (NC_002978.6), *Pseudomonas
180 entomophila* (NC_008027.1), *Commensalibacter intestine* (NZ_AGFR00000000.1),
Acetobacter pomorum (NZ_AEUP00000000.1), *Gluconobacter morbifer*
182 (NZ_AGQV00000000.1), *Providencia burhodogranariea* (NZ_AKKL00000000.1),
Providencia alcalifaciens (NZ_AKKM01000049.1), *Providencia rettgeri*
184 (NZ_AJSB00000000.1), *Enterococcus faecalis* (NC_004668.1), *Lactobacillus brevis*
(NC_008497.1), and *Lactobacillus plantarum* (NC_004567.2), using *bwa mem* (v. 0.7.15; Li
186 2013) with default parameters. We retained reads with mapping quality greater than 20 and
reads with no secondary alignment using *samtools* (Li *et al.* 2009). PCR duplicate reads
188 were removed using *Picard MarkDuplicates* (v.1.109; http://picard.sourceforge.net).
Sequences were re-aligned in the proximity of insertions-deletions (indels) with GATK (v3.4-
190 46; McKenna *et al.* 2010). We identified and removed any reads that mapped to the *D.
simulans* genome using a custom python script, following methods outlined previously
192 (Machado *et al.* 2019; Kapun *et al.* 2020; for a more in-depth analysis of D. simulans
contamination see Wallace *et al.* 2020).

Incorporation of the DGN dataset. We incorporated population allele frequency estimates
196 derived from inbred-line and haploid embryo sequencing data from populations sampled
throughout the world. These samples have been previously collected and sequenced by
198 several groups (Pool *et al.* 2012; Langley *et al.* 2012; Grenier *et al.* 2015; Kao *et al.* 2015;
Lack *et al.* 2015, 2016) and form the *Drosophila* Genome Nexus dataset (DGN; Lack *et al.*
200 2015, 2016). We included 25 DGN populations with ≥ 5 individuals per population, plus the
D. simulans haplotype built as part of the DGN dataset. The DGN populations that we used
202 are primarily from Africa (n=18) but also include populations from Europe (n=2), North
America (n=3), Australia (n=1), and Asia (n=1).
204   To incorporate the DGN populations into the DrosEU and DrosRTEC Pool-Seq
datasets, we used the pre-computed FASTA files ("Consensus Sequence Files" from
206 https://www.johnpool.net/genomes.html) and calculated allele frequencies at every site, for
each population, using custom *bash* scripts. We calculated allele frequencies per population
208 by summing reference and alternative allele counts across all individuals. Since estimates of
allele frequencies and total allele counts for the DGN samples only consider unambiguous

210   IUPAC codes, heterozygous sites or sites masked as N's in the original FASTA files were converted to missing data. We used *liftover* (Kuhn *et al.* 2013) to translate genome

212   coordinates to Drosophila reference genome release 6 (dos Santos *et al.* 2015) and formatted them to match the gSYNC format (described below).

SNP calling strategies. We used two complementary approaches to perform SNP calling.

The first was PoolSNP (Kapun *et al.* 2020), a heuristic tool which identifies polymorphisms based on the combined evidence from multiple samples. This approach is similar to other

common Pool-Seq variant calling tools (Koboldt *et al.* 2009, 2012; Kofler *et al.* 2011a, 2011b). PoolSNP integrates allele counts across multiple independent samples and applies

stringent minor allele count and minor allele frequency thresholds for variant detection. PoolSNP is expected to be good at detecting variants present in multiple populations, but is

not very sensitive to rare private alleles. The second approach was SNAPE-pooled (Raineri *et al.* 2012), a Bayesian tool which identifies polymorphic sites for each population

independently using pairwise nucleotide diversity estimates as a prior. SNAPE-pooled is expected to be more sensitive to rare private polymorphisms, but also might have a higher

false positive rate for variant detection.

gSYNC generation and filtering. Our pipeline utilizes a common data-format (SYNC; Kofler *et al.* 2011b) to encode allele counts for each population sample. A "genome-wide SYNC"

(gSYNC) file records the number of A,T,C, and G for every site of the reference genome. Because gSYNC files for all populations have the same dimension, they can be quickly

combined and passed to a SNP calling tool. They can be filtered and are also relatively small for a given sample (~500Mb), enabling efficient data sharing and access. The gSYNC

file is analogous to the gVCF file format as part of the GATK HaplotypeCaller approach (McKenna *et al.* 2010) but is specifically tailored to Pool-Seq samples.

To generate a Pool-SNP gSYNC file, we first converted BAM files to the MPILEUP format with *samtools mpileup* using the -B parameter to suppress recalculations of per-base

alignment qualities and filtered for a minimum mapping quality with the parameter -q 25. Next, we converted the MPILEUP file containing mapped and filtered reads to the gSYNC

format using custom python scripts, which are available at https://dest.bio. To generate a SNAPE-pooled gSYNC file, we ran the SNAPE-pooled version specific to Pool-Seq data for

each sample with the following parameters: θ=0.005, D=0.01, prior='informative', fold='unfolded' and nchr=number of flies (x2 for autosomes and x1 for the X chromosome)

following Guirao-Rico and Gonzalez (2021). We converted the SNAPE-pooled output file to a gSYNC file containing the counts of each allele per position and the posterior probability of

polymorphism as defined by SNAPE-pooled using custom python scripts. We only

considered positions with a posterior probability ≥ 0.9 as being polymorphic and with a posterior probability ≤ 0.1 as being monomorphic. In all other cases, positions were marked as missing data.

We masked gSYNC files for Pool-SNP and SNAPE-pooled using a common set of filters. Sites were filtered from gSYNC files if they had: (1) minimum read depth < 10; (2)

maximum read depth > the 95% coverage percentile of a given chromosomal arm and sample; (3) located within repetitive elements as defined by RepeatMasker; (4) within 5 bp

distance up- and downstream of indel polymorphisms identified by the GATK IndelRealigner. Filtered sites were converted to missing data in the gSYNC file. The location of masked

positions for every sample was recorded as a BED file.

VCF generation. We combined the masked PoolSNP-gSYNC files into a two-dimensional matrix, where rows correspond to each position in the reference genome and columns

describe chromosome, position and reference allele, followed by allele counts in SYNC format for every sample in the dataset. This combined matrix was then subjected to variant

calling using PoolSNP, resulting in a VCF formatted file. We performed SNP calling only for the major chromosomal arms (X, 2L, 2R, 3L, 3R) and the 4th (dot) chromosome.

We first evaluated the choice of two heuristic parameters applied to PoolSNP: global minor allele count (MAC) and global minor allele frequency (MAF). Using all 272 samples,

we varied MAF (0.001, 0.01, 0.05) and MAC (5-100) and called SNPs at a randomly selected 10% subset of the genome. We calculated p_N/p_S and used this value to tune our

choice of MAF and MAC. We found that a global MAF=0.001 and a global MAC=50 provided reasonable estimates of p_N/p_S for all populations. We therefore used these parameters for

genome-wide variant calling (see *Results*: Identification and quality control of SNPs). We kept a third heuristic parameter, the missing data rate, constant at a minimum of 50%.

We generated three versions of the variant files, which differ in their inclusion of the DGN samples and the SNP calling strategy. For PoolSNP variant calling, we generated two

variant tables: the first version incorporates all 272 samples of the Pool-Seq (DrosRTEC, DrosEU) and *in-silico* Pool-Seq populations (DGN). The second version only considers the

246 Pool-Seq samples. We combined masked SNAPE-pooled gSYNC files into a two-dimensional matrix, as described above, and generated a VCF formatted output based on

allele counts for any site found to be polymorphic in one or more populations. Based on this dataset we then generated a SNAPE-pooled VCF file, which included the 246 Pool-Seq

samples. Final VCF files were annotated with SNPeff (version 4.3; Cingolani *et al.* 2012) and stored in VCF and BCF (Danecek *et al.* 2011) file formats alongside an index file in TABIX

format (Li 2011). Besides VCF files, we also stored SNP data in the GDS file format using the *R* package SeqArray (Zheng *et al.* 2017).

Population genetic analyses. We estimated allele frequencies for each site across populations as the ratio of the alternate allele count to the total site coverage. We also calculated per-site averages for nucleotide diversity (π, Nei 1987), Watterson's θ (Watterson 1975) and Tajima's D (Tajima 1989) across all sites or in non-overlapping windows of 100 kb, 50 kb and 10 kb length. To estimate these summary statistics, we converted masked gSYNC files (with positions filtered for repetitive elements, low and high read depth, and proximity to indels; see *gSYNC generation and filtering*) back to the mpileup format using custom-made scripts. mpileup files were processed using npstat v.1 (Ferretti *et al.* 2013) with parameters -maxcov 10000 and -nolowfreq m=0 in order to include all filtered positions for analysis. We only considered sites identified as being polymorphic by PoolSNP or SNAPE-pooled for analysis, using the -snpfile option of npstat. For the DGN populations, chromosomes-wide summary statistics were estimated only for samples with less than 50% missing data per chromosome. Due to small sample sizes, Tajima's D was not estimated for 7 African DGN populations that consisted of only 5 haploid embryos. In addition, we calculated p_N/p_S ratios based on SNP annotations with SNPeff (Cingolani *et al.* 2012) using a custom-made python script. To compare population genetic estimates between the PoolSNP versus SNAPE-pooled datasets, we performed Pearson's correlations on the 210 populations present in both datasets (see Identification and quality control of SNPs) using the stats package of *R* v. 3.6.3. The effects of pool size (number of individuals sampled per population) on genome-wide estimates of π, Watterson's θ, Tajima's D and p_N/p_S estimates were examined for European and North American populations using the PoolSNP dataset and a generalized linear model (GLM) in *R* v3.6.3. Finally, for 48 European populations we estimated Pearson's correlations between π, Watterson's θ and Tajima's D as estimated from the PoolSNP dataset versus previous estimates by Kapun *et al.* (2020) using the stats package of *R* v3.6.3.

Next, we examined patterns of between-population differentiation by calculating window-wise estimates of pairwise F_{ST}, based on the method from Hivert *et al.* (2018) implemented in the computePairwiseFSTmatrix() function of the *R* package poolfstat (v1.1.1). This analysis was performed for the dataset composed of 271 samples processed with PoolSNP, focusing on SNPs shared across the whole dataset. Finally, we averaged pairwise F_{ST} within and among phylogenetic clusters (Africa [17 samples], North America [76 samples], Eastern Europe [83 samples] and Western Europe [93 samples]; not included: China and Australia). These F_{ST} tracks at windows sizes of 100kb, 50kb and 10kb are available at https://dest.bio (Supplemental Figures S2, S3).

To assess population structure in the worldwide dataset, we applied PCA, population clustering, and population assignment based on a discriminant analysis of principal

components (DAPC; Jombart *et al.* 2010) to all 271 PoolSNP-processed samples. For these

322   analyses, we subsampled a set of 100,000 SNPs spaced apart from each other by at least

500 bp. We optimized our models using cross-validation by iteratively dividing the data as

324   90% for training and 10% for learning. We extracted the first 40 PCs from the PCA and ran

Pearson's correlations between each PC and all loci. We subsequently extracted the top

326   33,000 SNPs with large and significant correlations to PCs 1-40. We chose the 33,000

number as a compromise between panel size and differentiation power. For example,

328   depending on the number of individuals surveyed, these 33,000 DIMs can discern

divergence (τ) between two populations with parametric F_{ST} of 0.001- 0.0001 for sample

330   sizes (n) of 10-1000. These estimates come from the phase change formula: $\tau \approx F_{ST} =$

$1/(nm)^{1/2}$ (Patterson *et al.* 2006). Here, the two populations were sampled for n/2 individuals

332   and genotyped at m=33,000 markers. Furthermore, we included SNPs as a function of the

%VE of each PC. PCAs, clustering, and assignment-based DAPC analyses were carried out

334   using the *R* packages FactoMiner (v. 2.3), factoextra (v. 1.0.7) and adegenet (v. 2.1.3),

respectively.

Web-based genome browser. Our HTML-based DEST browser (Supplemental Information

338   Figure S2) is built on a JBrowse Docker container (Buels *et al.* 2016), which runs under

Apache on a CentOS 7.2 Linux x64 server with 16 Intel Xeon 2.4 GHz processors and 32

340   GB RAM. It implements a hierarchical data selector that facilitates the visualization and

selection of multiple population genetic metrics or statistics for the 272 samples based on

342   the PoolSNP-processed dataset, taking into account sampling location and date.

Importantly, our genome browser provides a portal for downloading allelic information and

344   pre-computed population genetics statistics in multiple formats (Supplemental Information

Figures 2A+C, S3), a usage tutorial (Supplemental Information Figure S2B) and versatile

346   track information (Supplemental Information Figure S2D). Bulk downloads of full variation

tracks are available in BigWig format (Kent *et al.* 2010) and Pool-Seq files (in VCF format)

348   are downloadable by population and/or sampling date using custom options from the Tools

menu (Supplemental Information Figure S2C). All data, tools, and supporting resources for

350   the DEST dataset, as well as reference tracks downloaded from FlyBase (v.6.12) (dos

Santos *et al.* 2015), are freely available at https://dest.bio.

354   **Results and Discussion**

Integrating a worldwide collection of *D. melanogaster* population genomics

356   **resources.** We developed a modular and standardized pipeline for generating allele

frequency estimates from pooled resequencing of *D. melanogaster* genomes (Supplemental
358   Figure 1). Using this pipeline, we assembled a dataset of allele frequencies from 271 *D.
melanogaster* populations sampled around the world (Figure 1A, Supplemental Material,
360   Table S1). Many of these samples were collected at the same location, at different seasons
and over multiple years (Figure 1B). The nature of the genomic data for each population
362   varies as a consequence of biological origin (e.g., inbred lines or Pool-Seq), library
preparation method, and sequencing platform.

Figure 1. Sampling location, dates, and quality metrics. (A) Map showing the 271 sampling localities
366   forming the DEST dataset. Colors denote the datasets that were combined together. (B) Collection
dates for localities sampled more than once. (C) General sample features of the DEST dataset. The
368   x-axis represents the population sample, ordered by the average read depth.

370   To assess whether these features affect basic attributes of the dataset, we calculated six
basic quality metrics (Figure 1C, Supplemental Material, Table S2). On average, median
372   read depth across samples is 62X (DGN samples range: 1-190X; Pool-Seq samples range:
10-217X). Missing data rates were less than 7% for most (95%) of the samples. Excluding
374   populations with high missing data rate (>7%), the proportion of sites with missing data was
positively correlated with read depth (p=1.2x10^9, R^2=0.4). The positive correlation between

376   read depth and missing data rate is surprising and likely a consequence of masking sites
       with high coverage. The number of flies per sample varied from 40 to 205, with considerable
378   heterogeneity among the DrosRTEC samples (standard deviation [sd] = 30), but not among
       DrosEU samples (sd = 0.04). Variation in the number of flies and in sequencing depth is
380   reflected in the effective read depth, an estimate of the number of independent reads after
       accounting for double binomial sampling that occurs during PoolSeq (Eff. RD; Kolaczkowski
382   *et al.* 2011; Feder *et al.* 2012; Figure 1C). There was considerable variation in PCR
       duplicate rate among samples, with notable differences between batches of DrosEU
384   samples (~6% in 2014 vs. 18% in 2015/16; t-test *p*=1.8x10^{-19}) and DrosRTEC samples (~3%
       in samples collected as part of Bergland *et al.* (2014) vs. ~14% in samples collected as part
386   of Machado *et al.* (2019; p=6.37x10-3). Curiously, the 2015/2016 DrosEU samples were
       made with a PCR-free kit, suggesting that the observed PCR duplicates were optical
388   duplicates and not amplification artifacts. Contamination of samples by *D. simulans* varied
       among populations but was generally absent (<1% *D. simulans* specific reads).
390

       Identification and quality control of SNPs. In order to determine appropriate SNP calling
392   and filtering parameters, and to identify potentially problematic population samples, we first
       calculated the ratio of non-synonymous to synonymous polymorphism (p_N/p_S) for each
394   population sample. We chose this metric because it can reflect the presence of sequencing
       errors that would disproportionately inflate p_N relative to p_S.
396   For the PoolSNP dataset, we varied the global minor allele count (MAC) and global
       minor allele frequency (MAF) and then calculated p_N/p_S. We observed that p_N/p_S was
398   negatively correlated with MAC (linear regression; *p*<0.001; Figure 2A). MAC thresholds <50
       resulted in large variances of p_N/p_S caused by 36 populations characterized by unusually
400   high p_N/p_S ratios (Supplemental Material, Table S3; Figures 2A and 2C). Some (n=21) of
       these samples had previously been found to show positive values of Tajima's *D* across the
402   whole genome (Kapun *et al.* 2020) and are characterized by a large number of private
       polymorphisms (Supplemental Material, Table S3; see below), indicating that there may be
404   elevated numbers of sequencing errors in some samples. Applying a MAC threshold of 50
       reduced the elevated p_N/p_S ratios to values similar to the rest of the dataset, and suggesting
406   that the potential sequencing errors had been largely removed. To minimize false positive
       variant calling, we therefore conservatively chose MAC=50 and MAF=0.001 as threshold
408   parameters for SNP calling with PoolSNP. Using these parameters, we identified 4,381,144
       polymorphisms segregating among the 271 *D. melanogaster* samples (Pool-Seq plus DGN),
410   and 4,042,456 polymorphisms segregating among the 246 Pool-Seq samples (excluding
       DGN), using PoolSNP.

412   SNAPE calls variants in each sample separately using a probabilistic approach, in
contrast to PoolSNP, which integrates allelic information across all populations for heuristic
414   SNP calling. To quantify the amount of putative sequencing errors among low frequency
variants we varied the local MAF threshold per sample and calculated p_N/p_S for each sample
416   in the SNAPE-pooled dataset. Similar to PoolSNP, we found that elevated p_N/p_S was
negatively correlated with a local MAF threshold (linear regression; $p<0.001$; Figure 2B) and
418   that the 36 above-mentioned problematic samples also had a strong effect on the variance
and mean of p_N/p_S ratios. Accordingly, we removed these 36 samples and applied a
420   conservative MAF filter of 5% for the remainder of the SNAPE-pooled analysis. Our results
identified 8,541,651 polymorphisms segregating among the remaining 210 samples. Below,
422   we discuss the geographic distribution and global frequency of SNPs identified using these
two methods in order to provide insight into the stark discrepancy in the number of SNPs
424   that they identify.

Figure 2. The effect of heuristic minor allele count (MAC) and minor allele frequency (MAF)
428   thresholds on p_N/p_S ratios in SNP data based on PoolSNP (A) and SNAPE-pooled (B). Blue lines in
both panels show average genome-wide p_N/p_S ratios across 271 and 246 populations, respectively.
430   The blue ribbons depict the corresponding standard deviations. The bottom panels (C) and (D),
correspond to the top panels A and B but excluding 36 problematic samples, which are characterized
432   by elevated p_N/p_S, an exceptionally large number of private SNPs and genome-wide positive Tajima's
D. Note that the y-axes of the bottom and top panels differ in scale. The two arrows show the MAC
434   and MAF thresholds used for the final datasets.

436 **Patterns of polymorphism between PoolSNP and SNAPE-pooled.** We calculated three
metrics related to the amount of polymorphism discovered by our pipelines: the abundance

438 of polymorphisms segregating in *n* populations across each chromosome (Figure 3A), the
difference of discovered polymorphisms between SNAPE-pooled and PoolSNP (defined as

440 the absolute value of PoolSNP minus SNAPE-pooled; Figure 3B), and the amount of
polymorphism discovered per minor allele frequency bin (Figure 3C). We evaluated these

442 three metrics across a 2x2 filtering scheme: two MAF filters (0.001, 0.05) and two sample
sets (the whole dataset of 246 samples; and the 210 samples that passed the sequencing

444 error filter in SNAPE-pooled; see *Identification and quality control*). Notably, PoolSNP was
biased towards identification of common SNPs present in multiple samples, whereas

446 SNAPE-pooled was more sensitive to the identification of polymorphisms that appeared in
few populations only (Figure 3B). For example, at a MAF filter of 0.001, SNAPE-pooled

448 discovered more polymorphisms that were shared in less than 25 populations (relative to
PoolSNP), and these accounted for ~79% of all polymorphisms discovered by the pipeline.

450 Likewise, at a MAF filter of 0.05, SNAPE-pooled discovered more polymorphisms that were
shared in less than 97 populations; these accounted for ~71% of all discovered

452 polymorphisms. SNAPE-pooled identifies fewer polymorphic sites that are shared among a
large number of populations than PoolSNP does because SNAPE pooled does not integrate

454 information across multiple populations. As a consequence, it can fail to identify SNPs which
are overall at low frequencies and get called as monomorphic or missing in a subset of

456 populations given the posterior-probability thresholds that we employed (see Materials and
Methods).

458   We also compared allele frequency estimates between the two callers using the
aforementioned dataset of 210 populations applying a MAF filter of 0.05 (see Supplemental

460 Material, Table S2). Among the positions identified as polymorphic by both calling methods,
our frequency estimates were consistent for the great majority of SNPs in all samples

462 analyzed (> 97% of samples). A very small proportion differed in less than 5% frequency
among both methods (< 2.3% in all samples), and very few polymorphic SNPs differed by a

464 frequency of between 5-10% (< 0.15% in all samples) or greater than 10% (< 0.03% in all
samples) (Supplemental Material, Table S4). Positions with a discordant calling represented

466 less than a 25% of all common positions in all samples (Supplemental Material, Table S4),
the majority of them being called polymorphic by PoolSNP and classified as missing data by

468 SNAPE-pooled (Supplemental Material, Table S4). This is consistent with the SNAPE-
pooled method as well as the stringent parameters used (see Materials and Methods).

Figure 3. (A) Number of polymorphic sites discovered across populations. The x-axis shows the
472   number of populations which share a polymorphic site. The y-axis corresponds to the number of
polymorphic sites shared by any number of populations, on alog10 scale. The colored lines represent
474   different chromosomes, and are stacked on top of each other. (B) The difference of discovered
polymorphisms between SNAPE-pooled and PoolSNP. (C) Number of polymorphic sites as a function
476   of allele frequency and the number of populations the polymorphisms is present in. The color gradient
represents the number of variant alleles from low to high (black to green). The x-axis is the same as in
478   A, and the y-axis is the minor allele frequency. The 2x2 filtering scheme is shown on the right side of
the figure.

Mutation-class frequencies. We estimated the percentage of mutation classes (e.g., A→C,
482   A→G, A→T, *etc.*) accepted as polymorphisms in both our SNP calling pipelines, and
classified these loci as being either "rare" (i.e., allele frequency < 5% and shared in less than
484   50 populations) or "common" (allele frequency > 5% and shared in more than 150
populations). For this analysis, we classified the minor allele as the derived allele. Figure 4A
486   shows the percentage of each mutation class for the 210 populations which passed filters in
both SNAPE-pooled and PoolSNP. In addition, we overlaid, as a horizontal line, the
488   expected mutation frequencies for rare (blue; Assaf *et al.* 2017) and common (red; Mackay
et al. 2012) mutations. For example, A→C variants are expected to be more abundant as
490   common mutations than as rare mutations, and the opposite is true for C→A variants. In
general, our SNP discovery pipelines produced mutation-class relative frequencies of rare
492   and common mutations that are consistent with empirical expectations, however, there were
some exceptions to this pattern. For example, the frequencies of the C/G rare mutation-class
494   was consistently underestimated by both callers, a phenomenon that might be related to the
known GC bias of modern sequencing machines (Benjamini and Speed 2012). The

496   correlation between SNP calling pipelines was high across both common and rare mutation

classes, with marginal discrepancies observed for rare variants (Figure 4B).

500   **Figure 4.** Frequencies of observed nucleotide polymorphism in the DEST dataset (210 populations

common to PoolSNP and SNAPE-pooled). (A) Each panel represents a mutation type. The red color

502   indicates common mutations (AF > 0.05, and common in more than 150 populations) whereas the

blue color indicates rare mutations (AF < 0.05, and shared in less than 50 populations). The dark

504   colors correspond to the PoolSNP pipeline and the soft colors correspond to the SNAPE-pooled

pipeline. The hovering red and blue horizontal lines represent the estimated mutation rates for

506   common and rare mutations, respectively. (B) Correlation between the observed mutation frequencies

seen in SNAPE-pooled and PoolSNP. The one-to-one correspondence line is shown as a black-

508   dashed diagonal. Correlation estimates (Pearson's correlation) and *p*-values for common and rare

mutations are shown.

Comparison to previously published datasets. We compared the allele frequency and

512   read depth estimates from the DEST dataset (based on PoolSNP) to previously published

estimates by Bergland *et al.* (2014), Machado *et al.* (2019), and Kapun *et al.* (2020). For

514   these datasets we employed two types of correlations, the nominal correlation (i.e.,

Pearson's correlation; CO) and the concordance correlation coefficient (CCC; Lin 1989; Liao

516   and Lewis 2000). The CCC determines how much the observed data deviate from the line of

perfect concordance (i.e., the 45 degree-line on a square scatter plot).

518   Estimates of allele frequency were strongly correlated and consistent with previously

published data. The strongest correlation of DEST allele frequencies and previously

520   published allele frequencies was observed with the data of Kapun *et al.* (2020) (average CO

and CCC > 0.99; Figure 5, top row; Supplemental Material, Figure S4). Allele frequency

522   correlations with Machado *et al.* (2019) are also generally high (average CO and CCC >

0.98; Figure 5, top row; Supplemental Material, Figure S5). Allele frequency correlations with

the data from Bergland *et al.* (2014) were lower (0.94; Supplemental Material, Figure S6), likely reflecting differences in data processing and quality control.

We also examined two aspects of read depth, i.e., nominal coverage and effective coverage. Nominal coverage is the number of reads mapping to a site that has passed

quality control. Effective coverage is the approximate number of independent reads, after accounting for double binomial sampling, and is useful for obtaining unbiased estimates of

the precision of allele frequency estimates (Kolaczkowski *et al.* 2011; Kofler *et al.* 2011a; Feder *et al.* 2012; Schlötterer *et al.* 2014). Similar to allele frequency estimates, the Pearson

correlation coefficients for both coverage and effective coverage were large (0.92, 0.95, 0.90 for Machado *et al.* (2019), Kapun *et al.* (2020), and Bergland *et al.* (2014), respectively; see

Supplemental Material, Figures S7-12), indicating that sample identity was preserved appropriately. However, the concordance correlation coefficients were substantially lower

between the datasets (0.24, 0.88, 0.79, respectively), indicating systematic differences in read depth between the DEST dataset and previously published data. Indeed, read depth

estimates were on average ~12%, ~14% and ~20% lower in the DEST dataset as compared to the previously published data in Machado *et al.* (2019), Kapun *et al.* (2020), and Bergland

et al. (2014)(2014) respectively. The lower read depth and effective read depth estimates in the DEST dataset reflects our more stringent quality control and filtering.

Figure 5. Correlations between DEST dataset and previously published datasets. Correlations

546   between allele frequencies (AF), Nominal Coverage (COV), and Effective Coverage (N_{EFF}) between

the DEST dataset (using the PoolSNP method) and three previously *Drosophila* datasets: Machado *et*

548   *al.* (2019), Kapun *et al.* (2020), and Bergland *et al.* (2014). For each dataset, we show the distribution

of two types of correlation coefficients: the nominal (Pearson's) correlation (CO; dashed lines) and the

550   concordant correlation (CCC; solid lines). In addition to the actual correlations between the datasets

(red distributions), we show the distributions of correlations estimated with random population pairs

552   (green distributions).

554   **Genetic diversity.** We estimated nucleotide diversity (π), Watterson's θ and Tajima's D for

both the PoolSNP and SNAPE-pooled datasets (Supplemental Material, Table S5). Results

556   for the African, European and North American population samples are presented in Figure 6

(also see Supplemental Material, Figure S13 for estimates by chromosome arm). All

558   estimates were positively correlated between PoolSNP and SNAPE-pooled ($p<0.001$), with

Pearson's correlation coefficients of 0.88, 0.94 and 0.73 for π, Watterson's θ, and Tajima's

560   D, respectively. Higher values of genetic diversity were obtained for the SNAPE-pooled

dataset, probably due to its higher sensitivity for detecting rare variants (see *Patterns of*

562   *polymorphism between PoolSNP and SNAPE-pooled*). Pool size had no significant effect on

the four summary statistics in European or in North American populations (GLMs, all

564    $p>0.05$), suggesting that data from populations with heterogeneous pool sizes can be safely

merged for accurate population genomic analysis.

568   **Figure 6.** Population genetic estimates for African, European and North American populations.

Shown are genome-wide estimates of (A) nucleotide diversity (π), (B) Watterson's θ and (C) Tajima's

570   D for African populations using the PoolSNP data set, and for European and North American

populations using both the PoolSNP and SNAPE-pooled (SNAPE) datasets. As can be seen from the

572   figure, estimates based on PoolSNP versus SNAPE-pooled (SNAPE) are highly correlated (see main
        text). Genetic variability is seen to be highest for African populations, followed by North American and
574   then European populations, as previously observed (e.g., see Lack *et al.* 2016; Kapun *et al.* 2020).

576   The highest levels of genetic variability were observed for ancestral African populations
        (mean π = 0.0060, mean θ = 0.0059); North American populations exhibited higher genetic
578   variability (mean π = 0.0054, mean θ = 0.0054) than European populations (mean π =
        0.0049, mean θ = 0.0048). These results are consistent with previous observations based on
580   individual genome sequencing (e.g., see Lack *et al.* 2016; Kapun *et al.* 2020). Our
        observations are also consistent with previous estimates based on pooled data from three
582   North American populations (mean π = 0.00577, mean θ = 0.00597; Fabian *et al.* 2012) and
        48 European populations (mean π = 0.0051, mean θ = 0.0052; Kapun *et al.* 2020).
584   Estimates of Tajima's *D* were positive when using PoolSNP, and slightly negative using
        SNAPE. These results are expected given biases in the detection of rare alleles between
586   these two SNP calling methods. In addition, our estimates for π, Watterson's θ and Tajima's
        D were positively correlated with previous estimates for the 48 European populations
588   analyzed by Kapun *et al.* (2020) (all $p<0.01$). Notably, slightly lower levels of Tajima's *D* in
        North America compared to both Africa and Europe (Figure 6B) may be indicative for
590   admixture (Stajich and Hahn 2005) which has been identified previously along the North
        American east coast (Caracristi and Schlötterer 2003; Kao *et al.* 2015; Bergland *et al.* 2016).
592

        Phylogeographic clusters in *D. melanogaster*. We performed PCA on the PoolSNP
594   variants in order to include samples from North America (DrosRTEC), Europe (DrosEU), and
        Africa (DGN) datasets (excluding all Asian and Oceanian samples). Prior to analysis we
596   filtered the joint datasets to include only high-quality biallelic SNPs. Because LD decays
        rapidly in *Drosophila* (Comeron *et al.* 2012), we only considered SNPs at least 500 bp away
598   from each other. PCA on the resulting 100,000 SNPs revealed evidence for discrete
        phylogeographic clusters that correspond to geographic regions (Supplemental Material,
600   Figure S14B). PC1 (24% variance explained [VE]) partitions samples between Africa and the
        other continents (Figure 7A). PC2 (9% VE) separates European from North American
602   populations, and both PC2 and PC3 (4% VE) divide Europe into two population clusters
        (Figure 7B). Notably, these spatial relationships become evident when PCA projections from
604   each sample are plotted onto a world map (Figure 7C). Interestingly, the emergent clusters
        in Europe are not strictly defined by geography. For example, the western cluster (diamonds
606   in Figure 7D) includes Western Europe as well as Finland, Turkey, Cyprus, and Egypt. The
        eastern cluster, on the other hand, consists of several populations collected in previous
608   Soviet republics as well as Poland, Hungary, Serbia and Austria, raising the possibility that

recent geo-political division in Europe could have affected migration and population
610   structure. Whether this result arises as a relic of recent geopolitical history within Europe,
more ancient migration and colonization (e.g., following post-glacial range expansion, Kapun
612   *et al.* 2020), local adaptation, or sampling strategy (Novembre and Stephens 2008; cf.
Kapun *et al.* 2020) remains unknown. Future targeted sampling is needed to resolve these
614   alternative explanations.

618   **Figure 7.** Demographic signatures of the DrosEU, DrosRTEC, and DGN data (using the PoolSNP
pipeline). (A) PCA dimensions 1 and 2. The mean centroid of a country's assignment is labeled. (B)
620   PCA dimensions 1 and 3. (C) Projections of PC1 onto a World map. PC1 projections define the
existence of continental level clusters of population structure (indicated by the shapes circles: Africa;
622   triangles: North America; diamonds and squares: Europe). (D) Projections of PC3 onto Europe. These
projections show the existence of a demographic divide within Europe: the diamond shapes indicate a
624   western cluster, whereas the squares represent an eastern cluster. For panels C and D, the intensity
of the color is proportional to the PC projection. The black dashed line shows the two-cluster divide.

A unique feature of this dataset is that it contains a mixture of Pool-Seq and inbred (or
628   haploid) genome data. For some geographic regions, the DEST dataset contains both data
types. Inbred and Pool-Seq samples from nearby geographic regions clustered in the same
630   regions of PC space (Supplemental Material, Figure S15). Excluding the DGN-derived
African samples, no PC was significantly correlated with data type (PC1 *p* = 0.352, PC2 *p* =
632   0.223, PC3 *p* = 0.998).

634   **Geographic proximity analysis.** The geographic distribution of our samples allows
leveraging basic principles of phylogeography and population genetics to assess the
636   biological significance of rare SNPs (Wright 1943; Battey *et al.* 2020). Accordingly, we
expect to observe young neutral alleles at low frequencies among geographically close
638   populations. We tested this hypothesis by estimating the average geographic distance
among pairs of populations that share SNPs only occurring in these two populations
640   (doubletons), among three populations that share tripletons, and so forth. Without imposing
a MAF filter, both SNAPE-pooled and PoolSNP pipelines produced patterns concordant with
642   the expectation. Populations in close proximity were more likely to share rare mutations
relative to random chance pairings (Figure 8A). Notably, the PoolSNP dataset showed an
644   elevated number of rare alleles, which violate the phylogeographic expectation (Figure 8A);
however, this only affects 0.31% of all PoolSNP mutations. To further evaluate this pattern,
646   we estimated the probability that any given population pair belongs to a particular
phylogeographic cluster (Supplemental Material, Figure S16) as a function of their shared
648   variants. Our results indicate that rare variants, private to geographically proximate
populations, are strong predictors of phylogeographic provenance (see Figure 8B).

Figure 8. Geographic Proximity Analysis. (A) Average geographic distance between populations that
652   share a polymorphism at any given site for PoolSNP and SNAPE-pooled. The x-axis represents the
number of populations considered; the y-axis is the mean geographic distance among samples. The
654   yellow line represents the random expectation calculated as random pairings of the data. The band
around the lines is the standard deviation of the estimator. (B) Probability that all populations
656   containing a polymorphic site come from the same phylogeographic cluster (Supplemental Figure 14).
The y-axis is the probability of "x" populations belonging to the same phylogeographic cluster. The
658   axis only shows up to 40 populations; after that point, the probability approaches 0. The colors are
consistent across panels.

Demography-informative markers. An inherent strength of our broad biogeographic
662   sampling is the potential to generate a panel of core demography SNPs to investigate the
provenance of current and future samples. We created a panel of demography-informative
664   markers (DIMs) by conducting a DAPC to discover which loci drive the phylogeographic
signal in the dataset. We trained two separate DAPC models: the first utilized the four
666   phylogeographic clusters identified by principal components (PCs; Figure 6AB,
Supplemental Material, Figure S16, Table S1); the second utilized the geographic localities
668   where the samples were collected (i.e., countries in Europe and the US states). This
optimization indicated that the information contained in the first 40 PCs maximizes the
670   probability of successful assignment (Figure 9A). This resulted in the inclusion of 30,000
DIMs, most of which were strongly associated with PCs 1-3 (Figure 9B inset). Moreover, the
672   correlations were larger among the first 3 PCs and decayed monotonically for the additional
PCs (Figure 9B). Lastly, our DIMs were uniformly distributed across the fly genome (Figure
674   9C).

We assessed the accuracy of our DIM panel using a leave-one-out cross-validation
676   approach (LOOCV). We trained the DAPC model using all but one sample and then
classified the excluded sample. We performed LOOCV separately for the phylogeographic
678   cluster groups, as well as for the state/country labels. The phylogeographic model used all
DrosRTEC, DrosEU, and DGN samples (excluding Asia and Oceania with too few
680   individuals per sample); the state/country model used only samples for which each label had
at least 3 or more samples. Our results showed that the model is 100% accurate in terms of
682   resolving samples at the phylogeographic cluster level (Figure 9D) and 89% at the
state/country level (Figure 9E). We anticipate that this set of DIMs will be useful for future
684   analysis of geographic provenance of North American and European samples. We provide a
tutorial on the usage of the DIM in Supplemental Methods.

Figure 9. Demography-informative markers. (A) Number of retained PCs which maximize the DAPC model's capacity to assign group membership. Model trained on the phylogeographic clusters (dashed lines) or the country/state labels (solid line). (B) Absolute correlation for the 33,000 individual SNPs with highest weights onto the first 40 components of the PCA. Inset: Number of SNPs per PC. (C) Location of the 33,000 most informative demographic SNPs across the chromosomes. (D) LOOCV of the DAPC model trained on the phylogeographic clusters. (E) LOOCV of the DAPC model trained on the phylogeographic state/country labels. For panels D and E, the y-axis shows the highest posterior produced by the prediction model and the x-axis is the posterior assigned to the actual label classification of the sample. Also, for D and E, marginal histograms are shown.

Conclusions and Outlook

Here we have presented a new, modular and unified bioinformatics pipeline for processing, integrating and analyzing SNP variants segregating in population samples of *D. melanogaster*. We have used this pipeline to assemble the largest worldwide data repository of genome-wide SNPs in *D. melanogaster* to date, based both on previously published data (DGN: Africa; Lack *et al.* 2015, 2016) as well as on new data collected by our two collaborating consortia (DrosRTEC: mostly North America; Machado *et al.* 2019; DrosEU: mostly Europe; Kapun *et al.* 2020). We assembled this dataset using two SNP calling strategies that differ in their ability to identify rare polymorphisms, thereby enabling future work studying the evolutionary history of this species. We are dubbing this data repository and the supporting bioinformatics tools *Drosophila Evolution over Space and Time* (DEST).

One of the biggest challenges in the present "omics" era is the rapidly growing number of complex large-scale datasets which require technically elaborate bioinformatics know-how to become accessible and utilizable. This hurdle often prohibits the exploitation of already

available genomics datasets by scientists without a strong bioinformatics or computational background. To remedy this situation for the Drosophila evolution community, our

bioinformatics pipeline is provided as a Docker image (to standardize across software versions, as well as make the pipeline independent of specific operating systems) and a new

genome browser makes our SNP dataset available through an easy-to-use web interface (see Supplemental Information Figures S2, S3; available at https://dest.bio).

The DEST data repository and platform will enable the population genomics community to address a variety of longstanding, fundamental questions in ecological and evolutionary

genetics. The current dataset might for instance be valuable for providing a more accurate picture of the demographic history of *D. melanogaster* populations, in particular in Europe

and North America, and with respect to multiple bouts of out-of-Africa migration and recent patterns of admixture.

The DEST dataset will likewise be useful for an improved understanding of the genomic signatures underlying both global and local adaptation, including a more fine-grained view of

selective sweeps, their evolutionary origin and distribution (e.g., see Glinka *et al.* 2003; Beisswanger *et al.* 2006; Ometto 2010; Stephan 2016; Kapun *et al.* 2020). In terms of local

adaptation, the broad spatial sampling across latitudinal and longitudinal gradients on the North American and European continents, encompassing a broad range of climate zones

and areas of varying degrees of seasonality, will allow examining the parallel nature of local (clinal) adaptation in response to similar environmental factors in greater depth than possible

before (e.g., Turner *et al.* 2008; Kolaczkowski *et al.* 2011; Fabian *et al.* 2012; Reinhardt *et al.* 2014; Kapun *et al.* 2016, 2020; Machado *et al.* 2019; Bogaerts-Márquez *et al.* 2020;

Waldvogel *et al.* 2020).

       Another major opportunity provided by the DEST dataset lies in studying the temporal

736   dynamics of evolutionary change. Sampling at dozens of localities across the growing season and over multiple years will help to advance our understanding of the short-term

738   population and evolutionary dynamics of flies living in diverse environments, thereby providing novel insights into the nature of temporally varying selection (e.g., Wittmann *et al.*;

740   Bergland *et al.* 2014; Machado *et al.* 2019) and evolutionary responses to climate change (e.g., Umina 2005; Rodríguez-Trelles *et al.* 2013; Waldvogel *et al.* 2020).

742   Moreover, by integrating these worldwide estimates of allele frequencies, those from lab- and field-based 'evolve and resequence' (E&R; Turner *et al.* 2011; reviewed in Kofler

744   and Schlötterer 2014; Schlötterer *et al.* 2014; Flatt 2020) and mesocosm experiments (e.g.,

Rudman *et al.* 2019; Erickson *et al.* 2020), we might be able to gain deeper insights into the
746   genetic basis and evolutionary history of variation in fitness components (e.g., Flatt 2020).

The real value of the DEST dataset lies in the future: its long-term utility will grow as
748   natural and experimental populations are continually being sampled, resequenced and
added to the repository by the community of *Drosophila* evolutionary geneticists. The
750   pipeline that we have established will make future updates to the data-repository
straightforward. Furthermore, since it is not easily feasible for any single research group to
752   sample flies densely through time and across a broad geographic range, the growing value
of the DEST dataset will depend upon the synergistic collaboration among research groups
754   across the globe, as exemplified by the DrosRTEC and DrosEU consortia. Importantly, in an
era of rapidly decreasing sequencing costs, comprehensive population genomic analyses
756   are no longer limited by genetic marker density but by the availability of biological samples
from standardized, collaborative long-term collection efforts through space and time (e.g.,
758   Machado *et al.* 2019; Kapun *et al.* 2020). In this vein, the collaborative framework presented
here might allow us, as a global community, to fill some important gaps in the current data
760   repository: for example, many areas of the world (notably Asia and South America) remain
largely uncharted territory in *Drosophila* population genomics, and the addition of phased
762   sequencing data (e.g., providing information on haplotypes, LD, linked selection) will be
crucially important for future analyses of demography, selection and their interplay.
764   We are convinced that the DEST platform will become a valuable and widely-used
resource for scientists interested in *Drosophila* evolution and genetics, and we actively
766   encourage the community to join the collaborative effort we are seeking to build.

768   **Data availability**
All scripts to make figures and perform analyses associated with this manuscript are
770   available here: https://github.com/DEST-bio/data-paper. All scripts to build the dataset,
including the mapping pipeline, SNP calling scripts, and meta-data are available here:
772   https://github.com/DEST-bio/DEST_freeze1. All output from the DEST pipeline, including
intermediate output files, metadata, etc. can be found here: https://dest.bio. The genome
774   browser   associated   with   the   DEST   dataset   can   be   found   here:
http://dgvbrowser.uab.cat/dest/browser/. The mapping and SNP calling pipeline can be
776   found here: https://hub.docker.com/r/destbiodocker/destbiodocker

778   **Acknowledgements**
We are grateful to all the members of the DrosEU and DrosRTEC consortia for their long-
780   standing support, collaboration and for discussion. DrosEU is funded by a Special Topic

Networks (STN) grant from the European Society for Evolutionary Biology (ESEB). MK (M.
Kapun) was supported by the Austrian Science Foundation (grant no. FWF P32275); JG by
the European Research Council (ERC) under the European Union's Horizon 2020 research
and innovation programme (H2020-ERC-2014-CoG-647900) and by the Spanish Ministry of
Science and Innovation (BFU-2011-24397); TF by the Swiss National Science Foundation
(SNSF grants PP00P3_133641, PP00P3_165836, and 31003A_182262) and a Mercator
Fellowship from the German Research Foundation (DFG), held as a EvoPAD Visiting
Professor at the Institute for Evolution and Biodiversity, University of Mu□nster; AOB by the
National Institutes of Health (R35 GM119686); MK (M. Kankare) by Academy of Finland
grant 322980; VL by Danish Natural Science Research Council (FNU) grant 4002-00113B;
FS Deutsche Forschungsgemeinschaft (DFG) grant STA1154/4-1, Project 408908608; JP
by the Deutsche Forschungsgemeinschaft Projects 274388701 and 347368302; AU by FPI
fellowship (BES-2012-052999); ET Israel Science Foundation (ISF) grant 1737/17; MSV,
MSR and MJ by a grant from the Ministry of Education, Science and Technological
Development of the Republic of Serbia (451-03-68/2020-14/200178); AP, KE and MT by a
grant from the Ministry of Education, Science and Technological Development of the
Republic of Serbia (451-03-68/2020-14/200007); and TM NSERC grant RGPIN-2018-05551.

Author contributions

Martin Kapun: Conceptualization, Data curation, Formal Analysis, Funding acquisition,
Investigation, Methodology, Project Administration, Resources, Software, Supervision,
Visualization, Writing - original draft, Writing - review & editing. Joaquin Nunez: Formal
Analysis, Methodology, Software, Visualization, Writing - original draft, Writing - review &
editing. María Bogaerts-Márquez: Formal Analysis, Methodology, Software, Visualization,
Writing - original draft, Writing - review & editing. Jesús Murga-Moreno: Formal Analysis,
Methodology, Software, Visualization, Writing - original draft, Writing - review & editing.
Margot Paris: Formal Analysis, Methodology, Software, Visualization, Writing - original draft,
Writing - review & editing. Joseph Outten: Software, Writing - review & editing. Marta
Coronado-Zamora: Formal Analysis, Methodology, Software, Visualization, Writing - original
draft, Writing - review & editing. Aleksandra Patenkovic: Resources. Amanda Glaser-
Schmitt: Resources. Anna Ullastres: Resources. Antonio J. Buendía-Ruíz: Resources. Banu
S. Onder: Resources. Brian P Lazzaro: Resources, Writing - review & editing. Catherine
Montchamp-Moreau: Resources. Christopher W. Wheat: Resources, Writing - review &
editing. Cristina P. Vieira: Resources, Writing - review & editing. Daniel K. Fabian:
Resources. Darren J. Obbard: Resources. Dmitry V. Mukha: Resources. Dorcas J. Orengo:
Resources, Writing - review & editing. Elena Pasyukova: Resources. Eliza Argyridou:

Resources. <u>Emily L. Behrman</u>: Resources, Writing - review & editing. <u>Eran Tauber</u>:
818   Resources. <u>Eva Puerma</u>: Resources, Writing - review & editing. <u>Fabian Staubach</u>:
Resources, Writing - review & editing. <u>Francisco D Gallardo-Jiménez</u>: Resources. <u>Iryna</u>
820   <u>Kozeretska</u>: Resources. <u>J. Roberto Torres</u>: Resources. <u>Jessica K. Abbott</u>: Resources. <u>John</u>
<u>Parsch</u>: Funding acquisition, Resources, Writing - review & editing. <u>Jorge Vieira</u>: Resources,
822   Writing - review & editing. <u>M. Josefa Gómez-Julián</u>: Resources. <u>Katarina Eric</u>: Resources.
<u>Kelly A. Dyer</u>: Resources. <u>Lain Guio</u>: Resources. <u>Lino Ometto</u>: Writing - review & editing. <u>M.</u>
824   <u>Luisa Espinosa-Jimenez</u>: Resources. <u>Maaria Kankare</u>: Resources, Writing - review &
editing. <u>Mads F. Schou</u>: Resources, Writing - review & editing. <u>Maria P. García Guerreiro</u>:
826   Resources, Writing - review & editing. <u>Marija Savic Veselinovic</u>: Resources. <u>Marija</u>
<u>Tanaskovic</u>: Resources. <u>Marina Stamenkovic-Radak</u>: Funding acquisition, Resources.
828   <u>Mihailo Jelic</u>: Resources. <u>Miriam Merenciano</u>: Resources. <u>Oleksandr M. Maistrenko</u>: Writing
- review & editing. <u>Omar Rota-Stabelli</u>: Resources. <u>Sara Guirao-Rico</u>: Resources, Writing -
830   review & editing. <u>Sònia Casillas:</u> Resources, Writing - review & editing. <u>Sonja Grath</u>:
Resources. <u>Stephen W. Schaeffer</u>: Resources. <u>Subhash Rajpurohit</u>: Resources. <u>Svitlana V.</u>
832   <u>Serga</u>: Resources. <u>Thomas J.S. Merritt</u>: Resources. <u>Vivien Horváth</u>: Resources. <u>Vladimir E.</u>
<u>Alatortsev</u>: Resources. <u>Volker Loeschcke</u>: Resources. <u>Yun Wang</u>: Resources. <u>Antonio</u>
834   <u>Barbadilla</u>: Software, Writing - review & editing. <u>Dmitri Petrov</u>: Conceptualization, Funding
acquisition, Project Administration, Resources, Writing - review & editing. <u>Paul Schmidt</u>:
836   Conceptualization, Funding acquisition, Project Administration, Resources, Writing - review
& editing. <u>Josefa Gonzalez</u>: Conceptualization, Funding acquisition, Project Administration,
838   Resources, Supervision, Writing - original draft, Writing - review & editing. <u>Thomas Flatt</u>:
Conceptualization, Funding acquisition, Project Administration, Resources, Supervision,
840   Writing - original draft, Writing - review & editing. <u>Alan Bergland</u>: Conceptualization, Data
curation, Formal Analysis, Funding acquisition, Investigation, Methodology, Project
842   Administration, Resources, Software, Supervision, Visualization, Writing - original draft,
Writing - review & editing

Competing interests. The authors declare no competing interests.

References

848   Adams, M. D., 2000 The Genome Sequence of *Drosophila melanogaster*. Science 287:
        2185–2195.

850   Andrews, S., 2010 FastQC: A Quality Control Tool for High Throughput Sequence Data.
        Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

852   Arguello, J. R., S. Laurent, and A. G. Clark, 2019 Demographic History of the Human
               Commensal *Drosophila melanogaster*. Genome Biol Evol 11: 844–854.

854   Assaf, Z. J., S. Tilk, J. Park, M. L. Siegal, and D. A. Petrov, 2017 Deep sequencing of
               natural and experimental populations of *Drosophila melanogaster* reveals biases in
856   the spectrum of new mutations. Genome Res 27: 1988–2000.

         Bastide, H., A. Betancourt, V. Nolte, R. Tobler, P. Stöbe *et al.*, 2013 A Genome-Wide, Fine-
858   Scale Map of Natural Pigmentation Variation in *Drosophila melanogaster*. PLoS
               Genet 9: e1003534.

860   Battey, C. J., P. L. Ralph, and A. D. Kern, 2020 Space is the Place: Effects of Continuous
               Spatial Structure on Analysis of Population Genetic Data. Genetics 215: 193–214.

862   Behrman, E. L., V. M. Howick, M. Kapun, F. Staubach, A. O. Bergland *et al.*, 2018 Rapid
               seasonal evolution in innate immunity of wild *Drosophila melanogaster*. Proc Royal
864   Soc B 285: 20172599.

         Beisswanger, S., W. Stephan, and D. Lorenzo, 2006 Evidence for a Selective Sweep in the
866   *wapl* Region of *Drosophila melanogaster*. Genetics 172: 265–274.

         Benjamini, Y., and T. P. Speed, 2012 Summarizing and correcting the GC content bias in
868   high-throughput sequencing. Nucleic Acids Res 40: e72.

         Bergland, A. O., E. L. Behrman, K. R. O'Brien, P. S. Schmidt, and D. A. Petrov, 2014
870   Genomic Evidence of Rapid and Stable Adaptive Oscillations over Seasonal Time
               Scales in *Drosophila*. PLoS Genet 10: e1004775.

872   Bergland, A. O., R. Tobler, J. González, P. Schmidt, and D. Petrov, 2016 Secondary contact
               and local adaptation contribute to genome-wide patterns of clinal variation in
874   *Drosophila melanogaster*. Mol Ecol 25: 1157–1174.

         Bogaerts-Márquez, M., S. Guirao-Rico, M. Gautier, and J. González, 2020 Temperature,
876   rainfall and wind variables underlie environmental adaptation in natural populations
               of *Drosophila melanogaster.* Mol Ecol, in press (https://doi.org/10.1111/mec.15783).

878   Buels, R., E. Yao, C. M. Diesh, R. D. Hayes, M. Munoz-Torres *et al.*, 2016 JBrowse: a
               dynamic web platform for genome visualization and analysis. Genome Biol 17: 66.

880   Burke, M. K., 2012 How does adaptation sweep through the genome? Insights from long-
               term selection experiments. Proc Royal Soc B 279: 5029–5038.

882   Bushnell, B., J. Rood, and E. Singer, 2017 BBMerge – Accurate paired shotgun read
               merging via overlap. PLoS ONE 12: e0185056.

884   Campo, D., K. Lehmann, C. Fjeldsted, T. Souaiaia, J. Kao *et al.*, 2013 Whole-genome
               sequencing of two North American *Drosophila melanogaster* populations reveals
886   genetic differentiation and positive selection. Mol Ecol 22: 5084–5097.

         Capy, P., and P. Gibert, 2004 *Drosophila melanogaster, Drosophila simulans*: so similar yet
888   so different. Genetica 120(1-3): 5-16.

Caracristi, G., and C. Schlötterer, 2003 Genetic Differentiation Between American and

890   European *Drosophila melanogaster* Populations Could Be Attributed to Admixture of

African Alleles. Mol Biol Evol 20: 792–799.

892   Celniker, S. E., and G. M. Rubin, 2003 The *Drosophila melanogaster* genome. Annu Rev

Genomics Hum Genet 4: 89–117.

894   Cheng, C., B. J. White, C. Kamdem, K. Mockaitis, C. Costantini *et al.*, 2012 Ecological

Genomics of *Anopheles gambiae* Along a Latitudinal Cline: A Population-

896   Resequencing Approach. Genetics 190: 1417–1432.

Cingolani, P., A. Platts, L. Wang, M. Coon, T. Nguyen *et al.*, 2012 A program for annotating

898   and predicting the effects of single nucleotide polymorphisms, SnpEff. Fly 6: 80–92.

Comeron, J. M., R. Ratnappan, and S. Bailin, 2012 The Many Landscapes of Recombination

900   in *Drosophila melanogaster*. PLoS Genet 8: e1002905.

Corbett-Detig, R., and R. Nielsen, 2017 A Hidden Markov Model Approach for

902   Simultaneously Estimating Local Ancestry and Admixture Time Using Next

Generation Sequence Data in Samples of Arbitrary Ploidy. PLoS Genet 13:

904   e1006529.

Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks *et al.*, 2011 The variant call

906   format and VCFtools. Bioinformatics 27: 2156–2158.

David, J. R., and P. Capy, 1988 Genetic variation of *Drosophila melanogaster* natural

908   populations. Trends Genet 4: 106–111.

Deitz, K. C., G. A. Athrey, M. Jawara, H. J. Overgaard, A. Matias *et al.*, 2016 Genome-Wide

910   Divergence in the West-African Malaria Vector *Anopheles melas*. G3 6: 2867–2879.

Duchen, P., D. Živković, S. Hutter, W. Stephan, and S. Laurent, 2013 Demographic

912   Inference Reveals African and European Admixture in the North American

Drosophila melanogaster Population. Genetics 193: 291–301.

914   Duffy, J. B., 2002 GAL4 system in *Drosophila*: a fly geneticist's Swiss army knife. Genesis

34: 1–15.

916   Durmaz, E., C. Benson, M. Kapun, P. Schmidt, and T. Flatt, 2018 An inversion supergene in

Drosophila underpins latitudinal clines in survival traits. J Evolution Biol 31: 1354–

918   1364.

Durmaz, E., S. Rajpurohit, N. Betancourt, D. K. Fabian, M. Kapun *et al.*, 2019 A clinal

920   polymorphism in the insulin signaling transcription factor *foxo* contributes to life-

history adaptation in *Drosophila*. Evolution 73: 1774-1792.

922   Erickson, P. A., C. A. Weller, D. Y. Song, A. S. Bangerter, P. Schmidt *et al.*, 2020 Unique

genetic signatures of local adaptation over space and time for diapause, an

924   ecologically relevant complex trait, in *Drosophila melanogaster*. PLoS Genet 16:

e1009110.

926   Fabian, D. K., M. Kapun, V. Nolte, R. Kofler, P. S. Schmidt *et al.*, 2012 Genome-wide
                patterns of latitudinal differentiation among populations of *Drosophila melanogaster*
928   from North America. Mol Ecol 21: 4748–4769.

        Feder, A. F., D. A. Petrov, and A. O. Bergland, 2012 LDx: Estimation of Linkage
930   Disequilibrium from High-Throughput Pooled Resequencing Data. PLoS ONE 7:
                e48588.

932   Ferretti, L., S. E. Ramos-Onsins, and M. Pérez-Enciso, 2013 Population genomics from pool
                sequencing. Mol Ecol 22: 5561–5576.

934   Flatt, T., 2016 Genomics of clinal variation in *Drosophila*: disentangling the interactions of
                selection and demography. Mol Ecol 25: 1023–1026.

936   Flatt, T., 2020 Life-History Evolution and the Genetics of Fitness Components in *Drosophila
                melanogaster*. Genetics 214: 3–48.

938   Gautier, M., J. Foucaud, K. Gharbi, T. Cézard, M. Galan *et al.*, 2013 Estimation of population
                allele frequencies from next-generation sequencing data: pool-versus individual-
940   based genotyping. Mol Ecol 22: 3766–3779.

        Giesen, A., W. U. Blanckenhorn, M. A. Schäfer, K. K. Shimizu, R. Shimizu-Inatsugi *et al.*,
942   2020 Genomic signals of admixture and reinforcement between two closely related
                species of European sepsid flies. Preprint: bioRxiv 2020.03.11.985903.

944   Glinka, S., L. Ometto, S. Mousset, W. Stephan, and D. Lorenzo, 2003 Demography and
                natural selection have shaped genetic variation in *Drosophila melanogaster*: a multi-
946   locus approach. Genetics 165: 1269–1278.

        Gould, B. A., Y. Chen, and D. B. Lowry, 2017 Pooled Ecotype Sequencing Reveals
948   Candidate Genetic Mechanisms for Adaptive Differentiation and Reproductive
                Isolation. Mol Ecol 26: 163–177.

950   Grenier, J. K., J. R. Arguello, M. C. Moreira, S. Gottipati, J. Mohammed *et al.*, 2015 Global
                Diversity Lines–A Five-Continent Reference Panel of Sequenced *Drosophila
952   melanogaster* Strains. G3 5: 593–603.

        Guirao-Rico, S., and J. González, 2021 Benchmarking the performance of Pool-seq SNP
954   callers using simulated and real sequencing data. Mol Ecol Res, in press.

        Guirao-Rico, S., and J. González, 2019 Evolutionary insights from large scale resequencing
956   datasets in *Drosophila melanogaster*. Curr Opin Insect Sci 31: 70–76.

        Hales, K. G., C. A. Korey, A. M. Larracuente, and D. M. Roberts, 2015 Genetics on the Fly:
958   A Primer on the *Drosophila* Model System. Genetics 201: 815–842.

        Haudry, A., S. Laurent, and M. Kapun, 2020 Population genomics on the fly: recent
960   advances in *Drosophila*. Methods Mol Biol 290: 357–396.

Hijmans, R. J., S. E. Cameron, J. L. Parra, P. G. Jones, and A. Jarvis, 2005 Very high
962   resolution interpolated climate surfaces for global land areas. Int J Climatol 25:
          1965–1978.

964 Hivert, V., R. Leblois, E. J. Petit, M. Gautier, and R. Vitalis, 2018 Measuring Genetic
          Differentiation from Pool-seq Data. Genetics 210: 315–330.

966 Hoban, S., J. L. Kelley, K. E. Lotterhos, M. F. Antolin, G. Bradburd *et al.*, 2016 Finding the
          Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions, and Future
968   Directions. Am Nat 188: 379–397.

Hu, T. T., M. B. Eisen, K. R. Thornton, and P. Andolfatto, 2013 A second-generation
970   assembly of the *Drosophila simulans* genome provides new insights into patterns of
          lineage-specific divergence. Genome Res 23: 89–98.

972 Jennings, B. H., 2011 *Drosophila* – a versatile model in biology & medicine. Materials Today
          14: 190–195.

974 Jombart, T., S. Devillard, and F. Balloux, 2010 Discriminant analysis of principal
          components: a new method for the analysis of genetically structured populations.
976   BMC Genetics 11: 94.

de Jong, G., and Z. Bochdanovits, 2003 Latitudinal clines in *Drosophila melanogaster*: Body
978   size, allozyme frequencies, inversion frequencies, and the insulin-signalling pathway.
          J Genet 82: 207–223.

980 Kao, J. Y., A. Zubair, M. P. Salomon, S. V. Nuzhdin, and D. Campo, 2015 Population
          genomic analysis uncovers African and European admixture in *Drosophila*
982   *melanogaster* populations from the south-eastern United States and Caribbean
          Islands. Mol Ecol 24: 1499–1509.

984 Kapopoulou, A., M. Kapun, B. Pieper, P. Pavlidis, R. Wilches *et al.*, 2020 Demographic
          analyses of a new sample of haploid genomes from a Swedish population of
986   *Drosophila melanogaster*. Sci Rep 10: 22415.

Kapun, M., M. G. Barrón, F. Staubach, D. J. Obbard, R. A. W. Wiberg *et al.*, 2020 Genomic
988   Analysis of European *Drosophila melanogaster* Populations Reveals Longitudinal
          Structure, Continent-Wide Selection, and Previously Unknown DNA Viruses. Mol Biol
990   Evol 37: 2661–2678.

Kapun, M., D. K. Fabian, J. Goudet, and T. Flatt, 2016 Genomic Evidence for Adaptive
992   Inversion Clines in *Drosophila melanogaster*. Mol Biol Evol 33: 1317–1336.

Keller, A., 2007 *Drosophila melanogaster*'s history as a human commensal. Curr Biol 17:
994   R77–R81.

Kent, W. J., A. S. Zweig, G. Barber, A. S. Hinrichs, and D. Karolchik, 2010 BigWig and
996   BigBed: enabling browsing of large distributed datasets. Bioinformatics 26: 2204–
          2207.

998   Koboldt, D. C., K. Chen, T. Wylie, D. E. Larson, M. D. McLellan *et al.*, 2009 VarScan: variant
           detection in massively parallel sequencing of individual and pooled samples.
1000   Bioinformatics 25: 2283–2285.

       Koboldt, D. C., Q. Zhang, D. E. Larson, D. Shen, M. D. McLellan *et al.*, 2012 VarScan 2:
1002   Somatic mutation and copy number alteration discovery in cancer by exome
           sequencing. Genome Res 22: 568–576.

1004   Kofler, R., P. Orozco-terWengel, N. De Maio, R. V. Pandey, V. Nolte *et al.*, 2011a
           PoPoolation: A Toolbox for Population Genetic Analysis of Next Generation
1006   Sequencing Data from Pooled Individuals. PLoS ONE 6: e15925.

       Kofler, R., R. V. Pandey, and C. Schlotterer, 2011b PoPoolation2: identifying differentiation
1008   between populations using sequencing of pooled DNA samples (Pool-Seq).
           Bioinformatics 27: 3435–3436.

1010   Kofler, R., and C. Schlötterer, 2014 A guide for the design of evolve and resequencing
           studies. Mol Biol Evol 31: 474–483.

1012   Kolaczkowski, B., A. D. Kern, A. K. Holloway, and D. J. Begun, 2011 Genomic Differentiation
           Between Temperate and Tropical Australian Populations of *Drosophila
1014   melanogaster.* Genetics 187: 245–260.

       Kuhn, R. M., D. Haussler, and W. J. Kent, 2013 The UCSC genome browser and associated
1016   tools. Brief Bioinform 14: 144–161.

       Lachaise, D., M.-L. Cariou, J. R. David, F. Lemeunier, L. Tsacas *et al.*, 1988 Historical
1018   Biogeography of the *Drosophila melanogaster* Species Subgroup, pp. 159–225 in
           Evolutionary Biology, edited by M. K. Hecht, B. Wallace, and G. T. Prance. Springer
1020   US, Boston, MA.

       Lack, J. B., C. M. Cardeno, M. W. Crepeau, W. Taylor, R. B. Corbett-Detig *et al.*, 2015 The
1022   *Drosophila* Genome Nexus: A Population Genomic Resource of 623 *Drosophila
           melanogaster* Genomes, Including 197 from a Single Ancestral Range Population.
1024   Genetics 199: 1229–1241.

       Lack, J. B., J. D. Lange, A. D. Tang, C.-D. B Russell, and J. E. Pool, 2016 A Thousand Fly
1026   Genomes: An Expanded *Drosophila* Genome Nexus. Mol Biol Evol 33: 3308-3313.

       Langley, C. H., K. Stevens, C. Cardeno, Y. C. G. Lee, D. R. Schrider *et al.*, 2012 Genomic
1028   variation in natural populations of *Drosophila melanogaster.* Genetics 192: 533–598.

       Li, H., 2013 Aligning sequence reads, clone sequences and assembly contigs with BWA-
1030   MEM. Preprint: arXiv:1303.3997 [q-bio.GN].

       Li, H., 2011 Tabix: fast retrieval of sequence features from generic TAB-delimited files.
1032   Bioinformatics 27: 718–719.

       Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The Sequence
1034   Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079.

Li, H., and W. Stephan, 2006 Inferring the Demographic History and Rate of Adaptive
1036   Substitution in *Drosophila*. PLoS Genet 2: 10.

Liao, J. J. Z., and J. W. Lewis, 2000 A Note on Concordance Correlation Coefficient. PDA J
1038   Pharm Sci Technol 54: 23–26.

Lin, L. I.-K., 1989 A Concordance Correlation Coefficient to Evaluate Reproducibility.
1040   Biometrics 45: 255–268.

Lynch, M., D. Bost, S. Wilson, T. Maruki, and S. Harrison, 2014 Population-Genetic
1042   Inference from Pooled-Sequencing Data. Genome Biol Evol 6: 1210–1218.

Machado, H. E., A. O. Bergland, K. R. O'Brien, E. L. Behrman, P. S. Schmidt *et al.*, 2016
1044   Comparative population genomics of latitudinal variation in *Drosophila simulans* and
               Drosophila melanogaster. Mol Ecol 25: 723–740.

1046 Machado, H. E., A. O. Bergland, R. Taylor, S. Tilk, E. Behrman *et al.*, 2019 Broad
               geographic sampling reveals predictable, pervasive, and strong seasonal adaptation
1048   in *Drosophila*. Preprint: bioRxiv 337543.

Mackay, T. F. C., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles *et al.*, 2012 The
1050   *Drosophila melanogaster* Genetic Reference Panel. Nature 482: 173–178.

Markow, T. A., and P. M. O'Grady, 2005 *Drosophila: a guide to species identification and
1052   use*. Academic Press (Elsevier), Amsterdam, Boston.

Martin, M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing
1054   reads. EMBnet.journal 17: 10–12.

Mateo, L., G. E. Rech, and J. González, 2018 Genome-wide patterns of local adaptation in
1056   Western European *Drosophila melanogaster* natural populations. Sci Rep 8: 16143.

Mateo, L., A. Ullastres, and J. González, 2014 A transposable element insertion confers
1058   xenobiotic resistance in *Drosophila*. PLoS Genet 10: e1004560.

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis *et al.*, 2010 The Genome
1060   Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA
               sequencing data. Genome Res 20: 1297–1303.

1062 Novembre, J., and M. Stephens, 2008 Interpreting principal component analyses of spatial
               population genetic variation. Nat Genet 40: 646–649.

1064 Ometto, L., 2010 Inferring the Effects of Demography and Selection on *Drosophila
               melanogaster* Populations from a Chromosome-Wide Scan of DNA Variation. Mol
1066   Biol Evol 22: 2119–2130.

Orozco-terWengel, P., M. Kapun, V. Nolte, R. Kofler, T. Flatt *et al.*, 2012 Adaptation of
1068   *Drosophila* to a novel laboratory environment reveals temporally heterogeneous
               trajectories of selected alleles. Mol Ecol 21: 4931–4941.

1070   Paaby, A. B., A. O. Bergland, E. L. Behrman, and P. S. Schmidt, 2014 A highly pleiotropic
        amino acid polymorphism in the *Drosophila* insulin receptor contributes to life-history
1072   adaptation. Evolution 68: 3395–3409.

        Paaby, A. B., M. J. Blacket, A. A. Hoffmann, and P. S. Schmidt, 2010 Identification of a
1074   candidate adaptive polymorphism for *Drosophila* life history by parallel independent
        clines on two continents. Mol Ecol 19: 760–774.

1076   Pool, J. E., C.-D. B Russell, R. P. Sugino, K. A. Stevens, C. M. Cardeno *et al.*, 2012
        Population Genomics of Sub-Saharan *Drosophila melanogaster*: African Diversity
1078   and Non-African Admixture. PLoS Genet 8: e1003080.

        Raineri, E., L. Ferretti, E.-C. Anna, B. Nevado, S. Heath *et al.*, 2012 SNP calling by
1080   sequencing pooled samples. BMC Bioinformatics 13: 1–8.

        Ramaekers, A., A. Claeys, M. Kapun, E. Mouchel-Vielh, D. Potier *et al.*, 2019 Altering the
1082   Temporal Regulation of One Transcription Factor Drives Evolutionary Trade-Offs
        between Head Sensory Organs. Dev Cell 50: 780-792.

1084   Reinhardt, J., B. Kolaczkowski, C. Jones, D. Begun, and A. Kern, 2014 Parallel Geographic
        Variation in *Drosophila melanogaster*. Genetics 197: 361–373.

1086   Rodríguez-Trelles, F., R. Tarrío, and M. Santos, 2013 Genome-wide evolutionary response
        to a heat wave in *Drosophila*. Biol Lett 9: 20130228.

1088   Rudman, S. M., S. Greenblum, R. C. Hughes, S. Rajpurohit, O. Kiratli *et al.*, 2019
        Microbiome composition shapes rapid genomic adaptation of *Drosophila*
1090   *melanogaster*. Proc Natl Acad Sci USA 116: 20025-20032.

        dos Santos, G., A. J. Schroeder, J. L. Goodman, V. B. Strelets, M. A. Crosby *et al.*, 2015
1092   FlyBase: introduction of the *Drosophila melanogaster* Release 6 reference genome
        assembly and large-scale migration of genome annotations. Nucleic Acids Res 43:
1094   D690–D697.

        Schlötterer, C., R. Tobler, R. Kofler, and V. Nolte, 2014 Sequencing pools of individuals -
1096   mining genome-wide polymorphism data without big funding. Nat Rev Genet 15:
        749–763.

1098   Schneider, D., 2000 Using *Drosophila* as a model insect. Nat Rev Genet 1: 218–226.

        Sprengelmeyer, Q. D., S. Mansourian, J. D. Lange, D. R. Matute, B. S. Cooper *et al.*, 2020
1100   Recurrent Collection of *Drosophila melanogaster* from Wild African Environments
        and Genomic Insights into Species History. Mol Biol Evol 37: 627–638.

1102   Stajich, J. E., and M. W. Hahn, 2005 Disentangling the Effects of Demography and Selection
        in Human History. Mol Biol Evol 22: 63–73.

1104   Stephan, W., 2016 Signatures of positive selection: from selective sweeps at individual loci
        to subtle allele frequency changes in polygenic adaptation. Mol Ecol 25: 79–88.

1106 Turner, T. L., M. T. Levine, M. L. Eckert, and D. J. Begun, 2008 Genomic Analysis of
Adaptive Differentiation in *Drosophila melanogaster*. Genetics 179: 455–473.

1108 Turner, T. L., A. D. Stewart, A. T. Fields, W. R. Rice, and A. M. Tarone, 2011 Population-
Based Resequencing of Experimentally Evolved Populations Reveals the Genetic

1110 Basis of Body Size Variation in *Drosophila melanogaster*. PLoS Genet 7: e1001336.

Umina, P. A., 2005 A Rapid Shift in a Classic Clinal Pattern in *Drosophila* Reflecting Climate

1112 Change. Science 308: 691–693.

Waldvogel, A.-M., B. Feldmeyer, G. Rolshausen, M. Exposito-Alonso, C. Rellstab *et al.*,

1114 2020 Evolutionary genomics can improve prediction of species' responses to climate
change. Evol Lett 4: 4–18.

1116 Wallace, M. A., K. A. Coffman, C. Gilbert, S. Ravindran, G. F. Albery *et al.*, 2020 The
discovery, distribution and diversity of DNA viruses associated with *Drosophila*

1118 *melanogaster* in Europe. Virus Evolution, in revision (preprint: bioRxiv
2020.10.16.342956).

1120 Wittmann, M. J., A. O. Bergland, M. W. Feldman, P. S. Schmidt, and D. A. Petrov
Seasonally fluctuating selection can maintain polymorphism at many loci via

1122 segregation lift. Proc Natl Acad Sci USA 114: E9932–E9941.

Wright, S., 1943 Isolation by distance. Genetics 28: 114.

1124 Zheng, X., S. M. Gogarten, M. Lawrence, A. Stilp, M. P. Conomos *et al.*, 2017 SeqArray - a
storage-efficient high-performance data format for WGS variant calls. Bioinformatics

1126 33: 2251–2257.

Zhu, Y., A. O. Bergland, J. González, and D. A. Petrov, 2012 Empirical Validation of Pooled

1128 Whole Genome Population Re-Sequencing in *Drosophila melanogaster*. PLoS ONE
7: e41901.

1130