Cubam receptor-mediated endocytosis in hindgut-derived pseudoplacenta of a viviparous teleost *Xenotoca eiseni*

Atsuo Iida[1, *], Kaori Sano[2], Mayu Inokuchi[3], Takayuki Suzuki[1], Mao Kuriki[4], Maina Sogabe[4], Daichi Susaki[5], Kaoru Tonosaki[5], Tetsu Kinoshita[5], Eiichi Hondo[1]

1. Department of Animal Sciences, Graduate School of Bioagricultural Sciences, Nagoya University, Tokai National Higher Education and Research System, Nagoya, Aichi, Japan.

2. Department of Chemistry, Faculty of Science, Josai University, Sakado, Saitama, Japan.

3. Department of Life Sciences, Toyo University, Itakura, Gunma, Japan.

4. Department of Regeneration Science and Engineering, Institute for Frontier Life and Medical Sciences, Kyoto University, Kyoto, Kyoto, Japan.

5. Kihara Institute for Biological Research, Yokohama City University, Yokohama, Kanagawa, Japan.

Correspondence: Atsuo Iida

E-mail: tol2.4682@gmail.com

Author contribution: A.I. designed the study. A.I., K.S., and M.I. carried out the experiments. T.S., M.K., and M.S. contributed to fluorescent microscopy. D.S., K.T., T.K., and E.H. contributed to next-generation

20   sequencing and data analysis. A.I. wrote the manuscript.

22   **Competing interest statement**: The authors declare that they have no competing interests.

24   **Keywords**: Goodeidae, viviparity, trophotaenia, endocytosis, proteolysis

26   **This PDF file includes**:

27   Main Text

28   Figures 1 to 5

Abstract

Nutrient transfer from mother to the embryo is essential for reproduction in viviparous animals. Here, we

focused on the molecular mechanism of nutrient absorption in hindgut-derived pseudoplacenta

(trophotaenia) of the viviparous teleost *Xenotoca eiseni*. The candidate genes involved in the process

were identified by RNA-Seq, and then protein distribution and functions were investigated using molecular

biological methods. Our results suggested that Cubam (<u>Cub</u>ilin-<u>Am</u>nionless) receptor-mediated

endocytosis and cathepsin L-dependent proteolysis are involved in the maternal macromolecule

absorption via the trophotaenia. Such a nutrient absorption mechanism involving endocytosis is not a

specific trait to viviparous fish. Similar processes have been reported in the larval stage of oviparous fish

or the suckling stage of viviparous mammals. Our findings suggest that the viviparous teleost acquired

trophotaenia-based viviparity via a modification of the intestinal absorption system common in vertebrates.

This is a fundamental study to understand the strategic variation in the reproductive system of vertebrates.

Introduction

August Krogh wrote, "*For such a large number of problems there will be some animal of choice or a few such animals on which it can be most conveniently studied*" [1]. This study aimed to investigate the molecular mechanism of maternal nutrient absorption in a species-specific pseudoplacenta of a viviparous teleost species belonging to the family Goodeidae.

Viviparity is a reproduction system, whereby the oocyte is fertilized in the female body, and subsequent embryo growth occurs with maternal component supply. Each viviparous animal has acquired processes specialized to the gestation in both the mother and embryo [2]. The placenta and umbilical cords in viviparous mammals are major components of the process for mother-to-embryo material transport [3,4]. Other viviparous components such as the extended yolk sac or pseudoplacenta are known in several viviparous vertebrates, except mammals [5].

The family Goodeidae is a freshwater small teleost distributed in Mexico, which includes approximately 40 viviparous species [6]. They possess trophotaenia, which is a hindgut-derived pseudoplacenta that absorbs the maternal component [7,8]. Trophotaenia is a ribbon-like structure consisting of a single epithelial layer, internal blood vessels, and connective tissues [9,10]. The epithelial cell is like an enterocyte in the intestine. Electron microscopy indicated that microvilli form in the apical side of the cell and intracellular vesicles in the cytoplasm [11]. Since the 1980s, these structures have been believed to be involved in maternal component absorption [12]. The nature of the maternal component was predicted to be proteins or other macromolecules secreted into the serum or ovarian fluids; however, no one has experimentally determined its distinct component [13,14]. Recently, we demonstrated that a yolk protein vitellogenin is secreted into the ovarian lumen of pregnant females, and the intraovarian embryo absorbs the nutrient protein via the trophotaenia using a goodeid species *Xenotoca eiseni* [15]. In that study, enterocyte-like microvilli and intracellular vesicles were also observed in the epithelial cells of the

trophotaenia. We hypothesized that the epithelial layer cell in the trophotaenia absorbs the maternal

protein and/or other components as macromolecules because the ovarian lumen lacks proteolysis activity

like the digestive intestine. However, the molecules responsible for the trophotaenia-mediated

macromolecule absorption have not been reported.

Vacuolated enterocytes involved in macromolecule absorption have also been reported in other

vertebrate species, including suckling mammals and stomachless fish [16-18]. Park et al. [19] reported

that the scavenger receptor complex Cubam (Cubilin-Amnionless) and Dab2 are required for

macromolecule uptake in zebrafish juveniles. Furthermore, the conditional knockout of *Dab2* in mice led to

stunted growth and severe protein malnutrition at the suckling stage [19]. Based on this report, we discuss

the commonality of molecular process for the macromolecule absorption between the intestinal

enterocytes and the trophotaenia [15]. However, the molecules responsible for macromolecule absorption

in the trophotaenia are still elusive.

Here, we report candidate molecules for nutrient uptake and subsequent proteolysis in the trophotaenia

of *X. eiseni*. An RNA-Seq for trophoteniae indicated candidate receptor molecules and endocytosis-

associated proteases expressed in the trophotaenia. Immunohistochemistry and biochemical assays

suggested the presence and functions of the candidate factors in the trophotaenia.

Results

Gene expression in trophotaenia

Our previous findings and the hypothesis based on that are described in Figure 1A. In a viviparous

teleost, *X. eiseni*, the embryo is raised in the ovary while receiving maternal nutrients. The trophotaenia is

a pseudoplacenta that plays a role in the absorption of maternal nutrients consisting of proteins and other

supplements. Based on previous studies, we hypothesized that several maternal proteins are absorbed by

endocytosis-mediated proteolysis in the epithelial cells of the trophotaenia. To verify this hypothesis, we

explored candidate genes for receptor, adaptor, vesicle, and protease proteins that are highly expressed

in the trophotaenia. RNA-Seq analyses were performed using total RNA extracted from the trophotaeniae

of the 3rd- or 4th-week embryos (Figure 1B). We selected candidate genes and compared their predicted

expression level as transcription per million (TPM) values that were calculated using the known transcript

sequence of *P. reticulata* as reference (Dataset S1). RNA-Seq suggested that *cubilin* (*cubn*) and

amnionless (amn) were highly expressed in the trophotaeniae; however, other co-receptor genes (*lrp1aa*,

lrp2a) were considerably lower (Figure 1C, Table S1). Adaptor protein-2 (AP2) subunit genes (*ap2a1*,

ap2b1, *ap2m1a*, and *ap2s1*) were more highly expressed than the other family adaptor genes (*ldlrap1b*

and *numb*) (Figure 1C, Table S2). Two families of the vesicle coating protein genes (*clta*, *cltbb*, *cltc*; *flot1b*,

flot2a) were expressed higher than the vesicle proteins classified in other families (*cav2* and *cav3*) (Figure

1C, Table S3). The lysosomal endopeptidase enzyme gene (*ctsl.1*) exhibited a high TPM value (> 10,000)

rather than that of not only protease family genes but also most of the genes expressed in the

trophotaeniae (Figure 1C, Table S4). The trend of the predicted TPM values in each gene was not

noticeably different between the 3rd- and 4th- week samples.

Sequences for Cubam receptor genes

A Cubam receptor is known to be a membrane-bound multi-ligand receptor consisting of Cubn and Amn

(Figure 2A). The secreted protein Cubn specifically binds to the transmembrane protein Amn; thus, the

CUB domain that associates with the ligands are localized on the apical surface of the plasma membrane.

We obtained the amino acid sequences of *X. eiseni* Cubn and Amn by *de novo* assembly of NGS reads

from the trophotaeniae. The sequences for Cubn and Amn were compared between four vertebrate

species, *Homo sapiens*, *Danio rerio*, *P. reticulata*, and *X. eiseni.* The binding motifs were conserved in the

species (Figure 2B and C). The intracellular domain of Amn included two conserved motifs to bind the

adaptor proteins, including AP2. The motifs were also conserved in the species (Figure 2D).

Distribution of Cubilin and Amnionnless in trophotaenia

To validate the expression of *cubn* and *amn* in the trophotaenia, semi-quantitative RT-PCR analyses

were performed using total RNAs extracted from the whole-embryo including the trophotaeniae, isolated

trophotaeniae, and adult skeletal muscle. The muscle sample was used as a control for tissue with low

endocytosis activity. The gene expression patterns did not conflict between the results of RT-PCR and

RNA-Seq. In the embryo and trophotaenia, both *cubn* and *amn* were more highly expressed than *lrp2a*

(Figure 3A). Conversely, no or few expressions of the target genes except *actb* as a positive control were

detected in the adult muscle. Next, to detect protein localization, immunohistochemistry was performed

using antibodies against Cubn or Amn. In both proteins, strong signals were observed in the epithelial

monolayer of the trophotaenia, while few background noises were detected in the control assay (Figure

3B-D, see also Figure S1 and S2). Confocal microscopy revealed the cellular distribution of the anti-Cubn

signals. The signals in the apical surface of the epithelial cell were detected as a homogeneous

distribution, while almost all signals in the cytoplasm were captured as a dot pattern (Figure 3E, Figure

S3). Immunoelectron microscopy revealed that the microvilli were distributed on the apical surface of the

trophotaenia epithelium, and intracellular vesicles were observed in the cytoplasm; furthermore, anti-Cubn

signals were distributed in the intracellular vesicles and overlapped with the microvilli on the apical surface

(Figure 3F).

Proteolysis activity in trophotaenia

Cathepsin L is a lysosomal cysteine proteinase characterized by three conserved protease regions and

active sites consisting of cysteine, histidine, or asparagine (Figure 4A). The functional regions were

conserved in the *X. eiseni* Cathepsin L protein translated from the coding sequence of *ctsl.1*. RT-PCR

analysis revealed a strong expression of *ctsl.1* in the trophotaenia (Figure 4B). To identify which type of

cells include proteolysis activity in the trophotaenia, acidic organelles including lysosomes and

endosomes were detected using a fluorescent probe. The LysoTracker® indicated the presence of acidic

organelles in the epithelial layer cells (Figure 4C). The signals were distributed in the cytoplasm and were

not components in the nuclei (Figure 4D). According to the RNA-Seq analysis, *ctsl.1* was presumed to be

the highest expressed *cathepsin* gene in the trophotaenia; thus, we calibrated the proteolysis activity of

cathepsin L in the trophotaenia using a fluorescent substrate-based measurement system. The

fluorescence indicating substrate digestion was significantly higher in the trophotaenia lysate than in the

control at one h after the reagent mixture. Furthermore, the intensity of the lysate condition continued to

increase for 7 h (Figure 4E, Table S5). Conversely, the increase in intensity was strongly suppressed by a

cathepsin L inhibitor.

Adaptors and vesicle coating proteins

The expression of candidate genes for adaptors (*ap2a1*, *ap2b1*, *ap2m1a*, *ap2s1*, *ldlrap1b*, and *numb*)

and vesicle coating proteins (*clta* and *cltc*) were confirmed by RT-PCR (Figure S4). We determined an

incomplete transcript for *X. eiseni* Dab2 from the *de novo* assembly; however, it lacked internal sequences

in comparison with the proteins in other vertebrates (Figure S5). Furthermore, no amplicons were

obtained by RT-PCR using primers designed based on the Dab2-like sequence.

Discussion

RNA-Seq analyses revealed high expression of the genes for receptor-mediated endocytosis in the

trophotaenia of *X. eiseni* embryos. Cubn and Amn form a Cubam receptor complex that associates with

Vitamin-B12, albumin, transferrin, and other ligands [20,21]. The predicted amino acid sequences for *X.*

eiseni Cubn and Amn obtained from the *de novo* assembly both possessed a conserved motif which

allowed binding to each other, and Amn retained adaptor binding sites in the intracellular region [21, 22].

Immunohistochemical analysis indicated the presence of Cubn not only in the apical surface of the

epithelial layer but also in the intracellular vesicles in the cells, suggesting the incorporation and recycling

of endocytic receptors [23,24]. This evidence supports the idea that Cubam plays a role as a receptor for

the intraovarian nutrients and is involved in endocytosis. We also indicated the low presence of Lrp2 (also

known Megalin), which is a major co-receptor for Cubam [25]. This indicated that Cubam works

independently or in cooperation with other co-receptors, except Lrp2, in the trophotaenia. In zebrafish, a

previous study reported that Cubam-dependent endocytosis in the lysosome rich enterocytes (LREs) does

not require the presence of Lrp2 [19]. This supports our hypothesis; however, we do not exclude the

possibility that our sequencing and alignment processes could not catch other *lrp2* orthologous genes in

X. eiseni. Park et al. [19] also reported that Dab2 is an essential adaptor molecule not only in the zebrafish

larval intestine but also in the endocytic nutrient absorption in the ileum of suckling mice. However, we

obtained only an incomplete sequence for the *X. eiseni* Dab2-like protein without an internal region

compared with mammalian orthologs. This may be an alternative splice form with less endocytic activity

[26,27]. Furthermore, we did not detect Dab2-like expression by RT-PCR, and we could not exclude the

possibility that the predicted Dab2-like sequence is caused by an assembly error. Thus, the contribution of

Dab2 to endocytosis in the trophotaenia has not been confirmed in this study. The other adaptor or vesicle

coating proteins we detected were typical co-factors for receptor-mediated endocytosis. Conversely,

Caveolin is a vesicle coating protein involved in receptor-independent endocytosis [28]. Thus, a low

expression of *caveolin* genes (*cav2* and *cav3*) does not conflict with the activation of Cubam-mediated

endocytosis. Additionally, biochemical assays supported that cathepsin L is an active protease that

functions in the intracellular vesicles that is configured following membrane budding. This evidence

supports the idea that Cubam-mediated endocytosis and cathepsin L-dependent proteolysis is one of the

key mechanisms for the absorption of the maternal component in *X. eiseni* embryos.

Cubam is also known to be a scavenger receptor involved in nonspecific protein uptake [20]. In this

study, we did not determine the Cubam ligands in the trophotaenia. However, the candidates are limited to

intraovarian secreted proteins. In several goodeid species, previous studies indicated that the ovarian

fluids include various proteins in a pattern similar to that in the blood serum [13,29]. Therefore, several

serum proteins, albumen, transferrin, and others known as Cubam ligands, are potential targets in the

case of intraovarian nutrient absorption via the trophotaenia. Another possibility is that vitellogenin is also

a potential target because it is secreted into the ovarian fluids and is absorbed into the trophotaenia via

intracellular vesicles [15]. Another study suggested that the Cubn-Amn-Lrp2 receptor complex is related to

transport of yolk proteins, including vitellogenin, in endodermal epithelial cells during chicken yolk sac

growth [30]. These reports support the idea that Cubn and Amn are potent candidate molecules for a

receptor complex involved in maternal component uptake in the trophotaenia. However, functional

analyses such as gene knockout or transgenic technologies used in conventional model animals, such as

laboratory mice or small oviparous fish, have not been applied to goodeid fish. For a direct validation of

our hypothesis, these reverse genetic methods should also be developed in the viviparous teleost.

As described above, endocytosis-mediated nutrient absorption is not limited to viviparous fish; it has also

been reported in other animal species (Figure 5). In mammals, macromolecule absorption in the intestine

is limited to the suckling period [31]. In the stomachless fish, endocytosis-mediated absorption is

considered to continue in part of the intestine for life, because of low digestive activity in the enteric cavity

[18]. In most invertebrate species, because the extracellular digestive system is primitive, food particles

are absorbed into the intestinal cells by vesicle trafficking and degraded by intracellular digestion [32, 33].

Thus, we hypothesize that endocytic nutrient absorption and intracellular digestion are ancestral traits

common not only in vertebrates but also in invertebrates, and their importance has decreased in certain

vertebrates with development of the extracellular digestive system in the enteric cavity. The ancestor of

the goodeid species may have applied the endocytic process to the reproductive system, and then

configured the unique hind-gut derived pseudoplacenta for matrotrophic nutrition during gestation [7,34].

To validate our hypothesis, further exhaustive omics analyses using the goodeid species and comparative

research using the subfamily Empetrichthyinae, which is the oviparous species most closely related to the

viviparous goodeid fish, are required [6,35].

This study is an investigation of species-specific traits based on the transcriptome of a non-conventional

model species. The results revealed potential candidate molecules for nutrient absorption in the

pseudoplacenta of the viviparous teleost. As August Krogh wrote, this kind of study would be suitable for

investigation using the most appropriate species and is unsuitable for verification using alternative models

such as viviparous rodents or oviparous teleosts. We believe that this study is an important and

fundamental step in understanding the strategic variation of the reproductive system in vertebrates.

Methods

Animal experiments

This study was approved by the ethics review board for animal experiments at Nagoya University

(Approval number: AGR2020028). We sacrificed live animals in minimal numbers under anesthesia

according to the institutional guidelines.

Fish breeding

X. eiseni was purchased from Meito Suien Co., Ltd. (Nagoya, Japan). Adult fish were maintained in

freshwater at 27 °C under a 14:10-h light: dark photoperiod cycle. Fish were bred in a mass-mating

design, and approximately 30 adult fish were maintained for this study. The juveniles were fed live brine

shrimp larvae, and the adults were fed Hikari Crest Micro Pellets and ultraviolet-sterilized frozen

chironomid larvae (Kyorin Co., Ltd., Himeji, Japan). To accurately track the pregnancy period, the

laboratory-born fish were crossed in a pair-mating design, and the mating behavior was recorded.

Sample collection

Fish samples were anesthetized using tricaine on ice before the surgical extraction of tissues or

embryos. The obtained samples were stored on ice until subsequent manipulations. In this study, we

dissected approximately 10 pregnant females and extracted 15–30 embryos in each operation.

 RNA-Seq

Total RNA was extracted from trophotaenae of the 3rd or 4th week of the embryo extracted from the

pregnant female using the RNeasy Plus Mini kit (QIAGEN). A total of six samples were obtained from

three 3rd week embryos and three 4th week embryos. Next-generation sequencing (NGS) was outsourced

to Macrogen Japan Corp. (Kyoto, Japan) using NovaSeq6000 (Illumina, Inc., San Diego, CA, USA).

Approximately 60 million 150-bp paired-end reads were obtained in each sample. The NGS data was

deposited to the DNA Data Bank of Japan (DDBJ, ID: DRA011209). *De novo* assembly and mapping to

the reference sequence were performed by CLC Genomics Workbench .(Filgen, Inc., Nagoya, Japan)

The published transcript sequences of*Poecilia reticulata* (NCBI Genome, ID: 23338) was used as a

reference. The transcript sequences of *X. eiseni* were deposited into the DDBJ. The accession numbers

were listed in Table S6.

Reverse transcription (RT) PCR

Total RNA was extracted from tissues or whole embryos using the RNeasy Plus Mini kit and reverse-

transcribed using SuperScript IV reverse transcriptase (Thermo Fisher Scientific). PCR was carried out

using KOD-FX-Neo (Toyobo, Osaka, Japan) under the following conditions: 100 s at 94 °C, followed by 32

cycles of 20 s at 94 °C, 20 s at 60 °C, 60 s at 72 °C, and 120 s at 72 °C. Primer sequences are listed in

the Resource List in Table S6.

Antibodies and antiserums

258  Antiserums against Cubn and Amn were generated in this study. The antigen sequences were 6× His-

259 tagged peptide of 151 amino acids (aas) corresponding to the intermediate region of *X. eiseni* Cubn

260 (Accession#, LC595284; aa 691–841) and 6× His-tagged peptide of 247 aas corresponding to the C-

261 terminal of *X. eiseni* Amn (Accession#, LC595285). The experimental procedure has been described in

262 our previous study [15]. All antibodies and antiserums used in this study are listed in Table S7.

264 *Immunohistochemistry*

265  Tissue samples were fixed in 4.0% paraformaldehyde/phosphate-buffered saline (PFA/PBS) at 4 °C

266 overnight. Samples were permeabilized using 0.5% TritonX-100/PBS at room temperature for 30 min.

267 Endogenous peroxidase was inactivated by 3.0 % hydrogen peroxide/PBS for 10 min. Then, the sample

268 was treated with Blocking-One solution (Nacalai Tesque, Kyoto, Japan) at room temperature for 1 h.

269 Primary antibody or antiserums were used at 1:500 dilution with Blocking-One solution. Samples were

270 incubated with primary antibody or antiserum at 4 °C overnight. Secondary antibodies were used at a

271 1:500 dilution in 0.1% Tween-20/PBS. Samples were treated with the secondary antibody solution at 4 °C

272 for 2 h. We performed a 3,3'-Diaminobenzidine Tetrahydrochloride (DAB) color development using the

273 DAB Peroxidase Substrate Kit, ImmPACT (Vector Laboratories, Inc., Burlingame, CA, USA). Microscopic

274 observation was performed using an Olympus BX53 microscope and photographed using a DP25 digital

275 camera (Olympus, Shinjuku, Japan).

277 *Fluorescent Immunohistochemistry*

278  Tissue samples were fixed in 4.0% PFA/PBS at 4 °C overnight. Samples were permeabilized using 0.5%

TritonX-100/PBS at room temperature for 30 min. Endogenous peroxidase was inactivated by 3.0 %

hydrogen peroxide/PBS for 10 min. Then, the sample was treated with Blocking-One solution (Nacalai

Tesque, Kyoto, Japan) at room temperature for 1 h. Primary antibody (anti-Cubn) was used at a 1:500

dilution with Blocking-One solution. Samples were incubated with primary antibody at 4 °C overnight.

Secondary antibody was used at a 1:500 dilution in 0.1% Tween-20/PBS with 4',6-diamidino-2-

phenylindole (DAPI). Samples were treated with the secondary antibody solution at 4 °C overnight.

Microscopic observation was performed using a Leica TCS SP8 microscope (Leica Microsystems,

Wetzlar, Germany).

Immunoelectron microscopy

Embryo samples were fixed in 4.0% PFA/PBS. Fixed samples were washed in PBS, then reacted with a

primary antibody (anti-Cubn) at 4 °C overnight, and then reacted with biotinylated anti-rabbit IgG (Vector,

Burlingame, CA, USA) at room temperature for 2 h. Samples were performed with the avidin–biotin–

peroxidase complex kit (Vector), and visualized with 0.05% DAB (Dojindo Laboratories, Kumamoto,

Japan) and 0.01% hydrogen peroxide in 50 mM Tris buffer (pH 7.2) at room temperature for 10 min. The

procedure for electron microscopy is described in our previous study [15].

Labeling acidic organelles

The live embryos immediately after extraction from the pregnant female at the 4th-week post-mating were

incubated in PBS with a 1:1000 dilution of LysoTracker® Red (Thermo Fisher Scientific) for 1 h at room

temperature. The samples were fixed with 4.0 % PFA/PBS and stained with DAPI. Microscopic

observation was performed using a Leica DM5000 B microscope (Leica Microsystems, Wetzlar,

Germany).

Measurement of Cathepsin L activity

The trophotaenia lysate was prepared from six littermate embryos obtained from the pregnant females at

the 4th-week post-mating. The trophotaeniae were manually extracted from the embryo under a

microscope. Proteolysis detection was performed using the Cathepsin L Activity Fluorometric Assay Kit

(BioVision, Inc., Milpitas, CA, USA). The fluorescence intensities were measured using a Qubit 4

Fluorometer (Thermo Fisher Scientific).

Acknowledgments

This work was supported by research grants from the Nakatsuji Foresight Foundation and the Daiko

Foundation.

Data accessibility

The data that support the findings have been provided with the manuscript.

References

1. A. Krogh, The progress of physiology. *Am. J. Physiol.* **90**, 243-251 (1929).

2. D. Blackburn, "Viviparity and oviparity: Evolution and reproductive strategies" in: Encyclopedia of Reproduction, Vol. 4, E. Knobil, J. D. Neill, Eds. (Academic Press, 1999), pp. 994-1003.

3. G. J. Burton, A. L. Fowden, Review: The placenta and developmental programming: balancing fetal nutrient demands with maternal resource allocation. *Placenta*. **33**, S23-S27 (2012).

4. R. W. Gill, G. Kossoff, P. S. Warren, W. J. Garrett, Umbilical venous flow in normal and complicated pregnancy. *Ultrasound Med. Biol*. **10**, 349-363 (1984).

5. R. M. Roberts, J. A. Green, L. C. Schulz, The evolution of the placenta. Reproduction (Cambridge, England), **152**, R179–R189 (2016).

6. K. L. Foster, K. R. Piller, Disentangling the drivers of diversification in an imperiled group of freshwater fishes (Cyprinodontiformes: Goodeidae). *BMC Evol. Biol.* **18**, 116 (2018).

7. C. L. Turner, Pericardial sac, trophotaeniae, and alimentary tract in embryos of goodeid fishes. *J. Morphol.* **67**, 271-289 (1940).

8. S. M. Tinguely, A. Lange, C. R. Tyler, Ontogeny and Dynamics of the Gonadal Development, Embryogenesis, and Gestation in *Xenotoca eiseni* (Cyprinodontiformes, Goodeidae). *Sex Dev*. **13**, 297-310 (2019).

9. C. L. Turner, Viviparity Superimposed upon Ovo-Viviparity in the Goodeidae, a Family of

Cyprinodont Teleost Fishes of the Mexican Plateau. *J. Morphol.* **55**, 207- 251 (1933).

10. C. L. Turner, The trophotaeniae of the goodeidae, a family of viviparous cyprinodont fishes. *J.*

Morphol. **61**, 495- 523 (1937).

11. J. Lombardi, J. P. Wourms, The trophotaenial placenta of a viviparous goodeid fish. I.

Ultrastructure of the internal ovarian epithelium, the maternal component. *J. Morphol*. **184**, 277-

292 (1985).

12. F. Hollenberg, J. P. Wourms, Ultrastructure and protein uptake of the embryonic trophotaeniae of

four species of goodeid fishes (Teleostei: Atheriniformes). *J. Morphol*. **219**, 105-129 (1994).

13. F. Hollenberg, J. P. Wourms, Embryonic growth and maternal nutrient sources in goodeid fishes

(Teleostei: Cyprinodontiformes). *J. Exp. Zool.* **271**, 379–394 (1995).

14. A. Vega-López, E. Ortiz-Ordóñez, E. Uría-Galicia, E. L. Mendoza-Santana, R. Hernández-

Cornejo*, et al.,* The role of vitellogenin during gestation of *Girardinichthys viviparus* and *Ameca*

splendens; two goodeid fish with matrotrophic viviparity. *Comp. Biochem. Physiol. Part A Mol.*

Integr. Physiol. **147**, 731–742 (2007).

15. A. Iida, H. N. Arai, Y. Someya, M. Inokuchi, T. A. Onuma, *et al*., Mother-to-embryo vitellogenin

transport in a viviparous teleost *Xenotoca eiseni*. *PNAS*, **116**, 22359-22365 (2019).

16. J. P. Kraehenbuhl, C. Bron, C., B. Sordat, Transfer of humoral secretory and cellular immunity from mother to offspring. *Curr. Top. Pathol.* **66**, 105–157 (1979).

17. P. C. Moxey, J. S. Trier, Development of villus absorptive cells in the human fetal small intestine: a morphological and morphometric study. *Anat. Rec.* **195**, 463–482 (1979).

18. J. H. Rombout, C. H. Lamers, M. H. Helfrich, A. Dekker, J. J. Taverne-Thiele, Uptake and transport of intact macromolecules in the intestinal epithelium of carp (*Cyprinus carpio L.*) and the possible immunological implications. *Cell Tissue Res.* **239**, 519-530 (1985).

19. J. Park, D. S. Levic, K. D. Sumigray, J. Bagwell, O. Eroglu, *et al*., Lysosome-Rich Enterocytes Mediate Protein Absorption in the Vertebrate Gut. *Dev. Cell.* **51**, 7-20 (2019).

20. P. J. Verroust, H. Birn, R. Nielsen, R. Kozyraki, E. I. Christensen, The tandem endocytic receptors megalin and cubilin are important proteins in renal pathology. *Kidney Int.* 62, 745-756 (2002).

21. G. A. Pedersen, S. Chakraborty, A. L. Steinhauser, L. M. Traub, M. Madsen, AMN directs endocytosis of the intrinsic factor-vitamin B(12) receptor cubam by engaging ARH or Dab2. *Traffic.* **11**, 706-720 (2010).

22. C. Larsen, A. Etzerodt, M. Madsen, K. Skjødt, S. K. Moestrup, C. B. F. Andersen, Structural assembly of the megadalton-sized receptor for intestinal vitamin B12 uptake and kidney protein reabsorption. *Nat. Commun.* **9**, 5204 (2018).

23. E. I. Christensen, H. Birn, P. Verroust, S. K. Moestrup, Membrane receptors for endocytosis in the

    renal proximal tubule. *Int. Rev. Cytol.* **180**, 237-284 (1998).

24. B. D. Grant, J. G. Donaldson, Pathways and mechanisms of endocytic recycling. *Nat. Rev. Mol.*

    Cell Biol. **10**, 597-608 (2009).

25. E. I. Christensen, H. Birn, Megalin and cubilin: multifunctional endocytic receptors. *Nat. Rev. Mol.*

    Cell Biol. **4**, 256-266 (2002).

26. M. E. Maurer, J. A. Cooper, Endocytosis of megalin by visceral endoderm cells requires the Dab2

    adaptor protein. *J. Cell Sci*. **118**, 5345-55 (2005).

27. C. V. Finkielstein, D. G. Capelluto, Disabled-2: A modular scaffold protein with multifaceted

    functions in signaling. *Bioessays*. **38**, S45-55 (2016).

28. T. M. Williams, M. P. Lisanti, The caveolin proteins. *Genome Biol*. **5**, 214 (2004).

29. J. F. Schindler, Structure and function of placental exchange surfaces in goodeid fishes (Teleostei:

    Atheriniformes). *J. Morphol*. **276**, 991-1003 (2015).

30. R. Bauer, J. A. Plieschnig, T. Finkes, B. Riegler, M. Hermann, W. J. Schneider, The developing

    chicken yolk sac acquires nutrient transport competence by an orchestrated differentiation

    process of its endodermal epithelial cells. *J. Biol. Chem.* **288**, 1088-1098 (2013).

31. V. Muncan, J. Heijmans, S. D. Krasinski, N. V. Büller, M. E. Wildenberg, *et al.*, Blimp1 regulates

the transition of neonatal to adult intestinal epithelium. *Nat. Commun.* **2**, 452 (2011).

32. P. V. Fankboner, Digestive System of Invertebrates. In book: eLS (2001).

33. V. Hartenstein, P. Martinez, Phagocytosis in cellular defense and nutrition: a food-centered

approach to the evolution of macrophages. *Cell Tissue Res.* **377**, 527–547 (2019).

34. M. C. Uribe, H. J. Grier, S. A. Avila-Zúñiga, A. García-Alarcón, Change of lecithotrophic to

matrotrophic nutrition during gestation in the viviparous teleost *Xenotoca eiseni* (Goodeidae). *J.*

Morphol. **279**, 1336-1345 (2018).

35. R. Van Der Laan, W. N. Eschmeyer, R. Fricke, Family-group names of Recent fishes. *Zootaxa.*

3882, 1-230 (2014).

Figure 1. Exploring candidate genes for nutrient absorption

A. A working model and hypothesis of this study. In the goodeid viviparous fish (*X. eiseni*), intraovarian

embryo absorbs maternal nutrients via the trophotaenia. We hypothesized that endocytosis-mediated

proteolysis is related to nutrient absorption. Based on the scenario, potential target genes are for

endocytic receptors, adaptors, vesicle coating proteins, and proteases. **B**. An experimental scheme for the

RNA-Seq analysis. RNA samples were obtained from the trophotaeniae (white dotted line) of the single

intraovarian embryos extracted from the pregnant females of the 3rd- or 4th-week post-mating. The RNA-

Seq was performed using three samples at every stage. **C**. The graphs indicate the TPM values of the

genes selected from the RNA-Seq data that are involved in the endocytosis-mediated proteolysis pathway.

Figure 2. Structures and amino acid sequences for *X. eiseni* Cubn and Amn

A. The illustration indicates that a typical structure of Cubam receptor complex consists of Cubn and Amn.

Both proteins possess a conserved motif to bind each other in the N-terminal regions, and the Amn

possesses two adaptor binding motifs in the C-terminal intracellular domain. **B-D**. A comparison of amino

acid sequence around the Amn-binding motif in Cubn (**B**), Cubn-binding motif in Amn (**C**), and adaptor

binding motifs in Amn (**D**) between *X. eiseni* and three other vertebrate species. The gray filled texts are

conserved sequences among the four species.

Figure 3. Distribution of receptors involved in endocytosis

A. RT-PCR for the candidate genes for the receptors involved in the endocytosis. All amplicons were

detected as single band on the expected sizes based on the transcript sequences obtained from the *de*

novo assembly. **B**. The illustration indicates an internal structure of the trophotaenia. An epithelium cell

layer configures an outermost structure of the trophotaenia, which contacts the ovarian luminal fluids. The

layer consists of an enterocyte-like cell with microvilli on the apical surface and vesicles in the cytoplasm.

C-D. Immunohistochemistry using the Cubn antibody or Amn antiserum in the trophotaenia. In both

samples, DAB color development was observed in the epithelial layer. Ep, epithelial layer. Ct, connective

tissue. Scale bar, 50 µm. **E**. Confocal microscopy images of fluorescent immunohistochemistry using the

Cubn antibody in the epithelial layer cells. Yellow triangles indicate the dotted signals in the epithelial cells

of the trophotaenia. Green double-headed arrow indicates the apical surface defined by the flat signal.

Yellow double-headed arrow indicates the cytoplasmic region including the dotted signals. Scale bar, 5

µm. **F.** Immunoelectron microscopy using the Cubn antibody in the epithelial layer cells. Green double-

headed arrow indicates the microvilli on the apical surface. Yellow double-headed arrow indicates the

cytoplasmic region including the dotted signals. Asterisks indicate anti-Cubn signals in the microvilli.

Arrows indicate endocytic vesicles in the invagination phase. Arrowheads indicate intracellular vesicles

after endocytosis. Anti-Cubilin signals were observed in both vesicles. Scale bar, 1 µm.

Figure 4. Proteolysis activities in trophotaenia

A. The illustrations indicate a typical structure of cathepsin L and a comparison of the protease domains of

cathepsin L between *H. sapiens* and *X. eiseni*. The gray filled texts are conserved sequences among the

species. The black triangles indicate protease active sites. **B**. RT-PCR for *cstl.1* exhibited the highest TPM

value in the RNA-Seq analysis. The amplicons were detected as a single band on the expected size

based on the transcript sequences obtained from the *de novo* assembly. **C**. Labeling of acidic organelles

including lysosomal vesicle in the trophotaenia. The lysotracker treatment exhibited red fluorescence in

the epithelial layer (white dotted line). Ep, epithelial layer. Ct, connective tissue. Scale bar, 20 µm. **D**. High

magnification image of the lysotracker-treated epithelial cell layer of the trophotaenia. Yellow dotted line

indicates the epithelial layer and white dotted circles are the nuclei of the epithelial cells. The lysotracker

fluorescence was observed in the cytoplasm. Scale bar, 10 µm. **E**. Measurement of cathepsin L activity

based on fluorescent substrate degradation. The vertical line indicates the accumulation of cleaved

fluorescent substrates, which means cathepsin L activity in the sample solution. The fluorescent value of

487   the trophotaenia lysate sample was increased by 8 h after the reaction started.

	Viviparous mammal (*Mus musculus*)	**Stomachless fish** (*Danio rerio*)	**Viviparous fish** (*Xenotoca eiseni*)
Stage	Suckling	Larva ~ Adult	Intraovarian embryo
Tissue	Intestine (Ileum)	Intestine (LREs)	**Trophotaenia**
Factor	*Dab2*	*cubn*, *dab2*	*cubn*, *amn*, *ctsl.1*
Target	Oral food (Milk)	Oral food (Normal diet)	**Maternal supplement**

Figure 5. Comparison of endocytosis-mediated nutrient intake in vertebrates

Endocytosis-mediated nutrient intake has also been reported in a viviparous mammal and stomachless fish. In these species, oral ingestion macromolecules are absorbed from a region of the intestine via endocytosis. In contrast, intraovarian embryos of the viviparous fish (family Goodeidae) absorbs the maternal component in the ovarian fluids from the trophotaeniae via endocytosis. Endocytosis in each species is predicted to be driven by similar molecular process in the enterocytes of the intestine or the enterocyte-like cells of trophotaenia; however, the biological ontology would be divergent between the viviparous fish (intraovarian growth) and the others (feeding after birth).

Supplementary Information

Cubam receptor-mediated endocytosis in trophotaenia of the intraovarian embryo in a viviparous teleost

Xenotoca eiseni

Atsuo Iida, Kaori Sano, Mayu Inokuchi, Takayuki Suzuki, Mao Kuriki, Maina Sogabe, Daichi Susaki, Kaoru

Tonosaki, Tetsu Kinosita, Eiichi Hondo

Correspondence: Atsuo Iida

E-mail: tol2.4682@gmail.com

This PDF file includes:

      Figures S1 to S5
      Tables S1 to S7

Other supplementary materials for this manuscript include the following:

      Dataset S1

Additional Figures

Figure S1. Tissue structures of trophotaenia in *X. eiseni* embryo.

A. A HE-stained section (A) and illustration (A') of the trophotaenia of the 4th week embryo. Ep, epithelium

layer cells. Bd, blood vessel. Ct, connective tissue. Bar, 500 μm. **B**. Enlarged image of the epithelial layer

of the trophoatenia. Ep, epithelial layer. Nu, nucleus. Ct, connective tissue. Bar, 20 μm. **C**. Electron

microscopy for an epithelial layer cell of the trophoatenia. Microvilli are localized on the apical surface of the

cell. Arrows indicate endocytic vesicles in the invagination phase. Arrowheads indicate intracellular vesicles

after endocytosis. Scale bar, 1 μm.

Figure S2. Immunohistochemistry of Cubn or Amn in trophoatenia.

A-B. Immunohistochemistry using the Cubn antibody (A) or Amn antiserum (B) in the apical terminal of

trophotaenia. Scale bar, 100 μm. C-D. Immunohistochemistry using the Cubn antiserum in the apical

terminal (C) or enlarged image for the epithelial layer (D) of trophotaenia. Ep, epithelium layer. Ct,

connective tissue. Scale bar, 100 μm (C) or 50 μm (D).

Figure S3. Fluorescent immunohistochemistry using Cubn antibody in the trophotaenia.

Confocal microscopy for the trophotaenia of *X. eiseni* embryo at 4th week gestation. Anti-Cubn signals are

observed in the epithelial layer cells of the trophotaenia. Scale bar, 20 μm.

Figure S4. Expression analysis for the endocytic adaptor and vesicle coating protein genes.

```
H.sapiens DAB2 (NP_001334.2) MSNEVETSATN-GQPDQ--------------QAAPKAPSKKEKKKGPEK 34
X.eiseni  dab2              MSTEVENSVPATADPSSPSTASTTISPTTSPVTTTPKAQLKKEKKKVPEK 50
                            **.***.*..  .:*..                ::***  ****** ***

H.sapiens DAB2 (NP_001334.2) TDEYLLARFKGDGVKYKAKLIGIDDVPDARGDKMSQDSMMKLKGMAAAGR 84
X.eiseni  dab2              TDEYLLSRFQGDGVRYKAKLIGIDDVSEARGDKMCQDSMMKLKGMAVAAR 100
                            ******:**:****:***********.:******.**********.*.*

H.sapiens DAB2 (NP_001334.2) SQGQHKQRIWVNISLSGIKIIDEKTGVIEHEHPVNKISFIARDVTDNRAF 134
X.eiseni  dab2              SQGKHKQRIWVNISMSGLKIIDEKSGVIEHEHVVNKISFIARDVTDNRAF 150
                            ***:**********:**:******:******* *****************

H.sapiens DAB2 (NP_001334.2) GYVCGGEGQHQFFAIKTGQQAEPLVVDLKDLFQVIYNVKKKEEEK-KKIE 183
X.eiseni  dab2              GYVCGAEGQHQFFAIKTAQQAEPLVIDLKDLFQVIFNMRKKEAESSQKAE 200
                            *****.***********.*******:*********:*::*** *. :* *

H.sapiens DAB2 (NP_001334.2) EASKAVENGSEALMILDDQTNKLKSGVDQMDLFGDMSTPPDLNSPTESKD 233
X.eiseni  dab2              NGSAVVENGG-ALQSTDGES-KAAQPVEQLDLFGDITTPPDIRAP----- 243
                            :.* .****. **   *.:: *  . *:*:*****::****:.:*

H.sapiens DAB2 (NP_001334.2) ILLVDLNSEIDTNQNSLRENPFLTNGITSCSLPRPTPQASFLPENAFSAN 283
X.eiseni  dab2              --------------------------------------------------

H.sapiens DAB2 (NP_001334.2) LNFFPTPNPDPFRDDPFTQPDQSTPSSFDSLKSPDQKKENSSSSSTPLSN 333
X.eiseni  dab2              -------------------------------------NSGSS------ 248
                                                                 **.**

H.sapiens DAB2 (NP_001334.2) GPLNGDVDYFGQQFDQISNRTGKQEAQAGPWPFSSSQTQPAVRTQNGVSE 383
X.eiseni  dab2              --------------------------------------------------

H.sapiens DAB2 (NP_001334.2) REQNGFSVKSSPNPFVGSPPKGLSIQNGVKQDLESSVQSSPHDSIAIIPP 433
X.eiseni  dab2              --------------------------------------------------

H.sapiens DAB2 (NP_001334.2) PQSTKPGRGRRTAKSSANDLLASDIFAPPVSEPSGQASPTGQPTALQPNP 483
X.eiseni  dab2              ------------------DLFGTDLFVPPVS------SETSPADLFNNTP 274
                                              **:.:*:*.****      * *.. .: .*

H.sapiens DAB2 (NP_001334.2) LDLFKTSAPAPVGPLVGLGGVTVTLPQAGPWNTASLVFNQSPSMAPGAMM 533
X.eiseni  dab2              TNTVPSTIPALG--SLQLGPTATSVPAVGMWGTSPAVPAMFP--MPGIVT 320
                             : . :: **     : ** .:.::* .* *.*:. *    *   ** :

H.sapiens DAB2 (NP_001334.2) GGQPSGFSQPVIFGT--SPAVSGWNQPSPFAASTPPPVPVVWGPSASVAP 581
X.eiseni  dab2              PGLRPNFPQPTAFGVPMQPPVWAPQVVPQFSAAPLSPPHLQWGQPATS-- 368
                             *  ..*.**. **.  .*.* .:  . *:*:. .*  : ** .*:

H.sapiens DAB2 (NP_001334.2) NAWSTTSPLGNPFQSNIFPAPAVSTQPPSMHSSLLVTPPQPPPRAGPPKD 631
X.eiseni  dab2              ----------NPFQ-------AMGDHGPSRP------PPRPPVKETPPR- 394
                                      ****       *:. : **      **:** :  **:

H.sapiens DAB2 (NP_001334.2) ISSDAFTALDPLGDKEIKDVKEMFKDFQLRQPPAVPARKGEQTSSGTLS- 680
X.eiseni  dab2              VENSAFTALDPLGDKEKKTGKDMFKNFQIAKLPAIPARKGELMPSSTPPP 444
                            :...************ *  *:***:**: : **:****** .*.* .

H.sapiens DAB2 (NP_001334.2) ------AFASYFNSKVGIPQENADHDDFDANQL-LNKINEPPKPAPRQVS 723
X.eiseni  dab2              ANKESVPFDEYFSNKVGLAQDAADHDDFDINQMSILDGNDAPKQTPVQFT 494
                                 .*  .**..**:.*: *******  **:  :  . *:.** :* *.:

H.sapiens DAB2 (NP_001334.2) LPVTKS------TDNAFENP----------------FFKDSFGSSQASV 750
X.eiseni  dab2              APAAAPSFTPDLLDAAFSSAPVPNSSAPTLGQDLSHDMFDQAFGAPDPNP 544
                             *.:  .        * **...                :*.::**:.:..

H.sapiens DAB2 (NP_001334.2) ASSQPVSS----------EMYRDPFGNPFA 770
X.eiseni  dab2              FGAPPVAMNTVAQTSGSTDAFGDAFGNPFA 574
                             .: **:            : : *.******
```

Figure S5. Comparison amino acid sequences between *H. sapiens* DAB2 and *X. eiseni* dab2-like.

554 **Additional Tables**

		cubn	amn	lrp1aa	lrp2a
	#1	1421	745	6	12
	#2	768	966	4	6
3rd week	#3	1414	1269	10	10
	Average	**1201**	**993**	**7**	**9**
	#1	1528	1139	14	12
	#2	1464	818	10	7
4th week	#3	1250	1032	8	12
	Average	**1414**	**996**	**11**	**10**

555 **Table S1. The TPM values for endocytic receptor genes in the RNA-seq**

		ap2a1	*ap2b1*	*ap2m1a*	*ap2s1*	*ldlrapab*	*numb*
3rd week	#1	75	91	383	365	7	19
	#2	60	54	346	318	6	15
	#3	73	98	360	368	7	19
	Average	**69**	**81**	**363**	**350**	**7**	**18**
4th week	#1	78	84	349	359	8	17
	#2	89	83	374	370	8	15
	#3	82	85	385	371	6	13
	Average	**83**	**84**	**369**	**367**	**7**	**15**

558 **Table S2. The TPM values for adaptor genes in the RNA-seq**

		clta	cltb	cltc	cav2	cav3	flot1b	flot2a
3rd week	#1	851	67	337	1	0	229	91
	#2	680	76	209	0	0	196	50
	#3	744	73	308	2	0	221	89
	Average	**758**	**72**	**285**	**1**	**0**	**215**	**77**
4th week	#1	664	97	328	4	0	228	228
	#2	739	127	366	2	0	238	238
	#3	817	104	343	3	0	234	234
	Average	**740**	**109**	**346**	**3**	**0**	**233**	**233**

561 **Table S3. The TPM values for vesicle coating protein genes in the RNA-seq**

		ctsa	ctsb	ctsc	ctsh	ctsk	ctsla	ctsl.1	ctso	ctss2.1	ctsz
3rd week	#1	862	474	1867	52	11	831	12749	3	2144	66
	#2	691	234	1113	31	17	736	8412	5	1656	63
	#3	939	429	1907	52	14	939	11227	7	2042	63
	Average	**831**	**379**	**1629**	**45**	**14**	**835**	**10796**	**5**	**1947**	**64**
4th week	#1	751	2686	2273	47	24	960	17795	6	2494	530
	#2	749	2559	2320	55	24	809	18334	7	2500	551
	#3	720	1944	1884	48	18	924	15906	8	2342	523
	Average	**740**	**2396**	**2159**	**50**	**22**	**898**	**17345**	**7**	**2445**	**535**

564 **Table S4. The TPM values for protease genes in the RNA-seq**

		1 hour	2 hours	3 hours	4 hours	5 hours	6 hours	7 hours	8 hours
No lysate	#1	68.8	67.8	67.2	65.8	65.5	65.8	65.6	65.2
	#2	66.5	65.9	64.8	62.7	64.0	64.4	63.8	62.7
	#3	66.8	65.4	69.8	63.1	63.3	62.8	61.8	62.6
	Average	**67.4**	**66.3**	**67.3**	**63.8**	**64.3**	**64.3**	**63.8**	**63.5**
Lysate	#1	260.3	345.0	437.7	530.4	579.7	637.3	699.6	694.1
	#2	264.9	351.6	460.3	551.3	617.0	678.4	727.0	748.2
	#3	256.6	339.2	441.8	522.7	587.1	642.9	692.5	720.8
	Average	**260.6**	**345.3**	**446.6**	**534.8**	**594.6**	**652.9**	**706.3**	**721.0**
Lysate + inhibitor	#1	94.0	96.2	98.0	99.6	101.0	106.8	104.4	101.9
	#2	88.1	88.8	91.1	91.5	93.4	94.0	98.0	98.2
	#3	88.3	88.4	89.4	93.1	94.2	96.4	99.4	101.0
	Average	**90.1**	**91.2**	**92.8**	**94.7**	**96.2**	**99.0**	**100.6**	**100.4**

567 **Table S5. The fluorescent values indicated Cathepsin L-dependent proteolysis.**

Gene	Accession #		Primer (5'-3')	Related figures
cubn	LC595284	F	ATGGCTGTTACAGTGCAGCCTTCA	3A
		R	GGCATGTACATACAGGGAATGCTG	
amn	LC595285	F	TGCCCTTTACAAGCAGTGGATTCC	3A
		R	GCGGATTAGCACAACCACAATCAC	
lrp2a	LC595286	F	ATGTGGAGAACACCGCTGCTTCAA	3A
		R	CCACATGGAGCAGTCATCGAAATC	
ap2a1	LC595287	F	GCTGTGTCGAAGGGAGATGGAATG	S4
		R	AGAGCTAGATAGCGCAGATTGGTC	
ap2b1	LC595288	F	GTCATTGCTGCCATGACTGTTGGC	S4
		R	CCACAATAGCCTCCTGAACCACAT	
ap2m1a	LC595289	F	GACGTAATGACGGCCTACTTTGGC	S4
		R	CAGCAGCGGGTTTCATAGATGCCA	
ap2s1	LC595290	F	ATGATCCGCTTCATCCTCATCCAG	S4
		R	CTCTAGTGACTGAAGCATGAGGAG	
ldlrap1b	LC595291	F	GGACGCTTTAAAATCCGCTGGAAG	S4
		R	GCAGCTGAGTCTGTCTCATCTTGG	
numb	LC595296	F	GAATAAGCTACGGCAGAGCTTCCG	S4
		R	TTGGGCATTGGAGCGGAAGAGAAC	
clta	LC595292	F	GGATGATTTTGACATGCTGAACGC	S4
		R	TAACGGACCAGCGGGGACTGCTTT	
cltc	LC595293	F	CTCAGATCCTGCCAATTCGCTTTC	S4
		R	TCTTCTTCCACGCAAACAGACAGC	
ctsl.1	LC595294	F	CAGCATCTCTCTGGAAGATCTCGA	4B
		R	CAGACTAGTGGATAACTTGCTGCG	
actb	LC595295	F	ATGGAAGATGAAATCGCCGCACTG	3A, 4B
		R	GAAGCATTTACGGTGGACGATGGA	

570 **Table S6. Primers**

Name	Immunogen	Host	Supplier	ID	Related figures
CUBN Polyclonal Antibody (anti-Cubn)	Human CUBN	Rabbit	Invitrogen	PA5-83684	3C, 3E, 3G, S2, S3
Rabbit IgG Isotype Control (Control IgG)	N/A	Rabbit	Invitrogen	02-6102	3C, 3E, 3G, S2, S3
Cubn antiserum (anti-Cubn #2)	*X. eiseni* cubn	Mouse	N/A	N/A	S2
Amn antiserum (anti-Amn)	*X. eiseni* amn	Mouse	N/A	N/A	3D, S2
Preimmune serum (Control serum)	N/A	Mouse	N/A	N/A	3D, S2
Goat anti-Mouse IgG (H+L) Secondary Antibody, Alexa Fluor Plus 488	Rabbit IgG	Goat	Invitrogen	A32723	3E, S3
Horse Anti-Mouse IgG Antibody (H+L), Peroxidase	Rabbit IgG	Horse	VECTOR	PI-2000	3D, S2
Goat Anti-Rabbit IgG Antibody (H+L), Peroxidase	Mouse IgG	Goat	VECTOR	PI-1000	3C, S2
Goat Anti-Rabbit IgG Antibody (H+L), Biotinylated	Rabbit IgG	Goat	VECTOR	BA-1000	3G

573 **Table S7. Antibodies**

575