Title

Cubam receptor-mediated endocytosis in hindgut-derived pseudoplacenta of a

viviparous teleost *Xenotoca eiseni*

Running title

Endocytosis in pseudoplacenta of teleost

Authors

Atsuo Iida[1, *], Kaori Sano[2], Mayu Inokuchi[3], Jumpei Nomura[1], Takayuki Suzuki[1], Mao

Kuriki[4], Maina Sogabe[4], Daichi Susaki[5], Kaoru Tonosaki[5], Tetsu Kinoshita[5], Eiichi

Hondo[1]

Affiliations

1. Department of Animal Sciences, Graduate School of Bioagricultural Sciences,

Nagoya University, Tokai National Higher Education and Research System, Nagoya,

Aichi, Japan.

2. Department of Chemistry, Faculty of Science, Josai University, Sakado, Saitama,

Japan.

3. Department of Aquatic Bioscience, Graduate School of Agricultural and Life Sciences,

University of Tokyo, Bunkyo, Tokyo, Japan

4. Department of Regeneration Science and Engineering, Institute for Frontier Life and

Medical Sciences, Kyoto University, Kyoto, Kyoto, Japan.

5. Kihara Institute for Biological Research, Yokohama City University, Yokohama,

Kanagawa, Japan.

Correspondence: Atsuo Iida

E-mail: tol2.4682@gmail.com

Keywords

endocytosis, Goodeidae, proteolysis, pseudoplacenta, teleost, viviparity

Summary statement

33   Here, we report that an endocytic pathway is a candidate for nutrient absorption in

34   pseudoplacenta of a viviparous teleost. The trait may have developed from common

35   intestinal mechanism among vertebrates.

Abstract

Nutrient transfer from mother to the embryo is essential for reproduction in viviparous

animals. In the viviparous teleost *Xenotoca eiseni* belonging to the family Goodeidae,

the intraovarian embryo intakes the maternal component secreted into the ovarian fluid

via the trophotaenia. Our previous study reported that the epithelial layer cells of the

trophotaenia incorporate a maternal protein via vesicle trafficking. However, the

molecules responsible for the absorption were still elusive. Here, we focused on Cubam

(Cubilin-Amnionless) as a receptor involved in the absorption, and cathepsin L as a

functional protease in the vesicles. Our results indicated that the Cubam receptor is

distributed in the apical surface of the trophotaenia epithelium and then is taken into the

intracellular vesicles. The trophotaenia possesses acidic organelles in epithelial layer

cells and cathepsin L-dependent proteolysis activity. This evidence does not conflict with

our hypothesis that receptor-mediated endocytosis and proteolysis play roles in

maternal macromolecule absorption via the trohotaenia in viviparous teleosts. Such

nutrient absorption involving endocytosis is not a specific trait in viviparous fish. Similar

processes have been reported in the larval stage of oviparous fish or the suckling stage

of viviparous mammals. Our findings suggest that the viviparous teleost acquired

trophotaenia-based viviparity from a modification of the intestinal absorption system

common in vertebrates. This is a fundamental study to understand the strategic variation

of the reproductive system in vertebrates.

Introduction

August Krogh wrote, "*For such a large number of problems there will be some animal of choice or a few such animals on which it can be most conveniently studied*" [1]. This study aimed to investigate the molecular mechanism of maternal nutrient absorption in a species-specific pseudoplacenta of a viviparous teleost species belonging to the family Goodeidae.

Viviparity is a reproduction system, whereby the oocyte is fertilized in the female body, and subsequent embryo growth occurs with maternal component supply. Each viviparous animal has acquired processes specialized to the gestation in both the mother and embryo [2]. The placenta and umbilical cords in viviparous mammals are major components of the process for mother-to-embryo material transport [3,4]. Other viviparous components such as the extended yolk sac or pseudoplacenta are known in several viviparous vertebrates, except mammals [5].

The family Goodeidae is a freshwater small teleost distributed in Mexico, which includes approximately 40 viviparous species [6]. They possess trophotaenia, which is a hindgut-derived pseudoplacenta that absorbs the maternal component [7,8]. Trophotaenia is a ribbon-like structure consisting of a single epithelial layer, internal blood vessels, and connective tissues [9,10]. The epithelial cell is like an enterocyte in the intestine. Electron microscopy indicated that microvilli form in the apical side of the cell and intracellular vesicles in the cytoplasm [11]. Since the 1980s, these structures have been believed to be involved in maternal component absorption [12]. The nature of the maternal component was predicted to be proteins or other macromolecules secreted into the serum or ovarian fluids; however, no one has experimentally determined its distinct component [13,14]. Recently, we demonstrated that a yolk protein vitellogenin is

secreted into the ovarian lumen of pregnant females, and the intraovarian embryo

absorbs the nutrient protein via the trophotaenia using a goodeid species *Xenotoca*

eiseni [15]. In that study, enterocyte-like microvilli and intracellular vesicles were also

observed in the epithelial cells of the trophotaenia. We hypothesized that the epithelial

layer cell in the trophotaenia absorbs the maternal protein and/or other components as

macromolecules because the ovarian lumen lacks proteolysis activity like the digestive

intestine. However, the molecules responsible for the trophotaenia-mediated

macromolecule absorption have not been reported.

Vacuolated enterocytes involved in macromolecule absorption have also been reported

in other vertebrate species, including suckling mammals and stomachless fish [16-18].

Park et al. [19] reported that the scavenger receptor complex Cubam (Cubilin-

Amnionless) and Dab2 are required for macromolecule uptake in zebrafish juveniles.

Furthermore, the conditional knockout of *Dab2* in mice led to stunted growth and severe

protein malnutrition at the suckling stage [19]. Based on this report, we discuss the

commonality of molecular process for the macromolecule absorption between the

intestinal enterocytes and the trophotaenia [15]. However, the molecules responsible for

macromolecule absorption in the trophotaenia are still elusive.

Here, we report candidate molecules for nutrient uptake and subsequent proteolysis in

the trophotaenia of *X. eiseni*. An RNA-Seq for trophoteniae indicated candidate receptor

molecules and endocytosis-associated proteases expressed in the trophotaenia.

Immunohistochemistry and biochemical assays suggested the presence and functions

of the candidate factors in the trophotaenia.

Methods

Animal experiments

Methods

Animal experiments

This study was approved by the ethics review board for animal experiments at Nagoya

University (Approval number: AGR2020028). We sacrificed live animals in minimal

numbers under anesthesia according to the institutional guidelines.

Fish breeding

X. eiseni was purchased from Meito Suien Co., Ltd. (Nagoya, Japan). Adult fish were

maintained in freshwater at 27 °C under a 14:10-h light: dark photoperiod cycle. Fish

were bred in a mass-mating design, and approximately 30 adult fish were maintained for

this study. The juveniles were fed live brine shrimp larvae, and the adults were fed Hikari

Crest Micro Pellets and ultraviolet-sterilized frozen chironomid larvae (Kyorin Co., Ltd.,

Himeji, Japan). To accurately track the pregnancy period, the laboratory-born fish were

crossed in a pair-mating design, and the mating behavior was recorded.

Sample collection

Fish samples were anesthetized using tricaine on ice before the surgical extraction of

tissues or embryos. The obtained samples were stored on ice until subsequent

manipulations. In this study, we dissected approximately 10 pregnant females and

extracted 15–30 embryos in each operation.

Methods

Animal experiments

This study was approved by the ethics review board for animal experiments at Nagoya

University (Approval number: AGR2020028). We sacrificed live animals in minimal

numbers under anesthesia according to the institutional guidelines.

Fish breeding

X. eiseni was purchased from Meito Suien Co., Ltd. (Nagoya, Japan). Adult fish were

maintained in freshwater at 27 °C under a 14:10-h light: dark photoperiod cycle. Fish

were bred in a mass-mating design, and approximately 30 adult fish were maintained for

this study. The juveniles were fed live brine shrimp larvae, and the adults were fed Hikari

Crest Micro Pellets and ultraviolet-sterilized frozen chironomid larvae (Kyorin Co., Ltd.,

Himeji, Japan). To accurately track the pregnancy period, the laboratory-born fish were

crossed in a pair-mating design, and the mating behavior was recorded.

Sample collection

Fish samples were anesthetized using tricaine on ice before the surgical extraction of

tissues or embryos. The obtained samples were stored on ice until subsequent

manipulations. In this study, we dissected approximately 10 pregnant females and

extracted 15–30 embryos in each operation.

RNA-Seq

Total RNA was extracted from trophotaenae of the 3rd or 4th week of the embryo

extracted from the pregnant female using the RNeasy Plus Mini kit (QIAGEN). A total of

six samples were obtained from three 3rd week embryos and three 4th week embryos.

Next-generation sequencing (NGS) was outsourced to Macrogen Japan Corp. (Kyoto,

Japan) using NovaSeq6000 (Illumina, Inc., San Diego, CA, USA). Approximately 60

million 150-bp paired-end reads were obtained in each sample. The NGS data was

deposited to the DNA Data Bank of Japan (DDBJ, ID: DRA011209). *De novo assembly*

and mapping to the reference sequence were performed by CLC Genomics Workbench

(Filgen, Inc., Nagoya, Japan). The published transcript sequences of *Poecilia reticulata*

(NCBI Genome, ID: 23338) were used as a reference. *P. reticulata* is one of the species

related to *X. eiseni* for which the transcript sequences have been published, and it

belongs to the order Cyprinodontiformes. The transcript sequences of *X. eiseni* were

deposited into the DDBJ. The accession numbers are listed in Table 1.

Phylogenetic analysis

Amino acid sequences for Cubn, Amn, or Cathepsin L of *Homo sapiens* (human), *Mus*

musculus (house mouse), *Danio rerio* (zebrafish), *Oryzias latipes* (medakafish), *P.*

reticulata (guppy), *Xiphophorus maculatus* (platyfish), and *X. eiseni* were collected from

the NCBI protein database (https://www.ncbi.nlm.nih.gov/protein/). Phylogenetic trees

were constructed by the neighbor-joining method with 1,000 bootstrap replicates in

MEGAX (Ver.10.1.8) software (https://www.megasoftware.net/).

Reverse transcription (RT) PCR

Total RNA was extracted from tissues or whole embryos using the RNeasy Plus Mini kit

and reverse-transcribed using SuperScript IV reverse transcriptase (Thermo Fisher

Scientific). PCR was carried out using KOD-FX-Neo (Toyobo, Osaka, Japan) under the

following conditions: 100 s at 94 °C, followed by 32 cycles of 20 s at 94 °C, 20 s at

60 °C, 60 s at 72 °C, and 120 s at 72 °C. Primer sequences are listed in Table 1.

Antibodies and antiserums

Antiserums against Cubn and Amn were generated in this study. The antigen

sequences were 6× His-tagged peptide of 151 amino acids (aas) corresponding to the

intermediate region of *X. eiseni* Cubn (Accession#, BCN28455; aa 691–841) and 6×

His-tagged peptide of 247 aas corresponding to the C-terminal of *X. eiseni* Amn

(Accession#, BCN28456). The experimental procedure has been described in our

previous study [15]. All antibodies and antiserums used in this study are listed in Table

S1.

Immunohistochemistry

Tissue samples were fixed in 4.0% paraformaldehyde/phosphate-buffered saline

(PFA/PBS) at 4 °C overnight. Samples were permeabilized using 0.5% TritonX-100/PBS

at room temperature for 30 min. Endogenous peroxidase was inactivated by 3.0 %

hydrogen peroxide/PBS for 10 min. Then, the sample was treated with Blocking-One

solution (Nacalai Tesque, Kyoto, Japan) at room temperature for 1 h. Primary antibody

or antiserums were used at 1:500 dilution with Blocking-One solution. Samples were

incubated with primary antibody or antiserum at 4 °C overnight. Secondary antibodies

were used at a 1:500 dilution in 0.1% Tween-20/PBS. Samples were treated with the

secondary antibody solution at 4 °C for 2 h. We performed a 3,3'-Diaminobenzidine

Tetrahydrochloride (DAB) color development using the DAB Peroxidase Substrate Kit,

ImmPACT (Vector Laboratories, Inc., Burlingame, CA, USA). Microscopic observation

was performed using an Olympus BX53 microscope and photographed using a DP25

digital camera (Olympus, Shinjuku, Japan).

Fluorescent Immunohistochemistry

Tissue samples were fixed in 4.0% PFA/PBS at 4 °C overnight. Samples were

permeabilized using 0.5% TritonX-100/PBS at room temperature for 30 min.

Endogenous peroxidase was inactivated by 3.0 % hydrogen peroxide/PBS for 10 min.

Then, the sample was treated with Blocking-One solution (Nacalai Tesque, Kyoto,

Japan) at room temperature for 1 h. Primary antibody (anti-Cubn) was used at a 1:500

dilution with Blocking-One solution. Samples were incubated with primary antibody at

4 °C overnight. Secondary antibody was used at a 1:500 dilution in 0.1% Tween-20/PBS

with 4',6-diamidino-2-phenylindole (DAPI). Samples were treated with the secondary

antibody solution at 4 °C overnight. Microscopic observation was performed using a

Leica TCS SP8 microscope (Leica Microsystems, Wetzlar, Germany).

Immunoelectron microscopy

Embryo samples were fixed in 4.0% PFA/PBS. Fixed samples were washed in PBS,

then reacted with a primary antibody (anti-Cubn) at 4 °C overnight, and then reacted

with biotinylated anti-rabbit IgG (Vector, Burlingame, CA, USA) at room temperature for

2 h. Samples were performed with the avidin–biotin–peroxidase complex kit (Vector),

and visualized with 0.05% DAB (Dojindo Laboratories, Kumamoto, Japan) and 0.01%

hydrogen peroxide in 50 mM Tris buffer (pH 7.2) at room temperature for 10 min. The

procedure for electron microscopy is described in our previous study [15].

Labeling acidic organelles

The live embryos immediately after extraction from the pregnant female at the 4^{th}-week

post-mating were incubated in PBS with a 1:1000 dilution of LysoTracker® Red (Thermo

Fisher Scientific) for 1 h at room temperature. The samples were fixed with 4.0 %

PFA/PBS and stained with DAPI. Microscopic observation was performed using a Leica

DM5000 B microscope (Leica Microsystems, Wetzlar, Germany).

Measurement of Cathepsin L activity

The trophotaenia lysate was prepared from six littermate embryos obtained from the

pregnant females at the 4^{th}-week post-mating. The trophotaeniae were manually

extracted from the embryo under a microscope. Proteolysis detection was performed

using the Cathepsin L Activity Fluorometric Assay Kit (BioVision, Inc., Milpitas, CA,

USA). The fluorescence intensities were measured using a Qubit 4 Fluorometer

(Thermo Fisher Scientific).

Results

Gene expression in trophotaenia

Our previous findings and the hypothesis based on that are described in Figure 1A. In

a viviparous teleost, *X. eiseni*, the embryo is raised in the ovary while receiving maternal

nutrients. The trophotaenia is a pseudoplacenta that plays a role in the absorption of

maternal nutrients consisting of proteins and other supplements. Based on previous

studies, we hypothesized that several maternal proteins are absorbed by endocytosis-

mediated proteolysis in the epithelial cells of the trophotaenia. To verify this hypothesis,

we explored candidate genes for receptor, adaptor, vesicle, and protease proteins that

are highly expressed in the trophotaenia. RNA-Seq analyses were performed using total

RNA extracted from the trophotaeniae of the 3rd- or 4th-week embryos (Figure 1B). We

selected candidate genes and compared their predicted expression level as transcription

per million (TPM) values that were calculated using the known transcript sequence of *P.*

reticulata as reference. RNA-Seq suggested that *cubilin* (*cubn*) and *amnionless (amn*)

were highly expressed in the trophotaeniae; however, other co-receptor genes (*lrp1aa*,

lrp2a) were considerably lower (Figure 1C, Table 2). Adaptor protein-2 (AP2) subunit

genes (*ap2a1*, *ap2b1*, *ap2m1a*, and *ap2s1*) were more highly expressed than the other

family adaptor genes (*ldlrap1b* and *numb*) (Figure 1C, Table 2). Two families of the

vesicle coating protein genes (*clta*, *cltbb*, *cltc*; *flot1b*, *flot2a*) were expressed higher than

the vesicle proteins classified in other families (*cav2* and *cav3*) (Figure 1C, Table 2). The

lysosomal endopeptidase enzyme gene (*ctsl.1*) exhibited a high TPM value (> 10,000)

rather than that of not only protease family genes but also most of the genes expressed

in the trophotaeniae (Figure 1C, Table 2). In this investigation, we assumed that the 3rd

week is a good growth period (mid pregnancy) and that the 4th week is a poorer growth

(late pregnancy) period; thus, we expected that the transcriptomes related to the nutrient

transport would vary between these stages. However, the trend of the predicted TPM

values in most of the selected genes was not noticeably different between the samples.

In the 4th week, only one *clathrin* and four *cathepsin genes* were significantly higher than

levels in the 3rd week (Figure 1C).

Sequences for Cubam receptor genes

A Cubam receptor is known to be a membrane-bound multi-ligand receptor consisting

of Cubn and Amn (Figure 2A). The secreted protein Cubn specifically binds to the

transmembrane protein Amn; thus, the CUB domain that associates with the ligands are

localized on the apical surface of the plasma membrane. We obtained the amino acid

sequences of *X. eiseni* Cubn and Amn by *de novo* assembly of NGS reads from the

trophotaeniae. The assembled sequences of *X. eiseni* were calculated to be close to the

orthologous sequences of *P. reticulata* and *X. maculatus* (Figure 3A and B). The

sequences of Cubn and Amn were compared among four vertebrate species, *H.*

sapiens, *D. rerio*, *P. reticulata*, and *X. eiseni.* The binding motifs were conserved among

the species (Figure 2B and C). A phenylalanine-any-asparagine-proline-any-

phenylalanine (FXNPXF) amino acid sequence in the intracellular domain of Amn is

known to be a conserved motif used to bind adaptor proteins. The motifs were also

conserved among the species (Figure 2D).

Distribution of Cubilin and Amnionnless in trophotaenia

To validate the expression of *cubn* and *amn* in the trophotaenia, semi-quantitative RT-

PCR analyses were performed using total RNAs extracted from the whole-embryo

including the trophotaeniae, isolated trophotaeniae, and adult skeletal muscle. The

muscle sample was used as a control for tissue with low endocytosis activity. The gene

expression patterns did not conflict between the results of RT-PCR and RNA-Seq. In the

embryo and trophotaenia, both *cubn* and *amn* were more highly expressed than *lrp2a*

(Figure 4A). Conversely, no or few expressions of the target genes except *actb* as a

positive control were detected in the adult muscle. Next, to detect protein localization,

immunohistochemistry was performed using antibodies against Cubn or Amn. In both

proteins, strong signals were observed in the epithelial monolayer of the trophotaenia,

while few background noises were detected in the control assay (Figure 4B-F, see also

Figure S1 and S2). Confocal microscopy revealed the cellular distribution of the anti-

Cubn signals. The signals in the apical surface of the epithelial cell were detected as a

homogeneous distribution, while almost all signals in the cytoplasm were captured as a

dot pattern (Figure 4G, Figure S3). Immunoelectron microscopy revealed that the

microvilli were distributed on the apical surface of the trophotaenia epithelium, and

intracellular vesicles were observed in the cytoplasm; furthermore, anti-Cubn signals

were distributed in the intracellular vesicles and overlapped with the microvilli on the

apical surface (Figure 4H).

Proteolysis activity in trophotaenia

Cathepsin L is a lysosomal cysteine proteinase characterized by three conserved

protease regions and active sites consisting of cysteine, histidine, or asparagine (Figure

5A). The functional regions were conserved in the *X. eiseni* Cathepsin L protein

translated from the coding sequence of *ctsl.1*. The whole amino acid sequence of *X.*

eiseni Cathepsin L was calculated to be close to the orthologous proteins of *P. reticulata*

and *X. maculatus* (Figure 3C). RT-PCR analysis revealed strong expression of *ctsl.1* in

trophotaenia (Figure 5B). To identify the type of cells that have proteolysis activity in

trophotaenia, acidic organelles including lysosomes and endosomes were detected

using a fluorescent probe. LysoTracker® analysis indicated the presence of acidic

organelles in the epithelial layer cells (Figure 5C). The signals were distributed in the

cytoplasm and were not components of the nuclei (Figure 5D). According to the RNA-

Seq analysis, *ctsl.1* was presumed to be the highest expressed *cathepsin* gene in

trophotaenia; thus, we calibrated the proteolysis activity of cathepsin L in trophotaenia

using a fluorescent substrate-based measurement system. The fluorescence, indicating

substrate digestion, was significantly higher in the trophotaenia lysate than in the control

at 1 h after reagent mixing. Furthermore, the intensity for the lysate continued to

increase for 7 h (Figure 5E, Table 3). Conversely, the increase in intensity was strongly

suppressed by a cathepsin L inhibitor. The fluorescent values at each timepoint were

statistically different from each other.

Adaptors and vesicle coating proteins

The expression of candidate genes for adaptors (*ap2a1*, *ap2b1*, *ap2m1a*, *ap2s1*,

ldlrap1b, and *numb*) and vesicle coating proteins (*clta* and *cltc*) were confirmed by RT-

PCR (Figure 6). We determined an incomplete transcript for *X. eiseni* Dab2 from the *de*

novo assembly; however, it lacked internal sequences in comparison with the proteins in

other vertebrates (Figure S4). Furthermore, no amplicons were obtained by RT-PCR

using primers designed based on the Dab2-like sequence.

Discussion

RNA-Seq analyses revealed high expression of the genes for receptor-mediated

endocytosis in the trophotaenia of *X. eiseni* embryos. Cubn and Amn form a Cubam

receptor complex that associates with Vitamin-B12, albumin, transferrin, and other

ligands [20,21]. The predicted amino acid sequences for *X. eiseni* Cubn and Amn

obtained from the *de novo* assembly both possessed a conserved motif which allowed

binding to each other, and Amn retained adaptor binding sites in the intracellular region

[21, 22]. Immunohistochemical analysis indicated the presence of Cubn not only in the

apical surface of the epithelial layer but also in the intracellular vesicles in the cells,

suggesting the incorporation and recycling of endocytic receptors [23,24]. This evidence

supports the idea that Cubam plays a role as a receptor for the intraovarian nutrients

and is involved in endocytosis. We also indicated the low presence of Lrp2 (also known

Megalin), which is a major co-receptor for Cubam [25]. This indicated that Cubam works

independently or in cooperation with other co-receptors, except Lrp2, in the

trophotaenia. In zebrafish, a previous study reported that Cubam-dependent

endocytosis in the lysosome rich enterocytes (LREs) does not require the presence of

Lrp2 [19]. This supports our hypothesis; however, we do not exclude the possibility that

our sequencing and alignment processes could not catch other *lrp2* orthologous genes

in *X. eiseni*. Park et al. [19] also reported that Dab2 is an essential adaptor molecule not

only in the zebrafish larval intestine but also in the endocytic nutrient absorption in the

ileum of suckling mice. However, we obtained only an incomplete sequence for the *X.*

eiseni Dab2-like protein without an internal region compared with mammalian orthologs.

This may be an alternative splice form with less endocytic activity [26,27]. Furthermore,

we did not detect Dab2-like expression by RT-PCR, and we could not exclude the

possibility that the predicted Dab2-like sequence is caused by an assembly error. Thus,

the contribution of Dab2 to endocytosis in the trophotaenia has not been confirmed in

this study. The other adaptor or vesicle coating proteins we detected were typical co-

factors for receptor-mediated endocytosis. Conversely, Caveolin is a vesicle coating

protein involved in receptor-independent endocytosis [28]. Thus, a low expression of

caveolin genes (*cav2* and *cav3*) does not conflict with the activation of Cubam-mediated

endocytosis. Additionally, biochemical assays supported that cathepsin L is an active

protease that functions in the intracellular vesicles that is configured following

membrane budding. This evidence supports the idea that Cubam-mediated endocytosis

and cathepsin L-dependent proteolysis is one of the key mechanisms for the absorption

of the maternal component in *X. eiseni* embryos.

Cubam is also known to be a scavenger receptor involved in nonspecific protein uptake

[20]. In this study, we did not determine the Cubam ligands in the trophotaenia.

However, the candidates are limited to intraovarian secreted proteins. In several

goodeid species, previous studies indicated that the ovarian fluids include various

proteins in a pattern similar to that in the blood serum [13,29]. Therefore, several serum

proteins, albumen, transferrin, and others known as Cubam ligands, are potential targets

in the case of intraovarian nutrient absorption via the trophotaenia. Another possibility is

that vitellogenin is also a potential target because it is secreted into the ovarian fluids

and is absorbed into the trophotaenia via intracellular vesicles [15]. Another study

suggested that the Cubn-Amn-Lrp2 receptor complex is related to transport of yolk

proteins, including vitellogenin, in endodermal epithelial cells during chicken yolk sac

growth [30]. These reports support the idea that Cubn and Amn are potent candidate

molecules for a receptor complex involved in maternal component uptake in the

trophotaenia. However, functional analyses such as gene knockout or transgenic

technologies used in conventional model animals, such as laboratory mice or small

oviparous fish, have not been applied to goodeid fish. For a direct validation of our

hypothesis, these reverse genetic methods should also be developed in the viviparous

teleost.

As described above, endocytosis-mediated nutrient absorption is not limited to the

pseudoplacenta of the viviparous fish; it has also been reported in intestinal tissues

(Figure 7). In mammals, macromolecule absorption in the intestine is limited to the

suckling period [31]. In the stomachless fish, endocytosis-mediated absorption is

considered to continue in part of the intestine for life, because of low digestive activity in

the enteric cavity [18]. We found that Cubn and Amn are also distributed in the intestinal

epithelial cells of adult *X. eiseni.* (Figure S5). In most invertebrate species, because the

extracellular digestive system is primitive, food particles are absorbed into the intestinal

cells by vesicle trafficking and degraded by intracellular digestion [32, 33]. Thus, we

hypothesize that endocytic nutrient absorption and intracellular digestion are ancestral

traits common not only in vertebrates but also in invertebrates, and their importance has

decreased in certain vertebrates with development of the extracellular digestive system

in the enteric cavity. The ancestor of the goodeid species may have applied the

endocytic process to the reproductive system, and then configured the unique hind-gut

derived pseudoplacenta for matrotrophic nutrition during gestation [7,34]. To validate our

hypothesis, further exhaustive omics analyses using the goodeid species and

comparative research using the subfamily Empetrichthyinae, which is the oviparous

species most closely related to the viviparous goodeid fish, are required [6,35].

This study is an investigation of species-specific traits based on the transcriptome of a

non-conventional model species. The results revealed potential candidate molecules for

nutrient absorption in the pseudoplacenta of the viviparous teleost. As August Krogh

wrote, this kind of study would be suitable for investigation using the most appropriate

species and is unsuitable for verification using alternative models such as viviparous

rodents or oviparous teleosts. We believe that this study is an important and

fundamental step in understanding the strategic variation of the reproductive system in

vertebrates.

Acknowledgments

This work was supported by research grants from the Nakatsuji Foresight Foundation

and the Daiko Foundation.

Competing interest statement

The authors declare that they have no competing interests.

Data accessibility

397   The data that support the findings have been provided with the manuscript.

References

1. A. Krogh, The progress of physiology. *Am. J. Physiol.* **90**, 243-251 (1929).

2. D. Blackburn, "Viviparity and oviparity: Evolution and reproductive strategies" in: Encyclopedia of Reproduction, Vol. 4, E. Knobil, J. D. Neill, Eds. (Academic Press, 1999), pp. 994-1003.

3. G. J. Burton, A. L. Fowden, Review: The placenta and developmental programming: balancing fetal nutrient demands with maternal resource allocation. *Placenta*. **33**, S23-S27 (2012).

4. R. W. Gill, G. Kossoff, P. S. Warren, W. J. Garrett, Umbilical venous flow in normal and complicated pregnancy. *Ultrasound Med. Biol*. **10**, 349-363 (1984).

5. R. M. Roberts, J. A. Green, L. C. Schulz, The evolution of the placenta. Reproduction (Cambridge, England), **152**, R179–R189 (2016).

6. K. L. Foster, K. R. Piller, Disentangling the drivers of diversification in an imperiled group of freshwater fishes (Cyprinodontiformes: Goodeidae). *BMC Evol. Biol.* **18**, 116 (2018).

7. C. L. Turner, Pericardial sac, trophotaeniae, and alimentary tract in embryos of goodeid fishes. *J. Morphol.* **67**, 271-289 (1940).

8. S. M. Tinguely, A. Lange, C. R. Tyler, Ontogeny and Dynamics of the Gonadal Development, Embryogenesis, and Gestation in *Xenotoca eiseni*

(Cyprinodontiformes, Goodeidae). *Sex Dev*. **13**, 297-310 (2019).

9.   C. L. Turner, Viviparity Superimposed upon Ovo-Viviparity in the Goodeidae, a

Family of Cyprinodont Teleost Fishes of the Mexican Plateau. *J. Morphol.* **55**,

207- 251 (1933).

10. C. L. Turner, The trophotaeniae of the goodeidae, a family of viviparous

cyprinodont fishes. *J. Morphol.* **61**, 495- 523 (1937).

11. J. Lombardi, J. P. Wourms, The trophotaenial placenta of a viviparous goodeid

fish. I. Ultrastructure of the internal ovarian epithelium, the maternal component.

J. Morphol. **184**, 277-292 (1985).

12. F. Hollenberg, J. P. Wourms, Ultrastructure and protein uptake of the embryonic

trophotaeniae of four species of goodeid fishes (Teleostei: Atheriniformes). *J.*

Morphol. **219**, 105-129 (1994).

13. F. Hollenberg, J. P. Wourms, Embryonic growth and maternal nutrient sources in

goodeid fishes (Teleostei: Cyprinodontiformes). *J. Exp. Zool.* **271**, 379–394

(1995).

14. A. Vega-López, E. Ortiz-Ordóñez, E. Uría-Galicia, E. L. Mendoza-Santana, R.

Hernández-Cornejo*, et al.,* The role of vitellogenin during gestation of

Girardinichthys viviparus and *Ameca splendens*; two goodeid fish with

matrotrophic viviparity. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol*. **147**,

731–742 (2007).

15. A. Iida, H. N. Arai, Y. Someya, M. Inokuchi, T. A. Onuma, *et al.*, Mother-to-embryo

vitellogenin transport in a viviparous teleost *Xenotoca eiseni*. *PNAS*, **116**, 22359-

22365 (2019).

16. J. P. Kraehenbuhl, C. Bron, C., B. Sordat, Transfer of humoral secretory and cellular

immunity from mother to offspring. *Curr. Top. Pathol*. **66**, 105–157 (1979).

17. P. C. Moxey, J. S. Trier, Development of villus absorptive cells in the human fetal

small intestine: a morphological and morphometric study. *Anat. Rec*. **195**, 463–

482 (1979).

18. J. H. Rombout, C. H. Lamers, M. H. Helfrich, A. Dekker, J. J. Taverne-Thiele,

Uptake and transport of intact macromolecules in the intestinal epithelium of carp

(*Cyprinus carpio L.*) and the possible immunological implications. *Cell Tissue*

Res. **239**, 519-530 (1985).

19. J. Park, D. S. Levic, K. D. Sumigray, J. Bagwell, O. Eroglu, *et al.*, Lysosome-Rich

Enterocytes Mediate Protein Absorption in the Vertebrate Gut. *Dev. Cell.* **51**, 7-20

(2019).

20. P. J. Verroust, H. Birn, R. Nielsen, R. Kozyraki, E. I. Christensen, The tandem

endocytic receptors megalin and cubilin are important proteins in renal pathology.

Kidney Int. 62, 745-756 (2002).

21. G. A. Pedersen, S. Chakraborty, A. L. Steinhauser, L. M. Traub, M. Madsen, AMN

directs endocytosis of the intrinsic factor-vitamin B(12) receptor cubam by

engaging ARH or Dab2. *Traffic*. **11**, 706-720 (2010).

22. C. Larsen, A. Etzerodt, M. Madsen, K. Skjødt, S. K. Moestrup, C. B. F. Andersen,

Structural assembly of the megadalton-sized receptor for intestinal vitamin B12

uptake and kidney protein reabsorption. *Nat. Commun.* **9**, 5204 (2018).

23. E. I. Christensen, H. Birn, P. Verroust, S. K. Moestrup, Membrane receptors for

endocytosis in the renal proximal tubule. *Int. Rev. Cytol*. **180**, 237-284 (1998).

24. B. D. Grant, J. G. Donaldson, Pathways and mechanisms of endocytic recycling.

Nat. Rev. Mol. Cell Biol. **10**, 597-608 (2009).

25. E. I. Christensen, H. Birn, Megalin and cubilin: multifunctional endocytic

receptors. *Nat. Rev. Mol. Cell Biol.* **4**, 256-266 (2002).

26. M. E. Maurer, J. A. Cooper, Endocytosis of megalin by visceral endoderm cells

requires the Dab2 adaptor protein. *J. Cell Sci*. **118**, 5345-55 (2005).

27. C. V. Finkielstein, D. G. Capelluto, Disabled-2: A modular scaffold protein with

multifaceted functions in signaling. *Bioessays*. **38**, S45-55 (2016).

28. T. M. Williams, M. P. Lisanti, The caveolin proteins. *Genome Biol*. **5**, 214 (2004).

29. J. F. Schindler, Structure and function of placental exchange surfaces in goodeid

fishes (Teleostei: Atheriniformes). *J. Morphol*. **276**, 991-1003 (2015).

30. R. Bauer, J. A. Plieschnig, T. Finkes, B. Riegler, M. Hermann, W. J. Schneider,

The developing chicken yolk sac acquires nutrient transport competence by an

orchestrated differentiation process of its endodermal epithelial cells. *J. Biol.*

Chem. **288**, 1088-1098 (2013).

31. V. Muncan, J. Heijmans, S. D. Krasinski, N. V. Büller, M. E. Wildenberg, *et al.*,

Blimp1 regulates the transition of neonatal to adult intestinal epithelium. *Nat.*

Commun. **2**, 452 (2011).

32. P. V. Fankboner, Digestive System of Invertebrates. In book: eLS (2001).

33. V. Hartenstein, P. Martinez, Phagocytosis in cellular defense and nutrition: a food-

centered approach to the evolution of macrophages. *Cell Tissue Res*. **377**, 527–

547 (2019).

34. M. C. Uribe, H. J. Grier, S. A. Avila-Zúñiga, A. García-Alarcón, Change of

lecithotrophic to matrotrophic nutrition during gestation in the viviparous teleost

Xenotoca eiseni (Goodeidae). *J. Morphol*. **279**, 1336-1345 (2018).

35. R. Van Der Laan, W. N. Eschmeyer, R. Fricke, Family-group names of Recent

fishes. *Zootaxa*. **3882**, 1-230 (2014).

Figure 1. Exploring candidate genes for nutrient absorption

A. A working model and hypothesis of this study. In the goodeid viviparous fish (*X. eiseni*), intraovarian embryo absorbs maternal nutrients via the trophotaenia. We hypothesized that endocytosis-mediated proteolysis is related to nutrient absorption. Based on the scenario, potential target genes are for endocytic receptors, adaptors, vesicle coating proteins, and proteases. **B**. An experimental scheme for the RNA-Seq analysis. RNA samples were obtained from the trophotaeniae (white dotted line) of the single intraovarian embryos extracted from the pregnant females of the 3rd- or 4th-week post-mating. The RNA-Seq was performed using three samples at every stage. **C**. The graphs indicate the transcript per million (TPM) values of the genes selected from the RNA-Seq data that are involved in the endocytosis-mediated proteolysis pathway. Student's *t* test was used for statistical analyses. * $p < 0.05$. ** $p < 0.01$. Error bars represent the standard error.

Figure 2. Structures and amino acid sequences for *X. eiseni* Cubn and Amn

A. The illustration indicates that a typical structure of Cubam receptor complex consists of Cubn and Amn. Both proteins possess a conserved motif to bind each other in the N-terminal regions, and the Amn possesses two adaptor binding motifs in the C-terminal intracellular domain. **B-D**. A comparison of amino acid sequence around the Amn-binding motif in Cubn (**B**), Cubn-binding motif in Amn (**C**), and adaptor binding motifs in Amn (**D**) between *X. eiseni* and three other vertebrate species. The gray filled texts are conserved sequences among the four species.

Figure 3. Phylogenetic comparison of Cubn, Amn, and Cathepsin L amino acid

sequences

Phylogenetic tree indicating the relationship among the amino acid sequences of Cubn

(**A**), Amn (**B**), or Cathepsin L (**C**) in the seven vertebrate species. The sequences of *X.*

eiseni (Cyprinodontiformes: Goodeidae) identified in this study were calculated to be

closed to the sequences of orthologues in *P. reticulata* and *X. maculatus*

(Cyprinodontiformes: Poeciliidae).

Figure 4. Distribution of receptors involved in endocytosis

A. RT-PCR for the candidate genes for the receptors involved in the endocytosis. All

amplicons were detected as single band on the expected sizes based on the transcript

sequences obtained from the *de novo* assembly. **B**. The illustration indicates an internal

structure of the trophotaenia. An epithelium cell layer configures an outermost structure

of the trophotaenia, which contacts the ovarian luminal fluids. The layer consists of an

enterocyte-like cell with microvilli on the apical surface and vesicles in the cytoplasm. **C.**

Electron microscopy (left) and corresponding illustration (right) of an epithelial layer cell

of the trophotaenia. Microvilli are distributed on the apical surface of the cells. Arrows

indicate endocytic vesicles during the invagination phase. Arrowheads indicate

intracellular vesicles after endocytosis. Scale bar: 1 µm. **D**. The structure of the

trophotaeniae stained with HE (left) or fluorescent immunohistochemistry. A fibronectin

antibody was used to visualize the connective tissues inside the epithelial layer. Ep,

epithelial layer. Ct, connective tissue. Scale bar, 25 μm. **E-F**. Immunohistochemistry

using the Cubn antibody or Amn antiserum in the trophotaenia. In both samples, DAB

color development was observed in the epithelial layer. Ep, epithelial layer. Ct,

connective tissue. Scale bar, 50 μm. **G**. Confocal microscopy images of fluorescent

immunohistochemistry using the Cubn antibody in the epithelial layer cells. Yellow

triangles indicate the dotted signals in the epithelial cells of the trophotaenia. Green

double-headed arrow indicates the apical surface defined by the flat signal. Yellow

double-headed arrow indicates the cytoplasmic region including the dotted signals.

Scale bar, 5 μm. **H.** Immunoelectron microscopy using the Cubn antibody in the

epithelial layer cells. Green double-headed arrow indicates the microvilli on the apical

surface. Yellow double-headed arrow indicates the cytoplasmic region including the

dotted signals. Arrows indicate endocytic vesicles in the invagination phase. Arrowheads

indicate intracellular vesicles after endocytosis. Anti-Cubilin signals were observed in

both vesicles. The enlarged image is corresponding to the magenta square focused on

the microvilli. Blue arrows indicate anti-Cubn signals in the microvilli. Scale bar, 1 μm.

Figure 5. Proteolysis activities in trophotaenia

A. The illustrations indicate a typical structure of cathepsin L and a comparison of the protease domains of cathepsin L between *H. sapiens* and *X. eiseni*. The gray filled texts are conserved sequences among the species. The black triangles indicate protease active sites. **B**. RT-PCR for *cstl.1* exhibited the highest TPM value in the RNA-Seq analysis. The amplicons were detected as a single band on the expected size based on the transcript sequences obtained from the *de novo* assembly. **C**. Labeling of acidic organelles including lysosomal vesicle in the trophotaenia. The lysotracker treatment exhibited red fluorescence in the epithelial layer (white dotted line). Ep, epithelial layer. Ct, connective tissue. Scale bar, 20 μm. **D**. High magnification image of the lysotracker-treated epithelial cell layer of the trophotaenia. Yellow dotted line indicates the epithelial layer and white dotted circles are the nuclei of the epithelial cells. The lysotracker fluorescence was observed in the cytoplasm. Scale bar, 10 μm. **E**. Measurement of

cathepsin L activity based on fluorescent substrate degradation. The vertical line

indicates the accumulation of cleaved fluorescent substrates, which means cathepsin L

activity in the sample solution. The fluorescent value of the trophotaenia lysate sample

was increased by 8 h after the reaction started. Student's t test was used for statistical

analyses. ** $p < 0.01$.

Figure 6. Gene expression analysis of the endocytic adaptor and vesicle coating

genes

RT-PCR was performed for the candidate genes encoding the adaptors and vesicle

coating genes involved in endocytosis. The amplicons were detected as single bands of

the expected sizes based on the transcript sequences obtained from the *de novo*

assembly. M, marker.

	Viviparous mammal (*Mus musculus*)	**Stomachless fish** (*Danio rerio*)	**Viviparous fish** (*Xenotoca eiseni*)
Stage	Suckling	Larva ~ Adult	Intraovarian embryo
Tissue	Intestine (Ileum)	Intestine (LREs)	**Trophotaenia**
Factor	*Dab2*	*cubn*, *dab2*	*cubn*, *amn*, *ctsl.1*
Target	Oral food (Milk)	Oral food (Normal diet)	**Maternal supplement**

Figure 7. Comparison of endocytosis-mediated nutrient intake in vertebrates

Endocytosis-mediated nutrient intake has also been reported in a viviparous mammal and stomachless fish. In these species, oral ingestion macromolecules are absorbed from a region of the intestine via endocytosis. In contrast, intraovarian embryos of the viviparous fish (family Goodeidae) absorbs the maternal component in the ovarian fluids from the trophotaeniae via endocytosis. Endocytosis in each species is predicted to be driven by similar molecular process in the enterocytes of the intestine or the enterocyte-like cells of trophotaenia; however, the biological ontology would be divergent between the viviparous fish (intraovarian growth) and the others (feeding after birth).

Gene	Accession #		Primer sequence (5'-3')	Related figures
cubn	LC595284	F	ATGGCTGTTACAGTGCAGCCTTCA	4A
		R	GGCATGTACATACAGGGAATGCTG	
amn	LC595285	F	TGCCCTTTACAAGCAGTGGATTCC	4A
		R	GCGGATTAGCACAACCACAATCAC	
lrp2a	LC595286	F	ATGTGGAGAACACCGCTGCTTCAA	4A
		R	CCACATGGAGCAGTCATCGAAATC	
ap2a1	LC595287	F	GCTGTGTCGAAGGGAGATGGAATG	6
		R	AGAGCTAGATAGCGCAGATTGGTC	
ap2b1	LC595288	F	GTCATTGCTGCCATGACTGTTGGC	6
		R	CCACAATAGCCTCCTGAACCACAT	
ap2m1a	LC595289	F	GACGTAATGACGGCCTACTTTGGC	6
		R	CAGCAGCGGGTTTCATAGATGCCA	
ap2s1	LC595290	F	ATGATCCGCTTCATCCTCATCCAG	6
		R	CTCTAGTGACTGAAGCATGAGGAG	
ldlrap1b	LC595291	F	GGACGCTTTAAAATCCGCTGGAAG	6
		R	GCAGCTGAGTCTGTCTCATCTTGG	
numb	LC595296	F	GAATAAGCTACGGCAGAGCTTCCG	5
		R	TTGGGCATTGGAGCGGAAGAGAAC	
clta	LC595292	F	GGATGATTTTGACATGCTGAACGC	5
		R	TAACGGACCAGCGGGGACTGCTTT	
cltc	LC595293	F	CTCAGATCCTGCCAATTCGCTTTC	6
		R	TCTTCTTCCACGCAAACAGACAGC	
ctsl.1	LC595294	F	CAGCATCTCTCTGGAAGATCTCGA	5B
		R	CAGACTAGTGGATAACTTGCTGCG	
actb	LC595295	F	ATGGAAGATGAAATCGCCGCACTG	4A, 5B
		R	GAAGCATTTACGGTGGACGATGGA	

631 **Table 1. List of primers used in the study.**

632 F, forward, R, reverse.

		3 weeks				4 weeks			
		#1	#2	#3	Average	#1	#2	#3	Average
Receptor	*cubn*	1421	768	1414	**1201**	1528	1464	1250	**1414**
	amn	745	966	1269	**993**	1139	818	1032	**996**
	lrp1aa	6	4	10	**7**	14	10	8	**11**
	lrp2a	12	6	10	**9**	12	7	12	**10**
Adaptor	*ap2a1*	75	60	73	**69**	78	89	82	**83**
	ap2b1	91	54	98	**81**	84	83	85	**84**
	ap2m1a	383	346	360	**363**	349	374	385	**369**
	ap2s1	365	318	368	**350**	359	370	371	**367**
Vesicle	*ldlrapab*	7	6	7	**7**	8	8	6	**7**
	numb	19	15	19	**18**	17	15	13	**15**
	clta	851	680	744	**758**	664	739	817	**740**
	cltb	67	76	73	**72**	97	127	104	**109**
	cltc	337	209	308	**285**	328	366	343	**346**
	cav2	1	0	2	**1**	4	2	3	**3**
	cav3	0	0	0	**0**	0	0	0	**0**
	flot1b	229	196	221	**215**	228	238	234	**233**
	flot2a	91	50	89	**77**	228	238	234	**233**
Protease	*ctsa*	862	691	939	**831**	751	749	720	**740**
	ctsb	474	234	429	**379**	2686	2559	1944	**2396**
	ctsc	1867	1113	1907	**1629**	2273	2320	1884	**2159**
	ctsh	52	31	52	**45**	47	55	48	**50**
	ctsk	11	17	14	**14**	24	24	18	**22**
	ctsla	831	736	939	**835**	960	809	924	**898**
	ctsl.1	12749	8412	11227	**10796**	17795	18334	15906	**17345**
	ctso	3	5	7	**5**	6	7	8	**7**
	ctss2.1	2144	1656	2042	**1947**	2494	2500	2342	**2445**
	ctsz	66	63	63	**64**	530	551	523	**535**

633 **Table 2. The TPM values for endocytic genes in the RNA-seq analysis.**

		1 h	2 h	3 h	4 h	5 h	6 h	7 h	8 h
No lysate	#1	68.8	67.8	67.2	65.8	65.5	65.8	65.6	65.2
	#2	66.5	65.9	64.8	62.7	64.0	64.4	63.8	62.7
	#3	66.8	65.4	69.8	63.1	63.3	62.8	61.8	62.6
	Average	**67.4**	**66.3**	**67.3**	**63.8**	**64.3**	**64.3**	**63.8**	**63.5**
Lysate	#1	260.3	345.0	437.7	530.4	579.7	637.3	699.6	694.1
	#2	264.9	351.6	460.3	551.3	617.0	678.4	727.0	748.2
	#3	256.6	339.2	441.8	522.7	587.1	642.9	692.5	720.8
	Average	**260.6**	**345.3**	**446.6**	**534.8**	**594.6**	**652.9**	**706.3**	**721.0**
Lysate + inhibitor	#1	94.0	96.2	98.0	99.6	101.0	106.8	104.4	101.9
	#2	88.1	88.8	91.1	91.5	93.4	94.0	98.0	98.2
	#3	88.3	88.4	89.4	93.1	94.2	96.4	99.4	101.0
	Average	**90.1**	**91.2**	**92.8**	**94.7**	**96.2**	**99.0**	**100.6**	**100.4**

Table 3. The measured fluorescent signal values indicating cathepsin L-dependent proteolysis.