1   **A Reproducibility Analysis-based Statistical Framework for Residue-Residue**

2   **Evolutionary Coupling Detection**

3   Yunda Si, Yi Zhang and Chengfei Yan[*]

5   School of Physics, Huazhong University of Science and Technology, China

6   Correspondence: chengfeiyan@hust.edu.cn

<p>23   Abstract</p>

Direct coupling analysis (DCA) has been widely used to infer residue-residue

contacts from the multiple sequence alignment (MSA) of homologous sequences.

However, effectively selecting residue pairs for contact prediction according to

the result of DCA is a non-trivial task. In this study, we developed a general

statistical framework for significant evolutionary coupling detection, referred to as

IDR-DCA, which is based on reproducibility analysis of the coupling scores

obtained from DCA on manually created MSA replicates. IDR-DCA was applied to

select residue pairs for contact prediction for monomeric proteins, protein-

protein interactions and monomeric RNAs, in which three different versions of

DCA were applied. We demonstrated that with the application of IDR-DCA, the

residue pairs selected using a universal threshold always yielded stable

performance for contact prediction. Comparing with the application of carefully

tuned coupling score cutoffs, IDR-DCA always showed better performance. The

robustness of IDR-DCA was also supported through the MSA down-sampling

analysis. We further demonstrated the effectiveness of applying constraints

obtained from residue pairs selected by IDR-DCA to assist RNA secondary

structure prediction.

Introduction

Contacting residues in monomeric proteins/RNAs or between interacting proteins/RNAs often show covariance in the process of evolution to maintain the architectures and the interactions of these macromolecules, which allows us to infer the intra- or inter-protein/RNA residue-residue contacts through co-evolutionary analysis[1]. Direct coupling analysis (DCA) is a class of widely used methods for co-evolutionary analysis, which quantifies the direct coupling strength between two residue positions of a biological sequence through global statistical inference using maximum entropy models learned from large alignments of homologous sequences[2]. Comparing with local statistical methods like mutual information (MI) and correlated mutation analysis, DCA is able to disentangle direct couplings from indirect transitive correlations, thus showing much better performance in predicting residue-residue contacts[3,4]. A wide variety of algorithms at different levels of approximation for implementing DCA have been developed in recent years, with the focus being on improving the accuracy of DCA and increasing the computational efficiency. The developed algorithms for DCA include message passing DCA (mpDCA)[3], mean-field DCA (mfDCA)[4] and pseudo-likelihood maximization DCA (plmDCA)[2]. Among these developed algorithms, PlmDCA is currently the most popular algorithm for implementing DCA because of its high accuracy and moderate computational cost, which has been successfully applied to directly acquire contact constraints to assist the prediction of protein/RNA structures, interactions and dynamics[5-14], or been used

to provide major feature components for deep learning-based contact/distance

prediction methods[15–19].

Comparing with so many efforts made on improving DCA algorithms and

applying DCA to obtain structural information from sequence data, relatively less

attention has been made on how to quantify the number of residue pairs with

significant evolutionary couplings and select the predictive residue pairs from the

result of DCA. Generally, residue pairs with higher coupling scores obtained from

DCA tend to have higher probabilities to form contacts. Therefore, in most

previous studies, often a certain number (e.g. top 10 or top L/5, L as the sequence

length) of residue pairs with the highest coupling scores were selected for contact

prediction, or a coupling score cutoff was set empirically to select residue pairs

with coupling scores higher than the cutoff. However, both the number of

predictive residue pairs and the coupling score values are influenced by many

factors including the number and the length of the homologous sequences

forming the MSA, the detailed settings of the DCA algorithm, the functional

characteristics of the macromolecule[6,7]. Therefore, neither applying a "number"

cutoff nor a "coupling score" cutoff is an ideal protocol for selecting predictive

residue pairs from the result of DCA. In a previous work, for predicting residue-

residue contacts between interacting proteins, Ovchinnikov *et al.* first rescaled the

raw coupling scores from Gremlin (a software for implementing plmDCA) with an

empirical model to consider the influence of the number and the length of the

homologous sequences forming the MSA on the coupling scores, then

determined an optimal score cutoff based on the inter-protein residue-residue

contacts in the crystal structure of the 50S ribosome complex[6]. For the same

purpose, Hopf *et al.* did something similar but rescaled the coupling scores from

EVcouplings (another software for implementing plmDCA) with a different

empirical model[7]. Both the two methods achieve success in selecting inter-protein

residue pairs for contact prediction. However, since the parameters of these

empirical models were tuned on only a limited number of cases, whether they are

applicable for more general cases is questionable. Besides, Xu et al. proposed a

statistical approach referred to as inverse finite-size scaling (IFSS) to estimate the

significance of DCA results, which was later applied in epistasis detection of

microbial genomes[20–22]. However, to the best of our knowledge, the effectiveness

of this approach has never been shown in selecting evolutionary coupled residue

pairs. The lack of a general approach for detecting significant evolutionary

couplings from the result of DCA limits the appropriate application of this method.

For example, when applying DCA to infer inter-protein residue pairs with

significant evolutionary couplings to assist the protein-protein interaction

prediction at large scale, without appropriately measuring the coupling

significance, we may introduce false positive couplings or miss significant

couplings.

In this study, we develop a general statistical framework for significant residue-

residue coupling detection. The development of this statistical framework is

inspired by the quality control protocols in functional genomic experiments, in

which often reproducible signals in multiple experimental replicates are

considered as the genuine functional signal[23,24]. Here, given an MSA of

homologous sequences, two MSA (pseudo) replicates are created by randomly

assigning the sequences in the MSA into two groups. DCA is then performed on

both the original full MSA and the two MSA replicates. We assume that the

significant couplings are reproducible from DCA on the two MSA replicates.

Therefore, we perform reproducibility analysis on the coupling scores obtained

from DCA on the two MSA replicates, from which we assign each residue pair an

irreproducible discovery rate (IDR) calculated with the Gaussian copula mixture

modelling described in Li *et al.*[25], with the lower the IDR, the more reproducible

the residue-residue coupling. Then, we create an IDR signal profile for the residue

pairs under consideration, which represents the IDR variation with the ranking of

the residue pairs sorted descendingly according to the coupling scores obtained

from DCA on the full MSA. The residue pairs before the IDR signal profile reaching

a certain threshold are considered to be with significant evolutionary couplings.

This statistical framework, referred to as IDR-DCA, was applied to select residue

pairs for contact prediction for 150 monomeric proteins, 30 protein-protein

interactions and 36 monomeric RNAs, in which the DCA were performed with

three different versions of DCA including EVcouplings[26], Gremlin[5] and CCMpred[27].

The result shows that IDR-DCA can effectively select evolutionary coupled residue

pairs with a universal threshold (IDR cutoff=0.1), for that the numbers of residue

pairs selected by IDR-DCA vary dramatically for cases across the three datasets,

but the accuracies of the selected residue pairs for contact prediction are kept

stable. Comparing with the application of the DCA tool specific coupling score

cutoffs carefully tuned on each dataset to reproduce the overall accuracies of the

residue pairs selected by IDR-DCA, IDR-DCA is always able to select more residue

pairs, and provide effective contact predictions for more cases. We further

evaluated the robustness of IDR-DCA through the MSA downsampling analysis.

The result shows that as the numbers of homologous sequences forming the

MSAs getting smaller and smaller, generally IDR-DCA would select fewer and

fewer residue pairs to keep the accuracy of the selection, but the advantage of

IDR-DCA over the application of coupling score cutoffs are always kept at

different levels of the MSA down-sampling. Besides, we also evaluated IDR-DCA

on CASP14[28] free modeling targets, the result illustrates that IDR-DCA with the

universal cutoff can still achieve reasonable performance, but the previously tuned

coupling score cutoffs totally failed in the residue pair selection. Therefore, IDR-

DCA provides an effective and robust statistical framework for significant

evolutionary coupling detection. Finally, we applied constraints obtained from

residue pairs selected by IDR-DCA to assist RNA secondary structure prediction

as an application example of IDR-DCA. The result shows that the accuracy of RNA

secondary structure prediction can be dramatically increased with the application

of these constraints.

Results and Discussion

1. Overview of IDR-DCA

IDR-DCA includes three major stages: creating pseudo-replicates, performing

reproducibility analysis and detecting significant couplings, which are described

in detail in the following subsections (See Figure 1).

1.1 Creating pseudo-replicates

As it is shown in Figure 1A, given an MSA of homologous sequences, we first

perform DCA on the full MSA, from which we can obtain a coupling score x_i for

each residue pair. The residue pairs are then sorted descendingly according to the

coupling scores, in which the residue pairs with higher rankings (n_i) are more likely

to be with significant evolutionary couplings. After that, the sequences in the MSA

are randomly grouped into two subsets, and we then perform DCA on the two

MSA subsets separately, from which we can obtain a coupling score tuple

$(x_{i,1}, x_{i,2})$ and a ranking tuple $(n_{i,1}, n_{i,2})$ for each residue pair, with $x_{i,1}$, $x_{i,2}$

representing the coupling scores for the residue pair i from the DCA on the two

MSA subsets, and $n_{i,1}$, $n_{i,2}$ representing the rankings of residue pair i sorted

according to the coupling scores descendingly. Since the two MSA subsets can be

considered as (pseudo) biological replicates, the significant couplings are

expected to be reproducible from the DCA on the two MSA replicates. Therefore,

we perform reproducibility analysis on the coupling scores obtained from the

replicated DCA to evaluate the reproducibility of each residue-residue coupling.

1.2 Performing reproducibility analysis

175   The reproducibility of each residue-residue coupling is evaluated by calculating

176   the irreproducible discovery rate (IDR) for each residue-residue coupling with the

Gaussian copula mixture modelling described in Li *et al*[25]. Specifically, we assume

that there are two types of residue pairs (i.e. evolutionary coupled residue pairs

and evolutionary uncoupled residue pairs), for which the observed coupling score

tuples $(x_{i,1}, x_{i,2})$ are generated by a latent variable tuple (unobserved) (z_1, z_2)

following the Gaussian mixture distribution $(\pi_0 h_0(z_1, z_2) + \pi_1 h_1(z_1, z_2))$, with

$$h_0 \sim N\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right) \text{ and } h_1 \sim N\left(\begin{pmatrix} \mu_1 \\ \mu_1 \end{pmatrix}, \begin{pmatrix} \sigma_1^2 & \rho_1 \sigma_1^2 \\ \rho_1 \sigma_1^2 & \sigma_1^2 \end{pmatrix}\right) (\mu_1 > 0, \ \rho_1 > 0) \quad (1).$$

Where h_0 is the uncoupling component and h_1 is the coupling component.

Since evolutionary coupled residue pairs generally have higher and more

reproducible coupling scores, we require $(\mu_1 > 0, \rho_1 > 0)$. Because (z_1, z_2) are

not observable, and the relationship between (z_1, z_2) and the observable

coupling score tuples $(x_{i,1}, x_{i,2})$ is unknown, the association parameters $\theta =$

$(\pi_1, \mu_1, \sigma_1^2, \rho_1)$ of the Gaussian mixture model are determined through

maximizing the likelihood function of corresponding copula mixture model:

$$L(\theta) = \prod_{i=1}^{N} [\pi_0 h_0(G^{-1}(u_{i,1}), \ G^{-1}(u_{i,2})) + \pi_1 h_1(G^{-1}(u_{i,1}), G^{-1}(u_{i,2}))] \quad (2).$$

Where $(u_{i,1}, u_{i,2}) = \left(\frac{n_{i,1}}{N}, \frac{n_{i,2}}{N}\right)$ is the normalized ranking tuple of residue

pair i, with $n_{i,1}, n_{i,2}$ corresponding to the rankings of residue pair i according

to the coupling scores from replicated DCA, and N representing the total number

of residue pairs; $G(z_*) = \frac{\pi_1}{\sigma_1} \Phi(\frac{z_* - \mu_1}{\sigma_1}) + \pi_0 \Phi(z_*)$ is the cumulative marginal

distribution of z_1 and z_2, with z_* representing either z_1 or z_2, and Φ

representing the standard normal cumulative distribution function.

Given a set of parameters θ, the probability that a residue pair with the

normalized ranking tuple $(u_{i,1}, u_{i,2})$ being an evolutionary uncoupled residue pair

199 (local IDR) can be computed as:

200
$$\text{idr}(u_{i,1}, u_{i,2}) = \frac{\pi_0 h_0(G^{-1}(u_{i,1}), G^{-1}(u_{i,2}))}{\sum_{k=0,1} \pi_k h_k(G^{-1}(u_{i,1}), G^{-1}(u_{i,2}))} \quad (3).$$

The local IDR are then converted to (global) IDR for the multiple hypothesis

correction. The IDR of each residue pair represents the reproducibility of the

corresponding residue-residue coupling, with the lower the IDR value, the higher

the reproducibility (See Figure 1B).

1.3.   Detecting significant couplings

The rankings (n_i) and the reproducibilities (IDRs) of residue-residue couplings

are unified for the significant coupling detection. Specifically, we build an IDR

signal profile for all residue pairs under consideration, which represents the IDR

variation with the ranking of the residue pairs sorted descendingly according the

coupling scores obtained from DCA on the full MSA. Generally, the IDR of each

residue-residue coupling increases ($-\log 10(\text{IDR}) \downarrow$) with fluctuation when its

ranking goes down. After smoothing the IDR signal profile using a moving

average filter with a window size 5, the residue pairs before the IDR signal reaches

a specified cutoff are considered to be with significant evolutionary couplings (See

Figure 1C).

2. Detecting significant evolutionary couplings with variable IDR cutoffs

IDR-DCA was used to detect intra-protein residue-residue couplings for the

150 monomeric proteins in the original PSICOV contact prediction dataset[29], inter-

protein residue-residue couplings for 30 protein-protein interactions from

Ovchinnikov *el al.*[6]*,* and intra-RNA residue-residue couplings for 36 monomeric

RNAs from Pucci *et al.*[13], in which the DCA were performed with three widely used

plmDCA-based DCA software including EVcouplings[26], Gremlin[5] and CCMpred[27].

We first applied variable IDR cutoffs to select evolutionary coupled residue pairs.

The percentage of contacting residue pairs in the selected residue pairs was used

to evaluate the accuracy of the selection. Two intra-protein residues were

considered to be in contact if their $C\beta$-$C\beta$ distance ($C\alpha - C\alpha$ distance in the case

of glycine) is smaller than 8Å. For the inter-protein residues, the distance cutoff

was relaxed to 12 Å considering that the inter-protein residues have much lower

contact probability than the intra-protein residues. For the intra-RNA residues, a

contact was defined if their $C1'$-$C1'$ distance is smaller than 12 Å. In Figure 2A-

2C, we show the overall accuracies of the selected residue pairs from each dataset

with the application of variable IDR cutoffs. As we can see from Figure 2A-2C that

independent on the DCA tools, for all the three datasets, the accuracies drop at a

relative slow speed when increasing IDR cutoff until reaching 0.1, and after that

the accuracies drop dramatically. Therefore, 0.1 can be considered as a natural

IDR cutoff for selecting residue pairs for contact prediction when using the IDR-

DCA statistical framework.

For the purpose of comparison, we also selected residue pairs based on the

coupling score values. In Figure 2D-2F, we show the overall accuracies of the

residue pairs selected from each dataset with the application of variable coupling

score cutoffs. As we can see from Figure 2D-2F, the accuracy of the selected

residue pairs varies with the choice of the coupling score cutoff in DCA tool

specific and dataset dependent ways. Therefore, a universal coupling score cutoff

is not applicable for selecting residue pairs for contact prediction. For each dataset,

we actually can set empirical DCA tool specific coupling score cutoffs for the

residue pair selection, with which the selected residue pairs reproduce the

accuracies of the residue pairs selected by IDR-DCA with 0.1 as the IDR cutoff.

Specifically, for the monomeric protein dataset, the coupling score cutoffs for

EVcouplings, Gremlin and CCMpred were set as 0.42, 0.29 and 0.73; for the

protein-protein interaction dataset, the coupling score cutoffs were set as 0.16,

0.09 and 0.24; and for the monomeric RNA dataset, the coupling score cutoffs

were set as 0.29, 0.30 and 0.41. The dramatic variations of the coupling scores

cutoffs between different DCA tools and different datasets show that the coupling

score is not a good metric for the predictive residue pair selection.

It is easy to explain that the obtained coupling score cutoffs are tool

dependent. Since all the three versions of DCA use some sort of regularizations

to avoid the model overfitting, if the parameters of the regularizations are set

differently, or the regularizations are done in different ways, the scales of the

obtained coupling score values will vary. This is also the reason that the standard

practice in the DCA application relies more on the order of the prediction than

numeric values of the coupling scores. Besides, it is also reasonable that different

datasets show different scales of coupling scores, since the different types of

biophysical interactions can have different strengths. For example, the intra-

protein residue-residue interactions are generally stronger and more conserved

than the inter-protein residue-residue interactions, and may also show different

evolutionary characteristics from the RNA residue-residue interactions.

3. The performance of IDR-DCA with a universal IDR cutoff (0.1) on

evolutionary coupled residue pair selection

We analyzed the performance of IDR-DCA on selecting evolutionary coupled

residue pairs with 0.1 as the IDR cutoff. In Figure 3A-3C, we show the accuracies,

the numbers and the corresponding coupling score cutoffs (i.e. the smallest

coupling score) of the selected residue pairs for each case in the three datasets.

As we can see from Figure 3A-3C, the selected residue pairs yield quite stable

accuracies across cases in the three datasets independent on the DCA tools (e.g.

for most of the cases, the accuracies of selected residue pairs are higher than 50%.),

although the numbers of the selected residue pairs vary dramatically. We can also

see that the corresponding coupling score cutoffs of the selected residue pairs

vary dramatically not only between DCA tools, but also across cases in the three

datasets. This further supports that it is not appropriate to apply a universal

coupling score cutoff to select residue pairs for contact prediction.

We further analyzed the distance distribution of non-contacting intra-protein

residue pairs ($C\beta$-$C\beta$ distance $\geq$ 8Å) selected by IDR-DCA from the monomeric

protein dataset. The analysis was focused on the intra-protein residue pairs for

which have the largest sample size. We found that the distances of most of the

non-contacting residue pairs selected by IDR-DCA are actually just slightly larger

than 8Å (e.g. < 12Å), as it is shown in Figure S1A. Therefore, we suspect that

many of these "non-contacting" residue pairs by definition may also be "truly"

evolutionary coupled. Actually several slightly larger distance cutoffs (e.g. 8.5Å,

10Å, 12Å, etc.) had been used to define residue-residue contacts in previous

studies. We also noticed a tiny fraction of the selected residue pairs are in long

distance (e.g. $\geq 12\text{Å}$) in the crystal structure. These residue pairs can be

evolutionary coupled with the long distances caused by protein conformational

changes, as it is shown by Anishchenko et al., in which they found that most of

the evolutionary coupled residue pairs not in repeat proteins are actually in spatial

proximality in at least one biologically relevant conformation[11]. Besides, the

alignment errors in MSA and the approximations made in DCA can also be

responsible for these exceptions.

We also noticed that for several cases in the protein-protein interaction dataset

and the monomeric RNA dataset, IDR-DCA was not able to find any evolutionary

coupled residue pairs. The scatter plot of the coupling scores of all residue pairs

for these cases is shown in Figure S1B, in which the contacting residue pairs are

colored red, and the non-contacting residue pairs are colored black. As we can

see from the plot, for all the cases, very few top-ranked residue pairs based the

coupling scores from DCA are actually in contacts in the 3D crystal structure. This

means that the DCA for these cases actually failed to correctly model the residue-

residue couplings. Therefore, it is encouraging that IDR-DCA is able to avoid

selecting false positive residue pairs from these cases.

4. The performance comparison between the application of IDR-DCA and

coupling score cutoffs on evolutionary coupled residue pair selection

We compared the performance of IDR-DCA on evolutionary coupled residue

pair selection with the application of the DCA tool specific coupling score cutoffs

tuned on each dataset. As we have described in section 2, the coupling score

cutoffs were determined to reproduce the accuracies of the residue pairs selected

by IDR-DCA. As we can see from Figure 4A-4C, for all the three datasets, IDR-

DCA with a universal IDR cutoff (0.1) is always able to select more residue pairs

than the application of the carefully tuned DCA tool specific coupling score cutoffs,

although the accuracies of the selected residue pairs are almost the same.

Besides, IDR-DCA also shows a more stable performance across cases in each

dataset. For example, for most of the cases, the numbers of residue pairs selected

by IDR-DCA are very similar between different DCA tools, but the numbers of

residue pairs selected by applying the coupling score cutoffs are highly dependent

on the choice of the DCA tools (See Figure S2). Since the differences between the

three DCA tools are only on the detailed settings of the plmDCA algorithm (e.g.

the initial values for the optimization, the criterion for the convergence, the ways

of regularizations, etc.), DCA implemented with the three DCA tools on the same

MSA should provide similar number of evolutionary coupled residue pairs.

An effective contact prediction should provide enough residue pairs above a

certain level of accuracy to assist the structure prediction. Here, as rules of thumb,

for monomeric proteins or RNAs, we defined a prediction providing not fewer

than L/5 (L as the sequence length) residue pairs with an accuracy not lower than

50% as an effective contact prediction; for protein-protein interactions, an

effective contact prediction was defined if it can provide at least one residue pairs

with an accuracy not lower than 50%, considering even one inter-protein residue

contact constraint can significantly reduce the configuration space of the protein-

protein interactions. It should be noted that the "effective contact prediction"

defined here is only to make performance comparison between the application of

the universal IDR cutoff and the variable coupling score cutoffs quantitively, thus

other reasonable criteria can also be used in the analysis. In Figure 4D-4F, we

show the comparison of the numbers of cases with effective contact predictions

provided by applying IDR-DCA and by applying the coupling score cutoffs from

the three datasets respectively. As we can see that IDR-DCA is always able to

provide effective contact predictions for much more cases than the application of

the coupling score cutoffs. Besides, we can also see that the performance gap for

the protein-protein interaction dataset is much larger than those for the

monomeric protein dataset and the monomeric RNA dataset. This is mainly

because the coupling scales of DCA methods are also highly dependent on the

sequence length. For the monomeric protein and monomeric RNA dataset, the

variations of the sequence lengths (51~267 and 24~492) are significantly smaller

than that for the protein-protein interaction dataset (181~1453).

5. Evaluating the robustness of IDR-DCA through the MSA downsampling

analysis

We further evaluated the robustness of IDR-DCA through the MSA down-

sampling analysis on the monomeric protein dataset. Specifically, for each protein

in monomeric protein dataset, $\frac{1}{2}$, $\frac{1}{4}$, $\frac{1}{8}$ and $\frac{1}{16}$ of sequences in the original MSA

were randomly selected to form the MSAs with different levels of downsampling,

and then we applied IDR-DCA on the downsampled MSAs with 0.1 as the IDR

cutoff to select evolutionary coupled residue pairs. EVcouplings, Gremlin and

CCMpred were still applied respectively to perform the DCA.

In Figure 5A-5C, we show the accuracies, the numbers of residue pairs selected

by IDR-DCA for each case in the monomeric protein dataset with the application

of the three tools for DCA respectively. As we can see from the Figure 5A-5C, as

the size of MSA getting smaller and smaller, the numbers of the selected residue

pairs are also getting smaller and smaller, however, the accuracies of the selected

residue pairs for contact prediction are kept stable (See Figure S3 for an example).

Since DCA on MSA with fewer sequences tends to have lower statistical power to

accurately model the residue-residue couplings, it is reasonable that IDR-DCA

selected fewer residue pairs as we kept downsampling the MSA.

For the purpose of comparison, we also applied the previous determined

coupling score cutoffs to select residue pairs according to the coupling scores

obtained from DCA on the MSA with different levels of downsampling (See Figure

S4). In Figure 6A-6C, we show the comparison of the numbers and the accuracies

of the residue pairs selected by applying IDR-DCA and by applying the coupling

scores cutoffs for the three DCA tools respectively. As we can see from Figure 6A-

6C, the accuracies of the residue pairs selected by the two approaches are kept

comparable across different levels of the MSA downsampling, however, IDR-DCA

is always able to select more residue pairs. Besides, we also compared the

numbers of proteins with effective contact predictions provided by the two

approaches, which is shown in Figure 6D-6F. The definition of an effective contact

prediction for the monomeric protein is the same as before. As we can see from

Figure 6D-6F, for all the three DCA tools, IDR-DCA is always able to provide

effective contact predictions for more proteins across different levels of the MSA

downsampling.

6. Evaluating IDR-DCA on CASP14 free modelling targets

Recently, Alphafold2 achieved great success in protein structure prediction in

CASP14, in which the co-evolutionary relationships between residues were not

explicitly modelled[30]. Here, we applied IDR-DCA on CASP14 free modeling targets

to estimate the contribution of the co-evolutionary information to the structure

prediction. Specifically, we ran Alphafold2 locally to predict the structures for the

CASP14 free modelling targets with released experimental structures (fifteen

targets in total), from which we obtained the MSA and the predicted models for

each target. After removing the redundant sequences (sequence identity>90%) in

the MSA, we noticed that for three targets (T1033-D1, T1039-D1 and T1043-D1)

the obtained MSAs contain less than five sequences. It is obvious that these

targets are not appropriate for the IDR-DCA statistical framework, which were

therefore excluded from our study. We then applied IDR-DCA on the remaining

twelve targets, in which the three DCA tools (EVcouplings, Gremlin and CCMpred)

were still used for coevolutionary analysis respectively.

In Figure 7, we show the coupling score scattering plot for the twelve CASP14

free modelling targets, in which the contacting residue pairs are colored red and

the non-contacting residue pairs are colored black. We also highlight the

corresponding coupling score cutoff of the residue pairs selected by IDR-DCA

with 0.1 as the IDR cutoff for each target in blue dash. The numbers and the

precisions of the selected residue pairs are summarized in Table 1. As we can see

from the figure and the table, most of these targets (except for T1041-D1 and

T1037-D1) have very few (or even no) contacting residue pairs identified by DCA

(ranked among the top group according the coupling scores), for which IDR-DCA

can appropriately control the quality of the DCA results by selecting very limited

number of (or zero) residue pairs. For these cases, if a "number" cutoff (e.g. L/5)

is used for the residue pair selection, obviously a large number of non-contacting

residue pairs (false positive) will be selected. For T1037-D1 and T1041-D1, DCA

can identify a certain number of contacting residue pairs, as we can see from the

figure that most of these residue pairs can be selected by IDR-DCA. We also

include the TM-score of the top model predicted by Alpafold2 in the last column

of Table 1, a TM-score 0.7 or higher often corresponds a high accurate prediction.

As we can see from the table that Alphafold2 can predict high accurate models

for many targets with very few or even no residue pairs selected by IDR-DCA,

which means that Alphafold2 does go beyond the pure coevolutionary

information. Besides, for the purpose of comparison, the coupling score cutoffs

tuned previous on the PSICOV monomeric protein dataset (EVcouplings: 0.42,

Gremlin: 0.29, CCMpred: 0.73) were also applied for the residue pair selection, and

the number and the precision of the selected residue pairs are summarized in

Table S1. As we can see from the table that for most of the targets, no residue

pair can be selected, and for T1041-D1 and T1037-D1, only a tiny fraction of the

residue pairs identified by DCA can be selected. This illustrates that the pre-tuned

coupling score cutoffs do not work well on these CASP 14 targets at all, which

further supports that IDR-DCA has a much more robust performance on selecting

evolutionary coupled residue pairs.

7. Applying constraints obtained from IDR-DCA to assist RNA secondary

structure prediction

Since deep learning methods like Alphafold2 are now dominant methods for

protein structure prediction, we use RNA secondary structure prediction as an

application example to show the benefit of leveraging IDR-DCA statistical

framework.   Specifically,   the   webserver   2dRNAdca[31]

(http://biophy.hust.edu.cn/new/2dRNAdca/) was applied for the RNA secondary

structure prediction, which first applied a remove-and-expand algorithm to refine

residue pairs selected by IDR-DCA (0.1 as the IDR cutoff) to form the prior

constraints for RNA secondary structure prediction, and then the prior constraints

were further used to guide RNAfold[32] (a minimum free energy based RNA

secondary structure prediction method) to predict the RNA secondary structure.

26 RNAs without broken strands were selected from the RNA dataset for testing

the protocol. Since the result of IDR-DCA is not that dependent on the specific

DCA tools, only the IDR-DCA results based on CCMpred were employed in our

study. Besides, the prediction performances by RNAfold without prior constraints,

with prior constraints refined by the remove-and-expand algorithm from the top

L/5 (L as the sequence length) residue pairs and from residue pairs selected

according to the coupling scores (CCMpred coupling score cutoff: 0.41) were also

evaluated as the references. In Figure 8, we show the Matthews Correlation

Coefficients (MCC) between the experimental RNA secondary structure and the

predicted secondary structures by the four protocols for each of the 26 RNAs (The

RNAs are ordered according to the sequence length ascendingly, also see Table

S2). As we can see from the figure that the introduction of prior constraints by the

three protocols all dramatically improves the prediction performance for most of

the cases. However, the prediction protocol using the constraints refined from the

residue pairs selected by IDR-DCA yields a more stable performance, especially

for large RNAs. We also noticed that for short RNAs (e.g. sequence length<80),

the secondary structure prediction with the three types of prior constraints almost

makes no difference. This is mainly because that for short RNAs, the number of

residue pairs selected by the three approaches are very similar. However, for long

RNAs, IDR-DCA can more effectively select the residue pairs with significant

evolutionary couplings. It should be noted that since the variations of the RNA

sizes and MSA qualities (the numbers of effective sequences for all the MSAs are

larger than 70) of RNA dataset used in our study are not that large, in this case,

selecting residue pairs according to the coupling scores or trivially selecting the

top L/5 residue pairs to some extent can also produce reasonable results, the RNA

secondary prediction using the prior constraints from IDR-DCA only achieved

slightly higher overall accuracies (for RNAs with sequence lengths larger than 100,

the mean MCCs are 0.85, 0.82 and 0.82 for the IDR-DCA constraints, top L/5

constraints and coupling score cutoff constraints respectively). It is reasonable to

expect that performance gaps can be enlarged if they are tested on a more diverse

dataset.

Conclusion

DCA has been widely used to obtain residue-residue contact information to

assist the protein/RNA structure, interaction and dynamics prediction. Besides, the

coupling score matrices obtained from DCA also provide major feature

components for most of the deep learning methods for the protein residue-

residue contact/distance prediction, which has revolutionized the field of protein

structure prediction[17]. Given the MSA of homologous sequences, DCA can be

easily implemented with the state-of-art DCA software to provide the residue-

residue coupling scores. However, it is not easy to quantify the number of residue

pairs with significant evolutionary couplings and select these predictive residue

pairs from the result of DCA, because the number of predictive residue pairs and

the coupling score values from DCA are influenced by many factors including the

number and the length of the homologous sequences forming the MSA, the

detailed settings of the DCA algorithm, the functional characteristics of the

macromolecule, etc.

In this study, we presented a general statistical framework named IDR-DCA for

selecting residue pairs with significant evolutionary couplings. Benchmarked on

datasets of monomeric proteins, protein-protein interactions and monomeric

RNAs, we showed that IDR-DCA can effectively select predictive residue pairs with

a universal IDR cutoff (0.1). Comparing with the application of the DCA tool

specific coupling score cutoffs carefully tuned on each dataset to reproduce the

accuracies of the residue pairs selected by IDR-DCA, IDR-DCA is always able to

select more residue pairs and provide effective contact predictions for more cases.

Therefore, IDR-DCA provides an effective statistical framework for the

evolutionary coupled residue pair detection, which can also be considered as a

general approach for controlling the quality of the result of DCA. Besides, we also

used RNA secondary structure prediction as application example of IDR-DCA. Of

course, IDR-DCA can also be used in other application scenarios. Since the

statistical framework of IDR-DCA is actually not dependent on any detailed

implementation of the DCA algorithm, this statistical framework is also expected

to be applicable to performing quality control for other data-driven contact

prediction methods including deep learning.

Author Contributions

C.Y. designed the study, Y.S., Y.Z., C.Y. conducted the experiments, Y.S., C.Y. wrote

the paper.

Acknowledgements

This work is supported by the startup grant of Huazhong University of Science and

Technology.

Materials and Methods

1.   Preparing the three datasets

1.1 The protein dataset

The PSICOV contact prediction dataset[29] which contains 150 proteins was used

to evaluate the performance of IDR-DCA on detecting intra-protein residue-

residue couplings. The structures of the 150 proteins were obtained from

http://bioinfadmin.cs.ucl.ac.uk/downloads/PSICOV/suppdata/.   The   MSAs   of

homologous proteins for the 150 proteins were built through searching the

whole-genome   sequence   databases   Uniclust30[33]   and   UniRef90[34]   and   the

metagenome database (Metaclust)[35] using DeepMSA[36]. The redundant sequences

with sequence identity higher than 90% in the MSA were removed with HHfilter[37].

1.2 The protein-protein interaction dataset

The   PDB   benchmark   from   Ovchinnikov   *et al.*   was   used   to   evaluate   the

performance of IDR-DCA on detecting inter-protein residue-residue couplings[6].

The complex structures of the 30 protein-protein interactions were downloaded

direct from the Protein Data Bank[38]. The MSAs for protein-protein interologs of

the 30 protein-protein interactions were obtained from the supplementary data

of Ovchinnikov *et al.*[6]. The original PDB benchmark contains 32 protein-protein

interactions, however, 1IXR_B-1IXR_C was removed due to the interacting region

of 1IXR_B was missing in the MSA of protein-protein interologs; and 2Y69_B-

2Y69_C was removed due to the two chains actually do not directly interact with

each other in the crystal structure.

1.3 The RNA dataset

The D^{High} dataset from Pucci *et al.*[13] containing 36 RNAs associated to RNA

families with the number of effective sequences larger than 70 was used to

evaluate the performance of IDR-DCA on detecting intra-RNA residue-residue

couplings. The structures and MSAs of the 36 RNAs were obtained from

https://github.com/KIT-MBS/RNA-dataset. For each MSA, the columns with more

than 50% gaps were first removed, and then the redundant sequences with

sequence identity higher than 95% were removed with HHfilter.

1.4 The CASP14 free modeling targets

The free modeling targets in CASP14 with released experimental structures

(fifteen in total) were further applied to evaluate the performance of IDR-DCA.

First, we ran Alphafold2 locally to generate the MSA and the predicted models for

each target. For the MSA of each target, first the columns with more than 50%

gaps were removed, and then the redundant sequences with sequence identity

higher than 90% were removed with HHfilter. After removing the redundant

sequences in the MSA, we noticed that the MSAs of T1033-D1, T1039-D1 and

T1043-D1 only contained less than five sequences, which were excluded from our

study.

2. Performing the DCA

The three DCA tools: EVcouplings, Gremlin and CCMpred were applied

respectively in this study to perform the DCA. Evcouplings (only the plmc module)

was obtained from https://github.com/debbiemarkslabls/plmc; Gremlin was

obtained from https://github.com/sokrypton/GREMLIN; and CCMpred was

obtained from https://github.com/soedinglab/CCMpred. CCMpred and Gremlin

were run with their default settings, and EVcouplings was run with parameters "-

le 16.0 -lh 0.01 -m 100" for proteins and protein-protein interactions, and with

parameters "-a -ACGU -le 20.0 -lh 0.01 -m 50" for RNAs according to the

recommendations from the website.

3. Performing the reproducibility analysis

The   R   package   'idr'   obtained   from   https://cran.r-

project.org/web/packages/idr/index.html was employed for the reproducibility

analysis with the set of parameters "mu=1.0, sigma=1.0, rho=0.2, p=0.1, eps=1e-

5, max.iter=1000". For the monomeric proteins and RNAs, the residue pairs

separated by less than 6 residues were not considered in the IDR estimation. For

the protein-protein interactions, considering the contact probability of inter-

protein residues is much lower than that of intra-protein residues, the intra- and

inter-protein residue pairs were mixed together for the IDR calculation for the

purpose of better parametrization of the statistical model. However, only the IDRs

of inter-protein residue-residue couplings were used to build the IDR signal

profile for the inter-protein evolutionary coupling detection. For the purpose of

reducing the computational cost, we only perform the reproducibility analysis for

the top 10*L (L as the sequence length) couplings ranked based the coupling

score obtained from the DCA on the full MSA, since the number of evolutionary

coupled residue pairs is generally much smaller than this value.

4. Determining the coupling score cutoffs

For the purpose of comparison, we determined a DCA tool specific coupling

score cutoff on each dataset to reproduce the accuracy of the residue pairs

selected by IDR-DCA with 0.1 as the IDR cutoff. Specifically, for each DCA tool,

starting from 0, we kept increasing the coupling score cutoff for selecting residue

pairs from the corresponding dataset with a step size 0.01, until the accuracy of

the selected residue pairs exceeded the accuracy of the residue pairs selected by

IDR-DCA (0.1 as the IDR cutoff). Then the coupling score cutoff which yielded an

accuracy closest to accuracy of IDR-DCA was chosen as the empirical coupling

score cutoff for this DCA tool on the corresponding dataset.

5. Predicting RNA secondary structure

The 2dRNAdca webserver (http://biophy.hust.edu.cn/new/2dRNAdca/) were

employed to perform the constraints assisted RNA secondary structure prediction.

Ten RNAs with broken strands were removed from the RNA dataset in the

secondary structure prediction. The experimental secondary structure of each

RNA was calculated with X3DNA[28] without pseudoknotted base pairs. The

predicted secondary structures were evaluated by calculating the Matthews

Correlation Coefficient (MCC) between the predicted structure and the

experimental structure, which was calculated using the following formula:

$$\text{MCC} = \frac{TP \times TN - FP \times FN}{\sqrt{(TP + FP)(TP + FN)(TN + FP)(TN + FN)}} (4)$$

Where TP is the number of true positive base pairs; FP is the number of false

positive base pairs; TN is the number of true negative base pairs and FN is the

number of false negative base pairs. We show the secondary structure prediction

result for the RNA 4RYZ_A (chain A in PDB 4RYZ) in Figure S5.

Data availability

The PDB structures of PSICOV protein dataset were obtained from

http://bioinfadmin.cs.ucl.ac.uk/downloads/PSICOV/suppdata/.

The PDB structures of the protein-protein interaction dataset were directly

downloaded from the Protein Data Bank (https://www.rcsb.org/), and the MSAs

of the protein-protein interactions were obtained from

https://elifesciences.org/articles/02030.

The PDB structures and MSAs of the RNA dataset were obtained from

https://github.com/KIT-MBS/RNA-dataset.

Other data generated during and/or analyzed during the current study are

available from the corresponding author on reasonable request.

Code availability

The MSA generation and processing:

DeepMSA was obtained from https://zhanglab.ccmb.med.umich.edu/DeepMSA/.

HHFilter was obtained from https://github.com/soedinglab/hh-suite.

The DCA tools:

EVcouplings was obtained from https://github.com/debbiemarkslab/plmc.

Gremlin was obtained from https://github.com/sokrypton/GREMLIN.

CCMpred was obtained from https://github.com/soedinglab/CCMpred.

The reproducibility analysis:

The IDR package was obtained from

https://cran.rproject.org/web/packages/idr/index.html.

The script for IDR calculation was provided in

https://github.com/ChengfeiYan/IDR-DCA.

Alphafold2:

Alphafold2 was obtained from https://github.com/deepmind/alphafold.

RNA secondary structure prediction:

The webserver 2dRNAdca can be reached through

http://biophy.hust.edu.cn/new/2dRNAdca/.

X3DNA was obtained from https://x3dna.org/.

Bibliography

1. de Juan, D., Pazos, F. & Valencia, A. Emerging methods in protein co-

evolution. *Nat. Rev. Genet.* **14**, 249–261 (2013).

2. Ekeberg, M., Hartonen, T. & Aurell, E. Fast pseudolikelihood maximization

for direct-coupling analysis of protein structure from many homologous

amino-acid sequences. *J. Comput. Phys.* **276**, 341–356 (2014).

3.   Weigt, M., White, R. A., Szurmant, H., Hoch, J. A. & Hwa, T. Identification of

direct residue contacts in protein–protein interaction by message passing.

Proc. Natl. Acad. Sci. **106**, 67 LP – 72 (2009).

4.   Morcos, F. *et al.* Direct-coupling analysis of residue coevolution captures

native contacts across many protein families. *Proc. Natl. Acad. Sci.* **108**,

E1293 LP-E1301 (2011).

5.   Kamisetty, H., Ovchinnikov, S. & Baker, D. Assessing the utility of

coevolution-based residue–residue contact predictions in a sequence-

and structure-rich era. *Proc. Natl. Acad. Sci.* **110**, 15674 LP – 15679 (2013).

6.   Ovchinnikov, S., Kamisetty, H. & Baker, D. Robust and accurate prediction

of residue–residue interactions across protein interfaces using

evolutionary information. *Elife* **3**, e02030 (2014).

7.   Hopf, T. A. *et al.* Sequence co-evolution gives 3D contacts and structures

of protein complexes. *Elife* **3**, e03430 (2014).

8.   Sutto, L., Marsili, S., Valencia, A. & Gervasio, F. L. From residue coevolution

to protein conformational ensembles and functional dynamics. *Proc. Natl.*

Acad. Sci. **112**, 13567 LP – 13572 (2015).

9.   De Leonardis, E. *et al.* Direct-Coupling Analysis of nucleotide coevolution

facilitates RNA secondary and tertiary structure prediction. *Nucleic Acids*

Res. **43**, 10444–10455 (2015).

10. Ovchinnikov, S. *et al.* Large-scale determination of previously unsolved

protein structures using evolutionary information. *Elife* **4**, e09248 (2015).

11.   Anishchenko, I., Ovchinnikov, S., Kamisetty, H. & Baker, D. Origins of

662   coevolution between residues distant in protein 3D structures. *Proc. Natl.*

663   *Acad. Sci.* **114**, 9122 LP – 9127 (2017).

664   12.   Wang, J. *et al.* Optimization of RNA 3D structure prediction using

665   evolutionary restraints of nucleotide–nucleotide interactions from direct

666   coupling analysis. *Nucleic Acids Res.* **45**, 6299–6309 (2017).

667   13.   Pucci, F., Zerihun, M. B., Peter, E. K. & Schug, A. Evaluating DCA-based

668   method performances for RNA contact prediction by a well-curated data

669   set. *RNA* **26**, 794–802 (2020).

670   14.   Cuturello, F., Tiana, G. & Bussi, G. Assessing the accuracy of direct-

671   coupling analysis for RNA contact prediction. *RNA* **26**, 637–647 (2020).

672   15.   Wang, S., Sun, S., Li, Z., Zhang, R. & Xu, J. Accurate De Novo Prediction of

673   Protein Contact Map by Ultra-Deep Learning Model. *PLOS Comput. Biol.*

674   **13**, e1005324 (2017).

675   16.   Zeng, H. *et al.* ComplexContact: A web server for inter-protein contact

676   prediction using deep learning. *Nucleic Acids Res.* (2018)

677   doi:10.1093/nar/gky420.

678   17.   Senior, A. W. *et al.* Improved protein structure prediction using potentials

679   from deep learning. *Nature* **577**, 706–710 (2020).

680   18.   Li, Y. *et al.* Deducing high-accuracy protein contact-maps from a triplet of

681   coevolutionary matrices through deep residual convolutional networks.

682   *bioRxiv* (2020) doi:10.1101/2020.10.05.326140.

19.   Yang, J. *et al.* Improved protein structure prediction using predicted

interresidue orientations. *Proc. Natl. Acad. Sci. U. S. A.* (2020)

doi:10.1073/pnas.1914677117.

20.   Puranen, S. *et al.* SuperDCA for genome-wide epistasis analysis. *Microb.*

genomics **4**, (2018).

21.   Pensar, J. *et al.* Genome-wide epistasis and co-selection study using

mutual information. *Nucleic Acids Res.* **47**, (2019).

22.   Xu, Y., Puranen, S., Corander, J. & Kabashima, Y. Inverse finite-size scaling

for high-dimensional significance analysis. *Phys. Rev. E* **97**, (2018).

23.   Landt, S. G. *et al.* ChIP-seq guidelines and practices of the ENCODE and

modENCODE consortia. *Genome Res.* **22**, 1813–1831 (2012).

24.   Bailey, T. *et al.* Practical guidelines for the comprehensive analysis of ChIP-

seq data. *PLoS Comput Biol* **9**, e1003326 (2013).

25.   Li, Q., Brown, J. B., Huang, H. & Bickel, P. J. Measuring reproducibility of

high-throughput experiments. *Ann. Appl. Stat.* **5**, 1752–1779 (2011).

26.   Hopf, T. A. *et al.* The EVcouplings Python framework for coevolutionary

sequence analysis. *Bioinformatics* **35**, 1582–1584 (2019).

27.   Seemayer, S., Gruber, M. & Söding, J. CCMpred—fast and precise

prediction of protein residue–residue contacts from correlated mutations.

Bioinformatics **30**, 3128–3130 (2014).

28.   Colasanti, A. V., Lu, X. J. & Olson, W. K. Analyzing and building nucleic acid

structures with 3DNA. *J. Vis. Exp.* (2013) doi:10.3791/4401.

29. Jones, D. T., Buchan, D. W. A., Cozzetto, D. & Pontil, M. PSICOV: precise

structural contact prediction using sparse inverse covariance estimation on

large multiple sequence alignments. *Bioinformatics* **28**, 184–190 (2012).

30. Jumper, J. *et al.* Highly accurate protein structure prediction with

AlphaFold. *Nature* (2021) doi:10.1038/s41586-021-03819-2.

31. He, X., Wang, J., Wang, J. & Xiao, Y. Improving RNA secondary structure

prediction using direct coupling analysis. *Chinese Phys. B* **29**, (2020).

32. Hofacker, I. L. RNA secondary structure analysis using the Vienna RNA

package. *Current Protocols in Bioinformatics* (2009)

doi:10.1002/0471250953.bi1202s26.

33. Mirdita, M. *et al.* Uniclust databases of clustered and deeply annotated

protein sequences and alignments. *Nucleic Acids Res.* **45**, D170–D176

(2017).

34. Suzek, B. E., Huang, H., McGarvey, P., Mazumder, R. & Wu, C. H. UniRef:

comprehensive and non-redundant UniProt reference clusters.

Bioinformatics **23**, 1282–1288 (2007).

35. Steinegger, M. & Söding, J. Clustering huge protein sequence sets in linear

time. *Nat. Commun.* **9**, 2542 (2018).

36. Zhang, C., Zheng, W., Mortuza, S. M., Li, Y. & Zhang, Y. DeepMSA:

constructing deep multiple sequence alignment to improve contact

prediction and fold-recognition for distant-homology proteins.

Bioinformatics **36**, 2105–2112 (2020).

37.   Remmert, M., Biegert, A., Hauser, A. & Söding, J. HHblits: lightning-fast

iterative protein sequence searching by HMM-HMM alignment. *Nat.*

Methods **9**, 173–175 (2012).

38.   Berman, H. M. *et al.* The Protein Data Bank. *Nucleic Acids Res.* **28**, 235–

242 (2000).

738
Figure 1. The flowchart of IDR-DCA
(A) Creating pseudo MSA replicates for DCA. (B)Performing reproducibility analysis. (C)
Detecting significant couplings.

Figure 2. The overall accuracies of residue pairs selected from each dataset based on
IDR-DCA and coupling scores with the application of variable IDR and coupling score
cutoffs

(A)-(C) The overall accuracies of residue pairs selected by IDR-DCA with the application of
variable IDR cutoffs from the three datasets: (A) The monomeric protein dataset; (B) The
protein-protein interaction dataset; (C) The monomeric RNA dataset. (D)-(F) The overall
accuracies of residue pairs selected based on coupling scores with the application of variable
coupling score cutoffs from the three datasets: (D) The monomeric protein dataset; (E) The
protein-protein interaction dataset; (F) The monomeric RNA dataset. EVcouplings, Gremlin
and CCMpred were applied to perform the DCA for each case in the three datasets
respectively. The grey vertical dashed lines in (A)-(C) represent the natural IDR cutoff (0.1) for
IDR-DCA. The blue, green and red vertical dashed lines in (D)-(F) represent the empirical
coupling score cutoffs for EVcouplngs, Gremlin and CCMpred respectively, which were tuned
on each dataset to reproduce the accuracies of residue pairs selected by IDR-DCA with 0.1
as the IDR cutoff.

758
Figure 3. The performance of IDR-DCA on evolutionary coupled residue pair selection
with 0.1 as the IDR cutoff
(A)~(C) The accuracies, the numbers and the corresponding coupling score cutoffs of the
residue pairs selected by IDR-DCA with 0.1 as the IDR cutoff for each case in the three datasets:
(A) The monomeric protein dataset; (B) The protein-protein interaction dataset; (C) The
monomeric RNA dataset. For each case, EVcouplings, Gremlin and CCMpred were applied to
perform the DCA respectively. In the case that no residue pair is selected by IDR-DCA, the
corresponding accuracy and the corresponding score cutoff is not shown. For each dataset,
the cases (proteins, protein-protein interactions, RNAs) are ordered ascendingly in the plot
according to their sequence lengths.

Figure 4. The performance comparison between the application of IDR-DCA (0.1 as the IDR cutoff) and the coupling score cutoffs for the evolutionary coupled residue pair selection

(A)-(C) The comparison of the numbers of the residue pairs selected by applying IDR-DCA with 0.1 as the IDR cutoff and by applying the DCA tool specific coupling score cutoffs from the three datasets: (A) The monomeric protein dataset; (B) The protein-protein interaction dataset; (C) The monomeric RNA dataset. The coupling score cutoffs for the EVcouplings, Gremlin and CCMpred were tuned on each dataset respectively to reproduce the accuracies of residue pairs selected by IDR-DCA with 0.1 as the IDR cutoff. (D)-(F) The comparison of the numbers of cases with effective contact predictions provided by applying IDR-DCA with 0.1 as the IDR cutoff and by applying the DCA tool specific coupling score cutoffs for residue pair selection on the three datasets: (D) The monomeric protein dataset; (E) The protein-protein interaction dataset; (F) The monomeric RNA dataset.

Figure 5. The robustness evaluation of IDR-DCA (0.1 as the IDR cutoff) on evolutionary coupled residue pair selection through the MSA downsampling analysis

(A)-(C) The accuracies and the numbers of the residue pairs selected by IDR-DCA (0.1 as the IDR cutoff) for each protein in the monomeric protein dataset, in which the DCA were performed on the MSAs with different levels of downsampling with the application of the three DCA tools: (A) EVcouplings; (B) Gremlin; (C) CCMpred. In the case that no residue pair is selected, the corresponding accuracy is not shown in the plot. The proteins are ordered ascendingly in each plot according to their sequence lengths.

Figure 6. The performance comparison between the application of IDR-DCA (0.1 as the
IDR cutoff) and the coupling score cutoffs for the evolutionary coupled residue pair
selection from the monomeric protein dataset on the MSAs with different levels of
downsampling

(A)-(C) The comparison of the numbers and the accuracies of residue pairs selected by
applying IDR-DCA (0.1 as the IDR cutoff) and by applying the coupling score from the
monomeric protein dataset, in which the DCA were performed on the MSAs with different
levels of downsampling with the three DCA tools: (A) EVcouplings; (B) Gremlin; (C) CCMpred.
(D)-(F) The comparison of the numbers of cases with effective contact predictions provided
by applying IDR-DCA（0.1 as the IDR cutoff）and by applying the coupling score cutoffs for
residue pair selection from the monomeric protein dataset on the MSAs with different levels
of downsampling, in which the DCA were performed with the three DCA tools: (D)
EVcouplings; (E) Gremlin; (F) CCMpred.

Figure 7. The coupling score scattering plot for the twelve CASP14 free modeling targets, in which the contacting residue pairs are colored red, the non-contacting residue pairs are colored black, and the corresponding coupling score cutoff of the residue pairs selected by IDR-DCA for each target is shown in blue dash. EVcouplings (upper), Gremlin (middle) and CCMpred (bottom) were used respectively for direct coupling analysis.

Table 1. The performance of IDR-DCA on CASP14 free modeling targets

Targets	EVcouplings		Gremlin		CCMpred		MSA	Alphafold2
	#[a]	PRE[b]	#	PRE	#	PRE	# of seqs	TM-score[c]
T1027-D1	1	0.	4	0	13	0.15	177	0.60
T1029-D1	0	NA	0	1	1	0	304	0.47
T1031-D1	3	0.67	2	0.50	3	0.33	340	0.86
T1037-D1	7	0.71	11	0.64	20	0.70	745	0.95
T1038-D1	0	NA	0	NA	1	0	29	0.92
T1040-D1	0	NA	0	NA	0	NA	16	0.58
T1041-D1	31	0.81	26	0.81	26	0.81	1029	0.91
T1042-D1	4	0.75	9	0.56	5	0.60	275	0.85
T1049-D1	3	0.33	1	1.0	1	1.0	195	0.93
T1064-D1	0	NA	0	NA	1	0.0	12	0.67
T1074-D1	0	NA	0	NA	0	NA	15	0.93
T1090-D1	3	0	3	0	2	0	356	0.91

Note: a. The number of residue pairs selected by IDR-DCA for each target.

b. The precision of the selected residue pairs for contact prediction

c. The TM-score of the top model predicted by Alphafold2 locally.

Figure 8. The Matthews Correlation Coefficients between the experimental RNA secondary structure and the predicted RNA secondary structures by the four protocols for each of the 26 RNAs. The RNAs are ordered ascendingly according to their sequence lengths.