Article: Spatial integrated models foster complementarity between

monitoring programs in producing large-scale ecological indicators

Abstract (< 300 words)

Over the last decades, large-scale conservation projects have emerged that require

collecting ecological data over broad spatial and temporal coverage. Yet, obtaining relevant

information about large-scale population dynamics from a single monitoring program is

challenging, and often several sources of data, possibly heterogeneous, need to be integrated.

In this context, spatial integrated models combine multiple data types into a single

analysis to quantify population dynamics of a targeted population. Using available information

at different spatial or temporal scales, spatial integrated models have the potential to produce

detailed ecological estimates that would be difficult to obtain if data were analyzed separately.

So far, these models are available for open populations to estimate demographic parameters

(survival, recruitment), therefore requiring data collected in long-term monitoring programs. In

conservation biology however, we often need to quantify population abundance and density in

closed populations.

In this paper, we showcase the implementation of spatial integrated models to closed

populations in a conservation context. We analyzed spatial capture-recapture data together with

distance-sampling data to estimate abundance and density. Focusing on the Mediterranean

bottlenose dolphins (*Tursiops truncatus*) as a case study, we combined 21,464 km of photo-

identification boat surveys collecting spatial capture-recapture data with 24,624 km of aerial

line-transect following a distance-sampling protocol. We compared the performances of the

spatial integrated model, with that of the distance sampling model, and the spatial capture-

recapture model separated. We discussed the benefits of using a spatial integrated model in the

context of the assessment of French Mediterranean bottlenose dolphin conservation status to inform continental scale public policies.

Overall, we emphasize the usefulness of spatial integrated model to make the most of available datasets in a conservation context. Spatial integrated models are widely applicable and relevant to conservation research and biodiversity assessment at large spatial scales.

Introduction

Macro-institutions get increasingly involved in large-scale programs for biodiversity conservation over regional and continental areas. Whether these policies aim at assisting governments (e.g., the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services), or at implementing environmental management such as the European Union directives (Habitat Directive, 92/43/EEC, or Marine Strategy Framework Directive, MSFD, 2008/56/EC), conducting large-scale ecological monitoring is required to establish conservation status of targeted species and ecosystems, and to inform decision-making.

At large spatial scales, logistical and financial constraints often prevent a detailed coverage of the targeted population using a single collection effort, and different monitoring programs often coexist (Lindenmayer & Likens 2010; Zipkin & Saunders 2018; Isaac et al. 2019). The multiplication of monitoring programs over the same conservation context has fostered the development of statistical models that can estimate ecological indicators while accommodating several, possibly heterogeneous, datasets (Besbeas et al. 2002; Miller et al. 2019; Zipkin et al. 2019; Isaac et al. 2019; Farr et al. 2020). Integrating data from several monitoring protocols can give complementary insights on population structure and dynamics (Schaub & Abadi 2011), increase space and time coverage of the population (Schaub & Abadi 2011; Zipkin et al. 2019), and produce more precise estimate of ecological indicators (Isaac et al. 2019; Farr et al. 2020; Lauret et al. 2021).

An important objective of ecological monitoring programs is to estimate population abundance and density (Williams et al. 2002), for which distance sampling (DS, Buckland et al., 2005), and capture-recapture (CR, Williams et al. 2002) methods are widely used. DS and CR methods allows accounting for spatial variation in abundance and density (Camp et al., 2020; Miller et al., 2013; Royle et al., 2014), possibly at large scales (Bischof et al. 2020). Recent modelling tools have emerged to integrate both DS and CR methods into integrated population models (Kéry & Royle 2020). The extension to spatial integrated models (SIMs) has been proposed to account for spatial variation in abundance and demographic parameters while analyzing jointly DS data and SCR data (Chandler et al. 2018). To date however, SIM have never been applied at large spatial scales nor in a conservation context, although we suspect that using SIMs can be promising in these situations. Besides, SIM were developed and used on open populations to estimate temporal variation in population dynamics and vital rates such as survival and recruitment (Chandler & Clark 2014; Chandler et al. 2018; Sun et al. 2019). These applications rely on long-term datasets that are not always compatible with conservation objectives. Conservation biology being a crisis discipline (Soulé 1985), the agenda of biologists and managers is time constrained (Nichols & Williams 2006). In many cases, ecological information is needed, and needed quick (Nichols & Williams 2006; Lindenmayer & Likens 2010). Consequently, ecological inference is often restricted to closed-population indicators (e.g. abundance, density, distribution). When the temporal resolution of monitoring programs does not allow to quantify population dynamics, we argue that closed-population SIM can be an asset in numerous conservation contexts to deal jointly with existing monitoring programs and assess abundance and density.

In this paper, we showcase the use of SIM in a closed-population context demonstrating the benefit of combining DS and SCR to inform conservation policies. We consider a case study on Bottlenose dolphins (*Tursiops truncatus*) that is considered as "vulnerable" by the IUCN

Red List in the North-Western Mediterranean Sea (IUCN 2009). The protected status of

Bottlenose dolphins within the French seas (listed on Annex II of the European Habitats

Directive) led to the development of specific programs to monitor Mediterranean bottlenose

dolphins within the implementation of the European marine strategy framework directive,

which requires assessing the conservation status of this species every 6 years (Authier et al.

2017). We analyzed DS data collected by aerial line-transect surveys over a large area covering

coastal and pelagic seas (Laran et al. 2017). We also analyzed SCR data collected by a photo-

identification monitoring program restricted to coastal waters (Labach et al. 2019). We

compared the abundance and density of bottlenose dolphins estimated from DS model, SCR

model, and SIM to highlight the benefits of the integrated approach in a conservation context.

We discussed the promising opportunities and the conservation implications of using closed-

population SIM to make the best of available datasets.

Methods

Spatial integrated models for closed populations

To integrate DS and SCR data, we used the hierarchical model approach proposed by Chandler

et al. (2018). However, while initially developed for open populations, we built a model for

closed populations to estimate abundance and density (Fig 1). Our SIM is structured around

two layers with i) an ecological model that describes the latent density and spatial variation in

abundance based on an inhomogeneous point process (*Spatial abundance* section below), and

ii) two observation models that describe how the DS and SCR data arise from the latent

ecological model (*Capture-recapture data* and *Distance-sampling data* sections below).

Spatial abundance

For the ecological state model, we use a latent spatial point process modelling the

density of individuals and the overall abundance. Over the study area *S,* an intensity function

returns the expected number of individuals at location *s* in *S*. To account for spatial variation,

we model the latent density surface as an inhomogeneous point process. For every location *s* in

the study area *S,* the expected abundance λ is written as a log-linear function of an

environmental covariate, say habitat:

$$log\big(\lambda(s)\big) = \mu_0 + \mu_1 habitat(s) \tag{1}$$

105   where parameters to be estimated are μ_0 and μ_1 respectively the density asymptote and the

106   regression coefficient of the environmental covariate. Then, the expected abundance is derived

107   by integrating the intensity function over the study area:

$$E(N) = \int_S \lambda(s)ds. \tag{2}$$

The latent state process defined by Eq. 1 is an inhomogeneous point process that is common to

both the SCR and DS models, hence allowing integration and sharing SCR and DS data for

estimating density λ. To account for unseen individuals, we used the data augmentation

technique and augmented the observed datasets to reach M individuals (Royle & Dorazio 2012).

Each individual *i* is considered being ($z_i = 1$) or not ($z_i = 0$) a member of the population

according to a draw in a Bernoulli distribution of probability ψ, with $z_i \sim \text{Bernoulli}(\psi)$ where

ψ is the probability for individual *i* to be a member of the population, with $\psi = E(N)/M$ and

$N = \sum_{i=1}^{M} z_i$.

Capture-recapture data

To accommodate capture-recapture data, we built a SCR model (Royle et al. 2014). Detection

history of individuals were collected over T sampling occasions and captures were recorded at

detectors. We stored observations in a three-dimensional array *y* with y_{ijt} indicating whether

individual i was captured at detector j during sampling occasion t. We assume that observation

y_{ijt} is an outcome from a Bernoulli distribution with capture probability p_{ijt},

$y_{ijt} \sim \text{Bernoulli}(p_{ijt})$. We model capture probability with a half-normal detection function

$p_{ijt} = p_0 exp(-\frac{d_{ij}^2}{2\sigma^2})$ where d_{ij} is the Euclidian distance between the activity center of

individual i and the detector location j, σ is the scale parameter of the half-normal function, and

p_0 is the baseline encounter rate (Royle et al. 2014).

Distance-sampling data

To accommodate distance data, we built a hierarchical DS model (Kéry & Royle 2016). We

model the DS data conditional on the underlying density surface defined by Eqs (1) and (2).

We assume that the probability of detecting individual i in transect l is a decreasing function of

d_{il} the perpendicular distance between transect l and individual i, with $r_{il} = r_0 exp(-\frac{d_{il}^2}{2\eta^2})$,

where η is the scale parameter of the half-normal function, and r_0 is the probability of detection

on the transect, which we will set to 1. Because distance may not be estimated with perfection

by observers, we discretized the distance of observation in B distance bins. We assume that

density within each transect is uniform and that the number of individuals in each transect is

Poisson distributed. This leads to a Poisson model for the DS data in which the expected number

of detections at transect l in distance bin b is given by the expected number of individuals in

bin b multiplied by the average detection probability within each bin b.

Bottlenose dolphins case study

To illustrate the SIM, we focused on an area of 255,000 km^2 covering the North-Western

Mediterranean Sea within which we considered two monitoring programs about bottlenose

dolphins. We used SCR data from at-sea boat surveys over 21,464 km of the French continental

shelf. Observers performed monitoring aboard small boats to locate and photo-identify

bottlenose dolphins all year long between 2013 and 2015. Taking pictures of the dorsal fin of

each individual in the group makes possible the construction of detection history and hence the

analysis of the population through capture-recapture methods (Labach et al. 2019). Boat

surveys were restricted to the coastal waters of France, and is homogeneous in space and time.

We divided the duration of the monitoring programs into 8 equal sampling occasions as in

Labach et al. (2019). We also used DS data that were collected during winter and summer aerial

line-transect surveys covering 24,624 km of both coastal and pelagic NW Mediterranean Sea

between November 2011 to February 2012 and May to August 2012 (Laran et al. 2017). Two

trained observers collected cetacean data following a DS protocol (i.e. recording species

identification, group size, declination angle). Aerial surveys were conditional on a good weather

forecast and aerial sampling effort was homogeneous over the studied area. Although the SCR

and DS datasets were collected during separated time frames, we assumed that the bottlenose

dolphin abundance did not change much between 2011 and 2015 considering a long-lived

species such as bottlenose dolphins (Bearzi et al. 2009).

We divided the study area in 4356 contiguous pixel/sites creating a 5'x5' Mardsen grid

(WGS 84). To model spatial variation in the inhomogeneous point process, we used bathymetry

as an environmental covariate, which is expected to have a positive effect on bottlenose

dolphins' occurrence (Bearzi et al. 2009; Labach et al. 2019). To estimate the sampling effort

of aerial and boat surveys, we calculated the transect length (in km) prospected by each

monitoring protocol within each site during a time period. Sampling effort was therefore site

and occasion-specific in the case of the SCR model, and site specific for the DS model. For the

SCR model, we accounted for spatial and temporal variation in detection probability modeling

the scale parameter σ as a log-linear function of sampling effort E_{jt} at detector j during sampling

occasion t: $\log(\sigma_{jt}) = \beta_0 + \beta_1 E_{jt}$. We also modeled p_0 as a logit-linear function of E_{jt}:

$\text{logit}(p0_{ijt}) = \delta_0 + \delta_1 E_{jt}$. For the DS model, we accounted for spatial variation in the scale

parameter of the detection function modeling η as a log-linear function of sampling effort S_j

and of weather condition W_j in transect j: $log(\eta_j) = \alpha_0 + \alpha_1 S_j + \alpha_2 W_j$.

To highlight to benefit of data integration in estimating spatial density, we compared i)

the output of the spatial DS model, ii) the SCR model, and iii) the SIM.

Bayesian implementation

We ran all models with three Markov Chain Monte Carlo chains with 100,000 iterations

each in the NIMBLE R package (de Valpine et al. 2017). We checked for convergence

calculating the *R-hat* parameter (Gelman et al. 2013) and reported posterior mean and 80%

credible intervals (CI) for each parameter. We considered as important the effect of a regression

parameter whenever the 80% CI of its posterior distribution did not include 0. We also

calculated the predicted density of bottlenose dolphins (i.e. λ). Data and codes are available on

GitHub (*Link to be added after the double-blind peer review process*).

RESULTS

We detected 536 dolphins through aerial surveys clustered in 129 groups. We identified

927 dolphins over 1707 detections in photo-identification surveys, out of which 638 dolphins

were captured only once (68%), 144 were captured twice (15.5%), 149 were captured 3 times

and up to 8 times for one individual. The maximum distance between two sightings of the same

individual was 302 km and 115 km during the same sampling occasion.

We estimated 2450 dolphins (2276; 2631) with SIM over the study area (Table 1), 8470

dolphins (7620; 9329) with the DS model and 1756 dolphins (1645; 1872) with the SCR model

(Table 1). Density asymptotes of SIM (μ_0= -0.67 (-0.75; -0.60)) and SCR model (μ_0= -1.01 (-

1.37; -0.89)) were lower than asymptote of DS model (μ_0= 0.6 (0.50; 0.71)).

In SIM, estimated abundance increased when bathymetry increased ($\mu_1 = 0.43$

(0.37;0.48), Table 1), suggesting a preference for low-depth seafloors (Fig. 2). DS model also

estimated a positive effect of bathymetry ($\mu_1 = 0.34$ (0.28; 0.39), Table 1), while the SCR model

did not detect any effect of bathymetry on density ($\mu_1 = 0.02$ (-0.73; 0.83), Table 1). Then, SIM

and DS models predicted higher densities of bottlenose dolphins in the coastal seas than in the

pelagic seas, whereas the SCR model did not predict any variation in density between coastal

and pelagic waters (Fig. 2).

We detected a positive effect of aerial sampling effort on detection probability in both

DS model ($\alpha_1 = 0.61$ (0.54; 0.67)) and SIM ($\alpha_1 = 0.47$ (0.36; 0.57)). Boat sampling effort

exhibited a positive effect on detection probability for both the SCR model ($\beta_1 = 0.84$ (0.73;

0.96), $\square_1 = 0.76$ (0.71; 0.81)) and the SIM ($\beta_1 = -0.22$ (-1.34; 0.76), $\square_1 = 0.70$ (0.65; 0.75), Table

1). For the SIM and the DS model, the detection probability increased when the weather

condition improved (SIM: $\alpha_2 = 0.30$ (0.18; 0.42), DS: $\alpha_2 = 0.14$ (0.08; 0.20), Table 1).

DISCUSSION

Integrated models make the most of distance sampling and capture-recapture data

With our SIM, we estimated bottlenose dolphin abundance within the range of what was

found in previous studies in nearby areas (Gnone et al. 2011; Lauriano et al. 2014), and found

that densities were more likely to be higher in coastal areas (Bearzi et al. 2009).

A striking result was the large differences in abundance estimates between DS and SCR

models, which were also found in previous studies analyzing the same datasets in isolation.

Using capture-recapture data only, Labach et al. (2019) estimated 2647 dolphins (95%

confidence interval: 2059; 3528) inhabiting the French continental coast where our model

predicted 2450 dolphins (2276; 2631). Analyzing distance sampling data, Laran et al., (2017)

estimated 2946 individuals (95% confidence interval: 796; 11,462) during summer, and 10,233

(95% confidence interval: 4217; 24,861) during winter where our DS model estimated 8470

(7413; 9595) all year long.

We see several reasons that might explain these differences. First, the seasonal difference

in Laran et al. (2017) DS abundance estimates suggests an issue with the geographic closure

assumption that might explain the discrepancy in estimates obtained from SCR and DS models.

Although the Mediterranean bottlenose dolphins population is clustered in coastal sub-units

(Carnabuci et al. 2016), groups can be encountered offshore (Bearzi et al. 2009). In the DS

dataset, large dolphin groups were detected in the pelagic seas at the extreme south of sampling

design (Supporting information). These groups could either be i) occasional pelagic individuals

that belong to coastal populations and that are mainly resident outside our study area (e.g.

Balearic, South-Western Sardinia), or ii) resident pelagic populations that are not sampled by

coastal boat surveys (Louis et al. 2014). Second, DS and SCR models do not estimate the same

quantities. While DS models take a snapshot of abundance and density in the study area at the

moment of the sampling, SCR models estimate abundance and density of the sampled

population, whether or not the individuals are present in the study area during the sampling

period (Calambokidis & Barlow 2004). In our case study, SCR data were restricted to the

French continental coast and did not sample dolphin populations that exist elsewhere in the

study area, e.g. in Corsica, Liguria, and Tuscany (Carnabuci et al. 2016). Despite this

geographic sampling bias in the capture-recapture data, SCR models should predict the

existence of Corsican and Italian populations if the relationship between density and habitat in

Eq (1) was correct and consistent throughout the study area. As the capture-recapture survey

did not sample the lower range of bathymetry, our model underestimated abundance.

Overall, because groups of the Sardinian and Balearic populations and offshore groups can

be sampled in the distance-sampling surveys, the DS model drives upward abundance

compared to the SCR model that is unlikely to account for animals that are members of the

Southern neither the Eastern populations. To perform detailed analysis of the NW

Mediterranean bottlenose dolphin populations, one should consider additional environmental

covariates to better capture spatial variation in density (e.g., sea surface temperature, distance

to coast, or 200m contour, Lambert et al. 2017).

SIM overcomes the limitations of DS and SCR models, and makes the best of each

monitoring program. Using the DS data that were collected in both coastal and pelagic seas, we

informed the slope of the inhomogeneous point process (μ_1), and detected the effect of

bathymetry on density with SIM by correcting for the geographic sampling bias in the SCR

data. On the other hand, the SCR data were more informative (more detections, more

individuals) that the DS data to inform the asymptote of density (μ_0), making the SIM

abundance estimate closer to the SCR model estimate (Table 1).

Conservation implications

To date, the assessment of French Mediterranean bottlenose dolphin population required by

the EU are established using the DS data (Laran et al. 2017). Aerial surveys provide crucial

information on marine megafauna taxa, and on human pressures to fill several criteria of the

Marine Strategy Framework Directive (Laran et al. 2017; Pettex et al. 2017; Lambert et al.

2020). However, in the Mediterranean context, several monitoring programs are available and

SIM makes it possible to include existing datasets that have been discarded so far to inform

public policies (Cheney et al. 2013; Isaac et al. 2019). For bottlenose dolphins, at-sea photo-

identification programs collecting detailed data are an important asset to inform abundance.

Ecological indicators required by the EU directive for bottlenose dolphins would benefit from

integrating aerial line-transect with more data when available (Lauret et al. 2021).

We argue that the bottlenose dolphin is a relevant candidate for a real-life implementation

of the SIM. Between the aerial line-transects repeated every 6 years, many stakeholders

routinely collected detailed data about bottlenose dolphins in the French Mediterranean Sea.

The Marine Protected Areas network is currently developing long-term monitoring programs

to perform bottlenose dolphin photo-identifications, and the French Research Institute for

Exploitation of the Sea (i.e. IFREMER) collected yearly bottlenose dolphins' data during line

transects surveys for pelagic fisheries (Baudrier et al. 2018). Adding complementary long-term

datasets to the aerial-surveys would make possible to access the demographic parameters (*e.g.*

recruitments, survival (Chandler et al. (2018)), which would represent a major opportunity for

the knowledge about French Mediterranean bottlenose dolphin populations and to produce a

reliable EU ecological indicator. The use of SIM for the French Mediterranean bottlenose

dolphin population also opens perspectives to extend the modelling approach exploring

seasonality in density, and to measure immigration and dispersal between bottlenose dolphins

populations (Zipkin & Saunders 2018).

Spatial integrated models as a promising tool for conservation

When establishing species conservation status for large-scale environmental policies,

discarding some datasets from the analysis can reduce the reliability of the ecological estimation

(Bischof et al. 2016). Using multiple datasets into SIM help to overcome some limitations

present when using separated datasets (e.g. limited spatial or temporal survey coverage, Zipkin

& Saunders 2018; Isaac et al. 2019). However, caution should be taken as integrating data

requires additional modelling assumptions (Dupont et al., 2019; Farr et al., 2020; Fletcher et

al., 2019; Simmonds et al., 2020). Overall, SIM are flexible tools that can include more than 2

datasets (Zipkin & Saunders 2018), and various type of data that enlarge the scope of usable

information (presence-absence (Santika et al. 2017), count data (Chandler et al. 2018), citizen

science data (Sun et al. 2019)). Recent and current developments of SCR models open

perspectives to extend the SIM to account for unidentified individuals, or to better describe

animal movement (Milleret et al. 2019; Turek et al. 2020; Jiménez et al. 2020). Over the last

decades, the spatial scope of conservation efforts has greatly increased, and the analytical

methods have had to adapt accordingly (Zipkin & Saunders 2018). Spatial integrated models

are a promising tool that can make the most of available data to inform conservation policies,

especially for wide-range management decisions.

SUPPORTING INFORMATION

Supplementary information about study area, transects, and data is available online (Appendix

S1). The authors are solely responsible for the content and functionality of these materials.

Queries (other than absence of the material) should be directed to the corresponding author.

LITTERATURE CITED

Authier M, Commanducci FD, Genov T, Holcer D, Ridoux V, Salivas M, Santos MB, Spitz J.

2017. Cetacean conservation in the Mediterranean and Black Seas: Fostering

transboundary collaboration through the European Marine Strategy Framework

Directive. Marine Policy **82**:98–103. Available from

https://linkinghub.elsevier.com/retrieve/pii/S0308597X16307321 (accessed November

22, 2018).

Baudrier J, Lefebvre A, Galgani F, Saraux C, Doray M. 2018. Optimising French fisheries

surveys for marine strategy framework directive integrated ecosystem monitoring.

Marine Policy **94**:10–19. Available from

http://linkinghub.elsevier.com/retrieve/pii/S0308597X18301283 (accessed May 22,

2018).

Bearzi G, Fortuna CM, Reeves RR. 2009. Ecology and conservation of common bottlenose

dolphins *Tursiops truncatus* in the Mediterranean Sea. Mammal Review **39**:92–123.

Available from http://doi.wiley.com/10.1111/j.1365-2907.2008.00133.x (accessed November 28, 2018).

Besbeas P, Freeman SN, Morgan BJ, Catchpole EA. 2002. Integrating mark–recapture–recovery and census data to estimate animal abundance and demographic parameters. Biometrics **58**:540–547. Available from http://onlinelibrary.wiley.com/doi/10.1111/j.0006-341X.2002.00540.x/abstract (accessed October 18, 2016).

Bischof R et al. 2020. Estimating and forecasting spatial population dynamics of apex predators using transnational genetic monitoring. Proceedings of the National Academy of Sciences **117**:30531–30538. Available from http://www.pnas.org/lookup/doi/10.1073/pnas.2011383117 (accessed January 4, 2021).

Bischof R, Brøseth H, Gimenez O. 2016. Wildlife in a Politically Divided World: Insularism Inflates Estimates of Brown Bear Abundance: Transboundary wildlife populations. Conservation Letters **9**:122–130. Available from http://doi.wiley.com/10.1111/conl.12183 (accessed December 18, 2020).

Buckland ST, Anderson DR, Burnham KP, Laake JL. 2005. Distance sampling. Encyclopedia of biostatistics **2**. Wiley Online Library.

Calambokidis J, Barlow J. 2004. Abundance of blue and humpback whales in the eastern north pacific estimated by capture-recapture and line-transect methods. Marine Mammal Science **20**:63–85. Available from http://doi.wiley.com/10.1111/j.1748-7692.2004.tb01141.x (accessed December 2, 2020).

Camp RJ, Miller DL, Thomas L, Buckland ST, Kendall SJ. 2020. Using density surface models to estimate spatio-temporal changes in population densities and trend. Ecography. Available from http://doi.wiley.com/10.1111/ecog.04859 (accessed April

10, 2020).

Carnabuci M, Schiavon G, Bellingeri M, Fossa F, Paoli C, Vassallo P, Gnone G. 2016.

Connectivity in the network macrostructure of Tursiops truncatus in the Pelagos

Sanctuary (NW Mediterranean Sea): does landscape matter? Population Ecology

58:249–264. Available from http://doi.wiley.com/10.1007/s10144-016-0540-7

(accessed December 16, 2020).

Chandler RB, Clark JD. 2014. Spatially explicit integrated population models. Methods in

Ecology and Evolution **5**:1351–1360. Available from

http://doi.wiley.com/10.1111/2041-210X.12153 (accessed April 1, 2020).

Chandler RB, Hepinstall-Cymerman J, Merker S, Abernathy-Conners H, Cooper RJ. 2018.

Characterizing spatio-temporal variation in survival and recruitment with integrated

population models. The Auk **135**:409–426. Available from

https://academic.oup.com/auk/article/135/3/409-426/5148935 (accessed April 23,

2019).

Cheney B et al. 2013. Integrating multiple data sources to assess the distribution and

abundance of bottlenose dolphins *Tursiops truncatus* in Scottish waters: Abundance of

bottlenose dolphins around Scotland. Mammal Review **43**:71–88. Available from

http://doi.wiley.com/10.1111/j.1365-2907.2011.00208.x (accessed December 2, 2020).

Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of

wild fauna and flora. 1992. Page 206. Available from

http://data.europa.eu/eli/dir/1992/43/oj/eng (accessed August 21, 2019).

de Valpine P, Turek D, Paciorek CJ, Anderson-Bergman C, Lang DT, Bodik R. 2017.

Programming with models: writing statistical algorithms for general model structures

with NIMBLE. Journal of Computational and Graphical Statistics **26**:403–413. Taylor

& Francis.

Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008

establishing a framework for community action in the field of marine environmental

policy (Marine Strategy Framework Directive) (Text with EEA relevance). 2008. Page

164. Available from http://data.europa.eu/eli/dir/2008/56/oj/eng (accessed August 14,

2019).

Dupont P, Milleret C, Gimenez O, Bischof R. 2019. Population closure and the bias-precision

trade-off in spatial capture–recapture. Methods in Ecology and Evolution **10**:661–672.

Available from https://onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.13158

(accessed December 17, 2020).

Farr MT, Green DS, Holekamp KE, Zipkin EF. 2020. Integrating distance sampling and

presence-only data to estimate species abundance. Ecology. Available from

https://onlinelibrary.wiley.com/doi/10.1002/ecy.3204 (accessed October 15, 2020).

Fletcher RJ, Hefley TJ, Robertson EP, Zuckerberg B, McCleery RA, Dorazio RM. 2019. A

practical guide for combining data to model species distributions. Ecology:e02710.

Available from https://onlinelibrary.wiley.com/doi/abs/10.1002/ecy.2710 (accessed

April 15, 2019).

Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. 2013. Bayesian data

analysis. Chapman and Hall/CRC.

Gnone G et al. 2011. Distribution, abundance, and movements of the bottlenose dolphin

(Tursiops truncatus) in the Pelagos Sanctuary MPA (north-west Mediterranean Sea):

the bottlenose dolphin in the pelagos sanctuary mpa. Aquatic Conservation: Marine

and Freshwater Ecosystems **21**:372–388. Available from

http://doi.wiley.com/10.1002/aqc.1191 (accessed January 28, 2020).

Isaac NJB et al. 2019. Data Integration for Large-Scale Models of Species Distributions.

Trends in Ecology & Evolution:S0169534719302551. Available from

https://linkinghub.elsevier.com/retrieve/pii/S0169534719302551 (accessed November

5, 2019).

IUCN. 2009, November 18. Tursiops truncatus (Mediterranean subpopulation): Bearzi, G.,

Fortuna, C. & Reeves, R.: The IUCN Red List of Threatened Species 2012:

e.T16369383A16369386. International Union for Conservation of Nature. Available

from http://www.iucnredlist.org/details/16369383/0 (accessed October 14, 2017).

Jiménez J, Augustine B, Linden DW, Chandler R, Royle JA. 2020. Spatial capture–recapture

with random thinning for unidentified encounters. Ecology and Evolution:ece3.7091.

Available from https://onlinelibrary.wiley.com/doi/10.1002/ece3.7091 (accessed

December 16, 2020).

Kery M, Royle J. 2020. Applied hierarchical modeling in ecology: analysis of distribution,

abundance and species richness in r and bugs: volume 2: dynamic and advanced

models, 1st edition. Elsevier, Cambridge.

Kéry M, Royle JA. 2016. Applied hierarchical modeling in ecology: analysis of distribution,

abundance and species richness in R and BUGS. Elsevier/AP, Academic Press is an

imprint of Elsevier, Amsterdam ; Boston.

Labach H et al. 2019. Distribution and abundance of bottlenose dolphin (   )

over the French Mediterranean continental shelf. bioRxiv. Available from

http://biorxiv.org/lookup/doi/10.1101/723569 (accessed August 20, 2019).

Lambert C, Authier M, Dorémus G, Laran S, Panigada S, Spitz J, Van Canneyt O, Ridoux V.

2020. Setting the scene for Mediterranean litterscape management: The first basin-

scale quantification and mapping of floating marine debris. Environmental Pollution

263:114430. Available from

http://www.sciencedirect.com/science/article/pii/S0269749120305704 (accessed

January 5, 2021).

Lambert C, Virgili A, Pettex E, Delavenne J, Toison V, Blanck A, Ridoux V. 2017. Habitat

modelling predictions highlight seasonal relevance of Marine Protected Areas for

marine megafauna. Deep Sea Research Part II: Topical Studies in Oceanography

141:262–274. Available from

https://linkinghub.elsevier.com/retrieve/pii/S0967064517300942 (accessed January

27, 2021).

Laran S, Pettex E, Authier M, Blanck A, David L, Dorémus G, Falchetto H, Monestiez P,

Van Canneyt O, Ridoux V. 2017. Seasonal distribution and abundance of cetaceans

within French waters- Part I: The North-Western Mediterranean, including the Pelagos

sanctuary. Deep Sea Research Part II: Topical Studies in Oceanography **141**:20–30.

Available from https://linkinghub.elsevier.com/retrieve/pii/S096706451630409X

(accessed December 27, 2018).

Lauret V, Labach H, Authier M, Gimenez O. 2021. Using single visits into integrated

occupancy models to make the most of existing monitoring programs.

bioRxiv:848663. Cold Spring Harbor Laboratory. Available from

https://www.biorxiv.org/content/10.1101/848663v2 (accessed January 19, 2021).

Lauriano G, Pierantonio N, Donovan G, Panigada S. 2014. Abundance and distribution of

Tursiops truncatus in the Western Mediterranean Sea: An assessment towards the

Marine Strategy Framework Directive requirements. Marine Environmental Research

100:86–93. Available from

https://linkinghub.elsevier.com/retrieve/pii/S0141113614000737 (accessed December

2, 2020).

Lindenmayer DB, Likens GE. 2010. The science and application of ecological monitoring.

Biological Conservation **143**:1317–1328. Available from

http://linkinghub.elsevier.com/retrieve/pii/S0006320710000522 (accessed May 30,

2018).

Louis M, Fontaine MC, Spitz J, Schlund E, Dabin W, Deaville R, Caurant F, Cherel Y,

Guinet C, Simon-Bouhet B. 2014. Ecological opportunities and specializations shaped

genetic divergence in a highly mobile marine top predator. Proceedings of the Royal

Society B: Biological Sciences **281**:20141558. Available from

https://royalsocietypublishing.org/doi/10.1098/rspb.2014.1558 (accessed January 27,

2021).

Miller DAW, Pacifici K, Sanderlin JS, Reich BJ. 2019. The recent past and promising future

for data integration methods to estimate species' distributions. Methods in Ecology

and Evolution **10**:22–37. Available from

https://onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.13110 (accessed

September 18, 2019).

Miller DL, Burt ML, Rexstad EA, Thomas L. 2013. Spatial models for distance sampling

data: recent developments and future directions. Methods in Ecology and Evolution

4:1001–1010. Available from http://doi.wiley.com/10.1111/2041-210X.12105

(accessed April 3, 2020).

Milleret C, Dupont P, Bonenfant C, Brøseth H, Flagstad Ø, Sutherland C, Bischof R. 2019. A

local evaluation of the individual state-space to scale up Bayesian spatial capture–

recapture. Ecology and Evolution **9**:352–363. Available from

https://onlinelibrary.wiley.com/doi/abs/10.1002/ece3.4751 (accessed January 5, 2021).

Nichols, Williams. 2006. Monitoring for conservation. Trends in Ecology & Evolution

21:668–673. Available from

http://linkinghub.elsevier.com/retrieve/pii/S0169534706002680 (accessed March 5,

2018).

Pettex E et al. 2017. Using large scale surveys to investigate seasonal variations in seabird

distribution and abundance. Part I: The North Western Mediterranean Sea. Deep Sea

Research Part II: Topical Studies in Oceanography **141**:74–85. Available from

http://www.sciencedirect.com/science/article/pii/S0967064516303265 (accessed

January 5, 2021).

Royle JA, Chandler RB, Sollmann R, Gardner B, editors. 2014. Spatial capture-recapture.

Elsevier, Amsterdam.

Royle JA, Dorazio RM. 2012. Parameter-expanded data augmentation for Bayesian analysis

of capture–recapture models. Journal of Ornithology **152**:521–537. Available from

http://link.springer.com/10.1007/s10336-010-0619-4 (accessed November 3, 2020).

Santika T et al. 2017. First integrative trend analysis for a great ape species in Borneo.

Scientific Reports **7**:4839. Available from http://www.nature.com/articles/s41598-

017-04435-9 (accessed January 5, 2021).

Schaub M, Abadi F. 2011. Integrated population models: a novel analysis framework for

deeper insights into population dynamics. Journal of Ornithology **152**:227–237.

Available from http://link.springer.com/10.1007/s10336-010-0632-7 (accessed

December 17, 2020).

Simmonds EG, Jarvis SG, Henrys PA, Isaac NJB, O'Hara RB. 2020. Is more data always

better? A simulation study of benefits and limitations of integrated distribution

models. Ecography:ecog.05146. Available from

https://onlinelibrary.wiley.com/doi/abs/10.1111/ecog.05146 (accessed August 9,

2020).

Soulé ME. 1985. What is conservation biology? BioScience **35**:727–734. JSTOR.

Sun CC, Royle JA, Fuller AK. 2019. Incorporating citizen science data in spatially explicit

integrated population models. Ecology **100**. Available from

https://onlinelibrary.wiley.com/doi/abs/10.1002/ecy.2777 (accessed January 4, 2021).

Turek D, Milleret C, Ergon T, Brøseth H, de Valpine P. 2020. Efficient Estimation of Large-

Scale Spatial Capture-Recapture Models. preprint. Ecology. Available from

http://biorxiv.org/lookup/doi/10.1101/2020.05.07.081182 (accessed May 27, 2020).

Williams, Nichols, Conroy. 2002. Analysis and Management of Animal Populations.

Academic Press, San Diego, California, USA. Available from

https://www.elsevier.com/books/analysis-and-management-of-animal-

populations/williams/978-0-12-754406-9 (accessed May 25, 2018).

Zipkin EF, Inouye BD, Beissinger SR. 2019. Innovations in data integration for modeling

populations. Ecology:e02713. Available from

https://onlinelibrary.wiley.com/doi/abs/10.1002/ecy.2713 (accessed April 23, 2020).

Zipkin EF, Saunders SP. 2018. Synthesizing multiple data types for biological conservation

using integrated population models. Biological Conservation **217**:240–250. Available

from https://linkinghub.elsevier.com/retrieve/pii/S0006320717305141 (accessed

January 5, 2021).

TABLES

511
512   Table 1: Parameter estimates for the spatial integrated model (SIM), spatial capture-recapture
513   (SCR) model, and distance-sampling (DS) model. For each parameter, we display the posterior
514   mean and its 80% credible interval (CI).
515

Parameter	SIM		SCR model		DS model	
	Mean	**80% CI**	**Mean**	**80% CI**	**Mean**	**80% CI**
Estimated population size N	2450	2276, 2631	1756	1645, 1872	8470	7620, 9329
Asymptote of density μ_0	-0.67	-0.75, -0.60	-1.01	-1.37, -0.89	0.60	0.50, 0.71
Effect of bathymetry on density μ_1	0.43	0.37, 0.48	0.02	-0.73, 0.83	0.34	0.28, 0.39
SCR scale parameter: Intercept β_0	0.23	-0.82, 1.45	0.04	-1.01, 1.12		
SCR scale parameter: Effect of at-sea sampling-effort β_1	0.84	0.71, 0.95	0.84	0.73, 0.96		
SCR p_0 parameter: Intercept d_0	-6.65	-7.05, -6.27	-6.51	-6.93, -6.11		
SCR p_0 parameter: Effect of at-sea sampling-effort d_1	0.70	0.65, 0.75	0.76	0.71, 0.81		
DS scale parameter: Intercept α_0	-5.52	-6.50, -4.34			-7.91	-8.52, -7.24
DS scale parameter: Effect of aerial sampling-effort α_1	0.47	0.36, 0.57			0.61	0.54, 0.67
DS scale parameter: Effect of weather condition α_2	0.30	0.18, 0.42			0.14	0.08, 0.20

Figure 1: Graphical description of the Spatial Integrated Model (SIM) that combines Spatial
Capture Recapture (SCR), and Distance Sampling (DS). The SIM is a hierarchical model with
three processes: i) latent population size N and density informed by an inhomogeneous point
process, ii) DS observation process that link the line-transect dataset to the latent density
surface, iii) SCR observation process that links the detection histories to the latent density. The
observation process is stochastic according to detection probability. For DS model, the
observed group size is a Binomial draw in the latent abundance N at the sampling location. For
SCR model, observing an individual is a Bernoulli draw with a detection probability p. See text
for notation and more details.

Figure 2: Estimated density surface of bottlenose dolphins (*Tursiops truncatus*) for the 3
models. Lighter color indicates number of individuals per area unit. Both spatial integrated
model (SIM) and distance sampling (DS) predicted higher density in coastal seas, while spatial
capture-recapture (SCR) predicted homogeneous density across the study area. Note that
density scales are different between maps, indicating a higher overall population size for DS
model than for SIM, and SCR model.

539
540
Figure 1: Graphical description of the Spatial Integrated Model (SIM) that combines Spatial
Capture Recapture (SCR), and Distance Sampling (DS). The SIM is a hierarchical model with
three processes: i) latent population size N and density informed by an inhomogeneous point
process, ii) DS observation process that link the line-transect dataset to the latent density
surface, iii) SCR observation process that links the detection histories to the latent density. The
observation process is stochastic according to detection probability. For DS model, the
observed group size is a Binomial draw in the latent abundance N at the sampling location. For
SCR model, observing an individual is a Bernoulli draw with a detection probability p. See text
for notation and more details.
550

551
552
Figure 2: Estimated density surface of bottlenose dolphins (*Tursiops truncatus*) for the 3
models. Lighter color indicates number of individuals per area unit. Both spatial integrated
model (SIM) and distance sampling (DS) predicted higher density in coastal seas, while spatial
capture-recapture (SCR) predicted homogeneous density across the study area. Note that
density scales are different between maps, indicating a higher overall population size for DS
model than for SIM, and SCR model.
559