Sablefish (*Anoplopoma fimbria*) plasma biochemistry and hematology reference

intervals including blood cell morphology

Short title: Sablefish blood

Keywords: Sablefish; Anoplopoma fimbria; reference intervals; blood; plasma

biochemistry; hematology;

Carla B. Schubiger[1*], M. Elena Gorman[1], Jennifer L. Johns[1], Mary R. Arkoosh[2], Joseph

P. Dietrich[2]

[1]Carlson College of Veterinary Medicine, Oregon State University, Corvallis, Oregon,

USA

[2]Environmental & Fisheries Science Division, Northwest Fisheries Science Center,

National Marine Fisheries Service, National Oceanic and Atmospheric Administration,

Newport, Oregon, USA

*Corresponding author:

Email: Carla.schubiger@oregonstate.edu (CBS)

Abstract

Plasma biochemistry and hematology reference intervals are integral health assessment tools in all medical fields, including aquatic animal health. As sablefish (*Anoplopoma fimbria*) are becoming aquaculturally and economically more important, this manuscript provides essential reference intervals (RI) for their plasma biochemistry and hematology along with reference photomicrographs of blood cells in healthy, fasted sablefish. Blood cell counts and morphology can vary between species, precluding the use of RI's from other fish species for use in sablefish. Blood was collected for plasma biochemistry and hematology analysis between June 20 and July 18, 2019, from healthy, yearling sablefish, hatched and reared in captivity on a commercial diet. Overnight fast of 16-18 hours did not sufficiently reduce lipids in the blood, which lead to visible lipemia and frequent rupture of blood cells during analysis. Therefore, sablefish should be fasted for 24 to 36 hours before blood is collected to reduce hematology artifacts or possible reagent interference in plasma biochemistry analysis. RI's may also be influenced by feed, demographics, and production systems, warranting more studies on this topic. In sablefish, lymphocytes are the dominant leukocytes (98%), while eosinophils are rare, and basophils were not detected. Neutrophils are very large cells with Döhle bodies. These bodies are usually signs of toxic changes, but no such association was found in these fish.

Introduction

Sablefish (*Anoplopoma fimbria*), colloquially known as butterfish or black cod, are a highly valuable and fast-growing demersal species native to the continental slope of the

North Pacific Ocean. In 2012, U.S. harvest of wild sablefish was valued at

approximately $141 million (1). Wild sablefish populations have historically declined in

the late 1900s due to massive harvest by international fishing fleets, and as a result, the

catch is highly regulated to allow for currently stable wild populations (1–4). National

and international market demands are projected to continuously increase while harvest

of wild sablefish will likely remain limited, resulting in increased interest in an

aquaculture industry to meet sablefish demand.

Several research efforts and commercial projects are underway to establish sablefish

brood-stock, larval rearing techniques, feeds, as well as the development of net-pen

and landlocked recirculating aquaculture systems (RAS) (5–10). Investing in this

emerging aquaculture industry brings new opportunities to aquaculture communities

and a socioeconomically sensitive approach to satisfy market demands and create jobs

beyond what limited catch permits can achieve (11,12).

High stocking density within any culture system increases disease pressure and is a

ubiquitous challenge for producers and aquatic veterinarians in charge of maintaining

the health status of valuable finfish (13). Research efforts are ongoing to mitigate

disease pressure in these culturing systems (14,15). However, little is known about

normal physiological health parameters of sablefish. This knowledge gap leaves

professionals with insufficient tools to support the health needs of this industry.

This study's objective was to develop diagnostic references for normal ranges

(reference intervals; RI) of plasma biochemistry and hematology values and to

characterize blood cell morphology of cultured sablefish. Producers and aquaculture

healthcare professionals will be able to use these RI as a guideline and a diagnostic

tool.

Material and methods

Mixed-sex fish were obtained from a captive brood-stock and rearing program

conducted by the National Oceanic and Atmospheric Administration's

(NOAA) Northwest Fisheries Science Center (NWFSC) at the Manchester Research

Station (Port Orchard, WA). The fish were reared for one year in land-based tanks with

artificial recirculating seawater at the NWFSC (Seattle, WA) and transferred to NOAA's

Newport Research Station (Newport, OR) on April 18, 2019. At the start of holding, the

fish had an average weight of 186 g and grew to an average weight of 440 g on the final

day of sampling. In Newport, the sablefish were reared in a 2-m diameter fiberglass tank

in ca. 1500 L, with flow-through seawater at ca. 10-14.5 °C. The salinity ranged from 29

to 33.9 ppt and dissolved oxygen ranged from 6.9 to 9.6 mg/L. The indoor tank was lit

from above with a combination of natural (skylights) and artificial lighting. The artificial

lighting was on a daily timer, providing 9 hours of light, turning on after sunrise, and

turning off before sunset. Internal air temperatures ranged from 17.9 to 23.9 °C.

Fish were fed BioOregon's Biobrood (4 mm) diet at least twice per day five days per

week, at approximately 1% of body weight. This diet consists of marine fish oils and

fishmeal protein. The feed's composition, per manufacturer, includes 48% minimum

protein; 20% minimum oil; 8.5% maximum moisture; 1.0% maximum fiber; 11%

maximum ash; 18.2 MJ/kg digestible energy; and 60 ppm astaxanthin (a carotenoid).

The sablefish were visually inspected daily Monday through Friday. There were no

mortalities or signs of illness observed in the sablefish throughout the duration of the

study.

Preparation of reference animals, sample collection and handling

Pre-analytical procedures were consistent with methodologies applicable to health

assessments in aquatic medicine (16). In brief, fish were fasted for 24-36 hours prior to

terminal blood collection to avoid lipemia. Two fish were netted and transferred to 19-L

buckets containing 500 mg/L tricaine methanesulfonate (MS-222) for euthanasia. After

the movement of the operculum stopped, fish were weighed and measured and placed

in dorsoventral orientation. Blood was sampled within 3 minutes of euthanasia from the

caudal vein. 25 gauge (25 G X 5/8), heparinized needles were inserted at the midline

approximately 1-2 cm caudally of the anal fin and advanced craniodorsally at an

approximately 60-degree angle to just ventral of the spine. Approximately two to four

mLs of blood per fish were collected in the heparinized syringes. Immediately after

collection, blood smears were prepared, and non-heparinized microhematocrit capillary

tubes (Fisher Scientific) were filled to determine hematocrit and total plasma protein.

An aliquot of heparinized blood (1.5 mL) was centrifuged at 1200 x g at 14 °C for 10

minutes. Plasma was harvested and transferred to a sterile 2 mL empty tube.

Heparinized whole blood and plasma samples from each fish were stored in a chilled

container and transported within 2-4 hours to Oregon State University Veterinary

Diagnostic Laboratory in Corvallis, Oregon.

Blood was collected from a total of 87 sablefish. However, data from 6 fish (sampled on

June 20 and July 2, 2019) were excluded from analysis due to grossly high lipid content

in the blood that could potentially cause aberrant biochemical and hematological

analyses and skewing statistical results. Therefore, reference intervals were ultimately

determined from a total of 81 sablefish of the same lineage, age, and environmental

conditions. Fish were collected on June 24 (2 fish), 25 (3 fish), 26 (3 fish), and July 5 (1

fish), 10 (32 fish), and 17 (40 fish), 2019. This study was permitted by the animal care

and use committee at Oregon State University (ACUP 5154).

124 Hematology and Biochemistry Methods

Heparinized whole blood and plasma samples were immediately processed upon

receipt by the Veterinary Diagnostic Laboratory at the Oregon State University Carlson

College of Veterinary Medicine in Corvallis, Oregon. The lab evaluated the biochemical

and hematologic parameters.

Plasma biochemical profiles were selected based upon prior studies in other fish

species (17–19). Biochemical analysis was performed via the Beckman Coulter AU480

automated chemistry analyzer (Beckman Coulter Inc., Brea, CA, USA) and reported in

conventional units. The conventional units were then converted to the International

System of Units (SI units) (20). The biochemical parameters analyzed are listed in Table

1.

Table 1. Plasma biochemistry parameters.

Plasma Biochemistry	Description

Parameter & Unit of Measurement	
Albumin (Alb)	Protein synthesized by the liver, important for oncotic pressure in blood and carrier function of hormones, enzymes, vitamins, etc.
Globulins (Glob)	Proteins synthesized in liver and immune system, important in immune response, liver function, blood clotting, etc.
Total Protein (TP)	Albumin and Globulin combined
Glucose	Simple sugar important energy source
Cholesterol	Lipid synthesized in cells, serves as component of cell membranes and precursor of steroid hormones and vitamin D.
Creatinine	Breakdown product from muscle metabolism, excreted by kidneys
Urea nitrogen (BUN)	Waste product produced in liver from protein digestion, excreted by kidneys
Total Bilirubin	Breakdown product of red blood cells, excreted via liver, kidneys, and gastrointestinal tract
Alanine transaminase (ALT)	Formerly called serum glutamate-pyruvate transaminase (SGPT), enzyme present in normal liver and muscle cells
Aspartate Aminotransferase (AST)	Transport enzyme in amino acid metabolism, biomarker for liver health, also known as serum glutamic oxaloacetic transaminase (SGOT)
Creatine Kinase (CK)	Enzyme marker for muscle health
Sodium (Na)	Electrolyte to maintain fluid levels and pH
Potassium (K)	Electrolyte involved in muscle and nerve activity, fluid levels and more
Chloride (Cl)	Electrolyte involved in blood volume and pressure, and pH
Calcium (Ca)	Electrolyte stored in bones, if levels increase indicative of bone, kidney, or thyroid disease and others

Phosphorus (P)	Mineral in bones and teeth, involved in nerve signaling and muscle contractions, abnormal levels indicative of kidney, liver, or bone diseases
Total CO_2	Blood-gas waste product, part of bicarbonate involved in pH level
Anion Gap	Related to electrolyte panel, indicates pH levels

Measurement of all hematologic parameters was performed manually within 24 hours of

receipt. Heparinized whole blood was utilized to assess packed cell volume (PCV), total

protein (TP), and manual cell counting. PCV was obtained by the collection of whole

blood into plain hematocrit tubes and measurement post-5-minute centrifugation. Total

plasma protein was evaluated by a refractometer (Reichert, Depew, NY, USA).

Manual cell counts were performed by two board-certified clinical pathologists (EG and

JJ). Leukocyte and thrombocytes were stained by transferring 20 µL whole blood into

1.98 mL Natt-Herrick Solution (Vetlab supply, Palmetto Bay, FL, USA) for a 1:100

dilution. The mixture was allowed to incubate for 20-30 minutes at room temperature

then counted via Neubauer hemocytometer (Bright-Line, Hausser Scientific, Horsham,

PA) using a standard protocol (17,21,22). Leukocytes were counted in nine primary

squares, and the total was divided by nine and multiplied by 100 (dilution factor).

Thrombocyte counts were obtained by counting thrombocytes in the center main-square

of the hemocytometer then multiplying by 1000.

Blood smears were submitted on glass slides and stained with Modified Wright's stain

(Thermo Fisher Scientific) via an automated method (HemaTek 2000, Siemens

Healthcare, Erlangen, Germany). Leukocytes were identified based upon characteristics

described for other related marine species (23,24), and morphologic identity was initially

agreed upon by both pathologists (EG and JJ). Leukocyte differential counts were

performed by an individual pathologist (EG) to minimize inter-observer variability.

Reference interval calculations

Blood values were collected into Excel spreadsheets and reference intervals calculated

with the macro add-in Reference Value Advisor V 2.1 software (25). The software

follows recommendations and guidelines by the Clinical and Laboratory Standards

Institute (CLSI), the National Committee for Clinical Laboratory Standards (NCCLS),

and the International Federation of Clinical Chemistry (IFCC). The software determined

that the sample size was large enough to compute nonparametric reference intervals

and also determined the 90% confidence intervals of the limits of the nonparametric

reference intervals by a bootstrap method (25,26).

Results

Plasma biochemistry reference intervals

The reference intervals for measured plasma biochemistry parameters in the 81

sufficiently fasted sablefish is presented in Table 2. Overnight fasting of 16-18 hours

was not sufficient to reduce visible lipemia of the blood in three of the sablefish sampled

on June 20, 2019 (animals #32, #33, #34). Also, an unscheduled blood collection was

performed on July 2, 2019, with fish that were fasted for only 16-18 hours leading again

to grossly lipemic samples (animals #47, #48, #49). Consequently, the protocol was

adjusted to increase the fast time to 24-36 hours. Therefore, six of the 87 animals were

excluded from the reference interval calculations due to the severely lipemic condition of

the plasma samples. This condition displayed as visible turbidity in some samples (S1

Fig) and severe blood cell rupture in others.

Table 2. Reference intervals for plasma biochemistry parameters of fasted

sablefish.

Parameter; n = 81	SI units	Conventional units
Albumin (Alb)	6.1 – 13 g/L	0.61 – 1.3 g/dL
Globulin (Glob)	19.1 – 31.9 g/L	1.91 – 3.19 g/dL
Total Protein (TP)	25.2 – 44.9 g/L	2.52 – 4.49 g/dL
Glucose	0.95 – 4.91 mmol/L	17.1 – 88.4 mg/dL
Cholesterol	6.34 – 12.48 mmol/L	245.2 – 482.5 mg/dL
Creatinine	0 – 140.46 µmol/L	0 – 1.59 mg/dL
Urea nitrogen (BUN)	0.07 – 0.14 mmol/L	2.0 – 4.0 mg/dL
Total Bilirubin	1.71 – 3.42 µmol/L	0.1 – 0.20 mg/dL
Alanine aminotransferase (ALT, SGPT)	0 – 17.8 U/L	0 – 17.8 U/L
Aspartate aminotransferase (AST, SGOT)	0 – 1.88 µKat/L	0 – 110.4 U/L
Creatine kinase (CK)	0.09 – 1.49 µKat/L	5.1 – 87.4 U/L
Sodium (Na)	168.0 – 179.9 mmol/L	168.0 – 179.9 mEq/L
Potassium (K)	1.2 – 3.86 mmol/L	1.2 – 3.86 mEq/L
Chloride (Cl)	142.0 – 153.0 mmol/L	142.0 – 153.0 mEq/L
Calcium (Ca)	2.63 – 3.35 mmol/L	10.51 – 13.38 mg/dL
Phosphorus (P)	4.17 – 8.41 mmol/L	12.92 – 26.06 mg/dL
Total CO_2	4.64 – 8.70 mmol/L	4.64 – 8.70 mEq/L
Anion Gap	18.1 – 27.0 mmol/L	18.1 –27.0 mEq/L

185   Table 2 legend: Table 2 lists reference intervals for plasma biochemistry parameters of

186   sablefish that were fasted 36 to 48 hours. Data were collected in conventional units and

187   converted into SI units according to (20).

Hematology Reference Intervals

Hematologic parameters, including packed cell volume (PCV), plasma protein (PP), and

cell counts, were collected manually from the 72 fish sampled on July 10 and 17, 2019.

Thrombocyte counts were only available from 60 animals due to early technical

difficulties with staining. All data are presented in Table 3.

Table 3. Cell blood counts (CBC) were available from 72 animals, with the

exception that thrombocyte counts were only available from 60 animals.

Parameter	n = 72 (*n = 60)
Packed Cell Volume (PCV)	35.7 – 61.8 %
Plasma Protein (PP)	4.3 – 7.92 g/dL
Thrombocytes *	92,100 – 357,475 /µL
White Blood Cell (WBC)	4,333 – 33,083 /µL
Neutrophils	1.8 – 23.4 /µL
Lymphocytes	66 – 96 /µL
Monocytes	0 – 12.1 /µL
Eosinophils	0 - 2.2 /µL
Basophils	0 /µL

Table 3 legend: PCV, PP, and blood cell counts were manually performed from fish

sampled on July 10 and 17, 2019. Thrombocyte counts were manually performed from

60 fish sampled on July 10 and 17, 2019. Data present reference intervals.

Morphology of blood cells was similar to those noted in other fish species (23,24,27).

Erythrocytes are oval with a round to ovoid nuclei (Fig 1a,b,c). Polychromatic

erythrocytes were commonly found in sablefish, which indicates an active turnover of

erythrocytes (Fig 1a). Additionally, fish have a thin barrier between hematopoietic tissue

and the vasculature; hence, immature erythroid cells were common and may comprise

more than 10% of all erythrocytes in circulation (23,27) (Fig 1c). Immature erythrocytes

are smaller and more round than mature erythrocytes with basophilic cytoplasm (Fig

1a). Mitotic figures were not readily identified in blood smears of this population.

Fig 1a. Lymphocytes: Granular lymphocytes. Granular lymphocyte (GL), surrounded

by rounder, basophilic immature erythrocytes (IE) that are smaller than the more mature

erythrocytes (EC). Overall erythrocytes are in various stages of maturity contributing to

a distinct polychromatic appearance, ranging from slightly eosinophilic to basophilic

(Wright's stain x100 objective).

Fig 1b. Lymphocytes: Mature lymphocyte and monocyte. Mature lymphocyte (L)

and monocyte (M) with typical large oval eccentrically-positioned nucleus and a

moderate amount of basophilic cytoplasm that contains three discrete cytoplasmic

vacuoles. Mature lymphocytes are characterizes as round cells with large, round

nucleus with condensed chromatin and scant basophilic cytoplasm. Lymphocytes often

display blebbing of the cell borders (Wright's stain x100 objective).

Fig 1c. Lymphocytes: Reactive lymphocyte. Reactive lymphocyte (with Golgi body

left to the nucleus, RL), small lymphocyte (L) with blebbing and neutrophil (N). Reactive

lymphocytes are large cells, with a more open chromatin pattern and increased

amounts of basophilic cytoplasm (Wright's stain x100 objective).

In sablefish, lymphocytes predominated and comprised up to 98% of the total

leukocytes present in healthy individuals. Lymphocyte morphology was similar to that of

other species, and lymphocytes were identified as round cells with a high nucleus to

cytoplasm ratios (N:C ratios), round nuclei and condensed chromatin (Fig 1a-c). Cell

borders of lymphocytes were frequently blebbed (bulging plasma membrane) (Fig 1c).

Reactive lymphocytes were also similar to those in other species, with cells being larger

than mature lymphocytes, more open chromatin patterns, and increased basophilic

cytoplasm. Occasional Golgi bodies were noted (Fig 1c). Additionally, granular

lymphocytes were occasionally found (Fig 1a). Neutrophils were one of the largest

leukocytes identified in sablefish (Fig 1c, 2, 3). Neutrophils were oval with oval, clefted,

or segmented nuclei surrounded by moderate to abundant amounts of slightly

eosinophilic cytoplasm. Most neutrophils contained pale blue cytoplasmic inclusions

consistent with Döhle bodies. These bodies are often signs of toxic granulation,

inflammation or bacterial infection, but appeared to have no association with other toxic

changes in these fish (Fig 1c, 2, 3) (28–30).

Fig 2. Neutrophils. Neutrophils (N) are large leukocytes with large oval, clefted, or

segmented nuclei and moderate to large amounts of slightly eosinophilic cytoplasm that

often contains pale blue cytoplasmic inclusions consistent with Döhle bodies (Wright's

stain x100 objective).

Fig 3. Neutrophil (N) and Monocyte (M). Note similar size between these cells. Mature

erythrocytes (EC) also identified (Wright's stain x100 objective).

Monocytes were larger than lymphocytes and similar in appearance to monocytes in

other non-mammalian species (Fig 1b, 3, 4). Monocytes contained oval eccentrically-

positioned nuclei; amoeboid nuclear shapes were less frequent. Monocytes had small to

moderate amounts of variably basophilic cytoplasm and frequently contained discrete

cytoplasmic vacuoles (Fig 1b, 3, 4). Cell borders were distinct and non-blebbed, making

them easily distinguishable from lymphocytes. Eosinophils were rare in healthy

sablefish. These were small cells with a high N:C ratio (Fig 5). Nuclei of eosinophils

were oval and typically eccentric, and cytoplasm contained small, bright eosinophilic

granules (Fig 5). Basophils were not found. Thrombocytes were found individually and

in small aggregates (Fig 4, 6). Thrombocytes were small and typically oval to elongate

in shape. Few round forms were also noted. Nuclei of thrombocytes were pleomorphic,

ranging from round to oval to reniform (Fig 4, 6). The cytoplasm was clear with small

cytoplasmic granules occasionally seen.

Fig 4. Monocytes. Monocyte (M) contains several, variably sized, discrete cytoplasmic

vacuoles and distinct, smooth cell borders, in contrast to the two lymphocytes (L) that

display distinct blebbing. Also present is a thrombocyte (T) with a bi-lobbed nucleus

(Wright's stain x100 objective).

Fig 5. Eosinophils. Eosinophils (E) are slightly larger than mature lymphocytes (L) and

have large, ovoid, eccentric nuclei and small amounts of cytoplasm that contains small,

bright eosinophilic granules (Wright's stain x100 objective).

Fig 6. Thrombocytes (T). Thrombocytes are found individually and in small aggregates

in well-prepared blood smears. Thrombocytes are small and typically oval to elongate

and commonly displaying nuclear pleomorphism with round to oval to reniform nuclei.

The cytoplasm of thrombocytes is usually clear with occasional, small cytoplasmic

granules (Wright's stain x100 objective).

Discussion

This study addresses an urgent need associated with the development of a sustainable

and highly profitable emerging aquaculture species. Establishing a sablefish

aquaculture industry would contribute to the livelihoods of coastal (and inland)

communities along the northern Pacific Ocean. Species-specific biochemistry and

hematologic reference intervals (RI) derived from a healthy reference population are

crucial for adequate health monitoring of any species. Particularly crucial is the health

monitoring of species new to culturing. Management and feeding practices, and disease

treatment regimens can be appropriately adjusted in correlation to physiological

parameters of the fish determined from health monitoring. Blood cell counts are

valuable tools in assessing the general health status of the fish and can alert a health

care provider to infections and environmental or management issues, e.g., problems

with water quality or nutrition.

An important observation was that fasting for 24-36 hours was necessary to obtain

blood samples with lipid content low enough to not interfere with plasma biochemistry or

hematology analysis. High amounts of triglycerides in human blood has been found to

significantly interfere with measurements of phosphorus, calcium, creatinine, and total

protein due to an interaction with the analyzer and reagents (31,32). Plasma protein

measured via a refractometer was predictably increased relative to plasma total protein

measured by an automated method as lipids refract light and can falsely increase

refractometric measurement of protein. High lipid content is also suspected of causing

premature lysis of blood cells (33). If fasting is not possible, lipid removal by

centrifugation or lipid extraction with polar solvents is possible (34,35). The Clinical and

Laboratory Standards Institute (CLSI) recommends ultracentrifugation, but this

procedure is not always a practical option (36). Alternatives that might be more practical

for onsite blood diagnostics are dilution of the serum sample or usage of serum blanks

(35). However, if blood samples are necessary for diagnostics in disease events, versus

routine health checks, lipemic blood samples might be less common as inappetence is

often the first clinical sign of problems in a culturing system. For scheduled health

checks, extended fasting of at least 24 hours is recommended for sablefish based on

the results presented in this study. For our analysis, we excluded samples with grossly

visible lipemia (milky, turbid appearance), because of possible interference with

biochemistry values (animals #32-34, sampled on June 20) and blood samples with

moderate cell rupture that was likely caused by lipemia (animals #47-49, sampled on

July 2). To present how these exclusion criteria influenced the plasma biochemistry

intervals, the complete data set of all 87 fish is presented as supplemental material (S1

Appendix). The exclusion of these insufficiently fasted fish (16-18 hours) notably

widened the glucose range, but decreased the upper RI limits of cholesterol, creatinine,

bilirubin, and the anion gap. Changes to other parameters were smaller and often in the

decimals. A comparison of all values is presented as supplemental material (S1

Appendix).

The fish used in this study received a commercial diet at 1% of their body weight

distributed over at least two feeding events per day. Recent studies recommend feeding

sablefish to satiation once every other day to allow for adequate feed conversion and

consistent intake of the feed (37,38). Reduced feed intake has been observed after four

days of daily feeding, attributed to a prolonged gastrointestinal passage in this species

(37) (and personal communication with Ron Johnson). Likely that the delayed clearance

of blood lipids, experienced in this study, is a reflection of that observation.

A curious question is whether the observed affinity for high blood lipid concentration

reflects the general tender consistency of the meat in this species, which gave rise to

the colloquial name "butterfish". However, the energy density of sablefish is relatively

low compared with other pelagic fish from the North Pacific, and only low to moderate

amounts of the wet mass are attributed to lipids (39). A muscle lipid content of up to

14% has been reported in sablefish with higher amounts of lipids stored in the bones

(39–43). Nevertheless, sablefish seem to have a well-adapted fat metabolism. Earlier

studies have shown that prolonged starvation of five months in a laboratory setting did

not result in significant protein catabolism. Instead, fish survived by utilizing their fat

storage over muscle tissue, making sablefish generally well adapted to infrequent meals

in the wild (41,44). Sablefish are identified as opportunistic feeders, whose food sources

can alter by location and age (45). Older and spawning fish commonly reside in deeper

water, while the larvae hatch in the deep and then migrate up, and younger fish are

more commonly found in shallower waters (10,45,46). Besides the ability to travel very

long distances (an average of 191 km per year (47)), sablefish can also undergo an

impressive diel vertical migration in pursuit of some of their vertically migrating prey by

ascending to the surface at night and descending to waters of 750 m or greater before

sunlight (45). This exposes the animal to stark differences in temperature and pressure.

More detailed reasons for these migrations and the intricacy of sablefish lipid physiology

are still under further investigation.

Animal husbandry and management can have a substantial impact on blood reference

ranges. In a study of blood RIs in sunshine bass, the only biochemical reference

intervals that were similar in three different holding systems were glucose and cortisol

(48). Similarly, previous studies in sablefish describe that tank design and water

temperature can influence survival, anatomical deformities, meat quality, and ovarian

development (5–7). Moreover, studies found that blood values in sturgeon can differ

among different hatcheries (19). Stress is reported to alter the fish's ability for

osmoregulation, which will affect electrolyte concentrations in the blood, often resulting

in chloride loss and high glucose values (48,49). Fish sampled in this study had a wide

range of glucose levels (0.95–4.91 mmol/L, 17.1-88.4 mg/dL). This was mostly driven

by noticeably higher glucose levels in the last ten of 32 fish and 40 fish sampled on July

10 and July 17, respectively (S1 Appendix). Most of the glucose measurements were in

the upper teens to mid-forties mg/dl (approximately 1-2.5 mmol/L). The later upward

trend of glucose levels may be due to the repeated netting of fish from the holding tank

during the two to three hours of sampling. Chloride (142-153 mmol/L, 142-153 mEq/L)

was below what has been suggested as normal for marine teleosts (165-175 mmol/L,

165-175 mEq/L), but slightly higher than reported in Atlantic salmon (125-143 mmol/L,

125-143 mEq/L) which were regarded relatively stress-resistant, and much higher than

reported in Atlantic sturgeon (116-131 mmol/L, 116-131 mEq/L) (19,50,51). These

sources would also suggest that the sablefish RI's of monovalent sodium (168-179.9

mmol/L, 168-179.9 mEq/L) and potassium (1.2-3.86 mmol/L, 1.2-3.86 mEq/L), and

bivalent calcium (2.63–3.35 mmol/L, 10.51-13.38 mg/dL) are lower than in other marine

teleosts (175-200 mmol/L, 5-10 mmol/L and 3-5 mmol/L, respectively) (19,50). The

divergence of these parameters with reference values in older literature suggests that

further studies of fish species are warranted that take into account the demographics,

feeds, and production systems. Sex can also affect blood reference intervals (50). All-

female stocks are preferred in the production of many food fish species, including

sablefish because female fish grow larger than males (6,52). All-female stocks can be

achieved by mating females with neomales, which are genetic females who express

phenotypically male. This can either be achieved by the oral administration of

testosterone or by temperature management during the sexually undifferentiated phase

early in development (6). The mixed-sex population sampled in the present work

warrants further comparison and investigations into the physiology of this species. In

addition, the present sablefish blood RI should be compared to blood values of sablefish

that are cultured in other production systems like recirculating aquaculture systems

(RAS) and marine net pens, in different age groups, and from progeny that comes from

cultured rather than wild, captive brood-stock.

Acknowledgments

The authors thank Frank Sommers, Ron Johnson, and Rick Goetz of NOAA Fisheries,

NWFSC for supplying the cultured sablefish, and are grateful for Annabel Mendoza,

Serina Moheed, Brak Moczygemba, and Mary Beth Rew Hicks for animal husbandry

and fish necropsy assistance. This study was funded by an Oregon Sea Grant Program

Development Award #NA18OAR4170072 awarded to CBS and partially supported by

an Oregon State University Agricultural Research Foundation Grant #9026A awarded to

CBS.

References

1.   Sonu SC. Supply and market for Sablefish in Japan. NOAA Tech Memo NMFS.

2014;(October).

2.   Hanselman DH, Lunsford CR, Rodgveller CJ. Assessment of the sablefish stock

in Alaska. Stock Assess Fish Eval Rep Groundf Resour Gulf Alaska.

2015;2014(December 2015):576–717.

3.   National Oceanic and Atmospheric Administration. Pacific Coast Groundfish

Fishery 2019–20 Harvest Specifications, Yelloweye Rebuilding Plan Revisions,

and Management Measures: Environmental Assessment / Magnuson Stevens Act

Analysis / Regulatory Impact Review / Regulatory Flexibility Act Analysis.

2018;158.

4.   NOAA Fisheries. Annual Report to Congress on the Status of U.S. Fisheries.

Status Stock 2018. 2019;20.

5.   Cook MA, Massee KC, Wade TH, Oden SM, Jensen C, Jasonowicz A, et al.

Culture of sablefish (Anoplopoma fimbria) larvae in four experimental tank

designs. Aquac Eng. 2015;69:43–9.

6.   Huynh TB, Fairgrieve WT, Hayman ES, Lee JSF, Luckenbach JA. Inhibition of

ovarian development and instances of sex reversal in genotypic female sablefish

(Anoplopoma fimbria)exposed to elevated water temperature. Gen Comp

Endocrinol. 2019;279:88–98.

7.   Lee JSF, Cook MA, Luckenbach JA, Berejikian BA, Simchick CA, Oden SM, et al.

Investigation of long-term effects of larval rearing temperature on growth,

deformities, flesh quality, and phenotypic sex of cultured sablefish (Anoplopoma

fimbria). Aquaculture. 2017;479:91–9.

8.   Luckenbach JA, Fairgrieve WT. Gonadal sex differentiation and effects of dietary

methyltestosterone treatment in sablefish (Anoplopoma fimbria). Fish Physiol

Biochem. 2016;(42):233–48.

9.   Luckenbach JA, Fairgrieve WT, Hayman ES. Establishment of monosex female

production of sablefish (Anoplopoma fimbria) through direct and indirect sex

control. Aquaculture. 2017;479:285–96.

10.   Guzmán JM, Luckenbach JA, Middleton MA, Massee KC, Jensen C, Goetz FW,

432   et al. Reproductive life history of sablefish (Anoplopoma fimbria) from the U.S.

433   Washington coast. PLoS One. 2017;(September 8, 2017):1–24.

434   11.   Copes P. Adverse Impacts of Individual Quota Systems on COnservation and

435   Fish Harvest Productivity. Eighth Bienn Conf Int Inst Fish Econ Trade. 2000;(July

436   1996):1–16.

437   12.   Stearns G, Huppert D. Attitudes toward current and alternative managment

438   practices in Alaska's limited entry commercial herring roe fisheries. Washingt Sea

439   Grant. 2001;WSG-AV 02-.

440   13.   Lafferty KD, Harvell CD, Conrad JM, Friedman CS, Kent ML, Kuris AM, et al.

441   Infectious diseases affect marine fisheries and aquaculture economics. Ann Rev

442   Mar Sci. 2015;7:471–96.

443   14.   Arkoosh MR, Dietrich JP, Rew MB, Olson W, Young G, Goetz FW. Exploring the

444   efficacy of vaccine techniques in juvenile sablefish, Anoplopoma fimbria. Aquac

445   Res. 2018;49(1):205–16.

446   15.   Arkoosh MR, Dietrich JP. Pathogenicity of Members of the Vibrionaceae Family to

447   Cultured Juvenile Sablefish. J Aquat Anim Health. 2015;27(2):96–103.

448   16.   Nickum JG. Guidelines for the use of fishes in research. World Aquac. 2004;

449   17.   Arnold JE. Hematology of the sandbar shark, Carcharhinus plumbeus:

450   Standardization of complete blood count techniques for elasmobranchs. Vet Clin

Pathol. 2005;34(2):115–23.

18. Hrubec TC, Cardinale JL, Smith SA. Hematology and Plasma Chemistry

Reference Intervals for Cultured Tilapia (Oreochromis Hybrid). Vet Clin Pathol.

2000;29(1):7–12.

19. Matsche MA, Arnold J, Jenkins E, Townsend H, Rosemary K. Determination of

hematology and plasma chemistry reference intervals for 3 populations of captive

Atlantic sturgeon (Acipenser oxyrinchus oxyrinchus). Vet Clin Pathol.

2014;43(3):387–90.

20. Laposata M. Laboratory Medicine: The Diagnosis of Disease in the Clinical

Laboratory. In: Clinical Laboratory Reference Values [Internet]. 2nd ed. New York:

McGraw-Hill; 2014. Available from:

https://accessmedicine.mhmedical.com/content.aspx?bookid=1069§ionid=60

775149

21. Arnold JE, Matsche MA, Rosemary K. Preserving whole blood in formalin extends

the specimen stability period for manual cell counts for fish. Vet Clin Pathol.

2014;43(4):613–20.

22. Natt MP, Herrick CA. A new blood diluent for counting the erythrocytes and

leucocytes of the chicken. Poult Sci. 1952;31(4):735–8.

23. Grant KR. Fish hematology and associated disorders. Clin Lab Med.

2015;35(3):681–701.

24. Shigdar S, Cook D, Jones P, Harford A, Ward AC. Blood cells of murray cod

Maccullochella peelii peelii (Mitchell). J Fish Biol. 2007;70(3):973–80.

25. Geffré A, Concordet D, Braun JP, Trumel C. Reference Value Advisor: A new

freeware set of macroinstructions to calculate reference intervals with Microsoft

Excel. Vet Clin Pathol. 2011;40(1):107–12.

26. Friedrichs KR, Harr KE, Freeman KP, Szladovits B, Walton RM, Barnhart KF, et

al. ASVCP reference interval guidelines: Determination of de novo reference

intervals in veterinary species and other related topics. Vet Clin Pathol.

2012;41(4):441–53.

27. Witeska M. Erythrocytes in teleost fishes: A review. Zool Ecol. 2013;23(4):275–

81.

28. Weiner W, Topley E. Döhle bodies in the leucocytes of patients with burns. J Clin

Pathol. 1955;8(4):324–8.

29. Hine PM, Wain JM, Dunlop DM. Observations on granulocyte peroxidase in New

Zealand freshwater eels, Anguilla species. J Fish Biol. 1986;29(6):711–20.

30. Lamas J, Santos Y, Bruno DW, Toranzo AE, R A. Non-specific cellular responses

of rainbow trout to Vibrio anguillarum and its extracellular products (ECPs). J Fish

Biol. 45:839–54.

31. Calmarza P, Cordero J. Lipemia interferences in routine clinical biochemical tests.

2011;21(2):160–6.

32. Glick MR, Ryder KW, Jackson SA. Graphical comparisons of interferences in

clinical chemistry instrumentation. Clin Chem. 1986 Mar 1;32(3):470–5.

33. Dimeski G, Mollee P, Carter A. Increased lipid concentration is associated with

increased hemolysis. Clin Chem. 2005 Dec 1;51(12):2425–2425.

34. Nikolac N. Lipemia: Causes, interference mechanisms, detection and

management. Biochem Medica. 2014;24(1):57–67.

35. Soleimani N, Mohammadzadeh S, Asadian F. Lipemia interferences in

biochemical tests, investigating the efficacy of different removal methods in

comparison with ultracentrifugation as the gold standard. J Anal Methods Chem.

2020;2020(February 12):6 pages.

36. Parvin CA. Statistical quality control for quantitative measurement procedures:

Principles and definitions. 4th ed. CL. Wayne, PA: Clinical and Laboratory

Standards Institute; 2016.

37. Johnson RB, Nicklason PM, Armbruster LC, Sommers FC, Kim SK, Marancik D,

et al. Addition of the red macroalgae Turkish Towel Chondracanthus exasperates

and taurine improves the performance of alternative plant-based feeds for juvenile

sablefish Anoplopoma fimbria. Aquac Res. 2020;51(8):3191–204.

38. Nicklason PM, Johnson RB, Marancik DP. Effects of processing full-fat soya

beans and fish trim for sustainable sablefish Anoplopoma fimbria feeds. Aquac

Res. 2020;51(7):2667–75.

39. Van Pelt TI, Piatt JF, Lance BK, Roby DD. Proximate composition and energy

density of some North Pacific forage fishes. Comp Biochem Physiol - A Physiol.

1997;118(4):1393–8.

40. Karinen JF, Barnett HJ, Masuda M. Soft flesh in sablefish, Anoplopoma fimbria, of

southeastern Alaska: Relationships with depth, season, and biochemistry. Mar

Fish Rev. 2010;72(4):26–35.

41. Sullivan KM, Somero GN. Size and diet-related variations in enzymic activity and

tissue composition in the sablefish, Anoplopoma Fimbria. Biol Bull.

1983;164:315–26.

42. Ackman RG, Eaton CA, Ke PJ. Canadian Marine Oils of Low Iodine Value: Fatty

Acid Composition of Oils from Newfoundland Turbot (Greenland Halibut), Certain

Atlantic Herring, and a Sablefish. J Fish Res Board Canada. 1967;24(12):2563–

72.

43. Phleger CF, Soutar A, Schultz N, Duffrin E. Experimental sablefish fishing off San

Diego, California. Comm Fish Rev FWS Repr No 874. 1970;(874):31–40.

44. Sullivan KM, Smith KL. Energetics of Sablefish, Anoplopoma fimbria, under

Laboratory Conditions. Can J Fish Aquat Sci. 1982;39:1012–20.

45. Goetz FW, Jasonowicz AJ, Roberts SB. What goes up must come down: Diel

vertical migration in the deep-water sablefish (Anoplopoma fimbria) revealed by

pop-up satellite archival tags. Fish Oceanogr. 2018;27:127–42.

46.   Mason JC, Beamish RJ, McFarlane GA. Sexual maturity, fecundity, spawning,

532   and early life history of sablefish (Anoplopoma fimbria) off the Pacific Coast of

533   Canada. Can J Fish Aquat Sci. 1983;40(12):2126–34.

534   47.   Hanselman DH, Heifetz J, Echave KB, Dressel SC. Move it or lose it: Movement

535   and mortality of sablefish tagged in Alaska. Can J Fish Aquat Sci.

536   2014;72(2):238–51.

537   48.   Hrubec TC, Smith SA, Robertson JL, Bernard F, Veit HP, Libey G, et al. Blood

538   biochemical reference intervals for sunshine bass (Morone chrysops x Morone

539   saxatilis) in three culture systems. Am J Vet Res. 1996;(57):624–7.

540   49.   Silbergeld EK. Blood glucose: A sensitive indicator of environmental stress in fish.

541   Bull Environ Contam Toxicol. 1974;11(1):20–5.

542   50.   McDonald D, Milligan C. Chemical properties of the blood. In: Hoar W, Randall D,

543   Farrell A, editors. Fish Physi. San Diego, CA: Academic Press Inc.; 1992. p. 55–

544   133.

545   51.   Hunn J, Greer I. Influence of sampling on the blood chemistry of Atlantic Salmon.

546   Progress Fish-Culturist. 1991;53(3):184–7.

547   52.   Magerhans A, Hörstgen-Schwark G. Selection experiments to alter the sex ratio in

548   rainbow trout (Oncorhynchus mykiss) by means of temperature treatment.

549   Aquaculture. 2010;306(1–4):63–7.

Supporting information

S1 Fig. Lipemic blood. Photograph of lipemic blood sample from fish #34 on June

20th, 2019, after centrifugation. A packed cell fraction (bottom of tube) is overlaid by a

turbid, pale-beige plasma. The turbidity indicates a significant amount of blood lipids.

S1 Appendix. Raw data: Plasma biochemistry reference intervals. Data set

includes lipemic samples, RI calculation reports for each biochemistry parameter, and a

comparison of the RI of 87 fish (with lipemic samples) versus 81 fish (without lipemic

samples), plus all conversions from the conventional units to the SI units.

1c

T

6