Long noncoding RNA *VENTHEART* is required for cardiomyocyte specification and function

Albert Dashi[1,2,3], Wilson L.W. Tan[1,2,3], Chukwuemeka George Anene-Nzelu[1,2,3], Bangfen Pan[1,2,3], Autio

Matias Ilmari[2,3], Zenia Tiang[1,2,3], Robin J.G. Hartman[3,4], Justus Stenzig[9], Heming Wei[5], Chen Gao Bin[3],

Matthew Andrew Ackers-Johnson[1,2,3], Bing Lim[6], Anna Walentinsson[7], Vidhya Vardharajan Iyer[8,10],

Malin K.B. Jonsson[1, 8#], Roger S. Foo[1,2,3#]

[1]Cardiovascular Disease Translational Research Programme, Yong Loo Lin School of Medicine,

National University of Singapore, Singapore

[2]Cardiovascular Research Institute, National University Healthcare Systems, Singapore

[3]Genome Institute of Singapore, Agency for Science, Technology and Research, Singapore

[4]University of Utrecht, Heidelberglaan 8, 3584 CS Utrecht, the Netherlands.

[5]National Heart Research Institute Singapore (NHRIS), National Heart Centre Singapore

[6]Sana Biotechnology, 300 Technology Square, Cambridge, MA 02139, United State of America.

[7]Translational Science & Experimental Medicine, Research and Early Development, Cardiovascular,

Renal and Metabolism (CVRM), BioPharmaceuticals R&D, AstraZeneca, Gothenburg, Sweden.

[8]Bioscience Cardiovascular, Research and Early Development, Cardiovascular, Renal and Metabolism

(CVRM), BioPharmaceuticals R&D, AstraZeneca, Gothenburg, Sweden.

[9] University Medical Center Hamburg-Eppendorf, Hamburg, Germany

[10]Uppsala University, Uppsala, Sweden

[#]Shared senior authorship

Corresponding author:

Roger Foo

Genome Institute of Singapore

60 Biopolis Street

Singapore 138672

E: foosyr@gis.a-star.edu.sg

T: +44-66011036

F: +44-6808 8291

ABSTRACT

Rationale: Long noncoding RNAs (lncRNAs) control cardiac gene expression during heart

development and disease. It is accordingly plausible for the same lncRNA to regulate both cardiac

development, as well as play a role in adult heart disease progression. lncRNA regulators of early

cardiomyocyte (CM) lineage commitment have been identified and characterised, however those

controlling later CM specification remain unknown.

Objectives: In this study we identified a novel lncRNA required for CM specification, maturation and

function, and also discovered its suggested relevance to heart disease.

Methods and Results: We performed single cell RNA-seq on human embryonic stem cell derived

cardiomyocytes at 2, 6 and 12 weeks of differentiation. Weighted correlation network analysis

(WGCNA) identified core gene modules, including lncRNAs highly abundant and uniquely expressed

in the human heart. A lncRNA (we call *VENTHEART*, *VHRT*) co-expressed with cardiac maturation

and ventricular-specific genes *MYL2* and *MYH7*, as well as in adult human ventricular tissue. CRISPR-

mediated excision of *VHRT* led to impaired CM sarcomere formation, and loss of the CM

specification gene program. *VHRT* knockdown (KD) in hESC-CMs confirmed its regulatory role for

key cardiac contraction, calcium hemostasis and heart development genes, including *MYH6* and *RYR2*.

Functional evaluation after *VHRT* KD using impedance-based technology and action potential

recordings, proved reduced contraction amplitude and loss of the ventricular-like action potential in

CM, respectively. Through an integrative analysis of genome-wide association studies (GWAS),

expression quantitative trait locus (eQTL) and gene co-expression network, we found *VHRT* to be co-

regulated with core cardiac contractile genes, and the likely source of a heart failure genetic association

signal overlapping the *VHRT* gene locus. Finally, *VHRT* KD and human failing heart transcriptome

comparison validates the consistent downregulation again of cardiac contractile and calcium regulatory

genes ($P<0.05$).

Conclusion. We conclude that *VHRT* lncRNA is required for proper CM specification and function.

Furthermore, reduced *VHRT* may contribute to the development or progression of human heart disease.

INTRODUCTION

Cardiac development and heart disease progression are precisely coordinated at the transcriptome level

where gene expression programs are jointly controlled through signaling pathways and epigenetic

regulators. Human embryonic stem cell-derived cardiomyocytes (hESC-CMs) are a useful tool to study

cardiac commitment through their defined stages of cardiac differentiation[1,2]. This *in vitro* system can

model cardiac diseases including various forms of long QT syndrome[3], arrhythmogenic right ventricular

dysplasia[4], hypertrophic cardiomyopathy[5,6] and ischemic heart conditions[7]. Furthermore, hESC-CM are

routinely used to test novel drugs or therapeutic options. As examples, RNA interference (RNAi)

rescued the disease phenotype of hESC-CMs carrying a mutation causing long QT syndrome[8], or

carrying a mutation in phospholamban that results in dilated and arrhythmogenic cardiomyopathy[9].

As an *in vitro* cell model, many studies have investigated the mechanisms underlying early cardiac

commitment from hESC through to mesoderm and cardiac progenitor stages, and the signaling

pathways responsible for these transitions are relatively well-defined[10,11]. However, the later stages of

CM specification and maturation are less well-studied. Other than transcription factors, chromatin

modifiers and microRNAs that are well-known regulators of transcription, long noncoding RNAs

(lncRNAs) represent an additional layer of regulation that coordinate cardiac gene programs. LncRNAs

are >200 nucleotide long transcripts with limited protein coding potential, and often display species,

tissue- and disease-specific expression patterns[12]. They are diverse in their cellular expression patterns,

subcellular localization, evolutionary conservation, and mechanisms of action. Thousands of human

cardiac lncRNAs have been catalogued[13], but most remain to be functionally characterized. Previous

examples for cardiac development are *Braveheart (Bvht)*, necessary for the progression of nascent

mesoderm towards a cardiac fate through two different mechanisms of action[14,15], and *Fendrr*,

specifically expressed in the nascent lateral plate mesoderm and essential for proper development of the

heart and body wall[16]. More recently, another lncRNA was found to regulate murine mesodermal

specification by recruiting transcription factors eomes, trithorax group (TrxG) subunit WDR5, and

histone acetyltransferase GCN5 to the enhancer region of *Mesp1* gene, directly activating its

expression[17]. These studies were implemented using global transcriptome analysis focused on early

cardiac lineage commitment. Transcriptome profiling of the developing mammalian heart has also

revealed lncRNAs uniquely expressed in certain sets of cells and activated upon specific stimuli[18–20].

As with early differentiation, distinct lncRNA expression patterns in specific stages and cell states of

CM specification and maturation also underlie their essential biological function in heart disease.

Here, we have differentiated hESC to CM, and performed single cell RNA-seq analysis for co-

expression and transcriptional changes that occurred in hESC-CMs over time in culture. Hypothesizing

that key co-expressed lncRNAs could be drivers of CM specification, maturation and function, we

identified *VENTHEART* (*VHRT*) that was co-regulated with core cardiac development and maturation

genes and highly expressed in ventricular CMs. Additionally, an independent integrative analysis of

genome-wide association studies (GWAS), expression quantitative trait locus (eQTL) and human heart

tissue transcriptomes also pointed to the role of *VHRT* in cardiac function and heart failure disease

association.

RESULTS

Single cell RNA-seq of maturing hESC-CM identifies cardiac specific lncRNAs

A human ESC line with a *MYH6*-GFP reporter was generated and differentiated into CMs, using the

protocol published by Lian *et al*[21] with minor modifications (**Figure 1A, Supplementary Figure 1A-**

D). To map the single cell transcriptome of maturing hESC-CM, we captured cells from week 2 (W02),

week 6 (W06) and week 12 (W12) using the Fluidigm microfluidic C1 system. A total of 173 GFP+

(CMs) and 35 GFP- (non-CMs) cells from two biological repeats were sequenced. To keep only the

most optimum dataset, 21 samples were excluded from analysis due to either inadequate library size

(<1 M reads) or a low number of detected transcripts (<2,000 genes expressed) (**Supplementary Tables**

1 and 2). This resulted in a total of 36 CMs at W2, 60 CMs at W6 and 56 CMs at W12. To assess the

validity and quality of our data we undertook several further technical QC procedures (**Supplementary Figure 2A-F, Supplementary Table 1**). Only CMs (GFP+) were used for subsequent analysis for the remainder of this project. To confirm increased hESC-CM maturation, we firstly confirmed the consistent and timely upregulation of selected maturation genes (such as *TRDN*, *MYH7*, *TNNI3* and *MYL2*) (**Supplementary Figure 3A**), as well as calcium handling genes (*RYR2*, *ASPH*, *PLN* and *ATP2A2*). Notably, genes for cellular coupling gap junctions (*GJC1*), ion channel proteins (*KCNJ2*, *KCNA5*, *CACNA1G* and *HCN4*) and sarcomere assembly (*TCAP*) were also increased.

In order to explore for unsupervised co-regulated gene expression, we subjected the data to weighted correlation network analysis (WGCNA), and constructed gene-network modules that characterized expression changes across the time-course (**Supplementary Figure 4A, Supplementary Table 3**). Six modules differed significantly between the time points (**Figure 1B, Supplementary Figure 4B and Supplementary Table 4**). The "early-1" gene module consisted of genes highly expressed in W2 cells, and significantly downregulated in W6 and W12. Gene ontology (GO) functions for this "early-1" module was related to general tissue development and heart morphogenesis. The second module, "early-2", was also highly expressed at W2, and decreased at W6 and W12. The "early-2" module included pathways related to protein transport and localization. "Early-3" was another module that showed decreased expression over the time-course and contained GO pathways linked to cell cycle control. A "midstage" module showed the distinct opposite pattern of increasing expression from W02 to W06, and a further increase at W12. Genes in the "midstage" module were related to cardiac muscle development, maturation and cardiac functions, reflecting hESC-CMs strengthening their cardiac identity after W2 and mature and specify over time. Two further "late" modules showed a statistically significant but less marked difference during the time-course. "Late-1" included genes with general functions such as tau protein binding, "late-2" were those involved in general metabolic processes. Genes in all six modules reliably segregated cells into the three time points (**Figure 1C**), giving a comprehensive single cell transcriptional landscape of maturing hESC-CM over this time course.

Next, we curated for lncRNAs that correlated with transcriptional changes, and paid specific attention

to the "midstage" blue module with gene pathways involving cardiac muscle development and

maturation processes, including sarcoplasmic reticulum ion transport and actin mediated cell

contraction (**Figure 1D).** In this module we identified a set of lncRNAs highly co-regulated with other

important cardiac specific genes (**Figure 1E**). Among them, three (*RP11-532N4.2*, *LINC00881* and

RP11-432J24.5) are highly expressed in human hearts, and one (*RP1-46F2.2,* hg19 or *LINC01405*,

hg38) is limited to only cardiac ventricle and skeletal muscle, as shown from the publicly available

human tissue specific expression dataset (GTEx)[22,23] (**Supplementary Figure 4C**). Specifically,

LINC01405 we now call *VENTHEART* (*VHRT*), followed the tissue expression pattern of sarcomere

assembly and calcium regulatory genes, including myosin light chain 2 (*MYL2*), myosin heavy chain

beta (*MYH7*) and sarco/endoplasmic reticulum Ca2+-ATPase (*ATP2A2*), and showed the steepest

increase over the time-course compared to the other lncRNAs (**Figure 1F**). Human ESC-CMs in our

12-week time-course were largely negative for markers of the secondary heart field (*ISL1* and *KDR*),

and instead positive for primary heart field markers, *TBX5* and *CORIN*[18], and were MLC2v+ (encoded

by *MYL2*) for ventricular-like CMs (**Supplementary Figure 3B-C**). Indeed ranking all W12 hESC-CM

samples from low to high *VHRT* expression, confirmed that the expression of *MYL2*, *MYH7*, *ATP2A2*

and other genes of the same module ("midstage") was co-linear with *VHRT* (**Figure 1F**), suggesting

again that there are shared regulatory properties between genes in this module.

VENTHEART **is highly enriched in ventricular-like CMs.**

VHRT is highly conserved only among primates[24] (**Supplementary Figure 5À**), and the *VHRT* locus is

annotated to comprise of 6 isoforms (**Figure 2A, Supplementary Table 5**). To confirm differential

isoform expression, we performed primer specific RT-qPCR and identified isoform 6 as the most highly

expressed in hESC-CMs (**Figure 2B**). Isoform 6 *VHRT* was abundant in the cytoplasmic compartment,

whereas the other less expressed isoforms showed an unremarkable distribution across nuclear and

cytoplasmic compartments (**Figure 2C, Supplementary Figure 5B**). For cell and tissue expression

specificity, we found *VHRT* to be present in hESC-CM, but absent in human cardiac fibroblasts (hCF),

smooth muscle cells (hCASMC), human monocytic cell line (hTHP-1) and human coronary artery

endothelial cells (hCAEC) **(Figure 2D)**. Furthermore, human heart tissue expression analysis confirmed

that *VHRT* is absent or lowly expressed in atria, compared to ventricular tissue (**Supplementary Figure**

5C). This recapitulated data from GTEx portal, further underlining the specific *VHRT* expression

pattern in the heart. The Coding Potential Calculator (CPC)[25] algorithm suggested that among all

isoforms, isoform 6 may possess protein coding potential with a putative 76 aa long

peptide[26](**Supplementary Figure 6A-B**). We proceeded to test for the putative peptide empirically by

FLAG-tag at either C-terminus or N-terminus of *VHRT*-6 sORF (without introns), cloning it into a

vector for cell transfection. No bands were detected on Western blotting despite relevant positive

controls (**Supplementary Figure 6C**). This may reflect the possibility that the transcript is not

translated into a stable or functional peptide *in vitro*, and *VHRT* may function instead as a *bona fide*

non-coding RNA**.**

VHRT **deletion disrupts CM specification**

To test the hypothesis that *VHRT* regulates CM differentiation or specification, we performed *VHRT*

CRISPR/Cas9-directed knockout (KO) at hESC stage using a dual sgRNA system. To control for off-

target effects from CRISPR-targeting, we designed 2 unique sgRNA pairs that targeted different

locations of *VHRT (***Figure 3A-B).** Two independent homozygous knockout hESC lines were obtained

(clone 1: *VHRT* KO #1 and clone 16: *VHRT* KO #2). Untargeted cells were used as wild type control

(clone 9: WT) (**Supplementary Figure 8A-B).** Independent tri-lineage differentiation proved that

ectodermal, mesodermal and endodermal lineages were unaffected in both *VHRT* KO lines

(**Supplementary Figure 9A**). We then differentiated WT and KO lines to CM in parallel. Apart from

minimum variability between differentiations batches, both KO lines entered the initial stages of

differentiation efficiently (**Supplementary Figure 9B**). Key genes of the cardiac progenitor stage, such

as *NKX2-5,* were unchanged. The *MYH6*-GFP+ signal however was significantly reduced in KO at day

7-8, compared to WT (**Supplementary Figure 10A**), reflecting the likely importance of *VHRT* for the

transition from cardiac progenitor to early CM. Remarkably, upon continued culture, *VHRT* KO cells

showed reduced CM specification and maturation genes **(Figure 3C),** and an inability to adequately

develop sarcomere structure and organization from W6 to W12. Hence, while WT hESC-CM

strengthened their characteristic cellular organization from W2 to W12, the lack of proper sarcomere

assembly in *VHRT* KO #1 and *VHRT* KO #2 may reflect a failure to activate CM gene programs (**Figure**

3D, Supplementary Figure 10B-C).

***VHRT* deletion deregulates the transcriptome of maturing hESC-CM**

A small subset of *VHRT* KO cells (15% of *VHRT* KO #1, and 20% of *VHRT* KO #2) turned on the

MYH6-GFP reporter during the differentiation protocol (**Supplementary Figure 11A).** This raised the

question whether some *VHRT* KO cells developed general CM gene expression. We therefore isolated

GFP+ cells from WT and *VHRT* KO #1 at W2, W6 and W12 (**Supplementary Figure 11B**), and

performed RNA-seq to compare global gene expression. *VHRT* deletion resulted in a global increase in

differentially expressed genes over time in culture (**Figure 3E-G**). Across W2, W6 and W12, cardiac

contractile and calcium regulatory genes including *MYH7, MYL2, SLC8A1* and *IRX4* and failed to

upregulate in *VHRT* KO, compared with WT (**Figure 3C-E, Supplementary Table 6**). In contrast,

ectodermal and neuronal function genes, such as *SOX2, PAX6* and *RAX*, were significantly upregulated,

especially in W12 *VHRT* KO cells (**Figure 3G, Supplementary Table 6**). Furthermore, genes involved

in metabolic processes were persistently downregulated from W2 to W12, reflecting the lack of a

maturation metabolic switch in *VHRT* KO (**Supplementary Table 6**).

***VHRT* knockdown downregulates sarcomere assembly and calcium handling genes, and reduces**

CM contractility

Next, we sought to assess whether *VHRT* is also implicated in the regulation and maintenance of CM

function in hES-CM. We therefore performed *VHRT* knockdown (KD) using 2 independent GapmeRs,

specifically targeting *VHRT* isoform 6, since it is the most expressed isoform in hES-CMs (**Figure 4A**).

This was carried out in W6 hESC-CM, where *VHRT* gene expression was elevated compared to

preceding stages of differentiation. 75% *VHRT* KD was achieved with GapmeR#1, and 72.4% with

GapmeR#2, respectively. (**Figure 4B**).

VHRT KD CM showed significant differentially expressed (DE) genes belonging to biological

processes controlling cardiac electrophysiology, cardiac structure and heart development (**Figure 4C-**

D). There was significant downregulation of structural and sarcomere assembly genes including *MYH6,*

FHOD3 and *MYOM1,* and known markers of heart tissue morphogenesis *TBX20*, *HEY2* and *ALPK3.*

Remarkably, a strong repression was observed for calcium regulatory genes including *RYR2, SLC8A1*

and *CMYA5* (**Figure 4E-F).** Importantly, these findings were specific because the same significant

changes were not observed for other CM marker genes (**Figure 4B-D,** non-DE genes). To assess the

function on contractility and CM electrophysiology, we also performed an independent *VHRT* siRNA

KD, followed by impedance-based assays. *VHRT* KD significantly affected contractility performance,

with decreased relaxation velocity, decreased upstroke velocity, decreased amplitude and a prolonged

pulse width, compared to non-targeted siRNA (NT) and untreated (UT) controls (**Figure 4G,**

Supplementary Figure 11A). *VHRT* KD induced similar changes in contractility as KD of *ATP2A2,*

encoding for SERCA2a, consistent with altered calcium handling in *VHRT*-deficient CM. **(Figure 4G,**

Table 1). Patch clamp experiments showed that *VHRT*-KD cells had significantly more positive

maximum depolarization potential (MDP), compared to control (-54.2±6.5 vs -61.2±7.1 mV, n=28,

p≤0.01). *VHRT*-KD cells also displayed a more triangular action potential (AP) waveform, as assessed

by either the APD90/APD50 ratio (1.34±0.12 vs 1.23±0.14, p≤0.05) or the APD90-APD20 difference

(256±81 vs 166±50 ms, *p*≤0.001) (**Supplementary Figure 11B).** Representative AP from

spontaneously beating cells and cells paced at 1Hz are shown in **Supplementary Figure 11C**. Taken

together, these data confirmed that *VHRT* KD significantly compromised CM function.

Integrated GWAS and eQTL analysis implicates *VHRT* in human heart disease

Related to the unique expression profile of *VHRT* for differentiating CMs above, and the well-known

hallmark of "fetal gene reprogramming" in the heart disease stress-gene response, we also set out to

investigate a potential role for *VHRT* in human heart disease, via analysis of publicly available genome-

wide association study (GWAS) data sets[27–31]. Indeed, at least 2 independent GWAS signals were

located in or near the *VHRT* locus at chromosome 12q24.11. Significant signals (*p*-value<5e-8) for

coronary artery disease (CAD) and myocardial infarction (MI) overlapped with the genes *SH2B3* and

ATXN2 downstream of *VHRT*, but these were unlikely related to *VHRT* as they did not associate with

cardiac *VHRT* expression based on cis-eQTL data from human left ventricular (LV) samples.

Instead, a tightly linked cluster of SNPs (r2>0.8) overlapping with the *VHRT* gene and significantly

associated with *VHRT* expression in heart LV, showed suggestive association with non-ischemic heart

failure (HF) (*p*-value=5e-5) (**Figure 5A, B)**, indicating that the HF GWAS signal is most likely

mediated by altered *VHRT* cardiac expression. SNPs associated with reduced cardiac *VHRT* expression

were associated with increased risk for HF (OR=1.12-1.15) (**Supplementary Figure 12A-B**).

Importantly, none of the SNPs showed association with *MYL2* cardiac expression, suggesting that

VHRT rather than *MYL2* is the likely mediator of the observed cardiac disease GWAS signal. In line

with these findings, cardiac *VHRT* mRNA expression was found to be downregulated in a

transcriptomics study of patients with dilated cardiomyopathy, compared to healthy controls (adjusted

p-value<0.05)[31] (**Figure 5D)**. We also performed co-expression analysis using GeneNetwork v2.0[30],

which predicted pathway and human phenotype associations using 31,499 public human RNA-seq

samples. *VHRT* was once again tightly co-regulated with key CM sarcomeric genes, including *MYL2*,

MYL3, *MYL1*, *MYH7*, *TCAP*, *CSRP3* and *TNNC1,* replicating the empirical evidence from the gene

network analysis of single cell transcriptomes above. Furthermore, *VHRT* was predicted to participate

in heart disease pathways and phenotypes including hypertrophic and dilated cardiomyopathy (HCM

and DCM, respectively), as well as ventricular tachycardia **(Figure 5C, Supplementary Table 8)**.

***VHRT* KD transcriptome partially resembled that of gene expression changes in human heart**

failure

Building on the findings described above, we compared the *VHRT* KD transcriptome of DE genes

(N=1,100; 573 down- and 527 upregulated) with the expression profile of the same genes in human

failing hearts (Dilated Cardiomyopathy, DCM) (GEO accession: GSE141910). The latter study

comprised of RNA-seq from 61 left ventricle explants of idiopathic DCM failing (n=29) and non-failing

hearts (NF, n= 32) (**Supplementary Table 9). **Among *VHRT* KD DE genes, 73 (out of 1,100) were

also differentially expressed in DCM hearts (**Figure 6A, Supplementary Figure 13**, **Supplementary**

Table 10). Importantly, genes that were downregulated in *VHRT* KD, and recapitulated in DCM

("Down/Down" group) contained those enriched in pathways regulating cardiac muscle contraction,

hypertrophic cardiomyopathy (HCM) and dilated cardiomyopathy DCM (p ≤ 0.05), based on pathway

and molecular function enrichment analysis using Enricher[32] (**Figure 6B**). Functional terms related to

processes involved in actin filament binding and calcium handling (p ≤ 0.05), including genes such as

MYH6, *CAMK1D*, *MYO7A* and *CACNB2* (**Figure 6C, Supplementary Figure 13**), pointing again to

the potential mechanisms by which *VHRT* mediates disease progression in HF.

DISCUSSION

Our findings provide the first identification of the long non-coding RNA *VHRT* for CM specification,

maturation, and its potential involvement in HF disease pathophysiology. By tracking single-cell

resolution transcriptomes of hESC-CMs in prolonged culture, we have compiled an extensive catalogue

of hESC-CM expressed genes. Co-expression gene networks, and integrative GWAS and eQTL

analyses, highlighted *VHRT* as a novel ventricular-enriched lncRNA, whose expression is limited only

to primates.

The overall immaturity of hESC-CMs in culture is well acknowledged. Simple extended time in culture

is a useful means to achieve some additional maturity[1,19,33,34]. Our data reveals that key genes, *TNNI3*,

KCNJ2, *CASQ2*, and *TCAP* are upregulated during CM maturation, appearing at W6 or W12. Between

the 2 potassium channel genes: *KCNJ2,* which is ventricular-enriched, and *KCNA5,* which is

predominantly atrial, we found that *KCNJ2+* and *KCNA5+* hES-CMs are mutually exclusive.

Dichotomy for hESC-CMs in this differentiation protocol exists, at least in this respect[35,36]. We have

thus observed that with the 12-week time-course in this differentiation protocol, hESC-CMs come to

display largely primary heart field markers (*TBX5*, *HCN4* and *CORIN*), specifying mainly into

ventricular-like cells, also corroborated by upregulated ventricular-enriched *MYL2*, *MYH7* and *IRX4*.

This is consistent with the ventricular phenotype also described previously for this protocol[21] . Genes

in the "early-1" gene module, abundant at W2, and subsequent downregulated in W6 and W12, included

those that play a role in general organ development (*BMP2, FGF10, WNT11, TWIST1, SOX11* and

MEF2C), indicating that even though W2 hESC-CMs are committed to the cardiac lineage, they retain

an overall generic transcriptome similar to cells in other organs. Similarly, genes in the "early-2" and

"early-3" modules comprised of those for protein transport and localization, and cell cycle regulators

(*PSMA6*, *PSMB5*, *PSMA3* and *PSMB4*) that regulate mitotic cell cycle G1/S checkpoint and DNA

damage control. The downregulation of these genes coheres with the notion that hESC-CMs decrease

their proliferative capacity over the differentiation time-course. In contrast, genes in the "midstage"

module are significantly upregulated between W2 and W6, remaining stably abundant thereafter. These

are genes related to cardiac development and maturation, including those responsible for cardiac

contractile structures (*MYH7, TTN, MYL2, TNNI3, ASPH, MYBPC3*), higher-order myofibrillar

organization, cytoskeletal assembly (*MURC132* and *LDB3133*), and calcium handling and transport

(*RYR2, ATP2A2, CASQ1, ASPH, ANK2 and SLC8A1*). The "late-1"' and "late-2" modules contained

genes that are only expressed in subsets of W12 cells, where their expression contributed to significant

heterogeneity observed at the W12 stage. The "late-2" module comprised of a long list of ribosomal

protein pseudogenes associated with general metabolic processes. The significance of the latter will

need further investigation.

LncRNAs act as fine switches to modulate and orchestrate multiple aspects of cardiac development.

In recent years, many cardiac lncRNAs have been catalogued and functionally characterized,

especially for CM differentiation[37–39]. However, less is known for the later stages of CM specification

and maturation. Moreover, finding lncRNAs that exclusively regulate either atrial, ventricular or

pacemaker cells functions is challenging since different cardiac developmental stages are dictated by

different epigenetic mechanisms. The "midstage" module contained specific cardiac protein coding

genes known to support CM specification, maturation and function. Therefore, the transcriptional

profile of this module accordingly identified CM specific and enriched lncRNAs, among which *VHRT*

was highly co-regulated with *MYH7, MYL2, ATP2A2* and other cardiac maturation genes. Being

enriched in human ventricular tissue, but not expressed in atrial tissue or other cell types of the heart,

VHRT may indeed have a unique role in ventricular CM.

CM maturation and specification processes are promoted by pathways and gene programs for

metabolism, myofiber structure, electrophysiology and cell cycle[40]. Myosin heavy-chains (MYH6

and MYH7) and light-chains (MYL2) are myofiber proteins important for heart morphogenesis.

Pathogenic *MYH7* mutations are causal to hypertrophic cardiomyopathy[41]. Patient-specific iPSC-CMs

carrying the *MYH7*-E848G mutation, result in dysregulated myofibril alignment and contractile

impairment[42]. In the same way, Wei Zhou et al.[43] differentiated hiPSC derived CM from patients with

MYL2-R58Q[44,45] mutation and demonstrated that after 60 days in culture, *MYL2*-R58Q-iPSC-CMs

also showed sarcomeric disorganization, calcium dysregulation and contractility malfunction. *VHRT*

KO, starting from W6, displayed the similar myofilament disassembly and contractility defect,

concurrent with the downregulation of cardiac contractile and calcium regulatory genes. Indeed,

VHRT-KO phenocopies *MYL2*-R58Q and *MYH7*-E848G mutant CMs, implying that the overall

deregulated CM gene programme caused by *VHRT*-KO is at least in part, similar to that regulating

MYL2 and *MYH7*. Importantly, pluripotency, mesendoderm and cardiac mesoderm stages were not

affected by *VHRT* KO. This was also demonstrated by the finding that general CM genes were

normally expressed at W2, whereas with time in culture, ectodermal and neuronal function genes

became upregulated instead. However, unlike gene profiles of mutant *MYH7*[46,47] and *MYL2* cells[48], the

overall gene profile of W6-12 *VHRT*-KO (and KD) consistently show a downregulation of contractile

and calcium regulatory genes, and the lack of a metabolic switch towards "mature" CMs. These

results suggest a role for *VHRT* as an epigenetic switch for CM gene programmes during specification

and maturation, instead of merely conferring effector functions like those of the sarcomeric

components like *MYH7* or *MYL2*.

Other lncRNAs have also been shown to control cardiac gene expression previously. Many do so by

interacting with repressive protein complexes and transcription factors in the nucleus, thereby

regulating local gene expression through chromatin remodelling. Our characterization experiments

however, show *VHRT* to be localized in CM cytoplasm, indicating that *VHRT* may regulate gene

programs from the cytoplasmic compartment and not the nucleus, unlike mechanisms for the majority

of other lncRNAs. At least from our current assessment, *VHRT* does not appear to encode any detectably

stable micropeptide in hES-derived CM. Instead, *VHRT* KD by GapmeR was sufficient to downregulate

the expression of cardiac contraction, calcium transport and other ion channel genes. Whole cell patch

clamp recordings confirmed that *VHRT* KD cells showed the loss of a ventricular-like action potential.

Consistent with functional cytoplasmic *VHRT,* alternative siRNA-mediated *VHRT* KD also verified

reduced CM relaxation velocity, upstroke velocity and contractility amplitude. Some lncRNAs are

translocated out of the nucleus and shuttled to specific cytoplasmic locations to regulate protein

expression or function. Some do so by sponging miRNA[49] and other by directly binding to protein or

protein complexes. An example of the latter is the antisense lncRNA *ZFAS1*, which regulates Ca2+ flux

in the cells by directly inhibiting the activity of the protein SERCA2a (sarcoplasmic reticulum Ca2+-

ATPase 2a)[50]. Overexpression of *ZFAS1* altered Ca2+ flux leading to intracellular Ca2+ overload in

CM. Other lncRNAs, such as the *Sirt1* antisense lncRNA binds to *Sirt1* mRNA 3′-UTR and

subsequently increased *Sirt1* abundance[51]. Exemplified by these examples, *VHRT* repression that

negatively regulates the CM contraction and structural gene program, prompts the need for further

studies to understand how cytoplasmic *VHRT* supports and stabilizes the activity and abundance of

cardiac contractility genes expression.

To date many genetic variants have been associated with cardiovascular disease[28], but few have been

linked through lncRNAs function. An example is the cardioprotective lncRNA *Myocardial infarction*

associated transcript (*MIAT*). Six SNPs at the *MIAT* locus are associated with myocardial infarction

(MI)[52]. More recently, three SNPs (e.g. rs7140721, rs3729829 and rs3729825) at the *Myosin Heavy*

Chain Associated RNA Transcript (*MHRT*) locus are associated with increased risk of HF and

prognosis[53]. Thus far, we have concluded, thorough integrative analysis of GWAS, cis-eQTL and RNA-

seq from ~ 31,5K sample patients[30], that a series of SNPs in the *VHRT* locus also associated with heart

disease. The SNPs rs6489844-G and rs11065780-C are both associated with decreased *VHRT*

expression in human left ventricle and increased risk for HF. Moreover, *VHRT* is downregulated in

failing hearts from patients with DCM, reflecting the link between *VHRT* and HF. Indeed

downregulated contractile and calcium regulatory genes in failing DCM hearts, consistent with

downregulated genes in VHRT KD, again concur with the possible mechanism mediated the

contribution of VHRT to HF disease progression. The lack of species conservation for *VHRT*

complicates *in-vivo* validation since a homologue for *VHRT* in rodents cannot be identified.

Nevertheless, novel strategies to assess *VHRT* as a target for drug development are now attractive to

pursue.

In conclusion, we have shown that *VHRT* is a crucial regulator for CM specification and lies upstream

of key CM structural and contractile gene programs. Perturbation of *VHRT* results in the loss of CM

contractile function. Moreover, concordant HF genome-wide analyses and human heart transcriptomic

cross-comparisons motivate the need for follow up functional studies using models of heart disease.

ACKNOWLEDGEMENTS

Singapore International Graduate Award (A.D.); EMBO long-term fellowship 'EMBO ALTF 304-

2012' (M.K.B.J.). Funding from the Singapore National Medical Research Council (R.S.Y.F.) and

A*STaR Biomedical Research Council (R.S.Y.F., L.B.). Our gratitude to Dr. William Sun and Dr.

Kristy Purnamawati for the use of their multielectrode system.

AUTHOR CONTRIBUTIONS

AD, experimental planning and execution, figure preparation, manuscript drafting and writing

WT, bioinformatics analysis, figure preparation

CGA-N, experimental planning, figure preparation, manuscript writing

BP, cell culture and differentiation optimization

AM, CRISPR/Cas9 assay and experiments

CB, generation of MYH6 reporter line

MA, cell culture and RNA preparation

VV, contractility-based assay

RH, preparation of buffers and experimental assistance

ZT, library preparation for sequencing

HW, experimental planning and analysis, figure preparation

BL, senior supervisor, funding support

AW, bioinformatics analysis, figure preparation

MJ, study design, experimental planning and execution, analysis, figure preparation, manuscript

drafting and writing

RF, senior supervisor, funding support, manuscript writing

DISCLOSURE DECLARATION

The authors declare that they have no conflict of interest. M.J. and A.W. are employed by AstraZeneca.

METHODS AND MATERIALS

See Supplementary information online for a detailed description of all methods and materials.

REFERENCES

1.   Sartiani L, Bettiol E, Stillitano F, Mugelli A, Cerbai E, Jaconi ME. Developmental changes in

cardiomyocytes differentiated from human embryonic stem cells: a molecular and

electrophysiological approach. *Stem Cells* [Internet]. 2007;25:1136–1144. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/17255522

2.   Karakikes I, Ameen M, Termglinchan V, Wu JC. Human induced pluripotent stem cell-

derived cardiomyocytes: insights into molecular, cellular, and functional phenotypes. *Circ Res*

[Internet]. 2015;117:80–88. Available from: https://www.ncbi.nlm.nih.gov/pubmed/26089365

3.   Friedrichs S, Malan D, Voss Y, Sasse P. Scalable Electrophysiological Investigation of iPS

Cell-Derived Cardiomyocytes Obtained by a Lentiviral Purification Strategy. *J Clin Med*

[Internet]. 2015;4:102–123. Available from: https://www.ncbi.nlm.nih.gov/pubmed/26237021

4.   Kim C, Wong J, Wen J, Wang S, Wang C, Spiering S, Kan NG, Forcales S, Puri PL, Leone

TC, Marine JE, Calkins H, Kelly DP, Judge DP, Chen HS. Studying arrhythmogenic right

ventricular dysplasia with patient-specific iPSCs. *Nature* [Internet]. 2013;494:105–110.

Available from: https://www.ncbi.nlm.nih.gov/pubmed/23354045

5.   Wu H, Yang H, Rhee J-W, Zhang JZ, Lam CK, Sallam K, Chang ACY, Ma N, Lee J, Zhang

H, Blau HM, Bers DM, Wu JC. Modelling diastolic dysfunction in induced pluripotent stem

cell-derived cardiomyocytes from hypertrophic cardiomyopathy patients. *Eur Heart J*. 2019;

6.   Li X, Lu WJ, Li Y, Wu F, Bai R, Ma S, Dong T, Zhang H, Lee AS, Wang Y, Lan F. MLP-

deficient human pluripotent stem cell derived cardiomyocytes develop hypertrophic

cardiomyopathy and heart failure phenotypes due to abnormal calcium handling. *Cell Death*

Dis. 2019;10.

7.   Cardiovascular diseases [Internet]. [cited 2019 Nov 14];Available from:

https://www.who.int/health-topics/cardiovascular-diseases/

8.   Matsa E, Dixon JE, Medway C, Georgiou O, Patel MJ, Morgan K, Kemp PJ, Staniforth A,

Mellor I, Denning C. Allele-specific RNA interference rescues the long-QT syndrome

phenotype in human-induced pluripotency stem cell cardiomyocytes. *Eur Hear J* [Internet].

2014;35:1078–1087. Available from: https://www.ncbi.nlm.nih.gov/pubmed/23470493

9.   Stroik DR, Ceholski DK, Mleczko J, Thanel PF, Bidwell PA, Autry JM, Cornea RL, Thomas

DD. Viral expression of a SERCA2a-activating PLB mutant improves calcium cycling and

synchronicity in dilated cardiomyopathic hiPSC-CMs. *bioRxiv*. 2019;699975.

10.   Domian IJ, Chiravuri M, van der Meer P, Feinberg AW, Shi X, Shao Y, Wu SM, Parker KK,

Chien KR. Generation of functional ventricular heart muscle from mouse ventricular

progenitor cells. *Science (80-)* [Internet]. 2009;326:426–429. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/19833966

11.   Kattman SJ, Witty AD, Gagliardi M, Dubois NC, Niapour M, Hotta A, Ellis J, Keller G.

Stage-specific optimization of activin/nodal and BMP signaling promotes cardiac

differentiation of mouse and human pluripotent stem cell lines. *Cell Stem Cell* [Internet].

2011;8:228–240. Available from: https://www.ncbi.nlm.nih.gov/pubmed/21295278

12.   Ulitsky I, Bartel DP. XLincRNAs: Genomics, evolution, and mechanisms. Cell. 2013;

13.   van Heesch S, Witte F, Schneider-Lunitz V, Schulz JF, Adami E, Faber AB, Kirchner M,

Maatz H, Blachut S, Sandmann CL, Kanda M, Worth CL, Schafer S, Calviello L, Merriott R,

Patone G, Hummel O, Wyler E, Obermayer B, Mucke MB, Lindberg EL, Trnka F, Memczak

471  S, Schilling M, Felkin LE, Barton PJR, Quaife NM, Vanezis K, Diecke S, Mukai M, Mah N,

472  Oh SJ, Kurtz A, Schramm C, Schwinge D, Sebode M, Harakalova M, Asselbergs FW, Vink A,

473  de Weger RA, Viswanathan S, Widjaja AA, Gartner-Rommel A, Milting H, Dos Remedios C,

474  Knosalla C, Mertins P, Landthaler M, Vingron M, Linke WA, Seidman JG, Seidman CE,

475  Rajewsky N, Ohler U, Cook SA, Hubner N. The Translational Landscape of the Human Heart.

476  *Cell* [Internet]. 2019;178:242-260 e29. Available from:

477  https://www.ncbi.nlm.nih.gov/pubmed/31155234

478 14. Klattenhoff CA, Scheuermann JC, Surface LE, Bradley RK, Fields PA, Steinhauser ML, Ding

479  H, Butty VL, Torrey L, Haas S, Abo R, Tabebordbar M, Lee RT, Burge CB, Boyer LA.

480  Braveheart, a long noncoding RNA required for cardiovascular lineage commitment. *Cell*

481  [Internet]. 2013;152:570–583. Available from: internal-pdf://146.143.144.64/Braveheart, a

482  Long Noncoding.pdf

483 15. Xue Z, Hennelly S, Doyle B, Gulati AA, Novikova I V, Sanbonmatsu KY, Boyer LA. A G-

484  Rich Motif in the lncRNA Braveheart Interacts with a Zinc-Finger Transcription Factor to

485  Specify the Cardiovascular Lineage. *Mol Cell* [Internet]. 2016;64:37–50. Available from:

486  https://www.ncbi.nlm.nih.gov/pubmed/27618485

487 16. Sauvageau M, Goff LA, Lodato S, Bonev B, Groff AF, Gerhardinger C, Sanchez-Gomez DB,

488  Hacisuleyman E, Li E, Spence M, Liapis SC, Mallard W, Morse M, Swerdel MR, D'Ecclessis

489  MF, Moore JC, Lai V, Gong G, Yancopoulos GD, Frendewey D, Kellis M, Hart RP,

490  Valenzuela DM, Arlotta P, Rinn JL. Multiple knockout mouse models reveal lincRNAs are

491  required for life and brain development. *Elife* [Internet]. 2013;2:e01749. Available from:

492  https://www.ncbi.nlm.nih.gov/pubmed/24381249

493 17. Guo X, Xu Y, Wang Z, Wu Y, Chen J, Wang G, Lu C, Jia W, Xi J, Zhu S, Jiapaer Z, Wan X,

494  Liu Z, Gao S, Kang J. A Linc1405/Eomes Complex Promotes Cardiac Mesoderm

495  Specification and Cardiogenesis. *Cell Stem Cell* [Internet]. 2018;22:893-908 e6. Available

496  from: internal-pdf://113.73.8.13/linc1405 -exom regualtion.pdf

497 18. Zhang JZ, Termglinchan V, Shao NY, Itzhaki I, Liu C, Ma N, Tian L, Wang VY, Chang ACY,

498  Guo H, Kitani T, Wu H, Lam CK, Kodo K, Sayed N, Blau HM, Wu JC. A Human iPSC

Double-Reporter System Enables Purification of Cardiac Lineage Subpopulations with

Distinct Function and Drug Response Profiles. *Cell Stem Cell* [Internet]. 2019;24:802-811 e5.

Available from: https://www.ncbi.nlm.nih.gov/pubmed/30880024

19.   DeLaughter DM, Bick AG, Wakimoto H, McKean D, Gorham JM, Kathiriya IS, Hinson JT,

Homsy J, Gray J, Pu W, Bruneau BG, Seidman JG, Seidman CE. Single-Cell Resolution of

Temporal Gene Expression during Heart Development. *Dev Cell* [Internet]. 2016;39:480–490.

Available from: https://www.ncbi.nlm.nih.gov/pubmed/27840107

20.   Li G, Xu A, Sim S, Priest JR, Tian X, Khan T, Quertermous T, Zhou B, Tsao PS, Quake SR,

Wu SM. Transcriptomic Profiling Maps Anatomically Patterned Subpopulations among Single

Embryonic Cardiac Cells. *Dev Cell* [Internet]. 2016;39:491–507. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/27840109

21.   Lian X, Zhang J, Azarin SM, Zhu K, Hazeltine LB, Bao X, Hsiao C, Kamp TJ, Palecek SP.

Directed cardiomyocyte differentiation from human pluripotent stem cells by modulating

Wnt/beta-catenin signaling under fully defined conditions. *Nat Protoc* [Internet]. 2013;8:162–

175. Available from: https://www.ncbi.nlm.nih.gov/pubmed/23257984

22.   GTEx Portal [Internet]. [cited 2019 Nov 9];Available from: https://www.gtexportal.org/home/

23.   Carithers LJ, Moore HM. The Genotype-Tissue Expression (GTEx) Project. *Biopreserv*

Biobank [Internet]. 2015;13:307–308. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/26484569

24.   Ensembl genome browser 98 [Internet]. [cited 2019 Dec 28];Available from:

https://asia.ensembl.org/index.html

25.   Kong L, Zhang Y, Ye ZQ, Liu XQ, Zhao SQ, Wei L, Gao G. CPC: assess the protein-coding

potential of transcripts using sequence features and support vector machine. *Nucleic Acids Res*

[Internet]. 2007;35:W345-9. Available from: https://www.ncbi.nlm.nih.gov/pubmed/17631615

26.   Uncharacterized protein - Homo sapiens (Human) [Internet]. [cited 2019 Nov 10];Available

from: https://www.uniprot.org/uniprot/Q8IVN4

27.   Aragam KG, Chaffin M, Levinson RT, McDermott G, Choi SH, Shoemaker MB, Haas ME,

Weng LC, Lindsay ME, Smith JG, Newton-Cheh C, Roden DM, London B, Wells QS, Ellinor

PT, Kathiresan S, Lubitz SA. Phenotypic Refinement of Heart Failure in a National Biobank

Facilitates Genetic Discovery. *Circulation*. 2019;

28.   Nikpay M, Goel A, Won HH, Hall LM, Willenborg C, Kanoni S, Saleheen D, Kyriakou T,

Nelson CP, CHopewell J, Webb TR, Zeng L, Dehghan A, Alver M, MArmasu S, Auro K,

Bjonnes A, Chasman DI, Chen S, Ford I, Franceschini N, Gieger C, Grace C, Gustafsson S,

Huang J, Hwang SJ, Kim YK, Kleber ME, Lau KW, Lu X, Lu Y, Lyytikäinen LP, Mihailov E,

Morrison AC, Pervjakova N, Qu L, Rose LM, Salfati E, Saxena R, Scholz M, Smith A V.,

Tikkanen E, Uitterlinden A, Yang X, Zhang W, Zhao W, De Andrade M, De Vries PS, Van

Zuydam NR, Anand SS, Bertram L, Beutner F, Dedoussis G, Frossard P, Gauguier D, Goodall

AH, Gottesman O, Haber M, Han BG, Huang J, Jalilzadeh S, Kessler T, König IR, Lannfelt L,

Lieb W, Lind L, MLindgren C, Lokki ML, Magnusson PK, Mallick NH, Mehra N, Meitinger

T, Memon FUR, Morris AP, Nieminen MS, Pedersen NL, Peters A, Rallidis LS, Rasheed A,

Samuel M, Shah SH, Sinisalo J, EStirrups K, Trompet S, Wang L, Zaman KS, Ardissino D,

Boerwinkle E, Borecki IB, Bottinger EP, Buring JE, Chambers JC, Collins R, Cupples L,

Danesh J, Demuth I, Elosua R, Epstein SE, et al. A comprehensive 1000 Genomes-based

genome-wide association meta-analysis of coronary artery disease. *Nat Genet*. 2015;

29.   Roselli C, Chaffin MD, Weng LC, Aeschbacher S, Ahlberg G, Albert CM, Almgren P, Alonso

A, Anderson CD, Aragam KG, Arking DE, Barnard J, Bartz TM, Benjamin EJ, Bihlmeyer

NA, Bis JC, Bloom HL, Boerwinkle E, Bottinger EB, Brody JA, Calkins H, Campbell A,

Cappola TP, Carlquist J, Chasman DI, Chen LY, Chen YDI, Choi EK, Choi SH,

Christophersen IE, Chung MK, Cole JW, Conen D, Cook J, Crijns HJ, Cutler MJ, Damrauer

SM, Daniels BR, Darbar D, Delgado G, Denny JC, Dichgans M, Dörr M, Dudink EA, Dudley

SC, Esa N, Esko T, Eskola M, Fatkin D, Felix SB, Ford I, Franco OH, Geelhoed B, Grewal

RP, Gudnason V, Guo X, Gupta N, Gustafsson S, Gutmann R, Hamsten A, Harris TB,

Hayward C, Heckbert SR, Hernesniemi J, Hocking LJ, Hofman A, Horimoto ARVR, Huang J,

Huang PL, Huffman J, Ingelsson E, Ipek EG, Ito K, Jimenez-Conde J, Johnson R, Jukema JW,

Kääb S, Kähönen M, Kamatani Y, Kane JP, Kastrati A, Kathiresan S, Katschnig-Winter P,

Kavousi M, Kessler T, Kietselaer BL, Kirchhof P, Kleber ME, Knight S, Krieger JE, Kubo M,

Launer LJ, Laurikka J, Lehtimäki T, Leineweber K, Lemaitre RN, Li M, Lim HE, et al. Multi-

ethnic genome-wide association study for atrial fibrillation. *Nat Genet*. 2018;

30. Deelen P, van Dam S, Herkert JC, Karjalainen JM, Brugge H, Abbott KM, van Diemen CC,

van der Zwaag PA, Gerkes EH, Zonneveld-Huijssoon E, Boer-Bergsma JJ, Folkertsma P,

Gillett T, van der Velde KJ, Kanninga R, van den Akker PC, Jan SZ, Hoorntje ET, te Rijdt

WP, Vos YJ, Jongbloed JDH, van Ravenswaaij-Arts CMA, Sinke R, Sikkema-Raddatz B,

Kerstjens-Frederikse WS, Swertz MA, Franke L. Improving the diagnostic yield of exome-

sequencing by predicting gene–phenotype associations using large-scale gene expression

analysis. *Nat Commun*. 2019;

31. GEO Accession viewer [Internet]. [cited 2019 Dec 19];Available from:

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE1145

32. Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles G V., Clark NR, Ma'ayan A. Enrichr:

Interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics*.

2013;

33. Otsuji TG, Minami I, Kurose Y, Yamauchi K, Tada M, Nakatsuji N. Progressive maturation in

contracting cardiomyocytes derived from human embryonic stem cells: Qualitative effects on

electrophysiological responses to drugs. *Stem Cell Res* [Internet]. 2010;4:201–213. Available

from: https://www.ncbi.nlm.nih.gov/pubmed/20199896

34. Kuppusamy KT, Jones DC, Sperber H, Madan A, Fischer KA, Rodriguez ML, Pabon L, Zhu

WZ, Tulloch NL, Yang X, Sniadecki NJ, Laflamme MA, Ruzzo WL, Murry CE, Ruohola-

Baker H. Let-7 family of microRNA is required for maturation and adult-like metabolism in

stem cell-derived cardiomyocytes. *Proc Natl Acad Sci U S A* [Internet]. 2015;112:E2785-94.

Available from: https://www.ncbi.nlm.nih.gov/pubmed/25964336

35. Gaborit N, Le Bouter S, Szuts V, Varro A, Escande D, Nattel S, Demolombe S. Regional and

tissue specific transcript signatures of ion channel genes in the non-diseased human heart. *J

Physiol* [Internet]. 2007;582:675–693. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/17478540

36. Devalla HD, Schwach V, Ford JW, Milnes JT, El-Haou S, Jackson C, Gkatzis K, Elliott DA,

Chuva de Sousa Lopes SM, Mummery CL, Verkerk AO, Passier R. Atrial-like cardiomyocytes

from human pluripotent stem cells are a robust preclinical model for assessing atrial-selective

pharmacology. *EMBO Mol Med* [Internet]. 2015;7:394–410. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/25700171

37.   Klattenhoff CA, Scheuermann JC, Surface LE, Bradley RK, Fields PA, Steinhauser ML, Ding

H, Butty VL, Torrey L, Haas S, Abo R, Tabebordbar M, Lee RT, Burge CB, Boyer LA.

Braveheart, a long noncoding RNA required for cardiovascular lineage commitment. *Cell*

[Internet]. 2013;152:570–583. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/23352431

38.   Guo X, Xu Y, Wang Z, Wu Y, Chen J, Wang G, Lu C, Jia W, Xi J, Zhu S, Jiapaer Z, Wan X,

Liu Z, Gao S, Kang J. A Linc1405/Eomes Complex Promotes Cardiac Mesoderm

Specification and Cardiogenesis. *Cell Stem Cell*. 2018;22:893-908.e6.

39.   Ounzain S, Micheletti R, Arnan C, Plaisance I, Cecchi D, Schroen B, Reverter F, Alexanian

M, Gonzales C, Ng SY, Bussotti G, Pezzuto I, Notredame C, Heymans S, Guigo R, Johnson R,

Pedrazzini T. CARMEN, a human super enhancer-associated long noncoding RNA controlling

cardiac specification, differentiation and homeostasis. *J Mol Cell Cardiol* [Internet].

2015;89:98–112. Available from: https://www.ncbi.nlm.nih.gov/pubmed/26423156

40.   Maroli G, Braun T. The long and winding road of cardiomyocyte maturation. *Cardiovasc Res*.

2020;

41.   Viswanathan SK, Sanders HK, McNamara JW, Jagadeesan A, Jahangir A, Tajik AJ,

Sadayappan S. Hypertrophic cardiomyopathy clinical phenotype is independent of gene

mutation and mutation dosage. *PLoS One*. 2017;

42.   Yang KC, Breitbart A, De Lange WJ, Hofsteen P, Futakuchi-Tsuchida A, Xu J, Schopf C,

Razumova M V., Jiao A, Boucek R, Pabon L, Reinecke H, Kim DH, Ralphe JC, Regnier M,

Murry CE. Novel Adult-Onset Systolic Cardiomyopathy Due to MYH7 E848G Mutation in

Patient-Derived Induced Pluripotent Stem Cells. *JACC Basic to Transl Sci*. 2018;

43.   Zhou W, Bos JM, Ye D, Tester DJ, Hrstka S, Maleszewski JJ, Ommen SR, Nishimura RA,

Schaff H V, Kim CS, Ackerman MJ. Induced Pluripotent Stem Cell-Derived Cardiomyocytes

from a Patient with MYL2-R58Q-Mediated Apical Hypertrophic Cardiomyopathy Show

Hypertrophy, Myofibrillar Disarray, and Calcium Perturbations. *J Cardiovasc Transl Res*

[Internet]. 2019;Available from: https://www.ncbi.nlm.nih.gov/pubmed/30796699

44.   Greenberg MJ, Watt JD, Jones M, Kazmierczak K, Szczesna-Cordary D, Moore JR.

Regulatory light chain mutations associated with cardiomyopathy affect myosin mechanics

and kinetics. *J Mol Cell Cardiol* [Internet]. 2009;46:108–115. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/18929571

45.   Kampourakis T, Ponnam S, Irving M. Hypertrophic cardiomyopathy mutation R58Q in the

myosin regulatory light chain perturbs thick filament-based regulation in cardiac muscle. *J*

Mol Cell Cardiol [Internet]. 2018;117:72–81. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/29452157

46.   Cohn R, Thakar K, Lowe A, Ladha FA, Pettinato AM, Romano R, Meredith E, Chen YS,

Atamanuk K, Huey BD, Hinson JT. A Contraction Stress Model of Hypertrophic

Cardiomyopathy due to Sarcomere Mutations. *Stem Cell Reports*. 2019;

47.   Mosqueira D, Mannhardt I, Bhagwan JR, Lis-Slimak K, Katili P, Scott E, Hassan M,

Prondzynski M, Harmer SC, Tinker A, Smith JGW, Carrier L, Williams PM, Gaffney D,

Eschenhagen T, Hansen A, Denning C. CRISPR/Cas9 editing in human pluripotent stemcell-

cardiomyocytes highlights arrhythmias, hypocontractility, and energy depletion as potential

therapeutic targets for hypertrophic cardiomyopathy. *Eur Heart J*. 2018;

48.   Brodehl A, Ebbinghaus H, Deutsch MA, Gummert J, Gärtner A, Ratnavadivel S, Milting H.

Human induced pluripotent stem-cell-derived cardiomyocytes as models for genetic

cardiomyopathies. Int. J. Mol. Sci. 2019;

49.   Liu J, Li Y, Lin B, Sheng Y, Yang L. HBL1 Is a Human Long Noncoding RNA that

Modulates Cardiomyocyte Development from Pluripotent Stem Cells by Counteracting MIR1.

Dev Cell [Internet]. 2017;42:333-348 e5. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/28829943

50.   Zhang Y, Jiao L, Sun L, Li Y, Gao Y, Xu C, Shao Y, Li M, Li C, Lu Y, Pan Z, Xuan L, Zhang

Y, Li Q, Yang R, Zhuang Y, Zhang Y, Yang B. LncRNA ZFAS1 as a SERCA2a inhibitor to

cause intracellular Ca 2+ overload and contractile dysfunction in a mouse model of myocardial

infarction. *Circ Res*. 2018;

51. Li B, Hu Y, Li X, Jin G, Chen X, Chen G, Chen Y, Huang S, Liao W, Liao Y, Teng Z, Bin J.

Sirt1 antisense long noncoding rna promotes cardiomyocyte proliferation by enhancing the

stability of sirt1. *J Am Heart Assoc*. 2018;

52. Ishii N, Ozaki K, Sato H, Mizuno H, Susumu Saito, Takahashi A, Miyamoto Y, Ikegawa S,

Kamatani N, Hori M, Satoshi Saito, Nakamura Y, Tanaka T. Identification of a novel non-

coding RNA, MIAT, that confers risk of myocardial infarction. *J Hum Genet*. 2006;51:1087–

1099.

53. Zhang G, Dou L, Chen Y. Association of long-chain non-coding RNA MHRT gene single

nucleotide polymorphism with risk and prognosis of chronic heart failure. *Medicine*

(Baltimore). 2020;

54. Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Gliedt TP, Boehnke M, Abecasis

GR, Willer CJ, Frishman D. LocusZoom: Regional visualization of genome-wide association

scan results. In: Bioinformatics. 2011.

 FIGURE LEGENDS

Figure 1. Single cells RNA-seq analysis identifies cardiac enriched lncRNAs in week 2-12 hESC-

CM. **(A),** A schematic of the experimental plan. Differentiation of hESC to CM was performed over 7

days under insulin-free conditions and with the addition of small molecule CHIR99012 at day 0 and

IWP2 at day 3. Additional media change was performed on days as denoted (*). 36 cells were harvested

on week 2 (W02), 60 cells on week 6 (W06) and 56 cells on week 12 (W12). Immunofluorescence

staining for cardiac alpha actinin (ACTN2, red) mark sarcomeres, and green represents *MYH6*-GFP

reporter signal. **(B),** Violin plots showing the overall levels of gene expression within each gene module.

Each of the six modules (early-1, early-2, early-3, midstage, late-1, late-2) shows either overall increase

or decrease gene expression over time. Black lines track the median expression values across the time

points. **(C),** Heatmap showing the expression of all genes in the six modules with genes in columns.

Cells from each time point and replicates are in rows. **(D),** Histogram showing top 10 biological

processes of genes in the "midstage" blue module. **(E),** Plot showing the connectivity (KME values) of

lncRNAs associated with the other protein coding genes in the "midstage" module. Highly connected

lncRNAs with KME values closest to 1 represent those most highly co-regulated with other genes in

the module. Five highly co-regulated lncRNAs that are highly expressed in muscle heart are indicated.

RP1-46F2.2 (we call *VENTHEART, VHRT*; on chromosome 12) is annotated in red. **(F),** Plot showing

the expression of all "midstage" module genes (expression z-scores) in W12 hES-CMs ranked from 1

to 56 according to *VHRT* expression (red line). *VENTHEART* is correlated with the expression with core

cardiac genes *MYL2*, *MYH7*, *ATP2A2,* and other genes of the "midstage" module. Lines are fitted linear

for z-score values.

Figure 2. Characterisation of *VENTHEART* expression in human hES-CM. (A), Schematic

representing the 6 *VHRT* isoforms annotated on NONCODE and ENSEMBL. Top panel, red boxes

represent exons. Exon 1 is shared by all isoforms. Lower panel shows normalized FPKM peaks from

single cell RNA-seq for W02 (red), W06 (yellow) and W12 (blue), reflecting expression abundance of

VHRT at each time point. **(B),** Relative quantification of specific *VHRT* isoforms expression. Data are

represented as mean ± s.e.m., n=3; * *p*- value < 0.05 are represented; n.s.: not significant. Student's t-

test. **(C),** Graphs showing nuclear and cytoplasmic fractions. *GAPDH* and *18s* used as positive controls

for cytoplasmic compartment, and nuclear compartment with *U6* and *MALAT1*. Expression are shown

as percentage relative to nuclear fraction. n=3 biological replicates. * represents $p \leq 0.05$. Student's

paired t-test with a two-tailed distribution. **(D),** *VHRT* expression is present only in CM, but absent in

non-CM. Specific markers were used for the different cell types: *PDGFRA* for cardiac fibroblast (hCF),

TAGLN for smooth muscle cells (hCASMC), *MYL2* and *MYH7* for hESC-CM and *CDH5* for endothelial

cells (hCAEC). Data are represented as mean ± s.e.m., n=3. ** *p*-value < 0.01. Student's t-test.

Figure 3. CRISPR-mediated *VENTHEART* deletion results in impaired CM specification. (A),

Schematic showing CRISPR/eSpCas-9-mediated *VHRT* knockout (KO). First hESC KO line (*VHRT*

KO #1) was generating using sgRNA_2 and sgRNA_4, and the second (*VHRT* KO #2) using sgRNA_6

and sgRNA_10. **(B),** RT-qPCR validating loss of *VHRT* expression in *VHRT* KO #1 and #2. Expression

is represented as fold change relative to wild type (WT). **(C),** Significant downregulated expression of

cardiac genes *MYH7*, *MYL2* and *ATP2A2* in W12 *VHRT* KO cells. **(D),** Representative immunostainig

showing sarcomere disorganization for *VHRT* KO #1 and #2, compared to WT in W12 hES-CM. **(E-**

G), Distribution of gene expression log fold change (logFC), comparing between GFP+ cells in WT

and *VHRT* KO #1 for W02 **(E)**, W06 **(F)** and W12 **(G).** Differentially expressed (DE) genes at each

time point are highlighted in dark-grey. Non-DE genes in light-grey. Highlighted in blue are cardiac

development and maturation genes from the "midstage" module in Figure 1D, significantly

downregulated in W12. Orange are ectoderm and neuronal genes, significantly upregulated in W12.

Data are represented as expression fold change ± s.e.m., n=3. ** *p*- value < 0.01, ****p*- value < 0.001,

n.s.: not significant. Student's paired t-test with a two-tailed distribution.

Figure 4. *VENTHEART* knockdown decreases core cardiac gene expression and disrupts CM

function. (A), Schematic showing targeting GapmeRs. **(B),** RT-qPCR validation for *VHRT* knockdown

(KD) in W06 hESC-CM, transfected with GapmeR#1 (n=5), GapmeR#2 (n=3) or GapmeR non-

targeting control (CTR, n=5), and harvested for analysis after 4 days. **(C),** Volcano plot showing DE

genes for *VHRT* GapmR#1 (left panel) and *VHRT* GapmeR#2 (right panel). Highlighted in red are

significant DE genes. Non-DE genes are marked in grey. In black are annotated cardiac contraction, calcium signaling and cardiac development genes significantly downregulated by *VHRT* KD. **(D),** Histogram showing top biological process downregulated after *VHRT* GapmeR-mediated KD. **(E),** Significant downregulation of calcium handling genes *RYR2*, *SLC8A1* and validated by RT-qPCR. Expression are shown as relative expression to control (CTR). n=3 biological replicates. *p ≤ 0.05, ** p ≤ 0.01 and ***p ≤ 0.001. Student's paired t-test with a two-tailed distribution. **(F),** Immunostainig showing reduced ryanodine receptor 2 (RYR2) protein expression in *VHRT* GapmeR#1 KD and #2 KD, compared to CTR, in W06 hES-CM. **G**), Impedance-based contractility profile analysed using the CardioExcyte (Nanion). Reduction in relaxation velocity, upstroke velocity, amplitude and prolonged pulse width were observed for *VHRT* siRNA, similar to *ATP2A2* siRNA as positive control.

Figure 5. **Genetic evidence of a role for *VENTHEART* in heart disea**se. **(A),** LocusZoom[54] of 1-MB genomic region harbouring *VHRT* depicting significant *VHRT* cis-*eQTL* SNPs in heart left ventricle (HLV). **(B),** SNPs associated with coronary artery disease (CAD), myocardial infarction (MI), non-ischemic HF and atrial fibrillation (AF) from published GWAS[27–31]. A tightly linked cluster of SNPs (r2>0.8) located at and near the *VHRT* locus shows suggestive association with HF (*P*-value< 5e-5). The same SNPs are significantly associated with *VHRT,* but not *MYL2,* cardiac expression suggesting that *VHRT* rather than *MYL2* is the likely mediator of the observed cardiac disease GWAS signals. **(C),** *VHRT* in cardiac muscle contraction and disease based on co-expression gene network analysis using RNA-seq data from 31,499 human samples[30]. *P*-value < 0.05, Wilcoxon Rank-Sum Test. **(D),** Decreased *VHRT* expression in idiopathic dilated cardiomyopathic failing hearts, compared to healthy controls, based on a re-analysis of microarray data from GEO study GSE1145[31]. Benjamini-Hochberg adjusted *P*-value < 0.05 from univariate analysis with moderated t-statistic using the R limma package, R v3.2 and limma v3.26.0.

Figure 6. Transcriptome comparison of *VHRT* KD and human failing hearts proposes the gene program mechanism for disease causality. (A), Heatmap comparing *VHRT* KD DE genes with genes

739   that are differentially expressed in a panel of DCM failing hearts (GSE141910) (adjusted *P*-

740   value<0.05). **(B)**, KEGG pathway and (c) Gene Ontology analysis by Enrichr[32] (log10 (p-value) ≥ 1.3,

741   *p*-value ≤ 0.05) identifies cardiac contractile and calcium regulatory genes consistently downregulated

742   in *VHRT* KD and DCM ("Down/Down").

DETAILED MATERIAL AND METHODS

Stem cell maintenance and differentiation

Human embryonic stem cell line H1 was maintained using mTeSR medium (Stemcell Technologies, 85850) on 1:200 growth factor-reduced Geltrex (Thermo fisher, A1413202) coated tissue culture plates and passaged regularly as cell aggregates every 4-5 days using ReLeSR (Stemcell Technologies, 05872), an enzyme-free dissociation reagent specific for human pluripotent stem cells). Two days prior to starting differentiation, cells were dissociated using Accutase (Stemcell Technologies, 07922) and seeded as single cells in Geltrex -coated 12-well plates (passage ratio 1:2, between 500'000-600'000 cells). Differentiation was performed following the published protocol by Lian et al.[21] with modifications as follows. 10 μM of CHIR99021 (Stemcell Technologies, 72054) was added on day 0 and left for 24 hours followed by medium change. On day 3, 5uM IWP2 (Sigma Aldrich, I0536) was added using 50/50 mix of new fresh medium and conditioned medium collected from each well and left for 48 hours. Culture medium from day 0 until day 7 was RPMI1640 (HyClone, SH30027.01) plus B-27 serum-free supplement without insulin (Gibco, A1895601). From day 7 and onwards RPMI1640 with B-27 serum free supplement with insulin (Gibco, 17504044) was used and changed every 2-3 days.

Generation of *MYH6*-GFP reporter line

EGFP cassette with kanamycin selection was inserted into BACs for *MYH6* (RP11-834J17, BacPac) immediately before the initiating Methionine (ATG) using recombineering (Quick & Easy BAC Modification Kit, KD-001, Gene Bridges GmbH). The Tol2 transposon cassette with Ampicilin selection mark was inserted into the loxp site of the BAC in the backbone using recombineering. Ten million H1 cells were cultured in CF1 conditioned medium (20% KO serum replacement, 1 mMl-glutamine, 1% non-essential amino acids, 0.1 mM 2-mercaptoethanol and 8 ng ml^{-1} of basic fibroblast growth factor in DMEM:F12) for 6 days and dissociated into single cells with TrypLE™ Express (Thermo Fisher, 12604021) and electroporated with 20 micrograms of Tol2 transposes and 100 micrograms of Tol2/EGFP modified Transposon-BACs. After electroporation, cells were re-suspended in conditioned medium with

10 μM ROCK inhibitor Y276329 (Y27632 (Stemcell technology, 72302). ROCK inhibitor was added for the first 48 hours after electroporation. Fifty μg/ml geneticin (Gibco, 10131035) was added for selection of positive clones 72 hours post-electroporation. Fourteen days later after drug selection, single colonies were picked into 24 well plates for expansion. Fluorescent *in situ* hybridyzation (FISH) using non-modified BACs as probes was carried out to validate the incorporation of BAC construct into genome of ES cells (Cytogenetics Services, Genome Institute of Singapore). Karyotyping was performed to confirm a normal chromosome pattern.

Tri-lineage differentiation

STEMdiff™ Trilineage Differentiation Kit (Stemcell Technologies, 05230) was used to differentiate hESC-MYH6 reporter into the three germ layers: ectoderm (*NESTIN* and *PAX6*), mesoderm (*TBXT* and *CXCR4*) and endoderm (*SOX17*, *FOXA2* and *CXCR4*) following manufacturer's instructions. Differentiation was performed on a 24-well plate. At the end of the differentiation RNA was extracted and RTqPCR was run for respective lineage markers.

Immunofluorescent staining

Cells were fixed in 3.7% formaldehyde for 15 min at room temperature and stored in DPBS. They were permeabilized in 0.2% Triton X-100 for 15 min followed by a pre-blocking step with 2% BSA for 20 min. Primary antibody incubation was performed in DPBS + 10% goat serum (except for Nkx2.5 for which donkey serum was used) overnight at 4 degree and secondary antibody incubation for 2 hours at room temperature. DAPI was included during the final washing step. Antibodies used were cardiac troponin T (Lab Vision, ms-295-P0, mouse, 1:500 dilution), α-actinin (Sigma Aldrich, A7811, mouse, 1:1000 dilution), Nkx2.5 (Santa Cruz, sc-8697, goat, 1:200 dilution), MLC2v (ProteinTech, 10906-1-AP, rabbit, 1:200 dilution), ryanodine receptor (Thermo Fisher, MA3-925, mouse, 1:200), alexa fluor 594 goat-anti-mouse (Thermo Fisher, A-11020), alexa fluor 546 goat-anti-rabbit (Thermo Fisher, A-1107), alexa fluor

647 mouse-anti-goat (Thermo Fisher, A-21237).To improve signal and avoid unspecific binding, MLC2v was stained using cells that were fixed in ice cold methanol for 5 mins at 4 °C.

RNA and DNA isolation

RNA was extracted using Direct-zol™ RNA MiniPrep Kit (Zymo, R2060). Cells were directly lysed using Trizol reagent (Thermo Fisher, 15596026). DNA was purified using PureLink Genomic DNA Mini Kit (Thermo Fisher, K182001). All experiments were performed following the manufacturer's instructions.

PCR and reverse transcription quantitative PCR (RT-qPCR)

DNA or plasmid vector were PCR amplified using Q5 High-Fidelity 2X Master Mix (Bio Labs, M0492S) and target specific primers (IDT) following manufacturer's instructions. RNA (50-500 ng) was reverse transcribed to cDNA using qScript Flex cDNA Kit (Quantabio, 95049-025) with a combination of random primers and oligo (dT). The remaining cDNA (1:10) was mixed with PerfeCTa SYBR Green FastMix, low ROX (Quantabio, 95074-05K) and specific primers on a 384-well plate. Real time qPCR was run using ViiA 7 Real-Time PCR System (Thermo Fisher). Average Cq was recorded and ΔΔCq method was used to calculate relative gene expression changes. Expression levels of genes were normalized against two housekeeping genes, *GAPDH* and *PPIA*. List of primers and sequences used in this study are listed on **Supplementary Table 11.**

Single cells RNA-seq

Single cell preparation and cDNA preparation for RNA-seq

hESC-CM were dissociated into less dense monolayers > 3 days prior to RNA isolation by pre-treatment with 1 mg/ml collagenase IV for 1 hour at 37 ℃ followed by 5 minutes treatment with Accutase (Stemcell Technologies, 07922). Immediately prior to single cell capture, cells were dissociated using Accutase (Stemcell Technologies, 07922) for 4-5 minutes at 37 ℃ and re-suspended with a P1000 pipette to facilitate proper single cell formation. Cells were filtered using a 40 µM filter and kept in suspension in RPMI1640

with B-27 supplement and 5 uM ROCK inhibitor Y276329 (Stemcell technologies, 72302) inhibitor on ice and counted using the TC20 automated cell counter from Biorad. The cell size ranged between 16-22 µM and cell viability was above 95%. Fluidigm microfluidic C1 system was used for automated cell capture, conversion of polyA+ RNA into cDNA and a universal amplification step. After priming of the chip following Fluidigm manual, cells were loaded onto large sized C1 IFC chips (Fluidigm, 100-5761) at a concentration of 400 cells/ul (total volume of 20 ul with a 3:2 ratio of cell suspension:C1 suspension reagent). The mRNA Seq: Cell Load script was run followed by manual inspection to assess cell capture success. Pictures were taken from all 96 capture sites and a note was made whether 0, 1 or more cells were captured per site and if the captured cell was GFP negative or positive. Cell lysis, reverse transcription and PCR amplification was run overnight (using reagents from the SMARTer Ultra Low RNA Kit from ClonTech, 634833) and the cDNA products harvested and diluted in 10 µl of C1 DNA Dilution Reagent (from the C1 Reagent Kit for mRNA seq from Fluidigm, 100-6201) the next morning.

Library preparation and Quality Control

cDNA concentration was measured using the Quant-iT™ PicoGreen® dsDNA Assay Kit (LifeTech, P11496) and ranged from 0.1-1 ng/ul per cell. 3µl of cDNA was used for library preparation with the Nextera XT DNA Sample Preparation Kit (Illumina, FC-131-1096). Each library was barcoded and multiplexed 15 libraries into one sequencing lane. A total of 71 GFP positive (CM) and 31 GFP negative (non-CM) cells were sequenced on an Illumina HiSeq 2000 sequencer in 4 different sequencing runs. From the 71 GFP positive cells, 4 were duplicated in one run to control for technical variability. 3 samples (one from each time point) that were first ran in individual runs were subsequently re-sequenced together in the fourth run to control for batch effects. Three samples (one from each time point) also underwent a new round of library preparation and were sequenced together with the original library prepared sample to assess the influence of the library preparation step on the outcome.

Raw reads filtering, alignment, quality control and gene expression estimation

Illumina CASAVA version 1.8.2 was used to perform demultiplexing and generate FASTQ file. Quality of sequencing reads was examined with FASTQC[1]. Paired-end reads were aligned to human genome (*Homo*

Sapiens) hg19 assembly using state-of-the-art mapping tool, Tophat2[2] (version 2.0.11.Linux_x86_64) with Bowtie2 (version 2.2.1.0). The parameters that were set for Tophat2 alignment were –min-anchor 8, --splice-mismatches 1, --min-intron-length 50, --max-intron-length 500000, --min-isoform-fraction 0.15, --max-multihits 1, --segment-length 25, --segment-mismatches 2, --min-coverage-intron 50, --max-coverage-intron 20000,--min-segment-intron 50, --max-segment-intron 50000, --read-mismatches 3, --read-gap-length 3, --read-edit-dist 3, --read-realign-edit-dist 0, --max-insertion-length 3, --max-deletion-length 3, --mate-inner-dist 200, --mate-std-dev 20, --no-coverage-search and –library-type fr-unstranded. We also suppliedTophat2 with hg19 GENCODE (version 19) transcriptome annotation to Tophat2 with –G option. RSeQC (RNA-seq Quality Control Package version 2.6.1) was used to inspect sequence quality, transcript integrity number (TIN) which is analogous to RIN (RNA integrity number), duplication rate and mapped reads distribution[3]. Reads mapped to mitochondrial genes and ribosomal genes were also calculated using htseq-count (version 0.6.0).

Cuffdiff (version 2.2.1, with boost version 104700) was used to compute relative gene expression level in human genome (hg19) GENCODE (version 19) transcriptome annotating tf file with parameters: --library-norm-method classic-fpkm, --compatible-hits-norm, --frag-bias-correct and --max-frag-multihits 1. Ribosomal RNA and tRNA were masked in the gene expression level calculation. Gene expression level was reported in fragments per kilobase per million mapped reads (FPKM)[4]. We assumed genes with FPKM lower than 1 to be non-expressing. Only genes expressed with FPKM $\geq$ 1 in at least 2 samples were considered for downstream analyses.

Saturation Analysis

Using samtools version 0.1.19, saturation analysis was performed by randomly subsampling different number of reads from each sample, and re-calculating gene FPKM. The process of subsampling was repeated until there were at least 5 subsampled datasets per point, at library size of (0.1M, 0.5M, 1M, 2M, 3M, 4M, 5M). For each point, the average number of genes with FPKM greater than or equal to 1 was plotted.

Network construction and module detection using R package Weighted Correlation Gene Network Analysis (WCGNA).

Using WCGNA, we constructed a signed weighted correlation network by computing pairwise correlations, s between all genes across all single-cell RNA-seq samples[5]. Next, we chose soft thresholding power (β=3), in constructing an adjacency matrix using the formula, $a_{ij} = (0.5 + 0.5 \times s_{ij})^{\beta}$, where a_{ij} is defined as weighted correlation and s_{ij} is defined coefficient correlation between betweengene$_i$ and gene$_j$. We choose the power (β=3), which was the lowest power for which the scale-free topology fit index curve flattens out upon reaching a high value of 0.98. Using the adjacency matrix computed in the previous step, topological overlap was calculated to measure the network interconnectedness in a robust and biological meaningful way. The topological overlap was utilized to group highly correlated genes together using average linkage hierarchical clustering.

Modules were defined as the branches obtained by cutting the hierarchal tree using Dynamic Hybrid Tree Cut algorithm[6]. We defined the first principle component of a module as module eigengene, which was representative of the expression profile in each module. Genes in each module were removed if the correlation between the gene and module eigengene (kME) was less than 0.3. If a detected module did not have at least 30 genes with eigengene connectivity (kME) at least 0.5, the module was disbanded and its genes were unlabeled and returned to the pool of genes waiting for module detection. Modules whose eigengenes were highly correlated (correlation above 0.75) were merged. Construction of signed gene network and identification of modules was performed using R function, blockwise Modules with following parameters: soft thresholding power = 3, minimum module size = 30, mergeCutHeight = 0.25, corType = "Pearson", networkType = "signed", TOMType = "signed", minCoreKME = 0.5, minKMEtoStay = 0.3.

To identify modules that were significantly correlating with subgroups, we computed correlation of eigengenes of each module with cells in W02, W06 and W12. We picked modules which showed high correlation with each week and sub-population of cells for W12. In addition, we also computed Gene Significance (GS), which was defined as the correlation of each gene with each group of interest[5]. We also

calculated module membership, which was defined as the correlation between module eigengene and gene expression profile. GS and MM were important because it helped in the identification of genes with high significance for each week of development and high module membership in each week-/subtype-specific modules. Module membership was reported to be highly correlated to the intramodular connectivity, k_{ME}. Highly connected intramodular hub genes was observed to have high module membership values to the respective module.

Bulk RNA seq analysis

Total RNA sequencing was performed using Truseq Stranded Total RNA Library Prep kit (Illumina, RS-122-2201), which uses Ribo-Zero to remove cytoplasmic rRNA from total RNA. Remaining intact RNA is fragmented using a chemical mix, followed by first- and second-strand cDNA synthesis using random hexamer primers. "End-repaired" fragments were ligated with unique Illumina adapters. All individually indexed samples were subsequently pooled together and multiplexed for sequencing. Libraries were sequenced using the Illumina Hiseq 2000 sequencing system and paired-end 101 bp reads were generated for analysis. Quality control, gene alignment and gene expression estimation were performed as described in the single cell sequencing method.

GWAS and eQTL analysis and statistics

Genome-wide association studies (GWAS) summary statistics were accessed from the Cardiovascular Disease Knowledge Portal (CVDKP)[7], and included the following data sets: All-cause Heart Failure and Nonischemic Cardiomyopathy HRC GWAS in UK Biobank (394156 samples, European ancestry)[8], CARDIoGRAMplusC4D GWAS (184305 samples, mixed ancestry)[9], CAD exome chip analysis (120575 samples, mixed ancestry)[10] and 2018 AF HRC GWAS (588190 samples, mixed ancestry)[11]. Significant VHRT cis-eQTLs (FDR<0.05, +/- 1Mb window around the VHRT TSS) precalculated from genotype and RNA sequencing data from human heart left ventricular samples (n=386) were extracted from the Genotype-Tissue Expression (GTEx) Portal[12]. VHRT locus plots depicting GWAS and eQTL data were

created using LocusZoom[13] on genome build GRCh37/hg19 and LD population 1000 Genomes Phase 3, EUR.

Cross-interrogation of *VHRT* KD transcriptome with human DCM date sets

Significant DE genes in GapmeR#1 VHRT KD and GapmeR#2 VHRT KD vs CTRL (N=1,100; 573 down- and 527 upregulated at adjP-value<0.05, FC>|0.5| in both) were compared with public DCM transcriptomics study (GEO accession: GSE141910). This study comprises RNA sequencing datasets of 61 left ventricle explants (Idiopathic Dilated Cardiomyopathy and Non-Failing) from the MAGNet study (Myocardial Applied Genomics Network). Differential expression analysis was conducted on the DCM versus control samples using R package EdgeR, which applies generalized linear models (glm) on normalized RNA-seq gene count. Enrichment for pathway and molecular functions was then performed on the DE subgroup based on functional analysis using Enricher tools[14].

***VHRT* CRISPR/Cas9 knockout**

Plasmid pMIA3 (Addgene plasmid # 109399; http://n2t.net/addgene:109399; RRID: Addgene_109399) was used for CRISPR/Cas9 mediated KO. pMIA3 was generated and previously described by Dr. Matias Autio[15]. Two KO hESC lines were generated using dual single guide RNA (sgRNAs) to remove part of VHRT. Single guide RNA sequence were designed using cloud based software tools for digital DNA sequence editing Benchling[16] and CRISPOR[17,18]. Schematic representation of the pairs of sgRNAs and genomic regions to be removed are shown in the figure 5A. 10 µM sense and anti-sense oligonucleotides were ordered **(Supplemental Table 11)** and annealed to generate the 20 nucleotide spacer that defines the genomic target to be modified (VHRT 5'end and 3'end). pMIA3 was digested with BsmBI and sgRNA spacers cloned after human U6 promoter with T4 DNA ligase (New England Biolabs) following manufacturer's instruction. Ligated constructs underwent transformation using RapidTrans™ Chemically Competent Cells (Active motif). Plasmid was extracted from bacterial culture, purified and Sanger sequenced to confirm successful cloning. Prior to hESC targeting, cutting efficiency of pMIA3-sgRNAs

plasmids were tested on HEK293T using the EGxxFP plasmid (pCAG-EGxxFP was a gift from Masahito Ikawa, Addgene plasmid # 50716). Best combination of sgRNAs was then used for final targeting of hESCs. The best pair of guides were cloned into a single pMIA3 plasmid digested with NheI & XbaI, using NEBuilder isothermal assembly (New England Biolabs), according to manufacturer's instructions to make the final pMIA3 dual sgRNA plasmid. Primers for isothermal assembly are listed on **Supplemental Table 11.** Human ESCs were dissociated with Accutase (Stemcell Technologies, 07922) and ~1.5×10^6 cells were re-suspended in 100 μl P3 Primary Cell kit solution from Lonza (V4XP-3024) and mixed with 10 μg pMIA3dual sgRNA plasmid. To transfect hESC, nucleofection was performed using program CM-113 on the 4D-Nucleofector System (Lonza). Cells were then plated into Geltrex coated 6-well plate with mTeSR medium (Stemcell Technologies, 85850) containing 5 μM Y-27632. After 2 days in culture, cells were dissociated, FACS sorted for RFP positive cells and collected into a tube containing TeSR medium (Stemcell Technologies, 85850) with 5 μM Y-27632. 500 to 2000 cells were plated into wells of 6-well plate containing the above media. Single cell clones were monitored and upon sizeable growth, colonies were picked and passaged. Genomic DNA was extracted for genotyping and screened for successful KO. Location of primer pairs used for the genotyping are represented in **Supplemental Figure 7A-C** and primer sequences are listed in **Supplemental Table 11**. RT-qPCR was performed to validated KD of *VHRT* transcript.

Cellular fractionation

An average of 10^7 hESC-CMs were dissociated into single cell suspension as described above. RNA was extracted from nuclear and cytoplasmic fractions using a rapid, phenol-free, small scale protein and RNA isolation system PARIS kit (Thermo Fisher, AM1921) following manufacturer's instructions. During the fractionation procedure, Trypan blue solution was used to monitor proper separation of the cytoplasmic and nuclear fractions (**Supplemental Figure 5B**). To remove traces of genomic DNA, eluted RNA was treated with DNA-free™ DNase Treatment and Removal Reagents (Thermo Fisher, AM1906) following manufacturer's instructions. RNA concentration and purity were measured using NanoDrop. An Agilent

RNA 6000 Pico kit (5067-1513) was run to observe the RIN value, distribution and intensity of the signal of all RNAs, up to about 7000 nt, hence including the distinctive peak for tRNA, 18S and 28S. *VHRT* compartmentalization was detected with RT-qPCR along with cytoplasmic markers (*GAPDH* and 18S ribosomal RNA) and nuclear markers (*U6* and *MALAT1*).

In-vitro translation

Protein coding potential was assessed via web-based prediction algorithm (CPC: coding potential calculator). An open reading frame (ORF) was identified on *VHRT* isoform 6. 1x FLAG Tag (motif: D YKDDDDK) was inserted either at N-terminus (after the START codon) or the C-terminus (before STOP codon) of the ORF. Plasmids were transfected in HEK293T cells. Successful transfection was confirmed using a plasmid control containing sequence for blue fluorescent protein (BFP: ~29 kDa) tagged with FLAG at the C-terminus. 48 h after transfection, cells were lysed for protein extraction and Western blot. Protein analysis was performed using pre-casted gradient gel 4-20% mini-PROTEAN TGX Stain-Free (Bio-rad, 4568096). 10 to100 μg of protein was loaded onto the gels alongside a protein ladder (1st Base, Asia). The electrophoresis setting for initial running for 20 mins at 50 V and subsequently 100 V for 60-90 mins. The running buffer contained 25 mM Tris base, 192 mM glycine and 0.1 % SDS. Protein transfer was performed on 0.22 μm polyvinylidene diflouride (PVDF, BioRad, 1620260) membrane at 100 V for 1 h in transfer buffer (25 mM Tris base, 192 mM glycine, 20 % methanol). After transfer, the membrane was blocked in 5 % BSA/TBS-T (20 mM Tris-HCl pH 7.6, 140 mM NaCl supplemented with 0.1 % Tween 20, Merck, P1379) for 1 h on a rocking platform at room temperature (RT). Membranes were subsequently incubated with primary anti- mouse FLAG antibody (Sigma Aldrich, B3111) diluted in 5 % BSA/T-BST blocking buffer overnight at 4°C. Membranes were washed 3 times in TBS-T for 5 min. Horse radish peroxidase (HRP)-conjugated secondary antibody (Thermo Fisher, MA5-15367) was diluted in 5 % BSA/T-BST at 1:10 000 dilution and added to the membrane for 1h at RT. After membranes washing with TBS-T, chemiluminescence solution (West Femto, Thermo Fisher, 32106) was added to the membrane following the manufacturer's instructions and exposed to chemidoc (Biorad) for protein exposure.

Human heart tissue collection

Gene expression level was assessed by qPCR from explanted hearts of patients undergoing transplantation for end-stage heart failure. All donors provided written informed consent and samples were fully anonymized. 30-60 mg of cardiac tissue (Ventricle Tissue, VT and Atrial Tissue, AT) were mechanically powdered on liquid nitrogen and further disrupted in 350 µl RLT buffer at room temperature using a Tissue Lyzer device (Qiagen, Hilden, Germany; 2 min, 30 Hz). DNA extraction was performed with the RNeasy Plus kit (Qiagen) according to the manufacturer's instructions. 100-250 ng of RNA were reversely transcribed with the High Capacity cDNA Reverse Transcription kit (Applied Biosystems). qPCR was carried out on an ABI Prism 7900 HT Sequence Detection System (Applied Biosystems). Transcript abundance was calculated using the the delta-delta-CT method with GAPDH as reference transcript.

VHRT knockdown

GapmeR transfection

Anti-sense LNA - GapmeR –based technology (Qiagen) was used for gene KD. With the use of a web-based algorithm provided by Qiagen, GapmeRs were designed to target *VHRT* transcript (RP1-46F2.2, hg19 or LINC01405, hg38). 100 nM of each GapmeR was transfected using 6 ul Lipofectamine 2000 reagent (Invitrogen, 11668030) in a total volume of 1mL hESC-CM media (RPMI 1640 with B27 supplement with insulin, Thermo Fisher Scientific, A1895601) in a 12-well plate with 85-90% cell confluency (n= 4 biological replicates). Negative A–GapmeR was transfected as negative control. Sequence of GapmeRs are listed in **Supplemental Table 11.**

Fresh and new media was replaced after 48 h. Day 4 after transfection, RNA was purified and relative expression of *VHRT* and other important cardiac genes (*MYH6, MYH7, RYR2, ATP2A2, SCN5A* and *PLN*) were assayed via RT-qPCR in each group. Expression levels were normalized to housekeeping genes *GAPDH* and *PPIA*. % KD was calculated relative to Negative A-GapmeR and genes that had significant changes in expression (p-value ≤ 0.05, Student's t test) were considered statistically significant.

SiRNA transfection

In-vitro iCell cardiomyocytes (CDI, 01434) were maintained in culture according manufacturer instruction. SiRNA for *VHRT* was purchased from Riboxxx life science (lot 2017d0113) and for *ATP2A2* from Dharmacon (ON TARGETplus SMARTpool siRNA). They were dissolved in H_2O to a stock concentration of 10 µM**.** 10 nM siRNA was transfected using 0.1 uL Lipofectamine RNAiMax per well. Media was changed after 24h and cells were used in the corresponding assay after 72h. RNA was extracted using the RNeasy micro kit (Qiagen, 74004) in the Qiacube instrument. cDNA was prepared using reverse transcription according to High-Capacity RNA-to-cDNA™ Kit (Thermo Fisher, 4387406). qPCR was performed using TaqMan Fast Advanced Master mix (Thermo Fisher) together with specific gene expression assays Hs04405003_m1 (*RP1-46F2.2*) and housekeeping gene (*GAPDH*). Quantitative RT-PCR was performed with QuantStudio™ 7 Flex Real-Time PCR instrument (Thermo Fisher) and data was recorded during 40 PCR cycles and normalized against untreated samples. Sequence of siRNAs are listed in **Supplemental Table 11.**

Fluorescence-activated cell sorting (FACS) of hESC-CM

Cells were sorted using BD FACSAria II 5 laser from Flow Cytometry Core or (A*STAR, Singapore Immunology Network). Prior dissociation, cells were cultured with corresponding media containing 5uM ROCK inhibitor (Y27632, Stemcell Technologies, 72302) for 30 min at 37∘C. Media was aspirated and cells were treated with Accutase (Stemcell Technologies, 07922) (1mg/mL) for 5-8 min at 37∘C. For hESCs, Accutase (Stemcell Technologies, 07922) was neutralized using mTeSR medium (Stemcell Technologies, 85850) whereas passaging media was used for hESC-CMs. Cells were collected in 15 ml tube and spin at 300 g for 5 min. Human ESC were re-suspended in DPBS and hESC-CMs in FACS buffer (DPBS w/o Ca and w/o Mg, 2% HI-FB, 11550356, + 1mM EDTA, 5uM Y27632) and pipetted through 40 um cell strainer to avoid cell clumps. Due to their increase in cell size over time, older hESC-CMs were pipetted through a 70um cell strainer instead. During FACS, cells were collected in 2mL tube containing

corresponding media and re-plated on a new coated plate (1:200 Geltrex, if hESC or 0.1% gelatin, if hESC-CM). To avoid contamination 100x Penicillin-Streptomycin mixture was added at each step and sample were kept on ice at all time to reduce cell death.

Electrophysiological assays

Multielectrode array recordings

MEA chips were coated with 0.1% gelatin and hESC-CM (8 weeks after differentiation) were seeded on the chip at least 4 days prior to the recording to allow for proper attachment and coupling. Field potentials were recorded from spontaneously beating cells using a USB-MEA60-Up-BC-System (Multichannel systems) and analyzed using MC_Rack. Recordings were made in regular RPMI-1640 (Thermo Fisher, 11875093) +B-27 supplement media (Thermo Fisher, A3582801) with our without the addition of 100 µM noradrenaline.

Contractility and electrophysiology recordings

The Cardioexcyte 96 instrument (Nanion) was used to record and analyze impedance-based contractility. The stim plates were coated with fibronectin for >1h in the incubator. iCell cardiomyocytes (01434 from CDI) were thawed and plated at 20k cells/well following recommendation from the manufacturer. Experiments were performed from day 12 post-plating and onwards. At least 5 replicates were used for each condition. *ATP2A2* was used as positive control and non-targeted siRNA as well as untreated cells as negative controls. Readings for base impedance were measured before and after treatment with siRNAs, both for spontaneously beating and electrically paced cells. The data was analyzed using the data control software from Nanion. One-way ANOVA, multiple comparisons, compared to control siRNA.

Action potential (AP) recordings

Whole cell configuration of the patch-clamp technique was used to measure AP on two different groups of cells, Negative A-GapmeR and *VHRT* KD GapmeR. Both groups were co-transfected with BLOCK-iT™

Alexa Fluor ® Red Fluorescent Control (Invitrogen), therefore only cells marked in red and green (*MYH6-*GFP reporter) were used to record AP. The signal was amplified using an axon 700B patch clamp amplifier (axon instrument, Sunnyvale, USA) and low-pass filtered at 5 kHz. Patch pipettes were fabricated from glass capillaries (O.D, 1.5mm; I.D, 0.9mm) using a Sutter P-97 microelectrode puller (Novato, CA, USA) and the tips were heat polished with a microforge (NARISHIGE MF-900) to gain a resistance of 2-4 MΩ. The electrical signals were sampled at 2.5-10 kHz and filtered at 2 kHz using a low-pass filter. Data acquisition was achieved using the Digidata 1440A (axon instrument). Data analysis and fit were performed using clamp fit 10.2 (axon instrument) and Origin 7.0 software (Origin Lab Corporation). A pClamp software (Version8.1; Axcon Instrument) was used to generate voltage-pulse protocols, acquire and analyze data. APs were recorded under current-clamp mode at 35 ºC. Pipette solution contained (in mM): KCl 130, NaCl 5, MgCl2 1, MgATP 3, EGTA 10, and HEPES 10, with pH adjusted to 7.2 with KOH. Extracellular solution (Tyrode's solution) containing contained (in mM) NaCl 140, KCl 5.4, CaCl2 1.8, MgCl2 1, glucose 10, HEPES 10, with pH adjusted to 7.4 with NaOH. The parameters of APs include AP durations (APD) at 20%, 50% and 90% of repolarization (APD20, APD50, and APD90), AP amplitude (APA), maximal diastolic potential (MDP), and beating frequency (for spontaneous contractions) were analyzed.

Statistics

Excluding the NGS seq results, all gene expression analysis was performed using Excel (Microsoft office). Unless stated otherwise, data are represented as mean ± standard error of the mean (s.e.m). Statistical significance between two groups was done using student's t. P-values of ≤ 0.05 were considered as significant, unless otherwise indicated.

REFERENCES

1.   Anders S, Huber W. Differential expression analysis for sequence count data. *Genome Biol* [Internet]. 2010;11:R106. Available from: https://www.ncbi.nlm.nih.gov/pubmed/20979621

2.   Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol* [Internet]. 2013;14:R36. Available from: https://www.ncbi.nlm.nih.gov/pubmed/23618408

3.   Wang JT, Li JT, Zhang XF, Sun XW. Transcriptome analysis reveals the time of the fourth round of genome duplication in common carp (Cyprinus carpio). *BMC Genomics* [Internet]. 2012;13:96. Available from: https://www.ncbi.nlm.nih.gov/pubmed/22424280

4.   Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* [Internet]. 2012;7:562–578. Available from: https://www.ncbi.nlm.nih.gov/pubmed/22383036

5.   Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* [Internet]. 2008;9:559. Available from: https://www.ncbi.nlm.nih.gov/pubmed/19114008

6.   Langfelder P, Zhang B, Horvath S. Defining clusters from a hierarchical cluster tree: the Dynamic Tree Cut package for R. *Bioinformatics* [Internet]. 2008;24:719–720. Available from: https://www.ncbi.nlm.nih.gov/pubmed/18024473

7.   Cardiovascular Disease Knowledge Portal [Internet]. Available from: http://www.broadcvdi.org/

8.   Aragam KG, Chaffin M, Levinson RT, McDermott G, Choi SH, Shoemaker MB, Haas ME, Weng LC, Lindsay ME, Smith JG, Newton-Cheh C, Roden DM, London B, Wells QS, Ellinor PT, Kathiresan S, Lubitz SA. Phenotypic Refinement of Heart Failure in a National Biobank Facilitates Genetic Discovery. *Circulation*. 2019;

9.   Nikpay M, Goel A, Won HH, Hall LM, Willenborg C, Kanoni S, Saleheen D, Kyriakou T, Nelson

CP, CHopewell J, Webb TR, Zeng L, Dehghan A, Alver M, MArmasu S, Auro K, Bjonnes A, Chasman DI, Chen S, Ford I, Franceschini N, Gieger C, Grace C, Gustafsson S, Huang J, Hwang SJ, Kim YK, Kleber ME, Lau KW, Lu X, Lu Y, Lyytikäinen LP, Mihailov E, Morrison AC, Pervjakova N, Qu L, Rose LM, Salfati E, Saxena R, Scholz M, Smith A V., Tikkanen E, Uitterlinden A, Yang X, Zhang W, Zhao W, De Andrade M, De Vries PS, Van Zuydam NR, Anand SS, Bertram L, Beutner F, Dedoussis G, Frossard P, Gauguier D, Goodall AH, Gottesman O, Haber M, Han BG, Huang J, Jalilzadeh S, Kessler T, König IR, Lannfelt L, Lieb W, Lind L, MLindgren C, Lokki ML, Magnusson PK, Mallick NH, Mehra N, Meitinger T, Memon FUR, Morris AP, Nieminen MS, Pedersen NL, Peters A, Rallidis LS, Rasheed A, Samuel M, Shah SH, Sinisalo J, EStirrups K, Trompet S, Wang L, Zaman KS, Ardissino D, Boerwinkle E, Borecki IB, Bottinger EP, Buring JE, Chambers JC, Collins R, Cupples L, Danesh J, Demuth I, Elosua R, Epstein SE, et al. A comprehensive 1000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet*. 2015;

10.   Stitziel NO, Stirrups KE, Masca NGD, Erdmann J, Ferrario PG, König IR, Weeke PE, Webb TR, Auer PL, Schick UM, Lu Y, Zhang H, Dube MP, Goel A, Farrall M, Peloso GM, Won HH, Do R, Van Iperen E, Kanoni S, Kruppa J, Mahajan A, Scott RA, Willenborg C, Braund PS, Van Capelleveen JC, Doney ASF, Donnelly LA, Asselta R, Merlini PA, Duga S, Marziliano N, Denny JC, Shaffer CM, El-Mokhtari NE, Franke A, Gottesman O, Heilmann S, Hengstenberg C, Hoffmann P, Holmen OL, Hveem K, Jansson JH, Jöckel KH, Kessler T, Kriebel J, Laugwitz KL, Marouli E, Martinelli N, McCarthy MI, Van Zuydam NR, Meisinger C, Esko T, Mihailov E, Escher SA, Alver M, Moebus S, Morris AD, Müller-Nurasyid M, Nikpay M, Olivieri O, Perreault LPL, AlQarawi A, Robertson NR, Akinsanya KO, Reilly DF, Vogt TF, Yin W, Asselbergs FW, Kooperberg C, Jackson RD, Stahl E, Strauch K, Varga T V., Waldenberger M, Zeng L, Kraja AT, Liu C, Ehret GB, Newton-Cheh C, Chasman DI, Chowdhury R, Ferrario M, Ford I, Jukema JW, Kee F, Kuulasmaa K, Nordestgaard BG, Perola M, Saleheen D, Sattar N, Surendran P, Tregouet

D, Young R, Howson JMM, Butterworth AS, Danesh J, Ardissino D, et al. Coding variation in ANGPTL4, LPL, and SVEP1 and the risk of coronary disease. *N Engl J Med*. 2016;

11.   Roselli C, Chaffin MD, Weng LC, Aeschbacher S, Ahlberg G, Albert CM, Almgren P, Alonso A, Anderson CD, Aragam KG, Arking DE, Barnard J, Bartz TM, Benjamin EJ, Bihlmeyer NA, Bis JC, Bloom HL, Boerwinkle E, Bottinger EB, Brody JA, Calkins H, Campbell A, Cappola TP, Carlquist J, Chasman DI, Chen LY, Chen YDI, Choi EK, Choi SH, Christophersen IE, Chung MK, Cole JW, Conen D, Cook J, Crijns HJ, Cutler MJ, Damrauer SM, Daniels BR, Darbar D, Delgado G, Denny JC, Dichgans M, Dörr M, Dudink EA, Dudley SC, Esa N, Esko T, Eskola M, Fatkin D, Felix SB, Ford I, Franco OH, Geelhoed B, Grewal RP, Gudnason V, Guo X, Gupta N, Gustafsson S, Gutmann R, Hamsten A, Harris TB, Hayward C, Heckbert SR, Hernesniemi J, Hocking LJ, Hofman A, Horimoto ARVR, Huang J, Huang PL, Huffman J, Ingelsson E, Ipek EG, Ito K, Jimenez-Conde J, Johnson R, Jukema JW, Kääb S, Kähönen M, Kamatani Y, Kane JP, Kastrati A, Kathiresan S, Katschnig-Winter P, Kavousi M, Kessler T, Kietselaer BL, Kirchhof P, Kleber ME, Knight S, Krieger JE, Kubo M, Launer LJ, Laurikka J, Lehtimäki T, Leineweber K, Lemaitre RN, Li M, Lim HE, et al. Multi-ethnic genome-wide association study for atrial fibrillation. *Nat Genet*. 2018;

12.   GTEx Portal [Internet]. [cited 2019 Nov 9];Available from: https://www.gtexportal.org/home/

13.   Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Gliedt TP, Boehnke M, Abecasis GR, Willer CJ, Frishman D. LocusZoom: Regional visualization of genome-wide association scan results. In: Bioinformatics. 2011.

14.   Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles G V., Clark NR, Ma'ayan A. Enrichr: Interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics*. 2013;

15. Ang LT, Tan AKY, Autio MI, Goh SH, Choo SH, Lee KL, Tan J, Pan B, Lee JJH, Lum JJ, Lim CYY, Yeo IKX, Wong CJY, Liu M, Oh JLL, Chia CPL, Loh CH, Chen A, Chen Q, Weissman IL, Loh KM, Lim B. A Roadmap for Human Liver Differentiation from Pluripotent Stem Cells. *Cell Rep* [Internet]. 2018;22:2190–2205. Available from: internal-pdf://113.214.84.44/1-s2.0-S2211124718301517-main.pdf

16. Cloud-Based Informatics Platform for Life Sciences R&D | Benchling [Internet]. [cited 2019 Dec 2];Available from: https://www.benchling.com/

17. CRISPOR [Internet]. [cited 2019 Dec 2];Available from: http://crispor.tefor.net/crispor.py

18. Concordet J-P, Haeussler M. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids Res* [Internet]. 2018 [cited 2020 Jan 3];46:W242–W245. Available from: https://academic.oup.com/nar/article/46/W1/W242/4995687

19. Deelen P, van Dam S, Herkert JC, Karjalainen JM, Brugge H, Abbott KM, van Diemen CC, van der Zwaag PA, Gerkes EH, Zonneveld-Huijssoon E, Boer-Bergsma JJ, Folkertsma P, Gillett T, van der Velde KJ, Kanninga R, van den Akker PC, Jan SZ, Hoorntje ET, te Rijdt WP, Vos YJ, Jongbloed JDH, van Ravenswaaij-Arts CMA, Sinke R, Sikkema-Raddatz B, Kerstjens-Frederikse WS, Swertz MA, Franke L. Improving the diagnostic yield of exome- sequencing by predicting gene–phenotype associations using large-scale gene expression analysis. *Nat Commun*. 2019;

20. GEO Accession viewer [Internet]. [cited 2019 Dec 19];Available from: https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE1145

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1

(A), Karyotyping of the *MYH6*-GFP reporter cell line shows a normal karyotype. **(B),** FISH identified 1 signal incorporated on each allele of chromosome 14. **(C),** Immunofluorescence confirms protein expression of α-actinin (ACTN1), cardiac troponin T (cTnT) and NKX2-5 in hESC-derived cardiomyocytes (CMs) indicating high differentiation efficiency. Striated patterns with ACTN1and cTnT are typical for cardiac muscle. Overlap with GFP shows that all GFP positive cells expressed all three cardiac proteins. **(D),** A multielectrode array was used to record field potentials from hESC-CM under baseline conditions and upon administration of adrenergic stimulant noradrenaline. Physiological field potentials are as displayed showing that noradrenaline induced an increase in beating frequency, increased field potential amplitude and a shortening of the field potential duration.

Supplementary Figure 2

(A), Single cells captured in the Fluidigm C1 microfluidic system. Top, GFP positive (GFP+) and bottom, GFP negative (GFP-) cells, respectively. **(B),** Saturation plot for all genes expressed at FPKM ≥ 1 shows that for all samples, the maximum number of expressed genes was attained by around 1 million sequencing reads. **(C),** Histoplot showing the number of expressed genes (FPKM >1) detected in each single cell sample at the different time points. **(D),** Scatter plots for correlations between technical repeats show strong reproducibility and excludes any predominant effect of technical noise or batch effect. First, three libraries (one from each time point) were re-sequenced together in one sequencing lane and the results compared to the original sequencing of the same library (i-iii). Second, cDNA from previously sequenced samples were subjected to a new round of library preparation and re-sequenced together with initially prepared libraries (iv). Third, freshly prepared and sequenced libraries were compared to sequencing from the initial corresponding libraries (v). **(E),** Spike-in concentration vs FPKM values show strong linear correlation (r=0.96). Spike-in values in 2 weeks vs 6 weeks samples, 2 weeks vs 12 weeks, and 6 weeks vs 12 weeks

show high technical consistency. **(F),** Scatterplot comparing averaged gene expression in W02 to W06 cells, showing a high correlation r value (0.91), the expected lack of differential expression for stable cardiac genes (green dots: *TNNT2*, *ACTC1*, *TPM1*), but indistinct differential expression of variable cardiac genes (red dots: *MYH6*, *NKX2*-5, *DMD*) at this averaged pooled level of analysis. **(G),** Scatterplot comparing gene expression in a single W06 cell to another single W06 cell, now showing a lower correlation r value (0.6) than (E), the same lack of differential expression of stable cardiac genes (green dots), but instead distinct differential expression of the 3 variable cardiac genes (red dots).

Supplementary Figure 3

(A), Heatmap of cardiac genes that have either (a) constantly high expression in all cells at all-time points, (b) decreasing, or (c) increasing abundance in expression over time, or (d) highly variable expression across all time points. **(B),** Line plots showing mean gene expression across cells for first and second heart field markers. Data are represented as mean FPKM ± s.e.m. **(C),** Immunostaining confirms protein expression at all-time points for cardiac alpha actinin (α-act, top panel) and displaying sarcomere organization (α-act, bottom panel). **(D),** Immunostaining showing MLC2v protein expression appearing only from W06.

Supplementary Figure 4

(A), Module dendrogram showing gene-network modules that were derived using weighted gene co-expression network analysis (WGCNA). Each vertical black line represents a uniquely expressed gene, grouped according to co-expression gene modules (colored bars). Correlation between gene modules and the time points are represented in the 3 horizontal bars (blue for negative-correlation and red for positive-correlation). Six gene modules differed significantly between at least 2 of the time points. **(B),** Top 50 expressed genes in each of the 6 modules. KME refers to module membership of each gene where values closest to 1 indicate genes that are most highly co-regulated with the other genes in the same module. **(C),** Expression of 4 annotated lncRNAs associated with blue (midstage) module are represented. Line plots showing increased mean lncRNA expression across single cells in W02, W06 and W12 hESC-CM. Values

are represented as mean gene FPKM ± s.e.m. Right panel shows violin plot with expression profile across different tissues from GTEx data set. Data are represented as log Transcripts Per Kilobase Million (TMP).

Supplementary Figure 5

(A), Schematic representation showing genomic sequence alignment for *VHRT* extracted from Ensembl Genome Browser[30]. *VHRT* is primarily conserved among the primates.

(B), Schematic representing the workflow for cellular fractionation. Fractionation efficiency was visualized using trypan blue staining. RNA was extracted from each fraction and treated with DNAse and profiled using Bioanalyzer assay.

(C), Histogram showing qPCR results for human atrial and ventricular heart tissue for *VHRT*, *MYH7, MYL2* and *NPPA*. n=3. ** $p \leq 0.01$.

Supplementary Figure 6

(A), Table reflecting the coding potential for each *VHRT* isoform and positive controls using known protein coding genes *MYL2* and *MYH7*. Values were generated using the publicly available coding potential calculator (CPC). HIT ID shows the Uniprot ID related to the gene. A higher HIT score indicates higher likely for protein coding. For a true protein-coding isoform, most hits likely reside within one frame, whereas for a true noncoding isoform, even if it matches certain known protein sequence segments by chance, these random hits are likely to scatter between any of the three reading frames. Thus, FRAME score measures the distribution of the HSPs among three reading frames, the higher the FRAME score, the more concentrated the hits are and the more likely the transcript is protein-coding.

(B), Schematic showing the small open reading frame (sORF) region on *VHRT6* (*VHRT* isoform 6) and the sequence of the putative peptide with a predicted length of 76 amino acids (aa) and suggested mass of ca. 8 kDa. **(C),** Cartoon showing plasmid and inserts used for *in-vitro* translation. *VHRT6* (without introns) insert was generated with FLAG-tag at the N-terminus or C-terminus of the sORF (in red). Positive control plasmid contained TagBFP (Blue Fluorescent Protein) with FLAG (in blue). All inserts were cloned into

lentivirus plasmid (pLenti-EF1α) and used for cell transfection. **(D),** Western blotting using anti-FLAG antibody detecting only a band of ~ 28kDa for TagBFP positive control plasmid. No bands were detected for *VHRT6*. BFP: blue florescent protein, BSD: blasticidin selection cassette, F: 1xFLAG-tag.

Supplementary Figure 7

(A), Sequence alignment for the deletion of 1110 bp with sgRNA_2/4 set, and 1210 bp with sgRNA_6/10 set. Primers sets used for genotyping are schematically represented. **(B),** Genotyping results proving successful deletion of the *VHRT* locus using sgRNA_2/4 set. Validation primer set p1+p4 and p2+p3, were used and clone #1 resulted in complete *VHRT* KO. **(C),** Genotyping results proving successful deletion of *VHRT* locus using sgRNA_6/10 set. Validation primer set p5+p8 and p6+p7 were used and clone #16 resulted in complete *VHRT* KO.

Supplementary Figure 8

(A), Graphs showing the absence of changes in expression of key differentiation markers (*OCT4*, *NANOG*, *ISL1* and *NKX2-5*) in *VHRT* KO cells. **(B),** Tri-lineage differentiation following *VHRT* deletion. Graphs showing absence of significant changes in expression of specific lineage markers, confirming that *VHRT* KO cells were able to successfully specify into ectodermal (*NES* and *PAX6*), mesodermal (*TBXT* and *CXCR4*) and endodermal (*SOX17* and *FOXA2*) lineages despite *VHRT* deletion. Data are represented as fold change expression ± s.e.m, n=3 biological replicates. Student's paired t-test with a two-tailed distribution.

Supplementary Figure 9

(A), Representative images of GFP expression signal at d7-8 post hES differentiation in WT, and *VHRT* KO #1 and #2. **(B),** Immunostaining shows sarcomere disorganization for *VHRT* KO #1 and #2, compared

to WT, following W02 and W06 of the CM differentiation protocol. α-act: alpha-actinin 2, and MLC2v: Myosin regulatory light chain 2.

Supplementary Figure 10

(A), Histogram showing the differentiation efficiency as *MYH6*-GFP %positive cells between WT and *VHRT* KO at W02. Error bars represented ± s.d. (n=4 for WT, n=4 *VHRT* KO #1, n=3 *VHRT* KO #2). **(B)** Assay for *MYH6*-GFP positive (+) cells in WT and *VHRT* KO #1 by flow cytometry at W02, W06 and W12.

Supplementary Figure 11

(A), Histogram showing qPCR results for *VHRT* and *ATP2A2* siRNA KD (n=2). **(B),** Table showing action potentials recordings. 14 cells transfected with *VHRT* GapmeR#1(*VHRT* knockdown) and 15 cells transfected with Negative A GapmeR (Negative Ctrl) were recorded. hESC-CMs were co-transfected with BLOCK-iT™ Alexa Fluor ® Red Fluorescent to identified positively transfected hESC-CMs. Recordings were made from spontaneously beating cells and with electrical stimulation at 1 Hz. After *VHRT* GapmR#1 knockdown, hESC-CMs showed a less negative maximal diastolic potential (MDP) and a significantly higher APD90/APD50 ratio and APD90-APD20 than Negative Ctrl, indicating an action potential that was less ventricular in nature, compared to negative control cells. Parameters of APs include: AP amplitude (APA), maximal diastolic potential (MDP), overshoot, upstroke velocity of depolarization (Vmax), AP durations (APD) at 20%, 50% and 90% of repolarization (APD20, APD50, and APD90). * $p<0.05$ compared to Negative A control cells. Student's t test. **(C),** Representative action potential profile from W06 hESC-CMs after *VHRT* GapmeR#1 KD (lower panel) compared to CTR (non-targeting GapmeR) cells (lower panel).

Supplementary Figure 12

(A), LocusZoom of the 1-MB genomic region harboring *VHRT* depicting significant *VHRT* cis-eQTL SNPs in heart left ventricle (HLV) (upper panel), as well SNPs significantly and/or suggestively associated with CAD, MI, HF (non-ischemic) and/or AF (lower panel). The latter data is taken from published GWAS[8,9,11,19,20]. **(B),** Top panel showing an example of eQTL rs3184504-C associated with decreased MI and CAD risk, but not with *VHRT* expression. Mid panel displaying the eQTL rs6489844-G, associated with decreased expression of *VHRT* in HLV tissue and suggestively associated with increased risk for HF, but decreased risk of MI. Similarly, on the bottom panel the eQTL rs11065780-C is significantly associated with low *VHRT* expression in HLV and increased HF, but suggestively associated with decreased CAD risk (bottom panel). NES: normalized effect size. P < 0.05. **(C)**, *VHRT* (probeset 230195_at) is significantly downregulated in left ventricular (LV) tissue in patients with Idiopathic Dilated Cardiomyopathy (N=15) versus healthy controls (N=11), but not in Hypertrophic- (N=5), Idiopathic (N=12) or Ischemic (N=31) cardiomyopathy based on reanalysis of microarray data from GEO study GSE1145 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE1145).

Supplementary Figure 13

A), Heatmap of normalized expression values of 1100 significantly differentially expressed (DE) genes in *VHRT* KD, compared to the RNAseq from Non-Failing (NF) versus failing (Dilated Cardiomyopathy, DCM) hearts. The subset that were consistent for downregulation in the 2 comparisons are represented in Figure 6.

SUPPLEMENTARY TABLES

Supplementary Table 1: Single cell RNAseq and data quality.

Supplementary Table 2: Tabulation of gene expression in each week.

Supplementary Table 3: WGCNA module and gene kME values.

Supplementary Table 4: Gene ontology of 6 modules.

Supplementary Table 5: VHRT transcripts description and annotation**.**

Supplementary Table 6: Gene ontology for genes differentially regulated in VHRT WT vs VHRT KO #1 in 02W,06W,12W old hES-CM.

Supplementary Table 7: Gene ontology for genes differentially regualted in VHRT CTRL vs VHRT GapmeR KD.

Supplementary Table 8: GeneNetwork data analysis.

Supplementary Table 9: Human Dilated cardiomyopathy (DCM) and non-failing (NF) tissue biodata.

Supplementary Table 10: Combined transcriptome analysis of VHRT GapmeR KD and human Dilated cardiomyopathy.

Supplementary Table 11: Oligonucleotides.

Figure 1

A

Stage: hESC   Mesendoderm   Cardiac mesoderm   Cardiac progenitor   "Early" hESC-CM   "Mid" hESC-CM   "Late" hESC-CM

Day: -2   -1   0   1   3   5   7   14   42   84

CHIR 99021   *   *   IWP2   *   *

2 weeks (W02)   6 weeks (W06)   12 weeks (W12)

hESC-CM

20 um   α-act/DAPI   20 um   MYH6-GFP/DAPI

GFP + cell n=36   GFP + cell n=60   GFP + cell n=56

B

Modules: Early-1, Early-2, Early-3, Midstage, Late-1, Late-2

W02, W06, W12

Gene Expression Level (FPKM)

C

Expression z-score   0   2   4

hESC-CM
batch 1 / batch 2: W02, W06, W12

Gene Modules: Early-1, Early-2, Early-3, Midstage, Late-1, Late-2

D

GO: biological processes of genes in the "midstage" module

- Anatomical structure morphogenesis
- Anatomical structure formation involved in morphogenesis
- Muscle contraction
- Muscle structure development
- Muscle system porcess
- Cardiovascualr system development
- Circulatory system development
- Muscle tissue development
- Regulation of system process
- Heart development

-log10(P-value)

E

connectivity between lncRNA in module (KME)

NPPA-AS1, LINC00881, RP11-532N4.2, RP11-432J24.5, VHRT

Chr: 1 2 3 4 5 6 7 8 9 10 11 12 14 15 16 18 21 X

Chromosome Position

F

W12

Expression z-score

VHRT, MYL2, MYH7, ATP2A2, NPPA-AS1, LINC00881, RP11-532N24.2, RP11-432J24.5, all other genes of the midstage module

Rank by VHRT expression across cells

Figure 2

A chr12 l111,375,000   111,385,500 l   111,395,000 l

VHRT
Isoform 1:
Isoform 2:
Isoform 3:
Isoform 4:
Isoform 5:
Isoform 6:

FPKM peak

W02
W06
W12

B hESC-CM

Relative expression

Isoform: All 1 2 3 4 5 6

n.s.

C

■ Nuclear RNA   □ Cytoplasmic RNA

Relative to nuclear RNA

All isoforms Isoform 1 Isoform 3 Isoform 4 Isoform 5 Isoform 6 GAPDH 18S U6 MALAT1

VHRT

D

VHRT

Relative expression

MYL2

MYH7

TAGLN

■ Ventricular myocyte marker
■ Smooth muscle cell marker
■ Cardiac fibroblast marker
■ Endothelial cell marker
■ Moncytic cell marker

PDGFRA

CDH5

CD45

hCF hCSMC hCAEC hTHP-1 W02 W06 W12

non-CM   hESC-CM

Figure 3

Figure 4

A

B

C

TISSUES		
TISSUE	SAMPLES	AVERAGE
Muscle	447	2.80
Heart	91	2.23
Thymus	9	0.82
Testis	14	0.73
Skin	866	0.62
Intestine	134	0.24
Esophagus	60	0.02
Brain	1281	0
Breast	32	-0.03
Epithelial tissue	201	-0.15
Lung	349	-0.24
Spleen	21	-0.24
Prostate	135	-0.29
Ovary	24	-0.30
Kidney	99	-0.31
Stomach	19	-0.31
Bladder	68	-0.34
Thyroid	60	-0.35
Pancreas	355	-0.36
Adrenal gland	23	-0.36
Liver	141	-0.37
Bone marrow	114	-0.38
Placenta	111	-0.39
Blood	3003	-0.40
Adipose tissue	276	-0.42
Gall bladder	8	-0.50

PATHWAYS & PROCESSES

GO BIOLOGICAL PROCESSES (BP)

TERM	P-VALUE
Cardiac muscle contraction	1.3 x 10e-12
Muscle filament sliding	5.9 x 10e-12
Cardiac myofibril assembly	3.6 x 10e-10
Muscle contraction	5.2 x 10e-09
Regulation of the force of heart contraction	6.8 x 10e-09
Strieated muscle contraction	3.2 x 10e-08
Sarcomere organization	3.0 x 10e-08

GO BP Cardiac muscle contraction

MYLK2, TTN, TCAP, TNNI2, MYL1, MYL3, VHRT, NKX2-5, TNNI1, TNNC1, MYH7, MYL2, MYBPC3, ACTC1, MYH6, CSRP3, TNNI3, TNNT2, CASQ2, MYL4, ATP1A2

KEGG PATHWAY

TERM	P-VALUE
Cardiac muscle contraction	2.3 x 10e-07
Hypertrophic cardiomyopathy (HCM)	3.9 x 10e-06
Dilated cardiomyopathy (DCM)	1.2 x 10e-05

HUMAN PHENOTYPES (HPO)	
TERM	P-VALUE
Ventricular tachycardia	2.3 x 10-15
Muscle fiber cytoplasmatic inclusion	4.3 x 10-13
Myopathy	6.8 x 10-13
Supraventricular tachycardia	1.2 x 10-11
Tachycardia	1.4 x 10-11
Nemaline bodies	4.7 x 10-11
Muscle fiber inclusion bodies	5.1 x 10-11
EMG abnormality	9.2 x 10-11
Supraventricular arrhythmia	1.4 x 10-10
Abnormality of muscle fibers	2.1 x 10-10
Abnormality of circulating enzyme level	1.1 x 10-09
Primary atrial arrhythmia	1.1 x 10-09
Abnormal levels of creatine kinase in blood	1.5 x 10-09
Elevated serum creatine phosphokinase	1.7 x 10-09
Atrial fibrillation	3.6 x 10-09
Type 1 muscle fiber predominance	4.1 x 10-09
Dilated cardiomyopathy	5.2 x 10-09
Lipoatrophy	2.5 x 10-08
Neck muscle weakness	4.5 x 10-08
Asymmetric septal hypertrophy	7.5 x 10-08

D

VHRT

logFC=-1.48, adjP-value=0.002

Figure 6

A

VHRT KD			GSE141910	
CTR	Gap#1	Gap#2	NF	DCM

Expression z-score
−2 −1 0 1 2

Down-regulated gene by VHRT GapmeR knockdown

MYH6, TMTC1, MPP3, GPR133, PVRL1, CRISPLD2, BIRC3, HIF3A, CALCOCO2, GREB1L, FGF7, LIN7A, SYNE2, CACNB2, AKR1C3, SREBF1, CAMK1D, RAB8B, MYO7A, PPP1R13B, MTUS2, FAM208B, SLC11A2, FOXN3, TTLL5, ZBTB44, ABCC1, SNX29

Down in *VHRT* KD
Down in DCM

TPT1−AS1, CGNL1, SOX6, ENOX1, FXYD6, KIF7, SESN3, PLXNC1, MYPN, KIF26A, FLRT2, LRRC10, MSS51

Down in *VHRT* KD
Up in DCM

Up-regulated gene by VHRT GapmeR knockdown

MGST1, MT1G, RASD1, MT1X, NMB, SERPINB8, MEDAG, ART5, COQ10A, ARRDC4, TM6SF1, TYRO3, GNPNAT1, ZNF267, CHST3, DDIT4, SHOC2, AHNAK, TNFRSF1A, NDRG2, SLC35C1, GPR56

Up in *VHRT* KD
Down in DCM

LMO2, DDIT3, IFITM1, C14orf79, KLF5, RP11−18l14.10, RP11−111M22.3, RP11−1017G21.5, SOX8, S100A2

Up in *VHRT* KD
Up in DCM

B

KEGG Pathway (Enricher) — p ≤ 0.05

Down in *VHRT* KD **Down** in DCM

- Cardiac muscle contraction
- Hypertrophic cardiomyopathy (HCM)
- Dilated cardiomyopathy (DCM)
- Adrenergic signaling in cardiomyocytes
- Oxytocin signaling pathway
- Vitamin digestion and absorption
- Folate biosynthesis
- Ferroptosis

−log10 (p-value)

C

GO Molecular Function (Enricher) — p ≤ 0.05

- Microfilament motor activity (GO:0000146)
- Actin binding (GO:0003779)
- Motor activity (GO:0003774)
- Ketosteroid monooxygenase activity (GO:0047086)
- Manganese ion transmembrane transporter activity (GO:0005384)
- Iron ion transmembrane transporter activity (GO:0005381)
- Sphingolipid transporter activity (GO:0046624)
- Protein-glutamic acid ligase activity (GO:0070739)

−log10 (p-value)

	Base Impedance [ohm]	Beat rate [bpm]	Amplitude [ohm]	Pulse width 50% [ms]	Upstroke velocity [ohm/s]	Downstroke velocity [ohm/s]
UT	577±18[**]	60±0	7.1±0.4[**]	162±10	223±29[***]	-56±5
NT	523±39	60±0	6.0±0.6	164±14	176±22	-50±7
VHRT **siRNA**	403±8[****]	60±0	4.4±0.2[****]	176±8	108±3[****]	-36±2[***]
ATP2A2 **siRNA**	405±26[****]	60±0	3.3±0.5[****]	260±10[****]	63±9[****]	-17±2[****]

Table 1: CardioExocyte profile from CM after *VHRT* siRNA knockdown.

Contractility profile was measured using Cardioexcyte 96 instrument (Nanion). Base impendance, beat rate, amplitude, pulse width, upstroke velocity and down velocity are shown. *ATP2A2* siRNA was used as positive control and non-targeted siRNA (NT) as well as un-treated (UT) as negative controls. The data was analyzed using the data control software from Nanion. One-way ANOVA, multiple comparisons, compared to control siRNA. Data are shown as mean ± s.e.m. * $p<0.05$, ** $p<0.0$, *** $p<0.001$ and **** $p<0.0001$.

Supplementary Figure 1

A

Karyotyping

B

FISH

C

	CM marker	MYH6-GFP	merged with DAPI
ACTN1			
CTNTT			
NKX2-5			

D

MEA recording

Control

100 uM noradrenaline

Field potential (µV)

10 s

Supplementary Figure 2

A

B

Frst heart field marker (FHF)

TBX5

CORIN

Second heart field marker (SHF)

ISL1

KDR

Mean gene expression across cells (FPKM ± SEM)

C

D

Supplementary Figure 5

A

VHRT genomic allinemnt

primates

non-primates

Expanded Alignments

Species:

SpeciesName: low quality assembly

SpeciesName: high quality assembly

Aligned seq:

☐ gap

■ aligned

B

hESC-CM (n=3)

Cell Fractionation Buffer

Nuclei   Cytoplasm

Cell Disruption Buffer

2X Lysis/Binding Solution and RNAqueous®

Nuclear RNA   Cytoplasmic RNA

Nanodrop   Bioanalyzer

qPCR

Pre-fractionation Buffer

LIVE

45 min post -fractionation Buffer

Nucleus

After DNAse treatment

tRNA   18S

cytoplasmic RNA (4,368 pg/ul)   Sample 9

total RNA (1,770 pg/ul)

nuclear RNA (640 pg/ul)

C

VHRT

MYH7

MYL2

NPPA

Atrial tissue   Ventricular tissue

A

Gene ID	Isoform ID [Ensembl or NONCODE]	Sequence length [bp]	Coding / non-coding	Coding Potential Score	BLAST summary				
					HIT ID [Uniprot]	HIT score	FRAME score	ORF length [bp]	Protein length [AA]
VHRT	ENST00000331096	364	coding	1.17211	Q8IVN4_human	27.5	252.5	229	77
	ENST00000553177	524	noncoding	-0.510149	-	-	-	-	-
	ENST00000547607	674	noncoding	-0.995032	-	-	-	-	-
	ENST00000548368	705	noncoding	-0.686213	-	-	-	-	-
	NONHSAT030711.2	872	coding	0.914967	Q8IVN4_human	27	243	357	119
	NONHSAT162760.1	937	noncoding	-0.961563	-	-	-	-	-
MYL2	ENSG00000111245	664	coding	3.56504	P10916 (MLRV_human)	31.6	347.6	457	153
MYH7	ENST00000355349	6087	coding	37.1071	P12883 (MYH7_human)	187.3	989.1	5806	1936

B

VHRT

Isoform 6 : Exon1 — Exon 2 (ORF)

Q8IVN4: MYLAQSLSVRAAFLQTAQQGAVLPRLRAATASRWRTKGAESGEGEEHLSSHLSPARSARPVSSVLLLLKIRRVRMK

Length: 76 aa

Mass: 8.356 kDa

C

BSD
pLenti- EF1α

START
EF1a | 1 | F | 2 — ORF+F N-terminus

VHRT6

STOP
EF1a | 1 | 2 | F — ORF+F C-terminus

Positive control

EF1a | TagBFP | F — BFP+F

EF1a | TagBFP — BFP

F = 1x FLAG tag

D

VHRT6

ORF+F N-terminus | ORF+F C-terminus | BFP+F | BFP

kDa
245 -
180 -
75 -
25 -
5 -

Supplemental Figure 7

A

1110 bp

p1   p2   p3   p4

WT: CCCCCTCCCCTTGCTATTCTCGTGATAGAGCTCTCCTCAC*****CAAACGACCTAGGAGGTTAGAAGTTGATACCTTCTATGGAGT

clone: #1: CCCCTCCCCCTTT--GTTAGAAGTTGATACCTTCTATGGAGT

sgRNA_2   sgRNA_4

1210 bp

p5   p6   p7   p8

WT: CTTGCTATTCCTTTCTCCCCGTAGATTGCCC****GCAGCCAGCTTCCCGCTGGCGAACCAAGGGCGCGGAATCTGGAGAGGGC

clone: #16: CTTGCTATTCCTTTC--CCAAGGGCGCGGAATCTGGAGAGGGC

sgRNA_6   sgRNA_10

B

sgRNA_2+4   Product gel size [bp]

L   p1+p4   p2+p3   p1+p4   p2+p3   p1+p4   p2+p3

10000 -

2000 -

1000 -   ◄ 1319

◄ 833

250 -   ◄ 211

clone: #1   #9

VHRT KO#1   WT

c

sgRNA_6+10   Product gel size [bp]

L   p5+p8   p6+p7   p5+p8   p6+p7   p5+p8   p6+p7

10000 -

2000 -   ◄ 1558

1000 -

◄ 687

250 -   ◄ 347

#16   #9

VHRT KO#2   WT

Supplemental Figure 9

A

B

- : negative GFP
+ : positive GFP

GFP

A

VHRT

ATP2A2

B

Condition	Tot. no. cells	APA [mV]	MDP [mV]	Overshoot [mV]	Vmax [dV/dt]	APD90 [ms]	APD50 [ms]	APD20 [ms]	APD 90/50 ratio	APD 90-20
VHRT GapmR#1 [mean± SD]	14	92.4 ± 13.6	-53.7 ± 6.6*	38.7 ± 14.9	9.4 ± 6.3	444.2 ± 118.5	330.4 ± 95.2	192.8 ± 60.9	1.36 ± 0.1*	251.4 ± 79.9*
Negative CTR [mean± SD]	15	99.1 ± 10	-61.2 ± 7.1	37.9 ± 8.0	8.2 ± 4.6	411.5 ± 100.6	379 ± 109.6	245.4 ± 103.5	1.23 ± 0.1	166.0 ± 40.2

C

A

ipsum

C

B

rs3184504

- **rs3184504-C** significantly associated with **decreased MI risk** (p= 7.04e-14, OR=0.93).
- **rs3184504-C** significantly associated with **decreased CAD** risk (p=1.03e-09, OR=0.94).
- **rs3184504-C** not significantly associated with VHRT or MYL2 cardiac expression.
- **rs3184504-C** significantlty associated with deacreased ALDH2 expression in multiple tissue, excluding heart.

rs6489844

- **rs6489844-G** suggestively associated with **decreased MI risk** (p= 7.90e-07, OR=0.95)
- **rs6489844-G** suggestively associated with **increased HF (non-ischemic) risk** (p= 5.55e-05, OR=1.15)
- **rs6489844-G** significantly associated with lower VHRT HLV expression (MYL2 expression is n.s.)

rs11065780

- **rs11065780-C** suggestively associated with **increased HF (non-ischemic) risk** (p= 5.25e-05, OR=1.15)
- **rs11065780-C** suggestively associated with **decreased CAD risk** (p= 1.76e-03, OR=0.97)
- **rs11065780-C** significantly associated with lower VHRT HLV expression (MYL2 expression is n.s.)

Supplementary Figure 13