1

# Impact of a human gut microbe on *Vibrio cholerae* host

# colonization through biofilm enhancement

Kelsey Barrasso[1,2], Denise Chac[3], Meti D. Debela[6], Catherine Geigel[4], Jason B. Harris[6,7], Regina

C. LaRocque[6], Firas S. Midani[8], Firdausi Qadri[9], Jing Yan[4,5], Ana A. Weil[3,*], Wai-Leung Ng[1,2,*]

**Affiliations:**

1. Department of Molecular Biology and Microbiology, Tufts University School of Medicine, Boston, MA

2. Program of Molecular Microbiology, Graduate School of Biomedical Sciences, Tufts University School of Medicine, Boston, MA

3. Department of Medicine, University of Washington, Seattle, WA

4. Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT

5. Quantitative Biology Institute, Yale University, New Haven, CT

6. Division of Infectious Diseases, Massachusetts General Hospital, Boston, MA

7. Department of Pediatrics, Harvard Medical School, Boston, MA

8. Department of Molecular Virology and Microbiology at Baylor College of Medicine, Houston, TX

9. International Center for Diarrheal Disease Research, Bangladesh, Dhaka, Bangladesh

*   Co-corresponding authors/equal contributions

Corresponding authors: Ana A. Weil and Wai-Leung Ng

**Email:** anaweil@uw.edu wai-leung.ng@tufts.edu

**Author Contributions:**

- •    Designed research (KB, DC, JY, AAW, WLN)

- •    Performed research (KB, DC, MDD, CG, JY)

- •    Contributed new reagents or analytic tools (FSM, JY, JBH, RCL, FQ)

- •    Analyzed data (KB, DC, FSM, JY, AAW, WLN)

- •    Wrote the paper (KB, DC, JY, AAW, WLN)

**Competing Interest Statement:** No competing interests.

**Keywords:** Bacterial pathogenesis, gut microbiome, mixed species biofilm

**Abstract**

Recent studies indicate that the human intestinal microbiota could impact the outcome of infection by *Vibrio cholerae,* the etiological agent of the diarrheal disease cholera. A commensal bacterium, *Paracoccus aminovorans,* was previously identified in high abundance in stool collected from individuals infected with *V. cholerae* when compared to stool from uninfected persons. However, if and how *P. aminovorans* interacts with *V. cholerae* has not been experimentally determined; moreover, whether any association between this bacterium alters the behaviors of *V. cholerae* to affect the disease outcome is unclear. Here we show that *P. aminovorans* and *V. cholerae* together form dual-species biofilm structures at the air-liquid interface, with previously uncharacterized novel features. Importantly, the presence of *P. aminovorans* within the murine small intestine enhances *V. cholerae* colonization in the same niche that is dependent on the *Vibrio* exopolysaccharide (VPS) and other major components of mature *V. cholerae* biofilm. These studies illustrate that dual-

species biofilm formation is a plausible mechanism used by a gut microbe to increase the virulence

of the pathogen, and this interaction may alter outcomes in enteric infections.

**Significance Statement**

While ample evidence suggests that the outcome of some enteric infections can be affected by the

intestinal microbiota, how specific gut microbes change the behaviors of a pathogen is unclear.

Here we characterize the interaction between *Vibrio cholerae* and *Paracoccus aminovorans*, a gut

microbe known to increase in abundance in the intestines during active *V. cholerae* infection in

humans. These two bacteria form a dual-species biofilm structure at the air-liquid interface, and the

gut microbe increases the host colonization efficiency of *V. cholerae*. Importantly, our study

identifies a previously unknown mechanism of gut microbe-pathogen interaction that has the

potential to alter the disease outcome.

**Introduction**

*Vibrio cholerae* (*Vc*) causes an estimated 3 million infections and 120,000 deaths each year, and larger and more deadly outbreaks have increased during the last decade (1, 2). A wide range of clinical outcomes occur in persons exposed to *Vc*, ranging from asymptomatic infection to severe secretory diarrhea. It is nearly certain that many behaviors of *Vc* in the aquatic environment and inside the host are significantly affected by the presence of other microbes (3), and recent studies provide evidence that the gut microbiota may impact the severity of cholera (4-7).

Several functions of the gut microbiota influence the growth or colonization of enteric pathogens, including production of anti-microbial compounds, maintenance of the intestinal barrier, regulation of the host immune response, and modulation of available nutrients (8). Gut microbes have been shown to have an important role in *Vc* infection in various animal models. For instance, disruption of the commensal microbiota with antibiotics is required to allow successful *Vc* colonization in adult rodent models (9, 10). Conversely, *Vc* actively employs a Type VI secretion system to attack host commensal microbiota to enhance colonization of the gut in infant mice (11). Moreover, specific microbial species have a profound impact on *Vc* colonization. *Blautia obeum*, an anaerobic Gram-positive bacterium, decreases *Vc* intestinal colonization presumably by producing a signaling molecule that induces *Vc* into a high cell-density quorum sensing state (5) in which virulence gene expression is repressed (12, 13). Certain microbiota species reduce *Vc* colonization by producing the enzyme bile salt hydrolase that degrades the host-produced virulence-activating compound taurocholate (4, 5). Through metabolizing host glycans into short chain fatty acids that suppresses *Vc* growth, a prominent commensal species, *Bacteroides vulgatus,* reduces *Vc* proliferation within the intestine (14).

While the above studies exemplify how a single microbe or a group of microbes can protect the host from *Vc* infection, the mechanisms used by certain gut microbes to promote *Vc* virulence, thereby increasing the likelihood of individuals to develop cholera and worsen disease outcomes,

are less well understood. We have previously studied household contacts of cholera patients to

understand how gut microbes impact on susceptibility to cholera and identified bacteria associated

with increased or decreased susceptibility to *Vc* infection (6, 7). We also observed that the gut

microbial species *Paracoccus aminovorans* (*Pa*) was more likely to be present and more abundant

in the gut microbiota during *Vc* infection (7). The association between *Pa* and *Vc* is highly unusual

because most of the native gut microbiota is typically displaced by secretory diarrhea during cholera

(5, 15). To determine the underlying mechanisms driving these correlative clinical findings, we

evaluated the relationship between *Pa* and *Vc* in co-culture and determined the effects of *Pa* on *Vc*

infection outcomes with *in vivo* models. Here we show that *Pa* interacts directly with *Vc* to form

dual-species biofilm structures with previously uncharacterized features. Moreover, *Vc* colonization

inside the animal host is enhanced by the presence of *Pa* in the small intestine, and this effect is

dependent upon *Vc* biofilm production. Our findings demonstrate a plausible mechanism by which

a gut microbe specifically associates with *Vc,* and this reinforces our microbiome analysis in

humans that identified *Pa* as highly associated with infected individuals. Our findings also

demonstrate that interactions between these two species have the potential to directly impact *Vc*

pathogenesis and alter outcomes of *Vc* infection in humans.

**Results**

**_P. aminovorans_ is differentially abundant in individuals with active _V. cholerae_ infection**

_Paracoccus_ is a genus of soil microbes found in low abundance in the gut microbiome of humans (16, 17). Our previously published analysis of stool gut microbes from household contacts of cholera patients identified _P. aminovorans (Pa)_ as an unexpectedly abundant gut microbe during active _V. cholerae (Vc)_ infection, and this organism was rarely found in uninfected participants (7). In this prior study, we used a support vector machine model with recursive feature elimination to learn patterns of relative abundance of operational taxonomic units (OTUs) that distinguished infected (defined as _Vc_ DNA detected in stool, or culture positive) from uninfected persons (_Vc_ DNA undetected in stool). The model was trained on a subset of study participants and tested on another subset in a hold-out validation. Here, we have extracted data from this prior study to examine separately the _Pa_ OTUs in infected compared to uninfected persons. _Pa_ abundance was significantly higher as a proportion of the total sequencing reads in the stool of infected participants (6/22, 27%) of infected household contacts had detectable _Pa_ compared to only 5.6% (2/36) of uninfected individuals (Figure 1A). The ratio of _Pa_ to _Vc_ abundance present during infection was variable and averaged 1:1 (Figure 1B). These findings were particularly interesting because typically there is a drastic reduction of nearly all gut microbes during active _Vc_ infection (5, 15) due to secretory diarrhea, oral rehydration solution ingestion, and _Vc_ infection itself, and yet here _Pa_ was found in an increased abundance in some actively infected participants. Based on these findings, we hypothesized that _Pa_ may be resistant to displacement from the gut during infection. While our previous study demonstrates a positive correlation between _Pa_ in human stool and _Vc_ infection, a causal relationship between this gut microbiota species and _Vc_ infection had not been previously established.

**_Pa_ increases _Vc_ host colonization**

We modified a well-established infant mouse colonization model (18) to assess whether the presence of _Pa_ in the small intestine would promote _Vc_ host colonization. First, we isolated a

spontaneous streptomycin resistant (Strep$^R$) mutant derived from the ATCC type strain of *Pa* for

selection and enumeration of *Pa* following host colonization. Infant mice (3-day old) were

intragastrically inoculated with *Pa* ($10^7$ colony forming units [CFUs]) every 12 hours for 4 doses (0,

12, 24, 36 hours). At 24 and 48 hours after the first inoculation, small intestines from these animals

were dissected and homogenized. Gut homogenates were serially diluted and plated on medium

containing streptomycin to assess *Pa* colonization. Strep$^R$ *Pa* colonies (>$10^6$ CFUs/small intestine)

were recovered at these two time points (Figure 2A), and no Strep$^R$ colonies was detected in the

mock treated group, indicating that *Pa* successfully and stably colonized the small intestines of

these animals using these methods. Unlike previous studies (9, 10), pretreatment with antibiotics

did not change the outcome of *Pa* colonization (not shown). Sequencing analysis of the mouse

small intestines demonstrated no significant change in the microbial composition and diversity with

and without *Pa* colonization (Supplemental Figure 1 A and B).

We then evaluated if pre-colonization by *Pa* would influence *Vc* colonization in the small intestine.

*Pa* pre-colonization in the infant mice was established over a 36-hour period as described above.

Negative control animals were inoculated with sterile media in place of *Pa* over the same dosing

schedule. Twelve hours after the last *Pa* inoculation (i.e., 48 hours after the first *Pa* inoculation),

these animals were infected with *Vc* ($10^6$ CFU) to evaluate whether pre-colonization with *Pa* had

an impact on *Vc* colonization. Although we do not fully understand the exact composition and

growth dynamics of *Vc* and *Pa* inside the human gut, the pre-colonization/infection scheme was

aimed to closely simulate the ratio of *Pa* to *Vc* observed in the gut microbiota of *Vc* infected humans

(Figures 1 and 2A). Comparing *Pa* pre-colonized mice to the control group, there was a significant

increase (~10-fold, p≤0.0001) of *Vc* colonization in the mice pre-colonized with *Pa* (Figure 2B) 24

hours after infection. This enhanced intestinal colonization by *Vc* in the *Pa*-colonized mice was

observed as early as 6 hours after infection and maintained throughout the colonization period

(Supplementary Figure 2).

We reasoned that it was also possible for *Vc* and *Pa* to encounter one another in the environment

before entering the host. To model this scenario, *Vc* was mixed with *Pa* in 1:1 ratio, and the mixture

was used immediately for animal infection. In agreement with the results obtained with the *Pa* pre-

colonization model, *Vc* intestinal colonization was significantly higher when coinfected with *Pa* than

without *Pa* (Figure 2C). Given *Pa* colonization did not overtly change the overall composition of the

gut microbiota (Supplementary Figure 1), collectively, our results demonstrate that the presence of

a single gut microbiota species is sufficient to increase *Vc* host colonization. Our findings also

illustrate that our approach to microbiome studies in humans (6, 7) can be used as a predictive tool

to identify gut microbes that alter *Vc* virulence.

**Pa promotes Vc biofilm formation**

To investigate if the increased *Vc* intestinal colonization is due to direct interactions between *Vc*

and *Pa,* these two species were co-cultured and allowed to propagate for three days where both

planktonic growth and pellicle formation (i.e., biofilm formation at the air-liquid interface) of both

species was monitored. There was a small difference (< 2-fold) in growth in the planktonic phase

of either *Vc* or *Pa* in the co-cultures when compared to the cultures containing a single species

(Figure 3A). However, the *Vc/Pa* co-culture formed a pellicle that was visibly thicker and more

robust than that formed by *Vc* monoculture (photo shown in Figure 3B). The *Pa* monoculture did

not form a visible pellicle. The co-culture pellicle samples were carefully lifted and removed from

the culture medium, washed, and agitated to release single cells for enumeration of each species.

Compared to *Vc* monoculture, the co-culture samples contained over 50-fold more *Vc* cells while

the ratio of *Vc* to *Pa* approached to approximately 1:1 (Figure 3C). Moreover, only a small fraction

(0.01%) of *Vc* and *Pa* could be washed off from the isolated pellicles (Figure 3D), suggesting that

these species are tightly integrated into the pellicle.

Based on the above data, we hypothesize that *Vc* and *Pa* form dual-species biofilms at the air-

liquid interface. This is unexpected because *Vc* is known to form a clonal community in both *in vitro*

and *in vivo* biofilms and these are known to exclude other species including even planktonic *Vc*

cells (19, 20). To test this hypothesis, we transferred the co-culture pellicles onto coverslips for

imaging with confocal microscopy (Figure 4).  All cells in the pellicle were stained with FM 4-64

membrane dye, and *Vc* cells were differentiated from *Pa* using a constitutively produced

mNeonGreen reporter (21) expressed from a neutral *Vc* locus (22). In the *Vc/Pa* co-culture pellicles,

we observed a continuous film structure spanning the entire pellicle (Figure 4A). Notably, cocci-

shaped *Pa* cells were clearly visible in the co-culture pellicle (Figure 4B-C), consistent with the CFU

quantification in Figure 3A. Interestingly, *Pa* cells were found throughout the pellicle, with a higher

abundance in the bottom layer (Figure 4D-F), and always in close association with *Vc* cells. In

summary, we found that *Vc* and *Pa* coexist stably in the pellicle structure and this relationship may

explain the mechanism by which *Pa* resists displacement in humans during active *Vc* infection.

Next, we used a standard crystal violet (CV) microtiter plate assay (23) to quantitatively evaluate

how *Vc* and *Pa* interact under pellicle forming conditions. *Vc* and *Pa* were simultaneously

inoculated into the wells of microplates in two different *Vc:Pa* ratios (1:1 and 1:10). We also tested

if the viability of *Pa* was crucial for this interaction by using heat-killed *Pa* as a control. Consistent

with our pellicle compositional analysis, *Vc* formed a more robust biofilm than *Pa* under these

conditions as demonstrated by increased CV staining in wells containing *Vc* only compared to wells

containing *Pa* only (Figure 5A). Importantly, CV staining was increased in wells containing *Vc* and

live *Pa* compared to wells with *Vc* only, in a concentration-dependent manner (Figure 5A). In

contrast, CV staining was not different in wells containing *Vc* and heat-killed *Pa* compared to wells

with *Vc* only (Figure 5A).

To replicate our mouse experiments (Figure 2), we also tested if the order in which the two species

encounter one another is critical for the *Vc* biofilm enhancement phenotype. *Pa* was grown in wells

24 hours before the addition of *Vc*. As in our previous results in the co-inoculation experiment, an

increase in CV staining was observed in wells in which the two species were added sequentially,

but not in the wells with *Vc* only (Supplementary Figure 3). Moreover, wells pre-incubated with heat-

killed *Pa* and subsequently inoculated with *Vc* had no increase in CV staining compared to wells

inoculated with *Vc* alone (Supplementary Figure 3). Together, our biofilm quantification data

suggests that the presence of *Pa*, regardless of the order of encounter, results in an enhanced

biofilm formation of *Vc*.

**Vibrio exopolysaccharide is essential for a stable biofilm structure formed by *Vc* and *Pa***

To understand what biofilm component is required for the enhancement of biofilm production in

*Vc/Pa* co-culture, we repeated the above experiments with a Δ*vpsL Vc* mutant that cannot produce

the *Vibrio* exopolysaccharide (VPS) necessary for mature biofilm formation (24). In contrast to what

we observed with a *vpsL*$^+$ strain, there was no significant increase in CV staining in wells with both

Δ*vpsL* mutants and *Pa*, when compared to wells with the Δ*vpsL* mutants only (Figure 5B).

To further investigate the role of VPS in promoting co-culture biofilms, we stained the co-culture

pellicle *in situ* with Wheat Germ Agglutinin (WGA), a common stain for VPS (which contains GlcNAc

moieties, (25)). To avoid spatial overlap with the membrane stain (excited at 561 nm), the *Vc* cells

used in this experiment express a cyan-fluorescent protein SCFP3A cytosolically (excited at 445

nm), and the WGA is conjugated to Oregon Green (excited at 488 nm). Figure 6A shows a 3D view

of a large area of a *Vc/Pa* co-culture pellicle with WGA staining. We then compared the intensity of

WGA staining in areas with different compositions of *Vc* and *Pa*. Area 1 in Figure 6A represents a

location where *Vc* is the predominant species and *Pa* abundance is low (i.e., FM 4-64 staining

overlapped entirely with SCFP3A signal), while areas 2 and 3 in Figure 6A represent locations

where *Vc* and *Pa* coexist (i.e., contain regions with FM 4-64 staining but no SCFP3A signal).

Surprisingly, the WGA signal intensity was elevated in the *Pa*-rich regions rather than in the *Vc*-

rich regions (Figure 6B). Zoom-in view of the *Vc*-enriched region shows the characteristic sub-

envelope structures around clusters of *Vc* cells (Figure 6C), consistent with the known VPS

morphology in submerged *Vc* biofilms (26). In *Pa*-rich regions, the WGA-signal was stronger, and

the VPS structures were more compact (Figure 6C). Importantly, the VPS structure we observed

enclose both *Pa* and *Vc* cells, providing an intuitive explanation for how *Pa* cells are incorporated

into the pellicle. Together, these results suggested that the physical presence of *Pa* in co-culture

pellicle augments the production of VPS in *Vc* cells, leading to increased *Vc* biofilm formation; the

*Pa* cells, in turn, rely on VPS to be integrated into the 3D structure of the pellicle.

**Enhancement of *Vc* host colonization by *Pa* depends on biofilm exopolysaccharide**

Biofilm-grown *Vc* cells are known to be more infectious in humans due to increased resistance to

gastric pH and higher expression of virulence factors (e.g., such as the toxin co-regulated pilus,

which mediates host colonization) compared to planktonically grown cells (27-29). We hypothesize

that because *Vc* biofilm formation is enhanced in the presence of *Pa*, this results in increased

virulence inside the host, in a VPS-dependent manner.

To test our hypothesis and measure if the effect of the *Vc/Pa* biofilm interaction impacts host

colonization, we compared the colonization efficiency between wild-type (WT) or the $\Delta vpsL$ mutants

in infant mice with and without *Pa* pre-colonization. As shown previously (30), the $\Delta vpsL$ mutant

was able to colonize the mouse small intestine equally as well as the WT $vpsL^+$ strain, confirming

that the VPS is not absolutely required for host colonization when *Vc* was administered to the

animals alone. In contrast, while *Pa* increased WT $vpsL^+$ *Vc* colonization, the $\Delta vpsL$ mutant did not

exhibit the enhanced colonization phenotype in the *Pa* pre-colonized mice (Figure 7A). Similar

results were observed using the coinfection model; when *Vc* Δ*vpsL* mutants were coinfected with

*Pa*, there was no increase in host colonization (Figure 7B). Together, we concluded that the

enhancement of *Vc* intestinal colonization in the presence *Pa* is dependent on the *Vibrio*

exopolysaccharide, in line with our *in vitro* data.

**Accessary biofilm matrix proteins are involved in *Pa* and *Vc* interaction**

Mature *Vc* biofilm is stabilized with a variety of accessary matrix proteins in addition to the VPS

(26, 31, 32). To interrogate the roles of these components in the interactions between *Vc* and *Pa*,

we tested mutants lacking the cell-cell adhesion proteins RbmA (26, 31, 33) and mutants lacking

surface adhesion redundantly conferred by RbmC and Bap1 (26, 32, 33) for their ability to increase

biofilm formation in the presence of *Pa* using CV assays (Figure 7C). When compared to the wells

containing the Δ*rbmA* mutant alone, the CV staining was higher in the wells with both the Δ*rbmA*

mutant and *Pa.* However, the increase was not as high in the Δ*rbmA* mutant when compared to

that in WT *Vc/Pa* co-culture (Figure 7C). Furthermore, the presence of *Pa* did not increase CV

staining in the wells containing the Δ*rbmC* Δ*bap1* mutants (Figure 7C).

We then performed the infant mouse colonization experiments with *Vc* biofilm matrix protein

mutants using the *Pa* coinfection model to test the roles of these proteins *in vivo*. For this series of

experiments, each *Vc* biofilm mutant was coinfected into the animals with *Pa* in 1:1 ratio. In

agreement with our *in vitro* results, while host colonization was significantly higher for WT *Vc*

coinfected with *Pa* than without *Pa* (Figure 2C), Δ*rbmA* mutants did not show any increase in host

colonization when coinfected with *Pa*, and Δ*rbmC* Δ*bap1* mutants demonstrated reduced

enhancement in colonization during coinfection with *Pa* (Figure 7B). These results indicate that the

ability of *Vc* to form a structurally intact biofilm is important for the enhancement of colonization

facilitated by the presence of *Pa.*

**Discussion**

Evidence that the composition of the gut microbiota influences the clinical outcomes of enteric infections in humans is accumulating (34, 35). Several studies have identified commensal species and underlying colonization resistance mechanisms that could be protective against *V. cholerae* (*Vc*) infection. While these studies suggest that microbiota species reduce *Vc* virulence through various mechanisms during the early stages of infection (4, 5, 14), the precise role of these colonization resistance mechanisms in impacting susceptibility to cholera in humans has only begun to be appreciated. For instance, a bacterium in the genus *Blautia* was recently found to encode functions that confer colonization resistance (e.g., bile salt hydrolase) to *Vc* infection (4). Consistent with this finding, our previous stool microbiome study has also independently identified that one of the species in the genus *Blautia* is correlated with decreased susceptibility to *Vc* infection (6, 7).

While previous studies have identified microbiota-associated mechanisms that are protective against *Vc* infection, examples of interactions between *Vc* and a human-associated microbiota species that increases *Vc* pathogenicity are scarce. Although *Escherichia coli* and *Vc* are believed to reside in different intestinal niches, one previous study showed that an atypical *E. coli* isolated from a mouse that does not ferment lactose can increase the virulence of a quorum-sensing (QS) defective *Vc* strain N16961 (36). How QS-proficient *Vc* strains, which are prevalent in toxigenic clinical isolates (37), respond to typical *E. coli* in the human gut remains to be studied. In contrast, a recent study showed that *E. coli* motility facilitates aggregation of these two organisms in a dual-species biofilm, but there was no impact of such aggregation on *Vc* intestinal colonization (38). Indeed, coaggregation between *Vc* and other microbiota species has been observed (39), but these associations are not known to have a direct influence on *Vc* pathogenicity. This is consistent with our prior human studies in which *E. coli* species were present in the gut microbiota of persons during active *Vc* infection, but these were not correlated with active *Vc* infection (7). Our findings highlight the importance of coupling mechanistic studies (*in vitro* and animal models) with human

microbiome data analysis to pinpoint the relevant species and interactions involved in enteric

infections.

Here we show that the presence of a human gut microbe *Pa* promotes *Vc* host colonization, which

is consistent with our prior human study in which *Pa* was more likely to be present in persons

infected with *Vc*. This raises the possibility that uncharacterized interactions between *Vc* and

members of the gut microbiota may exacerbate *Vc* virulence and contribute to increased morbidity.

Our current study also establishes a plausible mechanism used by *Pa*, and perhaps other gut

microbes, to increase the virulence of *Vc* through induction of biofilm formation, a physiological

state in which *Vc* is known to increase expression of other virulence factors critical for human

infection and disease (27, 28). *Vc* biofilms have also been demonstrated to deform and even

damage tissue-engineered soft epithelia mimicking the host tissue (40), suggesting that *in vivo*-

formed biofilm structures could negatively impact host gut physiology.

While VPS and other biofilm components are not usually considered critical host colonization

factors, we found that these macromolecular structures were essential for the enhancement of *Vc*

host colonization induced by *Pa*. Whether these components mediate other *Vc*-gut microbe

interaction has not been studied. Interestingly, many gut microbes appear to predominantly exist

in the form of mixed-species biofilms on mucosal surfaces (41), suggesting microbiota-induced

biofilm enhancement could play a major role in modulating virulence of other pathogens. Many

structural components, regulatory factors, and signaling transduction pathways that control biofilm

formation in *Vc* have been well characterized (42), and these factors could be targeted for

manipulation by other gut microbes that modulate *Vc* virulence. For example, 3,5-dimethylpyrazin-

2-ol (DPO) was recently discovered as a new class of *Vc* quorum-sensing autoinducer that binds

to the transcription factor VqmA to activate expression of *vqmR*, which encodes a small regulatory

RNA that downregulates *Vc* biofilm formation. The VqmA-VqmR system can be activated both *in*

*vitro* by *E. coli* and *in vivo* by *B. obeum* (5, 43), and results in suppression of biofilm formation.

Interestingly, *Pa* demonstrates the opposite tendency by promoting *Vc* biofilm formation, with

implications for the enhancement of *Vc* colonization, in contrast to other commensal bacteria.

Many aspects of the *Vc*/*Pa* interaction are still unclear. What is the selective advantage that fosters

the formation of dual-species biofilm? Investigation of the structure-function relationship in other

multispecies biofilms, such as dental biofilms, demonstrates a coordinated organization of each

species that allows for optimal nutrient and oxygen usage, as well as mechanical stability (44, 45).

While we did not observe any growth yield enhancement in the planktonic phase of the co-culture,

there was a significant increase of *Vc* and *Pa* abundance in the co-culture pellicle at the air-liquid

interface. Thus, a possible driving force of this interaction could be the optimization of nutrient

sharing and distribution, or removal of toxic metabolites accumulated during growth.  The exact

mechanism used by these two species to detect and coordinate with each other remains unclear.

Secreted small molecules produced by *Pa* do not appear to impact *Vc* as evidenced by our prior

studies in which *Vc* cultured in *Pa* spent-cell supernatant did not yield result in increased biofilm

formation (7). Therefore, we surmised that the close physical association between *Vc* and *Pa* cells

in space in the co-culture pellicles is required for the enhanced biofilm formation. This hypothesis

is supported by our microscopy analysis. The *Vc/Pa* interaction has two reciprocal aspects: First,

*Pa* activates the production of VPS in *Vc* cells, leading to enhanced pellicle formation. Future

characterizations of *Pa* could potentially elucidate the underlying molecular mechanism of this

effect *Pa* has on *Vc*.  Second, in order to be integrated into the pellicle structure, *Pa* cells seem to

physically interact with VPS. Interestingly, we have shown that VPS staining signals are stronger

in the *Pa*-rich regions than in the *Vc*-rich regions. This could be explained either by a stronger

attraction between VPS and *Pa* than between VPS and *Vc* cells, or by activation of VPS

biosynthesis genes in *Vc* cells in the vicinity of the *Pa* cells, or both. Future biochemical and

biophysical studies to investigate this relationship may provide new insights about the interactions

between *Pa* and *Vc* biofilm, and about pathogen-gut microbe interactions in general.  Other

members of the *Paracoccus* genus are known to form biofilms and encode adhesins to facilitate

surface attachment (46, 47), and the potential role of these adhesins in facilitating interaction with

*Vc* remains to be studied.

In conclusion, we describe a novel interaction between *Vc* and a gut microbe found in high

abundance in *Vc*-infected persons that leads to a significant change in *Vc* biofilm behaviors, as well

as an increase in the virulence of the pathogen. Our findings are also consistent with other

observations that rare gut microbial species can have significant impacts on microbial ecosystems

(48). This study adds to the growing number of pathogen-gut microbial species interactions that

may impact outcomes in human diseases.

**Materials and Methods**

**Prior published study sample collection and analysis**

In a prior study, we enrolled household contacts of persons hospitalized with cholera at the

International Centre for Diarrheal Disease Research, Bangladesh (icddr,b)(7). Briefly, in this

previously published study, household contacts were followed prospectively with rectal swab

sampling, 30 days of clinical symptom report, and vibriocidal titer measurements, and 16S rRNA

sequencing was performed on rectal swab sampling from the day of enrollment in the study (7).

Persons with evidence of *V. cholerae* (*Vc*) infection at the time of enrollment in the study were

compared to those who did not have evidence of infection in a model to detect gut microbes that

were differentially abundant during *Vc* infection (7). *Vc* infection was defined as *Vc* DNA identified

on 16S rRNA sequencing or a positive *Vc* stool culture. In this previously published study, we used

a machine learning method called a support vector machine (SVM), which utilizes patterns of OTU

relative abundance to detect OTUs associated with infected compared to uninfected persons. This

SVM was used with a recursive feature elimination algorithm that simplifies models and increases

accuracy of the identification of differentially associated OTUs by removing uninformative bacterial

taxa (7). For the present study, we re-examined the microbiome data from household contacts at

the time of enrollment to quantify the abundance of 16s rRNA reads that mapped to *Pa* OTUs

between uninfected study participants and infected participants.

**Strains and culture conditions**

All *V. cholerae* (*Vc*) strains used in this study are streptomycin-resistant derivatives of C6706, a

1991 El Tor O1 clinical isolate from Peru (49). The in-frame Δ*vpsL* deletion mutants used in various

assays were previously described (50). The Δ*rbmA* and Δ*rbmC* Δ*bap1* mutants were constructed

by allelic exchange (51) using specific suicide vectors described before (20, 52). *Vc* strains used

for microscopy experiments, Δ*vc1807*::$P_{tac}$–*mNeonGreen* and Δ*vc1807*::$P_{tac}$ –*SCFP3A-spec$^R$*,

were constructed using natural transformation as previous described (22). The *P. aminovorans*

(*Pa*) used in our experiments is a Strep$^R$ isolate derived from the ATCC type strain (ATCC #49632).

*Vc* and *Pa* overnight cultures were grown with aeration in LB at 30°C. Heat-killed strains were

incubated at 60°C for 2 hours prior to experimentation. Unless specified, media was supplemented

with streptomycin (Sm, 100 μg/ml) and chloramphenicol (Cm, 10 μg/ml) when appropriate.

**Animal studies**

For establishing colonization of the microbiota species, 3-day old suckling CD-1 mice (Charles

River Laboratories) were fasted for 1 hour, then orally dosed with *Pa* at a concentration of $10^7$ CFU

using 30-gauge plastic tubing, after which the animals were placed with a lactating dam for 10-12

hrs and monitored in accordance with the regulations of Tufts Comparative Medicine Services. This

inoculation scheme was followed an additional 3 times, for a total of 4 inoculations of *Pa* over the

course of 48hrs. After 48hrs, mice were infected with $10^6$ CFU of *Vc*, WT C6706 or mutant strain,

or LB as a vehicle control in a gavage volume of 50 μl to evaluate the effect of *Pa* pre-colonization

on *Vc* host colonization. At 18-24hrs post-infection, animals were sacrificed, and small intestine

tissue samples were collected and homogenized for CFU enumeration. WT *Vc* is *lac*$^+$ and appears

blue on medium containing X-gal while *Pa* appears white on the same medium. For coinfection

experiments, cultures of *Vc* and *Pa* strains were mixed in a 1:1 ratio and mice were orally dosed

with a final bacterial count of $10^6$ CFU. Mice were sacrificed 20-24 hours post-infection and small

intestine samples were processed as outlined above to evaluate the colonization efficiency of both

species.

**Ethics Statement**

All animal experiments were performed at and in accordance with the rules of the Tufts

Comparative Medicine Services (CMS), following the guidelines of the American Veterinary Medical

Association (AVMA) as well as the Guide for the Care and Use of Laboratory Animals of the

National Institutes of Health. All procedures were performed with approval of the Tufts University

CMS (Protocol# B 2018-99). Euthanasia was performed in accordance with guidelines provided by

the AVMA and was approved by the Tufts CMS. The previously published study from which Figure

1 is derived (7) received approval from the Ethical Review Committee at the icddr,b and the

institutional review boards of Massachusetts General Hospital and the Duke University Health

System. Participants or their guardians provided written informed consent.

**Pellicle composition analysis**

To assess pellicle composition, overnight cultures of *Vc* and *Pa* were inoculated into glass culture

tubes (18 x 150 mm) containing 2mL LB media in a ratio of 1:10 *Vc* ($10^6$) to *Pa* ($10^7$) CFU, and co-

cultures were allowed to grow statically at room temperature for 3 days. Following static growth,

floating pellicles were carefully transferred into sterile 1.5mL Eppendorf tubes containing 1mL LB,

and samples were gently spun down to wash away any planktonic bacteria. Planktonic cells were

removed, and cell pellets of pellicle samples were resuspended in 1mL of fresh LB media. All

samples including supernatant from the pellicle wash step, were serial diluted and plated on Sm/X-

Gal media to differentiate *Vc* (blue) and *Pa* (white) colonies.

**Crystal violet biomass assays**

Crystal violet biofilm assays were performed as described previously in 96-well flat bottom clear,

tissue-culture treated polystyrene microplates (ThermoFisher) (23). In each well, *Vc* ($10^6$ CFUs)

and/or *Pa* ($10^6$ or $10^7$ CFUs) were inoculated into 200µL of medium. For experiments involving

spent culture supernatants, *Vc* ($10^6$ CFUs) were inoculated into each well containing 200 µL of

reconditioned supernatants (80% (v/v) filtered spent culture medium and 20% (v/v) 5× LB). Plates

were then sealed using a gas permeable sealing film (BrandTech) and incubated at 37°C.

Planktonic culture was removed after 24 hours of incubation, plates were washed with distilled

water once. Attached biofilms were stained with 0.1% crystal violet at room temperature for 15-20

min. The amount of biomass adhered to the sides of each well was quantified by dissolving the

crystal violet in 95% ethanol and the absorbance of the resulting solution was measured at 550 nm

or 570 nm using a plate reader.

**Microscopy**

Liquid LB culture of *Vc*, *Pa*, and co-cultures (*Pa*:*Vc* = 10:1) were prepared according to procedures described above. To image pellicles, we used a modified literature procedure (53). Monocultures and co-culture pellicles were first prepared following the procedure described above, except that 3 mL of the culture was incubated in a 5 mL culture tube. After 3 days of incubation at room temperature, the pellicles were carefully picked up by the large end of a 200 µL pipette tip, transferred to a coverslip (22 x 60 mm, No. 1.5), and immediately covered with another square coverslip to prevent drying. The LB medium contained 4 µg/mL FM 4-64 stain (ThermoFisher) to stain all cells. To stain VPS, the LB medium additionally contained 4 µg/mL of Wheat Germ Agglutinin conjugated to Oregon Green (ThermoFisher). The stained biofilms were imaged with a Nikon-W1 confocal microscope using 60× water objective (numerical aperture = 1.20). The imaging window was $221 \times 221\ \mu m^2$. For large-scale view, a 5x5 tiling was performed. For zoom-in view, the *z*-step size was 0.5 µm and the pixel size was 108 nm. For large-scale view, the *z*-step size was 1 µm and the pixel size was 216 nm.  The mNeonGreen (or SCFP3A) expressed by *Vc* was imaged at 488 nm (or 445 nm) excitation, FM 4-64 at 561 nm, and WGA-Oregon Green at 488 nm with the corresponding filters. All presented images are raw images processed from Nikon Element software.

**Statistics**

Error bars in the figures depict the median with a 95% confidence interval as indicted. Based on the experimental design, either standard *t*-test or Mann-Whitney test were used to compare treatment groups as indicated in each figure legend.

**Acknowledgments**

We thank members in the Ng and Weil Labs for helpful discussions. We acknowledge Ed Ryan for

his assistance in reviewing the manuscript. A.A.W and W-L.N. received support from a Rozan

Award from Tufts University School of Medicine for this project. A.A.W. support was provided by

AI123494 from the National Institute of Allergy and Infectious Diseases (NIAID). W-L.N was

supported by AI121337. J.Y. holds a Career Award at the Scientific Interface from the Burroughs

Wellcome Fund.

**References**

1. Camacho A*, et al.* (2018) Cholera epidemic in Yemen, 2016-18: an analysis of
surveillance data. *Lancet Glob Health* 6(6):e680-e690.

2. Luquero FJ*, et al.* (2016) Mortality Rates during Cholera Epidemic, Haiti, 2010-
2011. *Emerg Infect Dis* 22(3):410-416.

3. Weil AA & Ryan ET (2018) Cholera: recent updates. *Curr Opin Infect Dis*
31(5):455-461.

4. Alavi S*, et al.* (2020) Interpersonal Gut Microbiome Variation Drives Susceptibility
and Resistance to Cholera Infection. *Cell* 181(7):1533-1546.e1513.

5. Hsiao A*, et al.* (2014) Members of the human gut microbiota involved in recovery
from *Vibrio cholerae* infection. *Nature* 515(7527):423-426.

6. Levade I*, et al.* (2020) Predicting *Vibrio cholerae* infection and disease severity
using metagenomics in a prospective cohort study. *J Infect Dis*.

7. Midani FS*, et al.* (2018) Human Gut Microbiota Predicts Susceptibility to *Vibrio*
*cholerae* Infection. *J Infect Dis*.

8. McKenney PT & Pamer EG (2015) From Hype to Hope: The Gut Microbiota in
Enteric Infectious Disease. *Cell* 163(6):1326-1332.

9. Freter R (1955) The fatal enteric cholera infection in the guinea pig, achieved by
inhibition of normal enteric flora. *J Infect Dis* 97(1):57-65.

10. Nygren E, Li BL, Holmgren J, & Attridge SR (2009) Establishment of an Adult
Mouse Model for Direct Evaluation of the Efficacy of Vaccines against *Vibrio*
*cholerae*. *Infection and Immunity* 77(8):3475.

11. Zhao W, Caro F, Robins W, & Mekalanos JJ (2018) Antagonism toward the
intestinal microbiota and its effect on *Vibrio cholerae* virulence. *Science*
359(6372):210-213.

12. Jung SA, Chapman CA, & Ng WL (2015) Quadruple quorum-sensing inputs control
*Vibrio cholerae* virulence and maintain system robustness. *PLoS Pathog*
11(4):e1004837.

13. Watve S*, et al.* (2020) Parallel quorum-sensing system in *Vibrio cholerae* prevents
signal interference inside the host. *PLoS Pathog* 16(2):e1008313.

14. You JS*, et al.* (2019) Commensal-derived metabolites govern *Vibrio cholerae*
pathogenesis in host intestine. *Microbiome* 7(1):132.

15. David LA*, et al.* (2015) Gut microbial succession follows acute secretory diarrhea
in humans. *mBio* 6(3):e00381-00315.

16. Yatsunenko T*, et al.* (2012) Human gut microbiome viewed across age and
geography. *Nature* 486(7402):222-227.

17. Urakami T, Araki H, Oyanagi H, Suzuki K, & Komagata K (1990) *Paracoccus*
*aminophilus* sp. nov. and *Paracoccus aminovorans* sp. nov., which utilize N,N-
dimethylformamide. *International journal of systematic bacteriology* 40(3):287-291.

| 535 | 18. | Klose KE (2000) The suckling mouse model of cholera. *Trends Microbiol* 8(4):189-191. |
| 536 | | |

535    18.    Klose KE (2000) The suckling mouse model of cholera. *Trends Microbiol* 8(4):189-
536            191.

537    19.    Millet YA*, et al.* (2014) Insights into *Vibrio cholerae* intestinal colonization from
538            monitoring fluorescently labeled bacteria. *PLoS Pathog* 10(10):e1004405.

539    20.    Nadell CD, Drescher K, Wingreen NS, & Bassler BL (2015) Extracellular matrix
540            structure governs invasion resistance in bacterial biofilms. *ISME J* 9(8):1700-1709.

541    21.    Shaner NC*, et al.* (2013) A bright monomeric green fluorescent protein derived
542            from *Branchiostoma lanceolatum*. *Nat Methods* 10(5):407-409.

543    22.    Dalia AB, McDonough E, & Camilli A (2014) Multiplex genome editing by natural
544            transformation. *Proc Natl Acad Sci U S A* 111(24):8937-8942.

545    23.    O'Toole GA (2011) Microtiter dish biofilm formation assay. *J Vis Exp* (47).

546    24.    Yildiz FH & Schoolnik GK (1999) *Vibrio cholerae* O1 El Tor: identification of a gene
547            cluster required for the rugose colony type, exopolysaccharide production, chlorine
548            resistance, and biofilm formation. *Proc Natl Acad Sci U S A* 96(7):4028-4033.

549    25.    Yildiz F, Fong J, Sadovskaya I, Grard T, & Vinogradov E (2014) Structural
550            characterization of the extracellular polysaccharide from *Vibrio cholerae* O1 El-Tor.
551            *PLoS One* 9(1):e86751.

552    26.    Berk V*, et al.* (2012) Molecular architecture and assembly principles of *Vibrio
553            cholerae* biofilms. *Science* 337(6091):236-239.

554    27.    Tamayo R, Patimalla B, & Camilli A (2010) Growth in a biofilm induces a
555            hyperinfectious phenotype in *Vibrio cholerae*. *Infect Immun* 78(8):3560-3569.

556    28.    Gallego-Hernandez AL*, et al.* (2020) Upregulation of virulence genes promotes
557            *Vibrio cholerae* biofilm hyperinfectivity. *Proc Natl Acad Sci U S A* 117(20):11010-
558            11017.

559    29.    Zhu J & Mekalanos JJ (2003) Quorum sensing-dependent biofilms enhance
560            colonization in *Vibrio cholerae*. *Dev Cell* 5(4):647-656.

561    30.    Fong JCN, Syed KA, Klose KE, & Yildiz FH (2010) Role of Vibrio polysaccharide
562            (*vps*) genes in VPS production, biofilm formation and *Vibrio cholerae*
563            pathogenesis. *Microbiology (Reading)* 156(Pt 9):2757-2769.

564    31.    Fong JC, Karplus K, Schoolnik GK, & Yildiz FH (2006) Identification and
565            characterization of RbmA, a novel protein required for the development of rugose
566            colony morphology and biofilm structure in *Vibrio cholerae*. *J Bacteriol*
567            188(3):1049-1059.

568    32.    Fong JC & Yildiz FH (2007) The *rbmBCDEF* gene cluster modulates development
569            of rugose colony morphology and biofilm formation in *Vibrio cholerae*. *J Bacteriol*
570            189(6):2319-2330.

571    33.    Absalon C, Van Dellen K, & Watnick PI (2011) A communal bacterial adhesin
572            anchors biofilm and bystander cells to surfaces. *PLoS Pathog* 7(8):e1002210.

573    34.    Ubeda C, Djukovic A, & Isaac S (2017) Roles of the intestinal microbiota in
574            pathogen protection. *Clinical & Translational Immunology* 6(2):e128.

35. Weil AA, Becker RL, & Harris JB (2019) *Vibrio cholerae* at the Intersection of
Immunity and the Microbiome. *mSphere* 4(6):e00597-00519.

36. Yoon MY*, et al.* (2016) A single gene of a commensal microbe affects host
susceptibility to enteric infection. *Nat Commun* 7:11606.

37. Wang Y*, et al.* (2011) The Prevalence of Functional Quorum-Sensing Systems in
Recently Emerged *Vibrio cholerae* Toxigenic Strains. *Environ Microbiol Rep*
3(2):218-222.

38. Wang H*, et al.* (2021) Contributions of *Escherichia coli* and its motility to the
formation of dual-species biofilms with *Vibrio cholerae*. *Appl Environ*
*Microbiol*:AEM0093821.

39. Toh YS*, et al.* (2019) Role of coaggregation in the pathogenicity and prolonged
colonisation of *Vibrio cholerae*. *Med Microbiol Immunol* 208(6):793-809.

40. Cont A, Rossy T, Al-Mayyah Z, & Persat A (2020) Biofilms deform soft surfaces
and disrupt epithelia. *Elife* 9:e56533.

41. Sadiq FA*, et al.* (2021) Synergistic interactions prevail in multispecies biofilms
formed by the human gut microbiota on mucin. *FEMS Microbiol Ecol* 97(8).

42. Teschler JK*, et al.* (2015) Living in the matrix: assembly and control of *Vibrio*
*cholerae* biofilms. *Nat Rev Microbiol* 13(5):255-268.

43. Papenfort K*, et al.* (2017) A *Vibrio cholerae* autoinducer-receptor pair that controls
biofilm formation. *Nat Chem Biol* 13(5):551-557.

44. Christensen BB, Haagensen JA, Heydorn A, & Molin S (2002) Metabolic
commensalism and competition in a two-species microbial consortium. *Appl*
*Environ Microbiol* 68(5):2495-2502.

45. Mark Welch JL, Rossetti BJ, Rieken CW, Dewhirst FE, & Borisy GG (2016)
Biogeography of a human oral microbiome at the micron scale. *Proc Natl Acad Sci*
*U S A* 113(6):E791-800.

46. Yoshida K, Toyofuku M, Obana N, & Nomura N (2017) Biofilm formation by
*Paracoccus denitrificans* requires a type I secretion system-dependent adhesin
BapA. *FEMS Microbiol Lett* 364(4).

47. Srinandan CS, Jadav V, Cecilia D, & Nerurkar AS (2010) Nutrients determine the
spatial architecture of *Paracoccus* sp. biofilm. *Biofouling* 26(4):449-459.

48. Jousset A*, et al.* (2017) Where less may be more: how the rare biosphere pulls
ecosystems strings. *ISME J* 11(4):853-862.

49. Thelin KH & Taylor RK (1996) Toxin-coregulated pilus, but not mannose-sensitive
hemagglutinin, is required for colonization by *Vibrio cholerae* O1 El Tor biotype
and O139 strains. *Infect Immun* 64(7):2853-2856.

50. Waters CM, Lu W, Rabinowitz JD, & Bassler BL (2008) Quorum sensing controls
biofilm formation in *Vibrio cholerae* through modulation of cyclic di-GMP levels and
repression of *vpsT*. *J Bacteriol* 190(7):2527-2536.

51. Skorupski K & Taylor RK (1996) Positive selection vectors for allelic exchange.
*Gene* 169(1):47-52.

52.    Yan J, Sharo AG, Stone HA, Wingreen NS, & Bassler BL (2016) *Vibrio cholerae*
617           biofilm growth program and architecture revealed by single-cell live imaging. *Proc*
618           *Natl Acad Sci U S A* 113(36):E5337-5343.

619    53.    Fiebig A (2019) Role of Caulobacter Cell Surface Structures in Colonization of the
620           Air-Liquid Interface. *J Bacteriol* 201(18).

**Figures and Tables**

**Figure 1. *Pa* is more abundant in persons with *Vc* infection compared to**

**uninfected persons.** In a prior study of household contacts of cholera patients in

Bangladesh (7), *Pa* was identified as differentially abundant using a support vector

machine model with recursive feature elimination in order to discriminate patterns of

microbial taxa relative abundance that distinguished infected from uninfected persons.

The microbiota was assessed using 16S rRNA in rectal swabs collected from individuals

with *Vc* infection ($n$ = 22) compared to uninfected individuals ($n$ = 36). In this study, total

sum normalization was applied to OTU counts from each sample, and a median of

37,958 mapped reads per sample were generated (7). Based on this sequencing data,

the estimated limit of detection for a *Pa* OTU is $2.0\times10^{-5}$. **(A)** Relative abundance *Pa* in

infected and uninfected individuals and **(B)** ratio of *Vc* to *Pa* in six *Vc*-infected persons.

All data points are shown and boxes indicate interquartile range. Bars mark the

maximum and minimum values. These values were compared with non-parametric

unpaired Mann-Whitney U testing, $P$ <0.01.

.

**Figure 2. The presence of *Pa* enhances *Vc* colonization in the infant mouse**

**intestine. (A)** 3 day-old infant mice were intragastrically inoculated with $10^7$ CFU of *Pa*

every 12 hours for a period of 24 hr (2 doses) or 48 hr (4 doses). At each time point mice

were sacrificed and CFU were enumerated by plating serial dilutions of small intestine

samples on selective media. **(B)** 3 day-old infant mice were intragastrically inoculated 4

times with LB or $10^7$ CFU of *Pa* for every 12 hours, and subsequently infected with $10^6$

CFU of WT *Vc*. Mice were sacrificed 20-24 hr post-infection and the small intestine

samples were taken to enumerate *Vc*. Bars on graphs depict median value with 95%

confidence interval (CI) and individual data points plotted. Unpaired non-parametric *t*-test

(Mann-Whitney); *** $P \leq 0.0001$. (C) *Vc* was inoculated intragastrically into the animals

alone or together with *Pa* in a 1:1 ratio. After 24 hours, enumeration of *Vc* was

656    performed as described above. Unpaired non-parametric $t$-test (Mann-Whitney); ** $P \leq$

657    0.005.

**Figure 3. *Pa* promotes biofilm formation of *Vc*. (A)** Planktonic cell counts from

cultures used for pellicle analysis of *Vc* and *Pa* grown together or in monoculture. **(B)**

Representative images of pellicles formed by *Vc* grown in monoculture and in co-culture

with *Pa*. CFU counts of each strain in **(C)** pellicle samples and **(D)** spent medium used to

wash the pellicle. Bars on graphs depict median value with 95% confidence interval (CI)

and individual data points plotted. Unpaired non-parametric *t*-test (Mann-Whitney): ** *P* ≤

0.005.

**Figure 4. Representative microscopy images of *Vc* and *Pa* dual-species pellicles.**

**(A)** Large-scale cross-sectional image of the internal structure in a co-culture pellicle. All

cells are stained with FM 4-64 and *Vc* cells constitutively express mNeonGreen.

Therefore, the red signal in the overlay image corresponds primarily to *Pa* cells. **(B)**

Zoom-in view of the region highlighted in A. **(C)** Zoom-in view of the region highlighted in

B. **(D)** Cross-sectional views of the region shown in C, at the bottom of the pellicle. *Pa*

cells exist mainly at the pellicle-liquid interface, with clusters of *Pa* cells penetrating into

the interior of the pellicle. **(E-F)** *Top* (E) and *Bottom* (F) view of the co-culture structure

shown in B, rendered in 3D.

**Figure 5. *Pa* increases biofilm production in *Vc*.** Crystal violet assays were performed

in 96-well microtiter plates to quantify biofilm formation. Overnight-grown **(A)** wild-type

*Vc* or **(B)** Δ*vpsL* mutant and *Pa* cultures were diluted to a final concentration of $10^6$ CFU

in a total volume of 200 μL/well. In samples containing a 1:10 ratio of *Vc*/*Pa*, *Pa* was

diluted to a final concentration of $10^7$ CFU. Samples with heat-killed (HK) *Pa* are

specified in the x-axis. Microtiter plates were incubated at 37 °C for 24 hr. Crystal violet

staining and ethanol solubilization were performed as previously described (23).

Absorbance of the crystal violet stain was measured at 550 nm using a Biotek Synergy

HTX plate reader. Data is represented with horizontal lines indicating the mean with

standard deviation. Unpaired *t*-test (Mann-Whitney); **$P \le 0.005$.

**Figure 6. *Vc*/*Pa* co-culture biofilms depend on VPS. (A)** Representative top view of a
co-culture pellicle with WGA staining, rendered in 3D. *Vc* cells constitutively express
SCFP3A cytosolically; all cells were stained with FM 4-64 membrane stain; WGA is

conjugated to Oregon Green and shown in yellow. Note that WGA also stained dead

cells with an exposed peptidoglycan layer, corresponding to the bright spots in the

image. Scale bar: 100 µm. **(B)** Zoom-in view of the three regions 1-3 indicated by the

white boxes in A. Shown from *left* to *right* are *Vc* cell fluorescence (SCFP3A), membrane

staining (FM 4-64), WGA staining (Oregon Green), and the overlay of the three

channels. The overlaid images additionally show the cross-sectional view in the *xz* and

*yz* planes. Region 1 contains primarily *Vc* cells, and regions 2 and 3 contain *Pa*-enriched

regions. *Pa*-enriched regions demonstrate elevated WGA signal. Intensities in each

channel were kept consistent through region 1-3 for comparison.  Scale bars: 20 µm. **(C)**

Zoom-in view of the highlighted regions in B (white boxes, WGA channel only).

Intensities are adjusted to similar level for visualization of the internal structure.   Scale

bar: 5 µm.

**Figure 7. Enhanced *Vc* intestinal colonization in the presence of *Pa* is dependent**

**on VPS and accessory matrix proteins. (A)** 3 day-old infant mice were intragastrically

inoculated with LB or $10^7$ CFU of *Pa* every 12 hours for a period of 48 hours, and

subsequently infected with $10^6$ CFU of a *Vc* strain defective for extracellular matrix

production (Δ*vpsL*). Mice were sacrificed 20-24 hr post-infection and small intestine

samples were taken to enumerate *Vc* CFU. Data from infection with the wild-type *Vc*

strain (Figure 2B) is shown again here for comparison purposes. **(B)** *Vc* wild-type (WT)

or different biofilm mutants were mixed with *Pa* in 1:1 ratio, and the mixture was used

immediately for animal infection. Mice were sacrificed 20-24 hr post-infection and small

intestine samples were taken to enumerate *Vc* CFU. Each symbol represents an

individual mouse and data is represented with horizontal lines indicating the median with

a 95% confidence interval. Unpaired non-parametric $t$-test (Mann-Whitney); ns $P > 0.05$,

*** $P \leq 0.001$. Data from infection with the wild-type $Vc$ strain (Figure 2C) is shown again

here for comparison purposes. **(C)** Crystal violet assays performed in 96-well plates to

quantify pellicle formation. Overnight cultures of $Vc$ $rbmA^-$, $rbmC^-$ $bap1^-$ and $Pa$ were

diluted in fresh LB and plated as 200 μL/well. Samples were co-cultured in either 1:10

ratios of Vc/Pa and incubated at 37 °C for 24 hr. Crystal violet staining was then

performed and absorbance of the stain was measured at 570 nm. Horizontal lines

indicating mean with standard deviation are shown. Unpaired $t$-test; ****$P \leq 0.0001$, **$P$

$\leq 0.05$.

**Supplemental Figures**

**Supplemental Figure 1.** *P. aminovorans* **colonization does not significantly alter**

**the mouse gut microbial diversity.** Three day-old infant mice were intragastrically

inoculated with $10^7$ CFU of *Pa* or sterile LB medium as control (Ctrl) for a period of 12

hours to mimic the length of time between the final *Pa* inoculation and *Vc* infection in

mouse experiments. Small intestines were removed, homogenized, and DNA was

extracted and sequenced using shallow shotgun sequencing. **(A)** Relative abundance of

phylum-level microbes in the small intestines of mice and **(B)** principal component of

analysis using Bray Curtis Dissimilarity. N=3 per group. *Pa* is in the Proteobacteria

phylum. Two-way ANOVA testing of phylum level-abundance was performed, and all

comparisons are not significantly different (P>0.05).

**Supplemental Figure 2. *P. aminovorans* (*Pa*) colonization significantly increases *V.***
**cholerae (*Vc*) small intestine colonization.** Three day-old infant mice were
intragastrically inoculated with $10^7$ CFU of *Pa* or sterile LB medium as control for a
period of 36 hours. These animals were then infected with *Vc* 48 hours after the first
inoculation. At different time points, small intestines were removed, homogenized, and
*Vc* were enumerated. Student t-tests were used to analyze these samples. P values are
0.14, 0.035, and 0.004 at 6, 12, and 24 hours respectively (Unpaired non-parametric *t*-
test).

756
757
**Supplemental Figure 3. Established *Pa* cultures increase biofilm production in *Vc*.**

Biofilm formation assays were performed in 96-well microtiter plates. *Pa* was diluted in

LB to a concentration of $10^7$ CFU and grown for 24hr before the addition of WT *Vc*. WT

*Vc* was then diluted in LB to a concentration of $10^6$ CFU and added to wells. Plates were

incubated for an additional 24hr before crystal violet staining to quantify biofilm biomass.

Crystal violet staining and ethanol solubilization were performed as previously described

(23). Absorbance of the crystal violet stain was measured at 550 nm using a Biotek

Synergy HTX plate reader. Samples with heat-killed (HK) *Pa* are delineated by hatched

bars. Data is represented with horizontal lines indicating the median with 95%

confidence interval. Unpaired non-parametric t-test (Mann-Whitney); * $P \leq 0.05$.