Combined influence of food availability and agricultural intensification on a declining aerial

insectivore

Daniel R. Garrett*[1,2], Fanie Pelletier[1], Dany Garant[1], and Marc Bélisle[1]

Author's Affiliations

[1]Département de biologie, Université de Sherbrooke, 2500 Boulevard de l'Université,

Sherbrooke, Québec J1K2R1 Canada

[2]daniel.garrett@usherbrooke.ca

*Corresponding author: Université de Sherbrooke, 2500 boulevard de l'Université, Sherbrooke,

Qc, Canada, J1K 2R1, daniel.garrett@usherbrooke.ca

Running Head: Food, agriculture, and a declining bird

Abstract

Aerial insectivores show worldwide population declines coinciding with shifts in agricultural practices. Increasing reliance on certain agricultural practices is thought to have led to an overall reduction in insect abundance that negatively affects aerial insectivore fitness. The relationship between prey availability and the fitness of insectivores may thus vary with the extent of agricultural intensity. It is therefore imperative to quantify the strength and direction of these associations. Here we used data from an 11-year study monitoring the breeding of Tree Swallows (*Tachycineta bicolor*) and the availability of Diptera (their main prey) across a gradient of agricultural intensification in southern Québec, Canada. This gradient was characterized by a shift in agricultural production, whereby landscapes composed of forage and pastures represented less agro-intensive landscapes and those focusing on large-scale arable row crop monocultures, such as corn (*Zea mays*) or soybean (*Glycine max*) that are innately associated with significant mechanization and agro-chemical inputs, represented more agro-intensive landscapes. We evaluated the landscape characteristics affecting prey availability, and how this relationship influences the fledging success, duration of the nestling period, fledgling body mass, and wing length as these variables are known to influence the population dynamics of this species. Diptera availability was greatest within predominately forested landscapes, while within landscapes dominated by agriculture, it was marginally greater in less agro-intensive areas. Of the measured fitness and body condition proxies, both fledging success and nestling body mass were positively related to prey availability. The impact of prey availability varied across the agricultural gradient as fledging success improved with increasing prey levels within forage landscapes yet declined in more agro-intensive landscapes. Finally, after accounting for prey availability, fledging success was lowest, nestling periods were the longest, and wing length

of fledglings were shortest within more agro-intensive landscapes. Our results highlight the

interacting roles that aerial insect availability and agricultural intensification have on the fitness

of aerial insectivores, and by extension how food availability may interact with other aspects of

breeding habitats to influence the population dynamics of predators.

Key words: aerial insectivores; agricultural intensification; Diptera; fledging success; food

resources; nestling condition; resource availability; *Tachycineta bicolor*; Tree Swallows.

Introduction

Food availability is a key determinant of population size and dynamics (Hailman and Lack 1955, McCamey and Slobodkin 1963, Newton 1998). The amount of food resources an individual acquires from a given foraging effort affects the amount of energy it can allocate to both its survival and reproduction, thereby affecting vital rates (Martin 1987, Newton 1998, 2002, Stephens et al. 2007). Food reduction or supplementation experiments are often used to quantify the impact of food availability on reproductive output (Martin 1987, Seward et al. 2014, Ruffino et al. 2014, Seress et al. 2020). For example, *Bacillus thuringiensis* (Bt) insecticides are used to reduce insect abundance and both experimental and observational studies typically report reduced fitness of insectivorous predators following their application (Rodenhouse and Holmes 1992, Nagy and Smith 1997, Marshall et al. 2002, Poulin et al. 2010). Other studies, conducted under unaltered levels of food availability have reported no link with fitness outcomes of consumers (Dawson and Bortolotti 2000, Imlay et al. 2017, McClenaghan et al. 2019). Such contrasting results can be explained by food being overly abundant and thus other environmental (e.g., extreme temperature or elevated precipitation) or endogenous factors (e.g., parental condition, care, and past experience), having a greater influence than food availability in limiting reproductive output and subsequent population growth (Martin 1987, Newton 1998, White 2008, Winkler et al. 2013, Quinney 2019). Studies of naturally occurring predator-prey interactions have frequently been conducted within single landscapes, ones already suitable for elevated fitness, or have been conducted without reference to human dominated landscapes (Martin 1987, Fazey et al. 2005, Robb et al. 2008, Martin et al. 2012). Habitat induced variability in prey from such studies may thus not be great enough to influence the fitness of predators.

Human-driven landscape modifications are known to affect both the magnitude and direction of

a broad spectrum of ecological processes occurring over a multitude of spatio-temporal scales

(Rolstad 1991, Miller and Adams 1995, Fischer and Lindenmayer 2007, Davis 2007). Therefore,

studies disregarding such impacts may overlook confounding or interacting factors of foraging

within anthropogenic landscapes. For instance, populations of pest insects may be higher in

landscapes with an abundance of large scale monocultures (Andow 1983, 1991). Yet in these

same landscapes, the availability of other crucial resources may be limited, possibly influencing

either the lifetime or annual fitness of animals continuing to perceive them as suitable breeding

or foraging habitats (Robertson 2012, Hale et al. 2015, Letten et al. 2017). The historical removal

of forests and replacement of grasslands and wetlands with agricultural lands, including the

simplification of agricultural landscapes and a shift from mixed crop farming to livestock or

highly-mechanized row-cropping supported by significant agro-chemical inputs, a process

known as agricultural intensification, is one of the most prominent driver of human-induced

landscape change (Lambin et al. 2001, Fischer and Lindenmayer 2007). In North America, this

process is exemplified by the trend of agricultural landscapes transitioning from small scale

perennial crops and low impact farming practices, to being increasingly composed of only a few

arable row crop monocultures (e.g., corn (*Zea mays*) or soybean (*Glycine max*)), and the near

ubiquitous use of the aforementioned agro-intensive practices (Stanton et al. 2018).

Despite agricultural intensification being linked to the decline of many vertebrates (Ceballos

2002, Rosenberg et al. 2019), the relationship between observed prey availability and the fitness

of predators has rarely been investigated with respect to agricultural cover. For example, several

species of grassland and farmland birds have displayed significant declines since the 1970s

(Vickery et al. 2004, Boatman et al. 2004, Stanton et al. 2018, Rosenberg et al. 2019). In

particular, steep declines are observed in many aerial insectivores, a guild of birds foraging

nearly exclusively on aerial insects (Nebel et al. 2010, Michel et al. 2016, NABCI 2019). Due in

part to the spatio-temporal correlations between these phenomena, a common hypothesis is that

aerial insectivore declines are due in part to agricultural intensification influencing the

abundance and species composition of the insects consumed by these birds (Stanton et al. 2018,

Twining et al. 2018, Spiller and Dettmers 2019). Multiple lines of evidence exist linking changes

in specific insect communities, abundance, behavior or phenology to agricultural intensification

(Benton et al. 2002, Thomas 2004, Pisa et al. 2015, Stanton et al. 2018, Montgomery et al. 2020).

Yet, the link between agricultural intensification and aerial insectivore declines, through a

reduction in the availability of food resources, has so far yielded inconsistent results (Evans et al.

2007, Nocera et al. 2012, Imlay et al. 2017, Stanton et al. 2017, McClenaghan et al. 2019).

Within agricultural landscapes, practices such as large-scale, simplified arable row-cropping,

increased reliance on agro-chemical inputs (i.e., pesticides and fertilizers) as well as the removal

of both natural and marginal habitats are suspected to be the primary drivers of changes of insect

populations (Tscharntke et al. 2005, Grüebler et al. 2008, Attwood et al. 2008, Montgomery et al.

2020). Furthermore, within heavily cultivated landscapes, the same mechanisms reducing insect

abundance, may also impact insectivorous birds, either irrespective of, or in conjunction with,

levels of prey. For example, due to the increase of pesticides, insectivorous birds breeding within

landscapes dominated by agro-intensive monocultures may not only be exposed to decreased

prey availability, but also to elevated toxicological loads of food items (Morrissey et al. 2015,

Montiel-León et al. 2019, Malaj et al. 2020, Poisson et al. 2021). Breeding within agro-intensive

landscapes may thus have the potential to impact fitness not only through a trophic pathway, via

variation in prey availability, but also through other indirect pathways linked to agricultural

intensification (Stanton et al. 2018).

Here, we present results of an 11-year study monitoring both the reproductive attempts of Tree

Swallows (*Tachycineta bicolor*) and the availability of their aerial insect prey throughout a nest

box network located along a gradient of agricultural intensification in southern Québec, Canada.

Tree Swallows breed over much of North America and demonstrate population declines similar

to those of most other aerial insectivores, especially in the north eastern portion of their breeding

range (Shutler et al. 2012, Michel et al. 2016). Our principal objectives were to assess evidence

with respect to three hypotheses. The first was that a gradient of agricultural intensification

influences the availability of aerial insectivore prey, and we predicted that landscapes with

increasing levels of arable row crops would result in a reduction in prey availability (Benton et

al. 2002, Thomas 2004, Pisa et al. 2015, Stanton et al. 2018, Montgomery et al. 2020). The

second hypothesis was that, as aerial insectivores, breeding Tree Swallows are food limited

(Twining et al. 2018, Spiller and Dettmers 2019). Therefore, increased prey availability should

result in greater fitness, including the number of nestlings surviving until fledging, reduced

nestling duration (due to increased growth rates), and elevated nestling condition. Our third

hypothesis was that Tree Swallow fitness is influenced by agricultural landcover through more

than just the overall abundance of food, for example through pesticide exposure (Poisson et al.

2021) or variation in diet quality (Stanton et al. 2018, Twining et al. 2018). We therefore

predicted that, after controlling for prey availability, breeding attempts within landscapes

occupied by increasing row crops should result in decreased fitness.

Population dynamics of short lived species are thought to be driven by adult and juvenile return

rates and to a lesser extent fecundity (Sæther and Bakke 2000). Within Tree Swallow

populations, adult survival appears to be relatively stable throughout their distribution (Weegman

et al. 2017, Clark et al. 2018). Furthermore, recent evidence showed that, due to their high

elasticity, population viability of this species could largely be determined by fledgling survival

and recruitment rates (Jones et al. 2017, Weegman et al. 2017, Cox et al. 2018). Although food

limitation can be influential throughout the life cycle (Hostetler et al. 2015, Knight et al. 2019),

given the above, we focus primarily on the nestling period, because factors influencing fledgling

number and post fledging survival have a significant impact on population growth (Cox et al.

2018). We therefore assess the influence of prey availability on the fledging success, the duration

of the nestling period, and the morphological traits of fledglings (hitherto referred collectively as

fitness and body condition proxies). Evidence suggests food availability and agricultural

landscapes can be drivers of both nestling growth and condition, and yet they have received less

attention than annual reproductive output as a possible driver of aerial insectivore declines

(Smith and Bruun 1998, Granbom and Smith 2006, Pigeon et al. 2013, Almasi et al. 2015,

Kusack et al. 2020, Houle et al. 2020). Therefore, we assess the possibility that links between

agricultural intensification and declining populations of aerial insectivores are not only through a

reduction in the number of fledglings, but also through reduced offspring condition, as this may

carry over and affect future recruitment and productivity (Stutchbury et al. 2011, Naef-Daenzer

and Grüebler 2016, Saino et al. 2018, Evans et al. 2020, Jones and Ward 2020).

Methods

Study area and nest box system

Between 2006 and 2016, we monitored breeding attempts of Tree Swallows within 400 nest

boxes dispersed throughout a gradient of agricultural intensification in southern Quebec, Canada.

Ten nest boxes were spaced approximately 50 m apart along the field margins of 40 different

farms, located in various agricultural landscape contexts (Figure 1; see Ghilain and Bélisle 2008

for details). The gradient of agricultural intensification was characterized by an east-west shift of

agricultural production. While the eastern portion of the system was composed primarily of

pastures and forage crops (e.g., hay, alfalfa (*Medicago sativa*), and clover (*Trifolium spp.*))

interspersed with large expanses of forest cover, the west was dominated by large-scale

monocultures (principally corn, soybean, and wheat (*Triticum spp.*)) and was denuded of forest

cover (Bélanger and Grenier 2002, Jobin et al. 2005, Ruiz and Domon 2009). This increased use

of monocultures has resulted in a near reliance on fertilizers, pesticides, mechanization and

drainage of surface waters and wetlands (Jobin et al. 2003). Furthermore, between 2011 to 2019,

approximately 100% of the corn and 60% of the soybean were sown as neonicotinoid-coated

seeds in our study area (MDDELCC 2015). Subsequently, within the water bodies of the western

part of our study area, neonicotinoids, as well as many other pesticides, are frequently detected at

levels threatening to aquatic life if chronically exposed (Giroux 2019, Montiel-León et al. 2019).

Nest box monitoring and nestling condition

Each year, nest boxes were monitored prior to the arrival of Tree Swallows and then every two

days throughout the breeding season (Appendix S1: Figure S1). This detailed monitoring

allowed us to accurately estimate the dates of laying, incubation, hatching, and fledging, as well

as record the number of eggs, nestlings, and fledglings of each breeding event. We caught and

banded adult females during incubation and adult males during food provisioning. We were 99%

and 80% successful, respectively, at capturing targeted adults. Once caught, adults received an

aluminum US Geological Survey (USGS) band containing a unique identification code. At 12

days of age, nestlings also received a USGS band. To prevent premature fledging, following 16

days of age, the number of nestlings remaining within nest boxes were only counted, thus the

exact date each nestling fledged is unknown.

Insect samples and prey availability

Throughout the duration of this study, two insect traps were placed on each of the 40 farms

(N=80). Traps were spaced at least 250 meters apart along the central portion of each nest box

transect. The content of traps was collected every two days throughout each breeding season.

Insect traps were a combined window/water-pan flight trap. Traps were made by bisecting two

plexiglass sheets (11.5 cm x 30 cm), and then fixing them on top of a yellow bucket 15 cm deep

and 21 cm in diameter (Rioux Paquette et al. 2013, Bellavance et al. 2018). Traps were then

attached to a stake one meter above the ground and filled with approximately one liter of a

saltwater and soap solution. The content of each trap was collected by straining out insects into a

tube filled with 70% ethanol. Samples were then stored in closed boxes at room temperature until

processing.

Our interest being in prey availability during the nestling rearing period, we focused the

processing of insect samples collected between 1 June and 15 July, covering over 96% of

breeding attempts (Appendix S1: Figure S1). The processing of insect samples started by first

removing items and insects not found in the diet of Tree Swallows. This included removing

gastropods, large spiders, June bugs (Coleoptera: Melolonthinae), large Lepidoptera, and

bumblebees [(*Bombus* spp.), (Winkler et al. 2020)]. The remaining content of a trap was

composed of a wide range of insects stemming from many different habitat types (Laplante

2013) and overlapped significantly with the insects found within food boluses provided to

nestlings (Bellavance et al. 2018). In our system, 74% of the biomass and 77% of the insects

found within food boluses provided to nestlings are Diptera (Bellavance et al. 2018 and this

study), and this pattern is shared with other Tree Swallows study systems (McCarty and Winkler

1999a, Johnson and Lombardo 2000, Mengelkoch et al. 2004, Twining et al. 2018). We thus

used the biomass of Diptera within each sample as a proxy of prey availability (using the number

of Diptera resulted in qualitatively similar results; analyses not shown). We evaluated the

performance of prey biomass and fitness models when treating prey availability as all insects or

only Diptera (38% of the biomass from samples) and found models performed substantially

worse when treating prey availability as the biomass of all insects (Appendix S2: Table S1).

Moreover, we did not limit processing of Diptera to any specific size limit as nearly all major

groups of Diptera from our system can be found within the food boluses of Tree Swallows

(Bellavance et al. 2018). Once extracted, the number of individual Diptera was recorded and the

sample was placed in an oven at 60℃ for over 24 hours, ensuring no further change in biomass

occurred. Once dried, samples were weighed without delay ($\pm$ 0.0001 g). The dried biomass of

Diptera (hereinafter referred to as Diptera biomass) from these samples thus represents an

estimate of prey availability on a specific farm and year over the two days prior to the sample

collection.

Fitness and body condition proxies

We focused on four proxies of fitness and the body condition of fledglings; the proportion of a

brood that fledged (fledging success), the duration of the nestling period, and the body mass and

wing length of fledglings. Field methods did not allow us to define the fledging dates of specific

nestlings. Consequently, the duration of the nestling phase was represented by the number of

days between the maximum hatching date and mean Julian date between the first and last

nestling to leave the nest. At 16 days of age, we measured nestling mass ($\pm$ 0.01 g) and wing

lengths (± 1 mm) using a wing ruler. Mean fledging age was 19.69 ± 1.77 (SD) and the mean

number of days between the first and last nestling to leave the nest was 1.58 ± 1.08 (SD). We

therefore assumed morphological measures at 16 days of age were representative of fledgling

morphology (Zach and Mayoh 1982, McCarty 2001).

Landscape composition

Landscape composition (i.e., the relative coverage of habitats composing a given landscape)

focused on habitats Tree Swallows are hypothesized to use as indicators of nest-site selection,

and the principal habitats composing landscapes throughout this system (Rendell and Robertson

1989, 1990, Winkler et al. 2020). The relative cover of habitats was calculated within 500 m of

nest boxes. This spatial scale represents the area within which nestling-rearing female Tree

Swallows spend ~80% of their time (Elgin et al. 2020, Garrett et al. 2021) and is the largest

extent for which exhaustive annual characterization of landscapes has been performed in our

system since 2006 (Porlier et al. 2009). Moreover, Tree Swallows are often associated with

aquatic habitats and will travel distances greater than 500 m to forage over them (Elgin et al.

2020, Garrett et al. 2021). However, the relative cover of aquatic habitats within 500 m of nest

boxes and in these agricultural contexts was low 0.66% ± 1.07%. We therefore considered these

habitats as being potentially influential to the fitness of Tree Swallows (Berzins et al. 2021) and

calculated their relative cover at greater distances following the characterization of landscapes

within 500 m. Parcels representing different habitats and agricultural fields within each 500-m

buffer were delineated using orthophotos (scale 1:40 000) in QGIS (QGIS 2020). Content of

parcels were then characterized *in situ* at the end of the breeding season but before harvest in

order to facilitate their identification. This included characterizing which cultures, if any, were in

agricultural fields. Habitats and cultures were then reclassified into one of five higher order

categories. These higher order categories are forested, corn and soybean, forage fields (including

hay fields, other grasses, alfalfa, clover, pastures, and old fields), and cereals (other than corn

and soybean). We then calculated the mean percent cover of higher order habitats across the 10

nest boxes on each farm and for every year independently.

To obtain an integrative measure of the percent cover of all higher order habitats, defined as the

landscape context, we used a robust principal components analysis (PCA) for compositional data

(Filzmoser et al. 2009) to assign "site scores" to each of the farms during each year. Site scores

were defined by the values along the components of the resulting compositional PCA of each

farm during each year as fitted using the robCompositions package (Templ et al. 2011) in R

version 3.6.2 (R Core Team 2019). This method takes into consideration the principle that values

of compositional data are inherently related to one another given they must sum to one and

resulted in the calculation of 440 different landscape contexts observed throughout the duration

of the study (i.e., 40 farms x 11 years). Site scores were then assigned to each breeding attempt

and insect sample and used in all subsequent analyses as they allow for comparative

measurements of landscape contexts.

The first component (Comp.1) of the PCA explained 80.34% of the variance in landcover and

correlated positively with corn and soybean and negatively with both forage fields and forest

cover. The second component (Comp.2) explained 14.69% of the variance in landcover and

correlated negatively with forage fields and positively with forest cover (Figure 2). The first two

components of the PCA explained over 95% of the variance in our landscape characterization

dataset. We attempted to keep models as simple as possible and because forest cover has been

shown to be influential to the nest site selection and foraging behavior of this species (Rendell

and Robertson 1989, Courtois et al. 2021, Garrett et al. 2021), we included only these two

components to represent the landscape context. Landscapes expressed by maximizing Comp.1

and minimizing Comp.2 represent ones for which there is a mixture of corn, soybean, and cereals

and denuded of forest cover, and are thus referred to as mixed-row crop landscapes. Landscapes

expressed by minimizing Comp.1 and with negative Comp.2 values represent ones for which

there is an abundance of forage and pasture fields with average forest cover and are thus referred

to as forage landscapes.

Finally, to calculate the relative cover of aquatic habitats, we performed a post-hoc model

comparison determining the spatial extent to which the percent cover of open water was

influential in predicting fledging success. Each model had an identical set of covariates and

random effect structure (model: Base + Food * Land in Appendix S1: Table S1), varying only by

the percent cover of open water calculated at increasing spatial extents between 500 m to 10 km,

at 500-m increments. The model resulting in the lowest second-order Akaike information

criterion (AICc; Burnham and Anderson 2002) occurred when considering the percent cover of

open water within 3 km of a nest box. We thus included the percent cover of open water within

3 km around each breeding attempt as another landscape variable. Open water data were derived

from the annual crop inventory produced by Agriculture and Agri-Food Canada (AAFC; AAFC

2018).

Weather variables

Due to the ectothermic nature of insects, measurements of prey availability can be strongly

influenced by weather conditions during each sampling event (Winkler et al. 2013, Shipley et al.

2020). Furthermore, Tree Swallow breeding success may also be influenced by weather

conditions independently of its effects on prey availability through, for example, the induction of

hypothermic conditions of nestlings (Ardia et al. 2010). We wanted to control for these effects

by including parameters indicative of temperature and precipitation during insect sampling and

during the pertinent time windows of each breeding attempt. We therefore collected hourly

temperature with iButtons (model DS1922L; Embedded Data Systems, Lawrenceburg,

Kentucky, USA) attached to the underside of a single nest box on each farm. Temperatures from

data loggers placed on the outside of nest boxes were highly correlated to those recorded inside

of nest boxes ($r = 0.97$), and therefore we used the former as they are likely more representative

of the temperatures experienced by aerial insects and foraging adult Tree Swallows. Hourly

temperatures were recorded throughout the entirety of each breeding season and started prior to

the arrival of Tree Swallows onto the system. From these data, we derived daily summaries of

the minimum, mean, and maximum daily temperatures. We measured total farm-specific

precipitation every two days throughout the breeding season using a single pluviometer placed

on each farm and recorded measurements to the nearest 0.5 ml during each farm visit.

Statistical analyses

Seasonal influence of weather and landscape composition on prey availability

We assessed the influence of covariates on prey availability using generalized linear mixed-

effects models (GLMMs). We assessed a series of models representing alternative hypotheses in

order to evaluate if agricultural landcover is related to prey availability either additively or

through interactions with seasonality and/or weather. We calculated the mean temperature values

between the day of and day preceding the collection of insect samples, to ensure that weather

covariates were representative of the estimate of prey availability. We treated precipitation the

same way, as the bi-daily recordings already represented the accumulated precipitation during

each insect sampling period. The conditional distribution of Diptera biomass was right-skewed,

therefore GLMMs incorporated a Gamma distribution and a log link function. Fixed effects

within competing models included the site scores (Comp.1 and Comp.2) and their interaction,

Julian date, and weather variables associated with the collection of each sample. We treated the

effect of Julian date as a second-degree polynomial, expecting insect abundance to reach a peak

around the middle of the breeding season (Rioux Paquette et al. 2013, Bellavance et al. 2018).

Ad-hoc model comparison using full models (model: Base + Rain * Land + Time * Land;

Appendix S1: Table S1) highlighted that the most predictive temperature of Diptera biomass was

the mean maximum temperature over the two-day period (AICc w = 0.94). We therefore treated

temperature as the mean maximum during this period. Because we did not expect Diptera

biomass to respond linearly with temperature, as it typically rises at an increasing rate with

temperature before plateauing and subsequently declining, indicative of a third degree

polynomial, we treated the effect of temperature as such (Winkler et al. 2013). Finally,

preliminary exploration highlighted that the phenology of Diptera biomass varied markedly

among both years and farms. We hence performed further ad-hoc model comparisons of two full

models, differing only in the presence of a variable slope of Julian date, treated as a second-

degree polynomial, across years and farms through random effects. AICc weights greatly favored

this variable slope model (AICc w >0.99). All implementation of polynomial terms was with the

poly() function in R.

Evaluating the strength of evidence for the hypothesis that prey availability is related to

landscape context took an information theoretic and multimodel approach (Burnham and

Anderson 2002). Alternative hypotheses were thus evaluated using AICc and the associated

Akaike weight (w) of competing models. We hypothesized that the weather dependent nature of

insects may result in the influence of the landscape context on prey availability being either

masked by or vary with the timing of sampling and weather conditions around sampling time

(i.e., in the two days prior to insect collection). We accordingly defined a set of six candidate

models including a null model exempt of all fixed effects (Appendix S1: Table S1). Two models

treated fixed effects as strictly additive, including one where prey availability varied only with

sampling time and weather (Base), and one with the landscape context, while controlling for both

sampling time and weather (Base + Land). We also considered models where the effect of

landscape context on prey availability varied either with the sampling time (Base + Time *

Land), the amount of precipitation (Base + Rain * Land) or both the sampling time as well as the

amount of precipitation (Base + Rain * Land + Time * Land).

Influence of prey availability and agricultural landcover on fitness and body condition proxies

We reduced analyses to first breeding attempts (90% of all attempts and ranging between 80%

and 95% across years), as reproductive success can vary greatly between first and second

attempts, and as second attempts occur nearly exclusively following the failure of a nest during

laying or incubation. Fledging success was modeled as the number of fledglings over brood size

(i.e., number of successes over trials) via GLMMs with a binomial distribution and logit link

function (Douma and Weedon 2019). The duration of the nestling period as well as nestling body

mass and wing length were modeled using linear mixed effects models (LMMs). We included

year, farm and nest box identification as random factors for fledging success and duration of the

nestling period. Since multiple observations of nestling body mass and wing length for each

brood were observed and nesting brood identification within nest box identification led to

convergence failures, we nested brood identification within farm identification, along with year,

as random factors.

Model covariate summaries can be found in Table 1 and were averaged across a 12-day window

post-hatching for fledging success and a 16-day window post-hatching for the duration of the

nestling period, nestling body mass and wing length. Twelve days of age represents the period

when Tree Swallow nestlings become homeothermic, reach peak body mass, and during which

most nestling mortality occurs, thus representing the period where food availability is

presumably most crucial (McCarty 1995, Lamoureux 2010, Houle et al. 2020). Estimates of prey

availability during the time window of each breeding attempt was represented by predictions

from generalized additive models (GAMs) in which raw values of prey availability (i.e., Diptera

biomass) were regressed against the Julian date of sample collection for each farm and year

separately. We modeled Diptera availability to capture the general phenology throughout each

season and avoid biases caused by more punctual or local disturbances or phenomena such as the

capture of an insect swarm. GAMs were fitted as a tensor product smoother using the mgcv

package in R (Wood 2015) with an identical degree of smoothness (k=10) and using a Gamma

distribution with a log link function. From these predictions, we calculated an estimate of the

mean availability of prey during the respective temporal window of each fitness and body

condition proxy for each breeding attempt.

In order to evaluate our hypothesis regarding trophic links between agricultural landscapes and

fitness and body condition proxies, we compared a set of six models, each representing

biologically relevant alternative hypotheses predicting the outcome of each proxy and took an

information theoretic and multimodel approach (Appendix S1: Table S1). Besides a null model

only containing random effects, we started with a model including only confounding factors

(Base). Confounding factors remained identical across all fitness and body condition proxies and

were included to control for their known effects. Confounding factors included the mean

maximum daily temperature (Winkler et al. 2013) and mean daily precipitation (Cox et al. 2019,

2020) during the same post hatching and proxy-specific temporal window as prey availability.

We also included the morphological age of the breeding female (second year (SY) vs. after

second year (ASY); Hussell 1983), brood size, and the percent cover of open water around a nest

box. Finally, to account for seasonal variation in the species composition of Diptera (Bellavance

et al. 2018), and the trend of seasonal declines in reproductive success observed by Tree

Swallows (Winkler and Allen 1996), we also included a brood's hatching date. In the case of

nestling body mass and wing length, brood size was replaced with the summation of the daily

brood size during the 16-day period (Brood days), as this accounts for intra-brood competition in

the event of nestling mortality (Martin 1987). All model terms within Base were included in all

subsequent models of the candidate set. We included three models hypothesizing the effect of

prey availability and landscape context were influential in a strictly additive capacity, including a

model with only prey availability (Base + Food), a model with only site scores and their

interaction (Base + Land), and a model with both prey availability and site scores (Base + Food

+ Land). The final model is one hypothesizing that the effect of prey availability varied with

landscape context and thus included an interaction between prey availability and both sites score

values (Base + Food * Land).

In all analyses of both Diptera availability and fitness and body condition proxies, unless a top

performing model could be determined (AICc $w > 0.90$), the effects of key individual variables,

including interactions, were estimated via multi-model inference whereby model predictions

were calculated by model-averaging with shrinkage and shown with their 95% unconditional

confidence intervals (Burnham and Anderson 2002). All quantitative covariates were z-

transformed, r-squares were calculated following Nakagawa and Schielzeth (2013), and all

analyses were performed in R using the glmmTMB (Magnusson et al. 2020) and AICcmodavg

(Mazerolle 2020) packages. Model validation included evaluation of normally distributed

residuals (simulated), heteroskedasticity, and checks of variance inflation factors (VIF) following

Zuur et al. (2009) and using the DHARMa package (Hartig 2020).

Results

Seasonal influence of weather and an agricultural gradient on prey availability

We collected and processed 15,916 insect samples, resulting in information on 8,614 farm visits

over 11 years. Overall mean Diptera availability ($\pm$ SD) was 0.030 ± 0.044 g per trap and per

farm for each two-day sampling period. While it did not vary greatly between years (range of

means: 0.019 g - 0.037 g), the variance within each year was high (range of SD: 0.023 g -

0.059 g; Appendix S1: Figure S2). Diptera availability peaked near Julian day 183 (June 24;

Figure 3a), decreased following increasing precipitation, and rapidly increased with mean

maximum temperature, peaking at 21.7°C (Table 2; Appendix S1: Figure S3). While the

interaction between site score values indicated Diptera availability was highest within

predominantly forested landscapes (range percent forest cover: 0.0% to 69.1%), it was

marginally greater in predominantly forage landscapes than in ones dominated by corn and

soybean fields (Figure 3f). Compared to forage landscapes, predicted Diptera availability was

14.0% lower in corn and soybean landscapes as well as mixed-row crop landscapes, and nearly

200% greater in forested landscapes. Furthermore, the interaction between site scores and Julian

date highlighted that, at the start and end of the food provisioning period (Julian date 152 and

196, respectively), Diptera availability within corn and soybean landscapes was 59.2% and

62.9% lower than within forage landscapes (Appendix S1: Figure S4). While during the middle

of the food-provisioning period (Julian date 174), predicted Diptera availability was only 5.2%

lower within corn and soybean landscapes versus forage landscapes. Finally, we did not find

strong support that the influence of precipitation on Diptera biomass varied with landscape

context.

Fledging success

In total, we monitored the breeding activity of 1,897 breeding attempts producing at least one

nestling across the 40 farms and 11 breeding seasons. Overall mean fledging success ($\pm$ SD) was

0.74 $\pm$ 0.38 per brood and ranged between 0.63 and 0.88 among years, with the exception of the

2016 breeding season, where mean fledging success was 0.49 due to a period of prolonged cold

temperatures in June resulting in many brood failures (Appendix S1: Figure S2). As expected,

fledging success increased with prey availability (Table 2, Figure 3b). An increase of 17.0% in

fledging success was predicted between broods experiencing the minimum and maximum

observed levels of prey availability. A residual effect of land cover indicated that, under average

prey availability, fledging success was greatest within landscapes minimizing site score values,

and thus lowest within more agro-intensive landscapes (Figure 3g). Compared to forage

landscapes, fledging success was 10.3% lower in corn and soybean landscapes and 13.3% lower

within mixed-row crop landscapes. Interaction terms between site scores and prey availability

indicated under low prey conditions (half of the mean), fledging success within corn and soybean

landscapes or mixed-row crop landscapes was 8.6% and 9.7% lower than in forage landscapes,

respectively. Furthermore, under typically high levels of prey availability (twice the mean),

fledging success within these same two agro-intensive landscapes was 13.4% and 20% lower

than in forage landscapes, respectively (Appendix S1: Figure S5). Hatching date, younger aged

females (SY), brood size, and mean precipitation were all negatively related to fledging success

(Table 2). Both the mean maximum daily temperature and percent cover of open water were

positively related to fledging success (Table 2).

Nestling duration

We measured the duration of the nestling period for 1,556 broods from which at least one

nestling fledged. Mean nestling duration ($\pm$ SD) was 19.7 $\pm$ 1.8 days per brood and ranged

between 18.9 days and 21.1 days across years (Appendix S1: Figure S2). While we found no

effect of prey availability on nestling duration, the latter was 6.9% longer between maximum and

minimum Comp.1 scores and decreased slightly along Comp.2 (Figure 3h). Nestling duration

was thus longest within landscapes dominated by row crops and shortest within those occupied

by a mixture of forage fields and tree cover. We found that nestling duration increased with both

the younger age of the female (SY) and brood size but decreased with higher mean maximum

daily temperature. However, we found no strong relationship with hatching day, mean

precipitation, or percent cover of open water (Table 2).

Nestling body mass

We weighed 6,011 nestlings from 1,400 broods at 16 days of age that survived to fledging. Mean

nestling body mass ($\pm$ SD) was 21.2 $\pm$ 2.0 g, ranging between 20.4 g and 21.5 g across years

(Appendix S1: Figure S2). We found that nestling body mass increased with prey availability

(Figure 3d). An increase of 3.9% in nestling body mass was predicted between the minimum and

maximum observed levels of prey availability. After controlling for prey availability, we

observed no residual effect of landscape context on body mass (Figure 3i). We further found

nestling body mass decreased with high levels of precipitation, throughout the breeding season

and with increasing number of brood days. However, nestling body mass was not related to the

percent cover of open water or age of breeding females (Table 2).

Nestling wing length

We measured the wing length of 6,065 nestlings from 1,413 broods at 16 days of age that

survived to fledging. Mean nestling wing length ($\pm$ SD) was 45.8 $\pm$ 7.0 mm, ranging between

38.7 mm and 49.2 mm across years (Appendix S1: Figure S2). We found wing length was

unrelated to prey availability (Figure 3e). However, we observed a residual effect of landscape

context on wing length whereby wing length was up to 10.1% longer within forage landscapes

compared to corn and soybean landscapes (Figure 3j). We further found wing length increased

with increasing mean maximum daily temperatures and was shorter for nestlings of young (SY)

mothers and with increasing brood days. However, no relationship was observed between wing

length and either precipitation, percent cover of open water, or hatching date (Table 2).

Discussion

Peak prey availability for Tree Swallows occurred in landscapes dominated by forest cover and a

mixture of agricultural habitats, suggesting greater landscape heterogeneity results in greater

prey availability for this species throughout southern Québec's gradient of agricultural

intensification. We also observed that prey availability is an important determinant of Tree

Swallow reproductive success in this geographical context. Indeed, greater prey availability was

associated with higher fledging success and greater nestling body mass but unrelated to the time

taken by nestlings to fledge or their wing length upon fledging. Interestingly, we found landscape

context was associated with fledging success, nestling duration, and nestling wing length only

after controlling for the effect of prey availability. This residual effect of landcover suggests

other factors related to the agricultural landscape context affect breeding Tree Swallows.

Seasonal influence of weather and agricultural landcover on prey availability

The availability of prey was marginally greater in forage dominated landscapes than within corn

or soybean and mixed-row crop landscapes. These results are in agreement with those found in

the literature and our own predictions suggesting aerial insect abundance would be greater within

less agro-intensive habitats (Wickramasinghe et al. 2004, Evans et al. 2007, Grüebler et al.

2008). Our finding that prey availability was greatest within agricultural areas harboring elevated

forest cover is likely explained by habitat heterogeneity resulting in a more abundant and diverse

community of Diptera (Benton et al. 2003, Fahrig et al. 2011). Perhaps the most interesting

finding was that the phenology of prey availability as Diptera varied across this agricultural

gradient. Diptera biomass in the middle of the breeding season was similar between forage

dominated landscapes and landscapes dominated by more agro-intensive cultures. However, at

the beginning and ends of the breeding season prey availability was greater in less agro-intense

areas. This pattern was observed in other studies from this system yet on much shorter sequences

of years (Rioux Paquette et al. 2013, Bellavance et al. 2018). A possible cause of this spatio-

temporal variation in Diptera biomass is that the phenology and species composition of Diptera

varied across the agricultural gradient. Such a variation in species composition was indeed

observed in previous work from this system (Bellavance et al. 2018 and Appendix S2: Figure S1-

5). Tree Swallows, similar to many other passerine species, display seasonal declines in

reproductive success thought to be related to changes in the availability or adequacy of resources

(Winkler and Allen 1996, Nooker et al. 2005, Bourret et al. 2015). Therefore, the landscape

driven differences in the phenology of food resources reported here suggests that the severity of

seasonal declines in reproductive success of farmland birds could be landscape dependent.

Effects of prey availability on fitness and body condition proxies

Fledging success and nestling body mass were positively related to prey availability. These

findings support previous studies documenting similar relationships stemming from artificial

changes in local food availability (Boutin 1990, Granbom and Smith 2006, Robb et al. 2008,

Ruffino et al. 2014). For instance, multiple lines of evidence suggest declines in fitness proxies

and changes in foraging behaviors of avian species following the decrease of prey availability

through the application of various agro-chemical agents (Martin et al. 1998, 2000, Brickle et al.

2000, Hart et al. 2006, Poulin et al. 2010). Our results also agree with previous evidence from

other swallow populations observing unaltered fluctuations in prey availability being positively

related to fledging success and nestling condition (Quinney et al. 1986, McCarty and Winkler

1999b, Winkler et al. 2013, Teglhøj 2017, Twining et al. 2018). Yet, our results run counter to

recent studies from other swallow populations where measurable variation in aerial insect prey

was suggested to have no impact on seasonal reproductive success and morphological condition

of offspring (Dunn et al. 2011, Imlay et al. 2017, McClenaghan et al. 2019). A frequent claim of

these studies is that prey availability was overly abundant during periods of observation, and thus

other resources limited fitness, highlighting the importance of conducting food availability

studies across a broad enough range of habitat qualities to reduce the availability of resources

below such a critical threshold (Fazey et al. 2005). We found evidence that, within agricultural

systems, the growth of Tree Swallow populations could be food limited. However, to address the

role of prey availability on the population dynamics of this species more fully, the availability of

food resources will also need to be quantified with reference to both clutch size and egg

condition, as these factors partly determine brood size and fledgling number (Meijer et al. 1989,

Ardia et al. 2006, Gow et al. 2019). Indeed, food resources may limit clutch size through a

reduction in resource allocation (Wilson et al. 2017) and impact incubation capacity and hatching

by regulating the duration and time between foraging bouts (McCarty 1995, Monaghan and

Nager 1997, Thomson et al. 2007).

We found no indication that wing length was related to the availability of prey, perhaps

suggesting that prey availability during this study was adequate for allocating resources towards

the growth of flight feathers. We also found that the duration of the nestling period was unrelated

to prey availability, a result running counter to that of other swallow studies (Teglhøj 2017). One

plausible explanation is that the supply of food resources was sufficient to meet the basal

metabolic demand for the rate of nestling growth given the observed brood sizes (Leonard 2000,

Stodola et al. 2010). The duration of the nestling period is likely a function of the nestlings'

growth rate given that nestlings should remain within nests until reaching a sufficient condition

(Nilsson and Svensson 1993, Seki and Takano 1998, Rossmanith et al. 2007). Recognizing this,

the lack of a relationship between the duration of the nestling period and prey availability leads

us to suggest that, in our system, the duration of the nestling period may be driven by something

other than the accumulation of food resources.

Residual effects of agricultural landcover on fitness and body condition proxies

An important finding of our study is that relationships between fitness and body condition

proxies and agricultural landcover was observed even after controlling for prey availability.

Fledging success was lower, nestling period durations were longer, and nestling wing lengths

were indeed shorter within areas dominated by large-scale row-crop monocultures, principally

corn and soybean. Among the several hypotheses explaining such results, we discuss three non-

mutually exclusive alternatives whereby differences in landscape context could lead to variation

in the contamination of food by pesticides, in the nutritive value of food, and in the rate at which

food is provisioned.

During the years covered by this study, areas intensively-managed for growing corn and soybean

harbored elevated levels of pesticides in their surface waters, notably several herbicides (e.g.,

glyphosate, atrazine, S-metolachlor, imazethapyr) and neonicotinoid insecticides [e.g.,.

clothianidin, thiacloprid, thiametoxam, (Giroux 2019, Montiel-León et al. 2019)]. Many of those

pesticides were also found, even simultaneously, in the insects fed to nestlings by Tree Swallows

in our system (Poisson et al. 2021) and may hence have caused lethal and sublethal toxicological

effects on nestlings. For instance, many pesticides, such as the neonicotinoids, can alter the

neurological, endocrine and immunological systems of non-target animals, including vertebrates,

via skin contact, breathing or food/water consumption (Grue et al. 1997, Mayne et al. 2005,

Mineau and Palmer 2013, Gibbons et al. 2014, Lopez-Antia et al. 2015). Such disruptions can

induce, among many observed effects, anorexia and reduced mass gain, and even result in lower

nestling survival, delayed fledging, and lower fledging mass, irrespective of the amount of food

provided (Grue et al. 1997, Eng et al. 2017, 2019, Addy-Orduna et al. 2019). Pesticide exposure

of Tree Swallow broods within areas dominated by corn and soybean may thus have been severe

enough to result in nestling mortality or retarded nestling growth. Moreover, this may explain the

lack of relationship between prey availability and the duration of the nestling period we

observed, because exposure to some pesticides can also lessen parental care in birds (Grue et al.

1997, Custer 2011, Mineau and Palmer 2013, Gibbons et al. 2014).

Variation in landscape contexts may also have resulted in Tree Swallows feeding on diets

differing not only in prey species composition, but also in nutritive value. Indeed, the diet of

Tree Swallows varies across the agricultural intensification gradient covered by our study area

(Bellavance et al. 2018; see also Nocera et al. 2012, Michelson et al. 2018). In the period during

which Tree Swallows provide food to nestlings, they consume a diet with a greater biomass of

insects with aquatic larval stages (e.g., Ephemeroptera and Diptera: Nematocera) in landscapes

devoted mostly to forage crops and pastures (Bellavance et al. 2018 and Appendix S2: Figure

S1-5). This is despite these insects representing a minute portion of what is available across the

agricultural gradient studied here (Bellavance et al. 2018 and Appendix S2: Figure S1-5).

Increased provisioning of these types of insects may lead to a more nutritive diet in less

intensively cultivated landscapes as they typically contain high levels of highly unsaturated

omega-3 fatty acids (HUFA), which have been found to improve the mass gain,

immunocompetence, and fledging success of Tree Swallow nestlings (Twining et al. 2018). That

said, insects with an aquatic larval stage can also be loaded with various environmental

contaminants, such as pesticides and heavy metals, that may impair the breeding behavior and

success of swallows as well as their survival (Custer 2011, Alberts et al. 2013, Twining et al.

2021). The costs and benefits of feeding on aquatic insects rich in HUFA within an agricultural

context will thus require further attention before we can determine its real impact on Tree

Swallow fitness components and demography.

Landscapes dominated by row crop monocultures express landscape simplification in which

large swaths of areas are occupied by only a handful of habitats (Benton et al. 2003). This

phenomenon may make finding and exploiting suitable food resources more difficult for animals

relying upon "residual" marginal habitats as landscape simplification lowers the functional

connectivity of agricultural landscapes (Hinsley 2000, Bélisle 2005, Rainho and Palmeirim

2011). If the amount and quality of suitable foraging habitats are reduced and disconnected, we

may expect a negative impact of such landscape simplification on nestling growth if parents

cannot compensate by increasing foraging effort. There is currently insufficient evidence to

evaluate if differing compositions of agricultural landscapes consistently impact foraging adults

negatively, yet the evidence available does suggest this is the case (Poulin et al. 2010,

Staggenborg et al. 2017), albeit weakly in Tree Swallows (Stanton et al. 2016). In our study

system, we observed that the biomass of food boluses delivered to nestlings does not vary with

landscape context, yet their delivery rate is lowest within landscapes dominated by row crops

(Garrett et al. 2021). Assuming parents are maximizing provisioning rate of food items to their

offspring, given the availability of prey, this may provide evidence that adults are having to

spend more time foraging within these types of landscapes (Charnov 1976, Pyke et al. 1977). If

sustained, this increased foraging effort could come at a cost to parents and lead to carry-over

effects eventually impacting their future fitness (Saino et al. 1999, 2018, Harrison et al. 2011,

Stanton et al. 2017).

Conclusion

Taken together, our results provide evidence that food availability during the nestling period has

a positive effect on fitness components of Tree Swallows breeding in farmlands. Yet, more

importantly, after controlling for food resources, a residual effect of agricultural landcover

persisted. The interactive effect of prey availability and agricultural intensification on fitness and

body condition proxies likely means that, for breeding attempts within less agro-intensive areas,

increased prey availability has a positive influence. Yet, similar increases in prey availability

within more intensively cultivated landscapes potentially result in an increased toxicological load

of food to nestlings as well as a reduced dietary quality. Future work should expand on the

relationship between nestling condition, fledging success, and post-fledgling survival in

landscapes harboring elevated levels of agricultural habitats. More specifically, it is unknown

how the availability of food resources early in the breeding season is influenced by agricultural

land cover and if this relationship impacts settlement patterns and clutch size (Rendell and

Robertson 1990, Orians and Wittenberger 1991, Grüebler et al. 2010, Robillard et al. 2013).

However, it is certain that these are important features of the seasonal reproductive success and

the population dynamics of aerial insectivores. Furthermore, nestlings reared under reduced prey

availability conditions may be unfit for post-fledgling survival or pre-migration fueling (Saino et

al. 2012, Jones et al. 2017). For example, the body condition of nestling Barn Swallows

(*Hirundo rustica*) is positively correlated to post fledging survival and fledglings in better

condition initiated migration sooner (Evans et al. 2020). The poor body condition expressed by

fledglings within more agro-intensive areas may thus result in elevated post-fledging mortality, a

field readily understudied (Cox et al. 2014) yet a key part in the annual cycle of this species and

another missing link in the understanding of aerial insectivore declines.

Acknowledgments

We are indebted to the farm owners who kindly accepted to partake in our long-term study since 2004. Sincere thanks to the many graduate students, field and lab assistants who helped collect, enter, and proof data throughout the years, notably with respect to insect sample processing. This work was conducted under the approval of the animal care committee of the Université de Sherbrooke and was financially supported by Natural Sciences and Engineering Research Council of Canada (NSERC) discovery grants to FP, DG and MB and a Steacie fellowship to FP, two team research grants from the Fonds de recherche du Québec—Nature et technologies (FRQNT) to FP, DG and MB, by the Canada Research Chairs program to FP and MB, as well as by New Opportunities Funds of the Canadian Foundation for Innovation (FCI) to FP, DG and MB, the Canadian Wildlife Service of Environment and Climate Change Canada, and the Université de Sherbrooke.

Literature cited.

Addy-Orduna, L. M., J. C. Brodeur, and R. Mateo. 2019. Oral acute toxicity of imidacloprid,

thiamethoxam and clothianidin in eared doves: A contribution for the risk assessment of

neonicotinoids in birds. Science of The Total Environment 650:1216–1223.

Agriculture and Agri-Food Canada (AAFC). 2018. Annual Crop Inventory.

https://open.canada.ca/data/en/dataset/.

Alberts, J. M., S. M. P. Sullivan, and A. Kautza. 2013. Riparian swallows as integrators of

landscape change in a multiuse river system: Implications for aquatic-to-terrestrial transfers

of contaminants. Science of The Total Environment 463–464:42–50.

Almasi, B., P. Béziers, A. Roulin, and L. Jenni. 2015. Agricultural land use and human presence

around breeding sites increase stress-hormone levels and decrease body mass in barn owl

nestlings. Oecologia 179:89–101.

Andow, D. 1983. The extent of monoculture and its effects on insect pest populations with

particular reference to wheat and cotton. Agriculture, Ecosystems & Environment 9:25–35.

Andow, D. 1991. Vegetational Diversity And Arthropod Population Response. Annual Review

of Entomology 36:561–586.

Ardia, D. R., J. H. Pérez, and E. D. Clotfelter. 2010. Experimental cooling during incubation

leads to reduced innate immunity and body condition in nestling tree swallows. Proceedings

of the Royal Society B: Biological Sciences 277:1881–1888.

Ardia, D. R., M. F. Wasson, and D. W. Winkler. 2006. Individual quality and food availability

determine yolk and egg mass and egg composition in tree swallows Tachycineta bicolor.

Journal of Avian Biology 37:252–259.

Attwood, S. J., M. Maron, A. P. N. House, and C. Zammit. 2008. Do arthropod assemblages

display globally consistent responses to intensified agricultural land use and management?

Global Ecology and Biogeography 17:585–599.

Bélanger, L., and M. Grenier. 2002. Agriculture intensification and forest fragmentation in the

St. Lawrence valley, Québec, Canada. Landscape Ecology 17:495–507.

Bélisle, M. 2005. Measuring landscape connectivity: the challenge of behavioral landscape

ecology. Ecology 86:1988–1995.

Bellavance, V., M. Bélisle, J. Savage, F. Pelletier, and D. Garant. 2018. Influence of agricultural

intensification on prey availability and nestling diet in Tree Swallows (Tachycineta bicolor).

Canadian Journal of Zoology 96:1053–1065.

Benton, T. G., D. M. Bryant, L. Cole, and H. Q. P. Crick. 2002. Linking agricultural practice to

insect and bird populations: a historical study over three decades. Journal of Applied

Ecology 39:673–687.

Benton, T. G., J. A. Vickery, and J. D. Wilson. 2003. Farmland biodiversity: is habitat

heterogeneity the key? Trends in Ecology & Evolution 18:182–188.

Berzins, L. L., A. K. Mazer, C. A. Morrissey, and R. G. Clark. 2021. Pre-fledging quality and

recruitment in an aerial insectivore reflect dynamics of insects, wetlands and climate.

Oecologia.

Boatman, N. D., N. W. Brickle, J. D. Hart, T. P. Milsom, A. J. Morris, A. W. A. Murray, K. A.

Murray, and P. A. Robertson. 2004. Evidence for the indirect effects of pesticides on

farmland birds. Ibis 146:131–143.

Bourret, A., M. Bélisle, F. Pelletier, and D. Garant. 2015. Multidimensional environmental

influences on timing of breeding in a tree swallow population facing climate change.

Evolutionary Applications 8:933–944.

Boutin, S. 1990. Food supplementation experiments with terrestrial vertebrates: patterns,

problems, and the future. Canadian Journal of Zoology 68:203–220.

Brickle, N. W., D. G. C. Harper, N. J. Aebischer, and S. H. Cockayne. 2000. Effects of

agricultural intensification on the breeding success of corn buntings Miliaria calandra.

Journal of Applied Ecology 37:742–755.

Burnham, K. P., and D. R. Anderson. 2002. Model Selection and Multimodel Inference. second.

Springer.

Ceballos, G. 2002. Mammal Population Losses and the Extinction Crisis. Science 296:904–907.

Charnov, E. L. 1976. Optimal foraging, the marginal value theorem. Theoretical Population

Biology 9:129–136.

Clark, R. G., D. W. Winkler, R. D. Dawson, D. Shutler, D. J. T. Hussell, M. P. Lombardo, P. A.

Thorpe, P. O. Dunn, and L. A. Whittingham. 2018. Geographic variation and environmental

correlates of apparent survival rates in adult tree swallows Tachycineta bicolor. Journal of

Avian Biology 49:jav-012514.

Courtois, E., D. Garant, F. Pelletier, and M. Bélisle. 2021. Nonideal nest box selection by tree

swallows breeding in farmlands: evidence for an ecological trap? Authorea.

Cox, A. R., R. J. Robertson, B. C. Fedy, W. B. Rendell, and F. Bonier. 2018. Demographic

drivers of local population decline in Tree Swallows (Tachycineta bicolor) in Ontario,

Canada. The Condor 120:842–851.

Cox, A. R., R. J. Robertson, Á. Z. Lendvai, K. Everitt, and F. Bonier. 2019. Rainy springs linked

to poor nestling growth in a declining avian aerial insectivore (Tachycineta bicolor).

Proceedings of the Royal Society B: Biological Sciences 286:20190018.

Cox, A. R., R. J. Robertson, W. B. Rendell, and F. Bonier. 2020. Population decline in tree

swallows (Tachycineta bicolor) linked to climate change and inclement weather on the

breeding ground. Oecologia 192:713–722.

Cox, W. A., F. R. I. Thompson, A. S. Cox, and J. Faaborg. 2014. Post-fledging survival in

passerine birds and the value of post-fledging studies to conservation. The Journal of

Wildlife Management 78:183–193.

Custer, C. 2011. Swallows as a Sentinel Species for Contaminant Exposure and Effect Studies.

Pages 267–300 *in* J. E. Elliott, C. A. Bishop, and C. A. Morrissey, editors. Wildlife

Ecotoxicology: Forensic Approaches. Springer New York.

Davis, R. 2007. Habitat Fragmentation and Landscape Change: An Ecological and Conservation

Synthesis. Austral Ecology 32:477–478.

Dawson, R. D., and G. R. Bortolotti. 2000. Reproductive Success of American Kestrels: The

Role of Prey Abundance and Weather. The Condor 102:814–822.

Douma, J. C., and J. T. Weedon. 2019. Analysing continuous proportions in ecology and

evolution: A practical introduction to beta and Dirichlet regression. Methods in Ecology and

Evolution 10:1412–1430.

Dunn, P. O., D. W. Winkler, L. a. Whittingham, S. J. Hannon, and R. J. Robertson. 2011. A test

of the mismatch hypothesis: How is timing of reproduction related to food abundance in an

aerial insectivore? Ecology 92:450–461.

Elgin, A. S., R. G. Clark, and C. A. Morrissey. 2020. Tree Swallow selection for wetlands in

agricultural landscapes predicted by central-place foraging theory. The Condor 122:1–12.

Eng, M. L., B. J. M. Stutchbury, and C. A. Morrissey. 2017. Imidacloprid and chlorpyrifos

insecticides impair migratory ability in a seed-eating songbird. Scientific Reports 7:1–9.

Eng, M. L., B. J. M. Stutchbury, and C. A. Morrissey. 2019. A neonicotinoid insecticide reduces

fueling and delays migration in songbirds. Science 365:1177–1180.

Evans, D. R., K. A. Hobson, J. W. Kusack, M. D. Cadman, C. M. Falconer, and G. W. Mitchell.

2020. Individual condition, but not fledging phenology, carries over to affect post☐fledging

survival in a Neotropical migratory songbird. Ibis 162:331–344.

Evans, K. L., J. D. Wilson, and R. B. Bradbury. 2007. Effects of crop type and aerial invertebrate

abundance on foraging barn swallows Hirundo rustica. Agriculture, Ecosystems &

Environment 122:267–273.

Fahrig, L., J. Baudry, L. Brotons, F. G. Burel, T. O. Crist, R. J. Fuller, C. Sirami, G. M.

Siriwardena, and J.-L. Martin. 2011. Functional landscape heterogeneity and animal

biodiversity in agricultural landscapes. Ecology Letters 14:101–112.

Fazey, I., J. Fischer, and D. B. Lindenmayer. 2005. What do conservation biologists publish?

Biological Conservation 124:63–73.

Filzmoser, P., K. Hron, and C. Reimann. 2009. Principal component analysis for compositional

data with outliers. Environmetrics 20:621–632.

Fischer, J., and D. B. Lindenmayer. 2007. Landscape modification and habitat fragmentation: a

synthesis. Global Ecology and Biogeography 16:265–280.

Garrett, D. R., F. Pelletier, D. Garant, and M. Bélisle. 2021. Negative effects of agricultural

intensification on energetic gain rates of a declining aerial insectivore.

bioRxiv:10.1101/2021.03.24.436831.

Gibbons, D., C. A. Morrissey, and P. Mineau. 2014. A review of the direct and indirect effects of

neonicotinoids and fipronil on vertebrate wildlife. Environmental Science and Pollution

Research 22:103–118.

Giroux, I. 2019. Présence de pesticides dans l'eau au Québec : Portrait et tendances dans les

zones de maïs et de soya – 2015 à 2017, Québec, ministère de l'Environnement et de la

Lutte contre les changements climatiques, Direction générale du suivi de l'état de l'environ.

Gow, E. A., S. M. Knight, D. W. Bradley, R. G. Clark, D. W. Winkler, M. Bélisle, L. L. Berzins,

T. Blake, E. S. Bridge, L. Burke, R. D. Dawson, P. O. Dunn, D. Garant, and G. Holroyd.

2019. Effects of Spring Migration Distance on Tree Swallow Reproductive Success Within

and Among Flyways 7:1–10.

Granbom, M., and H. G. Smith. 2006. Food Limitation During Breeding in a Heterogeneous

Landscape. The Auk 123:97–107.

Grue, C. E., P. L. Gibert, and M. E. Seeley. 1997. Neurophysiological and Behavioral Changes

in Non-Target Wildlife Exposed to Organophosphate and Carbamate Pesticides:

Thermoregulation, Food Consumption, and Reproduction. American Zoologist 37:369–388.

Grüebler, M. U., F. Korner-Nievergelt, and J. Von Hirschheydt. 2010. The reproductive benefits

of livestock farming in barn swallows Hirundo rustica: quality of nest site or foraging

habitat? Journal of Applied Ecology 47:1340–1347.

Grüebler, M. U., M. Morand, and B. Naef-Daenzer. 2008. A predictive model of the density of

airborne insects in agricultural environments. Agriculture, Ecosystems & Environment

123:75–80.

Hailman, J. P., and D. Lack. 1955. The Natural Regulation of Animal Numbers. AIBS Bulletin

5:12–12.

Hale, R., R. Coleman, V. Pettigrove, and S. E. Swearer. 2015. Identifying, preventing and

mitigating ecological traps to improve the management of urban aquatic ecosystems.

Journal of Applied Ecology 52:928–939.

Harrison, X. A., J. D. Blount, R. Inger, D. R. Norris, and S. Bearhop. 2011. Carry-over effects as

drivers of fitness differences in animals. Journal of Animal Ecology 80:4–18.

Hart, J. D., T. P. Milsom, G. Fisher, V. Wilkins, S. J. Moreby, A. W. A. Murray, and P. A.

Robertson. 2006. The relationship between yellowhammer breeding performance, arthropod

abundance and insecticide applications on arable farmland. Journal of Applied Ecology

43:81–91.

Hartig, F. 2020. DHARMa: Residual Diagnostics for Hierarchical Regression Models.

Hinsley, S. A. 2000. The costs of multiple patch use by birds. Landscape Ecology 15:765–775.

Hostetler, J. A., T. S. Sillett, and P. P. Marra. 2015. Full-annual-cycle population models for

migratory birds. The Auk 132:433–449.

Houle, C., F. Pelletier, M. Bélisle, and D. Garant. 2020. Impacts of environmental heterogeneity

on natural selection in a wild bird population. Evolution 74:1142–1154.

Hussell, D. J. T. 1983. Age and Plumage Color in Female Tree Swallows. Journal of Field

Ornithology 54:312–318.

Imlay, T. L., H. A. R. Mann, and M. L. Leonard. 2017. No effect of insect abundance on nestling

survival or mass for three aerial insectivores. Avian Conservation and Ecology

12:10.5751/ACE-01092–120219.

Jobin, B., J. Beaulieu, M. Grenier, L. Bélanger, C. Maisonneuve, D. Bordage, and B. Filion.

2003. Landscape changes and ecological studies in agricultural regions, Québec, Canada.

Landscape Ecology 18:575–590.

Jobin, B., M. Grenier, and P. Laporte. 2005. Using satellite imagery to assess breeding habitat

availability of the endangered loggerhead shrike in Québec. Biodiversity and Conservation

14:81–95.

Johnson, M. E., and M. P. Lombardo. 2000. Nestling Tree Swallow (Tachycineta bicolor) Diets

in an Upland Old Field in Western Michigan. The American Midland Naturalist 144:216–

219.

Jones, T. M., and M. P. Ward. 2020. Pre☐ to post☐fledging carryover effects and the adaptive

significance of variation in wing development for juvenile songbirds. Journal of Animal

Ecology 89:2235–2245.

Jones, T. M., M. P. Ward, T. J. Benson, and J. D. Brawn. 2017. Variation in nestling body

condition and wing development predict cause-specific mortality in fledgling dickcissels.

Journal of Avian Biology 48:439–447.

Knight, S. M., E. A. Gow, D. W. Bradley, R. G. Clark, M. Bélisle, L. L. Berzins, T. Blake, E. S.

Bridge, L. Burke, R. D. Dawson, P. O. Dunn, D. Garant, G. L. Holroyd, D. J. T. Hussell, O.

Lansdorp, A. J. Laughlin, M. L. Leonard, F. Pelletier, D. Shutler, L. Siefferman, C. M.

Taylor, H. E. Trefry, C. M. Vleck, D. Vleck, L. A. Whittingham, D. W. Winkler, and D. R.

Norris. 2019. Nonbreeding season movements of a migratory songbird are related to

declines in resource availability. The Auk 136:1–13.

Kusack, J. W., G. W. Mitchell, D. R. Evans, M. D. Cadman, and K. A. Hobson. 2020. Effects of

agricultural intensification on nestling condition and number of young fledged of barn

swallows (Hirundo rustica). Science of The Total Environment 709:136195.

Lambin, E. F., B. L. Turner, H. J. Geist, S. B. Agbola, A. Angelsen, J. W. Bruce, O. T. Coomes,

R. Dirzo, G. Fischer, C. Folke, P. S. George, K. Homewood, J. Imbernon, R. Leemans, X.

Li, E. F. Moran, M. Mortimore, P. S. Ramakrishnan, J. F. Richards, H. Skånes, W. Steffen,

G. D. Stone, U. Svedin, T. A. Veldkamp, C. Vogel, and J. Xu. 2001. The causes of land-use

and land-cover change: moving beyond the myths. Global Environmental Change 11:261–

269.

Lamoureux, S. 2010. Impact de l'intensification agricole sur l'effort parental, la croissance et la

survie des oisillons chez l'hirondelle bicolore (Tachycineta bicolor). Université de

Sherbrooke.

Laplante, L. 2013. Influence de l'hétérogénéité et de la composition du paysage agricole sur la

communauté de diptères schizophores du sud du québec (canada). Université de

Sherbrooke.

Leonard, M. L. 2000. The Role of Development, Parental Behavior, and Nestmate Competition

in Fledging of Nestling Tree Swallows. The Auk 117:996–1002.

Letten, A. D., P.-J. Ke, and T. Fukami. 2017. Linking modern coexistence theory and

contemporary niche theory. Ecological Monographs 87:161–177.

Lopez-Antia, A., M. E. Ortiz-Santaliestra, F. Mougeot, and R. Mateo. 2015. Imidacloprid-treated

seed ingestion has lethal effect on adult partridges and reduces both breeding investment

and offspring immunity. Environmental Research 136:97–107.

Magnusson, A., H. J. Skaug, A. Nielsen, C. W. Berg, K. Kristensen, M. Maechler, K. J. van

Benthem, B. M. Bolker, N. Sadat, D. Lüdecke, R. Lenth, J. O'Brien, and M. E. Brooks.

2020. Package ' glmmTMB .'

Malaj, E., K. Liber, and C. A. Morrissey. 2020. Spatial distribution of agricultural pesticide use

and predicted wetland exposure in the Canadian Prairie Pothole Region. Science of The

Total Environment 718:134765.

Marshall, M. R., R. J. Cooper, J. A. DeCecco, J. Strazanac, and L. Butler. 2002. Effects of

Experimentally Reduced Prey Abundance on the Breeding Ecology of the Red-Eyed Vireo.

Ecological Applications 12:261.

Martin, L. J., B. Blossey, and E. Ellis. 2012. Mapping where ecologists work: biases in the

global distribution of terrestrial ecological observations. Frontiers in Ecology and the

Environment 10:195–201.

Martin, P. A., D. L. Johnson, D. J. Forsyth, and B. D. Hill. 1998. Indirect effects of the

pyrethroid insecticide deltamethrin on reproductive success of chestnut-collared longspurs.

Ecotoxicology 7:89–97.

Martin, P. A., D. L. Johnson, D. J. Forsyth, and B. D. Hill. 2000. Effects of two grasshopper

control insecticides on food resources and reproductive success of two species of grassland

songbirds. Environmental Toxicology and Chemistry 19:2987–2996.

Martin, T. E. 1987. Food as a Limit on Breeding Birds: A Life-History Perspective. Annual

Review of Ecology and Systematics 18:453–487.

Mayne, G. J., C. A. Bishop, P. A. Martin, H. J. Boermans, and B. Hunter. 2005. Thyroid

Function in Nestling Tree Swallows and Eastern Bluebirds Exposed to Non-Persistent

Pesticides and p, p-DDE in Apple Orchards of Southern Ontario, Canada. Ecotoxicology

14:381–396.

Mazerolle, M. J. 2020. AICcmodavg: model selection and multimodel inference based on

(Q)AIC(c).

McCamey, F., and L. B. Slobodkin. 1963. Growth and Regulation of Animal Populations. Bird-

Banding 34:52.

McCarty, J. P. 1995. Effects of short-term changes in environmental conditions on the foraging

ecology and reproductive success of tree swallows, Tachycineta bicolor. Cornell Univeristy.

McCarty, J. P. 2001. Variation in Growth of Nestling Tree Swallows Across Multiple Temporal

and Spatial Scales. The Auk 118:176–190.

McCarty, J. P., and D. W. Winkler. 1999a. Foraging Ecology and Diet Selectivity of Tree

Swallows Feeding Nestlings. The Condor 101:246–254.

McCarty, J. P., and D. W. Winkler. 1999b. Relative importance off environmental variables in

determining the growth off nestling Tree Swallows Tachycineta bicolor. Ibis 141:286–296.

McClenaghan, B., K. C. R. Kerr, and E. Nol. 2019. Does prey availability affect the reproductive

performance of Barn Swallows (Hirundo rustica) breeding in Ontario, Canada? Canadian

Journal of Zoology 97:979–987.

MDDELCC. 2015. Ministère du Développement durable, de l'Environnement et de la Lutte

contre les changements climatiques. Stratégie québécoise sur les pesticides 2015-2018.

http://www.mddelcc.gouv.qc.ca/.

Meijer, T., D. Masman, and S. Daan. 1989. Energetics of Reproduction in Female Kestrels. The

Auk 106:549–559.

Mengelkoch, J. M., G. J. Niemi, and R. R. Regal. 2004. Diet of the nestling tree swallow. The

Condor 106:423–429.

Michel, N. L., A. C. Smith, R. G. Clark, C. A. Morrissey, and K. A. Hobson. 2016. Differences

in spatial synchrony and interspecific concordance inform guild-level population trends for

aerial insectivorous birds. Ecography 39:774–786.

Michelson, C. I., R. G. Clark, and C. A. Morrissey. 2018. Agricultural land cover does not affect

the diet of Tree Swallows in wetland-dominated habitats. The Condor 120:751–764.

Miller, E. J., and L. W. Adams. 1995. Urban Wildlife Habitats: A Landscape Perspective. The

Journal of Wildlife Management 59:194.

Mineau, P., and C. Palmer. 2013. The impact of the nation's most widely used insecticides on

birds. Cambridge University Press.

Monaghan, P., and R. G. Nager. 1997. Why don't birds lay more eggs? Trends in Ecology &

Evolution 12:270–274.

Montgomery, G. A., R. R. Dunn, R. Fox, E. Jongejans, S. R. Leather, M. E. Saunders, C. R.

Shortall, M. W. Tingley, and D. L. Wagner. 2020. Is the insect apocalypse upon us? How to

find out. Biological Conservation 241:108327.

Montiel-León, J. M., G. Munoz, S. Vo Duy, D. T. Do, M. Vaudreuil, K. Goeury, F. Guillemette,

M. Amyot, and S. Sauvé. 2019. Widespread occurrence and spatial distribution of

glyphosate, atrazine, and neonicotinoids pesticides in the St. Lawrence and tributary rivers.

Environmental Pollution 250:29–39.

Morrissey, C. A., P. Mineau, J. H. Devries, F. Sanchez-Bayo, M. Liess, M. C. Cavallaro, and K.

Liber. 2015. Neonicotinoid contamination of global surface waters and associated risk to

aquatic invertebrates: A review. Environment International 74:291–303.

NABCI. 2019. NABCI [Canada, North American Bird Conservation Initiative]. Page North

American Bird Conservation Initiative Canada. 2019. The State of Canada's Birds, 2019.

Environment and Climate Change Canada, Ottawa, Canada.

Naef-Daenzer, B., and M. U. Grüebler. 2016. Post☐fledging survival of altricial birds: ecological

determinants and adaptation. Journal of Field Ornithology 87:227–250.

Nagy, L. R., and K. G. Smith. 1997. Effects of Insecticide-Induced Reduction in Lepidopteran

Larvae on Reproductive Success of Hooded Warblers. The Auk 114:619–627.

Nakagawa, S., and H. Schielzeth. 2013. A general and simple method for obtaining R 2 from

generalized linear mixed-effects models. Methods in Ecology and Evolution 4:133–142.

Nebel, S., A. Mills, J. D. McCracken, and P. D. Taylor. 2010. Declines of Aerial Insectivores in

North America Follow a Geographic Gradient. Avian Conservation and Ecology

5:10.5751/ACE-00391–050201.

Newton, I. 1998. Population Limitations in Birds. Academic Press.

Newton, I. 2002. The Role of Food in Limiting Bird Numbers. Ardea 38–90:11–30.

Nilsson, J.-Å., and M. Svensson. 1993. Fledging in altricial birds: parental manipulation or
sibling competition? Animal Behaviour 46:379–386.

Nocera, J. J., J. M. Blais, D. V. Beresford, L. K. Finity, C. Grooms, L. E. Kimpe, K. Kyser, N.
Michelutti, M. W. Reudink, and J. P. Smol. 2012. Historical pesticide applications
coincided with an altered diet of aerially foraging insectivorous chimney swifts.
Proceedings of the Royal Society B: Biological Sciences 279:3114–3120.

Nooker, J. K., P. O. Dunn, and L. A. Whittingham. 2005. Effects of food abundance, weather,
and female condition on reproduction in tree swallows (Tachycineta bicolor). Auk
122:1225–1238.

Orians, G. H., and J. F. Wittenberger. 1991. Spatial and Temporal Scales in Habitat Selection.
The American Naturalist 137:S29–S49.

Pigeon, G., R. Baeta, M. Bélisle, D. Garant, and F. Pelletier. 2013. Effects of agricultural
intensification and temperature on immune response to phytohemagglutinin in Tree
Swallows (Tachycineta bicolor). Canadian Journal of Zoology 91:56–63.

Pisa, L. W., V. Amaral-Rogers, L. P. Belzunces, J. M. Bonmatin, C. A. Downs, D. Goulson, D.
P. Kreutzweiser, C. Krupke, M. Liess, M. McField, C. A. Morrissey, D. A. Noome, J.
Settele, N. Simon-Delso, J. D. Stark, J. P. Van der Sluijs, H. Van Dyck, and M. Wiemers.
2015. Effects of neonicotinoids and fipronil on non-target invertebrates. Environmental
Science and Pollution Research 22:68–102.

Poisson, M., D. R. Garrett, A. Sigouin, M. Bélisle, D. Garant, L. Haroune, J. Bellenger, and F.

Pelletier. 2021. Assessing pesticides exposure effects on the reproductive performance of a

declining aerial insectivore. Ecological Applications 0:1–13.

Porlier, M., M. Bélisle, and D. Garant. 2009. Non-random distribution of individual genetic

diversity along an environmental gradient. Philosophical Transactions of the Royal Society

B: Biological Sciences 364:1543–1554.

Poulin, B., G. Lefebvre, and L. Paz. 2010. Red flag for green spray: adverse trophic effects of Bti

on breeding birds. Journal of Applied Ecology 47:884–889.

Pyke, G. H., H. R. Pulliam, and E. L. Charnov. 1977. Optimal Foraging: A Selective Review of

Theory and Tests. The Quarterly Review of Biology 52:137–154.

QGIS. 2020. QGIS Geographic Information System. QGIS Association. http://www.qgis.org.

Quinney, T. E. 2019. Wilson Ornithological Society. Pages 976–976 The Grants Register 2019.

Palgrave Macmillan UK.

Quinney, T. E., D. J. T. Hussell, and C. D. Ankney. 1986. Sources of Variation in Growth of

Tree Swallows. The Auk 103:389–400.

R Core Team. 2019. R: A language and environment for statistical computing.

Rainho, A., and J. M. Palmeirim. 2011. The Importance of Distance to Resources in the Spatial

Modelling of Bat Foraging Habitat. PLoS ONE 6:e19227.

Rendell, W. B., and R. J. Robertson. 1989. Nest-Site Characteristics, Reproductive Success and

Cavity Availability for Tree Swallows Breeding in Natural Cavities. The Condor 91:875.

Rendell, W. B., and R. J. Robertson. 1990. Influence of Forest Edge on Nest-Site Selection by

Tree Swallows. Wilson Journal of Ornithology 102:634–644.

Rioux Paquette, S., D. Garant, F. Pelletier, and M. Bélisle. 2013. Seasonal patterns in Tree

Swallow prey (Diptera) abundance are affected by agricultural intensification. Ecological

Applications 23:122–133.

Robb, G. N., R. A. McDonald, D. E. Chamberlain, and S. Bearhop. 2008. Food for thought:

supplementary feeding as a driver of ecological change in avian populations. Frontiers in

Ecology and the Environment 6:476–484.

Robertson, B. a. 2012. Investigating Targets of Avian Habitat Management to Eliminate an

Ecological Trap. Avian Conservation and Ecology 7:art2.

Robillard, A., D. Garant, and M. Bélisle. 2013. The Swallow and the Sparrow: how agricultural

intensification affects abundance, nest site selection and competitive interactions.

Landscape Ecology 28:201–215.

Rodenhouse, N. L., and R. T. Holmes. 1992. Results of Experimental and Natural Food

Reductions for Breeding Black-Throated Blue Warblers. Ecology 73:357–372.

Rolstad, J. 1991. Consequences of forest fragmentation for the dynamics of bird populations:

conceptual issues and the evidence. Pages 149–163 Metapopulation Dynamics: Empirical

and Theoretical Investigations. Elsevier.

Rosenberg, K. V., A. M. Dokter, P. J. Blancher, J. R. Sauer, A. C. Smith, P. A. Smith, J. C.

Stanton, A. Panjabi, L. Helft, M. Parr, and P. P. Marra. 2019. Decline of the North

American avifauna. Science 366:120–124.

Rossmanith, E., K. Höntsch, N. Blaum, and F. Jeltsch. 2007. Reproductive success and nestling

diet in the Lesser Spotted Woodpecker (Picoides minor): the early bird gets the caterpillar.

Journal of Ornithology 148:323–332.

Ruffino, L., P. Salo, E. Koivisto, P. B. Banks, and E. Korpimäki. 2014. Reproductive responses

of birds to experimental food supplementation: a meta-analysis. Frontiers in Zoology 11:80.

Ruiz, J., and G. Domon. 2009. Analysis of landscape pattern change trajectories within areas of

intensive agricultural use: case study in a watershed of southern Québec, Canada.

Landscape Ecology 24:419–432.

Sæther, B.-E., and O. Bakke. 2000. Avian Life History Variation and Contribution of

Demographic Traits to the Population Growth Rate. Ecology 81:642–653.

Saino, N., R. Ambrosini, D. Rubolini, M. Romano, M. Caprioli, A. Romano, and M. Parolini.

2018. Carry-over effects of brood size on morphology, reproduction, and lifespan in barn

swallows. Behavioral Ecology and Sociobiology 72:30.

Saino, N., S. Calza, P. Ninni, and A. P. Møller. 1999. Barn swallows trade survival against

offspring condition and immunocompetence. Journal of Animal Ecology 68:999–1009.

Saino, N., M. Romano, R. Ambrosini, D. Rubolini, G. Boncoraglio, M. Caprioli, and A.

Romano. 2012. Longevity and lifetime reproductive success of barn swallow offspring are

predicted by their hatching date and phenotypic quality. Journal of Animal Ecology

81:1004–1012.

Seki, S., and H. Takano. 1998. Caterpillar abundance in the territory affects the breeding

performance of great tit Parus major minor. Oecologia 114:514–521.

Seress, G., K. Sándor, K. L. Evans, and A. Liker. 2020. Food availability limits avian

reproduction in the city: An experimental study on great tits Parus major. Journal of Animal Ecology 89:1570–1580.

Seward, A. M., C. M. Beale, L. Gilbert, T. H. Jones, and R. J. Thomas. 2014. The Impact of Increased Food Availability on Reproduction in a Long-Distance Migratory Songbird: Implications for Environmental Change? PLoS ONE 9:e111180.

Shipley, J. R., C. W. Twining, C. C. Taff, M. N. Vitousek, A. Flack, and D. W. Winkler. 2020. Birds advancing lay dates with warming springs face greater risk of chick mortality. Proceedings of the National Academy of Sciences 117:25590–25594.

Shutler, D., D. J. T. Hussell, D. R. Norris, D. W. Winkler, R. J. Robertson, F. Bonier, W. B. Rendell, M. Bélisle, R. G. Clark, R. D. Dawson, N. T. Wheelwright, M. P. Lombardo, P. A. Thorpe, M. a. Truan, R. Walsh, M. L. Leonard, A. G. Horn, C. M. Vleck, D. Vleck, A. P. Rose, L. a. Whittingham, P. O. Dunn, K. a. Hobson, and M. T. Stanback. 2012. Spatiotemporal Patterns in Nest Box Occupancy by Tree Swallows Across North America. Avian Conservation and Ecology 7:art3.

Smith, H. G., and M. Bruun. 1998. The effect of egg size and habitat on starling nestling growth and survival. Oecologia 115:59–63.

Spiller, K. J., and R. Dettmers. 2019. Evidence for multiple drivers of aerial insectivore declines in North America. The Condor 121:1–13.

Staggenborg, J., H. M. Schaefer, C. Stange, B. Naef-Daenzer, and M. U. Grüebler. 2017. Time and travelling costs during chick-rearing in relation to habitat quality in Little Owls Athene noctua. Ibis 159:519–531.

Stanton, R., R. G. Clark, and C. A. Morrissey. 2017. Intensive agriculture and insect prey

1036 availability influence oxidative status and return rates of an aerial insectivore. Ecosphere

1037 8:10.1002/ecs2.1746.

1038 Stanton, R., C. A. Morrissey, and R. G. Clark. 2016. Tree Swallow (Tachycineta bicolor)

1039 foraging responses to agricultural land use and abundance of insect prey. Canadian Journal

1040 of Zoology 94:637–642.

1041 Stanton, R., C. A. Morrissey, and R. G. Clark. 2018. Analysis of trends and agricultural drivers

1042 of farmland bird declines in North America: A review. Agriculture, Ecosystems &

1043 Environment 254:244–254.

1044 Stephens, D. W., J. S. Brown, and R. C. Ydenberg. 2007. Foraging: behavior and ecology.

1045 University of Chicago Press.

1046 Stodola, K. W., D. A. Buehler, D. H. Kim, K. E. Franzreb, and E. T. Linder. 2010. Biotic and

1047 Abiotic Factors Governing Nestling-period Length in the Ovenbird (Seiurus aurocapilla).

1048 The Auk 127:204–211.

1049 Stutchbury, B. J. M., E. A. Gow, T. Done, M. MacPherson, J. W. Fox, and V. Afanasyev. 2011.

1050 Effects of post-breeding moult and energetic condition on timing of songbird migration into

1051 the tropics. Proceedings of the Royal Society B: Biological Sciences 278:131–137.

1052 Teglhøj, P. G. 2017. A comparative study of insect abundance and reproductive success of barn

1053 swallows Hirundo rustica in two urban habitats. Journal of Avian Biology 48:846–853.

1054 Templ, M., K. Hron, and P. Filzmoser. 2011. robCompositions: An R-package for Robust

1055 Statistical Analysis of Compositional Data. Pages 341–355 Compositional Data Analysis.

1056 John Wiley & Sons, Ltd.

Thomas, J. A. 2004. Comparative Losses of British Butterflies, Birds, and Plants and the Global

Extinction Crisis. Science 303:1879–1881.

Thomson, D. L., P. Monaghan, and R. W. Furness. 2007. The demands of incubation and avian

clutch size. Biological Reviews 73:293–304.

Tscharntke, T., A. M. Klein, A. Kruess, I. Steffan☐Dewenter, and C. Thies. 2005. Landscape

perspectives on agricultural intensification and biodiversity - ecosystem service

management. Ecology Letters 8:857–874.

Twining, C. W., N. R. Razavi, J. T. Brenna, S. A. Dzielski, S. T. Gonzalez, P. Lawrence, L. B.

Cleckner, and A. S. Flecker. 2021. Emergent Freshwater Insects Serve as Subsidies of

Methylmercury and Beneficial Fatty Acids for Riparian Predators Across an Agricultural

Gradient. Environmental Science & Technology 55:5868–5877.

Twining, C. W., J. R. Shipley, and D. W. Winkler. 2018. Aquatic insects rich in omega-3 fatty

acids drive breeding success in a widespread bird. Ecology Letters 21:1812–1820.

Vickery, J. A., R. B. Bradbury, I. G. Henderson, M. A. Eaton, and P. V. Grice. 2004. The role of

agri-environment schemes and farm management practices in reversing the decline of

farmland birds in England. Biological Conservation 119:19–39.

Weegman, M. D., T. W. Arnold, R. D. Dawson, D. W. Winkler, and R. G. Clark. 2017.

Integrated population models reveal local weather conditions are the key drivers of

population dynamics in an aerial insectivore. Oecologia 185:119–130.

White, T. C. R. 2008. The role of food, weather and climate in limiting the abundance of

animals. Biological Reviews 83:227–248.

Wickramasinghe, L. P., S. Harris, G. Jones, and N. Vaughan Jennings. 2004. Abundance and

Species Richness of Nocturnal Insects on Organic and Conventional Farms: Effects of

Agricultural Intensification on Bat Foraging. Conservation Biology 18:1283–1292.

Wilson, K. M., M. Kem, and N. T. Burley. 2017. Diet history effects on Zebra Finch incubation

performance: Nest attendance, temperature regulation, and clutch success. The Auk

134:295–307.

Winkler, D. W., and P. E. Allen. 1996. The Seasonal Decline in Tree Swallow Clutch Size:

Physiological Constraint or Strategic Adjustment? Ecology 77:922–932.

Winkler, D. W., K. K. Hallinger, D. R. Ardia, R. J. Robertson, B. J. M. Stutchbury, and R. R.

Cohen. 2020. Tree Swallow (Tachycineta bicolor). Page *in* A. F. Poole, editor. Birds of the

World. Cornell Lab of Ornithology.

Winkler, D. W., M. K. Luo, and E. Rakhimberdiev. 2013. Temperature effects on food supply

and chick mortality in tree swallows (Tachycineta bicolor). Oecologia 173:129–138.

Wood, S. N. 2015. Package 'mgcv.'

Zach, R., and K. R. Mayoh. 1982. Weight and feather growth of nestling tree swallows.

Canadian Journal of Zoology 60:1080–1090.

Zuur, A. F., E. N. Ieno, N. Walker, A. A. Saveliev, and G. M. Smith. 2009. Mixed effects models

and extensions in ecology with R. Page Smart Society. Springer New York.

1096

Tables

Table 1: Mean (±SD) of covariates used to model response variables. Covariates for prey

availability were averaged over two days prior to the collection of each insect sample, whereas

those for fitness and body condition proxies were averaged over either a 12-day (fledging

success) or 16-day (nestling period duration and nestling body mass and wing length) window

post hatching (presented are values for fledging success). Dashes imply the covariates were not

used in the modeling of the response variable. Model groups identify which covariates were

present within models found in Appendix S1: Table S1.

Covariate	Acronym	Unit	Prey availability	Fitness and body condition proxy	Model group
Julian day	JJ	(day)	172.20 ± (12.21)	-	Time
Maximum temperature	MAX	(°C)	25.61 ± (3.75)	25.36 ± (2.31)	Base
Precipitation	RN	(ml)	7.16 ± (11.13)	7.44 ± (4.92)	Base
Site score component 1	Comp.1	-	0.42 ± (1.77)	0.18 ± (1.81)	Land
Site score component 2	Comp.2	-	-0.24 ± (0.91)	-0.18 ± (0.92)	Land
Open water	WT	(%)	-	0.01 ± (0.01)	Base
Prey availability (Diptera dry biomass)	FDM	(g)	-	0.03 ± (0.02)	Food
Hatching day	HD	(day)	-	159.90 ± (5.76)	Base
Brood size	NN	(nestlings)	-	4.83 ± (1.27)	Base
Brood days	BD	(summation of daily brood size)	-	85.63 ± (18.19)	Base

1105

1106 **Table 2**. Standardized coefficient estimates, and their 95% confidence intervals of the top model

1107 for each response variable. See Table 1 for definitions of covariates and Appendix S1: Table S1

1108 for outcome of model selection. Coefficients in bold indicate confidence intervals not

1109 overlapping zero. Numbers next to acronym refer to the order of the polynomial term.

Covariate	Biomass	Fledging success	Duration	Body mass	Wing length
FDM	-	**0.11** (**0.02, 0.19**)	-	**0.10** (**0.01, 0.20**)	-
Comp.1	**-0.11** (**-0.17, -0.06**)	-0.02 (-0.10, 0.06)	**0.17** (**0.08, 0.26**)	-	**-0.58** (**-0.92, -0.24**)
Comp.2	**0.11** (**0.04, 0.17**)	**-0.15** (**-0.28, -0.01**)	0.05 (-0.11, 0.21)	-	-0.20 (-0.81, 0.40)
Comp.1:Comp.2	**-0.04** (**-0.07, -0.01**)	**0.08** (**0.01, 0.15**)	0.04 (-0.05, 0.12)	-	-0.08 (-0.41, 0.25)
Comp.1:FDM	-	**-0.05** (**-0.10, -0.01**)	-	-	-
Comp.2:FDM	-	**0.08** (**0.00, 0.17**)	-	-	-
WT	-	**0.24** (**0.09, 0.39**)	-0.08 (-0.24, 0.07)	0.08 (-0.08, 0.23)	0.49 (-0.10, 1.07)
HD	-	**-0.28** (**-0.35, -0.20**)	-0.05 (-0.15, 0.06)	**-0.26** (**-0.36, -0.16**)	0.11 (-0.27, 0.49)
MAX	-	**0.30** (**0.21, 0.39**)	**-0.33** (**-0.45, -0.21**)	0.04 (-0.08, 0.17)	**1.82** (**1.35, 2.30**)
RN	**-0.07** (**-0.08, -0.05**)	**-0.11** (**-0.20, -0.03**)	0.03 (-0.08, 0.15)	-0.12 (-0.23, 0.00)	0.08 (-0.38, 0.54)
AGSY	-	**-0.25** (**-0.44, -0.06**)	**0.36** (**0.09, 0.63**)	0.01 (-0.25, 0.28)	**-1.38** (**-2.39, -0.38**)
NN	-	**-0.18** (**-0.25, -0.11**)	**0.26** (**0.18, 0.34**)	-	-
BD	-	-	-	**-0.22** (**-0.29, -0.14**)	-0.06 (-0.35, 0.22)
JJ1	**17.77** (**6.40, 29.14**)	-	-	-	-
JJ2	**-10.69** (**-20.31, -1.06**)	-	-	-	-
JJ1:Comp.1	0.40 (-2.99, 3.79)	-	-	-	-
JJ2:Comp.1	**-5.44** (**-8.19, -2.68**)	-	-	-	-
MAX1	**-11.86** (**-14.33, -9.39**)	-	-	-	-
MAX2	**-9.72** (**-11.92, -7.52**)	-	-	-	-
MAX3	1.26 (-0.79, 3.31)	-	-	-	-

 Figure legends

Figure 1: Map of the 40 different farms across a gradient of agricultural intensification within

southern Québec, Canada. Each point represents the location of a farm. Underlying raster

represents the agricultural gradient derived from the Annual crop inventory of 2013 (AAFC

2018), for which each pixel represents either one of the five higher order categories or open

water. The rectangle within inset map represents the extent of the main map and thus the spatial

extent of the study system. Graticules are latitude and longitude.

Figure 2: Robust compositional principal components analysis (PCA) of landscape habitat

composition surrounding each of the 40 farms between 2006 and 2016. Arrows indicate the

eigenvalue and loadings of each higher order habitat. Points within the background represent the

site scores assigned to each farm and year combination and used to define landscape context

(N=440 farms*years)

Figure 3: Predictions for the main covariates of interest. All predictions are unconditional

predictions that have been model averaged, with shrinkage, according to Appendix S1: Table S1.

Each row are predictions for the same response variable. Each plot within the first column,

except for Figure a, is one fitness or body condition proxy plotted against the full range of

observed prey availability. Figure a is the predicted Diptera biomass against the range of Julian

dates. All error bars are unconditional 95% confidence intervals. Each plot within the second

column is a prediction surface of each response variable against the first two components of the

robust PCA describing landscape context. The prediction surface has been reduced to the convex

hull of site scores observed within this study. The range of colors represent the minimum (deep

blue) and maximum (yellow) predicted values of each response variable given site score values.

Individual plots are the predicted Diptera biomass against Julian date of collection (a), the predicted fledging success (b), duration of the nestling period (c), body mass (d) and wing length (e) of nestlings at 16 days of age against the full range of prey availability. The predicted Diptera biomass (f), fledging success (g), duration of the nestling period (h), body mass (i) and wing length (j) of nestlings at 16 days of age against the first two components of the compositional PCA. Predictions and their covariates have been back transformed to either the response or unstandardized scale, respectively. All covariates not explicitly presented in figures were fixed to their mean values prior to predictions.

1141 **Figures**

1142

1143 **Figure 1**

1144

1145 **Figure 2**

1146
1147　**Figure 3**