Full title: Systematic analysis of brain lactate and pH levels in 65 animal models related to

neuropsychiatric conditions

Short title: Brain lactate and pH in neuropsychiatric disorder models

Authors: Hideo Hagihara[1,2], Hirotaka Shoji[1,2], International Brain pH Project Consortium[2],

and Tsuyoshi Miyakawa[1,2]*

Affiliations:

[1] Division of Systems Medical Science, Institute for Comprehensive Medical Science,

Fujita Health University, Toyoake, Aichi 470-1192, Japan

[2] A full list of the authors in the International Brain pH Project Consortium is provided

at the end of the paper along with their affiliations and contributions.

*Corresponding Author: Tsuyoshi Miyakawa (miyakawa@fujita-hu.ac.jp)

Keywords: lactate, pH, brain, neuropsychiatric disorders, animal models, working memory

Abstract

Altered brain energy metabolism associated with increase in lactate levels and the resultant decrease in pH have been increasingly implicated in multiple neuropsychiatric disorders, such as schizophrenia, bipolar disorder, autism spectrum disorder and neurodegenerative disorders. Although it is controversial, change of pH/ lactate level as a primary feature of these diseases, rather than a result of confounding factors such as medication and agonal state, has been evidenced. Animal models that can be studied without such confounding factors inherent to humans are a suitable alternative to understand the controversy. However, the knowledge in animal models regarding brain pH and lactate and their relation to behavioral outcomes is limited in the context of neuropsychiatric disease conditions. In this study, we investigated the common occurrence of changes in the pH and lactate levels in the brain in animal models by analyzing 65 animal models related to neuropsychiatric and neurodegenerative diseases with 1,239 animals. Additionally, we evaluated the behavioral phenotypes relative to the chemical changes in the brain. Among the models, 27 had significant upregulation or downregulation in brain pH and 24, including Hivep2 KO, Camk2a KO, Nrgn KO, Ppp3r1 KO, Chd8 KO, Shank2 KO, Clock mutant, Fmr1 KO, BTBR, APP-J20 Tg, social defeat stress-exposed, corticosterone-treated, and streptozotocin-induced diabetic mice, demonstrated changes in lactate levels. Meta-analysis of the data revealed a highly significant negative correlation between brain pH and lactate levels, suggestive of

increased lactate levels causing decreased brain pH. Statistical learning algorithm based

on the comprehensive data has revealed that the increased brain lactate levels can be

predominantly predicted by the indices for the percentage of correct response in working

memory test, with a significant simple, negative correlation. Our results suggest that brain

energy metabolism is commonly altered in many animal models of neuropsychiatric and

neurodegenerative diseases, which may be associated with working memory performance.

We consider our study to be an essential step suggesting that the brain endophenotypes

serve as a basis for the transdiagnostic characterization of the biologically heterogeneous

and debilitating cognitive illnesses. Based on these results, we are openly accepting

collaborations to extend these findings and to test the hypotheses generated in this study

using more animal models. We welcome any mice/rat models of diseases with or without

any behavioral phenotypes.

Introduction

Neuropsychiatric disorders, such as schizophrenia (SZ), bipolar disorder (BD), major

depressive disorder (MDD), autism spectrum disorder (ASD), and Alzheimer's disease (AD),

are common with a prevalence of more than one-third of the population in most countries

being diagnosed with at least one such disorder at some point in their life (1). Although

these diseases clinically fall into different diagnostic categories, some biological features,

such as genetic mutations, molecular changes, and brain activity alterations, are common

among them (2–6), suggesting a common underlying biological basis. Increasing evidence

suggests that metabolic alterations in the brain are shared by the multiple

neuropsychiatric disorders. Increases in the levels of lactate, an end-product of glycolysis

pathway, have been observed in the brain of patients with SZ, BD, ASD, MDD, and epilepsy

(7–15). Increased lactate levels is considered to lead to decreased pH and are associated

with brain energy deficits (12). Recent large-scale meta-analyses have confirmed

increased brain lactate and decreased pH in SZ and BD (16,17). Such increased lactate and

decreased pH have also been observed in the brains of patients with AD (18–24). However,

the observed phenomena are potentially confounded by secondary factors inherent in

human studies, such as antipsychotic treatments (10). Agonal experiences associated with

these disorders may also complicate the interpretation of postmortem study results (25–

27). Although some human studies suggest that medication use is not a major factor for

regulating brain pH and lactate levels (7,10,11,15,28), excluding the effects of other

potential confounding factors in human studies especially using postmortem brain

samples is technically difficult. Animal models, devoid of such confounding factors, may

help to confirm whether increased brain lactate and decreased pH levels are associated

factors.

Recently, increased brain lactate and decreased pH levels were demonstrated to

be commonly found in five strains of neurodevelopmental mouse models of psychiatric

disorders (29). As all of the mice used in the study were drug-naïve, with equivalent

agonal states, postmortem intervals, and ages within each strain, those findings in mouse

models suggest that increased lactate and decreased pH reflect an underlying

pathophysiology, rather than mere artifacts, in at least a subgroup of patients with these

mental disorders. However, the knowledge of brain pH and lactate in the animal models is

limited to small numbers of models and systematic evaluations using the same platform

have not been conducted so far in animal models. Therefore, we have extended our

previous study (29) to a larger variety of animal models of neuropsychiatric disorders, as

well as of neurodegenerative disorder, AD, and peripheral diseases or insults that are

comorbid with psychiatric conditions (e.g., diabetes mellitus (DM), colitis, and peripheral

nerve injury). Those animal models include 65 strains or conditions of mice and rats with

genetic modifications, drug treatments, and other experimental manipulations (Table 1).

Combining the large-scale brain lactate data with behavioral data (e.g., working memory,

locomotor activity, anxiety-like behavior, and depression-like behavior), we also sought to

investigate the relations between alterations in brain lactate levels and behavioral

outcomes.

Results

Altered brain pH and lactate levels in animal models

The raw data of brain pH and lactate, and detailed information of animals (age, sex, and

treatment methods) are included in Supplementary Table 1. Among the 65

strains/conditions, 27 demonstrated significant changes in pH (5 increased, 22 decreased)

and 24 in lactate (19 increased, 5 decreased) in comparison with the corresponding

control animals (P <0.05; Supplementary Figure 1 and Supplementary Table 2).

Hierarchical clustering based on effect size and direction of changes classified those 65

models into four groups: high lactate/low pH group, moderate high lactate/moderate low

pH group, low lactate/high pH group, and a group with minimal to no changes in lactate or

pH, consisting of 16, 6, 15, and 28 models, respectively (Figure 1), where high and low

mean higher and lower in mutant/experimental animals related to the corresponding wild

type/control animals, respectively. High lactate/low pH group included, for example, SZ

model Ppp3r1 KO mice and Nrgn KO mice, SZ/intellectual disability (ID) model Hivep2 (also

known as Shn2) KO mice, AD model APP-J20 Tg mice, and ASD model Chd8 KO mice. Low

lactate/high pH group included mainly mouse models for ASD or developmental delay,

such as Shank2 KO mice, Fmr1 KO mice, BTBR mice, Stxbp1 KO mice, Dyrk1 KO mice, Auts2

KO mice, and patDp mice (Figure 1).

The Z-score-based meta-analysis of 1,239 animals analyzed in this study

revealed a highly significant negative correlation between brain lactate and pH levels

individually (Figure 2, Supplementary Figure 2), supporting the idea that decreased pH is

due to increased lactate levels in the pathological conditions related to neuropsychiatric

disorders.

Poorer working memory performance predicts higher brain lactate levels

Most of the animal models we examined are known to show a wide range of behavioral

abnormalities, such as deficits in learning and memory, and increased depression-like,

anxiety-like behaviors or impaired sensorimotor gating. Thereafter, with our

comprehensive lactate data, we examined potential relation of lactate alterations to their

behavioral phenotypes. Therefore, we examined whether behavioral patterns could

predict brain lactate levels by applying a statistical learning algorithm, which could

discover intrinsic links between the chemical signatures in the brain and behaviors. Of the

65 animal models, we collected comprehensive behavioral data of 24 mouse models,

which were available in public source (e.g., published papers and database repository) or

in-house studies (see Methods and Materials; Supplementary Table 3). We constructed an

effect size-based model for predicting the brain lactate levels from behavioral data using

leave-one-out cross-validation method. Statistical evaluation of the prediction accuracy of

the model revealed a significant correlation between the actual and the predicted brain

lactate levels (Figure 3a), indicating that behavioral measures have a potential to predict

the brain lactate levels of individual models.

The prediction analysis was implemented to evaluate the behavioral measures

most useful to characterize the brain lactate levels of individual strains. The prediction

algorithm used identified behavioral signatures related to brain lactate levels by weighting

behavioral measures according to their individual predictive strength. Thus, we identified

the behavioral measures accompanying changes in brain lactate levels by examining the

weighted behavioral measures used for the prediction in linear regression. Three out of

nine behavioral measures were selected to build the successful prediction model and in

those measures for working memory was the most selected (Figure 3b). According to

simple correlation analysis, the measures for working memory were negatively correlated

with the brain lactate levels (Figure 3c). These results suggest that higher lactate levels in

the brain are related to lower performance in working memory tests in mouse models of

neuropsychiatric disorders.

Effects of age and sex on the brain pH and lactate levels

Ages at sampling were matched within each strain/condition, but varied among

strains/conditions, ranging from 5 to 103 weeks old in mice (Supplementary Table 1). No

significant correlation was found between pH and age in wild type/control mice. Brain

lactate levels had a significant negative correlation with age (Supplementary Figure 5),

consistent with a previous MRS study in mice (30). However, limitations need to be

considered in interpretating our results, such as, differences in genetic background and

handling conditions before sampling (some mice had received repeated intraperitoneal

injections or behavioral tests, and others had been kept undisturbed until sampling)

among strains/conditions. We further examined the effects of sex on the brain pH and

lactate levels. To minimize the effects of the limitations mentioned above, we used

Z-scores that were calculated within each strain/condition and focused on

strains/conditions with mixed gender. Female had significantly higher pH and lower

lactate levels than male in wild type/control animals (Supplementary Figure 6).

Discussion

We performed a comprehensive analysis of brain pH and lactate in 65 animal models. The

data suggested the diversity of brain-energy-metabolism among these model animals.

Some mouse strains considered to model different diseases were found to exhibit similar

pattern of changes in pH and lactate levels. Specifically, SZ models (Ppp3r1 KO and Nrgn

KO mice), SZ/ID model (Hivep2 KO mice), BD/ID model (Camk2a KO mice), ASD model

(Chd8 KO mice), depression models (mice exposed to social defeat stress,

corticosterone-treated mice and Sert KO mice), AD model (APP-J20 Tg mice), and DM

model (STZ-treated mice) commonly exhibited decreased brain pH and increased lactate

levels. A BD model Polg1 Tg mice showed no differences in pH or lactate levels. However,

other BD model (Clock mutant mice) and ASD models, such as Shank2 KO (31), Fmr1 KO,

Dyrk1 KO (32), Auts2 KO (33), and patDp mice (34), were classified into a group with

opposite changes, or decreased lactate and increased pH group. Animal models with

different patterns of changes in brain pH and lactate levels may represent subpopulations

of patients or specific states of the diseases (13). While increased brain lactate levels in

neuropsychiatric conditions are almost consistent in the literature, decreased lactate

levels has also been found in a cohort of patients with SZ (35) and in euthymic state of BD

(36). Our results from animal studies may also support the idea that the patients

categorized based on the symptoms to particular neuropsychiatric disorders are

biologically heterogeneous (37) from a brain-energy-metabolism viewpoint.

178   The present animal studies revealed an extraordinarily high negative correlation

between brain lactate and pH levels, strengthening our previous findings from small-scale

animal studies (29). Negative correlation between them has been found in human

postmortem study (10). These results suggest that brain lactate is a main regulator of the

tissue pH (12), although we could not exclude the possibility that other factors such as

neuronal activity-regulated production of carbon dioxide, another metabolic acid, may

also contribute to the changes in brain pH (38,39).

We observed no significant correlation between age and brain pH in wild

type/control mice. In human studies, inconsistent results have been obtained with regard

to correlation between brain pH and age. Some studies showed no significant correlation,

(40,41), whereas other studies showed a negative correlation (42,43). Sex effects on brain

pH is also inconsistent in human studies (40,41). Systematic analysis focusing on the

effects of age and sex on the brain pH using animal models may help explain the

inconsistency found in the human studies.

Does brain lactate exert favorable or unfavorable effects on learning and

memory functioning? Our prediction analysis highlighted that poorer working memory

performance may be predominantly associated with higher lactate levels in animal models

of neuropsychiatric disorders (Figure 3). Additionally, in human studies, higher lactate has

been associated with lower cognition in the individuals with SZ (14) and mild cognitive

impairment (44). Given these observations, lactate production may be expected to exert

negative impacts on brain functions, especially memory formation. However, lactate

production stimulated by learning tasks has been suggested as requisite for memory

formation. Lactate production by the astrocytic glycogenolysis and its transport to

neurons serves as an energy substrate for neuronal activity, referred to as

astrocyte-neuron lactate shuttle (ANLS). Animal studies have demonstrated that the

pharmacological disruption of learning task-stimulated lactate production and transport

via the ANLS immediately before the testing impaired memory formation as assessed by

the plus-shaped maze spontaneous alteration task (testing short-term memory) (45) and

in the inhibitory avoidance task (testing long-term memory) (46,47). Collectively,

considering that brain lactate levels increase during stimulations in a temporally (and

spatially) restricted manner under physiological conditions (48,49), pathologically

persistent elevation of brain lactate levels may exert negative impact on brain functions

including memory processing, although the causality is unknown. Other possibility is that

decreased consumption of lactate for energy production due to mitochondrial dysfunction

in neurons may underline the impaired learning and memory functioning in the disease

conditions. Mitochondrial dysfunction has been thought to lead to lactate accumulation

because of insufficient capacity of mitochondrial metabolism to metabolize lactate that

was produced (16,50,51). Mitochondrial dysfunction has been consistently implicated in

multiple neuropsychiatric disorders, including SZ, BD, MDD, ASD, and AD (52–54), among

which working memory deficits are common symptoms (55). In addition, given that lactate

rise reflects neuronal activation (29) and multiple brain regions are abnormally activated,

activation in the brain regions other than frontal cortex, one of the brain regions critical

for working memory (56), interfere with working memory performance, as proposed that

activity of core brain region could be interfered with noise from the rest on cognitive tasks

in patients with SZ (57). There is also the possibility that increased lactate may have a

beneficial effect to compensate for the impaired memory and cognition, as lactate

administration that increases brain lactate levels has been shown to attenuate cognitive

deficits in human patients (58) and rodent model (59) of traumatic brain injury.

Additionally, lactate administration has also been shown to exert antidepressant effects in

depression model mice (60). We also cannot exclude the possibility that increased lactate

is also involved in behavioral alterations other than memory deficit per se, such as anxiety,

as we have found that increased brain lactate levels were associated with altered

anxiety-like behaviors in social defeat stress model of depression (61). Further studies are

required to address these issues, for example, by chronically inducing deficits of

mitochondria function to manipulate endogenous lactate levels in a brain-region-specific

manner and analyzing its effects on working memory.

As we used whole brain samples to measure the pH and lactate levels, we could

not determine whether the observed changes in pH/lactate levels occur ubiquitously in

the entire brain or selectively in specific brain region(s) in each strain or condition of the

models. Indeed, brain region-specific increase in lactate levels was observed in human

patients with ASD in the MRS study (8). The brain region-specific changes may occur even

in animal models in which significant changes were not detected in the present study and,

if so, such differences could be masked in the analysis using whole brain samples. Further

studies are required to address this issue, for example, by means of the measurements in

micro-dissected brain samples and in vivo analyses using pH- or lactate-sensitive biosensor

electrode (45,62) and MRS (63).

In conclusion, the present study demonstrated that altered brain pH and lactate

levels were commonly observed in many animal models of SZ, BD, ASD, AD and other

neuropsychiatric disorders. These findings provide further evidence supporting the idea

that altered brain pH and lactate levels are not mere artifacts such as medication

confounding, but rather implicated in the underlying pathophysiology of, at least

subpopulations of, patients with the diseases. Alteration in the brain-energy-metabolism

or hyper- or hypo-activity of neurons in the brain leading to abnormal lactate and pH

levels may serve as a potential therapeutic target of neuropsychiatric disorders (17). In

addition, detection of brain lactate, such as by MRS, may help to diagnose and

subcategorize such biologically heterogeneous diseases, as shown in mitochondrial

disease (64). Future studies to identify the effective treatment strategies specific to the

sets of animal models that could recapitulate diversity of brain-energy-metabolism in

human disease conditions may contribute to development of improved treatments for the

biologically defined subgroups of patients or disease states of the debilitating illnesses

beyond the clinically defined borders.

Table 1. Animal models used in this study

	Name	Description	Related diseases/conditions
1	APP Tg	Mice expressing familial Alzheimer's disease-mutant human amyloid beta precursor protein (PDGF-hAPP$_{swe/Ind}$, line J20) (65)	AD(66,67)
2	Arid1b KO	Mice with heterozygous knockout of the AT-rich interaction domain 1b (68)	ASD(69,70)
3	Auts2 KO	Mice with heterozygous knockout of the Autism susceptibility candidate 2 (33)	ASD(71–73), ID(74), SZ(75)
4	Barp KO	Voltage gated calcium channel beta-anchoring and -regulatory protein KO mice (76)	

5	Bdnf KO	Brain derived neurotrophic factor KO mice* (JAX, 004339)	
6	BTBR	Inbred mouse strain BTBR T+ tf/J (77,78)	ASD
7	Camk2a KO	Mice with heterozygous knockout of the calcium/calmodulin-dependent protein kinase II alpha (79–81)	BD(82–84), SZ(85)
8	Camkk1 KO	Mice with forebrain-specific constitutively active form of calcium/calmodulin kinase kinase 1 (86)	
9	Ccnd2 KO	Cyclin D2 KO mice (87)	
10	CFA treatment	Mouse model of chronic inflammatory pain induced by complete Freund's adjuvant (CFA) (88,89)	Chronic pain
11	Chd8 KO	Mice with heterozygous knockout of the	ASD(91–95)

		long isoform of chromodomain helicase DNA-binding protein 8 (90)	
12	Chn1 KO	Chimerin 1 (α-chimerin) KO mice (96)	ASD(96)
13	Clock mutant	Mice with N-ethyl-N-nitrosourea-induced mutation in circadian locomotor output cycles kaput (JAX, 002923) (97,98)	BD(99,100), SZ(101)
14	Corticosterone treatment	Mice chronically treated with corticosterone (102,103)	MD(104–106)
15	Crmp2 KO	Collapsin response mediator protein 2 KO mice (107)	AD(108), SZ(109)
16	Dextran treatment	Mice treated with dextran sulfate sodium (110)	Colitis
17	Disc1-L100P mutant	Mice with N-ethyl-N-nitrosourea-induced L100P amino acid exchange mutation in	SZ(112–114)

		exon 2 of Disrupted-in-Schizophrenia 1 (111)	
18	Disc1-Q31L mutant	Mice with N-ethyl-N-nitrosourea-induced Q31L amino acid exchange mutation in exon 2 of Disrupted-in-Schizophrenia 1 (111)	SZ(112–114)
19	Dyrk1a KO	Mice with heterozygous knockout of the dual specificity tyrosine phosphorylation regulated kinase 1a (32)	ASD/ID(70,115,116)
20	ECS treatment	Mice treated with electroconvulsive stimulation (117,118)	Treatment for MDD(119,120)
21	Fmr1 KO	Fragile X mental retardation protein translational regulator 1 KO mice (121)	ASD, FMR, SZ(85)
22	Gasc1	Gene amplified in squamous cell	ASD(124)

		carcinoma 1 hypomorphic mutant mice (122,123)	
	hypomorph		
23	Glra4 KO	Glycine receptor alpha 4 KO mice (125)	ID(126)
24	Grin1 KO (postnatal)	GABArgic neuron-specific glutamate receptor, ionotropic, NMDA1 KO mice (Protein phosphatase 1, regulatory subunit 2-cre; Grin1$^{loxP/loxP}$) (127)	SZ(128,129)
25	Grin1 KO (adult)	GABArgic neuron-specific glutamate receptor, ionotropic, NMDA1 KO mice (Protein phosphatase 1, regulatory subunit 2-cre; Grin1$^{loxP/loxP}$) (127)	SZ(128,129)
26	Gunn rat	Gunn rats (Gunn/Slc-j/j) (130)	SZ(131)
27	Hivep2 KO	Human immunodeficiency virus type 1 enhancer binding protein 2 (Schnurri-2) KO	ID(133,134), SZ(132)

		mice (132)	
28	Hyponatremia	Mice treated with 1-deamino-8-D-arginine vasopressin and fed with a liquid formula (135–137)	DS(138,139)
29	Il18 KO	Interleukin 18 KO mice (140,141)	DM(142)
30	Ketamine treatment	Mice treated with ketamine (143)	Psychosis(144)
31	Lurasidone treatment	Mice treated with lurasidone (145)	Atypical antipsychotic(146,147)
32	Mdga1 KO	MAM domain containing glycosylphospatidylinositol anchor 1 KO mice (148)	SZ(149–151)
33	Mdga2 KO	Mice with heterozygous knockout of the	ASD(153,154)

		MAM domain containing glycosylphospatidylinositol anchor 2 (152)	
34	Methamphetamine treatment	Mice treated with methamphetamine (155)	Psychosis(156)
35	Nhe5 KO	Na$^+$/H$^+$ exchanger 5 KO mice (157)	
36	Nlgn3-R451C KI	Mice with R451C amino acid exchange mutation in neuroligin 3 (77,158)	ASD(159,160)
37	Nr3c1 Tg	Mice overexpressing glucocorticoid receptor under the Camk2a promoter	MD(161)
38	Nrgn KO	Neurogranin KO mice (162–164)	SZ(165,166)
39	Oxamate treatment	Mice treated with sodium oxamate, an inhibitor of lactate dehydrogenase	
40	Pacap KO	Pituitary adenylate cyclase-activating	MD(168), SZ(169)

		polypeptide KO mice (167)	
41	patDp	Mice with a paternal duplication of human chromosome 15q11-13 (34)	ASD(170–173)
42	Phencyclidine treatment	Subchronic phencyclidine-treated mice (145,174)	SZ(175)
43	PCP+Lur	Phencyclidine (PCP)- and lurasidone (Lur)-treated mice (145,174)	
44	Polg1 Tg	Forebrain-specific catalytic subunit of mitochondrial DNA polymerase KO mice (176)	BD(177)
45	Ppp3r1 KO	Forebrain-specific protein phosphatase 3, regulatory subunit B, alpha isoform (calcineurin B, type 1) KO mice (178,179)	SZ(180)
46	Quinpirole	Mice treated with quinpirole, a dopamine	OCD(182)

		treatment	D2 receptor agonist (181)	
47	Reln Tg		Mice lacking the C-terminal region of Reelin (183)	ASD(184–186), BD(187), SZ(188)
48	Restraint stress		Mice exposed to chronic restraint stress (189)	Chronic stress
49	Sciatic nerve cuffing		The sciatic nerve cuffing mouse model of neuropathic pain (190,191)	Chronic pain
50	Scn2a KO		Mice with heterozygous knockout of the sodium voltage-gated channel alpha subunit 2 (192)	ASD(193,194), EP(195–197), ID(198,199)
51	Sert KO		Serotonin transporter KO mice (200)	ASD(201,202)
52	Shank2 KO		SH3 and multiple ankyrin repeat domain 2 KO mice (31)	ASD(154)
53	Shank3 KO		SH3 and multiple ankyrin repeat domain	ASD(204–206)

		3b KO mice (JAX, 017688) (203)	
54	Snap25-S187A KI	Mice with S187A amino acid exchange mutation in synaptosomal-associated protein of 25 kDa	ADHD(207–212), EP(213,214), SZ(215,216)
55	Social defeat stress (acute)	Mice exposed to social defeat stress (217,218)	Acute stress
56	Social defeat stress (chronic)	Mice exposed to social defeat stress (219,220)	Chronic stress
57	Streptozotocin treatment	Mice treated with streptozotocin (221)	DM(222)
58	Streptozotocin + restraint stress	Mice treated with streptozotocin and exposed to chronic restraint stress (189,221)	DM and DS comorbidity(223)

59	Stxbp1 KO	Mice with heterozygous knockout of the syntaxin-binding protein 1 (224)	ASD/ID(116,199,225), EP(226,227)
60	Syngap1 KO	Mice with heterozygous knockout of the synaptic Ras GTPase-activating protein 1 (228,229)	ID, SZ, ASD(154), EP(226)
61	Thalidomide treatment	Rats prenatally exposed to thalidomide (230,231)	ASD(232)
62	Tnx KO	Tenascin X KO mice (233,234)	EDS(235), SZ(236–238)
63	Trx1 KO	Rats with heterozygous knockout of the thioredoxin 1	EP
64	Tsc1 KO	Astrocyte-specific tuberous sclerosis complex 1 KO mice (Glial fibrillary acidic protein-cre; Tsc1$^{loxP/loxP}$) (239)	TSC(240)

65	Valproic acid treatment	Mice prenatally exposed to valproic acid (241)	ASD(242)

AD, Alzheimer's disease; ADHD, attention-deficit/hyperactivity disorder; ASD, autism

spectrum disorders; BD, bipolar disorder; DM, diabetes mellitus; EDS, Ehlers-Danlos

syndrome; DS, depression symptom; EP, epilepsy; FMR, Fragile X mental retardation; ID,

intellectual disability, KI, knock-in; KO, knock out; MD, major depressive disorder; OCD,

obsessive-compulsive disorder; SZ, schizophrenia; Tg, transgenic; TSC, tuberous sclerosis

complex. *Mice with off-target deletion of conditional Bdnf allele derived from Bdnf^{2lox}

mouse line.

Materials and Methods

Experimental animals and ethical statement

Mice and rats used in this study are listed in Table 1. Animal experiments were approved

by the Institutional Animal Care and Use Committee of Fujita Health University, based on

the Law for the Humane Treatment and Management of Animals and the Standards

Relating to the Care and Management of Laboratory Animals and Relief of Pain. Every

effort was made to minimize the number of animals used.

Sampling and handling of the brain samples

Upon the study, a standardized protocol regarding sampling and handling of the brain

samples has been established to minimize potential confounding effects because of the

technical differences among laboratories and performing blind studies, as follows:

Animals and samples

• Animals: Mouse and rat. For genetically engineered animals, mutants and their

wild-type littermates should be used.

• Number of animals: ≧6 per group (identical genetic background, littermate),

preferably.

• Sex of animals: All males, all females, or balanced among groups if mixed.

• Samples: Fresh-frozen whole brain.

Blind study

pH measurements were blinded: Upon sampling, the researchers were supposed to

randomize the animals regarding genotype and collect brain samples into tubes labeled

with serial numbers. The researchers were asked to provide the genotype information and

the corresponding serial numbers for the following statistical analyses, after the

measurements.

Brain sampling procedures

1. Sacrifice mouse/rat by cervical dislocation followed by decapitation, and remove

whole brain from the skull. Do not immerse the brain in any buffer solutions or

water.

2. Cut the brain along the longitudinal fissure of the cerebrum.

3. Collect the left and right hemispheres into a tube that can be tightly capped like

Cryotube and seal the caps with Parafilm (to minimize the effect of carbon dioxide

from dry ice on the tissue pH during transportation.).

4. Snap freeze in liquid N2, and store at -80C until the shipment.

5. Transport the frozen brain on dry ice.

The   protocol   is   also   publicly   available   at

http://www.fujita-hu.ac.jp/~cgbb/en/collaborative_research/index.html.

Measurements of tissue pH and lactate levels

Whole brain was used to measure pH and lactate levels as previously described (29).

Briefly, snap-frozen tissues were homogenized in ice-cold distilled H_2O (5 ml per 500 mg of

tissue). The pH of the homogenates was measured using a pH meter (LAQUA F-72,

HORIBA, Ltd., Kyoto, Japan) equipped with a Micro ToupH electrode (9618S-10D, HORIBA,

Ltd.) after a three-point calibration at pH 4.0, pH 7.0, and pH 9.0. Subsequently, the

concentration of lactate in the homogenates was determined using a multi-assay analyzer

(GM7 MicroStat, Analox Instruments, London, UK) after calibration with 8.0 M lactate

standard solution (GMRD-103, Analox Instruments). A 20-µl aliquot of centrifuged

supernatant (14,000 rpm, 10 min) was used for the measurement.

Effect size (d) was calculated for each strain/condition and each measure (e.g., pH, lactate

value, and behavioral index), as followed:

323            $d = (M_{mutants} - M_{controls})/S_{pooled}$

324    $S_{pooled} = [(S^2_{mutant} + S^2_{control})/2]^{1/2}$

325 The heat map was depicted using the R (version 3.5.2) gplots package.

326   Z-score transformation, a traditional method of data normalization for direct

327 comparison between different samples and conditions, was applied for each pH or lactate

328 value using individual animal data within each of strain, according to the following

329 formula:

330   Z-score = (value$_P$ – mean value$_{P1...Pn}$)/standard deviation$_{P1...Pn}$  ,

331 where P is any pH or lactate and $P1...P_n$ represent the aggregate measure of all pH or

332 lactate values.

334 **Prediction analysis**

335 We collected the comprehensive behavioral data as much as of animal models whose

336 brain pH and lactate levels were examined in this study. The following behavioral data of

337 24 animal models were obtained from published papers, Mouse Phenotype Database

(http://www.mouse-phenotype.org/), or in-house studies (Supplementary Table 3):

number of transitions in the light-dark transition test, percentage of immobile in the

forced swim test, time spent in open arm in the elevated plus maze test, prepulse

inhibition at 78-110 dB and 74-110 dB, startle response at 120 dB, distance traveled in the

open field test, and correct percentage in the T-maze, Y-maze, or other maze test from

APP Tg mice, Arid1b KO mice mice, Barp KO mice, BTBR mice, Camk2a KO mice, complete

Freund's adjuvant-treated mice, Chd8 KO mice, corticosterone-treated mice, Disc1-L100P

mutant mice, Disc1-Q31L mutant mice, Gasc1 hypomorphic mutant mice, Hivep2 KO mice,

Nhe5 KO mice, Nr3c1 Tg mice, Nrgn KO mice, Pacap KO mice, patDp mice, Ppp3r1 KO mice,

Reln Tg mice, Scn2a KO mice, Sert KO mice, Snap25-S187A KI mice, social defeat

stress-exposed mice, and Syngap1 KO mice. The literature search was conducted in

PubMed and Google Scholar using relevant key words: name of strain or experimental

condition, species (mice or rats), and name of behavioral tests. Among the top hits at the

search, we used the data that were presented in actual values of the mean and SD or SEM,

as priority. For some behavioral measures, possible mean and SD values were estimated

from the graph presented in the paper. In the matrix of strains/conditions and behavioral

measures, those with any missing values were excluded, resulting in obtaining nine

behavioral measures from 24 strains/conditions of mouse models. Effect size was

calculated for each strain/condition and each measure and used for the prediction

analysis.

358   Leave-one-out cross-validation was employed to determine whether behavioral

measures can predict brain lactate levels for individual strain of mice. From the analyzed

behavioral dataset of 24 mouse strains, one sample was selected and excluded to serve as

test data of the cross-validation. Thereafter, a multivariate linear regression model was

trained on the remaining 23 samples using a stepwise variable selection procedure with

EZR software (version 1.38; Saitama Medical Center, Jichi Medical University, Saitama,

Japan) (243), and the test sample was predicted. This was repeated 24 times, in which all

samples were chosen once as the test data. Behavioral measures selected at least one

time in the prediction model were considered as predictive behavioral measures. The

prediction performance was analyzed by evaluating correlation between the predicted

and actual values for the 24 mouse strains.

Statistical analysis

The pH and lactate data were analyzed using unpaired t-test, or one-way analysis of

variance (ANOVA) or two-way ANOVA followed by *post hoc* Tukey's multiple comparison

test using GraphPad Prism 8 (version 8.4.2; GraphPad Software, San Diego, CA).

Consortium

Project conception: Hideo Hagihara[1] and Tsuyoshi Miyakawa[1]. **Preparation of animals and tissues:** Hideo Hagihara[1], Hirotaka Shoji[1], Yoshihiro Takamiya[1], Mika Tanaka[1], Yoko Hagino[2], Hiroko Kotajima-Murakami[2], Kazutaka Ikeda[2], Hikari Otabi[3,4], Atsushi Toyoda[3,4,5], Mohamed Darwish[6], Hirofumi Nishizono[6], Keizo Takao[6,7], Kei Hori[8], Mikio Hoshino[8], Hyopil Kim[9], Bong-Kiun Kaang[9], Takao Kohno[10], Mitsuharu Hattori[10], Ken-ichi Matsumoto[11], Shigeo Okabe[12], Michiru Ida-Eto[13], Masaaki Narita[13], Haruki Fujisawa[14], Yoshihisa Sugimura[14], Tadahiro Numakawa[15], Hiroshi Kunugi[15], Katsuhiko Tabuchi[16], Yuta Katayama[17], Keiichi I Nakayama[17], Masayuki Matsushita[18], Johji Inazawa[19], Tohru Yamamoto[20], Haruko Nakamura[21], Yoshio Goshima[21], Hikari Hatakama[22], Nozomi Asaoka[23], Shuji Kaneko[22], Tetsuya Tatsukawa[25], Matthieu Raveau[25], Kazuhiro Yamakawa[24,25], Masafumi Ihara[26], Kyosuke Yamanishi[27], Kiran Sapkota[28], Kazutoshi Nakazawa[28], Isabella A Graef[29], Shuji Wakatsuki[30], Toshiyuki Araki[30], Kota Tamada[31,32], Toru Takumi[31,32], Iori Ohmori[33], Nanette Deneen Hannah[34], Lakshmi Rajagopal[34], Herbert Y Meltzer[34], Ikuo Nobuhisa[35], Tetsushi Kagawa[35], Tetsuya Taga[35], Akito Nakao[36], Yasuo Mori[36], Shota

Katori[37], Takuya Sato[37], Takuji Iwasato[37], Noboru H. Komiyama[38], Seth G. N. Grant[38], Anja

Urbach[39], Léa J Becker[40], Ipek Yalcin[40], Tsuyoshi Takagi[41], Takaoki Kasahara[42], Tadafumi

Kato[42,43], Yoshio Hoshiba[45], Ryuhei Miyake[45], Kisho Obi-Nagata[44], Akiko Hayashi-Takagi[44,45],

Mihiro Shibutani[46], Izuho Hatada[46], Shunsuke Ishii[47], Atsuko Hayata-Takano[48,49], Hitoshi

Hashimoto[48-52], Noriko Takahashi[53,54], Haruo Kasai[53,55], Emiko Okuda-Ashitaka[56], Freesia L

Huang[57], Tadayuki Shimada[58], Kanato Yamagata[58]. **Biochemical and statistical analyses:**

Hideo Hagihara[1] and Giovanni Sala[1]. **Behavioral analysis:** Hideo Hagihara[1], Hirotaka Shoji[1],

Giovanni Sala[1], Yoshihiro Takamiya[1]. **Writing of the manuscript:** Hideo Hagihara[1] and

Tsuyoshi Miyakawa[1]. All authors approved the final manuscript.

1. Division of Systems Medical Science, Institute for Comprehensive Medical Science,

Fujita Health University, Toyoake, Aichi, Japan

2. Addictive Substance Project, Tokyo Metropolitan Institute of Medical Science, Tokyo,

Japan

3. College of Agriculture, Ibaraki University, Ami, Ibaraki, Japan

4. United Graduate School of Agricultural Science, Tokyo University of Agriculture and

Technology, Fuchu, Tokyo, Japan

5. Ibaraki University Cooperation between Agriculture and Medical Science (IUCAM), Ami,

Ibaraki, Japan

6. Department of Behavioral Physiology, Graduate School of Innovative Life Science,

University of Toyama, Toyama, Japan

7. Life Science Research Center, University of Toyama, Toyama, Toyama, Japan

8. Department of Biochemistry and Cellular Biology, National Institute of Neuroscience,

National Center of Neurology and Psychiatry, Kodaira, Tokyo, Japan

9. Department of Biological Sciences, College of Natural Sciences, Seoul National

University, Seoul, South Korea

10. Department of Biomedical Science, Graduate School of Pharmaceutical Sciences,

Nagoya City University, Nagoya, Aichi, Japan

11. Department of Biosignaling and Radioisotope Experiment, Interdisciplinary Center for

Science Research, Organization for Research and Academic Information, Shimane

University, Izumo, Shimane, Japan

12. Department of Cellular Neurobiology, Graduate School of Medicine, The University of

Tokyo, Tokyo, Japan

13. Department of Developmental and Regenerative Medicine, Mie University, Graduate

School of Medicine, Mie, Japan

14. Department of Endocrinology & Metabolism, Fujita Health University School of

Medicine, Toyoake, Aichi, Japan

15. Department of Mental Disorder Research, National Institute of Neuroscience, National

Center of Neurology and Psychiatry, Tokyo, Japan

16. Department of Molecular & Cellular Physiology, Shinshu University School of Medicine,

Matsumoto, Nagano, Japan

17. Department of Molecular and Cellular Biology, Medical Institute of Bioregulation,

Kyushu University, Fukuoka, Japan

18. Department of Molecular and Cellular Physiology, Graduate School of Medicine,

University of the Ryukyus, Nishihara, Okinawa, Japan

19. Department of Molecular Cytogenetics, Medical Research Institute, Tokyo Medical and

Dental University, Tokyo, Japan

20. Department of Molecular Neurobiology, Faculty of Medicine, Kagawa University,

Kagawa, Japan

21. Department of Molecular Pharmacology and Neurobiology, Yokohama City University

Graduate School of Medicine, Yokohama, Kanagawa, Japan

22. Department of Molecular Pharmacology, Graduate School of Pharmaceutical Sciences,

Kyoto University, Kyoto, Kyoto, Japan

23. Department of Pharmacology, Kyoto Prefectural University of Medicine, Kyoto, Kyoto,

Japan

24. Department of Neurodevelopmental Disorder Genetics, Institute of Brain Sciences,

Nagoya City University Graduate School of Medical Sciences, Nagoya, Aichi, Japan

25. Laboratory for Neurogenetics, RIKEN, Center for Brain Science, Wako, Saitama, Japan

26. Department of Neurology, National Cerebral and Cardiovascular Center, Suita, Osaka,

Japan

27. Department of Neuropsychiatry, Hyogo College of Medicine, Nishinomiya, Hyogo,

Japan

28. Department of Neuroscience, Southern Research, AL, USA

29. Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA

30. Department of Peripheral Nervous System Research, National Institute of

Neuroscience, National Center of Neurology and Psychiatry, Kodaira, Tokyo, Japan

31. Department of Physiology and Cell Biology, Kobe University School of Medicine, Kobe,

Hyogo, Japan

32. RIKEN Brain Science Institute, Wako, Saitama, Japan

33. Department of Physiology, Okayama University Graduate School of Medicine,

Dentistry and Pharmaceutical Sciences, Okayama, Okayama, Japan

34. Department of Psychiatry and Behavioral Sciences, Northwestern University Feinberg

School of Medicine, Chicago, IL, USA

35. Department of Stem Cell Regulation, Medical Research Institute, Tokyo Medical and

Dental University, Tokyo, Japan

36. Department of Synthetic Chemistry and Biological Chemistry, Graduate School of

Engineering, Kyoto University, Kyoto, Japan

37. Laboratory of Mammalian Neural Circuits, National Institute of Genetics, Mishima,

Shizuoka, Japan

38. Genes to Cognition Program, Centre for Clinical Brain Sciences, University of Edinburgh,

Edinburgh, UK

39. Hans Berger Department of Neurology, Jena University Hospital, Jena, Germany

40. Institut des Neurosciences Cellulaires et Intégratives, Centre National de la Recherche

Scientifique, Université de Strasbourg, Strasbourg, France

41. Institute for Developmental Research, Aichi Developmental Disability Center, Kasugai,

Aichi, Japan

42. Laboratory for Molecular Dynamics of Mental Disorders, RIKEN Center for Brain

Science, Wako, Saitama, Japan

43. Department of Psychiatry and Behavioral Science, Juntendo University Graduate

School of Medicine, Tokyo, Japan

44. Laboratory for Multi-scale Biological Psychiatry, RIKEN Center for Brain Science, Wako,

Saitama, Japan

45. Laboratory of Medical Neuroscience, Institute for Molecular and Cellular Regulation,

Gunma University, Maebashi, Gunma, Japan

46. Laboratory of Genome Science, Biosignal Genome Resource Center, Institute for

Molecular and Cellular Regulation, Gunma University, Maebashi, Gunma, Japan

47. Laboratory of Molecular Genetics, RIKEN Tsukuba Institute, Tsukuba, Japan

48. Laboratory of Molecular Neuropharmacology, Graduate School of Pharmaceutical

Sciences, Osaka University, Suita, Osaka, Japan

49. United Graduate School of Child Development, Osaka University, Kanazawa University,

Hamamatsu University School of Medicine, Chiba University and University of Fukui, Japan

50. Division of Bioscience, Institute for Datability Science, Osaka University, Suita, Osaka,

Japan

51. Transdimensional Life Imaging Division, Institute for Open and Transdisciplinary

Research Initiatives, Osaka University, Suita, Osaka, Japan

52. Department of Molecular Pharmaceutical Science, Graduate School of Medicine,

Osaka University, Suita, Osaka, Japan

53. Laboratory of Structural Physiology, Center for Disease Biology and Integrative

Medicine, Faculty of Medicine, The University of Tokyo, Tokyo, Japan

54. Department of Physiology, Kitasato University School of Medicine, Sagamihara,

Kanagawa, Japan

55. International Research Center for Neurointelligence (WPI-IRCN), UTIAS, The University

of Tokyo, Tokyo, Japan

56. Department of Biomedical Engineering, Osaka Institute of Technology, Osaka, Japan

57. Program of Developmental Neurobiology, NICHD, NIH, Bethesda, MD, USA

58. Synaptic plasticity project, Tokyo Metropolitan Institute of Medical Science, Tokyo,

Japan

Acknowledgements

This work was supported by MEXT KAKENHI Grant Number JP16H06462, MEXT Promotion

of Distinctive Joint Research Center Program Grant Number JPMXP0618217663, JSPS

KAKENHI Grant Number JP20H00522, JP18K07378, and JP16H06276 (AdAMS), and AMED

Strategic Research Program for Brain Sciences Grant Number JP18dm0107101.

Disclosures

515 Authors declare no conflict of interest regarding this article.

References

1. Cross-national comparisons of the prevalences and correlates of mental disorders. WHO
International Consortium in Psychiatric Epidemiology. (2000): *Bull World Health Organ*
78: 413–426.
2. Cardno AG, Owen MJ (2014): Genetic relationships between schizophrenia, bipolar
disorder, and schizoaffective disorder. *Schizophr Bull* 40: 504–515.
3. Hall J, Trent S, Thomas KL, O'Donovan MC, Owen MJ (2015): Genetic risk for
schizophrenia: convergence on synaptic pathways involved in plasticity. *Biol Psychiatry*
77: 52–58.
4. Forero DA, Herteleer L, De Zutter S, Norrback K-F, Nilsson L-G, Adolfsson R, *et al.*
(2016): A network of synaptic genes associated with schizophrenia and bipolar disorder.
Schizophr Res 172: 68–74.
5. Douaud G, Groves AR, Tamnes CK, Westlye LT, Duff EP, Engvig A, *et al.* (2014): A
common brain network links development, aging, and vulnerability to disease. *Proc Natl*
Acad Sci 111: 17648–17653.
6. Argyelan M, Ikuta T, DeRosse P, Braga RJ, Burdick KE, John M, *et al.* (2014):
Resting-state fMRI connectivity impairment in schizophrenia and bipolar disorder.
Schizophr Bull 40: 100–110.

7. Dager SR, Friedman SD, Parow A, Demopulos C, Stoll AL, Lyoo IK, *et al.* (2004):

Brain metabolic alterations in medication-free patients with bipolar disorder. *Arch Gen*

Psychiatry 61: 450–458.

8. Goh S, Dong Z, Zhang Y, DiMauro S, Peterson BS (2014): Brain imaging evidence that

mitochondrial dysfunction is a neurobiological subtype of Autism Spectrum Disorder.

JAMA Psychiatry 71: 665–671.

9. Greene AE, Todorova MT, Seyfried TN (2003): Perspectives on the metabolic

management of epilepsy through dietary reduction of glucose and elevation of ketone

bodies. *J Neurochem* 86: 529–537.

10. Halim ND, Lipska BK, Hyde TM, Deep-Soboslay A, Saylor EM, Herman M, *et al.*

(2008): Increased lactate levels and reduced pH in postmortem brains of schizophrenics:

medication confounds. *J Neurosci Methods* 169: 208–213.

11. Machado-Vieira R, Zanetti MV, Otaduy MC, De Sousa RT, Soeiro-de-Souza MG,

Costa AC, *et al.* (2017): Increased brain lactate during depressive episodes and reversal

effects by lithium monotherapy in drug-naive bipolar disorder: A 3T 1H-MRS study. *J Clin*

Psychopharmacol 37: 40–45.

12. Prabakaran S, Swatton JE, Ryan MM, Huffaker SJ, Huang JT-J, Griffin JL, *et al.*

(2004): Mitochondrial dysfunction in schizophrenia: evidence for compromised brain

metabolism and oxidative stress. *Mol Psychiatry* 9: 684–697, 643.

13. Rossignol DA, Frye RE (2012): Mitochondrial dysfunction in autism spectrum

disorders: a systematic review and meta-analysis. *Mol Psychiatry* 17: 290–314.

14. Rowland LM, Pradhan S, Korenic S, Wijtenburg SA, Hong LE, Edden RA, Barker PB

(2016): Elevated brain lactate in schizophrenia: a 7 T magnetic resonance spectroscopy

study. *Transl Psychiatry* 6: e967–e967.

15. Soeiro-de-Souza MG, Pastorello BF, Leite C da C, Henning A, Moreno RA, Garcia

Otaduy MC (2016): Dorsal anterior cingulate lactate and glutathione levels in euthymic

bipolar I disorder: 1H-MRS study. *Int J Neuropsychopharmacol* 19.

https://doi.org/10.1093/ijnp/pyw032

16. Dogan AE, Yuksel C, Du F, Chouinard V-A, Öngür D (2018): Brain lactate and pH in

schizophrenia and bipolar disorder: a systematic review of findings from magnetic

resonance studies. *Neuropsychopharmacology* 43: 1681–1690.

17. Pruett BS, Meador-Woodruff JH (2020): Evidence for altered energy metabolism,

increased lactate, and decreased pH in schizophrenia brain: A focused review and

meta-analysis of human postmortem and magnetic resonance spectroscopy studies.

Schizophr Res S092099642030459X.

18. Lehéricy S, Marjanska M, Mesrob L, Sarazin M, Kinkingnehun S (2007): Magnetic

resonance imaging of Alzheimer's disease. *Eur Radiol* 17: 347–362.

19. Liguori C, Stefani A, Sancesario G, Sancesario GM, Marciani MG, Pierantozzi M

(2015): CSF lactate levels, τ proteins, cognitive decline: a dynamic relationship in

Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 86: 655–659.

20. Liguori C, Chiaravalloti A, Sancesario G, Stefani A, Sancesario GM, Mercuri NB, *et al.*

(2016): Cerebrospinal fluid lactate levels and brain [18F]FDG PET hypometabolism within

the default mode network in Alzheimer's disease. *Eur J Nucl Med Mol Imaging* 43: 2040–

2049.

21. Lyros E, Ragoschke-Schumm A, Kostopoulos P, Sehr A, Backens M, Kalampokini S,

et al. (2019): Normal brain aging and Alzheimer's disease are associated with lower

cerebral pH: an in vivo histidine 1H-MR spectroscopy study. *Neurobiol Aging*.

https://doi.org/10.1016/j.neurobiolaging.2019.11.012

22. Mullins R, Reiter D, Kapogiannis D (2018): Magnetic resonance spectroscopy reveals

abnormalities of glucose metabolism in the Alzheimer's brain. *Ann Clin Transl Neurol* 5:

262–272.

23. Paasila PJ, Davies DS, Kril JJ, Goldsbury C, Sutherland GT (2019): The relationship

between the morphological subtypes of microglia and Alzheimer's disease neuropathology.

Brain Pathol 29: 726–740.

24. Youssef P, Chami B, Lim J, Middleton T, Sutherland GT, Witting PK (2018): Evidence

supporting oxidative stress in a moderately affected area of the brain in Alzheimer's disease.

Sci Rep 8: 1–14.

25. Li JZ, Vawter MP, Walsh DM, Tomita H, Evans SJ, Choudary PV, *et al.* (2004):

Systematic changes in gene expression in postmortem human brains associated with tissue

pH and terminal medical conditions. *Hum Mol Genet* 13: 609–616.

26. Tomita H, Vawter MP, Walsh DM, Evans SJ, Choudary PV, Li J, *et al.* (2004): Effect

of agonal and postmortem factors on gene expression profile: quality control in microarray

analyses of postmortem human brain. *Biol Psychiatry* 55: 346–352.

27. Vawter M, Tomita H, Meng F, Bolstad B, Li J, Evans S, *et al.* (2006):

Mitochondrial-related gene expression changes are sensitive to agonal-pH state:

implications for brain disorders. *Mol Psychiatry* 11: 615–679.

28. Kato T, Murashita J, Kamiya A, Shioiri T, Kato N, Inubushi T (1998): Decreased brain

intracellular pH measured by 31P-MRS in bipolar disorder: a confirmation in drug-free

patients and correlation with white matter hyperintensity. *Eur Arch Psychiatry Clin*

Neurosci 248: 301–306.

29. Hagihara H, Catts VS, Katayama Y, Shoji H, Takagi T, Huang FL, *et al.* (2018):

Decreased brain pH as a shared endophenotype of psychiatric disorders.

Neuropsychopharmacology 43: 459–468.

30. Duarte JMN, Do KQ, Gruetter R (2014): Longitudinal neurochemical modifications in

the aging mouse brain measured in vivo by 1H magnetic resonance spectroscopy.

Neurobiol Aging 35: 1660–1668.

31. Won H, Lee H-R, Gee HY, Mah W, Kim J-I, Lee J, *et al.* (2012): Autistic-like social

behaviour in *Shank2*-mutant mice improved by restoring NMDA receptor function. *Nature*

486: 261–265.

32. Raveau M, Shimohata A, Amano K, Miyamoto H, Yamakawa K (2018):

DYRK1A-haploinsufficiency in mice causes autistic-like features and febrile seizures.

Neurobiol Dis 110: 180–191.

33. Hori K, Nagai T, Shan W, Sakamoto A, Abe M, Yamazaki M, *et al.* (2015):

Heterozygous disruption of autism susceptibility candidate 2 causes impaired emotional

control and cognitive memory. *PLOS ONE* 10: e0145979.

34. Nakatani J, Tamada K, Hatanaka F, Ise S, Ohta H, Inoue K, *et al.* (2009): Abnormal

behavior in a chromosome- engineered mouse model for human 15q11-13 duplication seen

in autism. *Cell* 137: 1235–1246.

35. Beasley CL, Dwork AJ, Rosoklija G, Mann JJ, Mancevski B, Jakovski Z, *et al.* (2009):

Metabolic abnormalities in fronto-striatal-thalamic white matter tracts in schizophrenia.

Schizophr Res 109: 159–166.

36. Brady RO, Cooper A, Jensen JE, Tandon N, Cohen B, Renshaw P, *et al.* (2012): A

longitudinal pilot proton MRS investigation of the manic and euthymic states of bipolar

disorder. *Transl Psychiatry* 2: e160–e160.

37. Insel TR, Cuthbert BN (2015): Brain disorders? Precisely. *Science* 348: 499–500.

38. Chesler M (2003): Regulation and modulation of pH in the brain. *Physiol Rev* 83:

1183–1221.

39. Zauner A, Bullock R, Di X, Young HF (1995): Brain oxygen, CO_2, pH, and

temperature monitoring: evaluation in the feline brain. *Neurosurgery* 37: 1168–1177.

40. Monoranu CM, Apfelbacher M, Grünblatt E, Puppe B, Alafuzoff I, Ferrer I, *et al.*

(2009): pH measurement as quality control on human post mortem brain tissue: a study of

the BrainNet Europe consortium. *Neuropathol Appl Neurobiol* 35: 329–337.

41. Preece P, Cairns NJ (2003): Quantifying mRNA in postmortem human brain: influence

of gender, age at death, postmortem interval, brain pH, agonal state and inter-lobe mRNA

variance. *Mol Brain Res* 118: 60–71.

42. Harrison PJ, Heath PR, Eastwood SL, Burnet PWJ, McDonald B, Pearson RCA (1995):

The relative importance of premortem acidosis and postmortem interval for human brain

gene expression studies: selective mRNA vulnerability and comparison with their encoded

proteins. *Neurosci Lett* 200: 151–154.

43. Forester BP, Berlow YA, Harper DG, Jensen JE, Lange N, Froimowitz MP, *et al.*

(2010): Age-related changes in brain energetics and phospholipid metabolism. *NMR*

Biomed 23: 242–250.

44. Weaver KE, Richards TL, Logsdon RG, McGough EL, Minoshima S, Aylward EH, *et*

al. (2015): Posterior cingulate lactate as a metabolic biomarker in amnestic mild cognitive

impairment. *BioMed Res Int* 2015. https://doi.org/10.1155/2015/610605

45. Newman LA, Korol DL, Gold PE (2011): Lactate produced by glycogenolysis in

astrocytes regulates memory processing. *PLOS ONE* 6: e28427.

46. Suzuki A, Stern SA, Bozdagi O, Huntley GW, Walker RH, Magistretti PJ, Alberini CM

(2011): Astrocyte-neuron lactate transport is required for long-term memory formation.

Cell 144: 810–823.

47. Descalzi G, Gao V, Steinman MQ, Suzuki A, Alberini CM (2019): Lactate from

astrocytes fuels learning-induced mRNA translation in excitatory and inhibitory neurons.

Commun Biol 2: 1–11.

48. Mangia S, Tkáč I, Gruetter R, Van de Moortele P-F, Maraviglia B, Uğurbil K (2007):

Sustained neuronal activation raises oxidative metabolism to a new steady-state level:

evidence from 1H NMR spectroscopy in the human visual cortex. *J Cereb Blood Flow*

Metab 27: 1055–1063.

49. Schaller B, Xin L, O'Brien K, Magill AW, Gruetter R (2014): Are glutamate and lactate

increases ubiquitous to physiological activation? A 1H functional MR spectroscopy study

during motor activation in human brain at 7Tesla. *NeuroImage* 93: 138–145.

50. Regenold WT, Phatak P, Marano CM, Sassan A, Conley RR, Kling MA (2009):

Elevated cerebrospinal fluid lactate concentrations in patients with bipolar disorder and

schizophrenia: implications for the mitochondrial dysfunction hypothesis. *Biol Psychiatry*

65: 489–494.

51. Stork C, Renshaw PF (2005): Mitochondrial dysfunction in bipolar disorder: evidence

from magnetic resonance spectroscopy research. *Mol Psychiatry* 10: 900–919.

52. Holper L, Ben-Shachar D, Mann JJ (2019): Multivariate meta-analyses of

mitochondrial complex I and IV in major depressive disorder, bipolar disorder,

schizophrenia, Alzheimer disease, and Parkinson disease. *Neuropsychopharmacology* 44:

837–849.

53. Manji H, Kato T, Di Prospero NA, Ness S, Beal MF, Krams M, Chen G (2012):

Impaired mitochondrial function in psychiatric disorders. *Nat Rev Neurosci* 13: 293–307.

54. Pei L, Wallace DC (2018): Mitochondrial etiology of neuropsychiatric disorders. *Biol*

Psychiatry 83: 722–730.

55. Millan MJ, Agid Y, Brüne M, Bullmore ET, Carter CS, Clayton NS, *et al.* (2012):

Cognitive dysfunction in psychiatric disorders: characteristics, causes and the quest for

improved therapy. *Nat Rev Drug Discov* 11: 141–168.

56. Andrés P (2003): Frontal cortex as the central executive of working memory: time to

revise our view. *Cortex* 39: 871–895.

57. Foucher JR, Vidailhet P, Chanraud S, Gounot D, Grucker D, Pins D, *et al.* (2005):

Functional integration in schizophrenia: too little or too much? Preliminary results on fMRI

data. *NeuroImage* 26. https://doi.org/10.1016/j.neuroimage.2005.01.042

58. Bisri T, Utomo BA, Fuadi I (2016): Exogenous lactate infusion improved

neurocognitive function of patients with mild traumatic brain injury. *Asian J Neurosurg* 11:

151–159.

59. Rice AC, Zsoldos R, Chen T, Wilson MS, Alessandri B, Hamm RJ, Ross Bullock M

(2002): Lactate administration attenuates cognitive deficits following traumatic brain injury.

Brain Res 928: 156–159.

60. Carrard A, Elsayed M, Margineanu M, Boury-Jamot B, Fragnière L, Meylan EM, *et al.*

(2016): Peripheral administration of lactate produces antidepressant-like effects. *Mol*

Psychiatry. https://doi.org/10.1038/mp.2016.179

61. Hagihara H, Shoji H, Otabi H, Toyoda A, Katoh K, Namihira M, Miyakawa T: Protein

lactylation induced by neural excitation. *Submitted to bioRxiv*

62. Marunaka Y, Yoshimoto K, Aoi W, Hosogi S, Ikegaya H (2014): Low pH of interstitial

fluid around hippocampus of the brain in diabetic OLETF rats. *Mol Cell Ther* 2.

https://doi.org/10.1186/2052-8426-2-6

63. Davidovic L, Navratil V, Bonaccorso CM, Catania MV, Bardoni B, Dumas M-E

(2011): A metabolomic and systems biology perspective on the brain of the Fragile X

syndrome mouse model. *Genome Res* 21: 2190–2202.

64. Lin DDM, Crawford TO, Barker PB (2003): Proton MR spectroscopy in the diagnostic

evaluation of suspected mitochondrial disease. *Am J Neuroradiol* 24: 33–41.

65. Mucke L, Masliah E, Yu G-Q, Mallory M, Rockenstein EM, Tatsuno G, *et al.* (2000):

High-level neuronal expression of Aβ1–42 in wild-type human amyloid protein precursor

transgenic mice: synaptotoxicity without plaque formation. *J Neurosci* 20: 4050–4058.

66. Mullan M, Crawford F, Axelman K, Houlden H, Lilius L, Winblad B, Lannfelt L

(1992): A pathogenic mutation for probable Alzheimer's disease in the APP gene at the

N-terminus of beta-amyloid. *Nat Genet* 1: 345–347.

67. Murrell J, Farlow M, Ghetti B, Benson MD (1991): A mutation in the amyloid

precursor protein associated with hereditary Alzheimer's disease. *Science* 254: 97–99.

68. Shibutani M, Horii T, Shoji H, Morita S, Kimura M, Terawaki N, *et al.* (2017): Arid1b

haploinsufficiency causes abnormal brain gene expression and autism-related behaviors in

mice. *Int J Mol Sci* 18: 1872.

69. D'Gama AM, Pochareddy S, Li M, Jamuar SS, Reiff RE, Lam A-TN, *et al.* (2015):

Targeted DNA sequencing from autism spectrum disorder brains implicates multiple

genetic mechanisms. *Neuron* 88: 910–917.

70. Fitzgerald TW, Gerety SS, Jones WD, van Kogelenberg M, King DA, McRae J, *et al.*

(2015): Large-scale discovery of novel genetic causes of developmental disorders. *Nature*

519: 223–228.

71. Kalscheuer VM, FitzPatrick D, Tommerup N, Bugge M, Niebuhr E, Neumann LM, *et*

al. (2007): Mutations in autism susceptibility candidate 2 (AUTS2) in patients with mental

retardation. *Hum Genet* 121: 501–509.

72. Bakkaloglu B, O'Roak BJ, Louvi A, Gupta AR, Abelson JF, Morgan TM, *et al.* (2008):

Molecular cytogenetic analysis and resequencing of contactin associated protein-like 2 in

autism spectrum disorders. *Am J Hum Genet* 82: 165–173.

73. Sultana R, Yu C-E, Yu J, Munson J, Chen D, Hua W, *et al.* (2002): Identification of a

novel gene on chromosome 7q11.2 interrupted by a translocation breakpoint in a pair of

autistic twins. *Genomics* 80: 129–134.

74. Beunders G, Voorhoeve E, Golzio C, Pardo LM, Rosenfeld JA, Talkowski ME, *et al.*

(2013): Exonic deletions in AUTS2 cause a syndromic form of intellectual disability and

suggest a critical role for the C terminus. *Am J Hum Genet* 92: 210–220.

75. Zhang B, Xu Y-H, Wei S-G, Zhang H-B, Fu D-K, Feng Z-F, *et al.* (2014): Association

study identifying a new susceptibility gene (AUTS2) for schizophrenia. *Int J Mol Sci* 15:

19406–19416.

76. Nakao A, Miki T, Shoji H, Nishi M, Takeshima H, Miyakawa T, Mori Y (2015):

Comprehensive behavioral analysis of voltage-gated calcium channel beta-anchoring and

-regulatory protein knockout mice. *Front Behav Neurosci* 9.

https://doi.org/10.3389/fnbeh.2015.00141

77. Isshiki M, Tanaka S, Kuriu T, Tabuchi K, Takumi T, Okabe S (2014): Enhanced

synapse remodelling as a common phenotype in mouse models of autism. *Nat Commun* 5:

4742.

78. McFarlane HG, Kusek GK, Yang M, Phoenix JL, Bolivar VJ, Crawley JN (2008):

Autism-like behavioral phenotypes in BTBR T+tf/J mice. *Genes Brain Behav* 7: 152–163.

79. Yamasaki N, Maekawa M, Kobayashi K, Kajii Y, Maeda J, Soma M, *et al.* (2008):

Alpha-CaMKII deficiency causes immature dentate gyrus, a novel candidate

endophenotype of psychiatric disorders. *Mol Brain* 1: 6.

80. Hagihara H, Horikawa T, Nakamura HK, Umemori J, Shoji H, Kamitani Y, Miyakawa

T (2016): Circadian gene circuitry predicts hyperactive behavior in a mood disorder mouse

model. *Cell Rep* 14: 2784–2796.

81. Hagihara H, Horikawa T, Irino Y, Nakamura HK, Umemori J, Shoji H, *et al.* (2019):

Peripheral blood metabolome predicts mood change-related activity in mouse model of

bipolar disorder. *Mol Brain* 12: 107.

82. Le-Niculescu H, Kurian SM, Yehyawi N, Dike C, Patel SD, Edenberg HJ, *et al.* (2009):

Identifying blood biomarkers for mood disorders using convergent functional genomics.

Mol Psychiatry 14: 156–174.

83. Ament SA, Szelinger S, Glusman G, Ashworth J, Hou L, Akula N, *et al.* (2015): Rare

variants in neuronal excitability genes influence risk for bipolar disorder. *Proc Natl Acad*

Sci 112: 3576–3581.

84. Li H, Zhou D-S, Chang H, Wang L, Liu W, Dai S-X, *et al.* (2019): Interactome

analyses implicated CAMK2A in the genetic predisposition and pharmacological

mechanism of bipolar disorder. *J Psychiatr Res* 115: 165–175.

85. Purcell SM, Moran JL, Fromer M, Ruderfer D, Solovieff N, Roussos P, *et al.* (2014): A

polygenic burden of rare disruptive mutations in schizophrenia. *Nature* advance online

publication. https://doi.org/10.1038/nature12975

86. Kaitsuka T, Li S-T, Nakamura K, Takao K, Miyakawa T, Matsushita M (2011):
Forebrain-specific constitutively active CaMKKα transgenic mice show deficits in
hippocampus-dependent long-term memory. *Neurobiol Learn Mem* 96: 238–247.
87. Goyal V, Zahra SA, Hennings C, Zimmer-Bensch G, Beetz C, Schmeer CW, *et al.*
(n.d.): A conditional knockout mouse model to study the specific role of cyclin D2 in brain
development, adult neurogenesis and human brain pathologies. *Soc Neurosci Annu Meet*
2019 227.19.
88. Urban R, Scherrer G, Goulding EH, Tecott LH, Basbaum AI (2011): Behavioral indices
of ongoing pain are largely unchanged in male mice with tissue or nerve injury-induced
mechanical hypersensitivity. *PAIN* 152: 990–1000.
89. Wang X, Guan S, Liu A, Yue J, Hu L, Zhang K, *et al.* (2019): Anxiolytic effects of
Formononetin in an inflammatory pain mouse model. *Mol Brain* 12: 36.
90. Katayama Y, Nishiyama M, Shoji H, Ohkawa Y, Kawamura A, Sato T, *et al.* (2016):
CHD8 haploinsufficiency results in autistic-like phenotypes in mice. *Nature* 537: 675–679.
91. O'Roak BJ, Vives L, Fu W, Egertson JD, Stanaway IB, Phelps IG, *et al.* (2012):
Multiplex targeted sequencing identifies recurrently mutated genes in autism spectrum
disorders. *Science* 338: 1619–1622.
92. O'Roak BJ, Vives L, Girirajan S, Karakoc E, Krumm N, Coe BP, *et al.* (2012):
Sporadic autism exomes reveal a highly interconnected protein network of de novo
mutations. *Nature* 485: 246–250.
93. Talkowski ME, Rosenfeld JA, Blumenthal I, Pillalamarri V, Chiang C, Heilbut A, *et al.*
(2012): Sequencing chromosomal abnormalities reveals neurodevelopmental loci that
confer risk across diagnostic Bboundaries. *Cell* 149: 525–537.
94. Neale BM, Kou Y, Liu L, Ma'ayan A, Samocha KE, Sabo A, *et al.* (2012): Patterns and
rates of exonic de novo mutations in autism spectrum disorders. *Nature* 485: 242–245.
95. Bernier R, Golzio C, Xiong B, Stessman HA, Coe BP, Penn O, *et al.* (2014): Disruptive
CHD8 mutations define a subtype of autism early in development. *Cell* 158: 263–276.
96. Iwata R, Ohi K, Kobayashi Y, Masuda A, Iwama M, Yasuda Y, *et al.* (2014): RacGAP
α2-chimaerin function in development adjusts cognitive ability in adulthood. *Cell Rep* 8:
1257–1264.

97. Vitaterna MH, King DP, Chang A-M, Kornhauser JM, Lowrey PL, McDonald JD, *et al.*

(1994): Mutagenesis and mapping of a mouse gene, clock, essential for circadian behavior.

Science 264: 719–725.

98. Roybal K, Theobold D, Graham A, DiNieri JA, Russo SJ, Krishnan V, *et al.* (2007):

Mania-like behavior induced by disruption of CLOCK. *Proc Natl Acad Sci* 104: 6406–

6411.

99. Shi J, Wittke-Thompson JK, Badner JA, Hattori E, Potash JB, Willour VL, *et al.*

(2008): Clock genes may influence bipolar disorder susceptibility and dysfunctional

circadian rhythm. *Am J Med Genet B Neuropsychiatr Genet* 147B: 1047–1055.

100. Soria V, Martínez-Amorós È, Escaramís G, Valero J, Pérez-Egea R, García C, *et al.*

(2010): Differential association of circadian genes with mood disorders: CRY1 and NPAS2

are associated with unipolar major depression and CLOCK and VIP with bipolar disorder.

Neuropsychopharmacology 35: 1279–1289.

101. Kishi T, Kitajima T, Ikeda M, Yamanouchi Y, Kinoshita Y, Kawashima K, *et al.*

(2009): Association study of clock gene (CLOCK) and schizophrenia and mood disorders

in the Japanese population. *Eur Arch Psychiatry Clin Neurosci* 259: 293.

102. Murray F, Smith DW, Hutson PH (2008): Chronic low dose corticosterone exposure

decreased hippocampal cell proliferation, volume and induced anxiety and depression like

behaviours in mice. *Eur J Pharmacol* 583: 115–127.

103. Zhao Y, Ma R, Shen J, Su H, Xing D, Du L (2008): A mouse model of depression

induced by repeated corticosterone injections. *Eur J Pharmacol* 581: 113–120.

104. Antonijevic IA, Steiger A (2003): Depression-like changes of the sleep-EEG during

high dose corticosteroid treatment in patients with multiple sclerosis.

Psychoneuroendocrinology 28: 780–795.

105. Brown ES, J. Woolston D, Frol A, Bobadilla L, Khan DA, Hanczyc M, *et al.* (2004):

Hippocampal volume, spectroscopy, cognition, and mood in patients receiving

corticosteroid therapy. *Biol Psychiatry* 55: 538–545.

106. Brown ES, Suppes T (1998): Mood symptoms during corticosteroid therapy: a review.

Harv Rev Psychiatry 5: 239–246.

107. Nakamura H, Yamashita N, Kimura A, Kimura Y, Hirano H, Makihara H, *et al.*
(2016): Comprehensive behavioral study and proteomic analyses of CRMP2-deficient mice.
Genes Cells 21: 1059–1079.

108. Yoshida H, Watanabe A, Ihara Y (1998): Collapsin response mediator protein-2 Is
associated with neurofibrillary tangles in Alzheimer's disease. *J Biol Chem* 273: 9761–
9768.

109. Nakata K, Ujike H, Sakai A, Takaki M, Imamura T, Tanaka Y, Kuroda S (2003): The
human dihydropyrimidinase-related protein 2 gene on chromosome 8p21 is associated with
paranoid-type schizophrenia. *Biol Psychiatry* 53: 571–576.

110. Nyuyki KD, Cluny NL, Swain MG, Sharkey KA, Pittman QJ (2018): Altered brain
excitability and increased anxiety in mice with experimental colitis: consideration of
hyperalgesia and sex differences. *Front Behav Neurosci* 12.
https://doi.org/10.3389/fnbeh.2018.00058

111. Shoji H, Toyama K, Takamiya Y, Wakana S, Gondo Y, Miyakawa T (2012):
Comprehensive behavioral analysis of ENU-induced Disc1-Q31L and -L100P mutant mice.
BMC Res Notes 5: 108.

112. St Clair D, Blackwood D, Muir W, Walker M, St Clair D, Muir W, *et al.* (1990):
Association within a family of a balanced autosomal translocation with major mental illness.
The Lancet 336: 13–16.

113. Millar JK, Wilson-Annan JC, Anderson S, Christie S, Taylor MS, Semple CAM, *et al.*
(2000): Disruption of two novel genes by a translocation co-segregating with schizophrenia.
Hum Mol Genet 9: 1415–1423.

114. Ekelund J, Hovatta I, Parker A, Paunio T, Varilo T, Martin R, *et al.* (2001):
Chromosome 1 loci in Finnish schizophrenia families. *Hum Mol Genet* 10: 1611–1617.

115. Krumm N, O'Roak BJ, Shendure J, Eichler EE (2014): A de novo convergence of
autism genetics and molecular neuroscience. *Trends Neurosci* 37: 95–105.

116. McRae JF, Clayton S, Fitzgerald TW, Kaplanis J, Prigmore E, Rajan D, *et al.* (2017):
Prevalence and architecture of de novo mutations in developmental disorders. *Nature* 542:
433–438.

117. Imoto Y, Segi-Nishida E, Suzuki H, Kobayashi K (2017): Rapid and stable changes in

maturation-related phenotypes of the adult hippocampal neurons by electroconvulsive

treatment. *Mol Brain* 10: 8.

118. Kobayashi K, Imoto Y, Yamamoto F, Kawasaki M, Ueno M, Segi-Nishida E, Suzuki

H (2016): Rapid and lasting enhancement of dopaminergic modulation at the hippocampal

mossy fiber synapse by electroconvulsive treatment. *J Neurophysiol* 117: 284–289.

119. Husain MM, Rush AJ, Fink M, Knapp R, Petrides G, Rummans T, *et al.* (2004):

Speed of response and remission in major depressive disorder with acute Electroconvulsive

therapy (ECT): A consortium for research in ECT (CORE) report. *J Clin Psychiatry* 65:

485–491.

120. Pagnin D, de Queiroz V, Pini S, Cassano GB (2004): Efficacy of ECT in depression: a

meta-analytic review. *J ECT* 20: 13–20.

121. Consorthium TD-BFX, Bakker CE, Verheij C, Willemsen R, Helm R van der,

Oerlemans F, *et al.* (1994): Fmr1 knockout mice: a model to study fragile X mental

retardation. *Cell* 78: 23–33.

122. Taga T, Yamaguchi Y, Kokubo Y, Hattori S, Takao K, Inazawa J, *et al.* (2011):

Establishment of a new mouse model of neurodevelopmental disorder. *J Pharmacol Sci*

115: 35–35.

123. Sudo G, Kagawa T, Kokubu Y, Inazawa J, Taga T (2016): Increase in GFAP-positive

astrocytes in histone demethylase GASC1/KDM4C/JMJD2C hypomorphic mutant mice.

Genes Cells 21: 218–225.

124. Kantojärvi K, Onkamo P, Vanhala R, Alen R, Hedman M, Sajantila A, *et al.* (2010):

Analysis of 9p24 and 11p12-13 regions in autism spectrum disorders: rs1340513 in the

JMJD2C gene is associated with ASDs in Finnish sample. *Psychiatr Genet* 20: 102–108.

125. Nishizono H, Darwish M, Endo TA, Uno K, Abe H, Yasuda R (2020): Glycine

receptor α4 subunit facilitates the early embryonic development in mice. *Reproduction* 159:

41.

126. Labonne JDJ, Graves TD, Shen Y, Jones JR, Kong I-K, Layman LC, Kim H-G

(2016): A microdeletion at Xq22.2 implicates a glycine receptor GLRA4 involved in

intellectual disability, behavioral problems and craniofacial anomalies. *BMC Neurol* 16:

132.

127. Belforte JE, Zsiros V, Sklar ER, Jiang Z, Yu G, Li Y, *et al.* (2010): Postnatal NMDA

receptor ablation in corticolimbic interneurons confers schizophrenia-like phenotypes. *Nat*

Neurosci 13: 76–83.

128. Begni S, Moraschi S, Bignotti S, Fumagalli F, Rillosi L, Perez J, Gennarelli M (2003):

Association between the G1001C polymorphism in the GRIN1 gene promoter region and

schizophrenia. *Biol Psychiatry* 53: 617–619.

129. Zhao X, Li H, Shi Y, Tang R, Chen W, Liu J, *et al.* (2006): Significant association

between the genetic variations in the 5′ end of the N-Methyl-D-Aspartate receptor subunit

gene GRIN1 and schizophrenia. *Biol Psychiatry* 59: 747–753.

130. Hayashida M, Miyaoka T, Tsuchie K, Yasuda H, Wake R, Nishida A, *et al.* (2009):

Hyperbilirubinemia-related behavioral and neuropathological changes in rats: A possible

schizophrenia animal model. *Prog Neuropsychopharmacol Biol Psychiatry* 33: 581–588.

131. Müller N, Schiller P, Ackenheil M (1991): Coincidence of schizophrenia and

hyperbilirubinemia. *Pharmacopsychiatry* 24: 225–228.

132. Takao K, Kobayashi K, Hagihara H, Ohira K, Shoji H, Hattori S, *et al.* (2013):

Deficiency of Schnurri-2, an MHC enhancer binding protein, induces mild chronic

inflammation in the brain and confers molecular, neuronal, and behavioral phenotypes

related to schizophrenia. *Neuropsychopharmacology* 38: 1409–1425.

133. Srivastava S, Engels H, Schanze I, Cremer K, Wieland T, Menzel M, *et al.* (2016):

Loss-of-function variants in *HIVEP2* are a cause of intellectual disability. *Eur J Hum Genet*

24: 556.

134. Steinfeld H, Cho MT, Retterer K, Person R, Schaefer GB, Danylchuk N, *et al.* (2016):

Mutations in HIVEP2 are associated with developmental delay, intellectual disability, and

dysmorphic features. *Neurogenetics* 17: 159–164.

135. Fujisawa H, Sugimura Y, Takagi H, Mizoguchi H, Takeuchi H, Izumida H, *et al.*

(2015): Chronic hyponatremia causes neurologic and psychologic impairments. *J Am Soc*

Nephrol ASN.2014121196.

136. Izumida H, Takagi H, Fujisawa H, Iwata N, Nakashima K, Takeuchi S, *et al.* (2017):

NMDA receptor antagonist prevents cell death in the hippocampal dentate gyrus induced

by hyponatremia accompanying adrenal insufficiency in rats. *Exp Neurol* 287, Part 1: 65–

74.

137. Kawakami T, Fujisawa H, Nakayama S, Yoshino Y, Hattori S, Seino Y, *et al.* (2020):

Vasopressin escape and memory impairment in a model of chronic syndrome of

inappropriate secretion of antidiuretic hormone in mice. *Endocr J.*

https://doi.org/10.1507/endocrj.EJ20-0289

138. Fan S-S, Lin L-F, Chen VC-H, Hsieh C-W, Hsiao H-P, McIntyre RS, *et al.* (2020):

Effects of lower past-year serum sodium and hyponatremia on depression symptoms and

cognitive impairments in patients with hemodialysis. *Ther Apher Dial* 24: 169–177.

139. Fujisawa C, Umegaki H, Sugimoto T, Samizo S, Huang CH, Fujisawa H, *et al.* (2021):

Mild hyponatremia is associated with low skeletal muscle mass, physical function

impairment, and depressive mood in the elderly. *BMC Geriatr* 21: 15.

140. Yamanishi K, Doe N, Mukai K, Ikubo K, Hashimoto T, Uwa N, *et al.* (2019):

Interleukin-18-deficient mice develop hippocampal abnormalities related to possible

depressive-like behaviors. *Neuroscience* 408: 147–160.

141. Yamanishi K, Hashimoto T, Miyauchi M, Mukai K, Ikubo K, Uwa N, *et al.* (2020):

Analysis of genes linked to depressive-like behaviors in interleukin-18-deficient mice:

Gene expression profiles in the brain. *Biomed Rep* 12: 3–10.

142. Kretowski A, Mironczuk K, Karpinska A, Bojaryn U, Kinalski M, Puchalski Z,

Kinalska I (2002): Interleukin-18 promoter polymorphisms in type 1 diabetes. *Diabetes* 51:

3347–3349.

143. Chatterjee M, Ganguly S, Srivastava M, Palit G (2011): Effect of 'chronic' versus

'acute' ketamine administration and its 'withdrawal' effect on behavioural alterations in

mice: Implications for experimental psychosis. *Behav Brain Res* 216: 247–254.

144. Lahti AC, Weiler MA, Tamara Michaelidis BA, Parwani A, Tamminga CA (2001):

Effects of ketamine in normal and schizophrenic volunteers. *Neuropsychopharmacology*

25: 455–467.

145. Huang M, Kwon S, Rajagopal L, He W, Meltzer HY (2018): 5-HT1A parital agonism

and 5-HT7 antagonism restore episodic memory in subchronic phencyclidine-treated mice:

role of brain glutamate, dopamine, acetylcholine and GABA. *Psychopharmacology (Berl)*

235: 2795–2808.

146. Meltzer HY, Cucchiaro J, Silva R, Ogasa M, Phillips D, Xu J, *et al.* (2011):

Lurasidone in the treatment of schizophrenia: a randomized, double-blind, placebo- and

olanzapine-controlled study. *Am J Psychiatry* 168: 957–967.

147. Ishibashi T, Horisawa T, Tokuda K, Ishiyama T, Ogasa M, Tagashira R, *et al.* (2010):

Pharmacological profile of lurasidone, a novel antipsychotic agent with potent

5-hydroxytryptamine 7 (5-HT7) and 5-HT1A receptor activity. *J Pharmacol Exp Ther* 334:

171–181.

148. Connor SA, Ammendrup-Johnsen I, Kishimoto Y, Karimi Tari P, Cvetkovska V,

Harada T, *et al.* (2017): Loss of synapse repressor MDGA1 enhances perisomatic inhibition,

confers resistance to network excitation, and impairs cognitive function. *Cell Rep* 21:

3637–3645.

149. Kähler AK, Djurovic S, Kulle B, Jönsson EG, Agartz I, Hall H, *et al.* (2008):

Association analysis of schizophrenia on 18 genes involved in neuronal migration:

MDGA1 as a new susceptibility gene. *Am J Med Genet B Neuropsychiatr Genet* 147B:

1089–1100.

150. Li J, Liu J, Feng G, Li T, Zhao Q, Li Y, *et al.* (2011): The MDGA1 gene confers risk

to schizophrenia and bipolar disorder. *Schizophr Res* 125: 194–200.

151. Hossain MR, Jamal M, Tanoue Y, Ojima D, Takahashi H, Kubota T, *et al.* (2020):

MDGA1-deficiency attenuates prepulse inhibition with alterations of dopamine and

serotonin metabolism: An ex vivo HPLC-ECD analysis. *Neurosci Lett* 716: 134677.

152. Connor SA, Ammendrup-Johnsen I, Chan AW, Kishimoto Y, Murayama C, Kurihara

N, *et al.* (2016): Altered cortical dynamics and cognitive function upon haploinsufficiency

of the autism-linked excitatory synaptic suppressor MDGA2. *Neuron* 91: 1052–1068.

153. Bucan M, Abrahams BS, Wang K, Glessner JT, Herman EI, Sonnenblick LI, *et al.*

(2009): Genome-wide analyses of exonic copy number variants in a family-based study

point to novel autism susceptibility genes. *PLOS Genet* 5: e1000536.

154. Pinto D, Pagnamenta AT, Klei L, Anney R, Merico D, Regan R, *et al.* (2010):

Functional impact of global rare copy number variation in autism spectrum disorders.

Nature 466: 368–372.

155. McClay JL, Adkins DE, Vunck SA, Batman AM, Vann RE, Clark SL, *et al.* (2013):

Large-scale neurochemical metabolomics analysis identifies multiple compounds

associated with methamphetamine exposure. *Metabolomics* 9: 392–402.

156. Glasner-Edwards S, Mooney LJ (2014): Methamphetamine psychosis: epidemiology

and management. *CNS Drugs* 28: 1115–1126.

157. Togashi K, Wakatsuki S, Furuno A, Tokunaga S, Nagai Y, Araki T (2013): Na+/H+

exchangers induce autophagy in neurons and inhibit polyglutamine-induced aggregate

formation. *PLOS ONE* 8: e81313.

158. Tabuchi K, Blundell J, Etherton MR, Hammer RE, Liu X, Powell CM, Südhof TC

(2007): A neuroligin-3 mutation implicated in autism increases inhibitory synaptic

transmission in mice. *Science* 318: 71–76.

159. Jamain S, Quach H, Betancur C, Råstam M, Colineaux C, Gillberg IC, *et al.* (2003):

Mutations of the X-linked genes encoding neuroligins NLGN3 and NLGN4 are associated

with autism. *Nat Genet* 34: 27–29.

160. Südhof TC (2008): Neuroligins and neurexins link synaptic function to cognitive

disease. *Nature* 455: 903–911.

161. van West D, Van Den Eede F, Del-Favero J, Souery D, Norrback K-F, Van Duijn C,

et al. (2006): Glucocorticoid receptor gene-based SNP analysis in patients with recurrent

major depression. *Neuropsychopharmacology* 31: 620–627.

162. Huang FL, Huang K-P (2012): Methylphenidate improves the behavioral and

cognitive deficits of neurogranin knockout mice. *Genes Brain Behav* 11: 794–805.

163. Huang FL, Huang K-P, Wu J, Boucheron C (2006): Environmental enrichment

enhances neurogranin expression and hippocampal learning and memory but fails to rescue

the impairments of neurogranin null mutant mice. *J Neurosci* 26: 6230–6237.

164. Pak JH, Huang FL, Li J, Balschun D, Reymann KG, Chiang C, *et al.* (2000):

Involvement of neurogranin in the modulation of calcium/calmodulin-dependent protein

kinase II, synaptic plasticity, and spatial learning: A study with knockout mice. *Proc Natl Acad Sci* 97: 11232–11237.

165. Stefansson H, Ophoff RA, Steinberg S, Andreassen OA, Cichon S, Rujescu D, *et al.* (2009): Common variants conferring risk of schizophrenia. *Nature* 460: 744–747.

166. Schizophrenia Working Group of the Psychiatric Genomics Consortium (2014): Biological insights from 108 schizophrenia-associated genetic loci. *Nature* 511: 421–427.

167. Hashimoto H, Shintani N, Tanaka K, Mori W, Hirose M, Matsuda T, *et al.* (2001): Altered psychomotor behaviors in mice lacking pituitary adenylate cyclase-activating polypeptide (PACAP). *Proc Natl Acad Sci* 98: 13355–13360.

168. Hashimoto R, Hashimoto H, Shintani N, Ohi K, Hori H, Saitoh O, *et al.* (2010): Possible association between the pituitary adenylate cyclase-activating polypeptide (PACAP) gene and major depressive disorder. *Neurosci Lett* 468: 300–302.

169. Hashimoto R, Hashimoto H, Shintani N, Chiba S, Hattori S, Okada T, *et al.* (2007): Pituitary adenylate cyclase-activating polypeptide is associated with schizophrenia. *Mol Psychiatry* 12: 1026–1032.

170. Bolton PF, Veltman MWM, Weisblatt E, Holmes JR, Thomas NS, Youings SA, *et al.* (2004): Chromosome 15q11-13 abnormalities and other medical conditions in individuals with autism spectrum disorders. *Psychiatr Genet* 14: 131–137.

171. Cook Jr EH, Scherer SW (2008): Copy-number variations associated with neuropsychiatric conditions. *Nature* 455: 919–923.

172. Dykens EM, Sutcliffe JS, Levitt P (2004): Autism and 15q11-q13 disorders: Behavioral, genetic, and pathophysiological issues. *Ment Retard Dev Disabil Res Rev* 10: 284–291.

173. Takumi T, Tamada K (2018): CNV biology in neurodevelopmental disorders. *Curr Opin Neurobiol* 48: 183–192.

174. Meltzer HY, Rajagopal L, Huang M, Oyamada Y, Kwon S, Horiguchi M (2013): Translating the N-methyl-d-aspartate receptor antagonist model of schizophrenia to treatments for cognitive impairment in schizophrenia. *Int J Neuropsychopharmacol* 16: 2181–2194.

175. Peterson RC, Stillman RC (1978): Phencyclidine: An overview. Phencyclidine Abuse:
An Appraisal (ed. by Petersen, R.C., Stillman, R.C.). *NIDA Res Monogr 21 US Gov Print*
Off Wash DC 1–17.
176. Kasahara T, Takata A, Kato TM, Kubota-Sakashita M, Sawada T, Kakita A, *et al.*
(2016): Depression-like episodes in mice harboring mtDNA deletions in paraventricular
thalamus. *Mol Psychiatry* 21: 39–48.
177. Kasahara T, Ishiwata M, Kakiuchi C, Fuke S, Iwata N, Ozaki N, *et al.* (2017):
Enrichment of deleterious variants of mitochondrial DNA polymerase gene (POLG1) in
bipolar disorder. *Psychiatry Clin Neurosci* 71: 518–529.
178. Zeng H, Chattarji S, Barbarosie M, Rondi-Reig L, Philpot BD, Miyakawa T, *et al.*
(2001): Forebrain-Specific Calcineurin Knockout Selectively Impairs Bidirectional
Synaptic Plasticity and Working/Episodic-like Memory. *Cell* 107: 617–629.
179. Miyakawa T, Leiter LM, Gerber DJ, Gainetdinov RR, Sotnikova TD, Zeng H, *et al.*
(2003): Conditional calcineurin knockout mice exhibit multiple abnormal behaviors related
to schizophrenia. *Proc Natl Acad Sci* 100: 8987–8992.
180. Gerber DJ, Hall D, Miyakawa T, Demars S, Gogos JA, Karayiorgou M, Tonegawa S
(2003): Evidence for association of schizophrenia with genetic variation in the 8p21.3 gene,
PPP3CC, encoding the calcineurin gamma subunit. *Proc Natl Acad Sci* 100: 8993–8998.
181. Asaoka N, Nishitani N, Kinoshita H, Nagai Y, Hatakama H, Nagayasu K, *et al.*
(2019): An adenosine A2A receptor antagonist improves multiple symptoms of repeated
quinpirole-induced psychosis. *eNeuro* 6. https://doi.org/10.1523/ENEURO.0366-18.2019
182. Stuchlik A, Radostová D, Hatalova H, Vales K, Nekovarova T, Koprivova J, *et al.*
(2016): Validity of quinpirole sensitization rat model of OCD: linking evidence from
animal and clinical studies. *Front Behav Neurosci* 10: 209.
183. Sakai K, Shoji H, Kohno T, Miyakawa T, Hattori M (2016): Mice that lack the
C-terminal region of Reelin exhibit behavioral abnormalities related to neuropsychiatric
disorders. *Sci Rep* 6: 28636.
184. Persico AM, D'Agruma L, Maiorano N, Totaro A, Militerni R, Bravaccio C, *et al.*
(2001): Reelin gene alleles and haplotypes as a factor predisposing to autistic disorder. *Mol*
Psychiatry 6: 150–159.

185. Serajee FJ, Zhong H, Mahbubul Huq AHM (2006): Association of Reelin gene

polymorphisms with autism. *Genomics* 87: 75–83.

186. Zhang H, Liu X, Zhang C, Mundo E, Macciardi F, Grayson DR, *et al.* (2002): Reelin

gene alleles and susceptibility to autism spectrum disorders. *Mol Psychiatry* 7: 1012–1017.

187. Goes FS, Willour VL, Zandi PP, Belmonte PL, MacKinnon DF, Mondimore FM, *et al.*

(2010): Sex-specific association of the reelin gene with bipolar disorder. *Am J Med Genet B*

Neuropsychiatr Genet 153B: 549–553.

188. Shifman S, Johannesson M, Bronstein M, Chen SX, Collier DA, Craddock NJ, *et al.*

(2008): Genome-wide association identifies a common variant in the Reelin gene that

increases the risk of schizophrenia only in women. *PLOS Genet* 4: e28.

189. Shoji H, Miyakawa T (2020): Differential effects of stress exposure via two types of

restraint apparatuses on behavior and plasma corticosterone level in inbred male

BALB/cAJcl mice. *Neuropsychopharmacol Rep* 40: 73–84.

190. Sellmeijer J, Mathis V, Hugel S, Li X-H, Song Q, Chen Q-Y, *et al.* (2018):

Hyperactivity of anterior cingulate cortex areas 24a/24b drives chronic pain-induced

anxiodepressive-like consequences. *J Neurosci* 38: 3102–3115.

191. Yalcin I, Megat S, Barthas F, Waltisperger E, Kremer M, Salvat E, Barrot M (2014):

The sciatic nerve cuffing model of neuropathic pain in mice. *J Vis Exp JoVE*.

https://doi.org/10.3791/51608

192. Tatsukawa T, Raveau M, Ogiwara I, Hattori S, Miyamoto H, Mazaki E, *et al.* (2019):

Scn2a haploinsufficient mice display a spectrum of phenotypes affecting anxiety,

sociability, memory flexibility and ampakine CX516 rescues their hyperactivity. *Mol*

Autism 10: 15.

193. Buxbaum JD, Daly MJ, Devlin B, Lehner T, Roeder K, State MW (2012): The autism

sequencing consortium: large-scale, high-throughput sequencing in autism spectrum

disorders. *Neuron* 76: 1052–1056.

194. Tavassoli T, Kolevzon A, Wang AT, Curchack-Lichtin J, Halpern D, Schwartz L, *et al.*

(2014): De novo SCN2A splice site mutation in a boy with Autism spectrum disorder. *BMC*

Med Genet 15: 35.

195. Allen AS, Berkovic SF, Cossette P, Delanty N, Dlugos D, Eichler EE, *et al.* (2013): De novo mutations in epileptic encephalopathies. *Nature* 501: 217–221.

196. Heron SE, Crossland KM, Andermann E, Phillips HA, Hall AJ, Bleasel A, *et al.* (2002): Sodium-channel defects in benign familial neonatal-infantile seizures. *The Lancet* 360: 851–852.

197. Sugawara T, Tsurubuchi Y, Agarwala KL, Ito M, Fukuma G, Mazaki-Miyazaki E, *et al.* (2001): A missense mutation of the Na+ channel αII subunit gene Nav1.2 in a patient with febrile and afebrile seizures causes channel dysfunction. *Proc Natl Acad Sci* 98: 6384–6389.

198. de Ligt J, Willemsen MH, van Bon BWM, Kleefstra T, Yntema HG, Kroes T, *et al.* (2012): Diagnostic exome sequencing in persons with severe intellectual disability. *N Engl J Med* 367: 1921–1929.

199. Rauch A, Wieczorek D, Graf E, Wieland T, Endele S, Schwarzmayr T, *et al.* (2012): Range of genetic mutations associated with severe non-syndromic sporadic intellectual disability: an exome sequencing study. *The Lancet* 380: 1674–1682.

200. Tanaka M, Sato A, Kasai S, Hagino Y, Kotajima-Murakami H, Kashii H, *et al.* (2018): Brain hyperserotonemia causes autism-relevant social deficits in mice. *Mol Autism* 9: 60.

201. Bacchelli E, Maestrini E (2006): Autism spectrum disorders: Molecular genetic advances. *Am J Med Genet C Semin Med Genet* 142C: 13–23.

202. Wiggins JL, Swartz JR, Martin DM, Lord C, Monk CS (2014): Serotonin transporter genotype impacts amygdala habituation in youth with autism spectrum disorders. *Soc Cogn Affect Neurosci* 9: 832–838.

203. Peça J, Feliciano C, Ting JT, Wang W, Wells MF, Venkatraman TN, *et al.* (2011): Shank3 mutant mice display autistic-like behaviours and striatal dysfunction. *Nature* 472: 437–442.

204. Durand CM, Betancur C, Boeckers TM, Bockmann J, Chaste P, Fauchereau F, *et al.* (2007): Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism spectrum disorders. *Nat Genet* 39: 25–27.

205. Gauthier J, Spiegelman D, Piton A, Lafrenière RG, Laurent S, St-Onge J, *et al.* (2009):

Novel de novo SHANK3 mutation in autistic patients. *Am J Med Genet B Neuropsychiatr*

Genet 150B: 421–424.

206. Moessner R, Marshall CR, Sutcliffe JS, Skaug J, Pinto D, Vincent J, *et al.* (2007):

Contribution of SHANK3 mutations to autism spectrum disorder. *Am J Hum Genet* 81:

1289–1297.

207. Barr CL, Feng Y, Wigg K, Bloom S, Roberts W, Malone M, *et al.* (2000):

Identification of DNA variants in the SNAP-25 gene and linkage study of these

polymorphisms and attention-deficit hyperactivity disorder. *Mol Psychiatry* 5: 405–409.

208. Brophy K, Hawi Z, Kirley A, Fitzgerald M, Gill M (2002): Synaptosomal-associated

protein 25 (SNAP-25) and attention deficit hyperactivity disorder (ADHD): evidence of

linkage and association in the Irish population. *Mol Psychiatry* 7: 913–917.

209. Mill J, Curran S, Kent L, Gould A, Huckett L, Richards S, *et al.* (2002): Association

study of a SNAP-25 microsatellite and attention deficit hyperactivity disorder. *Am J Med*

Genet 114: 269–271.

210. Kustanovich V, Merriman B, McGough J, McCracken JT, Smalley SL, Nelson SF

(2003): Biased paternal transmission of SNAP-25 risk alleles in attention-deficit

hyperactivity disorder. *Mol Psychiatry* 8: 309–315.

211. Mill J, Richards S, Knight J, Curran S, Taylor E, Asherson P (2004): Haplotype

analysis of SNAP-25 suggests a role in the aetiology of ADHD. *Mol Psychiatry* 9: 801–

810.

212. Feng Y, Crosbie J, Wigg K, Pathare T, Ickowicz A, Schachar R, *et al.* (2005): The

SNAP25 gene as a susceptibility gene contributing to attention-deficit hyperactivity

disorder. *Mol Psychiatry* 10: 998–1005.

213. Hamdan FF, Myers CT, Cossette P, Lemay P, Spiegelman D, Laporte AD, *et al.*

(2017): High rate of recurrent de novo mutations in developmental and epileptic

encephalopathies. *Am J Hum Genet* 101: 664–685.

214. Heyne HO, Singh T, Stamberger H, Abou Jamra R, Caglayan H, Craiu D, *et al.*

(2018): De novo variants in neurodevelopmental disorders with epilepsy. *Nat Genet* 50:

1048–1053.

215. Ayalew M, Le-Niculescu H, Levey DF, Jain N, Changala B, Patel SD, *et al.* (2012):

Convergent functional genomics of schizophrenia: from comprehensive understanding to

genetic risk prediction. *Mol Psychiatry* 17: 887–905.

216. Houenou J, Boisgontier J, Henrion A, d'Albis M-A, Dumaine A, Linke J, *et al.*

(2017): A multilevel functional study of a SNAP25 at-risk variant for bipolar disorder and

schizophrenia. *J Neurosci* 37: 10389–10397.

217. Kollack-Walker, Don, Watson, Akil (1999): Differential Expression of c- *fos* mRNA

Within Neurocircuits of Male Hamsters Exposed to Acute or Chronic Defeat. *J*

Neuroendocrinol 11: 547–559.

218. Martinez M, Phillips PJ, Herbert J (1998): Adaptation in patterns of c-fos expression

in the brain associated with exposure to either single or repeated social stress in male rats.

Eur J Neurosci 10: 20–33.

219. Golden SA, Covington HE, Berton O, Russo SJ (2011): A standardized protocol for

repeated social defeat stress in mice. *Nat Protoc* 6: 1183–1191.

220. Toyoda A (2017): Social defeat models in animal science: What we have learned from

rodent models. *Anim Sci J* 88: 944–952.

221. Furman BL (2015): Streptozotocin-induced diabetic models in mice and rats. *Curr*

Protoc Pharmacol 70: 5.47.1-5.47.20.

222. Lenzen S (2008): The mechanisms of alloxan- and streptozotocin-induced diabetes.

Diabetologia 51: 216–226.

223. Anderson RJ, Freedland KE, Clouse RE, Lustman PJ (2001): The prevalence of

comorbid depression in adults with diabetes: a meta-analysis. *Diabetes Care* 24: 1069–

1078.

224. Miyamoto H, Shimohata A, Abe M, Abe T, Mazaki E, Amano K, *et al.* (2017):

Potentiation of excitatory synaptic transmission ameliorates aggression in mice with Stxbp1

haploinsufficiency. *Hum Mol Genet* 26: 4961–4974.

225. Hoischen A, Krumm N, Eichler EE (2014): Prioritization of neurodevelopmental

disease genes by discovery of new mutations. *Nat Neurosci* 17: 764–772.

226. Carvill GL, Heavin SB, Yendle SC, McMahon JM, O'Roak BJ, Cook J, *et al.* (2013):

Targeted resequencing in epileptic encephalopathies identifies de novo mutations in CHD2

and SYNGAP1. *Nat Genet* 45: 825–830.

227. Saitsu H, Kato M, Mizuguchi T, Hamada K, Osaka H, Tohyama J, *et al.* (2008): De

novo mutations in the gene encoding STXBP1 (MUNC18-1) cause early infantile epileptic

encephalopathy. *Nat Genet* 40: 782–788.

228. Komiyama NH, Watabe AM, Carlisle HJ, Porter K, Charlesworth P, Monti J, *et al.*

(2002): SynGAP regulates ERK/MAPK signaling, synaptic plasticity, and learning in the

complex with postsynaptic density 95 and NMDA receptor. *J Neurosci* 22: 9721–9732.

229. Nakajima R, Takao K, Hattori S, Shoji H, Komiyama NH, Grant SGN, Miyakawa T

(2019): Comprehensive behavioral analysis of heterozygous Syngap1 knockout mice.

Neuropsychopharmacol Rep 39: 223–237.

230. Narita N, Kato M, Tazoe M, Miyazaki K, Narita M, Okado N (2002): Increased

monoamine concentration in the brain and blood of fetal thalidomide- and valproic acid–

exposed rat: Putative animal models for autism. *Pediatr Res* 52: 576–579.

231. Tsugiyama LE, Ida-Eto M, Ohkawara T, Noro Y, Narita M (2020): Altered neuronal

activity in the auditory brainstem following sound stimulation in thalidomide-induced

autism model rats. *Congenit Anom* 60: 82–86.

232. Strömland K, Nordin V, Miller M, Akerström B, Gillberg C (1994): Autism in

thalidomide embryopathy: a population study. *Dev Med Child Neurol* 36: 351–356.

233. Okuda-Ashitaka E, Kakuchi Y, Kakumoto H, Yamanishi S, Kamada H, Yoshidu T, *et*

al. (2020): Mechanical allodynia in mice with tenascin-X deficiency associated with

Ehlers-Danlos syndrome. *Sci Rep* 10: 6569.

234. Kawakami K, Matsumoto K (2011): Behavioral alterations in mice lacking the gene

for tenascin-X. *Biol Pharm Bull* 34: 590–593.

235. Burch GH, Gong Y, Liu W, Dettman RW, Curry CJ, Smith L, *et al.* (1997): Tenascin–

X deficiency is associated with Ehlers–Danlos syndrome. *Nat Genet* 17: 104–108.

236. Tochigi M, Zhang X, Ohashi J, Hibino H, Otowa T, Rogers M, *et al.* (2007):

Association study between the TNXB locus and schizophrenia in a Japanese population.

Am J Med Genet B Neuropsychiatr Genet 144B: 305–309.

237. Wang J, Sun S, Zhang L, Wang Z, Ye L, Liu L, *et al.* (2011): Further study of genetic
association between the TNXB locus and schizophrenia. *Psychiatr Genet* 21: 216.
238. Wei J, Hemmings GP (2004): TNXB locus may be a candidate gene predisposing to
schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* 125B: 43–49.
239. Shimada T, Sugiura H, Yamagata K (n.d.): Inhibition of Rheb improved abnormal
social behavior in astrocyte-specific Tsc1 knockout mice. *NEURO2019 42nd Annu Meet*
Jpn Neurosci Soc 62nd Annu Meet Jpn Soc Neurochem PB-126.
240. Crino PB, Nathanson KL, Henske EP (2006): The tuberous sclerosis complex. *N Engl*
J Med 355: 1345–1356.
241. Kotajima-Murakami H, Kobayashi T, Kashii H, Sato A, Hagino Y, Tanaka M, *et al.*
(2019): Effects of rapamycin on social interaction deficits and gene expression in mice
exposed to valproic acid in utero. *Mol Brain* 12: 3.
242. Christensen J, Grønborg TK, Sørensen MJ, Schendel D, Parner ET, Pedersen LH,
Vestergaard M (2013): Prenatal valproate exposure and risk of autism spectrum disorders
and childhood autism. *JAMA* 309: 1696.
243. Kanda Y (2013): Investigation of the freely available easy-to-use software "EZR" for
medical statistics. *Bone Marrow Transplant* 48: 452–458.

1223

Figure legends

Figure 1. Hierarchical clustering of 65 strains/conditions of animals regarding brain pH

and lactate levels. Effect size was calculated for each strain/condition and was used in this

analysis.

Figure 2. Highly significant negative correlations between brain pH and lactate levels.

Scatter plot showing correlations between pH and lactate levels of 1,239 animals. A

Z-score was calculated for each animal within the strain/condition and used in this study.

Figure 3. Poorer working memory predict higher brain lactate levels. (a) Prediction of

brain lactate levels from behavioral outcomes in the 24 mouse models related to

neuropsychiatric disorders. The scatter plot shows significant correlations between

predicted and actual lactate levels. (b) Feature preference for constructing the model to

predict brain lactate levels. (c) Scatter plot showing correlations between working

memory measures and actual brain lactate levels.

Supplementary Figure 1. Bar graphs showing the raw mean (± sem) of brain pH. Each plot

represents individual value. Asterisks indicate significant effects of the genotype/condition.

*$p < 0.05$, **$p < 0.01$; unpaired t-test, or one-way or two-way ANOVA followed by *post*

hoc Tukey's multiple comparison test. Detailed statistics are presented in Supplementary

Table 2. $^$The data of these mice have been reported previously (29).

Supplementary Figure 2. Bar graphs showing the raw mean (± sem) of brain lactate levels.

Each plot represents an individual value. Asterisks indicate significant effects of the

genotype/condition. *$p < 0.05$, **$p < 0.01$; unpaired t-test, or one-way or two-way ANOVA

followed by *post hoc* Tukey's multiple comparison test. Detailed statistics are presented in

Supplementary Table 2. [$]The data of these mice have been reported previously (29).

Supplementary Figure 3. Normal distribution of effect size values of 65 animal models (a,

pH: $D = 0.12$, $p = 0.30$; b, lactate: $D = 0.15$, $p = 0.093$)

Supplementary Figure 4. No significant correlations between the number of transitions in

the light/dark transition test (a), immobility in the forced swim test (b) and actual brain

lactate levels of 24 mouse strains used for prediction analysis

Supplementary Figure 5. Scatter plots showing correlations between age at sampling and

pH (a), and lactate levels (b) in wild type/control mice

Supplementary Figure 6. Dot plots showing pH (a) and lactate levels (b) of female and

male animals in 17 mixed gender strains/conditions. Bars indicate means

1256

Table 1. Animal models used in this study

Supplementary Table 1. Raw data of brain pH and lactate, and detailed information of the

animals (age, sex, and treatment methods)

Supplementary Table 2. Detailed statistics of pH and lactate measurements in 65 animal

models

Supplementary Table 3. Source of behavioral data used for prediction analysis

1263

1264   Figure 1

1267   Figure 2

1268

1269

1270   Figure 3