Title: Climate, latitude, and land cover predict flying insect biomass across a German malaise trap network

Authors: Ellen A.R. Welti[1]*, Petr Zajicek[1], Manfred Ayasse[2], Tim Bornholdt[3], Jörn Buse[4], Frank Dziock[5], Rolf A. Engelmann[6,7], Jana Englmeier[8], Martin Fellendorf[2], Marc I. Förschler[4], Mark Frenzel[9], Ute Fricke[10], Cristina Ganuza[10], Mathias Hippke[11], Günter Hoenselaar[12], Andrea Kaus-Thiel[13], Klaus Mandery[14], Andreas Marten[15], Michael T. Monaghan[16,17], Carsten Morkel[12], Jörg Müller[8], Stephanie Puffpaff[18], Sarah Redlich[10], Ronny Richter[6,7,19], Sandra Rojas Botero[20], Tobias Scharnweber[21], Gregor Scheiffarth[22], Paul Schmidt Yáñez[16], Rhena Schumann[23,24], Sebastian Seibold[25,26], Ingolf Steffan-Dewenter[10], Stefan Stoll[27], Cynthia Tobisch[28], Sönke Twietmeyer[29], Johannes Uhler[8], Juliane Vogt[20], Dirk Weis[30], Wolfgang W. Weisser[20], Martin Wilmking[21], Peter Haase[1,31]

Author affiliations:

[1] Department of River Ecology and Conservation, Senckenberg Research Institute and Natural History Museum Frankfurt, Gelnhausen 63571 Germany

[2] Institute of Evolutionary Ecology and Conservation Genomics, Ulm University, Ulm 89075 Germany

[3] Lower Oder Valley National Park, Schwedt/Oder, OT Criewen 16303 Germany

[4] Black Forest National Park, Freudenstadt 72250 Germany

[5] Animal Ecology, University of Applied Sciences HTW Dresden, Pillnitzer Platz 2, Dresden 01326 Germany

[6] German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig, Leipzig 04103 Germany

[7] Systematic Botany and Functional Biodiversity, Institute for Biology, Leipzig University, Berlin 04103 Germany

[8] Ecological Field Station, University of Würzburg, Rauhenebrach 96181 Germany

[9] Community Ecology, Helmholtz Centre for Environmental Research, Halle 06120 Germany

[10] Animal Ecology and Tropical Ecology, Biocenter, University of Würzburg, Würzburg 97074 Germany

[11] Biosphärenreservatsamt Schaalsee-Elbe, Zarrentin am Schaalsee 19246 Germany

[12] Kellerwald-Edersee National Park, Bad Wildungen 34537 Germany

[13] Hunsrück-Hochwald National Park, Birkenfeld 55765 Germany

[14] Institut für Biodiversitätsinformation e.V., Ebern 96106 Germany

[15] Harz National Park, Wernigerode 38855 Germany

[16] Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin 12587 Germany

[17] Institut für Biologie, Freie Universität Berlin, 14195 Germany

[18] Nationalparkamt Vorpommern, Born a. Darß 18375 Germany

[19] Geoinformatics and Remote Sensing, Institute for Geography, Leipzig University, Leipzig
04103 Germany

[20] Restoration Ecology Research Group, Department of Ecology and Ecosystem Management,
Technical University of Munich, Freising 85354 Germany

[21] Institute for Botany and Landscape Ecology, University Greifswald, Greifswald 17489
Germany

[22] Lower Saxon Wadden Sea National Park Authority, Wilhelmshaven 26382 Germany

[23] Biological Station Zingst, University of Rostock, Zingst 18374 Germany

[24] Mathematics and Natural Sciences, Institute of Biological Science, University of Rostock,
Rostock 18059 Germany

[25] Ecosystem Dynamics and Forest Management Research Group, Technical University of
Munich, Freising 85354 Germany

[26] Berchtesgaden National Park, Berchtesgaden 83471 Germany

[27] Environmental Planning and Technology, University of Applied Sciences Trier,
Environmental Campus Birkenfeld, Birkenfeld 55761 Germany

[28] Institute for Ecology and Landscape, Weihenstephan University of Applied Sciences, Freising
85354 Germany

[29] Eifel National Park, Schleiden-Gemünd 53937 Germany

[30] Biosphärenreservat Oberlausitzer Heide- und Teichlandschaft, Malschwitz OT Wartha 02694
Germany

[31] Faculty of Biology, University of Duisburg-Essen, Essen, Germany

*corresponding author, email: ellen.welti@senckenberg.de

ABSTRACT

We examine the effects of climate and land cover on flying insect biomass using an ecological

gradients approach and data from the first year of a new, ongoing, standardized German Malaise

Trap Program coordinated by the German Long-Term Ecological Research network. We

hypothesized that biomass would display a unimodal relationship with temperature, consistent

with thermal performance theory. We additionally predicted biomass would decrease with

precipitation due to reduced flying activity, and would decrease in areas with more heavily

human-modified land cover. Flying insect biomass was quantified from malaise traps at 84

locations across Germany throughout the 2019 growing season. We used an Akaike information

criterion approach to parse drivers of temperature, deviation in 2019 temperature from long-term

averages, precipitation, land cover, geographic coordinates, elevation, and sampling period. We

further examined how effects of temperature on insect biomass change across space by testing

for interactions between temperature and latitude. Flying insect biomass increased linearly with

monthly temperature across all samples. However, positive effects of temperature on flying

insect biomass declined with latitude, suggesting the warm 2019 summer temperatures in

southern Germany may have exceeded local insect optimums, and highlighting the spatial

variation in climate change-driven impacts on insect communities. Land cover explained less

variation in insect biomass, with the largest effect being lower biomass in forested sites. Large-

scale, standardized biodiversity monitoring like the German Malaise Trap Program are critical to

meet the challenge of disentangling the roles of multiple drivers on insect communities.

Keywords: ecological gradient, climate change, insect monitoring, observational, distributed,

malaise, pollinator, land use, thermal performance, LTER

INTRODUCTION

Amidst burgeoning evidence of widespread insect declines, there has been an increasing call for high quality, standardized, and large scale insect monitoring to improve estimates of trends, and identify drivers (Didham et al. 2020, Wagner 2020). Insects, and flying insects in particular, are hyper-diverse, and insect activity and life supports many other taxa both directly and indirectly. Insects pollinate of both the majority of wild plants and 75% of crop species (Losey and Vaughan 2006, Klein et al. 2007, Vanbergen and Insect Pollinators Initiative 2013, Reilly et al. 2020). Insect biomass is a key component of energy flows for many food webs (Stepanian et al. 2020). Biomass is a useful indicator of ecosystem function (Dangles et al. 2011, Barnes et al. 2016), and is representative of whole insect communities (Shortall et al. 2009, Hallmann et al. 2017). Climate change and anthropogenically-altered land cover are among the most commonly cited drivers of insect declines (Habel et al. 2019), but we have yet to understand their full impacts on insect biomass.

The effects of climate change are geographically pervasive (Wilson and Fox 2020) and may explain insect decline in natural areas (Rada et al. 2019, Janzen and Hallwachs 2019, Baranov et al. 2020, Welti et al. 2020a). Some insect taxa are currently benefiting from rising temperatures, which can increase local populations (Kaspari et al. 2019, Baker et al. 2021), diversity (Hofmann et al. 2018), and species' range sizes (Termaat et al. 2019). However, as temperatures continue to rise and increase more rapidly, temperature is expected to decrease insect productivity (Warren et al. 2018). Thermal performance theory captures this non-linearity in predicting a unimodal relationship between temperature and insect fitness, as measured by biomass or other performance indicators (Kingsolver and Huey 2008, Kühsel and Blüthgen 2015, Sinclair et al. 2016, Kaspari et al. 2019).

The predicted response of precipitation regimes to climate change varies with region, but

forecasts generally suggest increased frequency of both extreme events including both heavy

precipitation events and droughts (Myhre et al. 2019). Precipitation may directly and indirectly

affect flying insects. High precipitation may directly cause insect mortality, reducing population

sizes, but may also decrease the period of time insects are flying, reducing trap catch but not

necessarily reflecting true community biomass (Totland 1994). Indirect effects of precipitation

on flying insects mediated by plants (e.g. altering plant phenology or plant nutrition) are context-

dependent but increasing rainfall in average to wet climates is often detrimental (Lawson and

Rands 2019).

Changing land cover due to human activities is also a major threat to insects (Wagner

2020), with wide-ranging impacts from loss of resources and nesting locations at local scales, to

fragmented habitats at larger scales (Newbold et al. 2020). Heavily human modified landscapes

come with associated pressures, such as eutrophication and pesticide use with agricultural

intensification (Goulson et al. 2018, Carvalheiro et al. 2020), and light pollution from urban

environments (Owens et al. 2020). Urban land cover can result in declines of both insect

diversity (Fenoglio et al. 2020, Piano et al. 2020), and biomass (Macgregor et al. 2019,

Svenningsen et al. 2020).

Using data from the first year of monitoring (2019) of the German Malaise Trap

Program, we examine the effects of climate and surrounding land cover on flying insect biomass

across both the growing season (April-October) and a regional geographic gradient (ranging over

7° latitude, from the German Alps to the Baltic and North Sea). A recent study (Hallmann et al.

2017) reported flying insect biomass from 63 German locations over 27 years. However, not all

locations were sampled across years (the most sampling within an individual year comprised 23

locations), and the majority of sampling locations (58 out of 63) were clustered in central-west

Germany; thus the sites do not have representative coverage of Germany or represent an

extensive latitudinal gradient (coverage of 2° latitude). While our sampling does not yet have

multi-year coverage, the high number trap locations and broad spatial coverage allows us to

examine drivers of flying insect biomass using a macroecological gradients approach (Pianka

1966, Halbritter et al. 2013, Welti and Joern 2015, Peters et al. 2019).

Here we introduce the German Malaise Trap Program and provide a first examination

from this network of the drivers of variation in flying insect biomass. We hypothesize **(H1)** that

the effect of temperature on insect biomass will become less positive with increasing

temperatures, consistent with thermal performance theory. Spatially, we predict **(H2)** greater

increases in flying insect biomass with increasing temperature in colder sites and weaker to

negative effects of temperature on flying insect biomass at warmer sites. As elevation tends to

increase from northern to southern Germany and may affect temperature, we additionally test for

the effects of elevation on flying insect biomass. We hypothesize **(H3)** that flying insect biomass

will decline with increasing precipitation due to reduced flying activity. Finally, we predict **(H4)**

lower insect biomass in areas surrounded by more heavily human-modified land cover types

such as urban and agricultural areas.

MATERIALS AND METHODS

The German Malaise Trap Program

The German Malaise Trap Program currently comprises 31 LTER-D and National Natural

Landscape sites (biosphere reserves and national parks) (https://www.ufz.de/lter-

d/index.php?de=46285). The program was established in early 2019 with the aim of investigating

long-term trends in flying insect biomass and species composition using DNA metabarcoding. In

each site, one to six locations were selected and one malaise trap was installed per location. For

this study, we used the 2019 biomass data retrieved from 25 of the 31 sites; the remaining six

sites began sampling in 2020. To fill in spatial gaps, we included 8 sites in Bavaria from an

additional project but using the same malaise trap type and measurement methods. Overall, this

study includes 1039 samples from 84 malaise trap locations and 33 participating sites widely

distributed across Germany (Fig. 1). All traps were identically constructed with an opening area

of 1.16 m^2 on each of the two trap sides (Fig. S1). Traps predominantly ran from early April to

late October 2019 and were usually emptied every two weeks (14.03 days $\pm$ 0.06 SE; but ranging

7 - 29 days). Several samples were lost due to animal or wind damage. By sampling across all

times of day for the duration of the growing season, these data provide representative measures

of each location's flying insect community.

Lab procedures

166   Insect biomass was weighed while wet to preserve samples for future identification. Samples

167   were placed in a stainless steel sieve (0.8 mm mesh width) to remove excess alcohol, following

168   the weighing procedure in Hallmann et al. (2017), with one modification: instead of waiting until

169   filtered alcohol drops occurred >10 seconds apart, each sample was filtered for a standard five

170   minutes. Subsequently, insects were weighted to the nearest 0.01g.

Climate

Monthly means of maximum and minimum temperatures, and monthly cumulative precipitation

from 2019 were extracted from each sampling location using the Terraclimate dataset

(Abatzoglou et al. 2018). To compare 2019 with long-term climate averages, we extracted

monthly means of maximum and minimum temperature, and monthly cumulative precipitation

from 1960-2018 from each trap location using the CRU-TS 4.03 dataset (Harris et al. 2014)

downscaled with WorldClim 2.1 (Fick and Hijmans 2017). Data from both time periods (2019

and 1960-2018) were not available from either dataset alone. Both the Terraclimate and CRU-TS

4.03/ WorldClim 2.1 datasets have spatial resolutions of 2.5 arc minutes (~21 km^2) with our 84

trap locations occurring in 72 separate climate grid cells.

Maximum and minimum monthly temperatures in 2019 were generally higher than 1960-

2018 averages, especially during the hottest months (Supporting Information: Fig. S2a). While

one study suggested that minimum nightly temperatures are warming at a faster rate than

maximum temperatures, potentially limiting the ability of insects to recover from hot daytime

periods (Speights *et al.*, 2017), trap locations had slightly greater warming in their 2019

maximum monthly temperatures (2.062°C ± 0.047 SE), than minimum monthly temperatures

(1.938°C ± 0.048 SE) in comparison to 1960-2018 averages. As mean maximum and minimum

monthly temperatures were highly correlated ($F_{(1,512)}$=4442, R^2=0.9, P<0.001), we used only

mean maximum temperatures in our analyses. Cumulative annual precipitation was lower in

2019 (784 mm ± 32 SE) compared to the long-term 1960-2018 average (842 mm ± 32 SE).

There was no consistent difference between monthly cumulative precipitation in 2019 and the

long-term average, but the mid-summer months tended to be drier in 2019 (Supporting

Information: Fig. S2b). Both 2019 mean monthly maximum temperatures and precipitation

across our trap locations tended to decline with increasing latitude (Supporting Information: Fig.

S3).

Land cover

Land-cover categories in a 1-km buffer around each trap location were extracted using the

CORINE Land cover data from 2018 (European Union, Copernicus Land Monitoring Service

2018). A 1-km buffer was selected based on previous work suggesting land cover effects on

insect communities declined at larger scales (Seibold et al. 2019). Extracted land cover data was

comprised of 30 CORINE land cover categories, which we pooled into eight categories: urban

(7.5% of surrounding land cover), intensive agriculture (2.3%), non-irrigated agriculture

(15.9%), pasture/orchard (12.7%), forest (44.7%), grassland/shrubland (12.1%), freshwater

(3.9%), and saltwater (0.9%). For data visualization, trap locations were additionally categorized

by the dominant land cover within the 1-km buffer: urban (n=6), non-irrigated agriculture

(n=16), pasture/orchard (n=6), forest (n=44), grassland/shrubland (n=9), freshwater (n=2), and

saltwater (n=1).

Elevation

Elevation (m above sea level) was extracted using the Digital Terrain Model with 200-m grid

widths (DGM200) from the German Federal Agency for Cartography and Geodesy (GeoBasis-

DE / BKG 2013). Locations varied from elevations of 0 m on a barrier island in northeast

Germany, to 1413 m in the German Alps. Elevation across the 84 locations is negatively

correlated with latitude (Supporting Information: Fig. S4).

All GIS data extraction was conducted in QGIS ver. 3.14 (QGIS.org 2020).

AIC$_c$ analysis

To identify drivers of insect biomass, we used an Akaike Information Criterion corrected for

small sample sizes (AIC$_c$) modelling framework (Burnham and Anderson 2003); first building an

a priori full model, and then comparing all possible reduced models using the *dredge* function in

the R package MuMIn (Bartoń 2016). Mixed models were fit using the R package lme4 (Bates et

al. 2015). All analyses were conducted in R ver. 4.0.3 (R Core Team 2020). To reduce variance

inflation due to land cover categories being percentages, we sequentially removed land cover

categories from the model starting with the least common until the variance inflation factor was

<10; this removed the land cover types freshwater, intensive agriculture, and saltwater. Initial

analyses substituting the Land Use Index (LUI; Büttner 2014) for land cover percentages

resulted in no top models containing LUI; thus we opted to use individual land cover categories.

The 2nd degree polynomial of the sampling period was included in the model to capture the

hump-shaped pattern of flying insect biomass across the growing season. Sampling period refers

to the half-month period most overlapping the days of sampling, and is numerical (e.g. first half

of April = sampling period 1, second half of April = sampling period 2). Monthly temperature

and precipitation predictors correspond to the month in which the majority of sampling days

occurred. Monthly temperature was included as a 2nd degree polynomial to test our prediction of

a unimodal relationship. One approach to examine our hypothesis that temperature effects varied

spatially would be to include the interaction between temperature variables and latitude;

however, we did not include this interaction as it caused variance inflation. Precipitation and

elevation were scaled by dividing by 100. We did not include 2019 precipitation minus the long-

term average in our model because it caused inflated variance and 2019 precipitation was not

consistently different from the long-term average (Fig. S2b).

The full model contained the response variable of log10(x+1) transformed mg/day

biomass, the fixed predictors of the 2nd degree polynomial of 2019 average monthly maximum

temperature (poly(tmax,2)), the difference between 2019 average monthly maximum

temperature and the long-term average temperature (Δtemp), monthly cumulative precipitation

(precip), latitude, longitude, elevation, % cover of the five most dominant land cover categories

within a 1-km radius, a 2^{nd} degree polynomial of sampling period to account for seasonality

(poly(period,2)), and a random effect of trap identity (trap) to account for repeated observations.

The form of the mixed model was:

log10(mg/day+1) ~ poly(tmax,2) + Δtemp + precip + latitude + longitude + elevation + %forest

+ %grass/shrubland + %pasture/orchard + %non-irrigated crop + %urban + poly(period,2) +

(1|trap)

Temperature effects across latitudes

While variance inflation prohibited a temperature by latitude interaction term in the mixed

model, we wished to further examine our hypotheses that the effects of temperature on flying

insect biomass would decrease in the lower latitudes which had the highest temperatures in 2019.

For this analysis, we reduced the dataset to one average biomass value (in mg/day) for each

location and month. Calculating monthly values reduces complexity due to variation in when

samples are collected across locations, and was the first step towards calculating one value of

biomass across the growing season for each sampling location. We estimated the daily biomass

collected within each month, assuming traps caught the same amount of biomass each day within

a sample. We pooled samples within the same month and, for samples spanning more than one

month, we allocated the proportion of biomass from the sample to each month based on the

number of sampling days occurring within the month. With these assumptions, the average

biomass Bij (mg/day) of location i in month j in location is a weighted average of the n samples

occurring in the month according to the following formula:

Eq. 1

269
$$B_{i,j} = \frac{\sum_{k=1}^{n}(b_{ijk} \times D_{k,j} \div D_k)}{\sum_{k=1}^{n} D_{k,j}}$$

Where b$_{ijk}$ = the total biomass (mg) at location i occurring at least partially in month j for a

sample k, n= the total number of samples occurring at least partially in month j for location i, D$_{k,j}$

= the number of sampling days occurring in month j for a given sample k, D$_k$ = the total number

of sampling days for a given sample k

To examine how the effect of temperature changed over latitude, we averaged B$_{i,j}$ across

the seven sampling months (Apr. - Oct.) to calculate one monthly average of mg/day per

location. We used only the 48 locations which conducted sampling in all seven sampling months.

We calculated average monthly maximum temperature across the same period (Apr. - Oct.

2019), henceforth "growing season temperature". Then we examined the linear relationship

between monthly averages of mg/day and the interaction between growing season temperature

and latitude. We visualized the interaction between temperature and latitude using the R package

"effects" (Fox and Weisberg 2019).

Dominant land cover categories

To visualize changes in flying insect biomass in the dominant land-cover categories and across

the growing season, we plotted flying insect biomass/day over median sampling day (numerical

day of the year halfway between trap start and end day) for locations corresponding to each

dominant land cover. We used Welch's t-tests, which do not assume equal variance, to identify

significant differences between $\log 10(x + 1)$ transformed $B_{i,j}$ of sites dominated by forest,

grassland, pasture/orchard, non-irrigated agriculture, and urban landscapes, and averages of all

samples within each month. The locations with surrounding land cover dominated by saltwater

(n=1) and freshwater (n=2) were excluded due to low replication.

RESULTS

Mean flying insect biomass was 2,329 mg/day $\pm$ 79 SE across all sampling periods and trap

locations and varied from 0 to 17,543 mg/day. On average across locations, biomass increased

from 734 mg/day $\pm$ 98 SE in early April, to a peak of 5,356 mg/day $\pm$ 401 SE in late June,

declining to 568 mg/day $\pm$ 111 SE in late October. Flying insect biomass was strongly predicted

by temperature, latitude, precipitation, percent forest cover, and sampling period (Table 1). AIC_c

model comparison selected three top models which always included the 2^{nd} degree polynomial of

sampling period, the 2^{nd} degree polynomial of average monthly maximum 2019 temperature, and

negative effects of Δ temperature from the long-term average, latitude, monthly precipitation,

and percent forest cover (Supporting Information: Table S1). The second and third top models

additionally included a negative effect of percent urban cover and a positive effect of percent

pasture/orchard cover (Supporting Information: Table S2). However, these models did not

greatly improve R^2 or log-likelihood; thus these two land cover types provided only weak

additional explanatory power (Arnold 2010). Elevation did not appear in any top models

(Supporting Information: Table S1). The top model explained 51% of the variance in flying

insect biomass without location information (marginal R^2) and 74% of flying insect biomass was

accounted for when including the random effect of location identity (conditional R^2; Supporting

Information: Table S1).

Climate and latitude

Flying insect biomass increased with mean monthly maximum temperature within 2019

sampling (Table 1, Fig. 2a), and declined with increasing latitude (Table 1, Fig. 2b) and monthly

cumulative precipitation (Table 1, Fig. 2c). While 2019 mean monthly maximum temperature

was included in all top models, only the first order polynomial term was significant, suggesting

the effect was linear (Table 1; Supporting Information: Table S2). When included in the same

model as 2019 monthly maximum temperature, the Δ temperature had a negative effect on flying

insect biomass (Table 1; Fig. 2d). Across the 48 sites in which sampling occurred in all seven

months (Apr. - Oct.), an interaction between growing season temperature and latitude was

predictive of flying insect biomass (Supporting Information: Table S3; $F_{3,44} = 4.3$, $R^2 = 0.23$, *P* =

0.01). The effect of growing season temperature on flying insect biomass increased from a

negative temperature effect at lower latitudes to a positive effect at higher latitudes (Fig. 3).

Land cover

Flying insect biomass declined with increasing percent forest cover and tended to decrease with

percent urban cover and increase with percent pasture/orchard (Supporting Information: Table

S2). Categorizing trap locations by dominant land cover showed grassland/shrubland locations to

generally have the highest biomass in the early-mid growing season (Fig. 4c), and a moderate

increase in biomass in non-irrigated cropland across the growing season (Fig. 4e). However,

neither percent grassland/shrubland nor percent non-irrigated cropland explained variation in

biomass after accounting for seasonality and temperature (Supporting Information: Table S1).

Urban-dominated sites tended to have slightly higher catches of biomass within months (Fig. 4f),

but this may be due to the six urban-dominated sites all being located in southern Germany (Fig.

1). When included in the full model with temperature variables, flying insect biomass was

weakly negatively correlated with percent urban cover (Supporting Information: Table S2a).

DISCUSSION

In this first outcome of the German Malaise Trap Program, we found flying insects to be

extremely sensitive to temperature, with both a positive effect of monthly temperature and a

negative effect of 2019 temperatures exceeding long-term averages predicting daily sampled

biomass. Across all sampling, flying insect biomass increased linearly with temperature in

contrast to the unimodal relationship predicted by our first hypothesis (H1). However, in support

of H2, the relationship between temperature and flying insect biomass shifted from positive in

colder locations in northern Germany to negative in warmer locations in southern Germany.

Additionally, precipitation had a negative relationship with flying insect biomass, as predicted by

H3. Finally, we found mixed support for our H4 prediction that more heavily-human impacted

land use types would have the lowest biomass, with locations surrounded by pasture/orchard,

grassland, and non-irrigated agriculture tending to have higher insect biomass than those

surrounded by forests or urban areas.

Climate

Temperature effects on ectothermic insects can manifest directly on individuals' metabolism,

growth, reproduction, immunity, survival, behavior, and dispersal (Sinclair et al. 2016, Harvey et

al. 2020), and through effects on other trophic levels such as through altering plant biomass,

plant quality, or active time-periods of natural enemies (Laws and Joern 2013, Kwon et al. 2019,

Thakur 2020). Our results suggest an overall positive linear effect of temperature on flying insect

biomass across Germany. The lack of a unimodal relationship between temperature and insect

biomass may be a result of the coarse taxonomic (flying insects caught in malaise traps) and

temporal (two week) resolution of our samples in comparison to other studies (e.g. Kühsel and

Blüthgen 2015). However, we detected a negative effect of the Δ temperature (2019 temperatures

minus the colder long-term average). Additionally, the slope of the relationship between

temperature and flying insect biomass declined towards the lower latitudes, which tended to be

the warmest locations in 2019. Hallmann *et al.* (2017) found a positive effect of temperature on

insect biomass; however, biomass loss over time was greatest in the middle of the growing

season, when temperatures are highest. A decelerating benefit of temperature in regions with

greater increases in temperature is consistent with a study of two surveys of ant communities

across North America that were conducted 20 years apart, which found that sites with the largest

increases in average temperature had the largest declines in colony density (Kaspari et al. 2019).

We predict future monitoring will detect increasingly negative effects of temperature due to

ongoing climate warming, as temperature begins to exceed species' optimum temperature

ranges.

Climate change forecasts for Germany suggest slight increases in cumulative annual

precipitation but also shifts in the timing of rainfall with longer dry periods, especially in

summer (Bender et al. 2017). The 2019 growing season matched this prediction with June, July

and August being much drier than the long-term average and with the wettest month of the study

period being October. We detected an overall negative effect of precipitation on flying insect

biomass. As our study covered only one growing season, our result likely reflects direct insect

mortality from heavy rain or reduced flight activity. Insects can detect changes in barometric

pressure and will stop flying if they sense storms approaching (Pellegrino et al. 2013).

Land cover

With ~75% of global land significantly altered by human activities (IPBES 2019), land cover

change and land use intensification is a major contributor to insect decline (Potts et al. 2010,

Winfree et al. 2011, Díaz et al. 2019). However, we detected only weak effects of urban and

agricultural land cover on flying insect biomass. This may in part be due to a minority of heavily

human-impacted land cover surrounding our sites, especially with regards to intensive

agriculture. While percentage urban area had only a slight negative effect on flying insect

biomass, urban environments may affect insect communities in other ways, such as through

altering species composition (Theodorou et al. 2020). Higher temperatures in urban areas

compared to rural surroundings may partly compensate for lower habitat quality and explain the

above average biomass in spring and late summer/autumn in the urban areas investigated in our

study, while also making insects in urban areas more vulnerable to future warming.

Even within land-cover types, large variability exists in habitat quality. This is

particularly true of urban areas and agricultural land, which range from large paved urban

expanses and areas with intensive pesticide use to urban gardens and low intensity organic farms,

offering a wide range of habitat quality for insects (Bengtsson et al. 2005, Hausmann et al.

2020). While moderately impacted by human activity, non-irrigated agricultural areas, pasture

land, and orchards in this study tended to support higher biomass, suggesting these land use

types may provide suitable habitats for Germany's flying insects. Alternatively, fertilization and

the prevalence of monoculture on conventional farms may increase insect biomass through

alleviating nutrient limitation and providing high concentrations of host plants, while not

benefiting insect biodiversity (Root 1973, Haddad et al. 2000).

We detected reduced insect biomass in trap locations surrounded predominantly by

forests. This result is in contrast to previous work, albeit using different sampling methods,

showing high Lepidoptera biomass coming to light traps in forests compared to other land use

types in the UK (Macgregor et al. 2019), and higher flying insect biomass with increasing forest

cover in a study using car-mounted nets in Denmark (Svenningsen et al. 2020). Our malaise-

trapping methods are more directed toward Hymenoptera and Diptera, and towards the lower

canopy, potentially contributing to this discrepancy. Forests may also provide fewer floral

resources than open fields (Jachuła et al. 2017). Alternatively, the vegetation structure in forests

may limit insect movement through the landscape, reducing trap catch in comparison to more

open systems like grasslands (Cranmer et al. 2012).

Comparison with Hallmann et al. 2017

Average insect biomass captured in malaise traps of Hallmann *et al.* (2017) varied from 9,192

mg/day in 1989 to 2,531 mg/day in 2016 (mean of within-month sample means for May-Sept;

April excluded as no April 1989 sampling was conducted). In comparison, our traps collected a

monthly average of 2,404 mg/day in May-Sept. However, the malaise traps used in Hallmann *et*

al. (2017) were ~51% larger (Hallmann *et al.* [2017] trap opening dimensions per side: 1.79 m^2,

traps used in this study: 1.16 m^2). If trap opening has an appreciable positive effect on trap catch,

this suggests higher trap catch in this study compared to the last sampling year in Hallmann *et al.*

(2017). This discrepancy is most likely due to the difference in sampling locations between the

studies, but we cannot rule out an increase in biomass of flying insects in Germany.

Caveats

Insect biomass is a common currency ecosystem-level measure of insect productivity and is an

index of energy availability for higher trophic levels. Nonetheless, we cannot differentiate

variation in abundance, body size, species diversity, or dominance from biomass data alone.

High temperatures may reduce average insect body size either by causing smaller adult body

sizes within species (Atkinson 1994, Klockmann et al. 2017, Polidori et al. 2020) or by favoring

smaller species (Bergmann 1848, Daufresne et al. 2009, Merckx et al. 2018). While one long-

term study of flying insects in the Netherlands found no evidence of higher rates of decline in

larger species over the past two decades (Hallmann et al. 2020), larger-bodied species may have

become rare earlier in the last century (Seibold et al. 2015). Climate and land-cover change may

otherwise alter insect communities by favoring particular trophic levels (Agosta et al. 2018,

Welti et al. 2020b), invasive species (Ju et al. 2017), or pest species with outbreak dynamics

(Bernal and Medina 2018). Furthermore, insect responses to climate and land cover may vary

with microhabitat preference (Suggitt et al. 2018). For example, flying insects may be more

affected by rising temperatures than other insects as they cannot buffer high temperatures by

burrowing in soil or plant tissue (Baudier et al. 2015, Wagner 2020).

Future directions for insect monitoring

Limited large-scale, long-term standardized insect monitoring is one reason for low power in our

ability to disentangle different potential drivers of insect decline, and understand how this varies

with region and taxa. Empirical studies of insect communities often lack the spatial coverage

needed to be broadly representative or to test for variation due to land cover types (but see

Jeliazkov et al. 2016, Wepprich et al. 2019). Meta-analyses have large spatial coverage, but can

be subject to site and/or publishing bias and must reckon with variation in research goals and

methodology (Gurevitch and Mengersen 2010). Spatially distributed observational monitoring

efforts of multiple biological taxa have tended to target plants and vertebrates but not insects

(Eggleton 2020). However, several notable exceptions include mosquito and ground beetle

monitoring by the US National Ecological Observation Network (Thorpe et al. 2016, Blair et al.

2020), and regional-scale monitoring programs of butterflies and macro-moths (e.g. Kühn et al.

2008, Dennis et al. 2019, van Swaay et al. 2019). Additionally, several large scale malaise trap

programs including the Global Malaise Trap program operating since 2012

(http://biodiversitygenomics.net/projects/gmp/) and the Swedish Malaise Trap Program

(operational from 2003-2006; Karlsson et al. 2020) are taxonomic treasure troves. However,

neither measure insect biomass. More distributed, standardized, and continuous insect

monitoring programs are needed to fill gaps in current knowledge of temporal insect population

and community dynamics (Cardoso and Leather 2019).

The German Malaise Trap Program helps to fill this gap for Germany through both

continued monitoring of flying insect biomass and recently begun DNA metabarcoding. This

program provides a blueprint for standardized long-term biodiversity monitoring across Europe

(Haase et al. 2018), with malaise traps currently being considered as a standard component of

European insect biodiversity surveys. European LTER network (eLTER RI; Mirtl et al. 2018)

was recently accepted by the EU as part of the European Research Infrastructure Roadmap

(ESFRI), and the more than 400 European LTER sites are currently working to establish stronger

linkages and comparable monitoring schemes. Continued long-term monitoring by the German

Malaise Trap Program will advance our understanding of trends on spatial-temporal scales that

are lacking in previous studies on insects (Didham et al. 2020), and in productivity/biodiversity

trends more generally (McGill et al. 2015). As highlighted by the recent insect decline debate

(e.g. Wagner et al. 2021), comprehensive and standardized monitoring is critical to meet the

challenge of unravelling insect trends and drivers in the Anthropocene.

Data Archiving Statement

Flying insect biomass and trap location characteristics will be uploaded to the B2SHARE

depository upon acceptance of this manuscript.

References

Abatzoglou, J. T. et al. 2018. TerraClimate, a high-resolution global dataset of monthly climate
and climatic water balance from 1958–2015. - Scientific Data 5: 170191.
Agosta, S. J. et al. 2018. Upper thermal limits differ among and within component species in a
tritrophic host-parasitoid-hyperparasitoid system. - PLOS ONE 13: e0198803.
Arnold, T. W. 2010. Uninformative parameters and model selection using Akaike's Information
Criterion. - J. Wildl. Manage. 74: 1175–1178.
Atkinson, D. 1994. Temperature and organism size—A biological law for ectotherms? - In:
Begon, M. and Fitter, A. H. (eds), Advances in Ecological Research. Academic Press, pp.
1–58.
Baker, N. J. et al. 2021. Drivers of long-term changes in aquatic macroinvertebrate communities
within a near pristine mountainous stream. - Science of The Total Environment 785:
14685.
Baranov, V. et al. 2020. Complex and nonlinear climate-driven changes in freshwater insect
communities over 42 years. - Conservation Biology 34: 1241–1251.
Barnes, A. D. et al. 2016. Species richness and biomass explain spatial turnover in ecosystem
functioning across tropical and temperate ecosystems. - Philosophical Transactions of the
Royal Society B: Biological Sciences 371: 20150279.
Bartoń, K. 2016. MuMIn: Multi-Model Inference.
Bates, D. et al. 2015. Fitting linear mixed-effects models using lme4. - Journal of Statistical
Software 67: 1–48.
Baudier, K. M. et al. 2015. Microhabitat and body size effects on heat tolerance: implications for
responses to climate change (army ants: Formicidae, Ecitoninae). - J Anim Ecol 84:
1322–1330.
Bender, S. et al. 2017. Einfluss des Klimawandels auf die terrestrischen Wassersysteme in
Deutschland. Eine Analyse ausgesuchter Studien der Jahre 2009 bis 2013.
Bengtsson, J. et al. 2005. The effects of organic agriculture on biodiversity and abundance: a
meta-analysis. - Journal of Applied Ecology 42: 261–269.
Bergmann, C. 1848. Über die Verhältnisse der Wärmeökonomie der Thiere zu ihrer Größe.
Bernal, J. S. and Medina, R. F. 2018. Agriculture sows pests: how crop domestication, host
shifts, and agricultural intensification can create insect pests from herbivores. - Current
Opinion in Insect Science 26: 76–81.
Blair, J. et al. 2020. Robust and simplified machine learning identification of pitfall trap-
collected ground beetles at the continental scale. - Ecology and Evolution in press.
Burnham, K. P. and Anderson, D. R. 2003. Model selection and multimodel inference: A
practical information-theoretic approach. - Springer-Verlag.
Büttner, G. 2014. CORINE land cover and land cover change products. - In: Manakos, I. and
Braun, M. (eds), Land Use and Land Cover Mapping in Europe: Practices & Trends.
Remote Sensing and Digital Image Processing. Springer Netherlands, pp. 55–74.
Cardoso, P. and Leather, S. R. 2019. Predicting a global insect apocalypse. - Insect Conservation
and Diversity 12: 263–267.
Carvalheiro, L. G. et al. 2020. Soil eutrophication shaped the composition of pollinator
assemblages during the past century. - Ecography 43: 209–221.
Cranmer, L. et al. 2012. Landscape structure influences pollinator movements and directly
affects plant reproductive success. - Oikos 121: 562–568.
Dangles, O. et al. 2011. Predicting richness effects on ecosystem function in natural
communities: insights from high-elevation streams. - Ecology 92: 733–743.

Daufresne, M. et al. 2009. Global warming benefits the small in aquatic ecosystems. - PNAS
106: 12788–12793.
Dennis, E. B. et al. 2019. Trends and indicators for quantifying moth abundance and occupancy
in Scotland. - J Insect Conserv 23: 369–380.
Díaz, S. et al. 2019. Pervasive human-driven decline of life on Earth points to the need for
transformative change. - Science in press.
Didham, R. K. et al. 2020. Interpreting insect declines: seven challenges and a way forward. -
Insect Conservation and Diversity 13: 103–114.
Eggleton, P. 2020. The state of the world's insects. - Annual Review of Environment and
Resources 45: 61–82.
European Union, Copernicus Land Monitoring Service 2018. European Environment Agency
(EEA). in press.
Fenoglio, M. S. et al. 2020. Negative effects of urbanization on terrestrial arthropod
communities: A meta-analysis. - Global Ecology and Biogeography 29: 1412–1429.
Fick, S. E. and Hijmans, R. J. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces
for global land areas. - International Journal of Climatology 37: 4302–4315.
Fox, J. and Weisberg, S. 2019. An R Companion to Applied Regression. - Sage Publications,
Inc.
GeoBasis-DE / BKG 2013. Digital Terrain Model grid wisth 200 m (DGM200).
Goulson, D. et al. 2018. Rapid rise in toxic load for bees revealed by analysis of pesticide use in
Great Britain. - PeerJ 6: e5255.
Gurevitch, J. and Mengersen, K. 2010. A statistical view of synthesizing patterns of species
richness along productivity gradients: devils, forests, and trees. - Ecology 91: 2553–2560.
Haase, P. et al. 2018. The next generation of site-based long-term ecological monitoring: Linking
essential biodiversity variables and ecosystem integrity. - Science of The Total
Environment 613–614: 1376–1384.
Habel, J. C. et al. 2019. Mitigating the precipitous decline of terrestrial European insects:
Requirements for a new strategy. - Biodivers. Conserv. 28: 1343–1360.
Haddad, N. M. et al. 2000. The effects of long-term nitrogen loading on grassland insect
communities. - Oecologia 124: 73–84.
Halbritter, A. H. et al. 2013. How comparable are species distributions along elevational and
latitudinal climate gradients? - Global Ecology and Biogeography 22: 1228–1237.
Hallmann, C. A. et al. 2017. More than 75 percent decline over 27 years in total flying insect
biomass in protected areas. - PLoS One 12: e0185809.
Hallmann, C. A. et al. 2020. Declining abundance of beetles, moths and caddisflies in the
Netherlands. - Insect Conservation and Diversity 13: 127–139.
Harris, I. et al. 2014. Updated high-resolution grids of monthly climatic observations - the CRU
TS3.10 Dataset. - Int. J. Climatol. 34: 623–642.
Harvey, J. A. et al. 2020. Climate change-mediated temperature extremes and insects: From
outbreaks to breakdowns. - Global Change Biology in press.
Hausmann, A. et al. 2020. Toward a standardized quantitative and qualitative insect monitoring
scheme. - Ecology and Evolution 10: 4009–4020.
Hofmann, M. M. et al. 2018. Changes in the bee fauna of a German botanical garden between
1997 and 2017, attributable to climate warming, not other parameters. - Oecologia 187:
701–706.

IPBES 2019. Summary for policymakers of the global assessment report on biodiversity and ecosystem services. S. Díaz, J. Settele, E. S. Brondízio E.S., H. T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A. Purvis, J. Razzaque, B. Reyers, R. Roy Chowdhury, Y. J. Shin, I. J. Visseren-Hamakers, K. J. Willis, and C. N. Zayas (eds.).: 11.

Jachuła, J. et al. 2017. Validation of floral food resources for pollinators in agricultural landscape in SE Poland: Validation of floral food resources for pollinators. - Journal of the Science of Food and Agriculture in press.

Janzen, D. H. and Hallwachs, W. 2019. Perspective: Where might be many tropical insects? - Biological Conservation 233: 102–108.

Jeliazkov, A. et al. 2016. Large-scale semi-automated acoustic monitoring allows to detect temporal decline of bush-crickets. - Global Ecology and Conservation 6: 208–218.

Ju, R.-T. et al. 2017. Spring warming increases the abundance of an invasive specialist insect: links to phenology and life history. - Scientific Reports 7: 14805.

Karlsson, D. et al. 2020. The data of the Swedish Malaise Trap Project, a countrywide inventory of Sweden's insect fauna. - Biodiversity Data Journal 8: e56286.

Kaspari, M. et al. 2019. Species energy and Thermal Performance Theory predict 20-yr changes in ant community abundance and richness. - Ecology 100: e02888.

Kingsolver, J. and Huey, R. 2008. Size, temperature, and fitness: Three rules. - Evolutionary Ecology Research 10: 251–268.

Klein, A.-M. et al. 2007. Importance of pollinators in changing landscapes for world crops. - Proceedings of the Royal Society B: Biological Sciences 274: 303–313.

Klockmann, M. et al. 2017. Carried over: Heat stress in the egg stage reduces subsequent performance in a butterfly. - PLOS ONE 12: e0180968.

Kühn, E. et al. 2008. Getting the public involved in butterfly conservation: Lessons learned from a new monitoring scheme in Germany. - Israel Journal of Ecology and Evolution 54: 89–103.

Kühsel, S. and Blüthgen, N. 2015. High diversity stabilizes the thermal resilience of pollinator communities in intensively managed grasslands. - Nature Communications 6: 7989.

Kwon, E. et al. 2019. Geographic variation in the intensity of warming and phenological mismatch between Arctic shorebirds and invertebrates. - Ecological Monographs 89: e01383.

Laws, A. N. and Joern, A. 2013. Predator–prey interactions in a grassland food chain vary with temperature and food quality. - Oikos 122: 977–986.

Lawson, D. A. and Rands, S. A. 2019. The effects of rainfall on plant–pollinator interactions. - Arthropod-Plant Interactions 13: 561–569.

Losey, J. E. and Vaughan, M. 2006. The economic value of ecological services provided by insects. - BioScience 56: 311–323.

Macgregor, C. J. et al. 2019. Moth biomass increases and decreases over 50 years in Britain. - Nature Ecology and Evolution in press.

McGill, B. J. et al. 2015. Fifteen forms of biodiversity trend in the Anthropocene. - Trends in Ecology & Evolution 30: 104–113.

Merckx, T. et al. 2018. Body-size shifts in aquatic and terrestrial urban communities. - Nature 558: 113–116.

Mirtl, M. et al. 2018. Genesis, goals and achievements of Long-Term Ecological Research at the
global scale: A critical review of ILTER and future directions. - Science of The Total
Environment 626: 1439–1462.
Myhre, G. et al. 2019. Frequency of extreme precipitation increases extensively with event
rareness under global warming. - Scientific Reports 9: 16063.
Newbold, T. et al. 2020. Global effects of land use on biodiversity differ among functional
groups. - Functional Ecology 34: 684–693.
Owens, A. C. S. et al. 2020. Light pollution is a driver of insect declines. - Biological
Conservation 241: 108259.
Pellegrino, A. et al. 2013. Weather forecasting by insects: Modified sexual behaviour in response
to atmospheric pressure changes. - PloS one 8: e75004.
Peters, M. K. et al. 2019. Climate–land-use interactions shape tropical mountain biodiversity and
ecosystem functions. - Nature 568: 88–92.
Pianka, E. R. 1966. Latitudinal gradients in species diversity: A review of concepts. - The
American Naturalist 100: 33–46.
Piano, E. et al. 2020. Urbanization drives cross-taxon declines in abundance and diversity at
multiple spatial scales. - Glob Chang Biol 26: 1196–1211.
Polidori, C. et al. 2020. Climate change-driven body size shrinking in a social wasp. - Ecological
Entomology 45: 130–141.
Potts, S. G. et al. 2010. Global pollinator declines: trends, impacts and drivers. - Trends in
Ecology & Evolution 25: 345–353.
QGIS.org 2020. QGIS Geographic Information System. Open Source Geospatial Foundation
Project.
R Core Team 2020. R: A language and environment for statistical computing. - R Foundation for
Statistical Computing.
Rada, S. et al. 2019. Protected areas do not mitigate biodiversity declines: A case study on
butterflies. - Diversity and Distributions 25: 217–224.
Reilly, J. R. et al. 2020. Crop production in the USA is frequently limited by a lack of
pollinators. - Proceedings of the Royal Society B: Biological Sciences 287: 20200922.
Root, R. B. 1973. Organization of a plant-arthropod association in simple and diverse habitats:
The fauna of collards (Brassica oleracea). - Ecological Monographs 43: 95–124.
Seibold, S. et al. 2015. Association of extinction risk of saproxylic beetles with ecological
degradation of forests in Europe. - Conservation Biology 29: 382–390.
Seibold, S. et al. 2019. Arthropod decline in grasslands and forests is associated with landscape-
level drivers. - Nature 574: 671–674.
Shortall, C. R. et al. 2009. Long-term changes in the abundance of flying insects. - Insect
Conservation and Diversity 2: 251–260.
Sinclair, B. J. et al. 2016. Can we predict ectotherm responses to climate change using thermal
performance curves and body temperatures? - Ecology Letters 19: 1372–1385.
Stepanian, P. M. et al. 2020. Declines in an abundant aquatic insect, the burrowing mayfly,
across major North American waterways. - PNAS 117: 2987–2992.
Suggitt, A. et al. 2018. Extinction risk from climate change is reduced by microclimatic
buffering. - Nature Climate Change 8: 713–717.
Svenningsen, C. S. et al. 2020. Contrasting impacts of urban and farmland cover on flying insect
biomass. - bioRxiv: 2020.09.16.299404.

Termaat, T. et al. 2019. Distribution trends of European dragonflies under climate change. - Diversity and Distributions 25: 936–950.

Thakur, M. P. 2020. Climate warming and trophic mismatches in terrestrial ecosystems: the green–brown imbalance hypothesis. - Biology Letters 16: 20190770.

Theodorou, P. et al. 2020. Urban areas as hotspots for bees and pollination but not a panacea for all insects. - Nature Communications 11: 576.

Thorpe, A. S. et al. 2016. Introduction to the sampling designs of the National Ecological Observatory Network Terrestrial Observation System. - Ecosphere 7: e01627.

Totland, Ö. 1994. Intraseasonal variation in pollination intensity and seed set in an alpine population of Ranunculus acris in southwestern Norway. - Ecography 17: 159–165.

van Swaay, C. et al. 2019. The EU butterfly indicator for grassland species: 1990-2017.

Vanbergen, A. J. and Insect Pollinators Initiative 2013. Threats to an ecosystem service: pressures on pollinators. - Frontiers in Ecology and the Environment 11: 251–259.

Wagner, D. L. 2020. Insect declines in the Anthropocene. - Annual Review of Entomology 65: 457–480.

Wagner, D. L. et al. 2021. Insect decline in the Anthropocene: Death by a thousand cuts. - PNAS in press.

Warren, R. et al. 2018. The projected effect on insects, vertebrates, and plants of limiting global warming to 1.5°C rather than 2°C. - Science 360: 791–795.

Welti, E. A. R. and Joern, A. 2015. Structure of trophic and mutualistic networks across broad environmental gradients. - Ecology and evolution 5: 326–334.

Welti, E. A. R. et al. 2020a. Nutrient dilution and climate cycles underlie declines in a dominant insect herbivore. - Proc. Natl. Acad. Sci. U. S. A. 117: 7271–7275.

Welti, E. A. R. et al. 2020b. Salty, mild, and low plant biomass grasslands increase top-heaviness of invertebrate trophic pyramids. - Global Ecology and Biogeography 29: 1474–1485.

Wepprich, T. et al. 2019. Butterfly abundance declines over 20 years of systematic monitoring in Ohio, USA. - PLOS ONE 14: e0216270.

Wilson, R. J. and Fox, R. 2020. Insect responses to global change offer signposts for biodiversity and conservation. - Ecological Entomology in press.

Winfree, R. et al. 2011. Native pollinators in Anthropogenic habitats. - Annual Review of Ecology, Evolution, and Systematics 42: 1–22.

Table

Table 1. Top AIC$_c$ model of predictors of flying insect biomass. Model includes the random

variable of trap location. T-tests use Satterthwaite's method. Poly(X,1) indicates the first-order

term of the 2nd degree polynomial for variable X, while poly(X,2) indicates the second-order

term of the 2nd degree polynomial. Predictor variables are defined in the Materials and Methods.

Model characteristics include estimate (Est), standard error (SE), degrees of freedom (df), t-

value, and p-value (*P*).

	Est	**SE**	**df**	**t-value**	*P*
Intercept	7.84	0.95	89.46	8.24	< 0.001
%forest	-0.33	0.10	87.6	-3.31	0.001
latitude	-0.09	0.02	87.51	-4.79	< 0.001
poly(period,1)	-3.30	0.39	981.7	-8.56	< 0.001
poly(period,2)	-4.02	0.81	1011	-4.94	< 0.001
poly(tmax,1)	8.05	0.97	1017	8.34	< 0.001
poly(tmax,2)	0.21	0.35	961	0.59	0.55
precip	-0.13	0.04	1006	-2.96	0.003
Δtemp	-0.01	0.003	990.5	-3.81	< 0.001

Figures

Figure 1. Map of sampling locations. Malaise trap locations where samples were collected

beginning in 2019 are depicted as filled circles and color-coded by the dominant land cover in

the surrounding 1 km. Points coded as grey triangles indicate trap locations at which sampling

began in 2020 and are not included in the analyses. Overlapping locations were jittered

longitudinally to improve visualization.

Figure 2. Responses of flying insect biomass to the climate variables and latitude. Each point

represents the biomass at each site within each month ($B_{i,j}$ as calculated by Eq. 1). Across all

months and site combinations, flying insect biomass increased with mean monthly 2019

maximum temperature (a), and decreased with latitude (c) and cumulative monthly 2019

precipitation (c). The response of flying insect biomass to Δtemp shifted from positive when

examined alone to a negative when the effect of 2019 tmax was included in the model. We

therefore used the residuals of the relationship between biomass and 2019 tmax as our response

variable to show the negative relationship between insect biomass and Δtemp evident in the top

AIC$_c$ model (Table 1) (d).

Figure 3. The effect of temperature on flying insect biomass across latitude. The effect of

temperature shifted from negative to positive with increasing latitude across the 48 sites with

sampling in all 7 months (Apr.-Oct. 2019). Model coefficients are provided in Table S2.

Figure 4. Flying insect biomass over the growing season by dominant land cover. Biomass

(mg/day) over the median sampling day across all 84 trap locations (a), and comparisons

between all locations and locations with surroundings dominated by forests (b),

grassland/shrubland (c), pasture/orchard (d), non-irrigated cropland (e), and urban environments

(f). Point colors in panel (d) match the dominant land category following colors in panels B-F.

Mean and standard error are provided for biomass within each land cover category and month.

Stars indicate significant differences within each month based on Welch's t-tests between

biomass from dominant land cover categories and all site averages (* = $0.05 > P > 0.01$, ** =

$0.01 > P > 0.001$, *** = $P < 0.001$).

Supporting Information for:

Climate, latitude, and land cover predict flying insect biomass across a German malaise trap network

Contents:

Table S1-S3

Fig. S1-S4

Table S1. Top AIC models (ΔAICc<2) of predictors of flying insect biomass. All models included the random variable of trap identity. Int=intercept, poly(X,1) indicates the first-order term of the 2nd degree polynomial for variable X, while poly(X,2) indicates the second-order term of the 2nd degree polynomial. Predictor variables are defined in the Methods. Marg R^2= marginal R^2 or the percent variance explained by the fixed effects, Cond R^2= conditional R^2 or the percent variance explained by the fixed effects plus the random effect of trap, df= degrees of freedom, logLik= log likelihood, Δ= ΔAIC_c, w= model weight. For summary tables of the top model, see Table 1. For summary tables of the second and third top models, see Appendix 1: Table S2.

Int	latitude	poly (period,2)	poly (tmax,2)	Δtemp	precip	%forest	%urban	%pasture/ orchard	Marg R^2	Cond R^2	df	logLik	AICc	Δ	w
7.84	-0.088	+	+	-0.0124	-0.131	-0.328			0.51	0.74	11	-364.35	751	0	0.147
8.26	-0.095	+	+	-0.0126	-0.133	-0.402	-0.405		0.52	0.74	12	-363.55	751.4	0.44	0.118
7.81	-0.089	+	+	-0.0125	-0.131	-0.292		0.253	0.51	0.74	12	-364.3	752.9	1.94	0.056

Table S2. Summary tables of the second (a) and third (b) mixed effects models included in top AIC$_c$ models (Table S1). A summary table of the top model is provided in Table 1. All models contain the random variable of trap identity. T-tests use Satterthwaite's method. See Methods and Table 1 caption for variable explanations. Model characteristics include estimate (Est), standard error (SE), degrees of freedom (df), t-value, and p-value (*P*).

	Est	SE	df	t-value	*P*
a.) Model 2					
Intercept	8.26	0.98	87.66	8.43	< 0.001
%forest	-0.40	0.11	83.87	-3.7	< 0.001
latitude	-0.10	0.02	86.01	-5.07	< 0.001
poly(period,1)	-3.30	0.39	982.1	-8.55	< 0.001
poly(period,2)	-3.99	0.81	1011	-4.91	< 0.001
poly(tmax,1)	8.10	0.97	1016	8.39	< 0.001
poly(tmax,2)	0.22	0.35	960.2	0.63	0.53
precip	-0.13	0.04	1006	-2.99	0.003
Δtemp	-0.01	0.003	989.9	-3.86	< 0.001
%urban	-0.40	0.26	84.52	-1.59	0.12
b.) Model 3					
Intercept	7.81	0.95	88.53	8.21	< 0.001
%forest	-0.29	0.10	87.4	-2.80	0.006
latitude	-0.09	0.02	86.39	-4.81	< 0.001
%pasture/orchard	0.25	0.23	80.44	1.09	0.28
poly(period,1)	-3.32	0.39	979.9	-8.59	< 0.001
poly(period,2)	-3.98	0.81	1009	-4.89	< 0.001
poly(tmax,1)	8.10	0.97	1015	8.38	< 0.001
poly(tmax,2)	0.20	0.35	961	0.59	0.56
precip	-0.13	0.04	1006	-2.94	0.003
Δtemp	-0.01	0.003	989.5	-3.84	< 0.001

Table S3. Model coefficients for the interaction between monthly temperature and latitude (Fig. 3). While individual model coefficients were all marginally significant, the full model predicted 23% of the variation in average biomass (mg/day) averaged across the seven sampling months for the 48 sites with sampling in all months ($F_{3,44}$ = 4.3, R^2 = 0.23, P = 0.01). GST= growing season temperature, the average temperature for each site from Apr-Oct 2019. Model characteristics include estimate (Est), standard error (SE), degrees of freedom (df), t-value, and p-value (*P*).

	Est	**SE**	**t-value**	*P*
Intercept	43.1	21.65	1.99	0.053
GST	-2.1	1.17	-1.8	0.079
Latitude	-0.81	0.42	-1.91	0.063
GST * Latitude	0.04	0.02	1.86	0.07

Figure S1. Examples of traps running in 2019 as part of the German Malaise Trap Program. Photos show traps at the LTER site Tereno- Friedeburg (a; photo credit: Mark Frenzel), at the Harz National Park (b; photo credit: Andreas Marten), at the Black Forest National Park (c; photo credit: Jörn Buse), and at the LTER site Rhine-Main-Observatory (d; photo credit: Peter Haase).

Figure S2. Comparison of climate at the 84 trap locations between 2019 and long-term average (1960-2018) including the average maximum monthly temperatures (tmax) and minimum monthly temperatures (tmin) in °C ± standard error (a) and cumulative monthly precipitation in mm ± standard error (b). Period of the year in which malaise trap sampling occurred is shaded in grey.

Figure S3. Changes with latitude across our 84 trap locations in 2019 mean monthly maximum temperature (a), the 1960-2018 long-term average monthly maximum temperature (b), the change in 2019 mean monthly maximum temperature minus the 1960-2018 long-term average (c), and 2019 cumulative monthly precipitation (d). Each point represents one month at one location, and only month/location combinations from which flying insect biomass data were collected are included. Across all months, 2019 mean monthly maximum temperature showed a weak trend to decrease with latitude (a; $F_{1,520} = 3.6$, $R^2 = 0.01$, $P = 0.06$), while the 1960-2018 long-term average monthly maximum temperature did not vary with latitude (b; $F_{1,520} = 1.6$, $R^2 < 0.01$, $P = 0.21$). While varying with month, the average Δ °C of 2019 maximum temperature over the long-term average decreased with latitude (c; $F_{1,520} = 6$, $R^2 = 0.01$, $P = 0.01$), as did cumulative monthly precipitation (d; $F_{1,520} = 60.6$, $R^2 = 0.1$, $P < 0.001$).

Figure S4. Elevation declined with increasing latitude across our 84 trap locations ($F_{1,82} = 74.5$, $R^2 = 0.48$, $P < 0.001$).