1

Knockdown of Dnmt1 links Gene body DNA methylation to

regulation of gene expression and maternal-zygotic transition in

the wasp *Nasonia*

Deanna Arsala[1,2], Xin Wu[3], Soojin V. Yi[3], and Jeremy A. Lynch[1]

[1] Department of Biological Sciences, University of Illinois at Chicago, Chicago IL

[2] Dept. of Ecology & Evolution, University of Chicago, Chicago IL

[3] School of Biological Sciences, Georgia Institute of Technology, Atlanta GA

Abstract

23
Gene body methylation (GBM) is an ancestral form of DNA methylation

whose role in development has remained unclear. Unlike vertebrates, DNA

methylation is found exclusively in gene bodies in the wasp *Nasonia vitripennis*,

which provides a unique opportunity to interpret the role of GBM in development.

We confirmed that parental RNAi (pRNAi) knockdown of a DNMT1 ortholog (*Nv-*

Dnmt1a) in *Nasonia* leads to embryonic lethality and failures in cellularization and

morphogenesis. Using whole-genome bisulfite sequencing, we found a

widespread loss of GBM in *Nv-Dnmt1a* pRNAi embryos. Using RNAseq, we

found that methylated genes that lost GBM in the pRNAi samples were

exclusively downregulated during zygotic genome activation. Unexpectedly,

nearly all affected unmethylated genes were up-regulated after pRNAi. Lack of

proper clearance of mRNAs and abnormal activation drive this up-regulation,

indicating critical roles for *Nv-Dnmt1a* and GBM in the maternal-zygotic transition

(MZT) in the wasp, despite their absence in *Drosophila*.

Background

The maternal-zygotic transition (MZT) is an essential stage in multicellular

eukaryotic development and represents the transition of developmental control

<u>from</u> maternal genome products provided to the egg during oogenesis <u>to</u> the

products of newly formed zygotic genome [1, 2]. A major hallmark of the handover

to zygotic control is the coordination of the highly regulated decay of maternal

mRNAs with the activation of transcription from the zygotic genome.

The interactions between the maternal and zygotic genome have been

most comprehensively demonstrated in the insect model *Drosophila*

melanogaster. Maternal transcript destabilization is carried out in part by

maternally deposited mRNAs encoding RNA binding proteins (RBPs) such as

smaug and *brain tumor* (*brat*) [3, 4]. These factors are aided by the transcription

factor Zelda (Zld), which promotes miRNA expression that further destabilize

maternal transcripts. Under normal circumstances Zld also increases chromatin

accessibility and helps activate thousands of zygotic genes to their proper level [5,

6, 7]. However, if clearance of maternal transcripts is impeded, the zygotic genome

is not properly activated. Thus, disruption of either zygotic genome activation, or

maternal transcript clearance leads to embryonic lethality. In both cases death is

due to failures in the earliest developmental events that are dependent on the

zygotic genome: cellularization of the syncytial blastoderm, and gastrulation [5, 8] .

Most of the strategies for regulating the *D. melanogaster* ZGA (e.g.,

histone modification, chromatin packaging, action of miRNAs, mRNA

degradation), and many of the molecules involved, are conserved throughout the

animals [9]. However, *D. melanogaster* lacks a major class of genomic regulation,

in the form of methylation of cytosines in CpG dinucleotides (CpG methylation) [10,

11]. CpG methylation is crucial for regulating gene expression in most complex

eukaryotes [12, 13], and has a significant role in regulating the MZT in many species

[14, 15]. Thus, its absence in *D. melanogaster*, and in the other major invertebrate

model system *C. elegans*, makes it difficult to elucidate the molecular

mechanisms of DNA methylation in early development.

CpG methylation is catalyzed by DNA methyltransferases 1 and 3

(DNMT1 and DNMT3), which are conserved and ancient components of the

eukaryotic genomic toolkit [16], but have been lost in Dipterans, including *D.*

melanogaster [17, 18]. However, both DNMT1 and 3 orthologs are found in many

other insect taxa including hemimetabolous insects [19] and within the

holometabolous lineage Hymenoptera [20, 21, 22, 23].

The most well understood functional impact of CpG methylation,

elucidated from studies of mammalian model system, is in silencing gene

expression via methylation of cis-regulatory sequences, particularly of normally

unmethylated clusters of CpGs (CpG islands) [13, 24]. However it is becoming clear

that cis-regulatory regulation at CpG islands is not a ubiquitous mechanism [25]. A

more universal, and likely ancestral [20, 26], form of DNA methylation occurs

between the transcription start site and transcription end site of a gene, and is

known as gene body methylation (**GBM**) [27].

Previous studies have correlated GBM with stable and higher levels of

gene expression than genes without GBM [26, 28, 29, 30, 31], but its role in regulating

gene expression in specific developmental contexts such as the MZT is poorly

understood. Whole-genome bisulfite sequencing (WGBS) studies have shown

that DNA methylation patterns are dynamic during the MZT in vertebrate model

systems [32, 33, 34]. However, such methylation studies in vertebrates have focused

on cis-regulatory regions, including CpG islands in promoters and intergenic

regions. Disruption of methylation at CpG islands often has profound phenotypic

effects, while the role of GBM is less well understood. Because manipulating

DNA methylation machinery in vertebrate models affects both CpG islands and

GBM, it is difficult to identify functions of GBM that are independent of CpG

island methylation.

In contrast to vertebrates, DNA methylation occurs almost exclusively at

gene bodies in insects that have retained DNA methyltransferases 1 and 3

(DNMT1 and DNMT3) [19, 22, 35, 36]. Thus, such insect model systems have the

potential to make significant contributions in understanding the role of GBM in

development. A particularly attractive system with a full methylation toolkit and

exclusively gene body methylation is the wasp *Nasonia vitripennis* [36, 37, 38].

There are three DNMT1 paralogs in the *N. vitripennis* genome*, and one of

them, *Nv-DNA methyltransferase 1a* (*Nv-Dnmt1a*), was shown to be essential for

embryogenesis (the other two showed no obvious phenotype)[39]. Therefore, *N.*

vitripennis offers an exciting opportunity to functionally study the specific role of

gene body methylation in a well-defined developmental system. In this work, we

set out to experimentally investigate the role of *Nv-Dnmt1a* and thus identify a

role for gene body methylation in early embryogenesis of *N. vitripennis*.

Our detailed analysis of the developmental effects of *Nv-Dnmt1a* revealed

major defects in cellularization and morphogenesis of the early embryo, which

are phenotypes typically seen after disruption of the MZT in insects. Using

parental RNAi (pRNAi) and WGBS we found that the vast majority of gene body

methylation is lost when *Nv-Dnmt1a* is knocked down. We then show that loss of

gene body methylation during zygotic genome activation is correlated with

reduced gene expression from the affected locus during zygotic genome

activation. Furthermore, we found that genes that are not methylated, but whose

expression changed after *Nv-Dnmt1a* knockdown, were primarily upregulated in

knockdown embryos relative to control. *Nv-Dnmt1a* knockdown profoundly

altered the levels of expression of thousands of genes in the early embryo.

Results

Nv-Dnmt1a function is required for early embryonic events that require

proper regulation of the zygotic genome

The initial characterization of *Nv-Dnmt1a* showed that knockdown through

parental RNAi (pRNAi), where female embryos are injected with dsRNA [40], led to

lethality of embryos produced by injected females [39]. Understanding the precise

developmental failure(s) that lead to embryonic lethality should provide insight

into the functional importance of *Nv-Dnmt1a* for early embryonic development.

Therefore, we examined the embryonic development following *Nv-Dnmt1a*

pRNAi in more detail.

We first characterized the events of the first half of embryogenesis (0-15

hours, from egg lay to the completion of gastrulation) using time lapse imaging of

Nv-Dnmt1a pRNAi embryos. Similar to *D. melanogaster*, early development is

very rapid in *N. vitripennis.* During early embryogenesis, nuclei divide rapidly,

simultaneously, and without the formation of cell membranes to fill the large,

unicellular egg with syncytial nuclei. After 7 mitotic division cycles, nuclei migrate

to the cortex of the embryo, forming a syncytial blastoderm [41], consisting of a

single layer of nuclei populating the entire cortex of the egg. We observed no

obvious differences between control embryos and *Nv-Dnmt1a* pRNAi embryos

during the first 11 syncytial divisions. Syncytial blastoderm formation was

normal, and the last 4 divisions showed the characteristic increase in length

relative to earlier, pre blastoderm, cycles (Figure 1A).

The first obvious defect occurred at the 12[th] division cycle (Figure 1B, 1D).

There is normally a significant pause at this stage, as the nuclei are surrounded

by plasma membrane and thus transition from a syncytial to a cellular state, a

process referred to as cellularization (Figure 1A). Normally, cellularization

directly precedes the first morphogenetic movements of the embryo (gastrulation

and migration of the extraembryonic serosa) [41]. In this work, control embryos

began gastrulation 193 minutes after the onset of nuclear cycle 12, on average

(n=5, SD = 13.12, (Figure 1A)), similar to our previously published results in

untreated embryos [42]. Apparent morphogenetic movements in *Nv-Dnmt1a* pRNAi

embryos began significantly earlier, on average, 171 minutes (n=5, SD = 5.98)

into cycle 12 (Figure 1A).

In addition to being premature, these movements were abnormal. In *Nv-*

Dnmt1a pRNAi embryos, a line of cells at the anterior end of the embryo began

to ingress posteriorly (Figure 1D) which was not seen in control embryos (Figure

1B). In addition, internalization of the mesoderm and migration of the serosa from

the dorsal pole to cover the embryo both failed (Figure 1C, 1E).

Figure 1. Analysis of time-lapse imaging of control and *Nv-Dnmt1a* RNAi
embryos. **A:** Box and whisker plot displaying the length of blastoderm division
times that were measured during live imaging experiments. Nc = nuclear cycle #.
P values were determined by performing a 2-tailed t-test assuming unequal
variance between RNAi and control. N=5 control, N=5 *Nv-Dnmt1a* RNAi. **B-C:**
Still images from time-lapse of a control embryo at the onset of gastrulation (B)
and at post-gastrulation. Peach arrow in (B) point to the initial mesodermal folds
which are eventually internalized by post-gastrulation (C). The blue arrowhead in
B indicates the serosa. **D-E:** Still images from time-lapse of an RNAi embryo at
the onset of gastrulation and at the time equivalent to post-gastrulation. Defective
mesodermal folds are denoted by peach arrowheads in D and E. Abnormal
serosal movements are denoted by a blue arrowhead in E. Scale bars represent
50 μm.

To get a more detailed understanding of the cellular basis of these

defects, we examined high-resolution images of fixed and stained embryos at the

end of cycle 12 in control and knockdown samples. We imaged transverse

sections across the anterior-posterior axis of *Nv-Dnmt1a* pRNAi and control

embryos at the end of the 12th cell cycle, when the cellularization and the first

gastrulation movements should be occurring. Gastrulation in *N. vitripennis* is

different from what is seen in *D. melanogaster*. *N. vitripennis* embryos lack a

ventral furrow. Rather, the presumptive mesoderm on the ventral side of the

embryo forms a stiff plate. Subsequently, ectodermal cells separate from the

mesoderm at the lateral borders between the germ layers and crawl over this

mesodermal 'plate' before fusing over the mesoderm at the ventral midline [41].

During mid-cellularization, we observed membrane ingressing evenly

around each nucleus (Figure 2A) in control embryos. In contrast, membrane

ingression was uneven and did not wholly encapsulate nuclei in *Nv-Dnmt1a*

pRNAi embryos, leaving many nuclei completely or mostly uncovered prior to the

onset gastrulation (Figure 2C, white arrows point to examples of uncellularized

nuclei).

At the onset of morphogenesis, cell membranes fully enveloped each

blastoderm nucleus in control embryos, and a strong cable of actin was present

along the interior (basal) surface of the cellular blastoderm (Figure 2B-B', orange

arrowhead in 2B' marks the basal actin cable). Enlarged, round serosal cells

were present on the dorsal side, and the distinct population of presumptive

mesodermal cells was present on the ventral side (Figure 2C, pink arrowheads

mark the borders of the presumptive mesoderm), 24 out of 24 control embryos

are fully cellularized). In contrast, *Nv-Dnmt1a* pRNAi embryos lacked the cable

of actin along the interior surface of the blastoderm (Figure 2D-2D', orange

arrowhead marks uncellularized nuclei and obvious absence of the basal actin

cable), consistent with failed cellularization (23 out of 24 pRNAi embryos were

incompletely cellularized). In addition, the blastoderm was multilayered at many

locations, and clearly distinct serosal and mesodermal precursors were not

observed (Figure 2D).

Condition	Cellularized	Incomplete Cellularization
Control	24	0
Nv-dnmt1a RNAi	1	23***

Figure 2. Transverse views of control and *Nv-Dnmt1a* RNAi embryos during
cellularization and at the onset of gastrulation. Green, phalloidin (F-actin). white,
DAPI (DNA). (A) Control embryo during mid cellularization. (B) Control embryo at
the onset of gastrulation. Pink arrowheads demarcate the borders of the
presumptive mesoderm. (B') Zoomed in view of fully cellularized control embryo.
Orange arrowhead points to the basal actin cable. (C) *Nv-Dnmt1a* RNAi embryo
during mid-cellularization. White arrows point to examples of disordered nuclei.
(D) *Nv-Dnmt1a* RNAi embryo at the onset of gastrulation. White arrows point to
examples of uncellularized, disordered nuclei. (D') Zoomed in view of an *Nv-*
Dnmt1a RNAi embryo that fails to fully cellularize. Orange arrowhead points to an

uncellularized nucleus (note the absence of the basal actin cable). (E) Table with
n values of cellularized and incompletely cellularized control and *Nv-Dnmt1a*
RNAi embryos. *** = p value < 0.0001 determined by Fisher's Exact Test. All
scale bars (white lines in A-A''') represent 10 μm.
219
220

Nv-Dnmt1a is required for global gene body methylation

The failure of cellularization and morphogenesis are widely conserved

indicators of disrupted MZT [42, 43, 44]. Consequently, we hypothesized that strong

reduction of *Nv-Dnmt1a* by pRNAi would disrupt DNA methylation and

dysregulate zygotic genome activation in the early embryo of *N. vitripennis*. We

thus sought to characterize the DNA methylation patterns in the early embryo of

N. vitripennis and to examine how this pattern was affected when *Nv-Dnmt1a*

was depleted by pRNAi.

Sample	Non-conversion rate	Median CpG coverage	Mean CpG coverage	Methylated CpGs (mCGs)	mCGs within gene bodies
Control 1	0.0052	14	14.92	136,600	126,333
Control 2	0.0051	15	15.63	136,410	126,174
Control 3	0.0055	14	14.75	136,717	126,364
RNAi 1	0.0052	14	14.30	98,511	91,874
RNAi 2	0.0051	14	14.48	8,017	7,720
RNAi 3	0.0053	14	14.91	15,305	14,669

Table 1. Summary CpG statistics and non-conversion rates for each sample.

In control embryos (derived from mock-injected females) collected at the

late blastoderm stage (7-9 hours after egg-lay, corresponding to the time where

the developmental defects become evident) we identified 151,480 methylated

CpGs (mCGs) in a union set representing all of the different mCGs across three

biological replicates. Each sample contained very similar numbers of mCGs

across the three biological replicates (ranging from 136,410 to 136,717). The

vast majority of these mCGs were found within gene bodies (Table 1).

Figure 3. Global DNA methylation in control and *Nv-Dnmt1a* pRNAi late
blastoderm *N. vitripennis* embryos. (A) An example of distribution of fractional
DNA methylation in control embryo exhibiting the classical bimodal distribution.
We defined methylated genes as genes with a minimum of 0.02 gene body
methylation (shown as the dashed line) to classify methylated and unmethylated
genes. (B) DNA methylation is substantially reduced in *Nv-Dnmt1a* pRNAi
samples compared to the control samples. (C) Distributions of fractional
methylation of CpGs that are methylated at least one control sample demonstrate
that CpGs lose DNA methylation in RNAi embryos. (D) Distribution of mean
fractional methylation in the first and last four exons and introns of methylated
genes. Control samples shown in red, RNAi embryos shown in blue.

Mean fractional DNA methylation of gene bodies followed the canonical

bimodal distribution (Figure 3A) as seen in other invertebrate species including

insects [27]. Following the criteria used in a previous study we designated genes

with gene body fractional methylation levels of at least 0.02 as methylated genes

[27]. By this criterion we found 5423 methylated genes in the blastoderm embryo.

This is fewer than the 6008 methylated genes identified in adult *N. vitripennis*

tissues [36], where a slightly different method to define methylated genes was

used. This difference was to some degree, but not entirely, due to the methods

used because when we used the method described in [36], the number of

methylated genes identified in our samples increased to 5,685. Regardless of the

methods, 99% of the genes we designated as "methylated" in the blastoderm

embryo were also identified as methylated genes in adult wasps [36]. This

indicates that our results are highly compatible with those of Wang et al., 2013 [36],

and are consistent with the idea that GBM patterns are generally stable across

life stages.

We knocked-down *Nv-Dnmt1a* mRNA levels with pRNAi and examined

DNA methylation at the same developmental stage (7-9 hours after egg lay) as

the control. Based on our RNAseq results, *Nv-Dnmt1a* mRNA levels were

reduced by 5-fold in late blastoderm pRNAi samples relative to late blastoderm

control samples [45]. We observed a significant reduction in DNA methylation at

99.7% of methylated CpGs in the genome in pRNAi embryos. The magnitude of

the reduction was substantial, as we found an average of 81.5% DNA

methylation depletion per mCG (Figure 3B, 3C and Table 1).

While DNA methylation was strongly reduced genome-wide, the intragenic

pattern of GBM was maintained. For example, mCGs were enriched in exons

and were more numerous at the 5' end of the coding region of gene bodies in

control embryos (Figure 3D) similar to patterns found in other insects and other

developmental stages [23, 35, 46]. This pattern was maintained in pRNAi embryos,

despite the drastic overall reduction in DNA methylation levels (Figure 3D). This

suggests that *Nv-Dnmt1a* is required for gene body methylation but does not

govern the 5' and exon-biased pattern of DNA methylation that we see in

Nasonia and other hymenopteran insects.

Figure 4. Gene expression of methylated and unmethylated genes in control and
Nv-Dnmt1a RNAi *N. vitripennis* embryos. **(**A) Volcano plots showing differentially
expressed genes between control and *Nv-Dnmt1a* RNAi in embryos of different
developmental stages. X axis is $\log_2$ fold change in *Nv-Dnmt1a* RNAi samples
relative to control samples. Y axis is $\log_{10}$ adjusted p-values. Golden dots are
methylated, significantly differentially expressed (FDR-corrected $P \leq 0.1$). Purple
dots are significant and DEGs that are unmethylated (FDR-corrected $P \leq 0.1$).
Blue and grey are non-significant and represent methylated and unmethylated
genes, respectively. (B) Distribution of methylated and unmethylated genes for all
genes in the genome, differentially methylated genes (DMGs), down-regulated
differentially expressed genes (Down DEGs), and up-regulated differentially
expressed genes (Up DEGs). (C) The correlation between relative GBM change
and relative expression change (calculated as $\log_2$-fold change) in the RNAi
samples compared to the control samples was positive and statistically
significant (Pearson R = 0.27, $P < 0.001$).

Loss of Nv-Dnmt1a function disrupts the regulation of a large proportion of

the early embryonic transcriptome

Given the profound effects of *Nv-Dnmt1a* on global DNA methylation and

early developmental events, we sought to understand the effects of *Nv-Dnmt1a*

knockdown on gene expression over time in the early embryo.   Although the

developmental effects *Nv-Dnmt1a* pRNAi were not visible until cellularization and

the onset of morphogenesis, we hypothesized that changes in the transcriptome

in response to disrupted development would occur earlier in development,

leading up to the multifaceted failure of development at gastrulation.

To test this, we performed RNA-seq on timed egg collections from mock

injected (control) and *Nv-Dnmt1a* dsRNA injected wasps at time points that cover

early embryogenesis from egg laying to gastrulation. The collection time points

were: freshly laid eggs (0-2 hours after egg lay), early blastoderm (3-5 hours after

egg lay), middle blastoderm (5-7 hours after egg lay) and late blastoderm/onset

of gastrulation (7-9 hours after egg lay, when the corresponding WGBS data

were collected).

We observed that there is a massive increase in the number of genes

significantly affected by *Nv-Dnmt1a* pRNAi the middle blastoderm stage, and that

this increase is maintained into the late blastoderm (Fig 4A). We found 835 and

888 differentially expressed genes (DEGs) in the egg and early blastoderm

stages, respectively, while there were 2744 and 1831 DEGs in the middle and

late blastoderm stages, respectively. The middle blastoderm stage corresponds

to the most active part of the MZT, where maternal mRNAs are being finally

cleared and new transcription from the zygotic genome becomes the

predominant source of mRNAs.

Another intriguing pattern that appeared at the mid blastoderm stage is a

clear correlation between the direction of change of a DEG, and whether the

gene was methylated or not (unmethylated) at the late blastoderm stage (herein

referred to as 'mDEGs' for methylated DEGs, and 'uDEGs' for unmethylated

DEGs).   Specifically, down-regulated transcripts in *Nv-Dnmt1a* pRNAi middle

blastoderm samples were largely derived from methylated genes (1201 out of

1331 (90%)), while upregulated transcripts mostly came from unmethylated

genes (1227/1413, 87%) (Figure 4A). Both of these observations were

statistically significant (Fisher's exact test, $P < 0.001$). A nearly identical pattern

was maintained into the late blastoderm stage (Fig. 4A, B).

Aside from the directions of change, we also observed that the

magnitudes of change at the gene level also differed between methylated and

unmethylated genes after *Nv-dnmt1a* pRNAi. For example, 93% (731/787) of

mDEGs at the late blastoderm stage were downregulated 4-fold or less, with a

significant proportion (315/787, 40%) being downregulated by less than two-fold

(Supplementary Figure 1A). The relative handful of up-regulated mDEGs showed

an even smaller magnitude of change, with 112/164 (68%) changing by 2 fold or

less (Supplementary Figure 1B). In contrast, uDEGs tended to show larger

magnitude changes, as 52% (323/619) of up-regulated uDEGs, and 35%

(39/110) of the down-regulated uDEGS changed by greater than 4-fold after

pRNAi relative to control (Supplementary Figure 1).

The strong over-representation of methylated genes in down-regulated

genes following *Nv-Dnmt1a* pRNAi fits well with previous observations in insects

connecting DNA methylation with increased and stable gene expression [19, 36, 47].

Furthermore, we have found that the magnitude of the reduction in expression

was highly correlated with the degree of lost methylation after pRNAi (Fig. 4C,

Pearson's correlation coefficient = 0.27, P < 0.001).

These observations imply that GBM plays an important role in maintaining

normal levels of high expression from methylated loci, and its loss leads to

widespread reduction in transcription from these loci. Assuming that GBM is the

sole molecular function of *Nv-Dnmt1a*, the effects on uDEGs, and potentially

some mDEGs, may be due to indirect effects downstream of the initial reduction

of mDEGs after pRNAi.

While indirect effects on unmethylated genes could be predicted, the very

strong bias toward upregulation of this gene class was quite surprising. We

compared our raw RNA-seq data and DESeq2 normalized data, to investigate if

any technical bias due to the normalization method could have affected our

inference of gene expression changes. There were no concerning or obvious

differences between the distribution of raw read counts and the normalized

counts used for our analyses (Supplementary Figure 2). Further the size-factors

required for normalization were all relatively small (Supplementary Table 1). This

indicates that there was no bias introduced by our normalization methods, and

that our results reflect a real signal in the transcriptome data.

Figure 5. Violin plots of methylated and unmethylated differentially expressed
genes (DEGs) in control and *Nv-Dnmt1a* pRNAi eggs, early blastoderm embryos,
middle blastoderm embryos, and late blastoderm embryos.

Large-scale patterns of change differ between methylated and

unmethylated differentially expressed genes

Due to the chaotic and variable nature of the failures of cellularization and

morphogenesis, we hypothesized that the lethal phenotype is due to cascading

effects initiated by direct effects of *Nv-Dnmt1a* pRNAi on methylated genes which

could be amplified by the downstream effects on unmethylated genes. The

strong correlation of DNA methylation state with the direction of expression level

change (Figure 2C) is consistent with this hypothesis and indicates that there are

fundamental differences in how these classes of genes are regulated in the early

embryo. Thus, we further examined variation of gene expression among genes

that showed differential expression between control and *Nv-Dnmt1a* pRNAi

treatments in at least one stage (a total of 3904 genes).

We identified 1769 methylated differentially expressed genes (mDEGs),

which showed a remarkably stable pattern of expression across all stages,

consistent with our understanding of methylated genes in general [31, 36]. Most

mDEGs were clustered around the median and at relatively high expression

levels, with the frequency of higher and lower expressed genes tapering rapidly

and smoothly from this cluster. While the general shape of the distribution was

maintained after pRNAi, the average expression was shifted significantly

downward, particularly during the middle and late blastoderm stages (Figure 5).

When we isolated the relative handful of up-regulated mDEGs, we found that the

shape of the distribution shifted upward as expected, and also kept the same

shape with most genes clustered around the median expression level

(Supplementary Figure 3).

We identified 2135 unmethylated DEGs (uDEGs). In controls, uDEGs had

a low median expression level and there was no detectable enrichment in the

number of genes near this median level (Figure 5). Most noticeably, there was a

significant proportion of uDEGs that were expressed at very low levels, while

such low expression levels were almost absent in the mDEGs (Figure 5).

Nv-Dnmt1a pRNAi had little effect on global distribution of uDEG

expression levels in the egg and early blastoderm stages (Figure 5A, B),

consistent with the minor effects of pRNAi on these stages reported earlier

(Figure 4A). However, in the middle and late blastoderm stages (Figure 5C, D),

we observed that very few genes were expressed at the very low levels. This

indicates that uDEGs that were absent or expressed at very low levels in control

were present at significantly higher levels after *Nv-Dnmt1a* pRNAi. Similar, but

less pronounced patterns are observed when all genes, and non-DEGs are

plotted in similar way (Supplementary Figure 4).

Formally, there are two explanations for this pattern, which are not

mutually exclusive. Increased mRNA from a given gene could be due to excess

transcription from the locus. Alternatively, higher levels at a given time point

could be due to a disruption in degradation of mRNA from a previous stage,

leading to the presence of excess reads.

To identify potentially informative subsets of DEGs, that were not obvious

from the global comparisons above, we used hierarchical clustering to group

genes based on the similarity of their patterns of expression over time in control

embryos and maintained the order of genes in *Nv-Dnmt1a* pRNAi embryos to

facilitate direct comparisons. Gene expression levels at different time points and

treatments were then plotted according to these clusters (Figure 6A,B).

Among all the mDEGs identified across the four timepoints, we identified a

cluster of transcripts that show upregulation in the early blastoderm *Nv-Dnmt1a*

pRNAi embryos relative to control embryos (Fig. 6A, blue box, dotted lines). The

genes in this cluster were provided maternally at very high expression levels

(median log2 expression value of 10) in both control and pRNAi samples (Figure

6C). In control embryos, expression levels fell precipitously for these genes as

development progressed (median log2 expression value of 7, an 8-fold reduction

from control (Figure 6C)).

Figure 6. Differentially expressed methylated and unmethylated gene
expression in *Nv-Dnmt1a* pRNAi and control *N. vitripennis* embryos. (A)
Heatmap of 1769 methylated differentially expressed genes (mDEGs) and their
expression over time in control and *Nv-Dnmt1a* pRNAi samples represented as
Z-scores. Blue boxes on the heatmap highlight a cluster of mDEGs that are of
interest and discussed in the text. (B) Heatmap of 2135 unmethylated DEGs
(uDEGs) and their expression over time in control and *Nv-Dnmt1a* pRNAi
samples represented as Z-scores. Black and red boxes on the heatmap highlight
two clusters of uDEGs that are of interest and discussed in the text. (C) Violin
plots of 376 mDEGs in *Nv-Dnmt1a* pRNAi eggs and early blastoderm embryos.
These mDEGs are significantly enriched in the early blastoderm *Nv-Dnmt1a*

pRNAi samples relative to control early blastoderm embryos. Both comparisons
were subjected to a t-test. ns = not significant, *** P<0.001 by t-test.
(D) Violin plots of 213 uDEGs in control and *Nv-Dnmt1a* pRNAi early and middle
blastoderm embryos (B). ns = not significant, *** P<0.001 by t-test.
445
446

In *Nv-Dnmt1a* pRNAi embryos, the levels of this set of genes also fall

significantly from egg to early embryo, but the magnitude of reduction is smaller

(median log2 expression value of ~8, a 4-fold reduction in median level), leading

to the appearance of upregulation relative to control at this stage (Figure 6C).

Thus, it appears that mRNA degradation downstream of Nv-Dnmt1a function

plays an important role in regulating levels maternally expressed, methylated

genes.

Among the uDEGs, we identified two clusters of interest. The larger of

these (Figure 6B, see black box) show much higher expression in middle and

late blastoderm in *Nv-Dnmt1a* pRNAi embryos relative to control. Expression

levels of these uDEGs are at low to moderate levels in eggs and early

blastoderm embryos in both control and pRNAi and remain so in the later stages

of control embryos (Figure 6B, compare consistent "cool" colors as opposed to

increased warmth in the RNAi samples). When we plot genes corresponding to

this cluster (change less than 2-fold between early and middle blastoderm

control, and are significantly higher after pRNAi in the middle blastoderm), we

found that 437 genes are mostly expressed at very low levels in the early

blastoderm in both control and pRNAi. At the middle blastoderm stage, there is a

significant general increase of these genes in the pRNAi case, while they remain

mostly unchanged at low levels in the control (Figure 6D). The most likely

explanation for this pattern is that the increased levels in pRNAi embryos are due

to improper activation of these genes at the middle and late blastoderm stage.

There is another cluster of uDEGs whose relative levels in control

embryos increase markedly in the early blastoderm stage and then fall steeply in

the subsequent stages (Figure 6B, see red box). In *Nv-Dnmt1a* pRNAi embryos,

the initial increase is not affected, but levels do not fall as significantly at later

stages, leading to an apparent upregulation of their transcripts relative to control

embryos in later stages (Figure 6B). The 615 genes that fit this profile show

again very similar distributions between the control and pRNAi cases in the early

blastoderm. In contrast there is a severe drop in levels for the control embryos at

the middle blastoderm stage, while the pRNAi levels decrease less dramatically

(Supplementary figure 5). This pattern is similar for the 376 mDEGs that were

apparently upregulated in the early blastoderm stage (Figure 6A,C) and indicates

that regulation of mRNA degradation is affected downstream of *Nv-Dnmt1a*

pRNAi during the major phase of the MZT.

Many potential targets for mediating the developmental effects of Nv-

Dnmt1a pRNAi

While there is no single obvious candidate gene to explain our results,

there are many RNA binding and regulating factors, as well as transcription

factors, cyclins and cytoskeletal components that could plausibly play important

roles in producing the patterns of gene expression change and developmental

failure after pRNAi. We performed gene ontology analyses, and found that terms

involved in metabolism, and other basal cell functions were the most significantly

and consistently enriched in both in methylated and unmethylated DEGs

throughout development (Supplementary Table 2). In addition, we found that

RNA degradation is an enriched category of methylated genes, which could

explain part of the pattern of gene expression change (Supplementary Table 2).

495   Among the RNA binding factors affected by *Nv-Dnmt1a* pRNAi are

homologs of the well-known MZT factors *Smaug* and the CCR-NOT deadenlyase

complex. Fascinatingly, we also found that an ortholog of the *D. melanogaster*

nonsense mediated mRNA decay (NMD) component *nonC (Nv-nonC)* was down-

regulated after *Nv-Dnmt1a* pRNAi. While the *D. melanogaster* gene has no role

in embryogenesis [48], the mouse ortholog (named Smg1 after the *C. elegans*

gene (no relation to *smaug*), which is the founding member of this gene family [49])

is required for embryogenesis [50]. Furthermore, the human ortholog of

Smg1/*nonC* is regulated by DNA methylation in humans [51]. We decided to

functionally test whether *Nv-nonC* could be an important intermediary in

producing the defects in cellularization and gastrulation when *Nv-Dnmt1a* was

knocked down.

507   Live imaging of *Nv-nonC* knock-down embryos revealed that affected

embryos consistently arrested at nuclear cycle 11 and did not proceed to the final

syncytial division or gastrulation (compare Supplementary Movie 1 (control) to

Supplementary Movie 2 (*Nv-nonC* RNAi)). This is more severe than the strongest

Nv-Dnmt1a phenotypes, so we cannot say for certain the exact role of *Nv-nonC*

in the regulatory networks downstream of *Nv-Dnmt1a* at the moment. However,

the phenotype is consistent with *Nv-Smg1* having an important role in the

transition from egg to embryo, and its partial reduction and disturbance of its

potential interaction partners after *Nv-Dnmt1a* pRNAi might play a significant role

in the failures of cellularization and gastrulation.

Discussion

A previous study found that one of the DNMT1 orthologs in the wasp (*Nv-*

Dnmt1a) was required for *Nasonia* embryogenesis, while the other two DNMT1

paralogs (*Nv-Dnmt1b*, *Nv-Dnmt1c*) and the DNMT3 ortholog *Nv-dnmt3* were

apparently dispensable [39]. *Nv-Dnmt1a* pRNAi embryos failed to form segments

and the knockdown was lethal [39] which led us to further examine the phenotype

in more detail [52, 53].

We found that the first signs of developmental failure are nuclei that fail to

be correctly cellularized. This directly precedes failure of the morphogenetic

processes of mesoderm internalization, and migration of the serosa. These are

all developmental events that are strongly affected when the MZT is disrupted in

N. vitripennis and *D. melanogaster*. Consistent with this, *Nv-Dnmt1a* knockdown

was associated with major disruption of gene expression levels during early

embryogenesis. The most significant alterations to the transcriptome occurred at

the mid blastoderm stage, which correlates well with when the major wave of

zygotic genome activation occurs in *N. vitripennis*. This observation, in

combination with our finding that GBM was significantly reduced after *Nv-Dnmt1a*

pRNAi, strongly indicates that loss of gene body methylation disrupts the MZT in

N. vitripennis.

The exact mechanism by which Nv-Dnmt1a regulates the MZT and

downstream processes is not yet understood. However, our current results

provide a good basis to generate hypotheses. A crucial observation was that

there was an extremely strong correlation between methylation state of a gene

and the direction of expression change after *Nv-Dnmt1a* pRNAi: downregulated

genes in pRNAi embryos were almost all methylated in control embryos, and

upregulated genes were conversely almost exclusively unmethylated (Figure 4).

Previous studies posited that gene body methylation has an important role

in facilitating high, steady levels of gene expression [17, 36]. However, whether gene

body methylation directly regulates levels of gene expression has been

controversial [54, 55]. Phylogenetic analyses of methylated and unmethylated plant

species as well as analyses of within-species variation of gene body methylation

in plants [54, 55, 56] led to at most weak effects of gene body methylation on gene

expression, although the evolutionary conservation of gene body methylation

indicated associated functionality. Here we show that as a result of pRNAi of *Nv-*

Dnmt1a, the vast majority of methylated genes that are differentially expressed

show reduction of gene expression. Moreover, the degree of expression

reduction directly correlated with the degree of gene body methylation loss. As

far as we are aware, these are the first such demonstrations directly linking gene

body DNA methylation to gene expression levels.

Another novel finding is that a large proportion of the unmethylated

differentially expressed genes were upregulated in *Nv-Dnmt1a* pRNAi relative to

control. Presumably the effects of *Nv-Dnmt1a* pRNAi on unmethylated genes are

indirect and are more proximally caused by the disrupted expression of the

methylated genes. It is not clear why such indirect effects result in such a large

bias toward increased gene expression. However, we noticed that the bulk of

unmethylated genes cluster at very low expression levels in the blastoderm

stages (Figure 5), and that most of the change in unmethylated genes we

observe after *Nv-dnmt1a* knockdown is a subset of these lowly expressed genes

being significantly increased (Figure 5). We propose that one or more mDEG is

important for keeping the lowly expressed transcripts at low levels (by either

repressing transcription or encouraging transcript degradation). Future

experiments will employ techniques that can specifically detect changes in

mRNA regulation and transcription initiation to determine how these processes

affect indirect targets of Nv-Dnmt1a and potentially which direct targets mediate

these effects.

The roles of DNMT1 orthologs in methylation and development have been

explored in other insects. In the cockroach, *Blatella germanica*, *Dnmt1* RNAi

results in embryonic lethality visible both before and after gastrulation [19]. The

effects of disrupting the MZT are less well understood in hemimetabolous

insects, but strong *B. germanica zelda (BgZelda)* RNAi knockdown leads to early

embryonic lethality similar to that of *B. germanica Dnmt1* (*BgDnmt1*) [57].

BgDnmt1 RNA leads to reduced GBM, and significant changes in both

methylated and unmethylated genes. However, both sets of genes show equal

amounts of up- and down-regulation, which is in contrast to what we see at the

middle and late blastoderm stages [19]. This could simply be due to changes the

role of GBM and the regulatory networks downstream of methylated genes that

have accumulated since the wasp and roach lineages diverged. Alternatively,

while our samples were precisely focused on capturing events directly at the

MZT, the samples examined in *B. germanica* were necessarily less temporally

restricted and extended further in developmental time. This may have allowed

for downstream knock-on effects to overwhelm a potentially conserved signal of

increased uDEGs and decreased mDEGs at the critical point in the MZT. More

studies of Dnmt function with a focus on the events of the MZT in other insects

will be required to resolve these questions.

Results of *Dnmt1* knockdown in the milkweed bug *Oncopeltus fasciatus*

and the beetle *Tribolium castaneum* indicate that Dnmt proteins may have

important functions not related to their methylation function [35, 58, 59]. In *O.*

fasciatus, GBM methylation was shown to be dependent on its Dnmt ortholog in

adults, and pRNAi against it led to disruption of oogenesis. However, no

significant changes in gene expression were detected in ovaries, or in other adult

tissues, raising the question of whether GBM is functionally relevant for

oogenesis [35]. Subsequently, *O. fasciatus* Dnmt was shown to be crucial for

meiosis in male testis, but again in and apparently GBM independent manner [58].

Consistent with this, *T. castaneum* still retains a Dnmt1 ortholog (but lacks

Dnmt3), but shows no significant genomic cytosine methylation [59]. Despite this,

knockdown of *T. castaneum* Dnmt1 leads to early embryonic lethality, indicating

methylation independent functions for Dnmt1 orthologs in beetles and bugs.

We believe that the clear correlation between reduction of methylation and

reduction of expression level of methylated genes is strong evidence that the

methyltransferase function of Nv-Dnmt1a is the major reason *Nv-Dnmt1a* is an

essential gene in *N. vitripennis*. However, we certainly cannot exclude that

additional functions of the protein exist, and that they contribute to the complex

phenotypes we observed after knocking down *Nv-Dnmt1a* with pRNAi. A

significant portion of our future aims is to isolate the role of GBM from other

potential effects of the RNAi. We also look forward to studies in other insects

that will show whether additional functions of Dnmt1 orthologs are novel,

ancestral or some combination of both.

Finally, our results point to the middle blastoderm stage as the crucial

point where *Nv-Dnmt1a* has its strongest effects on the embryonic transcriptome.

Our data also show that the indirect effects of *Nv-Dnmt1a* pRNAi are mediated

by both disruption of mRNA degradation, and misregulation of transcription from

the zygotic genome. Thus, *Nv-Dnmt1a* appears to exert its strongest effects at

the time, and using the molecular processes, that are most characteristic of the

MZT. We therefore propose that GBM is a crucial component of the MZT in

Nasonia, and that this epigenetic mark is likely to play an important role in the

MZT in other insects.

Acknowledgements

We thank The University of Chicago Genomics Facility (RRID:SCR_019196), for

their very valuable assistance with generating the WGBS and RNAseq data.

This work was supported by grant MCB1615664 from NSF to SVY, and by grants

R01GM129153 and R03HD087476 from NIH to JAL.

Methods

Live imaging

All embryos were live-imaged as described in Arsala & Lynch 2017 [42]. Briefly, we

live-imaged all embryos at 28C for 15 hours on an Olympus BX-80 inverted

microscope under 30X silicone immersion and DIC optics.

Embryo Collection

All embryos used for fluorescence in situ hybridizations were collected and fixed

as described as in Buchta et al., 2013 [41].

Ovary Collection

Mothers were cold anesthetized and their ovaries were hand-dissected and

extracted using stainless-steel needles. The ovaries were fixed in 5%

formaldehyde for 30 minutes and subsequently used for fluorescence in-situ

hybridization.

<u>Phalloidin staining & transverse sectioning</u>

Control injected and *Nv-Dnmt1a* dsRNA (1ug/ul) injected mothers were allowed

to lay eggs on hosts for 2 hours at 25C. Freshly laid eggs were collected from

hosts and aged on 1% agarose PBS plates at 25C until they were 8-10 hours old,

so that all embryos were aged to correspond with the onset of gastrulation. We

hand dissected the chorions of fixed embryos. Afterward, the fixed embryos

were placed in 1X PBS and stained with AlexaFluor 488 phalloidin at a 1:250

dilution. The stained, fixed embryos were placed in vectashield with DAPI

overnight. Finally, the embryos were laid flat on a cover slip and were cut in half

along the transverse axis using a razor slotted into an embryo 'guillotine.' The

resulting embryo 'halves' were flipped 'up' on the cover slip and imaged on an

Andor Revolution WD spinning disc confocal system. The samples were

illuminated with 488nm (phalloidin) and 405nm (DAPI) diode laser and Z-stack

images were captured using a 30x objective at 1micron increments.

<u>RNA extraction</u>

Embryos were collected and aged as described above. We designed 3 non-

overlapping dsRNA constructs and injected 100 females per biological replicate

with either 1µg/mL dsRNA diluted in water or with water as a control. The

females were left unmated, so that all resulting progeny were male as the

variation in sex and ploidy could confound our analysis. The injected females

were allowed to lay eggs that were aged at 25C until they reached the following

four time points across the MZT: 0-2hours to generate a freshly laid egg

transcriptome, 3-5hours to generate an early blastoderm transcriptome, 5-7

hours to generate a middle blastoderm transcriptome, 7-9 hours to generate late

blastoderm embryos that are at the onset of gastrulation. Once the embryos

reached the time points, we isolated 100ng of total RNA using TRI Reagent for

all samples in biological triplicate. We confirmed the RIN value for all samples

were above 9.0 using a 2100 Agilent Bioanalyzer and immediately prepared 24

libraries for sequencing.

RNA library preparation and bulk RNA-sequencing

Using the 100ng of total RNA from each sample isolated above, we performed

poly-A tail selection using the NEBNext Poly (A) mRNA magnetic isolation

module (E7490) and immediately proceeded with library preparation. Libraries

were prepared for bulk RNA-sequencing using the NEBNext Ultra Directional

RNA Library Prep Kit for Illumina (E7420). The libraries were purified, validated

and pooled before sequencing. The libraries were subjected to 100bp paired-end

sequencing across 2 lanes on a HiSeq4000. Raw data from this analysis can be

found in the NCBI BioProject archive with accession number PRJNA701367.

RNA-seq analysis

To assess sequencing quality, we using FastQC and saw high quality reads

across all samples. We trimmed adapters and removed low quality reads using

Trim Galore-0.6.0 only using the --paired parameter. We reassessed the trimmed

read quality using FastQC and proceeded to map the reads to genome using

HISAT2-2.0.5 (hisat2 -p 8 --max-intronlen 10000 -q –x). Differential expression

analysis was performed using the R package DESeq2[60].

<u>DNA extraction and bisulfite library preparation</u>

We extracted and purified genomic DNA from embryos using the QIAamp DNA

Micro Kit (Qiagen 56304). 40ng of genomic DNA was bisulfite converted and

libraries were prepared at the University of Chicago Genomics Facility using the

Accel-NGS Methyl-Seq DNA Library Kit from Swift Biosciences. Raw bisulfite

sequencing reads can be found in the NCBI BioProject archive with accession

number PRJNA701143.

<u>Analysis of Bisulfite-sequencing data</u>

CpG sites were determined to methylated using a binomial test [22, 36] where the

rate of deamination representing the probability of success and total number of

reads representing the number of trials [22, 36]. P-values were then corrected for

multiple testing [61] and CpGs with adjusted P-values of < 0.05 were labeled as

methylated CpGs. Genes containing at least one methylated CpG were

subsequently labeled as methylated genes. Fractional methylation of CpGs was

calculated by dividing the number of methylated reads by the total number of

reads and the fractional methylation of genes was calculated by taking the

average fractional methylation of all CpGs within the gene body.

For the differential methylation analysis, only CpG sites that were methylated in

at least one control sample were retained similar to previous insect studies [47, 62].

The DSS package [63] was used to assess differentially methylated CpGs with

RNAi status as the sole predictor of methylation. P-values for each CpG site was

adjusted for multiple testing [61] with significance level set at an adjusted P-value

of < 0.1. Genes with at least one differentially methylated CpG were considered

as differentially methylated genes.

Supplementary Data:

Sample	Size Factor
Egg control rep 1	2.008
Egg control rep 2	2.618
Egg control rep 3	2.389
Egg RNAi rep 2	1.855
Egg RNAi rep 3	2.422
Early control rep 1	0.799
Early control rep 2	0.789
Early control rep 3	1.041
Early RNAi rep 1	0.945
Early RNAi rep 2	1.234
Early RNAi rep 3	1.004
Middle control rep 2	0.848
Middle control rep 3	0.609
Middle RNAi rep 1	0.709
Middle RNAi rep 2	0.835
Middle RNAi rep 3	0.843
Late control rep 1	0.763
Late control rep 2	0.891
Late control rep 3	0.689
Late RNAi rep 1	0.822
Late RNAi rep 2	0.461
Late RNAi rep 3	1.026

726   **Supplementary Table 1.** Size Factors used for normalization of RNAseq
727   libraries
728

Category	Enriched terms	Count	%	P-Value	Benjamini
KEGG_PATHWAY	RNA degradation	16	2.1	0.00018	0.014
KEGG_PATHWAY	Oxidative phosphorylation	28	2.4	0.00064	0.032
KEGG_PATHWAY	Proteasome	16	1.4	0.00061	0.059
KEGG_PATHWAY	Basal transcription factors	10	1.3	0.0032	0.083
KEGG_PATHWAY	Endocytosis	21	2.7	0.0016	0.063
COG_ONTOLOGY	Translation, ribosomal structure and biogenesis	10	1.3	0.0023	0.044

731 **Supplementary Table 2**. GO Analysis for down-regulated methylated genes in
732 middle and late blastoderm *Nv-Dnmt1a* RNAi embryos

Gene	Forward Primer	Reverse Primer
Nv-dnmt1a	ggccgcggAGCGTCGATGCCTACTTATG	cccggggcAGCTTGTTCTCCACCATCAA
Nv-dnmt1a	ggccgcggACGTATGACTGGGTTGAGTC	cccggggcGGAGCATGAAGTTGGAACAC
Nv-dnmt1a	ggccgcggttaatcactaccccctgcat	cccggggcatgtttcgcccaatactgtt
Nv-nonC	GGCCGCGGCCTTGAGAAACGGACAGAGA	CCCGGGGCGAAGACATTGGCTCGGTTTC

737 **Supplementary Table 3**. Primers names and sequences

740 **Supplementary Figure 1.** Bar graphs showing the number of upregulated
741 differentially expressed genes (UpDEGs) and downregulated differentially
742 expressed genes (DownDEGs).

743
744 **Supplementary Figure 2.** Density distributions of (A) raw transcriptome read
745 counts and (B) normalized counts from DESeq2. Data are log2-
746 transformed. Methylated and unmethylated genes exhibit distinctive patterns of
747 expression consistent with previous studies. The density distributions of the raw
748 and normalized counts are highly similar.
749

752 **Supplementary Figure 3.** Violin plots of methylated and unmethylated
753 upregulated differentially expressed genes during the late blastoderm stage in
754 control and *Nv-dnmt1a* pRNAi embryos.

755
756
757
758

759
760

761 **Supplementary Figure 4.** Violin plots of methylated and unmethylated genes
762 during the late blastoderm stage in control and *Nv-dnmt1a* pRNAi embryos
763 broken up into three categories: top, all differentially expressed genes (All

764 DEGs), middle, all genes, and bottom, all non-differentially expressed genes (All
765 nonDEGs).
766
767
768

769
770 **Supplementary Figure 5.** Violin plots of 615 genes that have a similar
771 expression profile between control and *Nv-dnmt1a* pRNAi early blastoderm
772 embryos and a significant difference in their expression in between control and
773 pRNAi middle blastoderm embryos.
774
775
776
777 **Supplementary Movie 1.** Time lapse movie of developing control, mock-
778 injected, embryo. 1 image taken per minute. Anterior is to left.

779 **Supplementary Movie 2.** Time lapse movie of developing *Nv-nonC* knockdown
780 embryo. 1 image taken per minute. Anterior is to left.
781
782 **References**

783 1. Tadros W, Lipshitz HD. The maternal-to-zygotic transition: a play in two
784 acts. *Development* **136**, 3033-3042 (2009).
785
786 2. Vastenhouw NL, Cao WX, Lipshitz HD. The maternal-to-zygotic transition
787 revisited. *Development* **146**, dev161471 (2019).
788
789 3. Tadros W*, et al.* SMAUG is a major regulator of maternal mRNA
790 destabilization in Drosophila and its translation is activated by the PAN GU
791 kinase. *Developmental cell* **12**, 143-155 (2007).
792
793 4. Laver JD*, et al.* Brain tumor is a sequence-specific RNA-binding protein
794 that directs maternal mRNA clearance during the Drosophila maternal-to-
795 zygotic transition. *Genome biology* **16**, 94 (2015).
796
797 5. Fu S, Nien C-Y, Liang H-L, Rushlow C. Co-activation of microRNAs by
798 Zelda is essential for early Drosophila development. *Development* **141**,
799 2108-2118 (2014).
800
801 6. Bushati N, Stark A, Brennecke J, Cohen SM. Temporal reciprocity of
802 miRNAs and their targets during the maternal-to-zygotic transition in
803 Drosophila. *Current Biology* **18**, 501-506 (2008).
804
805 7. Schulz KN*, et al.* Zelda is differentially required for chromatin accessibility,
806 transcription factor binding, and gene expression in the early Drosophila
807 embryo. *Genome research* **25**, 1715-1726 (2015).
808
809 8. Dahanukar A, Walker JA, Wharton RP. Smaug, a novel RNA-binding
810 protein that operates a translational switch in Drosophila. *Molecular cell* **4**,
811 209-218 (1999).
812
813 9. Hamm DC, Harrison MM. Regulatory principles governing the maternal-to-
814 zygotic transition: insights from Drosophila melanogaster. *Open Biol* **8**,
815 (2018).
816
817 10. Urieli-Shoval S, Gruenbaum Y, Sedat J, Razin A. The absence of
818 detectable methylated bases in Drosophila melanogaster DNA. *FEBS*
819 *letters* **146**, 148-152 (1982).
820
821 11. Simpson VJ, Johnson TE, Hammen RF. Caenorhabditis elegans DNA
822 does not contain 5-methylcytosine at any time during development or
823 aging. *Nucleic acids research* **14**, 6711-6719 (1986).

12. Jiang N, Wang L, Chen J, Wang L, Leach L, Luo Z. Conserved and Divergent Patterns of DNA Methylation in Higher Vertebrates. *Genome Biology and Evolution* **6**, 2998-3014 (2014).

13. Deaton AM, Bird A. CpG islands and the regulation of transcription. *Genes & development* **25**, 1010-1022 (2011).

14. Santos F, Hendrich B, Reik W, Dean W. Dynamic reprogramming of DNA methylation in the early mouse embryo. *Developmental biology* **241**, 172-182 (2002).

15. Yartseva V, Giraldez AJ. The Maternal-to-Zygotic Transition During Vertebrate Development: A Model for Reprogramming. *Current topics in developmental biology* **113**, 191-232 (2015).

16. Lyko F. The DNA methyltransferase family: a versatile toolkit for epigenetic regulation. *Nature Reviews Genetics* **19**, 81 (2018).

17. Bewick AJ, Vogel KJ, Moore AJ, Schmitz RJ. Evolution of DNA methylation across insects. *Molecular biology and evolution* **34**, 654-665 (2017).

18. Provataris P, Meusemann K, Niehuis O, Grath S, Misof B. Signatures of DNA methylation across insects suggest reduced DNA methylation levels in Holometabola. *Genome biology and evolution* **10**, 1185-1197 (2018).

19. Ventos-Alfonso A, Ylla G, Montanyes J-C, Belles X. DNA methylation in cockroaches is essential in early embryo development and reduces gene expression noise. *bioRxiv*, (2020).

20. Yi S. Birds do it, bees do it, worms and ciliates do it too: DNA methylation from unexpected corners of the tree of life. *Genome biology* **13**, (2012).

21. Bonasio R*, et al.* Genome-wide and caste-specific DNA methylomes of the ants Camponotus floridanus and Harpegnathos saltator. *Current Biology* **22**, 1755-1764 (2012).

22. Lyko F, Foret S, Kucharski R, Wolf S, Falckenhayn C, Maleszka R. The honey bee epigenomes: differential methylation of brain DNA in queens and workers. *PLoS biology* **8**, (2010).

23. Walsh TK*, et al.* A functional DNA methylation system in the pea aphid, Acyrthosiphon pisum. *Insect Molecular Biology* **19**, 215-228 (2010).

869 24. Bird A. Functions for DNA methylation in vertebrates. In: *Cold Spring*
870 *Harbor Symposia on Quantitative Biology*). Cold Spring Harbor Laboratory
871 Press (1993).
872

873 25. Keller TE, Han P, Yi SV. Evolutionary Transition of Promoter and Gene
874 Body DNA Methylation across Invertebrate–Vertebrate Boundary.
875 *Molecular Biology and Evolution* **33**, 1019-1028 (2016).
876

877 26. Zemach A, McDaniel IE, Silva P, Zilberman D. Genome-wide evolutionary
878 analysis of eukaryotic DNA methylation. *Science* **328**, 916-919 (2010).
879

880 27. Sarda S, Zeng J, Hunt BG, Yi SV. The evolution of invertebrate gene body
881 methylation. *Molecular biology and evolution* **29**, 1907-1916 (2012).
882

883 28. Hellman A, Chess A. Gene body-specific methylation on the active X
884 chromosome. *science* **315**, 1141-1143 (2007).
885

886 29. Maunakea AK*, et al.* Conserved role of intragenic DNA methylation in
887 regulating alternative promoters. *Nature* **466**, 253-257 (2010).
888

889 30. Hunt BG, Glastad KM, Yi SV, Goodisman MAD. The Function of
890 Intragenic DNA Methylation: Insights from Insect Epigenomes. *Integr*
891 *Comp Biol* **53**, 319-328 (2013).
892

893 31. Huh I, Zeng J, Park T, Yi SV. DNA methylation and transcriptional noise.
894 *Epigenetics & Chromatin* **6**, (2013).
895

896 32. Smith ZD*, et al.* DNA methylation dynamics of the human preimplantation
897 embryo. *Nature* **511**, 611-615 (2014).
898

899 33. Li L, Lu X, Dean J. The maternal to zygotic transition in mammals.
900 *Molecular aspects of medicine* **34**, 919-938 (2013).
901

902 34. Guo H*, et al.* The DNA methylation landscape of human early embryos.
903 *Nature* **511**, 606-610 (2014).
904

905 35. Bewick AJ, Sanchez Z, Mckinney EC, Moore AJ, Moore PJ, Schmitz RJ.
906 Dnmt1 is essential for egg production and embryo viability in the large
907 milkweed bug, Oncopeltus fasciatus. *Epigenetics & chromatin* **12**, 1-14
908 (2019).
909

910 36. Wang X*, et al.* Function and Evolution of DNA Methylation in Nasonia
911 vitripennis. *PLoS Genetics* **9**, e1003872 (2013).
912

913   37.   Park J, *et al.* Comparative analyses of DNA methylation and sequence
914   evolution using Nasonia genomes. *Molecular biology and evolution* **28**,
915   3345-3354 (2011).
916
917   38.   Werren JH, Richards S, Desjardins CA, Niehuis O, Gadau J, Colbourne
918   JK. Functional and Evolutionary Insights from the Genomes of Three
919   Parasitoid *Nasonia* Species. *Science* **327**, 343-348 (2010).
920
921   39.   Zwier M, Verhulst E, Zwahlen R, Beukeboom L, Van de Zande L. DNA
922   methylation plays a crucial role during early Nasonia development. *Insect*
923   *molecular biology* **21**, 129-138 (2012).
924
925   40.   Lynch JA, Desplan C. A method for parental RNA interference in the wasp
926   Nasonia vitripennis. *Nat Protoc* **1**, 486-494 (2006).
927
928   41.   Buchta T, Özüak O, Stappert D, Roth S, Lynch JA. Patterning the dorsal–
929   ventral axis of the wasp Nasonia vitripennis. *Developmental biology* **381**,
930   189-202 (2013).
931
932   42.   Arsala D, Lynch JA. Ploidy has little effect on timing early embryonic
933   events in the haplo-diploid wasp Nasonia. *Genesis (New York, NY : 2000)*
934   **55**, (2017).
935
936   43.   Liang H-L, Nien C-Y, Liu H-Y, Metzstein MM, Kirov N, Rushlow C. The
937   zinc-finger protein Zelda is a key activator of the early zygotic genome in
938   Drosophila. *Nature* **456**, 400-403 (2008).
939
940   44.   Laver JD, Marsolais AJ, Smibert CA, Lipshitz HD. Regulation and
941   Function of Maternal Gene Products During the Maternal-to-Zygotic
942   Transition in Drosophila. *Current topics in developmental biology* **113**, 43-
943   84 (2015).
944
945   45.   Arsala D, Wu X, Yi SV, Lynch JA. Supplemental item 1. (2020).
946
947   46.   Wang Y, *et al.* Functional CpG methylation system in a social insect.
948   *Science* **314**, 645-647 (2006).
949
950   47.   Wu X, Lindsey AR, Chatterjee P, Werren JH, Stouthamer R, Soojin VY.
951   Distinct epigenomic and transcriptomic modifications associated with
952   Wolbachia-mediated asexuality. *PLoS pathogens* **16**, e1008397 (2020).
953
954   48.   Chen Z, Smith KR, Batterham P, Robin C. Smg1 nonsense mutations do
955   not abolish nonsense-mediated mRNA decay in Drosophila melanogaster.
956   *Genetics* **171**, 403-406 (2005).
957

958   49.   Pulak R, Anderson P. mRNA surveillance by the Caenorhabditis elegans
959   smg genes. *Genes & development* **7**, 1885-1897 (1993).
960

961   50.   McIlwain DR*, et al.* Smg1 is required for embryogenesis and regulates
962   diverse genes via alternative splicing coupled to nonsense-mediated
963   mRNA decay. *Proceedings of the National Academy of Sciences* **107**,
964   12186-12191 (2010).
965

966   51.   Bhuvanagiri M*, et al.* 5-azacytidine inhibits nonsense-mediated decay in a
967   MYC-dependent fashion. *EMBO molecular medicine* **6**, 1593-1609 (2014).
968

969   52.   Eckersley-Maslin MA, Alda-Catalinas C, Reik W. Dynamics of the
970   epigenetic landscape during the maternal-to-zygotic transition. *Nature*
971   *Reviews Molecular Cell Biology* **19**, 436-450 (2018).
972

973   53.   Dunican DS, Ruzov A, Hackett JA, Meehan RR. xDnmt1 regulates
974   transcriptional silencing in pre-MBT Xenopus embryos independently of its
975   catalytic function. *Development* **135**, 1295-1302 (2008).
976

977   54.   Seymour DK, Gaut BS. Phylogenetic Shifts in Gene Body Methylation
978   Correlate with Gene Expression and Reflect Trait Conservation. *Molecular*
979   *Biology and Evolution* **37**, 31-43 (2020).
980

981   55.   Bewick AJ, Zhang YW, Wendte JM, Zhang XY, Schmitz RJ. Evolutionary
982   and Experimental Loss of Gene Body Methylation and Its Consequence to
983   Gene Expression. *G3-Genes Genom Genet* **9**, 2441-2445 (2019).
984

985   56.   Takuno S, Seymour DK, Gaut BS. The Evolutionary Dynamics of
986   Orthologs That Shift in Gene Body Methylation between Arabidopsis
987   Species. *Molecular Biology and Evolution* **34**, 1479-1491 (2017).
988

989   57.   Ventos-Alfonso A, Ylla G, Belles X. Zelda and the maternal-to-zygotic
990   transition in cockroaches. *The FEBS Journal* **286**, 3206-3221 (2019).
991

992   58.   Washington JT, Cavender KR, Amukamara AU, McKinney EC, Schmitz
993   RJ, Moore PJ. The essential role of Dnmt1 in gametogenesis in the large
994   milkweed bug Oncopeltus fasciatus. *bioRxiv*, 2020.2007.2023.218180
995   (2020).
996

997   59.   Schulz NK*, et al.* Dnmt1 has an essential function despite the absence of
998   CpG DNA methylation in the red flour beetle Tribolium castaneum.
999   *Scientific reports* **8**, 1-10 (2018).
1000

1001   60.   Love MI, Huber W, Anders S. Moderated estimation of fold change and
1002   dispersion for RNA-seq data with DESeq2. *Genome biology* **15**, 550
1003   (2014).

1004
1005   61.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical
1006   and powerful approach to multiple testing. *Journal of the Royal statistical*
1007   *society: series B (Methodological)* **57**, 289-300 (1995).
1008
1009   62.   Hu H*, et al.* In-depth proteome of the hypopharyngeal glands of honeybee
1010   workers reveals highly activated protein and energy metabolism in priming
1011   the secretion of royal jelly. *Molecular & Cellular Proteomics* **18**, 606-621
1012   (2019).
1013
1014   63.   Park Y, Wu H. Differential methylation analysis for BS-seq data under
1015   general experimental design. *Bioinformatics* **32**, 1446-1453 (2016).
1016
1017