Naive human B cells can neutralize SARS-CoV-2 through recognition of its receptor binding domain

Jared Feldman[1][*], Julia Bals[1][*] Kerri St. Denis[1], Evan C. Lam[1], Blake M. Hauser[1], Larance Ronsard[1], Maya Sangesland[1], Thalia Bracamonte Moreno[1], Vintus Okonkwo[1], Nathania Hartojo[1] Alejandro B. Balazs[1], Daniel Lingwood[1][#] and Aaron G. Schmidt[1,2][#]

[1]Ragon Institute of MGH, MIT and Harvard, Cambridge, MA, 02139, USA

[2]Department of Microbiology, Harvard Medical School, Boston, MA 02115, USA

*equal contribution

#corresponding

Key words: germline precursors, B cells, immunity, SARS-CoV-2, coronaviruses

Correspondence:

Daniel Lingwood
Tel:857-268-7180; E-mail: dlingwood@mgh.harvard.edu

Aaron G. Schmidt
Tel: 857-268-7118; E-mail: aschmidt@crystal.harvard.edu

Running title: Isolation of antibody germline precursors that bind and neutralize SARS-CoV-2

44   **ABSTRACT**
45
Exposure to a pathogen elicits an adaptive immune response aimed to control and eradicate. This
initial exposure is imprinted on the immune system, so that a subsequent encounter to the same
pathogen or a variant will result in a memory recall response that is often protective. Interrogating
the naive B cell repertoire in terms of both abundance and specificity to said pathogen may
contribute to an understanding of how to potentially elicit protective responses. Here, we isolated
naive B cells across 8 human donors, targeting the SARS-CoV-2 receptor-binding domain (RBD).
Single B cell sorting, and subsequent sequence analysis, showed diverse gene usage and pairing
with no apparent restriction on complementarity determining region length in either the heavy or
light chains. We show that recombinantly expressed IgGs and Fabs of these germline precursors
bind SARS-CoV-2 RBD. Importantly, a subset of these naive antibodies also bind SARS-CoV, an
emergent variant (501Y.V2) and a potential pandemic (WIV-1) coronavirus. Furthermore, naive
antibodies can also neutralize SARS-CoV-2 pseudoviruses in the absence of any somatic
hypermutation, suggesting that protective immunity to coronaviruses, more broadly, may be
genetically encoded. Future studies aimed at understanding the naive repertoire to other
coronaviruses may ultimately reveal shared specificities that could be leveraged to develop pan-
coronavirus vaccines aimed at priming encoded germline responses.
62
63

MAIN TEXT

Initial exposure to viral antigens by natural infection or vaccination primes an immune response and often establishes an immune memory which can prevent or control future infections. The naive repertoire contains potential B cell receptor (BCR) rearrangements capable of recognizing these antigens, often the surface-exposed glycoproteins. An early step in generating humoral immunity involves activation of these naive B cells through recognition of a cognate antigen (*1*) which in turn can lead to affinity maturation through somatic hypermutation (SHM) and subsequent differentiation (*2*). The initial engagement of the naive repertoire begins this cascade and often coincides with the eventual generation of a protective or neutralizing antibody response (*3, 4*).

For SARS-CoV-2, the etiological agent of COVID-19, the development of a neutralizing antibody response after primary infection or vaccination is associated with protection against reinfection in non-human primates (*5-8*). In humans, the presence of neutralizing antibodies can predict disease severity and survival after primary SARS-CoV-2 infection (*9*) and correlates with protection from secondary infection (*10, 11*). Further, memory B cells induced by natural infection in humans may persist and, in some cases, continue to affinity mature for greater than 6 months after primary infection (*12-15*) providing potentially durable long-term protection. Comparable levels of neutralizing antibody titers were present in convalescent COVID-19 subjects and vaccine recipients (*16-18*) further supporting the role of adaptive immune responses in helping to control and prevent disease severity. Understanding the origins of these protective responses is critical.

Both infection- and vaccine-elicited antibodies target the major envelope glycoprotein, spike, present on the virion surface (*18*). A substantial component of the neutralizing response engages the receptor binding domain (RBD) (*19-24*) and does so by sterically blocking interactions with the viral receptor ACE2 (*25-30*). Isolated RBD-directed monoclonal antibodies derive from diverse heavy- and light-chain variable gene segments suggesting that multiple biochemical solutions for developing RBD-directed antibodies are encoded within the human B-cell repertoire (*19, 21, 24, 31*). Potential immunogenicity of this antigenic site is based on the human naive B cell repertoire, and the overall frequency of naive BCRs that have some level of intrinsic affinity to stimulate their elicitation (*32-35*); in other words, to reliably amplify a protective SARS-CoV-2 response through vaccination or infection, the appropriate naive B cells must be both present (i.e.,

precursor frequency) and activatable (i.e., above an affinity threshold). However, antigen-

specificity of naive B cells is largely undefined. Here, we characterize the human naive B cell

repertoire specific for the SARS-CoV-2 RBD from healthy individuals in order to understand their

relative abundance, affinity, and potential for providing a protective (i.e., neutralizing) response

after natural infection or vaccination.

To measure the reactivity of naive human B cells specific for the SARS-CoV-2 RBD we adapted

an *ex vivo* B cell profiling approach used previously to study epitope-specific naive precursors

targeting neutralizing sites on HIV (*36-38*) and influenza virus surface glycoproteins (*32*). We first

designed a SARS-CoV-2 RBD construct that positions two glycans at residues 475 and 501 to

selectively block binding to ACE2 and the receptor-binding motif (RBM)-directed antibody, B38

(**fig. S1**) (*39*). Using this "ΔRBM" probe, in addition to wildtype SARS-CoV-2 spike, and RBD

probes, we isolated naive (CD19$^+$/IgD$^+$/IgG$^-$) B cells specific to the RBD and, more finely, the

RBM from the peripheral blood of 8 SARS-CoV-2 seronegative human donors (**Fig. 1A** and **fig.

S2**). We defined RBM-specificity as B cells that bound to fluorescently labeled spike and RBD,

but not the ΔRBM probe (**fig. S3A**). Although rare, all 8 donors had detectable populations of

RBM-specific naive B cells (**fig. S3B**). The median frequency of RBM-specific B cells among

total and naive B cells was 0.0021% and 0.0023%, respectively (**Fig. 1B**). Within spike-reactive,

naive cells, the median frequency of RBM-specific B cells was 3.2% (**Fig. 1C**); this potentially

suggests that a large proportion of spike epitopes targeted by naive responses reside outside of the

RBD. The majority of IgD$^+$ RBM-specific B cells were CD27$^-$ (mean frequency ~97%), in

agreement with the naive B cell phenotype (**fig. S3C**).

To understand in more detail the properties of this naive repertoire, we obtained 163 paired heavy-

and light-chain antibody sequences from 5 of the 8 donors (**Fig. 1D** and **fig. S4**). Sequence analysis

showed that all clones were unique with diverse gene usage for both heavy and light chains and

minimal gene pairing preferences (**fig. S4**). These data reflect the polyclonal gene usage observed

in RBD-specific memory B cells sequenced from COVID-19 convalescent individuals (*21, 24, 31*)

and vaccine recipients (*18*), suggesting that a diverse pool of antibody precursors can be activated

upon antigen exposure. In comparing this naive repertoire to gene usage distribution from non-

SARS-CoV-2-specific repertoires (*40*), we observed an increase in mean repertoire frequency of

~20% for IGHV3-9 in 4 out of 5 sequenced donors (**fig. S5A**). Notably, this enrichment of IGHV3-

9 was also observed in isolated memory B cells from convalescent individuals (*41*) and vaccine

recipients (*18*), as well as in expanded IgG$^+$ B cells sequenced from a cohort of COVID-19 subjects

during acute infection (*31*). These expanded clones detected shortly after symptom onset displayed

low levels of somatic hypermutation (SHM) (*31*), suggesting potential IGHV3-9 usage in an early

extrafollicular response in which naive B cells differentiate into short-lived plasma cells (*42*).

Additionally, IGHV3-53 and 3-30 gene segments, over-represented in RBD-specific antibodies

isolated from convalescent subjects (*22, 30, 43*), were recovered from three sequenced donors (13

total clones; ~8.0% of total). The amino acid length of heavy and light chain third

complementarity-determining regions (CDR3) ranged from 8 to 27 (average length ~16) for

HCDR3 and 4 to 13 (average length ~10) for LCDR3 (**Fig. 1E**). These lengths are normally

distributed relative to both unselected human repertoires (*40, 44*) and RBD-specific memory B

cell repertoires (*18, 21, 22, 24*); this is in contrast to antibody precursors targeting the influenza

and HIV receptor binding sites which have strict requirements for length (*45*) or gene usage (*46,*

47). These data suggest that overall HCDR3 length does not restrict precursor frequency and there

appears no inherent bias for CDR3 length conferring RBM-specificity. The majority of obtained

sequences were at germline in both the variable heavy (V_H) and light (V_L) chains. However, despite

sorting B cells with a naive phenotype, some sequences were recovered that deviated from

germline. Specifically, the V_H ranged from 91.4 to 100% identity to germline, with a median of

99.7%; the V_L ranged from 93.6 to 100%, with a median of 99.3% (**Fig. 1F**).

To obtain affinities of the isolated naive antibodies, we cloned and recombinantly expressed 44

IgGs selected to reflect the polyclonal RBD-specific repertoire with representatives from diverse

variable region gene segments (**Table S1**). Additionally, we ensured diversity in terms of HCDR3

length, kappa and lambda usage, as well as representation from all 5 donors. By ELISA, we

identified IgGs with detectable binding to SARS-CoV-2 RBD; we summarize these results for all

antibodies (**Fig. 2**) and parsed by donor (**fig. S6A**). Across 5 donors, 36 (~81%) bound to

monomeric SARS-CoV-2 RBD (**Fig. 2A**) with EC_{50} values ranging from 3.3 to 410 nM and a

mean of 62 nM (**Fig. 2B** and **fig. S6A**). These antibodies included 32 unique variable heavy and

light chain pairings (**Table S1**). We further defined the epitopic region of these IgGs using the

ΔRBM construct and the individual glycan variants, Δ501 and Δ475, both of which independently

block ACE2 cell-surface binding but are on opposite sides of the RBM (**fig. S1E, F**). 11 IgGs had

no detectable ΔRBM binding (e.g., ab079, ab119), while 21 IgGs had reduced ELISA binding

relative to wild-type RBD (**Fig. 2A, B**), reflected in the reduced ΔRBM median EC$_{50}$ values (**Fig.**

2C). We also identified examples of antibodies sensitive to only Δ475 (e.g., ab185) and only Δ501

(e.g., ab007) (**Fig. 2B** and **fig. S7**). Of the binding population, there is no apparent predisposition

for HCDR3 length or light chain pairing (**Fig. 2 D, E**).

We next asked whether these antibodies could recognize the SARS CoV-2 variant 501Y.V2

currently circulating in South Africa (*48*). This variant includes a combination of the three RBD

mutations: K417N, E484K, and N501Y (*48*). We find that 22 antibodies (50%) had detectable

binding for this variant (**Fig. 2B**). Interestingly one antibody, ab033, seemed to be specific to this

variant and did not bind SARS-CoV-2. We additionally tested the cross-reactivity of these

antibodies to both SARS-CoV and WIV-1, related coronaviruses previously circulating and of

pandemic potential (*49*), respectively. While the majority of antibodies (~86%) were specific to

the SARS-CoV-2 RBD, 5 antibodies were also cross-reactive to both SARS-CoV and WIV-1. Of

these cross-reactive antibodies, ab017 and ab114, derives from the same IGHV3-33 and IGVL2-

14 paring but were isolated from different donors, suggesting a shared or public clonotype.

Additionally, we found that ab189 was uniquely sensitive to the glycan at residue 475; this was

surprising given the lack of conservation between of SARS-CoV, CoV-2 and WIV-1 RBMs,

despite using the same ACE2 receptor (*50*). Collectively, these data demonstrate that coronavirus

-specific naive antibodies are present in distinct donors, are of unrestricted gene usage and when

as expressed as IgGs, have detectable affinity to RBDs from the currently administered vaccines,

a circulating variant of concern, and from at least two related viruses.

To obtain binding kinetics independent of avidity effects from bivalent IgGs, 11 antibodies were

selected for expression as Fabs to determine monovalent binding affinity (*K$_{DS}$*) by biolayer

interferometry (BLI). Using monomeric RBD as the analyte, 9 of the 11 Fabs had detectable

binding with *K$_{DS}$* ranging from ~6.5 to ~75 µM; the other two remaining Fabs (ab177, ab185),

gave unreliable affinity measurements (i.e., >100 µM) (**fig. S8**). Notably, all Fabs had

characteristically fast off rates (*k$_{off}$*). This observation is consistent for germline B cells where fast

off-rates are compensated by avidity due to overall BCR surface density (*51*); subsequent affinity

gains via somatic hypermutation often result in slowing of the off-rate and is a canonical

mechanism of improved antigen binding (*52-54*). While the biologically relevant antigen affinities

of naive B cells are uncertain, recent studies have suggested that B cell activation and affinity

maturation is not restricted by immeasurably low affinity BCR interactions (*55-57*), thus the

isolated naive antibodies described here, especially those with demonstrable affinity may very well

be engaged and elicited upon antigen exposure.

We next used a SARS-CoV-2 pseudovirus assay (*9*) to ask whether any of the isolated antibodies

were capable of blocking transduction of target cells. We found that of the 36 RBD-binding

antibodies tested in this assay, 5 had detectable levels of neutralization (~14%) (**Fig. 3A** and **fig.**

S9). These neutralizing antibodies, obtained from multiple donors, have few commonalities with

respect to their gene usages and HCDR3 lengths (**Fig. 3B**). While these naive antibodies were not

as potent compared to the control antibody B38, isolated from a memory B cell (*29*), the

observation, nevertheless, that the naive repertoire contains antibodies that have, at germline, the

potential to neutralize is noteworthy.

Here, we showed that isolated naive antibodies from 5 different donors can bind SARS-CoV-2

RBD and its variant 501Y.V2, a proportion of which are cross-reactive to SARS CoV and WIV-

1. These data suggest that RBM-specific precursors may be represented across a substantial

fraction of individual human naive repertories. This observation is consistent with longitudinal

studies of SARS-CoV-2 infected individuals in which the majority of convalescent individuals

seroconverted with detectable RBD serum antibodies and neutralization titers (*14, 58*). A subset

of these germline antibodies already shows neutralization activity, albeit weakly, against SARS-

CoV-2 pseudoviruses. They therefore have the potential to affinity mature into more potent

neutralizing antibodies after vaccination or natural infection. Indeed, recent studies have

characterized potent RBD-directed neutralizing antibodies with a limited level of somatic

hypermutation (*19, 21, 24, 31, 59*). The naive antibodies characterized here engage epitopes across

the RBM with a range of angles of approach as defined by our glycan variant probes; this is

consistent with other RBD-targeting antibodies characterized by epitope-mapping, deep

mutational scanning and structures (*25, 27, 60*) (**Fig. 4**). Having BCRs within the naive repertoire

that recognize distinct or partially overlapping epitopes across the RBM may be advantageous in

eliciting protective responses to variants of SARS-CoV-2 (*61-64*). Collectively, our observations
described here for SARS-CoV-2 are notably in contrast to other viruses such as HIV, whose
germline precursors targeting its receptor binding site lack demonstrable affinity, are highly
restricted in their gene usages and cannot at germline neutralize the virus (*65-67*). Future
investigations aimed at understanding the naive repertoire with respect to coronaviruses of
pandemic potential may reveal further commonalties in antigen-specific precursors, enabling the
development of pan-coronavirus vaccines aimed at priming specific germline responses of
protective potential.

METHODS

Donor Samples

PBMCs were isolated from blood donors obtained from the MGH blood donor center (8 donors
total). Prior to donating blood, subjects were required to sign a donor attestation/consent statement,
as per hospital requirements, stating ''I give permission for my blood to be used for transfusion to
patients or for research''. The gender and age are not recorded, however eligible donors are of at
least 16 years old and weigh a minimum of 110lbs. All experiments were conducted with MGH
Institutional Biosafety Committee approval (MGH protocol 2014B000035). Isolated PBMCs were
used for B cell enrichment and single cell sorting (described below); plasma was aliquoted and
stored at -80 ºC until further use. Additionally, the control convalescent sera used for ELISA was
obtained under the approved Partners Institutional Review Board (protocol 2020P000895) for use
of patient samples for the development and validation of SARS-CoV-2 diagnostic tests (*9*).

Expression and purification of recombinant CoV Antigens

Plasmids encoding the receptor binding domains (RBDs) were designed based on GenBank
sequences MN975262.1 (SARS-CoV-2), ABD72970.1 (SARS-CoV), and AGZ48828.1 (WIV-1).
Constructs were codon optimized and synthesized by IDT. Glycosylation sites at SARS-CoV-2
RBD residues 501 and/or 475 as well as 501Y.V2 mutations (K417N/E484K/N501Y) were
inserted by QuikChange Mutagenesis (Agilent) following the manufacturer's suggested protocol
and were sequence confirmed (Genwiz). SARS-CoV-2 spike contained a C-terminal foldon
trimerization domain and HRV 3C-cleavable 6xHis and 2xStrep II tags (*68*). All proteins were

transiently expressed in Expi293F cells (ThermoFisher). 5 to 7 days post-transfection, supernatants
were harvested by centrifugation and further purified using immobilized metal affinity
chromatography (IMAC) with cobalt-TALON resin (Takara) followed by Superdex 200 Increase
10/300 GL size exclusion column (GE Healthcare).
253

Expression and purification IgGs and Fabs
IgG and Fab genes for the heavy- and light-chain variable domains were synthesized and codon
optimized by IDT and subcloned into pVRC protein expression vectors and sequence confirmed
(Genewiz). Fabs and IgGs were similarly expressed and purified as described above for RBDs.
IgGs were buffer exchanged into PBS while Fabs were concentrated and further purified by
Superdex 200 Increase 10/300 GL size exclusion column.
260

ELISA
Both sera and monoclonal antibody reactivity to CoV antigens were assayed by ELISA. Briefly,
96-well plates (Corning) were coated with 5 µg/ml of monomeric RBDs diluted in PBS and
incubated overnight at 4ºC. Plates were blocked with 1% BSA in PBS containing 1% Tween-20
(PBS-T) for 1hr at room temperature (RT). Blocking solution was discarded and 4-fold serial
dilutions of human plasma (1:20 starting dilution) or isolated monoclonal antibodies (150 µg/ml
starting concentration) in PBS were added to wells and incubated for 1hr at RT. Plates were then
washed three times with PBS-T. Secondary, anti-human IgG-HRP (Abcam), was added to each
well at 1:20,000 dilution in PBS-T and incubated for 1hr at RT. Plates were then washed three
times with PBS-T and developed with 1-Step ABTS substrate (ThermoFisher) per manufacturer
recommendations. Absorbance was measured using a plate reader at 405nm. EC_{50} values were
determined for monoclonal antibodies by non-linear regression (sigmoidal) using GraphPad Prism
8.4.3 software. (GraphPad Software). ELISAs against 501Y.V2, SARS-CoV and WIV-1 RBDs
were done at a single IgG concentration (150 µg/ml) in replicate. Positive binding was defined by
an $OD_{405} \geq 0.30$.

ACE-2 cell binding assay

ACE-2 expressing 293T cells were incubated with 200 nM of RBD antigen in PBS for 1hr on ice. Cells were resuspended in 50µL of secondary stain containing streptavidin-PE (Invitrogen) at a 1:200 dilution and incubated for 30 min on ice. Cell binding was analyzed by flow cytometry using a Stratedigm S1300Exi Flow Cytometer equipped with a 96 well plate high throughput sampler. Rsulting data were analyzed using FlowJo (10.7.1).

Probe Generation

SARS-CoV-2 RBD and ΔRBM constructs were expressed as dimeric murine-Fc (mFc; IgG1) fusion proteins containing a HRV 3C-cleavable C-terminal 8xHis and SBP tags and purified as described above. SBP-tagged RBD- and ΔRBM-mFc dimers were individually mixed with fluorescently labeled streptavidin, SA-BV650 and SA-BV786 (BioLegend), to form RBD-mFc-BV650 and ΔRBM-mFc-BV786 tetramers. SARS-CoV-2 spike with a C-terminal Strep II tag was labeled separately with StrepTactin PE and APC (IBA) to form spike-PE and -APC tetramers, respectively. Both labeling steps were performed for 30 min at 4 ºC prior to sorting.

Single B Cell Sorting

Naive B cells were purified from PBMCs using the MACS Human B Cell isolation kit (Miltenyi Biotec) and incubated with 25nM of each SARS-CoV-2 probe (RBD-mFc-BV650, ΔRBM-mFc-BV786, spike-PE, and spike-APC) for 30 min at 4ºC. Cells were stained with anti-human CD19 (Alexa-700), CD3 (PerCP-Cy5), IgD (PE-Cy7), IgG (BV711), CD27 (BV510), LiveDead Violet (Invitrogen), and Calcien (Invitrogen) for an additional 30 min. RBM-specific naive B cells, defined as $CD19^+/CD3^-/IgG^-/IgD^+/$spike $PE^+/$spike $APC^+/RBD^+/\Delta RBM^{--}$, were single-cell sorted using BD FACS Aria II (BD Biosciences) into 96-well plates containing lysis buffer supplemented with 1% BME. Within the $CD19^+/IgG^-/IgD^+$ gated cells, we also confirmed that 97% of the events were CD27 negative. Plates were stored at -80 ºC for subsequent analysis. Flow cytometry data was analyzed using FlowJo software version 10.7.1.

BCR Sequencing

BCR Sequencing was carried out as described previously (*32*). Briefly, whole transcriptome
amplification (WTA) was performed on the sorted cell-lysates according to the Smart-Seq2
protocol (*69*). We then amplified heavy and light chain sequences from the WTA products utilizing
pools of partially degenerate pools of V region specific primers (Qiagen HotStar Taq Plus). Heavy
and light chain amplifications were carried out separately, with each pool containing pooled
primers against human IGHV and heavy chain constant region genes, or human IGLV, IGKV, and
light chain constant region genes. Cellular barcodes and index adapters (based on Nextera XT
Index Adapters, Illumina Inc.) were added using a step-out PCR method. Amplicons were then
pooled and sequenced using a 250x250 paired end 8x8 index reads on an Illumina Miseq System.
The data were then demultiplexed, heavy and light chain reads were paired, and overlapping
sequence reads were obtained (Panda-Seq) (*70*) and aligned against the human IMGT database
(*71*).

Interferometry binding experiments
Interferometry experiments were performed using a BLItz instrument (ForteBio). Fabs (0.1 mg/ml)
were immobilized on Ni-NTA biosensors. The SARS-CoV-2 RBD analyte was titrated (10µM,
5µM, 2.5µM, and 1µM) to acquire binding affinities; the K_D was obtained through global fit of the
titration curves by applying a 1:1 binding isotherm using vendor-supplied software.

Pseudotyped Neutralization Assay
SARS-CoV-2 neutralization was assessed using lentiviral particles pseudotyped as previously
described (*9, 72*). Briefly, lentiviral particles were produced via transient transfection of 293T
cells. The titers of viral supernatants were determined via flow cytometry on 293T-ACE2 cells
(*72*) and via the HIV-1 p24[CA] antigen capture assay (Leidos Biomedical Research, Inc.). Assays
were performed in 384-well plates (Grenier) using a Fluent Automated Workstation (Tecan). IgGs
starting at 150 µg/ml, were serially diluted (3-fold) in 20µL followed by addition of 20 µL of
pseudovirus containing 250 infectious units and incubated at room temperature for 1 hr. Finally,
10,000 293T-ACE2 cells (*72*) in 20 µL cell media containing 15 µg/ml polybrene were added to
each well and incubated at 37 °C for 60-72 hrs. Following transduction, cells were lysed using a
previously described assay buffer (*73*) and shaken for 5 min prior to quantitation of luciferase
expression using a Spectramax L luminometer (Molecular Devices). Percent neutralization was

determined by subtracting background luminescence measured from cells control wells (cells only)

from sample wells and dividing by virus control wells (virus and cells only). Data were analyzed

using Graphpad Prism.

Acknowledgements

We thank members of the Schmidt and Lingwood Labs for helpful discussions, especially Timothy

Caradonna and Daniel Maurer. We thank Goran Bajic for critical reading of the manuscript. We

thank Samuel Kazer, James Gatter and Alex Shalek for BCR sequencing advice, Jason McLellan

for the SARS-CoV-2 spike plasmid, and Nir Hacohen and Michael Farzan for ACE-2 expressing

293T cells. We acknowledge support from NIH (R01AI146779, R01AI124378, R01AI137057,

R01AI153098, R01AI155447, DP2DA042422, DP2DA040254, T32 AI007245), a Massachusetts

Consortium on Pathogenesis Readiness (MassCPR) grant to A.G.S. and a MGH Transformative

Scholars Program and Charles H. Hood Foundation to A.B.B.

Author Information

Author Contributions J.F., J.B., A.B.B., D.L. A.G.S. designed research; J.F., J.B., K.S.D., E.C.L,

B.M.H., L.R. M.S. T.B.M.,V.O., N.H., performed research; J.F., J.B., D.L. A.G.S analyzed data;

J.F. and A.G.S. wrote the paper. J.F., J.B., B.M.H, A.B.B., D.L. A.G.S edited and commented on

the paper.

Correspondence and requests for materials should be addressed to: Daniel Lingwood

(dlingwood@mgh.harvard.edu) or Aaron G. Schmidt (aschmidt@crystal.harvard.edu).

Competing financial interest: No competing financial interests.

FIGURES

Fig. 1. SARS-CoV-2-specific naive B cells isolation and characterization. (**A**) RBM-specific naive B cells from seronegative human donors were isolated by fluorescence-activated cell sorting gated on CD19⁺IgD⁺IgG⁻; representative plot from donors 1 and 2 is shown. ΔRBM is a sorting probe with N-linked glycans at residues 501 and 475. RBM-specific B cell frequency among (**B**) total, naive, and (**C**) spike-positive cells from each donor ($n = 8$). (**D**) Heat map showing variable-gene usage for all paired B cell sequences. Scale indicates percent of total sequences for each donor separately. (**E**) Heavy (H) and light (L) CDR3 amino acid length distribution determined using IMGT numbering. Red bars indicate median amino acid length. (**F**) Divergence from inferred germline gene sequences. Red bars indicate the median percent values.

Fig. 2. Binding properties and specificity of isolated naive antibodies. (**A**) ELISA binding of 44 IgGs to SARS-CoV-2 RBD and ΔRBM binding (left and right panels, respectively). 36 IgGs with reduced ΔRBM binding were defined by >1.5 fold-change in ELISA EC_{50} relative to wild-type RBD. (**B**) IgG binding to SARS-CoV-2 (wildtype and 501Y.V2), CoV-1, WIV-1 RBDs and individual RBD glycan variants. (**C**) ELISA EC_{50} values for IgGs with detectable SARS-CoV-2 RBD binding (n = 36). Red bars indicate the mean EC_{50} values. (**D**) Pearson correlation analysis of SARS-CoV-2 RBD affinities and HCDR3 length. (**E**) ELISA EC_{50}s for IgGs based on kappa or lambda gene usage.

Fig. 3. Naive antibodies neutralize SARS-CoV-2 (**A**) SARS-CoV-2 pseudovirus neutralization assay for 5 IgGs. The neutralizing monoclonal antibody, B38, was used as a positive control. Dashed lines indicate IC_{50} values and data represent means $\pm$ SD of two technical replicates. (**B**) Variable region gene usage and CDR3 sequences for the IgGs.

Fig. 4. SARS-CoV-2 RBD-specific naive antibodies and their epitope specificity. (**A**) Epitopic Region I defined by loss of binding to Δ475; structure of RBM-directed S2E12 (PDB 7K4N) and representative naive IgGs (italics) in this study shown. (**B**) Epitopic Region II, defined by loss of binding to ΔRBM but neither single glycan variant alone; structure of RBM-directed B38 (PDB 7BZ5) and representative naive IgGs. (**C**) Epitopic Region III, defined by loss of binding to Δ501 with representative naive IgGs (no structure is presently available to define this region). (**D**) Summary of gene usage of isolated naive antibodies from each Region (**A-C**); bars represent total number of antibodies and color-coded by each Region.

REFERENCES

1. N. E. Harwood, F. D. Batista, Early events in B cell activation. *Annual review of immunology* **28**, 185-210 (2009).

2. G. D. Victora, M. C. Nussenzweig, Germinal centers. *Annual review of immunology* **30**, 429-457 (2012).

3. C. Havenar-Daughton, R. K. Abbott, W. R. Schief, S. Crotty, When designing vaccines, consider the starting material: the human B cell repertoire. *Current opinion in immunology* **53**, 209-216 (2018).

4. F. Sallusto, A. Lanzavecchia, K. Araki, R. Ahmed, From vaccines to memory and back. *Immunity* **33**, 451-463 (2010).

5. W. Deng *et al.*, Primary exposure to SARS-CoV-2 protects against reinfection in rhesus macaques. *Science* **369**, 818-823 (2020).

6. A. Chandrashekar *et al.*, SARS-CoV-2 infection protects against rechallenge in rhesus macaques. *Science*, (2020).

7. N. B. Mercado *et al.*, Single-shot Ad26 vaccine protects against SARS-CoV-2 in rhesus macaques. *Nature* **586**, 583-588 (2020).

8. Q. Gao *et al.*, Development of an inactivated vaccine candidate for SARS-CoV-2. *Science* **369**, 77-81 (2020).

9. W. F. Garcia-Beltran *et al.*, COVID-19 neutralizing antibodies predict disease severity and survival. *Cell*, (2020).

10. A. Addetia *et al.*, Neutralizing Antibodies Correlate with Protection from SARS-CoV-2 in Humans during a Fishery Vessel Outbreak with a High Attack Rate. *Journal of Clinical Microbiology* **58**, e02107-02120 (2020).

11. S. F. Lumley *et al.*, Antibodies to SARS-CoV-2 are associated with protection against reinfection. *medRxiv*, 2020.2011.2018.20234369 (2020).

12. A. Sokal *et al.*, Maturation and persistence of the anti-SARS-CoV-2 memory B cell response. *bioRxiv*, 2020.2011.2017.385252 (2020).

13. C. Gaebler *et al.*, Evolution of antibody immunity to SARS-CoV-2. *Nature*, (2021).

14. J. M. Dan *et al.*, Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection. *Science*, eabf4063 (2021).

15. L. B. Rodda *et al.*, Functional SARS-CoV-2-specific immune memory persists after mild COVID-19. *Cell*, (2020).

16. E. J. Anderson *et al.*, Safety and immunogenicity of SARS-CoV-2 mRNA-1273 vaccine in older adults. *New England Journal of Medicine*, (2020).

17. E. E. Walsh *et al.*, Safety and Immunogenicity of Two RNA-Based Covid-19 Vaccine Candidates. *New England Journal of Medicine* **383**, 2439-2450 (2020).

18. Z. Wang *et al.*, mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants. *bioRxiv*, 2021.2001.2015.426911 (2021).

19. P. J. M. Brouwer *et al.*, Potent neutralizing antibodies from COVID-19 patients define multiple targets of vulnerability. *Science* **369**, 643-650 (2020).

20. B. Ju *et al.*, Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature* **584**, 115-119 (2020).

21. C. Kreer *et al.*, Longitudinal isolation of potent near-germline SARS-CoV-2-neutralizing antibodies from COVID-19 patients. *Cell* **182**, 843-854. e812 (2020).

22. D. F. Robbiani *et al.*, Convergent antibody responses to SARS-CoV-2 in convalescent individuals. *Nature* **584**, 437-442 (2020).

23. A. Z. Wec *et al.*, Broad neutralization of SARS-related viruses by human monoclonal antibodies. *Science* **369**, 731-736 (2020).

24. S. J. Zost *et al.*, Rapid isolation and profiling of a diverse panel of human monoclonal antibodies targeting the SARS-CoV-2 spike protein. *Nature Medicine* **26**, 1422-1427 (2020).

25.   C. O. Barnes *et al.*, SARS-CoV-2 neutralizing antibody structures inform therapeutic strategies.
Nature, (2020).
26.   C. O. Barnes *et al.*, Structures of Human Antibodies Bound to SARS-CoV-2 Spike Reveal
Common Epitopes and Recurrent Features of Antibodies. *Cell* **182**, 828-842.e816 (2020).
27.   L. Piccoli *et al.*, Mapping Neutralizing and Immunodominant Sites on the SARS-CoV-2 Spike
Receptor-Binding Domain by Structure-Guided High-Resolution Serology. *Cell* **183**, 1024-
1042.e1021 (2020).
28.   R. Shi *et al.*, A human neutralizing antibody targets the receptor binding site of SARS-CoV-2.
Nature, 1-8 (2020).
29.   Y. Wu *et al.*, A noncompeting pair of human neutralizing antibodies block COVID-19 virus binding
to its receptor ACE2. *Science* **368**, 1274-1278 (2020).
30.   M. Yuan *et al.*, Structural basis of a shared antibody response to SARS-CoV-2. *Science* **369**, 1119-
1123 (2020).
31.   S. C. A. Nielsen *et al.*, Human B Cell Clonal Expansion and Convergent Antibody Responses to
SARS-CoV-2. *Cell Host & Microbe* **28**, 516-525.e515 (2020).
32.   M. Sangesland *et al.*, Germline-encoded affinity for cognate antigen enables vaccine amplification
of a human broadly neutralizing response against influenza virus. *Immunity* **51**, 735-749. e738
(2019).
33.   P. Dosenovic *et al.*, Anti–HIV-1 B cell responses are dependent on B cell precursor frequency and
antigen-binding affinity. *Proceedings of the National Academy of Sciences*, 201803457 (2018).
34.   R. K. Abbott *et al.*, Precursor frequency and affinity determine B cell competitive fitness in
germinal centers, tested with germline-targeting HIV vaccine immunogens. *Immunity* **48**, 133-146.
e136 (2018).
35.   A. Amitai *et al.*, Defining and Manipulating B cell Immunodominance Hierarchies to Elicit Broadly
Neutralizing Antibody Responses Against Influenza Virus. *Cell Systems* **11**, 573-588. e579 (2020).
36.   C. Havenar-Daughton *et al.*, The human naive B cell repertoire contains distinct subclasses for a
germline-targeting HIV-1 vaccine immunogen. *Science translational medicine* **10**, eaat0381
(2018).
37.   J. G. Jardine *et al.*, HIV-1 broadly neutralizing antibody precursor B cells revealed by germline-
targeting immunogen. *Science* **351**, 1458-1463 (2016).
38.   J. M. Steichen *et al.*, A generalized HIV vaccine design strategy for priming of broadly neutralizing
antibody responses. *Science* **366**, (2019).
39.   B. M. Hauser *et al.*, Engineered receptor binding domain immunogens elicit pan-coronavirus
neutralizing antibodies. *bioRxiv*, 2020.2012.2007.415216 (2020).
40.   B. Briney, A. Inderbitzin, C. Joyce, D. R. Burton, Commonality despite exceptional diversity in
the baseline human antibody repertoire. *Nature* **566**, 393-397 (2019).
41.   B. Ju *et al.*, Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature* **584**, 115-
119 (2020).
42.   I. C. MacLennan *et al.*, Extrafollicular antibody responses. *Immunological reviews* **194**, 8-18
(2003).
43.   Y. Cao *et al.*, Potent Neutralizing Antibodies against SARS-CoV-2 Identified by High-Throughput
Single-Cell Sequencing of Convalescent Patients' B Cells. *Cell* **182**, 73-84.e16 (2020).
44.   C. Soto *et al.*, High frequency of shared clonotypes in human B cell receptor repertoires. *Nature*
566, 398-402 (2019).
45.   A. G. Schmidt *et al.*, Viral receptor-binding site antibodies with diverse germline origins. *Cell* **161**,
1026-1034 (2015).
46.   T. Zhou *et al.*, Multidonor analysis reveals structural elements, genetic determinants, and
maturation pathway for HIV-1 neutralization by VRC01-class antibodies. *Immunity* **39**, 245-258
(2013).

47. A. P. West, R. Diskin, M. C. Nussenzweig, P. J. Bjorkman, Structural basis for germ-line gene
usage of a potent class of antibodies targeting the CD4-binding site of HIV-1 gp120. *Proceedings*
of the National Academy of Sciences **109**, E2083-E2090 (2012).
48. H. Tegally *et al.*, Emergence and rapid spread of a new severe acute respiratory syndrome-related
coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa. *medRxiv*,
2020.2012.2021.20248640 (2020).
49. V. D. Menachery *et al.*, SARS-like WIV1-CoV poised for human emergence. *Proceedings of the*
National Academy of Sciences **113**, 3048-3053 (2016).
50. C. G. Rappazzo *et al.*, Broad and potent activity against SARS-like viruses by an engineered human
monoclonal antibody. *Science*, eabf4830 (2021).
51. D. Lingwood *et al.*, Structural and genetic basis for development of broadly neutralizing influenza
antibodies. *Nature* **489**, 566 (2012).
52. A. G. Schmidt *et al.*, Immunogenic stimulus for germline precursors of antibodies that engage the
influenza hemagglutinin receptor-binding site. *Cell reports* **13**, 2842-2850 (2015).
53. F. D. Batista, M. S. Neuberger, Affinity dependence of the B cell response to antigen: a threshold,
a ceiling, and the importance of off-rate. *Immunity* **8**, 751-759 (1998).
54. J. M. Tas *et al.*, Visualizing antibody affinity maturation in germinal centers. *Science* **351**, 1048-
1054 (2016).
55. C. Viant *et al.*, Antibody Affinity Shapes the Choice between Memory and Germinal Center B Cell
Fates. *Cell* **183**, 1298-1311. e1211 (2020).
56. R. Di Niro *et al.*, Salmonella infection drives promiscuous B cell activation followed by
extrafollicular affinity maturation. *Immunity* **43**, 120-131 (2015).
57. M. Kuraoka *et al.*, Complex antigens drive permissive clonal selection in germinal centers.
Immunity **44**, 542-552 (2016).
58. D. Stadlbauer *et al.*, Repeated cross-sectional sero-monitoring of SARS-CoV-2 in New York City.
Nature, 1-5 (2020).
59. S. I. Kim *et al.*, Stereotypic neutralizing VH antibodies against SARS-CoV-2 spike protein receptor
binding domain in COVID-19 patients and healthy individuals. *Science Translational Medicine*,
(2021).
60. A. J. Greaney *et al.*, Complete Mapping of Mutations to the SARS-CoV-2 Spike Receptor-Binding
Domain that Escape Antibody Recognition. *Cell Host & Microbe*, (2020).
61. A. J. Greaney *et al.*, Comprehensive mapping of mutations to the SARS-CoV-2 receptor-binding
domain that affect recognition by polyclonal human serum antibodies. *bioRxiv*, 2020.2012.
2031.425021 (2021).
62. A. S. Lauring, E. B. Hodcroft, Genetic Variants of SARS-CoV-2—What Do They Mean? *JAMA*,
(2021).
63. N. G. Davies *et al.*, Estimated transmissibility and severity of novel SARS-CoV-2 Variant of
Concern 202012/01 in England. *medRxiv*, (2020).
64. H. Tegally *et al.*, Emergence and rapid spread of a new severe acute respiratory syndrome-related
coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa. *medRxiv*,
(2020).
65. S. Hoot *et al.*, Recombinant HIV envelope proteins fail to engage germline versions of anti-CD4bs
bNAbs. *PLoS Pathog* **9**, e1003106 (2013).
66. J. F. Scheid *et al.*, Sequence and structural convergence of broad and potent HIV antibodies that
mimic CD4 binding. *Science* **333**, 1633-1637 (2011).
67. T. Zhou *et al.*, Structural basis for broad and potent neutralization of HIV-1 by antibody VRC01.
Science **329**, 811-817 (2010).
68. D. Wrapp *et al.*, Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science*
367, 1260-1263 (2020).
69. J. J. Trombetta *et al.*, Preparation of single-cell RNA-seq libraries for next generation sequencing.
Current protocols in molecular biology **107**, 4.22. 21-24.22. 17 (2014).

70.   A. P. Masella, A. K. Bartram, J. M. Truszkowski, D. G. Brown, J. D. Neufeld, PANDAseq: paired-
end assembler for illumina sequences. *BMC bioinformatics* **13**, 1-7 (2012).
71.   B. Shi *et al.*, Comparative analysis of human and mouse immunoglobulin variable heavy regions
from IMGT/LIGM-DB with IMGT/HighV-QUEST. *Theoretical Biology and Medical Modelling*
11, 1-11 (2014).
72.   M. J. Moore *et al.*, Retroviruses pseudotyped with the severe acute respiratory syndrome
coronavirus spike protein efficiently infect cells expressing angiotensin-converting enzyme 2.
Journal of virology **78**, 10628-10635 (2004).
73.   E. Siebring-van Olst *et al.*, Affordable luciferase reporter assay for cell-based high-throughput
screening. *Journal of biomolecular screening* **18**, 453-461 (2013).
575

Supplementary Information

Naive human B cells can neutralize SARS-CoV-2 through recognition of its receptor binding domain

Jared Feldman[1*], Julia Bals[1*] Kerri St. Denis[1], Evan C. Lam[1], Blake M. Hauser[1], Larance Ronsard[1], Maya Sangesland[1], Thalia Bracamonte Moreno[1], Vintus Okonkowo[1], Nathania Hartojo[1] Alejandro B. Balazs[1], Daniel Lingwood[1#] and Aaron G. Schmidt[1,2#]

[1]Ragon Institute of MGH, MIT and Harvard, Cambridge, MA, 02139, USA

[2]Department of Microbiology, Harvard Medical School, Boston, MA 02115, USA

fig. S1. Design and characterization of SARS-CoV-2 antigens. (A) SARS-CoV-2 RBD in

complex with viral receptor, ACE2 shown in blue and grey, respectively (PDB 6M0J). Wild-type

RBD with, the receptor binding motif (RBM), shown in orange (left panel). Structural model of

the ΔRBM probe designed to abrogate binding to ACE2 (right panel). Putative N-linked

glycosylation sites engineered onto the RBM are shown in red spheres at amino acid positions 501

and 475. (**B**) SDS-PAGE gel under reducing (R) and non-reducing (NR) conditions for monomeric

RBD, RBD-Fc and ΔRBM-Fc. (**C**) Wildtype RBD, ΔRBM and single glycan variant binding to

ACE2-expressing 293T cells by flow cytometry. Wild-type RBD binding shown in blue, glycan

variant binding shown in red. Streptavidin-PE was used to detect the relative intensity of antigen

binding to cell-surface ACE2. A PBS control (gray) indicates secondary-only staining. (**D**) Control

antibody ELISA binding to RBD and ΔRBM antigens. RBM-specific antibody, B38 (left). Non-

RBM-specific control antibody, CR3022 (right). (**E**) ΔRBM and Δ501 and Δ475 variants analyzed

by SDS-PAGE gel under reducing conditions; wildtype RBD is shown for comparison.

fig. S2. **ELISA binding reactivity of healthy donor sera to SARS-CoV-2 spike and RBD.**

Seronegative donors 1-8 were single-cell B sorted to isolate RBD-specific naive B cells. Sera from

a COVID-19 convalescent patient and control antibody, B38, were included as positive controls.

fig. S3. **PBMC flow cytometry analyses.** (**A**) Representative gating strategy used for FACS of

PBMCs pooled from donors 1 and 2. Gating was on naive B cells defined by single living

lymphocytes that were CD19$^+$CD3$^-$IgD$^+$IgG$^-$. Sorted cells were RBM-specific as defined by spike-

PE$^+$/spike-APC$^+$/RBD-Fc-BV650$^+$/ΔRBM-Fc-BC650-. Sort gate is denoted by the blue arrow. The

bottom right plot shows CD27 staining of sorted RBM-specific naive B cells. (**B**) Flow cytometry

showing the sort gate and percentage of RBM-specific B cells for the remaining 6 healthy human

donors. (**C**) RBM-specific B cell frequency among CD27$^+$ and CD27$^-$ cells. Each symbol

represents a different donor (n = 8).

fig. S4. Variable-gene usage of isolated naive antibodies. (**A**) Heatmap showing frequency of $V_H:V_L$ pairing. Overall frequency of variable (V) and joining (J) genes for (**B**) heavy chains, (**C**) kappa light chains, and (**D**) lambda light chains. Heatmap scales represent percent of total paired sequences.

fig. S5. **Repertoire comparison and germline identity by individual donor**. (**A**) Heatmap

showing V_H-gene usage of isolated antibodies derived from donors 1-5. Unselected repertoire gene

usage derive from a high-throughput sequencing data set of circulating B cells across 10 human

subjects (*1*). Heatmap scale represent percent of total paired sequence from each donor.

Divergence from inferred germline gene sequences separated by individual donor for (**B**) V_H and

(**C**) V_L. Red bars indicate the median percent values, and each dot represents an individual paired

sequence.

fig. S6. **ELISA specificity for isolated naive antibodies related to Figure 2**. Heatmap showing

IgG binding to RBDs (n = 44) sorted by donor.

fig. S7. **RBD glycan variant binding by ELISA**. Representative examples of epitopic regions as defined by patterns of RBD glycan variant binding by ELISA. Region 1 refers to IgGs with reduced binding to Δ475, region 2 refers to IgGs with reduced binding to both glycan variants, and region 3 refers to IgGs with reduced binding to Δ501.

fig. S8. **SARS-CoV-2 RBD-binding kinetics of isolated naive antibodies.** (**A**) Biolayer interferometry (BLI) binding kinetic analysis of titrated SARS-CoV-2 RBD to immobilized Fabs. Dotted line at 60 s denotes the start of the dissociation phase. (**B**) Kinetic and equilibrium constants for binding to RBD calculated from a 1:1 binding model using a global fit to all curves for each Fab using vendor supplied software. B38 Fab is used as a positive control.

fig. S9. **SARS-CoV-2 pseudotyped neutralization assay**. (**A**) Neutralization data for 44 purified

IgGs. Curves in color highlighted antibodies with neutralizing activity with (**B**) donor and

monovalent RBD affinity listed for this subset of antibodies. Dashed lines indicate IC_{50} values and

data represent means ± SD of two technical replicates.

SI REFERENCES

1.　B. Briney, A. Inderbitzin, C. Joyce, D. R. Burton, Commonality despite exceptional diversity
in the baseline human antibody repertoire. *Nature* **566**, 393–397 (2019).