Tig1 regulates proximo-distal identity during salamander limb regeneration

Catarina R. Oliveira[1], Dunja Knapp[1], Ahmed Elewa[2], Sandra G. Gonzalez Malagon[3,4], Phillip B. Gates[3], Andreas Petzhold[1], Hernan Arce[5,6], Rodrigo C. Cordoba[5] , Osvaldo Chara[5,6,7], Elly M. Tanaka[8], Andras Simon[2], Maximina H. Yun[1,9]*

Affiliations:

[1] Technische Universität Dresden, CRTD/Center for Regenerative Therapies Dresden, Dresden, Germany

[2] Department of Cell and Molecular Biology, Karolinska Institute, Stockholm, Sweden

[3] Institute of Structural and Molecular Biology, University College London, London, UK

[4] Department of Biomedical Research, Institute of Molecular Biology and Biotechnology/FORTH, Ioannina, Greece

[5] Systems Biology Group, Institute of Physics of Liquids and Biological Systems, National Scientific and Technical Research Council, University of La Plata, La Plata, Argentina

[6] Institute of Technology, Universidad Argentina de la Empresa, Buenos Aires, Argentina

[7] Technische Universität Dresden, Center for Information Services and High Performance Computing ZIH, Dresden, Germany

[8] Institute of Molecular Pathology, Vienna Biocenter, Vienna, Austria

[9] Max Planck Institute for Molecular Cell Biology and Genetics, Dresden, Germany

*Correspondence to: maximina.yun@tu-dresden.de

Abstract:

Salamander limb regeneration is an accurate process which gives rise exclusively to the missing structures, irrespective of the amputation level. This suggests that cells in the stump have an awareness of their spatial location, a property termed 'positional identity'. Little is known about how positional identity is encoded, in salamanders or other biological systems. Through single-cell RNAseq analysis, we identified Tig1/RARRES1 as a potential determinant of proximal identity. Tig1 encodes a conserved cell surface molecule, is regulated by retinoic acid and exhibits a graded expression along the proximo-distal axis of the limb. Its overexpression leads to regeneration defects in the distal elements and elicits proximal displacement of blastema cells, while its neutralisation blocks proximo-distal cell surface interactions. Critically, Tig1 reprogrammes distal cells to a proximal identity, upregulating Prod1 and inhibiting HoxA13 and distal transcriptional networks. Thus, Tig1 is a central cell surface determinant of proximal identity in the salamander limb.

Main Text:

Introduction:
The mechanisms by which cells determine their position within a biological structure are central to the correct development and regeneration of tissues and organs. Despite their importance, they remain poorly understood. Significant insights into this problem originate from the study of salamander limb regeneration, a powerful model for probing the mechanisms underlying positional information[1-3]. In salamanders, the amputation of a limb elicits the generation of a mass of progenitor cells -named blastema- which subsequently grows and re-patterns, invariably giving rise to the structures distal to its level of origin[4]. This indicates that mature cells within the limb stump retain a memory of their location along the proximo-distal (PD) axis which blastema cells are able to interpret in order to regenerate only the missing parts, a property termed positional identity[5]. Elegant blastema juxtaposition experiments offered first clues into the nature of PD identity, suggesting that it is encoded as a proximo-distal gradient of molecules along the limb and manifested at the cell surface level. First, grafting of a distal blastema onto the site of a proximal one leads to the displacement of the distal blastema to its level of origin along the PD axis as regeneration proceeds[6], indicating an initial recognition of PD disparity at the cell surface. Second, apposing proximal and distal blastemas in culture results in the engulfment of the distal blastema by the proximal one, indicating the existence of differential adhesive properties between cells of different PD identities and therefore underscoring the importance of cell surface interactions[7]. Third, grafting of a wrist blastema onto a shoulder stump elicits regenerative growth from the shoulder stump, while wrist blastema cells contribute to the hand structures[8]. Similarly, transplantation of distal blastema cells to proximal blastemas results in translocation of distal cells and their contribution to hand elements, indicating that PD identity is manifested on an individual cell basis and is a stable feature of cell identity[9]. Additional clues into the nature of PD identity came from classical experiments addressing the effect of retinoic acid (RA) and its derivatives on limb regeneration. Notably, exposing a wrist blastema to increasing doses of retinoic acid leads to regeneration of more proximal elements in a dose-dependent manner, resulting in tandem limb duplications at the highest retinoic acid concentrations[5,10]. Thus, retinoic acid is able to reprogram positional identity towards proximal values.

From these seminal findings, a set of principles for identifying a mediator of proximal identity was proposed[11]: it should exhibit a graded expression along the PD axis, be present at the cell surface, and be upregulated by retinoic acid. This led to the identification of the GPI-anchored cell-surface molecule Prod1[11], which fulfils all said criteria. Indeed, Prod1 was shown to mediate the engulfment activity of proximal blastemas when juxtaposed to distal ones and thus cell affinity, an important aspect of proximal identity[11]. Subsequently, Prod1 overexpression in early blastemas was shown to affect patterning of distal elements during limb regeneration and, notably, its focal overexpression in cells of distal domains led to their translocation to proximal positions within the limb[9]. Further, Prod1 expression was shown to be regulated by Meis homeobox transcription factors[12], critical targets of retinoic acid during proximo-distal patterning of the vertebrate limb[13]. Altogether, these experiments associated Prod1 to the acquisition of proximal identity. Given its PD distribution along the mature limb, it was proposed that Prod1 is part of the positional memory system that confers mature limb cells their spatial identity and determines the extent of regeneration[14]. As such a system would determine how much to regenerate based on the levels of PD factors, direct or indirect connections to growth control mechanisms are to be expected. Indeed, Prod1 was found to interact with a Schwann-cell derived factor of the anterogradient protein family, nAG, which acts as a mitogen for blastema cells[15], providing a link between positional determination and growth regulation.

To date, Prod1 is considered the best example of a molecule encoding positional information[3]. However, important questions remain unanswered. First, it is unclear whether Prod1 reprograms cells to a proximal identity, as it could be expected from a PD determinant, or simply regulates their adhesive properties such that they translocate to proximal locations. Second, axolotl Prod1 lacks a membrane anchor, despite exhibiting the ability to regulate aspects of cell affinity associated to proximalisation[16]. This raises the possibility that, in order to achieve a graded distribution along the PD axis and execute its functions, Prod1 interacts with additional, as yet unidentified components of the proximo-distal memory system.

In view of these considerations, we set out to identify novel components of the PD determination system. This led to the identification of Tig1, a RA responsive gene encoding a cell surface molecule which exhibits graded expression along the PD axis of the mature limb. We show that Tig1 fulfils functional criteria attributed to a proximal determinant, including the ability to reprogramme blastema cells to a proximal identity. Thus, our results implicate Tig1 in the regulation of proximo-distal identity during salamander limb regeneration.

Tig1, a potential cell surface determinant of proximal identity
To identify molecular components of the PD memory system, we performed Principal Component Analysis on a single-cell RNAseq dataset of connective tissue cells during axolotl limb regeneration[17]. This particular tissue type was selected given that it is implicated in the establishment of PD identity during limb regeneration, in contrast to other types such as muscle, which appears to be positionally naïve[18]. Through this approach, we identified a principal component (PC3) that accurately modelled changes along the proximo-distal axis, as exemplified by the clear separation between HoxA13[+] (distal) and Meis2[+] (proximal) cells (Fig.1a). By mining the transcriptome of single cells at the Meis2[+] end of this component, we identified the axolotl homologue of Tig1 (Tazarotene-induced gene 1), also known as Rarres1 (Retinoic Acid Receptor Responder 1), as a highly enriched gene in modelled proximal cells (Fig 1a, 1b). To experimentally test this notion, we performed qRT-PCR gene expression studies in mature and regenerating limbs. As predicted by our *in silico* analysis, Tig1 is expressed in a graded fashion along the PD axis of the mature limb, reaching the highest level in the upper arm and decreasing towards the hand (Fig.1c). Thus, similar to Prod1, Tig1 fulfils a key criterion associated with a molecular determinant of proximal identity, namely graded expression along the mature limb.

Human Tig1 is a RA-responsive putative tumour suppressor[19,20] which exhibits structural resemblance to CD38, a retinoid-responsive surface molecule found in immune cells[21]. Comparative protein sequence analysis (Supplementary Fig. 1) indicates that axolotl Tig1 exhibits substantial sequence similarity (57%) with its human counterpart. Further, the most significant protein features of human Tig1, namely a transmembrane domain, a latexin domain -present in the carboxypeptidase inhibitor Latexin and highly conserved among vertebrates-, and a hyaluronic acid binding motif [20,22], are all present in axolotl Tig1 (Supplementary Fig. 1). These observations raise the possibility that axolotl Tig1 is a RA-responsive gene encoding a cell surface protein, thus fulfilling important criteria[11] for a factor involved in the determination of proximo-distal identity. Indeed, axolotl Tig1 expression is upregulated by RA *in vivo*, in both normal and regenerating limb tissues (Fig. 1e). In agreement, several RA (RXRA; RXRB, RXRG; RARA, RARG) and Meis (Meis1 and Meis2 binding sites) regulatory elements are present in the axolotl Tig1 promoter (Supplementary data 1). Of note, RA-dependent Tig1 induction is observed in connective-tissue derived *Am*AL1 cells, but not in the myogenic line *Nv*A1 (Supplementary Fig. 2). To determine its cellular localisation, axolotl Tig1 was expressed in *Am*AL1 cells as a fusion protein carrying a myc tag in either its C- or N-terminus (Supplementary Fig. 3a), followed by live cell staining with anti-myc antibodies (Fig. 1d). While no myc-derived signal is detected upon expression of N-terminal myc-Tig1, a

distinct cell surface staining is observed following expression of the C-terminal tagged Tig1 (Fig. 1d). The opposite pattern is observed for myc-tagged forms of axolotl Amphiregulin, a known surface protein whose N-terminus faces the extracellular space (Fig. 1d). Similar results were observed in 293T cells (Supplementary Fig. 3b). Together, these observations indicate that axolotl Tig1 localises to the cell surface, with its C-terminus, which comprises the majority of its protein sequence, facing the extracellular space.

While little is known about the biological functions of Tig1 in any context, *in vitro* studies suggest that human Tig1 has a negative impact on cell proliferation[20]. This is particularly interesting in light of the fact that Prod1 has also been shown to impair proliferation[9]. To test whether this is also the case for axolotl Tig1, pN2-Tig1 or empty pN2 vectors were electroporated together with a fluorescent reporter into midbud blastemas, followed by systemic EdU injection and subsequent analysis of EdU incorporation (Supplementary Fig. 4a). As the latexin domain had been proposed to be important for Tig1 function[22], we generated a Tig1 mutant version (Tig1[P155A]) in which the critical cis-proline residue within the latexin domain was mutated to an alanine. Overexpression of either Tig1 or Tig1[P155A] led to significant decreases in EdU incorporation by blastema cells (Supplementary Fig. 4 b, c), suggesting that the function of Tig1 as a negative regulator of cell proliferation is conserved in the axolotl, and that an intact latexin domain is dispensable for it.   Together, the aforementioned observations draw important parallels between Tig1 and Prod1, and thus pinpoint Tig1 as a candidate regulator of proximal identity during regeneration.

Tig1 misexpression disrupts regeneration of distal structures
To gain insights into the functions of Tig1, we first examined its expression throughout limb regeneration. Notably, Tig1 expression is upregulated during the first 7 days post amputation (Fig 1f), coinciding with the establishment of proximal identities[23], followed by a strong downregulation that is maintained until the end of the process (Supplementary Fig. 5), in agreement with previous observations[17]. Next, we sought to disrupt the endogenous expression dynamics by overexpressing Tig1 from the mid-bud stage onwards, when it is normally downregulated. Significantly, increasing the levels of Tig1 via baculovirus(Bv)-mediated transduction, which targets connective tissue cells[24], leads to impaired regeneration of distal elements (Fig. 2a, b). Defects induced by Tig1 overexpression include shortening of digits (2/12), morphological alterations in the lower arm elements radius and ulna, namely pronounced reduction in length (5/12), elongated distal epiphysis (2/12) and uneven thickness of the elements (2/12), as well as malformations of carpals and metacarpals (5/12), while normal limb elements are observed in contralateral control limbs transduced with an mCherry Bv. In particular, there is a significant reduction in limb length (Fig. 2c) in agreement with the observed shortening of the lower arm. This phenotype implicates Tig1 in the establishment of PD patterning of the regenerating limb. Furthermore, it exhibits a striking resemblance to that resulting from overexpressing Prod1[9] or Meis2a[14]. Taken together, these observations strongly suggests that Tig1 may be implicated in the determination of proximal identity.

Tig1 induces proximal displacement of blastema cells
The few molecules implicated in the regulation of proximal identity to date, Prod1 and Meis1/2, are able to elicit the proximal translocation of cells when overexpressed in the early blastema[9,14]. To test whether this is also true for Tig1, we performed focal electroporations on the distal-most compartment of 4-day blastemas. Concurrent with previous reports[9], transfection of a fluorescent reporter results in the labelling of cells that contribute to distal-most elements, digits and carpals, whilst electroporation of Prod1 leads to the displacement of the labelled cells towards proximal regions, occasionally beyond the amputation plane (Fig. 3a, b). Notably, electroporation of either form of Tig1 results in the proximal displacement of

the transfected cells (Fig. 3a,b), with a stronger effect observed for Tig1[P155A], as evaluated by the percentage of animals exhibiting electroporated cells in the proximal segments, including lower and upper arm. These observations show that Tig1 induces proximal displacement of blastema cells that would otherwise contribute to distal domains. In order to track changes in the frequency and distribution of cells along the PD axis, we developed an image analysis algorithm termed Meandros. This software allows the automated detection and analysis of the distribution of cells within the limb based on their fluorescence (Fig. 3c and Supplementary Fig. 6). Analysis of the scaled limb width indicates that this parameter does not change across different experimental conditions (Supplementary Fig. 7). Quantitative insights into the final location of cells along the PD axis indicate a proximal shift in fluorescence distribution after overexpression of Tig1 and Tig1[P155A] when compared with RFP (Fig. 3d). While the majority of the cells electroporated with the control plasmid contribute to the distal hand elements, the overexpression of both wild-type and mutant Tig1 leads to a marked extension in the distribution of cells up to the lower (Tig1) and upper (Tig1 [P155A]) arm levels, reinforcing the notion that Tig1 overexpression induces proximal displacement.

Meandros analysis highlighted differences in PD distribution between Prod1 and Tig1 overexpressing cells (Supplementary Fig 8a). Furthermore, we observed that the combination of Prod1 and Tig1 expression has an additive effect on the extent of proximalisation of Tig1 alone (Supplementary Fig 8b). It remains an open question whether Prod1 and Tig1 act independently of each other or they synergise within the same pathway to promote proximal cell translocation. Expression data (see below) suggest that Tig1 acts upstream of Prod1. However, as dose-dependent effects remain possible, this is a notion that merits additional consideration.

While the displacement assay is indicative of the participation of Tig1 in determining the positional outcome of cells, it does not inform on the cellular processes that could mediate this relocation, such as proliferation, apoptosis, migration and/or adhesion. To address the contribution of cell proliferation to proximal displacement, we compared the effect of overexpressing Prod1, Tig1 and an anti-proliferative factor with no connection to proximalisation, p53[25]. Whereas all three molecules decrease EdU incorporation upon electroporation in blastema cells (Supplementary Fig. 9a), p53 overexpression does not lead to the proximal translocation of blastema cells (Supplementary Fig. 9b-e, Supplementary Fig. S6), demonstrating that the inhibition of cell proliferation *per se* is not sufficient to promote proximal displacement, and indicating that additional processes are at play. Given these observations, and considering that we and others[9,14] have not detected any significant changes in apoptosis in the context of proximal displacement, it seems likely that the preferential distribution of cells towards proximal limb elements is primarily explained by migration and/or active cell sorting arising from differential affinity properties along the PD axis.

Tig1 mediates proximo-distal cell surface interactions
The aforementioned observations led us to functionally test the involvement of Tig1 in mediating proximo-distal differences in cell affinity. To this end, we adapted the blastema engulfment assay[7,11] (Fig. 4a), consisting of juxtaposing blastema mesenchymal tissues derived from similar or different levels along the PD axis and analysing their behaviour *ex vivo* over 7 days. Recapitulating previous observations[7,11], we found that apposing proximal (P) and distal (D) blastemas resulted in engulfment of the distal blastema by its proximal counterpart, while no engulfment took place between blastemas of the same positional value (PP or DD), or between PD blastema pairs treated with anti-Prod1 neutralising antibodies (Fig. 4b, c). To test whether Tig1 mediates such cell affinity differences, we generated custom antibodies against

two non-overlapping regions of axolotl TIG1 (Figure S1, Supplementary data 2 and 3), and addressed the effects of apposing P and D blastemas in their presence. We found that anti-TIG1 neutralising antibodies fully abrogated the engulfment activity, indicating that Tig1 mediates PD differences in cell affinity. This implies that Tig1 underlies cell surface recognition between proximal and distal cells, providing a mechanistic explanation for its role in proximal displacement.

Tig1 reprogrammes cells to a proximal identity

Whether proximal displacement mediated by Tig1 is merely caused by a change in cell affinity or is a consequence of a broader reprogramming of cellular identity towards proximal values is a critical question[9] that, to date, remains unanswered. To address this issue, we investigated the impact of overexpressing Tig1 in cells which had already been specified to a distal identity. To exclusively label distal cells, we used a C-Ti$^{t/+}$(*HoxA13:HoxA13-T2a-mCherry*)^{ET} reporter line, *HoxA13:mCherry hereafter,* created by CRISPR/Cas9 technology[26]. In these axolotls, mCherry is fused to the C-terminus of the HOXA13, separated by a T2a self-cleaving peptide, and expressed from the endogenous Hoxa13 genomic locus. Validation studies show that mCherry is exclusively detectable in the distal-most part of the limb bud/ blastema during development and regeneration (Supplementary Fig. 10), labelling the field that gives rise to the hand segment in accordance with the model of progressive specification[23,27], thus offering the means to genetically identify blastema cells that have been specified to a distal fate based on their fluorescence. Upon appearance of the cohort of mCherry$^+$ distal cells in the regenerating limb, we electroporated Tig1 or Tig1^{P155A}T2A-GFP constructs, FACS sorted mCherry$^+$/GFP$^+$ (distal cells overexpressing Tig1) and mCherry$^+$/GFP$^-$ (distal control cells) cells at 3 or 8 days post electroporation, and performed bulk RNA sequencing analysis (Fig. 5a). PCA analysis indicates a clear segregation between control versus Tig1/Tig1^{P155A} overexpressing cells, and between timepoints (Fig 5b). Overexpression of Tig1 versus Tig1^{P155A} has limited impact on clustering, suggestive of phenotypic similarities and consistent with our experimental data. This is also reflected in the differential expression analysis (Fig. 5c). Notably, we found a substantial number of genes (c. 5000) whose expression significantly changed upon Tig1 overexpression (P<0.001), suggesting an extensive transcriptome remodelling (Fig. 5c, Supplementary Fig. 11). Using K-means hierarchical clustering we grouped the top 2000 differentially expressed genes into three main gene clusters. Gene Ontology enrichment analysis of the three groups (Fig. 5c), revealed main GO term categories for each cluster. Cluster A is enriched in GO terms related to developmental and tissue morphogenesis processes (Fig. 5d), whereas clusters B (genes activated by Tig1 at all timepoints) and C (genes activated by Tig1 but primarily at 8dpe) are enriched in immune system and actin cytoskeleton remodelling processes respectively (Supplementary Fig. 12, Supplementary data 4).

Critically, in depth analysis of gene expression changes caused by Tig1 overexpression revealed effects in genes related to the establishment of the proximo-distal axis. Tig1 overexpression results in a highly significant upregulation of the proximalisation factor Prod1, a moderate upregulation of the proximal regulator Meis1, and the downregulation of the distal marker HoxA13 compared to control cells (Fig. 5e). Consistent with the latter, a network of 11 genes known to be either co-regulated or direct targets of HoxA13 is also downregulated in response to either of the Tig1 forms (Fig. 5f). Among these are bone morphogenetic proteins (BMP) Bmp2 and Bmp7, important for sustaining progressive regeneration and formation of distal elements[27]. In addition, elevated Tig1 levels promote the downregulation of morphogenetic factors associated with blastema growth and patterning, including Lhx9, Lmo1, Spry1 and Lhx2 (Fig. 5g), which are known to be expressed in the distal portion of the developing and regenerating vertebrate limb and are downregulated in the axolotl limb upon proximalising doses of RA[28]. Further, we observe downregulation of Fgf8, a negative regulator

of Meis[29] associated with the specification of the distal limb domain[30]. This inhibitory effect on distal growth provides a molecular explanation for the defects caused by Tig1 overexpression in midbud blastemas (Fig. 2). Besides its effect on growth factors, Tig1 overexpression also promotes the downregulation of genes encoding factors that promote cell cycle progression, and the upregulation of genes encoding the tumour suppressors Cdkn1B and Retinoblastoma (Supplementary Fig. 13), consistent with the anti-proliferative effect of Tig1 (Supplementary Fig. 4). Also in agreement with the predicted functions of Tig1, a number of cell adhesion and ECM remodelling factors were found among the differentially expressed genes (Fig. 5h), suggestive of changes in cell affinity and migratory capabilities. These include the downregulation of Tnc, Fn1, Col3A1, Lsu, Col8A1, MMP18 and N-cadherin, and the upregulation of Hzal3, Sdc3, Cdh1 and MMP13. Of note, MMP13 upregulation has been shown to occur upon proximalising RA doses[28]. Further, Cdh1 and Sdc3 were found to be upregulated in proximal connective tissue cells during limb regeneration according to our PCA analysis (Fig. 1b). These observations are consistent with the acquisition of proximal ECM and cell affinity expression profiles upon Tig1 overexpression, and provide a mechanistic explanation for the effect of Tig1 in the engulfment and displacement assays. Interestingly, whereas our global analysis reveals dynamic changes in gene expression between 3 and 8dpe (Fig 5b, c), no significant changes are observed for the aforementioned PD-related genes (Fig 5e-h). This suggest that Tig1 elicits a rapid and stable transcriptome re-shaping towards proximal identity in originally distal cells, already manifest at 3dpe. Altogether, this dataset demonstrates that Tig1 confers blastema cells proximal values and hinders distalisation.

Tig1 controls the expression of key proximo-distal regulators
In order to validate the RNAseq and obtain further insights into the impact of Tig1 on the expression of PD-related factors, we performed qRT-PCR analysis on *HoxA13:mCherry* sorted cells after Tig1 electroporation, according to the conditions used for the RNAseq (Fig. 6a). We observed that overexpression of either Tig1 or Tig1[P155A] lead to the upregulation of the proximal factor Prod1, as well as the downregulation of HoxA11, HoxA13, the distal growth factor Lxh2, and two selected HoxA13-regulated genes, Enpp2 and Epha7 (Fig 6a), consistent with the results of our RNAseq analysis. Interestingly, we note that Tig1 overexpression not only leads to an increase in the relative levels of Tig1 mRNA, but also to an increase in Tig1 5'UTR (not present in the pN2-Tig1 vector), indicating that increased Tig1 levels result in the upregulation of Tig1 transcription, constituting a positive feedback loop (Fig 6a). It is noteworthy that Meis1 expression shows an increase upon Tig1 overexpression even though, due to the high variability between the replicates, it is not significant. A correlation between Meis and Tig1 gene expression is also indicated by our PCA analysis of connective tissue cells at 11dpa (Fig1b). Therefore it is conceivable that Tig1 activation is achieved through the upregulation of Meis factors, given that the Tig1 promoter contains Meis binding sites (Supplementary data 1). This is a notion that merits further exploration. Together, these results suggest that Tig1 is associated with and actively regulates the network of factors that modulate proximo-distal identity in the salamander limb.

Concluding remarks
The molecular and cellular basis of positional identity remains poorly understood[1,31]. Mechanistic knowledge is currently lacking, from the molecular factors that determine the PD nature of a cell, to their coordination and their re-specification in a regenerative context. Our study offers significant insights into this problem by identifying Tig1 as a new determinant of PD identity.
Whilst a system for the specification of PD identity has been proposed to operate at the cell surface level, Prod1 remained its only presumed component. Like Prod1[11], Tig1 is a RA-

responsive factor localised to the cell surface and exhibiting graded PD expression in the mature limb. Further, Prod1 expression is regulated by Meis[12] and, as our Tig1 promoter analysis suggests, Tig1 is also a putative Meis target. Yet unlike Prod1, which is salamander-specific[32] and lacks a membrane anchor in the axolotl[16], Tig1 is highly conserved through evolution and membrane bound, raising the possibility that it forms part of a conserved PD determination mechanism. Phenotypically, Tig1 resembles Prod1 in that it mediates cell behaviours associated with proximal identity, including the engulfment activity of apposed proximal and distal blastemas, and the displacement of blastema cells to proximal locations. Further, like Prod1[9] and Meis1[14], Tig1 overexpression leads to patterning defects during limb regeneration, primarily affecting distal elements. Altogether, this strongly suggest that Tig1 is involved in the determination of proximal identity. Yet, our study provides additional evidence for this notion, namely the demonstration that Tig1 is able to elicit the reprogramming of distal cells to a proximal identity.

To our knowledge, this is the first example of a cell surface molecule able to induce a global reprogramming of proximo-distal identity. Our observations suggest that the amount of Tig1 molecules at the cell surface can determine the expression levels of proximal factors such as Meis1/2 and Prod1 in order to promote proximalisation, supporting the notion that Tig1 is a critical component of a positional identity system encoded at the cell surface level. This, together with its graded expression along the PD axis of the mature limb, suggests that Tig1 confers spatial coordinates to adult tissues and contributes to the sensing of positional disparities upon amputation, an essential component of the regenerative response[2,33], as previously proposed for Prod1[11]. It would be of interest to address whether Prod1 and Tig1 act together in order to mediate PD disparity recognition. The existence of cell-autonomously encoded spatial information in adult cells has been well-documented in human fibroblasts, where it is epigenetically maintained[34,35]. It would thus be informative to address if Tig1 is also part of the mammalian code, and whether epigenetic mechanisms are responsible for the regulation of its expression as well as that of other PD factors.

Beside the regulation of proximal factors, Tig1-induced reprogramming involves a number of additional, noteworthy processes. First, it entails a marked downregulation of distal factors. Whether the prevention of distalisation is sufficient to proximalise cell identity is an important outstanding question. Second, it is associated with a marked anti-proliferative molecular signature, consistent with the fact that Tig1 counteracts proliferation. It is thus possible that a stable proximal identity is associated to lack of cell proliferation. This is in agreement with the decrease in proliferation observed in the proximal-most regions of the late blastema[36]. Importantly however, suppressing cell cycle progression is not sufficient to promote proximal translocation. Lastly, the mechanisms by which the levels of this cell surface factor determine broad transcriptional changes are as yet unknown. It is conceivable that cytoskeletal arrangements are involved, as Tig1 has been proposed as a modulator of microtubule dynamics[37-39], a notion that should be addressed by further research.

Given the importance of Tig1 for the specification of proximal identity, and taking into account its plastic responsiveness to RA treatment in both blastema and mature limb tissues, we propose that Tig1 is a likely mediator of the RA-dependent proximalisation effect[10,40] responsible for limb segment duplications. Meis1 has been proven necessary for the proximalising activity of RA in this context[14]. In light of our observations, in particular the upregulation of Tig1 upon RA treatment and the upregulation of Meis1 following an increase in Tig1 levels, forthcoming efforts should test whether Tig1 is part of the RA-responsive network regulating proximalisation of regenerating structures.

Together, the body of evidence hereby presented indicates that Tig1 is a central regulator of proximo-distal identity. While this clearly constitutes an important step towards understanding

the molecular network underlying PD determination, critical questions remain. These include the nature of the connection between Tig1 and other key factors, such as Prod1 and Meis, in the determination of PD identity, the functional relevance of Tig1 for the establishment of PD memory in the normal and regenerating limb, and its mechanism of action at the cellular and molecular levels, particularly whether and how it may act as a 'sensor' of positional disparity between cells. Future studies will provide answers to these questions, delivering fundamental insights into the mechanism underlying positional identity in a vertebrate context.

References

Flowers, G. P. & Crews, C. M. Remembering where we are: Positional information in salamander limb regeneration. *Dev Dyn* **249**, 465-482, doi:10.1002/dvdy.167 (2020).
Vieira, W. A. & McCusker, C. D. Hierarchical pattern formation during amphibian limb regeneration. *Biosystems* **183**, 103989, doi:10.1016/j.biosystems.2019.103989 (2019).
Wolpert, L. & Vicente, C. An interview with Lewis Wolpert. *Development* **142**, 2547-2548, doi:10.1242/dev.127373 (2015).
Brockes, J. P. & Kumar, A. Comparative aspects of animal regeneration. *Annu Rev Cell Dev Biol* **24**, 525-549, doi:10.1146/annurev.cellbio.24.110707.175336 (2008).
Stocum, D. L. & Thoms, S. D. Retinoic-acid-induced pattern completion in regenerating double anterior limbs of urodeles. *J Exp Zool* **232**, 207-215, doi:10.1002/jez.1402320208 (1984).
Crawford, K. & Stocum, D. L. Retinoic acid proximalizes level-specific properties responsible for intercalary regeneration in axolotl limbs. *Development* **104**, 703-712 (1988).
Nardi, J. B. S., D. Surface properties of regenerating limb cells: Evidence for gradation along the proximodistal axis. *Differentiation* **25**, 27-31, doi:https://doi.org/10.1111/j.1432-0436.1984.tb01334.x (1983).
Pescitelli, M. J., Jr. & Stocum, D. L. The origin of skeletal structures during intercalary regeneration of larval Ambystoma limbs. *Dev Biol* **79**, 255-275, doi:10.1016/0012-1606(80)90115-3 (1980).
Echeverri, K. & Tanaka, E. M. Proximodistal patterning during limb regeneration. *Dev Biol* **279**, 391-401, doi:10.1016/j.ydbio.2004.12.029 (2005).
Maden, M. Vitamin A and pattern formation in the regenerating limb. *Nature* **295**, 672-675, doi:10.1038/295672a0 (1982).
da Silva, S. M., Gates, P. B. & Brockes, J. P. The newt ortholog of CD59 is implicated in proximodistal identity during amphibian limb regeneration. *Dev Cell* **3**, 547-555, doi:10.1016/s1534-5807(02)00288-5 (2002).
Shaikh, N., Gates, P. B. & Brockes, J. P. The Meis homeoprotein regulates the axolotl Prod 1 promoter during limb regeneration. *Gene* **484**, 69-74, doi:10.1016/j.gene.2011.06.003 (2011).
Mercader, N. *et al.* Conserved regulation of proximodistal limb axis development by Meis1/Hth. *Nature* **402**, 425-429, doi:10.1038/46580 (1999).
Mercader, N., Tanaka, E. M. & Torres, M. Proximodistal identity during vertebrate limb regeneration is regulated by Meis homeodomain proteins. *Development* **132**, 4131-4142, doi:10.1242/dev.01976 (2005).

Kumar, A., Godwin, J. W., Gates, P. B., Garza-Garcia, A. A. & Brockes, J. P. Molecular basis for the nerve dependence of limb regeneration in an adult vertebrate. *Science* **318**, 772-777, doi:10.1126/science.1147710 (2007).

Blassberg, R. A., Garza-Garcia, A., Janmohamed, A., Gates, P. B. & Brockes, J. P. Functional convergence of signalling by GPI-anchored and anchorless forms of a salamander protein implicated in limb regeneration. *J Cell Sci* **124**, 47-56, doi:10.1242/jcs.076331 (2011).

Gerber, T. *et al.* Single-cell analysis uncovers convergence of cell identities during axolotl limb regeneration. *Science* **362**, doi:10.1126/science.aaq0681 (2018).

Nacu, E. *et al.* Connective tissue cells, but not muscle cells, are involved in establishing the proximo-distal outcome of limb regeneration in the axolotl. *Development* **140**, 513-518, doi:10.1242/dev.081752 (2013).

Jing, C. *et al.* Tazarotene-induced gene 1 (TIG1) expression in prostate carcinomas and its relationship to tumorigenicity. *J Natl Cancer Inst* **94**, 482-490, doi:10.1093/jnci/94.7.482 (2002).

Nagpal, S. *et al.* Tazarotene-induced gene 1 (TIG1), a novel retinoic acid receptor-responsive gene in skin. *J Invest Dermatol* **106**, 269-274, doi:10.1111/1523-1747.ep12340668 (1996).

Wang, X. *et al.* TIG1 promotes the development and progression of inflammatory breast cancer through activation of Axl kinase. *Cancer Res* **73**, 6516-6525, doi:10.1158/0008-5472.CAN-13-0967 (2013).

Aagaard, A. *et al.* An inflammatory role for the mammalian carboxypeptidase inhibitor latexin: relationship to cystatins and the tumor suppressor TIG1. *Structure* **13**, 309-317, doi:10.1016/j.str.2004.12.013 (2005).

Roensch, K., Tazaki, A., Chara, O. & Tanaka, E. M. Progressive specification rather than intercalation of segments during limb regeneration. *Science* **342**, 1375-1379, doi:10.1126/science.1241796 (2013).

Oliveira, C. R. *et al.* Pseudotyped baculovirus is an effective gene expression tool for studying molecular function during axolotl limb regeneration. *Dev Biol* **433**, 262-275, doi:10.1016/j.ydbio.2017.10.008 (2018).

Yun, M. H., Gates, P. B. & Brockes, J. P. Regulation of p53 is critical for vertebrate limb regeneration. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 17392-17397, doi:10.1073/pnas.1310519110 (2013).

Fei, J. F. *et al.* Application and optimization of CRISPR-Cas9-mediated genome engineering in axolotl (Ambystoma mexicanum). *Nat Protoc* **13**, 2908-2943, doi:10.1038/s41596-018-0071-0 (2018).

Vincent, E. *et al.* BMP signaling is essential for sustaining proximo-distal progression in regenerating axolotl limbs. *Development* **147**, doi:10.1242/dev.170829 (2020).

Nguyen, M. *et al.* Retinoic acid receptor regulation of epimorphic and homeostatic regeneration in the axolotl. *Development* **144**, 601-611, doi:10.1242/dev.139873 (2017).

Mercader, N. *et al.* Opposing RA and FGF signals control proximodistal vertebrate limb development through regulation of Meis genes. *Development* **127**, 3961-3970 (2000).

Mariani, F. V., Ahn, C. P. & Martin, G. R. Genetic evidence that FGFs have an instructive role in limb proximal-distal patterning. *Nature* **453**, 401-405, doi:10.1038/nature06876 (2008).

Stocum, D. L. & Cameron, J. A. Looking proximally and distally: 100 years of limb regeneration and beyond. *Dev Dyn* **240**, 943-968, doi:10.1002/dvdy.22553 (2011).

Garza-Garcia, A. A., Driscoll, P. C. & Brockes, J. P. Evidence for the local evolution of mechanisms underlying limb regeneration in salamanders. *Integr Comp Biol* **50**, 528-535, doi:10.1093/icb/icq022 (2010).

Endo, T., Bryant, S. V. & Gardiner, D. M. A stepwise model system for limb regeneration. *Dev Biol* **270**, 135-145, doi:10.1016/j.ydbio.2004.02.016 (2004).

Rinn, J. L., Bondre, C., Gladstone, H. B., Brown, P. O. & Chang, H. Y. Anatomic demarcation by positional variation in fibroblast gene expression programs. *PLoS Genet* **2**, e119, doi:10.1371/journal.pgen.0020119 (2006).

Rinn, J. L. *et al.* Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell* **129**, 1311-1323, doi:10.1016/j.cell.2007.05.022 (2007).

McCusker, C. D. *et al.* Positional plasticity in regenerating Amybstoma mexicanum limbs is associated with cell proliferation and pathways of cellular differentiation. *BMC Dev Biol* **15**, 45, doi:10.1186/s12861-015-0095-4 (2015).

Maimouni, S. *et al.* Tumor suppressor RARRES1 links tubulin deglutamylation to mitochondrial metabolism and cell survival. *Oncotarget* **10**, 1606-1624, doi:10.18632/oncotarget.26600 (2019).

Sahab, Z. J. *et al.* Tumor suppressor RARRES1 interacts with cytoplasmic carboxypeptidase AGBL2 to regulate the alpha-tubulin tyrosination cycle. *Cancer Res* **71**, 1219-1228, doi:10.1158/0008-5472.CAN-10-2294 (2011).

Sahab, Z. J., Hall, M. D., Zhang, L., Cheema, A. K. & Byers, S. W. Tumor Suppressor RARRES1 Regulates DLG2, PP2A, VCP, EB1, and Ankrd26. *J Cancer* **1**, 14-22, doi:10.7150/jca.1.14 (2010).

Stocum, D. L. & Crawford, K. Use of retinoids to analyze the cellular basis of positional memory in regenerating amphibian limbs. *Biochem Cell Biol* **65**, 750-761, doi:10.1139/o87-098 (1987).

Knapp, D. *et al.* Comparative transcriptional profiling of the axolotl limb identifies a tripartite regeneration-specific gene program. *PLoS One* **8**, e61352, doi:10.1371/journal.pone.0061352 (2013).

Tank, P. W., Carlson, B. M. & Connelly, T. G. A staging system for forelimb regeneration in the axolotl, Ambystoma mexicanum. *J Morphol* **150**, 117-128, doi:10.1002/jmor.1051500106 (1976).

LeCun, Y., Bengio, Y. & Hinton, G. Deep learning. *Nature* **521**, 436-444, doi:10.1038/nature14539 (2015).

K. He, G. G., P. Dollár and R. Girshick. Mask R-CNN. *2017 IEEE International Conference on Computer Vision (ICCV), Venice*, 2980-2988, doi:10.1109/ICCV.2017.322 (2017).

K. He, X. Z., S. Ren and J. Sun. in *2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR), Las Vegas, NV*   770-778 (2016).

Sobkow, L., Epperlein, H. H., Herklotz, S., Straube, W. L. & Tanaka, E. M. A germline GFP transgenic axolotl and its use to track cell fate: dual origin of the fin mesenchyme during development and the fate of blood cells during regeneration. *Dev Biol* **290**, 386-397, doi:10.1016/j.ydbio.2005.11.037 (2006).

Yun, M. H., Gates, P. B. & Brockes, J. P. Sustained ERK activation underlies reprogramming in regeneration-competent salamander cells and distinguishes them from their mammalian counterparts. *Stem Cell Reports* **3**, 15-23, doi:10.1016/j.stemcr.2014.05.009 (2014).

Corcoran, J. P. & Ferretti, P. RA regulation of keratin expression and myogenesis suggests different ways of regenerating muscle in adult amphibian limbs. *J Cell Sci* **112 (Pt 9)**, 1385-1394 (1999).

Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8 (2014).

Ge, S. X., Son, E. W. & Yao, R. iDEP: an integrated web application for differential expression and pathway analysis of RNA-Seq data. *BMC Bioinformatics* **19**, 534, doi:10.1186/s12859-018-2486-6 (2018).

Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* **25**, 25-29, doi:10.1038/75556 (2000).

Acknowledgments: We thank all members of the Yun Lab for critical comments, Jifeng Fei for advice on knock-in transgenesis, Jeremy Brockes for insightful discussions, Gabriel Waksman for institutional support, MPI-CBG Protein Expression Facility for protein/viral expression support, TUD-CMCB Light Microscopy Facility and TUD-CMCB Histology Facility for tissue processing and imaging support, and Beate Gruhl, Anja Wagner and Dominick Kruger for animal care.

Funding: A.E. was supported by a NIH Ruth Kirschstein postdoctoral fellowship (F32GM117806), H.A. by a CONICET-UADE scholarship, O.C. by CONICET (Argentina) and ANPCyT grants (PICT-2014-3469 & PICT-2017-2307), E.M.T by ERC Advanced grant (294324) and DFG grant (TA274/3-3); A.S. by Cancerfonden, Swedish Research Council, Knut and Alice Wallenberg Foundation, Stiftelsen Olle Engkvist Bryggmästare, and M.H.Y by Deutsche Forschungsgemeinschaft grants (DFG 22137416 & 450807335) and TUD-CRTD core funds.

Author contributions: C.O., D.K., S.G.M. and P.B.G. performed experiments; C.O., S.G.M., A.E., D.K., P.B.G., A.P, H.A., R. C., O.C., A.S. and M.H.Y. analysed and discussed data; C.O., D.K., S.G.M., P.B.G. and M.H.Y. designed experiments; E.M.T. contributed reagents; M.H.Y. supervised the project and wrote the manuscript with contributions from all authors.

Competing interests: Authors declare no competing interests.

Material & Correspondence: Correspondence and request for materials should be addressed to M.H.Y.

Figure Legends:

Fig. 1. The retinoic acid-responsive gene Tig1 encodes a cell surface protein and is differentially expressed along the proximo-distal axis of the axolotl limb. (a,b) Tig1 expression is segregated along the modelled proximodistal axis of the regenerating limb. (a) Principal component analysis (PCA) (PC3 vs PC4) of individual connective tissue cells at 11 days post amputation (dpa). PC3 models the proximodistal axis in the regenerating blastema, as exemplified by the expression of Meis2 (proximal) and HoxA13 (distal). (b) Contributions of axolotl genes to PC3. Genes associated with the proximal (red) and distal (blue) axes, or with cell adhesion (yellow) are indicated. The scRNAseq dataset does not contain expression data for Prod1. Note that Tig1 is found at the extreme Meis2$^+$ end of PC3. (c) Tig1 expression

along the proximo-distal axis of the mature axolotl limb, relative to Ef1-α (UA: upper arm; LA: lower arm; H: hand; qRT-PCR, n=4). Error bars represent standard deviation of the mean. (d) Representative images of AL1 cells after electroporation with the indicated myc-tagged constructs and live cell staining using anti-myc antibodies. Scale bar: 20μm. (e) Expression of axolotl Tig1 in mature or regenerating (midbud stage) limbs 48h after retinoic acid (RA) treatment (100 μg/g body weight), relative to DMSO treated samples, normalised against Ef1-α expression (qRT-PCR, n=4). (f) Microarray[41] analysis of Tig1 gene expression for the indicated bulk tissues and time points post-amputation and wound healing (lateral wound). Note the upregulation of Tig1 during the first 7 days after amputation. Shading represents standard deviation of the mean. ** p<0.001 ; * p<0.05 (paired-matched t-test, c and e).

Fig. 2. Tig1 overexpression disrupts regeneration of distal elements. (a) Schematic of the experimental approach. (b) Tig1 Bv transduction from 15dpa onwards leads to shortening of the regenerated limb through reduction of the lower arm (LA) and digit malformations, as analysed through Alcian blue/Alizarin red staining (n=12). (c) Quantification of limb length. Error bars represent standard deviation of the mean. * p<0.05 (paired-matched t-test).

Fig. 3. Tig1 induces distal-to-proximal displacement of blastema cells. (a) Representative images of the displacement assay. Cells in the distal third of a 4-day blastema give rise to distal elements[9]. Upon electroporation of a control vector (n=7) to enable tracing during regeneration, they contribute primarily to digits (first column). Overexpression of Prod1 (n=8), Tig1 or Tig1[P155A] (n=13) coupled to fluorescent proteins promotes autologous displacement of cells from distal to proximal positions during limb regeneration. dpe: days post electroporation. Scale bar: 1 mm. (b) Quantification of the percentage of animals with electroporated cells found in the indicated limb segments. (c) An image analysis algorithm for the automatised quantification of cells (by fluorescence intensity) along the PD axis. f indicates the number of pixels whose intensity is higher than the threshold; f/fmax indicates the ratio between each f and its maximal value along the anatomical axis. (d) Distribution of fluorescence intensity along the PD axis for the indicated conditions. Coloured shading indicates standard deviation of the *f/fmax*. Grey shading indicates standard deviation of the PD position corresponding to elbow (E) or wrist (W). A: amputation plane.

Fig. 4. Tig1 mediates proximo-distal cell-surface interactions. (a) Schematic of the engulfment assay. Juxtaposition of proximal (P) and distal (D) blastemas results in the engulfment of the distal one by its proximal counterpart. (b) Representative images of juxtaposed proximal (red) and distal (white) blastemas at the indicated times post juxtaposition, and after the indicated treatments. Note lack of engulfment following treatment with anti-TIG1 (α-TIG1) antibodies raised against two extracellular epitopes. [Antibody]=20μg/ml. (c) Quantification of engulfment for the indicated pairs (pooled data from 3 independent experiments).

Fig. 5. Tig1 reprograms distal cells to proximal identities. (a) Experimental design. Tig1-T2A-GFP or Tig1[P155A]-T2A-GFP were electroporated in the distal, mCherry[+] domain of late bud blastemas of *HoxA13:mCherry* axolotls; FACS isolation followed by bulk RNAseq analysis was carried out for the indicated cells at 3 or 8dpe. (b) tSNE plot using the first and second dimension shows consistent behaviour between replicates and a separation between treatments. (c) Heatmap of the 2000 most differentially expressed genes between conditions (d) GO enrichment analysis of Biological Process for three gene groups determined by K-means hierarchical clustering, displaying the top ten enriched terms for cluster A. The enriched pathways are listed with their corresponding adjusted p-values and number of genes. (e-h) Tig1

overexpression in distal cells promotes a transcriptome shift toward proximal identity. Bars represent ratios of gene expression in Tig1 overexpressing cells versus control cells. (e) Relative gene expression changes in PD markers. Note that Tig1 overexpression leads to the upregulation of proximal factors Meis1a and Prod1 and the downregulation of distal genes including the master regulator HoxA13 (e), and its downstream network (f). mCherry protein persists in distal cells even if Hoxa13 is downregulated. (g) Relative gene expression changes in genes associated with blastema growth. (h) Relative gene expression changes in cell adhesion/matrix remodelling factors. Adj. p-values: * $p<0.05$; ** $p<0.001$.

Fig. 6. Tig1 controls the expression of proximo-distal factors Meis1, Prod1 and HoxA13. (a) qRT-PCR quantification of the expression of the indicated factors in FACS-sorted blastema cells from *HoxA13:mCherry* axolotls at 3 or 8 days post-electroporation with the indicated constructs, relative to the control sample at 3dpe. Rpl4 was used as a normaliser. * $p<0.05$; ** <0.001 (unpaired T-test).

Additional information:

Supplementary information is available for this manuscript.

Supplementary information:

Supplementary Figures 1- 13

Supplementary Table 1

Supplementary data 1-4

Supplementary figure/table legends:

Supplementary Fig. 1. Axolotl TIG1 protein sequence. Sequence alignment between axolotl TIG1 and its human counterpart. Axolotl TIG1 exhibits 57% similarity at the aminoacid level, an intact latexin domain (IPDXXG,red square) and a hyaluronic acid binding motif (green square). The peptide sequences selected for subsequent antibody production are underscored in yellow. The hydrophobic tract corresponding to the transmembrane domain is underscored in blue.

Supplementary Fig. 2. Tig1 is upregulated by RA in cells of connective tissue origin. (a,b) qRT-PCR analysis of Tig1 expression in connective tissue- derived axolotl AL1 cells (a) or muscle-derived newt A1 cells (b). Data represents Tig1 gene expression at the indicated times following treatment with 1×10^{-7} M RA, relative to treatment with vehicle (DMSO). Results are normalised against Ef1-α (n=3). *$p<0.05$, **$p<0.001$.

Supplementary Fig. 3. Axolotl TIG1 localises to the cell surface exposing its C-terminus to the extracellular environment. (a) Western blot analysis of myc-tagged protein in *Am*AL1 extracts 48hs after nucleofection with the indicated vectors. Asterisk indicates the band corresponding to the TIG1-myc fusion (31kDa). (b) Representative images of 293T cells 24 hours after transfection with myc-tagged versions of axolotl amphiregulin or TIG1 followed by live cell staining using anti-myc antibodies. Scale bar: 20μm.

Supplementary Fig. 4. Tig1 overexpression impairs blastema cell proliferation *in vivo*. (a) Schematic of experimental strategy (b) Representative images of axolotl late-bud blastemas following anti-RFP immunohistochemistry and azide-mediated EdU detection, 72h after electroporation with the indicated constructs and after two EdU pulses at 24h and 48h post electroporation (n = 5). Insert shows a magnification of the blastema area. Arrowheads indicate EdU$^+$/RFP$^+$ cells. Scale bar: 100μm. (c) Percentage of EdU$^+$ cells among the RFP$^+$ cell population. **p<0.001.

Supplementary Fig. 5. Analysis of Tig1 gene expression during axolotl limb regeneration. Expression of Tig1 during axolotl limb regeneration following amputation at the upper arm level, relative to mature limb and normalised against Ef1-α (qRT-PCR, n=6). Similar results were obtained normalising against L-27.

Supplementary Fig. 6. Meandros *f/fmax* charts per experimental condition. Plots of fluorescence distribution along the PD axis as obtained by Meandros image analysis algorithm are given for each individual condition. Dashed line at PD position=0 indicates the location of the amputation plane (A). Coloured shading indicates standard deviation of *f/fmax*. Grey shading indicates standard deviation of the PD position corresponding to elbow (E) or wrist (W). Control (n=3), Tig1^{wt} (n=4), Tig1^{P155A} (n=4), Prod1 (n=5), Prod1-Tig1^{wt} (n=3), p53 (n=3).

Supplementary Fig. 7. Meandros scaled width analysis along the PD axis. Plots representing the variation of limb width (μm) along the PD axis as obtained by Meandros image analysis algorithm are given for each individual condition. Dashed line at PD position=0 indicates the location of the amputation plane (A). Coloured shading indicates standard deviation of scaled width. Grey shading indicates standard deviation of the PD position corresponding to elbow (E) or wrist (W).

Supplementary Fig. 8. Tig1 and Prod1 have an additive effect on distal-to-proximal blastema cell displacement. (a) Distribution of fluorescence intensity along the PD axis for the indicated conditions. Coloured shading indicates standard deviation. Grey shading indicates standard deviation of the PD position corresponding to elbow (E) or wrist (W). A: amputation plane. (b) Quantification of the number of limbs with electroporated cells in the indicated segments following a displacement assay. Control (n=7), Tig1^{wt} (n=7), Prod1 (n=8) and Prod1-Tig1^{wt} (n=12).

Supplementary Fig. 9. Impairment in cell cycle progression is not sufficient to promote distal-to-proximal blastema cell displacement. (a) Quantification of EdU$^+$/RFP$^+$ blastema cells following anti-RFP immunohistochemistry and azide-mediated EdU detection, 72 hours post electroporation with the indicated pN2 vectors and after two EdU pulses at 24h and 48h post electroporation (n = 5). (b) Quantification of the number of animals with electroporated cells in the indicated limb segments following a displacement assay using a control vector (n=7), p53 (n=7), Tig1^{wt} (n=7), and Prod1 (n=8). (c) Distribution of fluorescence intensity along the PD axis for the indicated conditions. Coloured shading indicates standard deviation. Note that p53 overexpression does not lead to distal-to-proximal displacement. Grey shading indicates standard deviation of the PD position corresponding to elbow (E) or wrist (W). A: amputation plane.

Supplementary Fig. 10. A HoxA13 knock-in reporter line allows distal cell labelling during salamander limb regeneration. (a) Representative images of limb bud development in the *HoxA13:HoxA13-T2A-mCherry* axolotl line. Scale bar: 100μm. (b) Representative

images of limb regeneration in *HoxA13:HoxA13-T2A-mCherry* axolotls. mCherry fluorescence appears in the distal domain of midbud blastema, and intensifies through the course of regeneration until digit stage. Scale bar: 500µm.

Supplementary Fig. 11. Tig1 overexpression promotes broad transcriptomic changes. (a-d) MA plot representation of differential expression analysis of genes expressed in control distal cells (mCherry$^+$) versus Tig1 (a, b) or Tig1^{P155A} (c, d) overexpressing distal cells (mCherry$^+$/GFP$^+$), at 3 (a,c) or 8 (b, d) days post electroporation. The number of differentially expressed genes (DGE) for each comparison is annotated in the top right corner. Red points mark the genes with significantly increased or decreased expression (FDR<0.05). The number of differentially expressed genes is indicated in the top left corner of each graph.

Supplementary Fig. 12. Gene ontology analysis of differentially expressed genes between control, Tig1 and Tig1^{P155A} overexpressing cells. GO enrichment analysis of the Biological process terms in the 3 groups determined by K-means hierarchical clustering (number of groups was chosen based on tSNE and Elbow method), displaying the top ten enriched terms for cluster B (a) and C (b). The enriched pathways are listed with their corresponding adjusted p-value and number of genes.

Supplementary Fig. 13. Tig1 overexpression promotes the downregulation of cell cycle progression genes and upregulation of tumour suppressors. Relative gene expression changes in genes associated with cell cycle progression among differentially expressed genes between control, Tig1 or Tig1^{P155A} overexpressing distal cells at 3 or 8 dpe. * p<0.05; ** p<0.001.

Supplementary Table 1. qRT-PCR primer sequences.

Supplementary data description:

Supplementary data 1. Predicted regulatory elements (RA-related, Hox and Meis1/2 binding sites) within 40kb upstream of the start site of Tig1 coding sequence.

Supplementary data 2. Elisa and SDS-PAGE analysis of purified antibodies against Tig1 peptide 154

Supplementary data 3. Elisa and SDS-PAGE analysis of purified antibodies against Tig1 peptide 154

Supplementary data 4. List of genes corresponding to the GO pathway enrichment analysis and corresponding adjusted p-values.

Methods:

Animal husbandry
Procedures for care and manipulation of all animals used in this study were performed in compliance with the Animals -Scientific Procedures- Act 1986 (United Kingdom Home Office), and the laws and regulations of the State of Saxony, Germany. Axolotls (*A.*

mexicanum) were obtained from Neil Hardy Aquatica (Croydon, UK) and from the axolotl Facility at TUD-CRTD Center for Regenerative Therapies Dresden (Germany). Animals were maintained in individual aquaria at approximately 18-20 °C, as previously described[25]. Axolotls of the leucistic (d/d) strain were used in all experiments.

Animal Procedures and tissue collection

Animals of length between 2.5 and 4 cm from snout to cloaca (that is, between 4 and 8 cm from snout to tail tip) were used in all experiments, except for the engulfment assay, for which animals 7-10 cm from snout to cloaca were utilised.

Axolotls were anesthetised in 0.03 % benzocaine (Sigma) prior to limb amputation at the mid-humerus or mid-radius/ulna level as indicated. Animals were allowed to regenerate at 20 °C and limbs or blastemas collected and processed for various assays as specified. Axolotl regeneration stages were defined as previously described[42]. All procedures were performed using Olympus SZX10 microscopes. Injections were carried out with a Narishige micromanipulator connected to a PV820 pressure injector (WPI).

***In silico* modelling of the proximodistal axis from scRNAseq data.** Single-cell RNA-seq data from genetically labelled (via LoxP-reporter and Prrx1 limb enhancer driving expression of the Cre recombinase) FACS sorted connective tissue cells from 3, 5, 8, 11 and 18 dpa limb blastemas[17] were analysed as follows: read counts were normalised and used as input for principal component analysis. The top twenty loadings of the positive and negative axes of the first nine principal components were inspected to identify principal components that segregated proximodistal markers on separate ends. Principal components 3, 5 and 9 in timepoint 11 dpa had the largest distance between MEIS2 (proximal) and HOXA13 (distal), followed by PC6 and PC3 of 18 dpa. We therefore adopted PC3 of 11dpa as a model for the proximodistal axis based on single-cell RNAseq data.

Tig1 cloning

For cell localisation studies, *A. mexicanum* (*Am*) Tig1 and *H. sapiens* Tig1 were amplified and cloned into a pEGFP-N2 vector (Clontech) in which the eGFP sequence was removed and a myc tag added in either C- or N- terminal positions, using standard cloning techniques. For expression studies, Am TIG1 or Prod1 were cloned into a pEGFP-N2 vector (Clontech), replacing the EGFP sequence. The p53-N2 vector was previously described (Yun et al, 2013). The *Am* TIG1^{P155A} mutant version was generated using site-directed mutagenesis kit (New England Biolabs). For baculovirus production, *Am* Tig1^{wt} and *Am* Tig1^{P155A} cDNA sequences were coupled with a T2a element followed by reporter gene, EGFP or mCherry, and inserted into a pOEMGED baculovirus-rescuing vector (Oliveira et al 2018). Two fragments corresponding to Tig1 gene and T2a-mCherry sequences were amplified from pN2-TIG1-myc vectors using standard PCR techniques, and integrated into a BmtI and EcoRI-HF-digested pOEMGED plasmid via Gibson assembly method (NEB), with 1:3:3 plasmid to amplicon ratio. The Tig1-T2a-EGFP cassette was then transferred to a small transfection vector derived from a SacI and SphI-HF – digested pGEMT plasmid and ligated using classic cloning techniques. The same strategy was used to generate pOEMGED and pGMT-derived vectors containing the Tig1^{P155A} sequence.

Displacement assay

Forelimbs of 4 cm (snout-tail length) axolotls were amputated at the distal end of the humerus. At 4 days post amputation, the distal region of blastemas was co-injected and electroporated with a reporter plasmid (pEGFP-N2 or pRFP-N2; Clontech) together with the Tig1, Tig1^{P155A}, Prod1 or a p53 gene plasmids alone or in combinations as indicated (Fig. 3, Supplementary

Fig. 6, 8 and 9), using a 1:3 reporter plasmid to plasmid or mix ratio, with a final DNA concentration of 2 µg/µl. Animals were carefully placed in a tightly fit 2% agarose mold in a petri dish with 0.8 PBS, and electroporated using the NepaGene Nepa II machine coupled with CUY611P3-1 electrodes (Xceltis). A bipolar poring pulse of 70v for 5ms and subsequent 5 bipolar transfer pulses of 50v for 50 ms, with an interval of 999ms and 5% decay were applied. Gene expression was accessed at 1, 7, 12, 28 dpe using light microscopy. Scoring for the localisation of cells according to upper arm, lower arm and hand levels used to calculate the frequency of displacement phenotypes was done manually. All samples were analysed at endpoint (28dpe). This analysis took into consideration any identifiable cell observed at a given PD level.

Image analysis of cell distribution along the PD axis
The spatial distribution of electroporated cells and their clones along the proximal-distal (PD) axis of the regenerating axolotl limb was quantified by using Meandros, a Python 3-based image analysis tool that we designed to extract intensity profiles from fluorescent signals spread over non-linear tissue shapes (full methods will be described elsewhere). Through a user-friendly graphical user interface (GUI), Meandros allows simultaneous handling of multiple images, analyse them and run statistical tests on the flight to finally export the analysed data and plots. Briefly, we used the following modules from Meandros: I) ROI detection: Meandros automatically detected the initial ROI contours delimiting the borders of the regenerating limb from the bright filed of the microscopy images by using a deep neural network approach[43], using the Mask R-CNN solution[44] as architecture and utilising ResNet101[45] as backbone. When necessary, we fine-tuned this initial automatic guess by using the Ramer–Douglas–Peucker (RDP) algorithm to manually correct the ROI contours and export the corrected limits with Bézier curves. II) PD axis determination: By using the image bright field, we identified the PD axis by drawing a Bézier curve crossing the amputation plane, passing by the elbow and the wrist joints together with the digits tip. We arbitrarily settled the amputation plane and the distal extreme of the longest digit tip as 0 and 100 % of the PD axis, respectively, and we annotated the elbow and wrist joints as anatomical landmarks for the ulterior analysis. III) Thresholds and exclusion areas: By using the image channels depicting the fluorescent signal of interest, we selected the image-dependent intensity threshold discriminating from the background the pixels that later were quantified. When necessary, we excluded from the analysis image areas in which unspecific signal was present, such as bone autofluorescence or tissue heterogeneity-derived light diffraction artefacts, by enclosing them within a polygon. IV) Collapsing 2D images into a 1D axis: We collapsed the 2D image information into the 1D PD axis obtained for each image. To achieve this, Meandros determined the orthogonal direction to the tangent line passing by each pixel of the PD axis. The tangent line in the pixel i from the PD axis was calculated from the least square line fitted to the $i - j$, $i - j - 1$, … , i, …, $i + j - 1$, $i + j$ pixels, where j was set in its default value of 20. Thus, for each pixel i of the PD axis, Meandros determined the number of pixels belonging to the orthogonal line intersecting the domain within the ROI (w_i), and the frequency of them with intensity higher than the previously settled threshold (f_i). Observables: We quantified each PD profile as the spatial distribution along the PD axis coordinate i of the ratio between each f_i and its maximal value along the PD axis f_{max} (f_i/f_{max}). We also quantified the local limb width profile as each w_i along the coordinate i of the PD axis.

Engulfment assay
To generate blastemas of distinct proximal-distal values, forelimbs of 8.5 cm (snout-vent length) animals were amputated either at the stylopod at the distal end of humerus, or at the

distal end of radius/ulna level. At 12 days post-amputation, blastemas along with 2 mm of the adjacent stump were collected and washed twice in A-PBS (80 % 1X PBS). The epidermis was carefully removed with the help of fine forceps.

When using exclusively DD white axolotls, one blastema of each pair was labelled with Fast-DiO or CM-DiI at a concentration of 0.15 mg/mL and 0.25 mg/ml respectively, diluted in serum-free MEM and incubated for 30 min at 25 ˚C, and finally washed twice with A-PBS. In the other cases, the pairs consisted in one white and one GFP (Caggs:EGFP[46]) -derived blastema.

The stump segment was removed, thereby creating a basal surface corresponding to the proximal-most area of the blastemas. Blastema pairs were then transferred to wells of 24 or 96 multi-well plates, and juxtaposed at their basal surface. Space in between wells was filled with water to maintain a moist environment, and the plate was incubated at 25 ˚C for 20 minutes. Blastemas retain a thin PBS layer on its surface that prevents them from drying for such a short period of time. MEM supplemented with 5 % FBS was carefully added to the wells to cover the blastema pair, and incubation proceeded for 7 days at 25 ˚C, 5 % CO_2. Images were acquired in Zeiss Axio Zoom V16 microscope (Zeiss).

Treatment with antibodies against *Am* TIG1, *Am* PROD1 or control IgG rabbit antibody (antibody section) was performed at a final concentration of 20 µg/ml in MEM supplemented with 5 % FBS. The solution was added to the juxtaposed blastema pairs after a 20 minute incubation period, and maintained until the end of the experiment.

Baculovirus-mediated transduction
Baculovirus-mediated transduction of axolotl tissues was performed as described elsewhere[24]. Briefly, Baculovirus was pseudotyped with *vsv-ged* gene, inserted into the rescue vector under the baculovirus polyhedrin promoter. Genes of interest (Tig1[wt]-T2a-EGFP, Tig1[P155A]-T2a-EGFP) were cloned into the baculovirus rescue vector under the *CMV* promoter using standard restriction enzyme methods. Pseudotyped baculoviruses carrying the gene of interest were generated by co-transfection of the rescue vector together with replication-incompetent baculovirus DNA into a modified *Spodoptera frugiperda* cell line (expresSF+, ProteinSciences Corporation). Upon culture expansion, recombinant baculoviruses were collected, concentrated and purified. Baculovirus titre was assessed by end-point dilution assay in SF-9 Easy Titer cells, a modified *Spodoptera frugiperda* cell line expressing a reporter *egfp* gene under the control of the baculovirus polyhedrin promoter.

Expression of the genes of interest in axolotl blastemas was achieved by injecting the corresponding baculovirus (2 µl each; Bv titre: c. $1x10^{12}$) into the upper arm blastemas as indicated. Injections were performed using a glass capillary needle pulled on a micropipette puller (Sutter Instrument Co. model P-97). Glass capillaries (1.2 mm OD, 0.9 mm ID borosilicate glass) were acquired from Harvard Apparatus.

Cell culture
Axolotl AL1 cells (initially derived by S. Roy, University of Montreal, Montreal, Canada) were grown on 0.75 % gelatine-coated plastic dishes in MEM (Gibco, UK) supplemented with 10 % heat-inactivated foetal calf serum (FCS, Gibco), 25 % H2O, 2 nM L-Glutamine (Gibco), 10 µg/ml insulin (Sigma, St Louis, MO) and 100 U/ml penicillin/streptomycin (Gibco) in a humidified atmosphere of 2.5 % CO2 at 25 °C. Cell subculture was performed as previously described[47]. 293T cells were grown on plastic dishes in DMEM (Gibco) supplemented with 10 % heat-inactivated foetal calf serum (FCS, Gibco), 2 nM L-Glutamine (Gibco), and 100 U/ml penicillin/streptomycin (Gibco) in a humidified atmosphere of 2.5 % CO2 at 37 °C. Retinoic acid (RA; all-trans retinoic acid, 50mg/ml stock solution, Sigma) treatment was performed as described[48].

AL1 cell electroporation

AL1 cell suspensions (10^5 cells in 0.1 ml) were mixed with 2 μg of the indicated constructs (pN2-amphiregulin-myc, pN2-myc-amphiregulin, pN2-Tig1-myc, pN2-myc-Tig1) and nucleofected using a Nucleofector II (Lonza) as per manufacturer's instructions. The nucleofected cells were added to 1.5 ml supplemented minimal medium (MEM, Gibco) and incubated at 25 °C for 20 minutes prior to plating into gelatin-coated dishes. Expression of myc-tagged Tig1 was confirmed through Western blot analysis, as previously described[25]. Live staining was carried out at 48 hours post nucleofection.

293T lipofection

293T cells were grown to 70 % confluency and lipofected using lipofectamine 2000 (ThermoFisher). Briefly, 8 μl Lipofectamine solution were mixed with 200 μl OPTI-MEM (Gibco), incubated for 5 minutes at room temperature and added to a pre-mix of 3 μg of the indicated constructs plus 200 μl Opti-MEM. After a 20 minute incubation, the 400 μl solution was added to the 293T cells, in their own growth media (1.5 ml), drop-by-drop. Cells were incubated for 24 hours prior to live staining.

qRT-PCR

RNA was isolated from axolotl AL1 cells or indicated tissues using Tri Reagent (Sigma) and random primed cDNA synthesised using Superscript II (Invitrogen). Gene expression was determined by quantitative real time PCR using the indicated primers (Supplementary table 1). RT-PCR was carried out using iQ SYBR Green supermix (Bio-rad, Hercules, CA) on a Chromo 4 instrument running Opticon 3 software (Bio-rad). All reactions were run in triplicate and at least 3 independent RNA preparations were analysed for each sample.

For FACS sorted blastema cells: three biological replicates per condition of the libraries used for the RNA seq experiment plus 3 additional replicates, prepared in the same way (see below), served as basis for the qPCR to confirm gene expression changes caused by Tig1 overexpression. The minimum number of biological replicates for each condition was 5. qPCR was carried out using iQ SYBR Green supermix (Bio-rad, Hercules, CA) on a Chromo 4 instrument running Opticon 3 software (Bio-rad). Gene expression was normalised to the ribosomal protein Rlp4. For visualising individual genes, all conditions were normalised to the mean value of the control condition at 3 dpe.

EdU incorporation assays

To determine the percentage of cells in S-phase, AL1 cells were sub-cultured and incubated in growth medium supplemented with 5 μM 5-ethynyl-2'-deoxyuridine (EdU) for 24 hours. Cells were then fixed in 4 % PFA for 10 minutes and stained using Click-iT Edu Alexa Fluor 594 Imaging kit (Life Technologies) according to manufacturer's instructions.

To detect EdU incorporation in salamander tissues, 10 mM EdU (20 μl per animal) were administered by i.p. injection at 24 and 48h after electroporation. Blastemas were collected 72h post-electroporation, fixed O/N in 4 % PFA, embedded and cryosectioned. EdU incorporation was determined using Click-iT Edu Alexa Fluor 488 Imaging kit (Life Technologies) on tissue sections. Estimation of EdU incorporation was performed by calculating the number of EdU$^+$ cells over the total number of transduced cells (RFP$^+$).

Generation of C-Ti$^{t/+}$(*HoxA13:HoxA13-T2a-mCherry*)^{ET} transgenic axolotls

The endogenous HOXA13 protein was tagged at the C-terminus according to the published protocol[26]. Briefly, a portion of the Hoxa13 gene including a part of the single intron and the remaining downstream part of the CDS was amplified from the axolotl genomic DNA and

inserted into the vector pGEM-T along with a DNA fragment encoding T2a-mCherry-3xnls using Gibson assembly cloning (BioLabs). The resulting vector was injected into fertilised eggs along with the CAS9 protein (MPI-CBG) and the gRNA targeting the sequence GGGGCTTTTGCGGGTTTTCC, which is present in the intron sequence within the vector and in the genome. The insertion is mediated by NHEJ and is thus expected to cause a small InDel within the intron, but it will result in a seamless Hoxa13 CDS. T2a-mCherry is followed by the pGEM-T poly(A) signal and the rest of the vector.

Validation: Ti$^{t/+}$(*HoxA13:mCherry*) axolotls exhibit expression of mCherry in the distal part of the developing limb bud throughout limb formation (Supplementary fig. 10). During regeneration, mCherry expression is observed in the distal portion of the blastema from mid-bud onwards, coinciding with the appearance of the HoxA13 domain[23] and remaining until digit stage. The animals used in the experiment were heterozygous for the T2a tag, to overcome any potential loss of function that could be originated from the knock-in.

Antibodies

Polyclonal anti *Am* TIG1 antibodies were manufactured by Eurogentec, by raising rabbit antisera against two non-overlapping peptides corresponding to residues MAQVKSVKQRLRND (154) and GSSGGRSEEGSASF (155) of the full length axolotl TIG1 protein, using KLH as carrier. Both reagents were analysed by solid phase ELISA and affinity purified using their respective immobilised peptides (Supplementary data 2 and 3). Note that while these reagents work for Elisa and western blotting, we did not manage to find conditions suitable for immunofluorescence applications. The two affinity-purified anti-PROD1 antibodies were generated as previously described[11]. Anti-myc (mouse monoclonal 9E10, Sigma-Aldrich) and anti-RFP (Rockland) were also used in this study.

Live cell staining and immunofluorescence

For live cell staining, 293T or AL1 cells were incubated for 3 hours with anti-Myc mouse monoclonal antibody (Sigma-Aldrich; 1:500), and secondary staining was performed using anti-mouse AlexaFluor568 antibodies (Invitrogen; 1:1000). Hoechst 33258 (2 µg/ml) was used for nuclei counterstaining. Samples were observed under a Zeiss Axioskop2 microscope and images were acquired with a Hamamatsu Orca camera using Openlab (Improvision) software. Whenever comparative analyses between samples were performed, all images were acquired with identical camera settings and illumination control. Image processing (contrast enhancement) was equally applied to all matched experimental and control samples using Openlab software.

Tissue dissociation for RNAseq

Limbs of HoxA13:Cherry transgenics (4.5 cm snout-vent length) were amputated at the stylopod level, near the elbow. The resulting blastemas were electroporated with Tig1-T2a-GFP or Tig1^{P155A}-T2a-GFP plasmids at 13 dpe, and harvested at 3 or 8 days post electroporation. 12 blastemas were pooled per condition, washed in 0.8 PBS, and incubated in 500 µl of 1x Liberase™ research grade (Sigma-Aldrich) for 5 minutes at room temperature. The epidermis was removed and the blastema mesenchyme dissociated with the help of very fine forceps. The incubation proceeded at room temperature for 30 minutes with rocking motion at 40 rpm. Cells in solution were gently pipetted up and down, filtered through a 70 µM Filcon (BD™ Medimachine) and collected into a falcon tube. Sytox blue (Thermo-Fisher) was added at 1:2000 and the solution was kept in ice until FACS sorting. After sorting, cells were kept in 1x freshly-prepared lysis buffer containing RNAse inhibitor, and kept at -80 °C. cDNA library preparation from 300 cells per sample (3 replicates per condition) followed by

SMARTseq2 sequencing was performed by CMCB-Deep sequencing facility/Dresden-concept Genome Center.

Bioinformatic analysis

A putative Tig1 promoter region corresponding to 40Kb upstream the start site (axolotl-omics.org) was analysed using the web interface https://molotool.autosome.ru/.

Differential gene expression analysis was performed using DESeq2 package[49]. The iDEP.92 application[50] was used for K-means clustering and GO enrichment. From the initial 44224 genes in 18 samples, 19761 genes passed the filter (0.5 counts per million in at least one sample), 12917 were converted to Ensembl gene IDs present in the iDEP database. The remaining 6844 genes were kept in the data using original IDs. K-means clustering across all samples was performed using top 2000 variable genes as ranked by standard deviation. Number of clusters, k=3 was chosen based on elbow and tSNE method. Pathway enrichment analysis for each cluster was conducted based on Gene Ontology (GO) Biological Process database[51]. P-values are corrected for multiple testing using false discovery rate (FDR). Gene correlations in Fig 6 and Supplementary Fig. 15 were carried out through R, using the scRNAseq dataset reported in Gerber et al.

Statistical analysis

Animals in each sample group were randomly selected. Sample group size (n) is indicated in each figure legend, while all experiments were carried out in at least three biological replicates. Statistical analyses were performed with Prism 4.0 software. The method used for each analysis is indicated in the corresponding section.

Fig 1

Supplementaty Fig. 1

Supplementaty Fig. 2

a

pEGFP-N2 pN2-Tig1-C-myc pN2-Tig1-N-myc

37kDa —

25kDa —

*

b

C-terminus   N-terminus

pN2-Amphiregulin

pN2-TIG-1

Supplementaty Fig. 3

a

pN2-RFP   pN2-RFP +
            pN2-Tig1 or Tig1^P155A

24h → IP EdU

48h → cryosectioning
+ anti-EdU/anti-RFP

IP EdU

b

pN2-RFP   pN2-Tig1+pN2-RFP   pN2-Tig1^P155A+pN2-RFP

EdU/RFP/Hoestch

c

% EdU$^+$/RFP$^+$ cells

pN2-RFP   pN2-RFP+   pN2-RFP+
              pN2-Tig1   pN2-Tig1^P155A

**   **

Supplementaty Fig. 4

Supplementaty Fig. 5

Fig 2

a

| | RFP | PROD1 | TIG1 WT | TIG1^{P155A} |

1 dpe

7 dpe

24 dpe

b

% limbs with cells in the indicated segment

Digits   Carpals   Lower arm   Upper arm

TIG1^{P155A}
PROD1
TIG1wt
RFP

b

PD Axis

W
A E

P ⟵⟶ D

A   E   W

$f/fmax$ (%)

PD position (%)

d

$f/fmax$ (%)

A   E   W

PD position (%)

— *Control*
— *Prod*1
— *Tig1^{wt}*
— *Tig1^{P155A}*

Fig 3

Supplementaty Fig. 6

Supplementaty Fig. 7

a

b

Supplementaty Fig. 8

Supplementaty Fig. 9

a

D/D
🟢🟢 → 🟢🟢

P/P
🔴🔴 → 🔴🔴

P/D
🔴🟢 → engulfment

P/D   α-PROD1
🔴🟢 → 🔴🟢

P/D   α-TIG1
🔴🟢 → ?

b

	24h	96h	168h
α-RFP			
α-TIG1			
α-PROD1			

c

blastema pair	Engulfment	
	Frequency	%
PD	12/17	70.5
PP	1/7	14.2
DD	0/3	0.0
PD + α RFP	6/8	75.0
PD + α TIG1	0/10	0.0
PD + α PROD1	3/10	30.0

Fig 4

Fig 5

Supplementaty Fig. 10

Supplementaty Fig. 11

a

adj.Pval	nGenes	Pathways
8.50E-90	325	Immune system process
1.40E-80	266	Immune response
6.30E-74	203	Cell activation
4.50E-71	188	Leukocyte activation
6.90E-58	128	Myeloid leukocyte activation
1.70E-54	165	Immune effector process
3.50E-51	189	Regulation of immune system process
2.50E-48	120	Leukocyte activation involved in immune response
1.40E-44	125	Leukocyte mediated immunity
1.50E-44	102	Leukocyte degranulation

GO Analysis - Cluster B

b

adj.Pval	nGenes	Pathways
5.60E-18	14	Muscle filament sliding
5.60E-18	14	Actin-myosin filament sliding
1.30E-17	60	Tissue development
2.60E-15	17	Actin-mediated cell contraction
1.90E-13	17	Actin filament-based movement
4.80E-13	13	Myofibril assembly
7.30E-12	31	Actin filament-based process
9.30E-12	40	Epithelium development
1.00E-11	31	Epithelial cell differentiation
1.00E-11	16	Striated muscle cell development

GO Analysis - Cluster C

Supplementaty Fig. 12

Supplementaty Fig. 13

Fig 6

gene	Forward (5' primer)	Reverse (3'primer)
Rlp4	TGAAGAACTTGAGGGTCATGG	CTTGGCGTCTGCAGATTTTTT
Hoxa13	ACTGGTTCATGAAGACAATTGCCT	CTGTTTTCACCGTCTTCCCCATTA
Hoxa11	GCCAAACAGGCTTGGTTTTA	CTTCCGGCGATTGTTTAGTC
Hoxa9	GAGACAAGCCTGCCATTGAC	GGTTCTGGAACCAGATCTTGAC
Meis1	GCCGTCGCCAACAATGATTT	GTGATGCGTACGGTTGATGC
Enpp2	CGATACCAGTCCCAACGCAT	GAGTTGCTCAATGTCCCGGA
Epha7	CGTTACAATCGGCATGAGGG	GCTAGGCCTTACTTCCCGTG
Lhx2	CCTCTTTCAAACACCACCAGTTG	TGTGTTTTCTTGGCGTAGGAGAT
Prod1	GATCTCAAAGGTCGGTTCAGA	CACAAGCCTGCTACTTCTAGA
Tig1CDS	TCCAATGAGCCCCTGAGAGA	GACACTCTTCACTTGGGCCA
Tig1-3'UTR	CTCCCTCAAACAATCCGGCT	GTTGTCGAGCGCTTCCATTC
ef1-α	AACATCGTGGTCATCGGCCAT	GGAGGTGCCAGTGATCATGTT

Supplementaty Table 1

ANTIBODY SERVICE

ANTIBODY PURIFICATION ANALYSIS BY INDIRECT ELISA

Customer :   **Dr.   YUN MAXIMINA**

Peptide :	EP104154		*S:*	Serum
Rabbit	1499		*FT:*	Flow-through
			PA:	Purified antibodies

2- Results

Dilution (x-fold)		Against Peptide									Against Carrier		
		serum (S)			col. flow-through1 (FT)			purif. Antibodies (PA)			S	FT	PA
		1	2	3	4	5	6	7	8	9	10	11	12
100	A	3,118	3,136	3,121	0,382	0,357	0,400	3,240	3,437	3,498	3,815	3,802	1,158
300	B	1,527	1,499	1,281	0,366	0,354	0,369	2,427	2,545	2,985	3,771	3,782	0,657
900	C	0,712	0,608	0,546	0,342	0,333	0,249	1,109	1,269	1,559	3,770	3,769	0,490
2700	D	0,490	0,354	0,320	0,256	0,292	0,249	0,719	0,586	0,779	3,727	3,733	0,392
8100	E	0,408	0,321	0,279	0,235	0,260	0,239	0,438	0,459	0,503	3,562	3,447	0,306
24300	F	0,435	0,281	0,317	0,231	0,241	0,232	0,375	0,415	0,397	3,025	2,857	0,308
72900	G	0,285	0,280	0,367	0,219	0,237	0,206	0,318	0,360	0,363	0,064	0,062	0,055	- contr.
218700	H	0,266	0,223	0,337	0,204	0,234	0,205	0,299	0,353	0,295	3,693	3,686	3,695	+ contr.

Peptide (straight line)   carrier (discontinous line)

Visit   **www.eurogentec.com/ELISA-interpretation.html**   for ELISA results interpretation

ANTIBODY SERVICE

SDS-PAGE ANALYSIS (Bioanalyser, Agilent)

Customer : **Dr.   YUN MAXIMINA**

Peptide : EP104154

Rabbit 1499

Size [kDa]	% Total	Observations
4,5	0,0	Lower Marker
8,2	0,0	System Peak
14,8	1,3	
25,5	23,7	LC
56,4	75,0	HC
240,0	0,0	Upper Marker

Purity (sum HC+LC)   **98,7 %**

LC:	Light Chain
HC:	Heavy Chain
rSA:	rabbit Serum albumin

Supplementary data 2

ANTIBODY SERVICE

ANTIBODY PURIFICATION ANALYSIS BY INDIRECT ELISA

Customer :　**Dr.　YUN MAXIMINA**

Peptide :	EP104155		*S:*	Serum
Rabbit	1499		*FT:*	Flow-through
			PA:	Purified antibodies

2- Results

Dilution (x-fold)		Against Peptide									Against Carrier		
		serum (S)			col. flow-through1 (FT)			purif. Antibodies (PA)			S	FT	PA
		1	2	3	4	5	6	7	8	9	10	11	12
100	A	3,460	3,461	3,417	0,712	0,949	0,725	3,664	3,681	3,661	3,805	3,781	0,994
300	B	2,361	2,236	2,219	0,471	0,491	0,425	3,632	3,654	3,662	3,790	3,758	0,770
900	C	1,201	0,984	1,036	0,421	0,489	0,424	2,992	3,085	3,128	3,764	3,758	0,558
2700	D	0,733	0,626	0,588	0,394	0,439	0,422	1,788	1,809	1,910	3,738	3,715	0,472
8100	E	0,635	0,616	0,396	0,382	0,398	0,390	0,886	0,885	0,944	3,658	3,544	0,448
24300	F	0,486	0,399	0,393	0,349	0,381	0,382	0,572	0,556	0,538	3,395	3,169	0,359
72900	G	0,376	0,381	0,389	0,347	0,331	0,330	0,319	0,370	0,326	0,058	0,053	0,062	- contr.
218700	H	0,279	0,259	0,262	0,280	0,283	0,329	0,329	0,287	0,298	3,646	3,621	3,631	+ contr.

Peptide (straight line)　　　　　　　　carrier (discontinous line)

Visit　**www.eurogentec.com/ELISA-interpretation.html**　　for ELISA results interpretation

ANTIBODY SERVICE

SDS-PAGE ANALYSIS (Bioanalyser, Agilent)

Customer :   **Dr.   YUN MAXIMINA**

Peptide :   EP104155

Rabbit   1499

Size [kDa]	% Total	Observations
4,5	0,0	Lower Marker
8,3	0,0	System Peak
25,3	13,0	LC
28,0	2,2	LC
56,8	79,1	HC
63,2	5,7	
240,0	0,0	Upper Marker

Purity (sum HC+LC)   **94,3 %**

LC:	Light Chain
HC:	Heavy Chain
rSA:	rabbit Serum albumin

Supplementary data 3