Visualizing cellular and tissue ultrastructure using Ten-fold Robust Expansion Microscopy (TREx)

Hugo G.J. Damstra[1], Boaz Mohar[2], Mark Eddison[2], Anna Akhmanova[1], Lukas C. Kapitein[1*], Paul W. Tillberg[2*]

[1]Cell Biology, Neurobiology and Biophysics, Department of Biology, Faculty of Science, Utrecht University, Utrecht, The Netherlands

[2]Janelia Research Campus, HHMI, Ashburn, Virginia 20147, USA

Correspondence: l.kapitein@uu.nl; tillbergp@janelia.hhmi.org

ABSTRACT

Expansion microscopy (ExM) is a powerful technique to overcome the diffraction limit of light microscopy

that can be applied in both tissues and cells. In ExM, samples are embedded in a swellable polymer gel to

physically expand the sample and isotropically increase resolution in x, y and z. The maximum resolution

increase is limited by the expansion factor of the polymer gel, which is four-fold for the original ExM

protocol. Variations on the original ExM method have been reported that allow for greater expansion

factors, for example using iterative expansion, but at the cost of ease of adoption or versatility. Here, we

systematically explore the ExM recipe space and present a novel method termed Ten-fold Robust

Expansion Microscopy (TREx) that, like the original ExM method, requires no specialized equipment or

procedures to carry out. We demonstrate that TREx gels expand ten-fold, can be handled easily, and can be

applied to both thick tissue sections and cells enabling high-resolution subcellular imaging in a single

expansion step. We show that applying TREx on antibody-stained samples can be combined with off-the-

shelf small molecule stains for both total protein and membranes to provide ultrastructural context to

subcellular protein localization.

INTRODUCTION

Expansion Microscopy (ExM) circumvents the diffraction limit of light microscopy by physically expanding the specimen four-fold in each dimension (Chen, Tillberg and Boyden, 2015; Tillberg *et al.*, 2016). While other super-resolution approaches require specialized optics (Hell and Wichmann, 1994), fluorophores (Rust, Bates and Zhuang, 2006), or software (Gustafsson, 2000), and are not readily compatible with thick tissue slices, ExM is compatible with any microscope (Zhang *et al.*, 2016; Gao *et al.*, 2019), including combinations with other super resolution modalities (Halpern *et al.*, 2017; Gao *et al.*, 2018; Xu *et al.*, 2019), and performs well in both cultured cells and thick tissue slices (Chen, Tillberg and Boyden, 2015; Tillberg *et al.*, 2016). The resolution increase of ExM depends on labeling efficiency, anchoring efficiency, and the expansion factor of the gel recipe used. Recently, method variants have been described that increase the expansion factor to increase the maximum resolution obtainable with ExM. For example, iterative ExM (iExM) uses sequential embedding in multiple expansion gels to achieve 15x and greater expansion but requires a complex sequence of gel re-embedding, link cleaving, and fluorophore transfer (Chang *et al.*, 2017), limiting broad adoption in biological labs.

The expansion factor of the gel itself can be increased by decreasing the concentration of crosslinker (Okay, 2009), usually bisacrylamide (bis), at the expense of gel mechanical integrity. For example, reducing the bis in the original ExM recipe from 1.5 ppt (parts per thousand) to 250 ppm (parts per million) produces a ~9x expanding gel (Chen, Tillberg and Boyden, 2015, SF5), but these gels are too soft to hold their shape under the force of gravity. As a result, they are difficult to handle without breaking and display non-uniform expansion. This tradeoff of expansion versus gel mechanical integrity has not been explored in a quantitative or systematic way.

Another gel recipe variant, using a high concentration of the monomer dimethylacrylamide (DMAA), has enough crosslinking through side reactions and polymer chain entanglement that the crosslinker can be omitted entirely, producing ~10-fold expansion in one step (Truckenbrodt *et al.*, 2018). This recipe has been used to expand cultured cells and thin cryosectioned tissue (Truckenbrodt *et al.*, 2019), but this recipe reportedly requires rigorous degassing to remove oxygen prior to gelation, making it laborious to adopt. Moreover, expansion of >50 μm thick tissue slices has not been demonstrated using this method. Thus, a robustly validated and easily adoptable method that is compatible with multiple sample types and enables single step expansion well over 4x without compromising gel integrity is lacking.

Here, we explored the expansion gel recipe space in a systematic manner, assessing the limits of single-round expansion using reagents and methods that would be familiar to labs already doing standard ExM. For any given choice of recipe parameters (monomer concentrations, gelation temperature, radical

initiation rate, etc.), varying the crosslinker alone yielded a family of recipes whose expansion factor and

mechanical quality vary smoothly from high expanding, poor quality to low expanding, excellent quality

gels. A range of crosslinker concentrations was tested because the optimal crosslinker concentration may

be different for each recipe family. Here we present an optimized ExM method that allows for robust ten-

fold expansion in a single step, termed TREx. We show TREx can be used to expand thick tissue slices and

adherent cells and is compatible with antibodies, as well as off-the-shelf small molecule stains for total

protein and membranes. Together, we show how TREx enables 3D nanoscopic imaging of specific

structures stained with antibodies in combination with cellular ultrastructure.

RESULTS

To systematically explore the expansion recipe space, we developed a streamlined method for

synthesizing dozens of gel recipes and characterizing their mechanical quality in parallel. For every set of

gel recipe parameters (component concentrations and gelation temperature, listed in Fig. 1A) we define a

recipe family as the set of recipes generated by varying the crosslinker (bisacrylamide) concentration. For

each family, we tested five recipes with crosslinker concentrations log-spaced from 1 ppt (parts per

thousand or mg/mL) to 10 ppm (parts per million), plus one with zero crosslinker. We also included the

original ExM recipe, with 1.5 ppt crosslinker. For each recipe, we cast three gel specimens, expanded them

fully in water, and measured the expansion factor (Fig. 1B). We found that resistance to deformation under

the force of gravity was a good proxy for the more subjective judgement of ease of gel handling. We

measured gel deformation by placing a semicircular punch from each gel upright on its curved edge and

allowing the gel corners to deflect under the force of gravity. We defined the deformation index as the

vertical displacement of each gel corner, divided by the gel radius (Fig. 1C), which ranges from 0 (for gels

that do not deform) to 1 (for gels that deform freely under their own weight). We manually calibrated this

measurement, finding that deformation indices between 0 and 0.125 corresponded to gels with excellent

ease of handling, 0.125 to 0.25 corresponded to acceptable ease of handling, and anything higher than 0.25

was unacceptable. While not as theoretically informative as elastic modulus and yield strength

measurements, this measurement can be repeated and extended to new gel recipes by any lab developing

expansion methods, without access to specialized equipment. We plotted the deformation index for each

gel as a function of its expansion factor (Fig. 1D).

Development of the TRex gel recipe

We began by characterizing a recipe family generated from the original ExM recipe (family A).

Consistent with (Chen, Tillberg and Boyden, 2015), reducing the crosslinker to 300 ppm increased the

expansion to ~9x, below which the gels fail to form consistently (Fig. 1B, purple). We next characterized a

high-monomer recipe family adapted from the 4x-expanding Ultra-ExM recipe (Gambarotto *et al.*, 2019)

(family B). Gambarotto et al found that a higher monomer concentration relative to the original ExM recipe

was necessary for high fidelity preservation of the shape of centrioles. This was offset with a lower

crosslinker concentration of 1 ppt bisacrylamide to achieve 4x expansion. Indeed, for this high-monomer

family of recipes, expansion as a function of crosslinker concentration was shifted leftward compared to

standard ExM (Fig. 1B, blue). As the crosslinker was decreased below 30 ppm the increase in expansion

factor saturated around 11.5x. The deformation index versus expansion factor curve for the high monomer

family ran below that for standard ExM, indicating that for a given expansion factor a high-monomer gel

holds its shape better than the corresponding standard ExM gel (Fig. 1D, blue).

Compared with standard ExM, this high monomer family uses a higher concentration of radical

initiator and accelerator to trigger polymerization (5 ppt each of APS and TEMED, versus 2 ppt in standard

ExM), which causes gels to form within minutes at room temperature. Because the rates of initiation and

polymerization increase with temperature, it is likely that specimens are not fully equilibrating to the

gelation temperature of 37 °C before the onset of gelation, introducing a potential source of experimental

variability. This rapid gelation makes the gelation chamber assembly step more time sensitive and presents

challenges for adapting the technique to thick tissue slices, as thick tissue slices require extra time for the

gelation solution to diffuse throughout the sample prior to polymerization. We tested the same high

monomer recipe family but with initiator reduced to 1.5 ppt (family C). For all families, the accelerator

(TEMED) was varied in tandem with the initiator. The expansion versus crosslinker curve for this family

was similar to family B for high crosslinker concentrations but displayed a slightly greater slope. Unlike

family B, the expansion factor did not saturate upon decreasing the crosslinker concentration. Instead, the

expansion factor continued to increase to 13x expansion at 10ppm crosslinker (Fig. 1B, red), with gels

failing to form without any crosslinker present. While deformation of these gels for the same expansion

factor was increased compared to family B, although better than for family A and acceptable up to an

expansion factor of 10 (30 ppm bis, Fig. 1D, red), in our judgement this was outweighed by the slower and

more controlled gelation kinetics achieved by lowering the concentration of initiator and accelerator.

The recipe families explored above (B, C) feature a high fraction of sodium acrylate relative to

acrylamide. Acrylate drives expansion of the gel but comes in widely varying purity levels and, we found

in some cases, causes tissue to shrink. The tissue shrinkage was modest compared to the gel expansion but

may not be uniform at all scales. We therefore tested an alternative recipe family (D) with higher acrylamide

to acrylate ratio (2.1:1). Increasing the acrylamide to acrylate ratio did not change the expansion factors

appreciably at a given crosslinker concentration, suggesting that the swelling effect of acrylate saturates at

high concentrations. At the maximum expansion factor of ~10, the deformation behavior was comparable

to family C, which biased our choice to family D due to the lower acrylate content of the gels.

We further tested an elevated gelation temperature of 50 °C (family E), in an attempt to increase

the initiation rate without introducing premature gelation at room temperature seen in recipe family B.

Compared to family D, the expansion factors were around 15% higher at 100 ppm (6x) and 300 ppm

crosslinker (9x), but gelation failed at lower concentrations, leaving family D as the family with a higher

maximum expansion factor (i.e., 10x at 30 ppm bis). The deformation versus expansion curve for family E

was similar to the other high monomer recipe families (Fig. 1D, green), but was found to be sensitive to

processing details, such as the gelation chamber construction and placement within the incubator. This

suggests that premature gelation prior to equilibrating at the higher temperature reduces the replicability of

this recipe family.

Considering all 5 recipe families, family B (high acrylate and high APS/TEMED) displayed the

best mechanical performance, i.e. the lowest deformation index for any given expansion factor. Family D

(high acrylamide and low APS/TEMED) displayed similar performance, with the deformation index

remaining well within the acceptable range for expansion factors up to 10. In handling high-expanding

(>8x) gels from all recipe families, we found that while those from the standard ExM family (A) were

extremely prone to breaking, those from any of the high monomer families could be handled more easily

(and even dropped from a height of several feet) without breaking. Because the reduced initiator

concentration of family D results in a slower and more controlled polymerization rate, and because we

preferred a lower acrylate content, we chose this recipe family to proceed to biological specimen expansion.

We found that the exact expansion factor varied for different gelation chamber geometries but could readily

be adjust by finetuning the crosslinker concentration (e.g., 50 ppm for Ashburn, VA, USA experiments

versus 90 ppm for Utrecht, The Netherlands experiments), We thus recommend that users test gels with a

range of crosslinker concentrations between 30 and 100 ppm to find a suitable recipe for their specimen

preparation. We name the resulting method Ten-fold Robust Expansion (TREx) microscopy.

Subcellular imaging of specific proteins and cellular ultrastructure in thick brain slices

A major advantage of ExM over other super-resolution techniques is its compatibility with thick

tissue slices. Tissue slices require the onset of gelation to be delayed, usually with low temperature

incubation and a radical inhibitor (e.g. 4-Hydroxy-TEMPO; 4HT), to give the monomer solution time to

diffuse into the tissue (Chen, Tillberg and Boyden, 2015; Tillberg *et al.*, 2016) . For TREx, we found that

in some conditions room oxygen was sufficient to inhibit gelation with shaking on ice indefinitely. To

increase robustness for thick tissue slices, we included 15 ppm 4HT, allowing ample time to mount

specimens in gelation chambers at room temperature. ExM enables uniform expansion by treating

embedded tissue with strong proteolytic digestion step. Proteins (including antibodies) are crosslinked to

the gel with acryloyl-X SE (AcX), allowing for retention of protein fragments after digestion. Replacing

proteolytic digestion with a high detergent, high temperature disruption step has been shown to enable

expansion with better overall retention of proteins (Ku *et al.*, 2016; Tillberg *et al.*, 2016; Gambarotto *et al.*,

2019; Zwettler *et al.*, 2020). We found that this gentler disruption approach also allowed the AcX anchor

treatment to be reduced from 100 µg/mL overnight (standard ExM) to 10 µg/mL for 1 hour while enabling

full expansion of mouse brain slices without the appearance of crack-like artifacts.

In electron microscopy, non-specific stains for proteins and membranes are commonly used to

provide structural detail at high spatial resolution. Recently, the use of non-specific NHS ester protein stains

and other small molecule probes has been combined with ExM (M'Saad and Bewersdorf, 2020; Mao *et al.*,

2020; Yu *et al.*, 2020). Expansion allows visualization of intracellular detail in densely stained samples,

which would not otherwise lead to meaningful contrast. These applications have the promise to bring

together the advantages of light microscopy (specific staining using antibodies and volumetric imaging)

with the advantage of seeing cellular context typically provided by electron microscopy. Because TREx

reaches single-step expansion factors at which small-molecule stains are expected to be useful, we set out

to explore this idea further. We applied NHS ester dyes to demonstrate total protein content in thick slices

(100 µm) of mouse brain cortex expanded with TREx (Fig. 2A and Supp. Movie 1). The neuropil region

outside the cell somas contained a rich profusion of fibers and structures visible in sharp relief. The nucleus

of each cell was easily identified, with especially strong staining in nucleoli-like structures. Surrounding

each nucleus, the nuclear envelope could be identified, with particularly dense total protein stain on the side

facing the nucleus. The nuclear envelope was punctuated by heavily stained spots that span the envelope,

consistent with nuclear pore complexes (Fig. 2A, inset). Within the cytosol, several organelles were

visualized by either heavy inner staining with a dim border or weak inner staining.

We next tested whether antibodies delivered using a standard immunofluorescence procedure

before embedding were also retained in the TREx gel. We stained mouse brain cortex tissue for Bassoon (a

marker for both excitatory and inhibitory pre-synaptic active zones), Homer (a marker for the excitatory

post-synaptic apparatus), and VGAT (a vesicular GABA transporter in the pre-synaptic compartments of

inhibitory synapses). After staining and anchoring with AcX, tissue was expanded with TREx and imaged

by light sheet microscopy. Numerous putative excitatory synapses were observed at high density, with

clearly separated Bassoon and Homer pre- and post-synaptic staining (Fig. 2B and Supp. Movie 2). VGAT

had a more extended staining pattern, consistent with the known distribution of synaptic vesicles throughout

pre-synaptic boutons. Elaborately shaped compartments with dense VGAT staining were seen with multiple

synaptic release sites marked by Bassoon (Fig. 2B, inset). As expected, these release sites were not paired

with the post-synaptic marker Homer. These results demonstrate the ability of TREx to preserve correct

synaptic staining while enabling sub-diffraction limited imaging of large tissue sections.

Subcellular imaging of specific proteins and cellular ultrastructure in cultured cells

We next explored the use of TREx for high-resolution imaging of specific proteins, NHS stains and

lipid membranes in cultured cells. For membranes, a custom-synthesized membrane probe compatible with

the ExM process has been shown to visualize membrane surfaces in fixed brain tissue (Karagiannis *et al.*,

2019). This probe relies on a peptide-modified lipid tail that intercalates in target membranes and provides

opportunities for anchoring to the gel through D-lysines its peptide sequence. We asked whether the

commercially available membrane-binding fluorophore-cysteine-lysine-palmitoyl group (mCLING) could

also be used for membrane staining and gel anchoring. mCLING has been developed as a fixable

endocytosis marker consisting of a fluorophore and a short polypeptide group with one cysteine and seven

lysines coupled to a palmitoyl membrane anchor (Revelo and Rizzoli, 2016). Due to the presence of

multiple lysines, we hypothesized that mCLING would be compatible with standard ExM anchoring

through AcX.

While the standard protocol for mCLING delivery relies on active endocytosis in living cells, we

asked if mCLING would stain intracellular membranes more uniformly when added to fixed cells, which

would have the added benefit of not perturbing intracellular membrane trafficking by long incubation in

live cells. To test this, we fixed activated Jurkat T cells, incubated the fixed cells with mCLING overnight,

and proceeded with the TREx protocol. We found that mCLING efficiently intercalates in both the plasma

membrane and internal organelles and is retained following standard AcX treatment (Fig. 3A and Supp.

Movie 3).

By carefully rendering the imaged volumes we could, with one probe, both appreciate the ruffled

morphology of the plasma membrane on top of the flattened part of the cell, and visualize the organelle

clustering typical of activated T-cells (Fig. 3A, B). In electron microscopy, distinct morphologies are used

to identify organelles. Similarly, we found that we could clearly identify different organelles based on

mCLING, suggesting that it could be used for automated segmentation of organelles. Indeed, we found that

mitochondria could be readily segmented using a trainable Weka segmentation algorithm (Fig. 3B)

(Arganda-Carreras *et al.*, 2017). While the resolution of subcellular structures is limited by the density of

mCLING moieties in the membrane, the efficiency of crosslinking to the gel, and the maximum expansion

factor, we found TREx allows sufficient single-step expansion to resolve individual mitochondrial cristae,

which are known from electron microscopy to be spaced 70 nm apart (Fig. 3C). Although mCLING is

thought to be membrane impermeable (due to multiple positively charged amino groups), it readily stained fixed and unpermeabilized cells following extended incubation. Because this approach does not require labeling live cells and is expected to reduce differences in uptake efficiency between intracellular compartments, we used this approach in all subsequent experiments.

We next tested if mCLING could also be used to visualize membranes in more complex cell types. To test this, we used differentiated Caco-2 cells grown to form an epithelial monolayer. Using TREx, we could expand the entire monolayer and visualize the membrane using mCLING (Fig. 3D-G and Supp. Movie 4). This enabled us to resolve individual microvilli as hollow membrane protrusions within the dense brush border. In addition, we were able to resolve the elaborate interdigitated cell-cell junctions that could previously only be clearly appreciated using electron microscopy (Drenckhahn and Franz, 1986). The advantage of optical, volumetric imaging is underscored by the fact that we can easily render one dataset in several ways, either resembling scanning electron microscopy to highlight volumetric surface morphology (Fig. 3D), or transmission electron microscopy to explore single planes in more detail (Fig. 3E). Together, these data illustrate that TREx can be combined with a commercially available membrane stain to provide rapid volumetric insights into the elaborate membranous architecture of cells.

We further tested whether TREx using general membrane stains could be combined with general protein stains and/or antibody stains. U2OS cells transfected with GFP-Sec61β were fixed, treated with mCLING, stained for GFP, and expanded with TREx followed by the NHS stain (Fig. 4A). Because we performed the NHS stain after disruption, we used temperature- and detergent-mediated disruption, rather than proteolytic digestion. We found that secondary antibodies that had been used to stain GFP before gelation withstood this disruption step. We observed a clear degree of overlap between mCLING and NHS, especially in the dense perinuclear region, but we could also identify distinct features of each stain (Fig. 4A). Interestingly, we observed strong nucleoplasmic staining in the NHS stain and clear puncta where the imaging plane intersected the nuclear envelope, which presumably are nuclear pores (Fig. 4A). These results demonstrate that general stains for membranes and proteins can be combined with antibody-based labeling to reveal specific proteins in their ultrastructural context.

Previously, we have used ExM to study the three-dimensional organization of microtubules (MT) in neurons and T cells (Hooikaas et al., 2020; Jurriens et al., 2020). For high-resolution imaging of the MT cytoskeleton, cells are usually pre-extracted with detergent and glutaraldehyde to remove the soluble pool of tubulin, followed by paraformaldehyde fixation (Tas et al., 2017). This reduces background but does not preserve membranes. We reasoned that the increased expansion of TREx would dilute the background due to soluble tubulin by the expansion factor cubed. Ten-fold expanded microtubules remain diffraction limited in width (i.e., 250-350 nm), so their signal should be reduced only by the expansion factor itself

(due to the expansion along their length). Therefore, we wondered whether the resulting relative boost in

signal over background would eliminate the need for pre-extraction to enable high-resolution imaging of

microtubules in combination with membranes. To test this, we fixed cells expressing GFP-Sec61β without

pre-extraction, treated them with mCLING, stained for GFP and tubulin, and then proceeded with TREx

(Fig. 4B and Supp. Movie 5). Subsequent confocal microscopy revealed the interplay between microtubules

and ER in three dimensions and revealed how other membranous organelles were connected to both

structures (Fig. 4C). Thus, TREx facilitates high-resolution three-dimensional mapping of specific

cytoskeletal and membranous structures in combination with markers that provide ultrastructural context.

DISCUSSION

We developed Ten-fold Robust Expansion (TREx) in order to expand biological specimens 10-fold

in a single round of expansion, without specialized equipment or procedures. In developing this method,

we established a framework for assessing gel recipes operating near this apparent limit of single-round

expansion. We found that the mechanical performance of gel recipes, i.e. resistance to deformation versus

gel expansion factor, varies smoothly with changes in crosslinker. For all high monomer (~3 M total

acrylamide and acrylate) gel recipe families, the relation between expansion factor and crosslinker

concentration fell close to a common curve. The high radical initiation rate of family B enabled gelation

without the inclusion of a crosslinker, suggesting that side reactions and polymer entanglement in these

conditions create sufficient effective crosslinks to form a gel. Gel deformation measurements versus

expansion factor, though less precise, also show high monomer recipe families falling close to a common

curve. Compared to the high monomer families, the original ExM recipe family is less resistant to

deformation for a given expansion factor and expands more for a given crosslinker concentration. The factor

determining gel properties is not crosslinker concentration in the gel recipe *per se*, but rather the density of

effective crosslinks formed between neighboring polymer chains in the gel. This suggests that the original,

monomer poor recipe less efficiently incorporates crosslinker molecules as effective polymer crosslinks.

This may be because the resulting lower rate of chain extension allows incorporated crosslinker molecules

to be re-reacted by the same polymer chain before they can react with neighboring polymers.

While the expansion factor of the original ExM recipe can be tuned by varying the crosslinker

concentration, it has been shown that increasing the monomer content is required to maintain nanoscale

isotropy, using centrioles as a convenient standard reference structure (Gambarotto *et al.*, 2019).

Considering gel quality versus expansion factor alone, the high monomer recipe family B derived from the

U-ExM recipe allows for a 10-fold expanding gel (at crosslinker concentration of 30 ppm) with a low

deformation index of 0.13. However, the high radical initiation rate used in this family (5 ppt APS and

TEMED) results in fast gelation. This increases the time sensitivity of mounting the specimen in the gelation

chamber and adds an additional challenge for adapting the method to thick tissues requiring extended
incubation in the gelation solution. Recipe families C and D solve this problem by reducing initiation rates,
at a slight expense of mechanical performance compared with family B. Like family B, family C has a high
acrylate content, which might contribute to imaging artifacts due to shrinkage of the sample prior to gelation
and inconsistent acrylate purity. Family D reduces the acrylate content by half while retaining similar
mechanical performance to family C, especially in the ten-fold expansion regime. Finally, with family E
we explored whether increasing the gelation temperature to 50ºC would produce the improved mechanical
performance of family B (through increased temperature-dependent radical initiation) without premature
gelation at room temperature. However, we found that this reduced the expansion factor and increased
susceptibility to experimental variation. Therefore, we based our TREx recipe on recipe family D. The
exact crosslinker concentration that produces 10-fold expanding gels was found to vary between labs (i.e.
50 ppm in Ashburn, VA, USA versus 90 ppm in Utrecht, The Netherlands, possibly due to differences in
gelation chamber design), so we recommend that each lab test a range of crosslinker concentrations between
30 and 100ppm using their choice of specimen, gelation chamber, and incubator.

We applied TREx to mouse brain tissue slices stained either for specific targets with antibodies, or for total
protein distribution with NHS ester dyes. Single round ten-fold expansion with TREx followed by total
protein staining was sufficient to reveal densely packed axons and dendrites running through the neuropil,
while individual organelles could be resolved within the neuronal soma. The nuclear envelope, along with
presumptive nuclear pore complexes, was also clearly resolved. The correct relative localizations of pre-
and post-synaptic markers and pre-synaptic neurotransmitter vesicles stained with standard
immunohistochemistry were also retained in TREx-expanded tissue slices.

We further demonstrated the utility of TREx for the study of cell biology through combinations
with several staining modalities in cultured cells, prepared in several culture formats. After a single round
of expansion with TREx, the commercially available membrane stain mCLING was able to clearly resolve
the internal structure of mitochondria and the detailed pattern of plasma membrane ruffling in activated
Jurkat T cells. While these structures would be readily resolved with electron microscopy, we were able to
do so in the context of complete cells, enabling concomitant detection and automated segmentation of
mitochondria clustered followed T cell activation. Caco-2 cells grown on permeable filters were also
successfully stained with mCLING and expanded with TREx to reveal the detailed structure of epithelial
microvilli and membrane interdigitations at the contacts of neighboring cells. These structures had
previously been known from electron microscopy but were now imaged with ease in the context of entire
cell monolayers. Successful application of TREx to filter-cultured Caco-2 cells further demonstrates the

robustness of TREx, because we had repeatedly failed to cleanly recover epithelial cultures using standard
ExM.

316   Combining mCLING with a total protein stain using NHS esters and an antibody marking the ER
in U2OS cells, shows close contacts between ER and other organelles. While the NHS ester and mCLING
staining patterns were similar in their overall contours, some clear differences in staining patterns were
noted, including the presence of presumptive nuclear pore complexes in the NHS ester channel. The strong
overlap between NHS ester and mCLING stains was not unexpected, given the reactivity of NHS esters
towards both unreacted lysines in the mCLING molecule and antibodies. However, the extent to which
NHS ester staining is truly unbiased over all proteins and how it may be modulated by the local environment
awaits further exploration.

324   In U2OS cells, TREx retained anti-tubulin antibody stain with high efficiency, maintaining
continuous microtubules with high signal-to-noise ratio after 1000-fold volumetric expansion. Combined
staining for ER and all membranes revealed close appositions along microtubules, ER and presumptive
mitochondria. In unexpanded cells, high quality antibody staining for microtubules requires monomeric
tubulin to be removed with a pre-extraction step, which destroys membranes and extracts other proteins.
For specimens expanded with TREx, this is not necessary as the monomeric tubulin signal is diluted 1000-
fold volumetrically, while the diameter of an expanded microtubule is still below the diffraction limit, so
the signal is only diluted by the the 10-fold linear expansion factor. This 1000-fold volumetric dilution is
also what enables the use of dense protein stains such as NHS ester dyes, which in unexpanded specimens
would be too dense to resolve any meaningful structure.

334   In summary, by systematically exploring the ExM recipe space we established a novel recipe using
standard ExM reagents that can be rapidly adopted by other labs. TREx allows for ten-fold expansion of
both thick tissue slices and cells in a single expansion step and has applications in tissue and high-resolution
subcellular imaging. Importantly, TREx of antibody-stained samples can be combined with off-the-shelf
small molecule stains for both membranes and total protein to provide ultrastructural context.

ACKNOWLEGDMENTS

We are grateful to Sven van IJzendoorn (UMCG) and Wilco Nijenhuis (UU) for providing the Caco2

monolayer samples. L.C.K. is supported by the European Research Council (ERC Consolidator Grant

819219). B.M. and P.W.T are supported by the Howard Hughes Medical Institute (HHMI).

REFERENCES

Arganda-Carreras, I. *et al.* (2017) 'Trainable Weka Segmentation: A machine learning tool for

microscopy pixel classification', *Bioinformatics*. doi: 10.1093/bioinformatics/btx180.

Chang, J.-B. *et al.* (2017) 'Iterative expansion microscopy', *Nature Methods*. Nature Research, 14(6), pp.

593–599. doi: 10.1038/nmeth.4261.

Chen, F., Tillberg, P. W. and Boyden, E. S. (2015) 'Expansion microscopy', *Science (New York, N.Y.)*,

347(6221), pp. 543–548. doi: 10.1126/science.1260088.

Drenckhahn, D. and Franz, H. (1986) 'Identification of actin-, α-actinin-, and vinculin-containing plaques

at the lateral membrane of epithelial cells', *Journal of Cell Biology*. doi: 10.1083/jcb.102.5.1843.

Gambarotto, D. *et al.* (2019) 'Imaging cellular ultrastructures using expansion microscopy (U-ExM)',

Nature Methods. Nature Publishing Group, 16(1), pp. 71–74. doi: 10.1038/s41592-018-0238-1.

Gao, M. *et al.* (2018) 'Expansion Stimulated Emission Depletion Microscopy (ExSTED)'. doi:

10.1021/acsnano.8b00776.

Gao, R. *et al.* (2019) 'Cortical column and whole-brain imaging with molecular contrast and nanoscale

resolution', *Science*. American Association for the Advancement of Science, 363(6424). doi:

10.1126/science.aau8302.

Gustafsson, M. G. L. (2000) 'Surpassing the lateral resolution limit by a factor of two using structured

illumination microscopy', *Journal of Microscopy*. John Wiley & Sons, Ltd, 198(2), pp. 82–87. doi:

10.1046/j.1365-2818.2000.00710.x.

Halpern, A. R. *et al.* (2017) 'Hybrid Structured Illumination Expansion Microscopy Reveals Microbial

Cytoskeleton Organization', *ACS Nano*, 11(12), pp. 12677–12686. doi: 10.1021/acsnano.7b07200.

Hell, S. W. and Wichmann, J. (1994) 'Breaking the diffraction resolution limit by stimulated emission:

stimulated-emission-depletion fluorescence microscopy', *Optics Letters*. The Optical Society, 19(11), p.

780. doi: 10.1364/ol.19.000780.

Hooikaas, P. J. *et al.* (2020) 'Kinesin-4 kif21b limits microtubule growth to allow rapid centrosome

polarization in t cells', *eLife*. eLife Sciences Publications Ltd, 9, pp. 1–75. doi: 10.7554/ELIFE.62876.

Jurriens, D. *et al.* (2020) 'Mapping the neuronal cytoskeleton using expansion microscopy', in *Methods in*
Cell Biology. doi: 10.1016/bs.mcb.2020.04.018.

Karagiannis, E. D. *et al.* (2019) 'Expansion Microscopy of Lipid Membranes', *bioRxiv*. doi:
10.1101/829903.

Ku, T. *et al.* (2016) 'Multiplexed and scalable super-resolution imaging of three-dimensional protein
localization in size-adjustable tissues', *Nature Biotechnology*, 34(9), pp. 973–981. doi: 10.1038/nbt.3641.

M'Saad, O. and Bewersdorf, J. (2020) 'Light microscopy of proteins in their ultrastructural context',
Nature Communications. doi: 10.1038/s41467-020-17523-8.

Mao, C. *et al.* (2020) 'Feature-rich covalent stains for super-resolution and cleared tissue fluorescence
microscopy', *Science Advances*. American Association for the Advancement of Science, 6(22), p.
eaba4542. doi: 10.1126/sciadv.aba4542.

Okay, O. (2009) 'General Properties of Hydrogels', in Gerlach G., A. K. (ed.) *Hydrogel Sensors and*
Actuators. Springer Series on Chemical Sensors and Biosensors (Methods and Applications). vol. 6.
Springer, Berlin, Heidelberg, pp. 1–14. doi: 10.1007/978-3-540-75645-3_1.

Revelo, N. H. and Rizzoli, S. O. (2016) 'The membrane marker mCLING reveals the molecular
composition of trafficking organelles', *Current Protocols in Neuroscience*. doi:
10.1002/0471142301.ns0225s74.

Rust, M. J., Bates, M. and Zhuang, X. W. (2006) 'Sub-diffraction-limit imaging by stochastic optical
reconstruction microscopy (STORM)', *Nat Methods*, 3(10), pp. 793–795. doi: Doi 10.1038/Nmeth929.

Tas, R. P. *et al.* (2017) 'Differentiation between Oppositely Oriented Microtubules Controls Polarized
Neuronal Transport', *Neuron*. doi: 10.1016/j.neuron.2017.11.018.

Tillberg, P. W. *et al.* (2016) 'Protein-retention expansion microscopy of cells and tissues labeled using
standard fluorescent proteins and antibodies', *Nature Biotechnology*. Nature Publishing Group, 34(9), pp.
987–992. doi: 10.1038/nbt.3625.

Truckenbrodt, S. *et al.* (2018) 'X10 expansion microscopy enables 25-nm resolution on conventional
microscopes.', *EMBO reports*. EMBO Press, 19(9), p. e45836. doi: 10.15252/embr.201845836.

Truckenbrodt, S. *et al.* (2019) 'A practical guide to optimization in X10 expansion microscopy', *Nature*
Protocols. doi: 10.1038/s41596-018-0117-3.

Xu, H. *et al.* (2019) 'Molecular organization of mammalian meiotic chromosome axis revealed by

expansion STORM microscopy', *Proceedings of the National Academy of Sciences of the United States of*

America. National Academy of Sciences, 116(37), pp. 18423–18428. doi: 10.1073/pnas.1902440116.

Yu, C.-C. (Jay) *et al.* (2020) 'Expansion microscopy of C. elegans', *eLife*, 9. doi: 10.7554/eLife.46249.

Zhang, Y. S. *et al.* (2016) 'Hybrid Microscopy: Enabling Inexpensive High-Performance Imaging

through Combined Physical and Optical Magnifications', *Scientific Reports*. Nature Publishing Group,

6(1), p. 22691. doi: 10.1038/srep22691.

Zwettler, F. U. *et al.* (2020) 'Molecular resolution imaging by post-labeling expansion single-molecule

localization microscopy (Ex-SMLM)', *bioRxiv*. Cold Spring Harbor Laboratory, p. 2020.03.12.988923.

doi: 10.1101/2020.03.12.988923.

MATERIAL AND METHODS

Gelation chambers for recipe space exploration

A glass slide served as the bottom piece of each gelation chamber. Four strips of 250 μm-thick adhesive silicone material (Digikey cat# L37-3F-320-320-0.25-1A), ~3 mm wide and running the width of the slide, were adhered to the slide to partition it into three separate chambers, each ~12 mm wide. A plus-charged glass slide was placed over the silicone strips to form the top of the gelation chamber and held in place with tape. Two sides of each chamber were open to air, providing a convenient fill port for adding ~100 uL of monomer solution after chamber construction.

Gel synthesis and characterization for recipe space exploration

Sodium acrylate was made by neutralizing acrylic acid (Sigma, 147230) by adding NaOH until the pH reached the range of 7.5-8. Initial neutralization (until pH ~7) was done with 10 N NaOH on ice and using a fume hood to avoid acrylic acid vapor. Neutralization was done in a volume of water calculated to yield a final concentration of 4 M sodium acrylate. The gel recipes for each family contained 1xPBS and the amounts of acrylamide (Sigma, A4058), sodium acrylate, and initiator (APS, Sigma, A3678) indicated in Fig. 1A. Each gel recipe contained the same amount of TEMED (Sigma, T7024) as APS. For each recipe family, gelation solution with crosslinker withheld (but including APS and TEMED) was premixed on ice, in one tube for each recipe family. This solution was then split into six tubes and mixed with serial dilutions of crosslinker (bisacrylamide, Sigma, M1533) to arrive at final crosslinker concentrations of 1 ppt, 300, 100, 30, and 10 ppm, and 0. Complete gelation solution thus obtained was pipetted into gelation chambers and incubated at 50 ºC for 1 hour or 37 ºC for 2 hours. Gels were then cooled for 15 minutes at room temperature and chamber tops carefully removed. Gels typically remained stuck exclusively to the top (plus charged) slide. Specimens of each gel were taken with a 6 mm biopsy punch, taking care to avoid material within ~2 mm of the chamber edges (due to oxygen exposure during gelation). Excess gel was scrapped away with a razor blade. A few drops of distilled water were pipetted onto each gel to help release them from the glass slide. Each 6 mm gel specimen was gently released from the slide with a razor blade, placed in a 9 cm Petri Dish and expanded by washing with excess water 2x 15 minutes followed by 2x 1 hour. Diameters of expanded gels were measured and divided by 6 mm to obtain the expansion factor. A semi-circle 25 mm in diameter was punched from each gel using a cookie cutter. Semi-circular gel punches were placed in a plastic tray, which was stood up on end so that the gel stood upright on its curved side. For stiff gels, the top edge remained straight and level while for weak gels the top edge deformed under the force of gravity to form a curve. Each gel was photographed, with a ruler positioned for scale. Using image J, each

top edge was described by seven manually chosen points, which were then fit to a circle. This best-fit circle

was used to calculate the vertical deviation of the gel corners, which was divided by the gel radius to obtain

the deformation index.

TREx gelation solution

Sodium acrylate was either purchased (Sigma, 408220) or made by neutralizing acrylic acid by adding

NaOH until the pH reached the range of 7.5-8. Initial neutralization (until pH ~7) was done with 10N NaOH

on ice and using a fume hood to avoid acrylic acid vapor. TREx gelation solution contains 1.1 M sodium

acrylate, 2.0 M acrylamide (AA), 50 ppm (for tissue slices prepared at Janelia) or 90ppm (for cultured cells

prepared at University of Utrecht) N,N'-methylenebisacrylamide (bis), PBS (1x), 1.5 ppt APS, 1.5 ppt

TEMED, and (optionally, for thick tissue slices) 15 ppm 4HT. Monomer solution was made by combining

all components of gelation solution except APS, TEMED, and 4HT. Monomer solution may be aliquoted

and stored at -20°C, but must be thawed at room temperature and thoroughly vortexed before use to

redissolve any acrylamide crystals that may have precipitated at low temperature before freezing. 4HT,

TEMED, and APS were added to monomer solution to produce gelation solution directly before use.

Tissue experiments

Mice were transcardially perfused with ice cold 4% formaldehyde in 100 mM sodium phosphate buffer, pH

7.4. Brains were dissected out and post-fixed in 4% formaldehyde overnight at 4°C, followed by washing

with PBS (1x) and slicing at 100 μm by vibratome. Slices were optionally stained with standard IHC

procedures. Primary antibodies (Chicken anti Bassoon Synaptic systems cat#141016, RRID:AB_2661779;

Rabbit anti Homer abcam cat#97593, RRID:AB_10681160; Ms IgG3 VGAT SySy cat#131011,

RRID:AB_887872) were used at 1:300 dilution in PBS with 0.1% Triton and 2% BSA (PBT) overnight at

4°C. Sections were washed 3x 30min in PBT and given at least 6h in secondary antibodies 1:500 in PBT at

room temperature (Goat Anti-Rabbit alexa 488 abcam cat#150077, RRID:AB_2630356; Goat anti-Mouse

IgG3 alexa 594 invitrogen cat# A-21155, RRID: AB_2535785; Goat anti chicken CF633 biotium

cat#20126, RRID: AB_10852831). After 3x 30min washes in PBS, slices were treated with 10 μg/mL AcX

in PBS for 1 hour at room temperature, followed by rinsing with PBS. Slices were incubated with TREx

gelation solution (using 50ppm bis, and with 4HT added up to 15ppm), for 20min on ice, with shaking.

Following incubation on ice, each tissue specimen was placed on a glass slide at room temperature. Four

dabs of vacuum grease were applied to the slide, with each dab at least several mm from the tissue specimen.

A coverslip was placed over the tissue and vacuum grease dabs, and pressed down until contacting the

tissue, taking care not to let the tissue slide around on the slide. The vacuum grease served to hold the

assembly in place, thus forming the gelation chamber. Gelation solution was pipetted into the chamber from

the side to fully surround the tissue. The chamber was incubated at 37 °C for 1 hour to complete gelation.

Following embedding, excess gel was removed with a razor blade, and gelled slices were recovered into

disruption buffer (5% SDS, 200 mM NaCl, 50 mM Tris pH 7.5), using a razor blade to gently release them

from the glass slide. Gels were incubated at 80 °C for 3 hours, followed by washing in water once and PBS

several times to remove residual SDS. Slices were optionally stained with BODIPY-FL NHS at 10-20 μM

(and optionally DAPI) in PBS for 1 hour at room temperature. Disrupted slices were placed in glass bottom

6-well plates and washed in milliQ water 3x 15 minutes followed by 2x 1 hour to fully expand. Gels with

total protein stain were imaged by confocal microscopy using a Zeiss LSM 800 with 40x/1.2NA, water

immersion objective. Gels with synaptic antibody stains were imaged on a Zeiss Z1 lightsheet microscope

with 10x/0.3NA illumination objectives and 20x/1.0NA water immersion detection objective. Scale bar

represent the expanded size divided by the approximate expansion factor of 10.

Cell culture and T cell activation

Jurkat T cells (clone E6.1) were grown in RPMI 1640 medium w/ L-Glutamine (Lonza) supplemented with

9% Fetal Bovine Serum and 1% penicillin/streptomycin. For T cell activation, 18 mm #1.5 coverslips

(Marienfeld, 107032) were coated with Poly-D-Lysine (Thermo Fisher Scientific, A3890401), washed with

phosphate buffered saline (PBS) and incubated overnight at 4 °C with a mouse monoclonal anti-CD3 (clone

UCHT1, StemCell Technologies, #60011) 10 μg/mL in PBS. Cells were spun down for 4 minutes at 1000

rpm and resuspended in fresh, prewarmed RPMI 1640 medium, after which cells were incubated on the

coated coverslips for 3 minutes prior to fixation. U2OS cells were cultured in DMEM medium

supplemented with 9% Fetal Bovine Serum and 1% penicillin/streptomycin. U2OS cells were transfected

with GFP-Sec61β (Addgene, 15108) using FuGENE6 (Promega). Caco2-BBE cells (a gift from S.C.D. van

IJzendoorn, University Medical Center Groningen, The Netherlands) were maintained in DMEM

supplemented with 9% FBS, 50 μg/μl penicillin/streptomycin and 2 mM L-glutamine. For imaging, cells

were seeded on 6.5 mm Transwell filters (3470; Corning) at a density of 1 x 10^5/cm^2 and cultured for 10-

12 days to allow for spontaneous polarization and brush border formation.

Immunofluorescence, mCLING treatment and antibodies

For all experiments, cells were fixed for 10 minutes with pre-warmed (37 °C) 4% paraformaldehyde + 0.1%

glutaraldehyde in PBS. For visualization of lipid membranes, cells were washed twice in PBS after fixation

and incubated in 5 μM either mCLING-Atto647N (Synaptic Systems, 710 006AT1) or mCLING-Atto488

(Fig. 3C, Synaptic Systems, 710 006AT3) in PBS overnight at RT. The following day, cells were fixed a

second time with pre-warmed (37 °C) 4% paraformaldehyde + 0.1% glutaraldehyde in PBS. Next, cells

were washed with PBS and permeabilized using PBS + 0.2% Triton X-100. Epitope blocking and antibody

labeling steps were performed in PBS + 3% BSA. For immunofluorescence staining, we used a rabbit

monoclonal antibody against α-tubulin (clone EP1332Y, Abcam, ab52866) and a chicken polyclonal

antibody against GFP (Aves Labs, GFP-1010) in combination with goat anti-rabbit IgG (H+L) Alexa Fluor

594 (Molecular Probes, a11037) and goat anti-chicken IgY (H+L) Alexa Fluor 488 (Molecular Probes,

a11039), respectively.

TREx protocol and NHS staining

For TREx Microscopy, samples were treated with 0.1 mg/mL acryloyl X-SE (AcX) (Thermo Fisher,

A20770) in PBS overnight at RT. TEMED and APS were added to monomer solution (1.5 ppt each) to

produce gelation solution. 170 µL of gelation solution was transferred to a silicone mold with inner diameter

of 13 mm (Sigma-Aldrich, GBL664107) attached to a parafilm-covered glass slide, with the sample put

cell-down on top to close off the gelation chamber. The sample was directly transferred to a 37 °C incubator

for 1 hour to fully polymerize the gel. All gels excluding samples that were processed for subsequent NHS

ester staining were transferred to a 12-well plate and digested with 7.5 U/mL Proteinase-K (Thermo Fisher,

EO0491) in TAE buffer (containing 40 mM Tris, 20 mM acetic acid and 1 mM EDTA) supplemented with

0.5% Triton X-100, 0.8 M guanidine-HCl, and DAPI for 4 hours at 37 °C. The gel was transferred to a Petri

dish, water was exchanged 2x 30 minutes and the sample was left in milliQ water to expand overnight.

For NHS-staining, gels were first treated in disruption buffer containing 200 mM SDS, 200 mM NaCl and

50 mM Tris pH 6.8 for 1.5 hours at 78°C. Gels were washed twice for 15 minutes in PBS and incubated

with 20 µg/mL Atto 594 NHS ester (Sigma-Aldrich, 08471) in PBS prepared from a 20 mg/mL stock

solution in DMSO for 1 hour at RT with shaking. After staining, gels were washed with excess of PBS,

transferred to a Petri dish, and expanded overnight. Prior to imaging the cells were trimmed using a scalpel

blade to fit in a Attofluor Cell Chamber (Molecular probes A-7816).

Image acquisition and analysis

All cultured cell ExM images were acquired using a Leica TCS SP8 STED 3X microscope equipped with

a HC PL APO 86x/1.20W motCORR STED (Leica 15506333) water objective. A pulsed white laser (80

MHz) was used for excitation. The internal Leica GaAsP HyD hybrid detectors were used with a time gate

of $1 \leq tg \leq 6$ ns. The set-up was controlled using LAS X and run without stimulated emission depletion

(STED).

All data processing and analysis was done using ImageJ and Arivis.

For Fig. 2A, raw data was drift corrected using Huygens Professional (SVI) and imported into ImageJ
where a sum-projection of 2 planes (z-spacing: 0.8 µm) was made. Fig. 2B is a maximum projection of 2
planes (z-spacing: 0.38 µm) and indicated zoom is a volumetric render of the raw data in Arivis. For Fig.
3A, raw data was imported in Arivis, a Discrete Gaussian Filter with smoothing radius of 2 was applied
and this dataset was used for volumetric renders and clipping. Gamma was adjusted manually to increase
visibility of plasma membrane ruffles and intracellular organelles in the same view. For Fig. 3B, the same
raw dataset was imported in ImageJ and a sum-projection of 3 planes (z-spacing: 0.35 µm) around the plane
of the immunological synapse was segmented for mitochondria using the trainable Weka segmentation
plugin in ImageJ. Fig. 3C is a sum-projection of 3 planes (z-spacing 0.35 µm). The linescan in Fig. 3C
was generated using ImageJ and processed using Graphpad Prism 8.

For Fig. 3D, raw data was imported in Arivis, a Discrete Gaussian Filter with smoothing radius of 2 was
applied and this dataset was volumetrically rendered with the opacity mapped to the z-axis. For Fig. 3E,
raw data was imported in ImageJ and a running sum projection of 5 slices (z-spacing 0.35 µm) was
generated using the RunningZProjector plugin and gamma was adjusted manually to increase intracellular
contrast. The resulting image was imported into Arivis for volumetric rendering. Fig. 3F and G are
maximum projections of 3 planes (z-spacing: 0.35 µm) and respective zooms.

Fig 4A is a maximum projection of 3 planes (z-spacing: 0.35 µm). For Fig 4B, raw data was imported into
Arivis, a Discrete Gaussian Filter with smoothing radius of 2 was applied and this dataset was used for
volumetric renders and clipping. Fig 4C is a sum projection of 3 slices (z-spacing 0.35 µm) of the same raw
data as Fig. 4B and was processed using ImageJ.

FIGURE 1: Development of TREx gel recipe

A) Parameters of gel recipe families explored, including component concentrations and gelation temperature. Each family was characterized by keeping these conditions constant while systematically varying the crosslinker concentration.

B) Expansion factor (mean ±s.d., n=3) versus crosslinker concentration (log scale) for each gel recipe family without biological specimens. Line colors correspond to recipe families as in 1A. Specific recipes (with specific crosslinker concentrations) are indicated as follows: filled purple dot - original ExM recipe, blue dot - UltraExM, and yellow dot - TREx. All recipe families were tested with crosslinker concentrations of 0, 10, 30, 100, 300, 1000 ppm, plus an addition condition with 1500 ppm for family A corresponding to the original ExM recipe. Only conditions in which gels formed are plotted.

C) Definition of gel deformation index. Example gels from recipe family A with high crosslinker and low

deformation (top panel, 1.5 ppt), and low crosslinker and high deformation (middle panel, 300 ppm).

Bottom panel, schematic illustrating deformation index measurement.

D) Deformation index (mean ±s.d., n=3) versus expansion factor for each gel recipe family without

biological specimens, with line colors and dots corresponding to specific recipes as in 1A and 1B.

Horizontal grey lines indicate thresholds for gels with mechanical quality deemed perfect (deformation <

0.125) and acceptable (deformation < 0.25). Ideal recipes would occupy the lower right quadrant,

corresponding to high expansion and low deformability.

BDP-FL-NHS

Homer/Bassoon/VGAT/DAPI

FIGURE 2: TREx in mouse brain tissue slices

A) Mouse brain tissue (cortex) expanded using TREx, stained for total protein content with BODPIY-FL

NHS, and imaged by confocal microscopy. Displayed contrast is inverted to show dense stained regions as

dark. Inset, zoom-in showing nuclear envelope with densely stained structures spanning the nuclear

envelope, consistent with nuclear pore complexes.

B) Mouse brain tissue (cortex) stained with antibodies against (purple) homer, (yellow) bassoon, and (blue)

VGAT, and DAPI (grey), and expanded using TREx. Inset, zoom-in showing paired bassoon and homer-

rich structures, consistent with excitatory synapses and complex structured pre-synaptic vesicle pool

marked by VGAT bearing several release sites marked by bassoon.

Scale bars (corrected to indicate pre-expansion dimensions): main ~2 µm, zooms ~400 nm

FIGURE 3: TREx can be used to visualize the ultrastructure of cellular membranes

A) Volumetric render of Jurkat T cell activated on anti-CD3 coated coverslip fixed and stained using mCLING. Colored clipping planes indicate portion clipped out to reveal intracellular detail.

B) Immunological synapse of activated T cell stained with mCLING in (A) revealing organelle clustering at the immunological synapse. Below: mitochondria segmented using the trainable Weka segmentation algorithm indicated in magenta.

C) Representative example of mitochondrion in T cells visualized with mCLING. Line profile along the orange dashed line indicates mitochondrial cristae.

D) Volumetric projection of Caco2 monolayer apical brush border as seen from above looking down on the cells.

E) Volumetric side projection of a portion of the cell shown in D as seen from the side.

F) Brush border x, y plane of the cell shown in D, with zooms revealing hollow microvilli.

G) Similar to F, representative plane below the apical surface revealing highly interdigitated cell-cell contacts.

Scale bars (corrected to indicate pre-expansion dimensions): A, B, D, E ~2 μm, C ~500 nm, F and G main ~2 μm and zooms ~500 nm

FIGURE 4: TREx Microscopy can combine antibody-based staining with NHS ester total protein

stain to provide cellular context

A) Merged plane of expanded U2OS cell expressing GFP-Sec61β stained for mCLING, GFP, NHS ester
and DAPI. Single planes of mCLING and NHS ester are displayed in inverted contrast.

B) Volumetric render of U2OS cell expressing GFP-Sec61β Sec61β stained for mCLING, GFP and tubulin.
Top portion of cell is clipped with clipping plane indicated in red. Volumetric render of entire volume for
GFP and tubulin in insert A and B, respectively.

C) Single planes displayed in inverted contrast of the top of cell in B revealing the tight spatial organization.

Scale bars (corrected to indicate pre-expansion dimensions): A, C ~1 μm, B ~2 μm

SUPPLEMENTAL MOVIES

Supplemental Movie 1: Z-stack of Fig. 2A

Supplemental Movie 2: 3D render of Fig. 2B

Supplemental Movie 3: 3D render of Fig. 3A

Supplemental Movie 4: 3D render of Fig. 3D

Supplemental Movie 5: 3D render of Fig. 4B