Enzyme kinetics of CRISPR molecular diagnostics

Ashwin Ramachandran and Juan G. Santiago
Department of Mechanical Engineering, Stanford University, California, USA 94305.
Email: ashwinrc@stanford.edu, juan.santiago@stanford.edu

Abstract

CRISPR diagnostic assays have gained significant interest in the last few years. This interest has grown rapidly during the current COVID-19 pandemic where CRISPR diagnostics have been frontline contenders for rapid testing solutions. This surge in CRISPR diagnostics research prompts the following question: What exactly are the achievable limits of detection and associated assay times enabled by the kinetics of Cas12 and Cas13 enzymes? To address this question, we here present a model based on Michaelis-Menten enzyme kinetics theory applied to Cas enzymes. We use the model to develop analytical solutions for reaction kinetics and develop back-of-the-envelope criteria to validate and check for consistency in reported enzyme kinetics parameters. Our analysis shows there is not a single published study which presents reliable and consistent values of kinetic parameters for Michaelis-Menten-type models of CRISPR-associated enzymes Cas12 and Cas13. We further use our model to explore ranges of reaction time scales and degree of reaction completion for practically relevant target concentrations applicable to CRISPR-diagnostic assays.

1. Introduction

There have been many interesting advancements in the field of CRISPR diagnostic assays in the last few years. This is particularly true during the current COVID-19 pandemic where CRISPR diagnostics have been frontline contenders for rapid testing solutions[1–4]. CRISPR diagnostic assays use subtypes of Cas12 or Cas13 enzymes for DNA or RNA detection, respectively[5–8]. The emerging and strong interest in CRISPR diagnostics research prompts the following question: What exactly are the achievable limits of detection (LODs) and associated assay times enabled by the kinetics of Cas12 and Cas13 enzymes (**Figure 1**)? As expected, the answer to this important question depends strongly on the CRISPR-associated enzyme kinetics as these establish and fundamentally limit the sensitivity of these diagnostics.

For ssDNA and dsDNA detection using CRISPR-Cas12, LbCas12a has been the most widely used Cas enzyme.[2,4,7,9–11] The study of Chen et al.[11] was the first to characterize and report the kinetic rates of LbCas12a. The latter study reported an enzyme turnover rate (k_{cat}) and Michalis-Menten constant (K_M) of 250 per second and 4.94×10^{-7} M, respectively, with a catalytic efficiency (k_{cat}/K_M) of 5.1×10^8 $M^{-1}s^{-1}$ for LbCas12a activated by target ssDNA. For a dsDNA activator, the study reported a k_{cat} and K_M of 1250 per second and 7.25×10^{-7} M, respectively, and a k_{cat}/K_M of 1.7×10^9 $M^{-1}s^{-1}$. The latter value reportedly approached the limit of diffusion.[11] For ssRNA detection using CRISPR-Cas13, the study of Slaymaker et al.[12] was the first to report enzyme turnover rates among all the subtypes and orthologs of Cas13. They reported a k_{cat} of 987 turnovers per second for PbuCas13b. Subsequently, Shan et al.[13] reported a k_{cat} of 4854 per second and K_M of 4.45×10^{-6} M using a 5 nt (of rU) ssRNA fluorescent reporter in an application of LbuCas13a for MicroRNA (miRNA) detection. Later, Zhou et al.[14] showed for LbuCas13a that a shorter reporter molecule which was composed of 2 nt (of rU) lowered the k_{cat} to 204 per second

and K_M of 2.62 x 10^{-6} M. Most recently, Fozouni et al.[1] presented a study of SARS-CoV-2 viral RNA detection using LbuCas13a and a set of 3 gRNAs, and used a model to fit their experimental data which used parameters of k_{cat} = 600 per second and K_M between 1 x 10^{-6} M and 3 x 10^{-6} M.

In this work, we provide back-of-the-envelope analyses and scaling arguments based on Michaelis-Menten-type enzyme kinetics theory applied to CRISPR-diagnostics. We present a set of simple rules based on fundamental principles of species conservation, maximum reaction velocities, and reaction time scales, and these rules can be used as a check on the reported parameters of enzyme kinetics. We apply these rules and perform calculations on several published studies. Surprisingly, we find several studies violate these simple validations. Lastly, we develop and present results from analytical and numerical models for predicting Cas enzyme kinetics and use these to discuss the effect of kinetic parameters on the assay time scales and limits of sensitivity of CRISPR-diagnostic assays.

2. Theory

In this section, we present an enzyme kinetics model for the trans-cleavage activity of CRISPR-associated enzymes. The mechanisms of relevant reactions are summarized in **Figure 1**. This model invokes Michaelis-Menten enzyme kinetics theory[15] and is used to develop closed-form approximations for product formation versus time. We also present simple back-of-the-envelope rules based on kinetics theory and fundamental principles of species conservations that can be used to check for accuracy of reported enzyme kinetics data.

2.1 Michaelis-Menten model for *trans*-cleavage activity

Upon recognition of the target nucleic acid, the CRISPR-enzymes are activated and exhibit non-specific *trans*-cleavage of single-stranded nucleic acids (c.f. **Fig. 1**). [9,11,16] This *trans*-cleavage activity is a multi-turnover enzymatic catalyzed by the target activated CRISPR enzyme and is the key process underlying nucleic acid detection assays. Cas12 can be activated by target ssDNA and dsDNA, while Cas13 is activated by ssRNA. Upon activation, Cas12 and Cas13 indiscriminately cleave ssDNA and ssRNA molecules, respectively. CRISPR assays typically use fluorescent nucleic acid molecules (with a quencher) as reporters, which fluoresce only when they are cleaved by the CRISPR enzyme in the presence of target.

The *trans*-cleavage CRISPR enzymatic reaction can be modeled, in general, as

$$E + S \overset{k_f, k_r}{\longleftrightarrow} C \overset{k_{cat}}{\longrightarrow} E + P \tag{1}$$

where, E is target-activated Cas12/13-gRNA complex, S is the uncleaved reporter, C is the reaction intermediate (enzyme-substrate reporter complex), P is the cleaved reporter molecules, k_f and k_r are the forward and reverse rate constant for the reaction between E and S, and k_{cat} is the catalytic turnover rate of the enzyme. The reaction rate laws for the individual species are as follows:

$$\frac{d[E]}{dt} = -k_f[E][S] + k_r[C] + k_{cat}[C] \tag{2}$$

$$\frac{d[S]}{dt} = -k_f[E][S] + k_r[C] \tag{3}$$

$$\frac{d[C]}{dt} = k_f[E][S] - k_r[C] - k_{cat}[C] \qquad (4)$$

$$\frac{d[P]}{dt} = k_{cat}[C]. \qquad (5)$$

These are subject to initial conditions $[E](0) = E_0, [S](0) = S_0, [C](0) = 0,$ and $[P](0) = 0.$ Next, we use the quasi-steady state assumption (see Briggs-Haldane approximation[15]) which models the concentration of the intermediate enzyme-substrate complex as nearly constant in time. Thus, we can write

$$\frac{d[C]}{dt} \approx 0 \rightarrow [C] = \frac{k_f[E][S]}{k_r + k_{cat}}. \qquad (6)$$

Further, species conservation yields

$$[E](t) + [C](t) = E_0, \text{ and } [C](t) + [S](t) + [P](t) = S_0, \qquad (7)$$

where E_0 and S_0 are respectively the initial amount of target-activated enzyme and substrate (reporter). We note that E_0 is at most equal to the amount of target nucleic acid present in the sample (usually the Cas-gRNA complex is present in abundance). However, at very low concentrations of target, target recognition and activation of the CRISPR enzyme (*cis*-cleavage) is limited by the diffusion of the target molecules to the active site of the enzyme.[17] For simplicity, we will here assume that E_0 is given by the concentration of the target nucleic acid.

Under these assumptions, the rate of product formation (reporter cleavage) is governed by the Michaelis-Menten equation

$$v = \frac{d[P]}{dt} = k_{cat}E_0 \frac{[S]}{K_M + [S]}. \qquad (8)$$

In traditional Michaelis-Menten kinetics analysis, $E_0 \ll S_0$ and the kinetics are studied over short time scales (relative to the overall product formation time-scale). Hence, in this limit, very little product is formed $P \ll S_0$ and we have $[S] \approx S_0$. Thus, the product is formed at a constant rate (e.g., linear fluorescence increase vs time), and the reaction velocity is given by Equation (8) with $[S] \approx S_0$.

In most practical applications of CRISPR-diagnostics, the concentration of the substrate is much smaller than the Michaelis-Menten constant of the enzyme, $[S] \ll K_M$. Typically, $S_0 \sim \mathcal{O}(100 \text{ nM})$ and $K_M \sim \mathcal{O}(1 \text{ μM})$. Thus, we can use this approximation to further derive an expression for the evolution of the product (reporter cleavage) versus time, over much longer time scales. Specifically, we can rewrite Equation (8) as

$$\frac{d[P]}{dt} = \frac{k_{cat}E_0}{K_M}(S_0 - [P]). \qquad (9)$$

Equation (9) can be rearranged into a linear first-order differential equation[18] as

$$\frac{d[P]}{dt} + \frac{k_{cat}E_0}{K_M}[P] = S_0. \qquad (10)$$

Equation (10) can be solved analytically for the concentration of product formed (reporters cleaves) versus time as

$$\frac{[P](t)}{S_0} \approx \left(1 - \exp\left(-\frac{t}{\tau}\right)\right) \qquad (11)$$

where $\tau = K_M/k_{cat}E_0$. Here τ is a time scale that governs the characteristic time to approximately complete the reaction, and $[P](t)/S_0$ is simply the fraction of cleaved reporters relative to the total initial uncleaved reporters. Physically, we see that the time scale of significant product formation

τ decreases with: (i) higher turnover rate k_{cat}, (ii) higher target concentration E_0, (iii) lower K_M (which represents a higher affinity of the activated enzyme to the substrate). Further, we also see that the ratio of k_{cat} to K_M is very important for the product formation rate and this ratio arises naturally from the current analysis.

Figure 1. Schematic for RNA and DNA detection using CRISPR-Cas13 and Cas12, respectively. The Cas13:gRNA complex is activated by a target ssRNA (*cis*-cleavage; single turnover), after which the activated enzyme cleaves ssRNA molecules including ssRNA reporters indiscriminately (*trans*-cleavage; multi turnover). Cas12:gRNA can be activated by both target ssDNA and dsDNA (*cis*-cleavage; here, shown only for dsDNA). After activation, the Cas12 enzyme cleaves ssDNA molecules including ssDNA reporter molecules indiscriminately (*trans*-cleavage activity; multi turnover). The *trans*-cleavage activity for Cas12 and Cas13 each follows Michaelis-Menten kinetics, where the target-activated Cas-gRNA complex is the enzyme (*E*), the fluorescent reporter molecule is the substrate (*S*), and the cleaved reporter molecule is the product (*P*).

2.2 Back-of-the-envelope checks on reported kinetics data

We here present three simple back-of-the-envelope hand calculations that can be used to verify the accuracy of reported kinetics data from experiments and/or simulations. The first rule derived here originates simply from species conservation: The amount of product formed during the initial linear portion t_{lin} of the kinetics measurements (c.f. **Figure 2**; early time scales where $\tau < 1$) cannot exceed the initial amount of substrate (before the start of the reaction). t_{lin} is a quantity that typically is easily apparent from experimental data (e.g. the duration of the initial, nearly linear increase of fluorescence versus time data). We can define the non-dimensional parameter α as the ratio of the total product formed at t_{lin} and the initial substrate concentration S_0. For most CRISPR diagnostics assays, α is the ratio of the amount of cleaved reporters at time t_{lin} to the initial concentration of the uncleaved reporters S_0. Simple conservation of species then limits this value to unity as in

$$\alpha = \frac{v t_{lin}}{S_0} < 1. \tag{12}$$

Here v is the reported reaction velocity (e.g., in nM/s).

The second rule is derived from the theoretical maximum reaction velocity possible as given by Michaelis-Menten kinetics. From Equation (8), the maximum rate of product formation occurs when the substrate concentration is far in excess compared to K_M, $[S] \gg K_M$, and the theoretical maximum velocity v_{max} is equal to $k_{cat} E_0$. One can therefore define a non-dimensional quantity, β, as the reported velocity v normalized by v_{max} as follows:

$$\beta = \frac{v}{v_{max}} = \frac{v}{k_{cat} E_0} \leq 1. \tag{13}$$

Third, we observe that the time scale of linear portion of the kinetics data t_{lin} should be at most on the same order as the reaction time scale τ. For this, we define a non-dimensional quantity, γ, defined as the ratio of t_{lin} and τ, and the third rule is given by

$$\gamma = \frac{t_{lin}}{\tau} = \frac{t_{lin} k_{cat} E_0}{K_M} \leq \mathcal{O}(1). \tag{14}$$

Lastly, we note that the first rule (condition on α) does not require *a priori* estimates of the reaction parameters k_{cat} and K_M. The first rule is therefore a good check on ensuring consistent fluorescence versus concentration calibration for experiments. For example, inconsistencies in calibration or conversion from fluorescence to concentration will result in unphysical values of v, and these artifacts can lead to an apparent violation of the first rule (we here assume the initial substrate concentrations and times are properly reported). The second and third rules rely on experimentally/numerically derived reaction parameters k_{cat} and K_M (e.g., using a curve fit to initial reaction velocity versus substrate concentration kinetics data). The latter two rules (involving β and γ) provide useful validations of self-consistency in kinetics data and the reported values of k_{cat} and K_M. Briefly, all data should obey Rules 1 and 2, and reactions for which $[S] \ll K_M$ (typically, less by at least about a factor of 5-fold) must obey Rule 3.

3. Results and Discussion

3.1 Back-of-the-envelope checks on published data

Presented in **Table 1** are the back-of-the-envelope checks for quantities α, β, and γ (c.f. Section 2.2) performed on published studies which present raw Michaelis-Menten-type

characterization experiments and kinetics data for different Cas12 and Cas13 enzymes. To the best of our knowledge, we have included all published studies to date which present any Michaelis-Menten kinetics data for CRISPR-Cas12/13. We found that all of the studies listed in the first column of **Table 1** violated at least one of the three checks involving α, β, and γ (see fifth, sixth, and seventh columns, respectively). For example, the studies of Chen et al.[11] for LbCas12a and Zhou et al.[14] for LbuCas13a both had values of α and γ significantly greater than 1. The study of Shan et al.[13] for LbuCas13a reported values of α, β, and γ which were each greater than 1, and so violated all the three rules. Slaymaker et al.[12] studied PbuCas13b, and the study violated at least two (α and β conditions) out of the three checks. These results in **Table 1** suggest that the kinetics parameters (second, third, and fourth columns of **Table 1**) published in the aforementioned studies (first column of **Table 1**) are not reliable and could be grossly incorrect. Also listed in **Table 1** are other relevant publications[1,10,19–22] (column 10) which cite the aforementioned studies and either directly rely (e.g. in making estimates or modeling experiments) on the published kinetics data or provide new experimental data which seems to corroborate reported kinetics parameter values which violated our checks.

Lastly, we present updated kinetics parameter values (columns 11, 12 and 13 of **Table 1**) which we recently received from the corresponding authors of Refs.[11,12] through personal communication[23,24]. Unlike the originally published data, the updated kinetic rates are consistent with (i.e. pass) our three back-of-the-envelope checks (see columns 14, 15, and 16 of **Table 1**). That is, for these updated values, the three quantities α, β, and γ are each less than unity as expected. If these updated kinetic parameters are correct, a major consequence is that the kinetics of LbCas12a is not catalytically efficient or diffusion-limited ($k_{cat}/K_M \sim 10^6$ to 10^7 $M^{-1}s^{-1}$), contrary to the understanding so far (e.g. see Refs. [4,10,11,19,21]). Also, turnover rates for LbCas12a and PbuCas13b, respectively, which are 2 to 3 orders of magnitude lower would imply that proportionally longer reaction time scales are required for *trans*-cleavage activities (Equation (11)).

3.2 Predictions of analytical and numerical enzyme kinetics model

We here use the model developed in Section 2.1 to explore the kinetics of Cas enzymes. **Figure 2** shows the fraction of reporters cleaved by the target-activated enzyme divided by the initial amount of uncleaved reporters present in the reaction versus time normalized by the reaction time scale τ. Results are shown for k_{cat} and K_M of 17 s^{-1} and 1.01 x 10^{-6} M, which are based on the updated kinetic parameters for LbCas12a enzyme and a dsDNA activator (**Table 1**). Also, we assumed 100 pM (E_0) of target dsDNA and 200 nM (S_0) of ssDNA reporters for **Figure 2**. Presented in **Figure 2** are results from both a numerical model that solves the complete set of coupled differential equations for the enzyme kinetics (Equations (2)-(5)) (solid line) and the analytical solution (shown as symbols in **Figure 2**) given by Equation (11). First, we see very good agreement between the analytical and numerical solution results. Also, indicated by the green dashed line is a tangent to the numerical solution at early reaction times ($t \rightarrow 0$ s), and the slope of this line is the scaled reaction velocity $v^*(= v\tau/S_0)$, where v is a quantity typically measured in Michaelis-Menten-type kinetics analysis (Equation (8)). Consistent with our normalization and species conservation (α condition; **Section 2.2**), note that the cleaved reporter fraction asymptotes to unity for large times. Also, note from **Figure 2** that τ ($= K_M/k_{cat}E_0$) represents the physically relevant product formation/reaction time scale. Consistent with our formulation (i.e. the γ condition; **Section 2.2**), we see that most of the early-time linear increase in the product amount

happens for $t/\tau < 1$. Lastly, we note that due to our choice of normalization variable, the curve showed in **Figure 2** can be fairly generalized to other concentrations of target and reporters (see Equation (11)) as long as $E_0 \ll S_0$, and if S_0 is lower than K_M by at least a factor of ~5.

Next, we study the fraction of cleaved reporters as a function of the target nucleic acid concentration and the concentration of the initial uncleaved reporter (**Figure 3**). We assume the same enzyme kinetics parameters as **Figure 2,** and study the product formed after 60 min of reaction time (commensurate with typical molecular diagnostic assays). First, note that, for initial reporter concentrations lower than about 1 μM, the fraction of cleaved reporters is nearly invariant with the initial reporter concentration (this limit is provided by the analytical solution in Equation (11)). Further, the fraction of cleaved reporters after 60 min reaction varies significantly from ~100% at target concentrations greater than ~100 pM to less than 1% for target concentrations lower than ~200 fM. Lastly, we note that the cleaved reporter fraction shown in **Figure 3** is an estimate measure of the signal-to-noise ratio requirement for assays which rely on end-point fluorescence readout for target detection. A higher initial (uncleaved) reporter concentration increases background signal, and a low fraction of cleaved reporter requires high sensitivity detectors (with low noise).

Figure 2. Fraction of reporters cleaved by the Cas enzyme versus time normalized by the reaction time scale, τ, as predicted by the numerical model (solid line) and analytical solutions (symbols). Results are shown for kinetic parameters k_{cat} and K_M of 17 s^{-1} and 1.01 x 10^{-6} M, and initial target-activated enzyme concentration of 100 pM and initial uncleaved reporter concentration of 200 nM.

Figure 3. Model predictions for the fraction of cleaved reporters $[P]$ relative to the initial amount of uncleaved reporters S_0. The cleaved reporter fraction is plotted as a function of target nucleic acid concentration and initial uncleaved reporter concentration, for a 60 min assay time, and k_{cat} and K_M of 17 s^{-1} and 1.01 x 10^{-6} M (same as **Figure 2**). Contour levels of 0 and 1 represents no reporter cleavage and complete reporter cleavage, respectively.

3.3 Scaling of assay time scales

Here, we present the variation of reaction time scale τ for varying amounts of target nucleic acid based on Michaelis-Menten kinetics analysis (Section 2.1). **Figure 4** shows contours of τ (in seconds) as a function of the Cas enzyme catalytic efficiency (k_{cat}/K_M) and the target concentration E_0. Note the inversely proportional relationship for the time scale between catalytic efficiency and target abundance: $\tau = (k_{cat}/K_M)^{-1} E_0^{-1}$. High catalytic efficiency and large target concentrations lead to shorter reaction times scales (hence faster reaction completion and fast detection). Notably, if the updated kinetics parameters in **Table 1** are true, the time scale for *trans*-cleavage reactions involving LbCas12a (for which $k_{cat}/K_M \sim 10^6$ to 10^7 M^{-1}s^{-1}) can vary from ~5 min for target concentrations greater an 1 nM to ~1 day for target concentrations less than ~1 pM. Note that, by definition, the reaction time scale τ denotes the time at which about ~63% of the reporters are cleaved.

Figure 4. Contours of the reaction time scale τ (in seconds) versus the Cas enzyme catalytic efficiency parameter k_{cat}/K_M and the target nucleic acid concentration E_0. Target concentrations of order 10,000 copies per µL or lower are likely very difficult to detect in a 60 min assay.

4. Summary

In this work, we presented a model for the *trans*-cleavage of CRISPR-associated enzyme kinetics based on Michaelis-Menten kinetics theory. We used this model to develop simple, back-of-the-envelope rules to check for consistency in experimentally derived and reported kinetic rate parameters. We applied these rules to data reported in published studies which characterize CRISPR-enzyme *trans*-cleavage rates. We found that all published studies which present Michaelis-Menten kinetics data violate these basic validation criteria. We further used our model to develop simple scaling laws relating the kinetic parameters to reaction time scales and the degree of reaction completion. We applied this theory, using updated kinetic rates for LbCas12a, to make predictions of enzyme kinetics across practically relevant ranges of target concentrations for CRISPR-detection.

Table 1. Summary of published enzyme kinetics parameters and our back-of-the-envelope checks for CRISPR-Cas12 and CRISPR-Cas13. Shaded boxes indicate violation of our back-of-the-envelope criteria for consistent kinetics data (Section 2.2).

Enzyme	Original kinetic parameters			Back-of-the-envelope checks			Checks violated[#]	Original reference	Relevant citations to the original reference	Updated kinetic paramaters**			Back-of-the-envelope checks		
	k_{cat} (1/s)	K_M (M)	k_{cat}/K_M (1/Ms)	α	β	γ				k_{cat} (1/s)	K_M (M)	k_{cat}/K_M (1/Ms)	α	β	γ
LbCas12a (ssDNA activator)	250	4.94×10^{-7}	5.1×10^8	>10	0.7	70	Rules 1, 3	Ref. 11	-	3	6.2×10^{-7}	5×10^6	0.14	0.66	0.7
LbCas12a (dsDNA activator)	1250	7.25×10^{-7}	1.7×10^9	>10	0.6	67	Rules 1, 3	Ref. 11	Refs. 10, 19, 20, 21, 22	17	1.01×10^{-6}	1.7×10^7	0.2	0.65	0.67
PbuCas13b (ssRNA activator)	987	n.r.	n.r.	>10,000	>20	-	Rules 1, 2, 3	Ref. 12	Ref. 1	1	10.86×10^{-6}	9.2×10^4	0.4	0.4	0.55
LbuCas13a (ssRNA activator)	4854	4.45×10^{-6}	1.1×10^9	>20	>7	7	Rules 1, 2, 3	Ref. 13	Ref. 14	-	-	-	-	-	-
LbuCas13a (ssRNA activator) *	204	2.62×10^{-6}	0.8×10^8	>10	0.7	46	Rules 1, 3	Ref. 14	-	-	-	-	-	-	-

** Updated values were provided to us by the corresponding authors of the original references (Refs. 23 and 24).

* Differences between parameters in Refs. 13 and 14 were attributed to the differences in number of rU in the reporter ssRNA sequences

Rule 1: $\alpha < 1$. Rule 2: $\beta \leq 1$ (maximum value of the reported measured velocity is used to compare with theoretical maximum velocity). Rule 3: $\gamma \leq O(1)$

n.r.: not reported

References

1. Fozouni, P. *et al.* Amplification-free detection of SARS-CoV-2 with CRISPR-Cas13a and mobile phone microscopy. *Cell* (2021). doi:10.1016/j.cell.2020.12.001
2. Broughton, J. P. *et al.* CRISPR–Cas12-based detection of SARS-CoV-2. *Nat. Biotechnol.* (2020). doi:10.1038/s41587-020-0513-4
3. Ackerman, C. M. *et al.* Massively multiplexed nucleic acid detection with Cas13. *Nature* (2020). doi:10.1038/s41586-020-2279-8
4. Ramachandran, A. *et al.* Electric field-driven microfluidics for rapid CRISPR-based diagnostics and its application to detection of SARS-CoV-2. *Proc. Natl. Acad. Sci. U. S. A.* (2020). doi:10.1073/pnas.2010254117
5. Sashital, D. G. Pathogen detection in the CRISPR-Cas era. *Genome Medicine* (2018). doi:10.1186/s13073-018-0543-4
6. Chertow, D. S. Next-generation diagnostics with CRISPR. *Science (80-.).* (2018). doi:10.1126/science.aat4982
7. Gootenberg, J. S. *et al.* Multiplexed and portable nucleic acid detection platform with Cas13, Cas12a and Csm6. *Science (80-.).* (2018). doi:10.1126/science.aaq0179
8. Myhrvold, C. *et al.* Field-deployable viral diagnostics using CRISPR-Cas13. *Science (80-.).* (2018). doi:10.1126/science.aas8836
9. Kellner, M. J., Koob, J. G., Gootenberg, J. S., Abudayyeh, O. O. & Zhang, F. SHERLOCK: nucleic acid detection with CRISPR nucleases. *Nat. Protoc.* (2019). doi:10.1038/s41596-019-0210-2
10. English, M. A. *et al.* Programmable CRISPR-responsive smart materials. *Science (80-.).* (2019). doi:10.1126/science.aaw5122
11. Chen, J. S. *et al.* CRISPR-Cas12a target binding unleashes indiscriminate single-stranded DNase activity. *Science (80-.).* (2018). doi:10.1126/science.aar6245
12. Slaymaker, I. M. *et al.* High-Resolution Structure of Cas13b and Biochemical Characterization of RNA Targeting and Cleavage. *Cell Rep.* (2019). doi:10.1016/j.celrep.2019.02.094
13. Shan, Y., Zhou, X., Huang, R. & Xing, D. High-Fidelity and Rapid Quantification of miRNA Combining crRNA Programmability and CRISPR/Cas13a trans-Cleavage Activity. *Anal. Chem.* (2019). doi:10.1021/acs.analchem.9b00073
14. Zhou, T. *et al.* CRISPR/Cas13a powered portable electrochemiluminescence chip for ultrasensitive and specific MiRNA detection. *Adv. Sci.* **7**, 1903661 (2020).
15. Chen, W. W., Niepel, M. & Sorger, P. K. Classic and contemporary approaches to modeling biochemical reactions. *Genes and Development* (2010). doi:10.1101/gad.1945410
16. Gootenberg, J. S. *et al.* Nucleic acid detection with CRISPR-Cas13a/C2c2. *Science (80-.).* (2017). doi:10.1126/science.aam9321
17. Jeon, Y. *et al.* Direct observation of DNA target searching and cleavage by CRISPR-Cas12a. *Nat. Commun.* (2018). doi:10.1038/s41467-018-05245-x
18. Metsky, H. C. *et al.* Efficient design of maximally active and specific nucleic acid diagnostics for thousands of viruses. *bioRxiv* (2020). doi:10.1101/2020.11.28.401877
19. Lee, R. A. *et al.* Ultrasensitive CRISPR-based diagnostic for field-applicable detection of Plasmodium species in symptomatic and asymptomatic malaria. *Proc. Natl. Acad. Sci. U. S. A.* (2020). doi:10.1073/pnas.2010196117

20.   Peng, S., Tan, Z., Chen, S., Lei, C. & Nie, Z. Integrating CRISPR-Cas12a with a DNA circuit as a generic sensing platform for amplified detection of microRNA. *Chem. Sci.* (2020). doi:10.1039/d0sc03084h

21.   Liang, M. *et al.* A CRISPR-Cas12a-derived biosensing platform for the highly sensitive detection of diverse small molecules. *Nat. Commun.* (2019). doi:10.1038/s41467-019-11648-1

22.   Nguyen, L. T., Smith, B. M. & Jain, P. K. Enhancement of trans-cleavage activity of Cas12a with engineered crRNA enables amplified nucleic acid detection. *Nat. Commun.* (2020). doi:10.1038/s41467-020-18615-1

23.   Doudna, J. A. Personal Communication. *Pers. Commun.* (2020).

24.   Zhang, F. & Slaymaker, I. M. Personal Communication. *Pers. Commun.* (2020).

23.   Doudna, J. A. Personal Communication. *Personal Communication* (2020). Personal Communication with the Communicating Author of Reference 11. We initially communicated our observations to the author on December 30, 2020, received a reply on January 4, 2021, and were provided with the updated kinetics rates on January 7, 2021.

24.   Zhang, F. & Slaymaker, I. M. Personal Communication. *Personal Communication* (2021). Personal Communication with the Communicating Authors of Reference 12. We initially communicated our observations to the authors on January 18, 2021, received a reply on January 18th, 2021, and were provided with the updated kinetic rates on January 27, 2021.