Tittle:

Ground cover presence in organic olive orchards affects the interaction of natural

enemies against *Prays oleae*, promoting an effective egg predation

Authors:

Hugo Alejandro Álvarez[a]*, Raquel Jiménez-Muñoz[a], Marina Morente[a], Mercedes

Campos[b], Francisca Ruano[a]

Affiliations:

[a] Department of Zoology, University of Granada, Granada, Spain

[b] Department of Environmental Protection, Zaidin Experimental Station (CSIC),

Granada, Spain

* Corresponding author at: Department of Zoology, Faculty of Sciences, University of

Granada, Av. Fuente Nueva s/n 18071, Granada, Spain.

E-mail addresses: hugoalvarez01@gmail.com, hugoalvarez01@correo.ugr.es (H.A.

Álvarez)

Abstract

The olive moth, *Prays oleae*, is one of the most common insects that damages olives in

the Mediterranean region. The establishment of ground cover within olive orchards has

been promoted in this region in recent years to avoid erosion and soil degradation.

Nevertheless, its role as a shelter for natural enemies of pests has been controversial. In

this study, we have investigated the effectiveness of the biological control of *P. oleae* in

organic olive orchards with ground cover (mowed) and without ground cover (tilled).

For this, (1) we assessed the relationship between predated eggs and the abundance of

natural enemies in both types of orchards; (2) we compared both the potential damage

of the pest and the egg hatching in the two types of orchards; and (3) we examined the

interaction amongst families of natural enemies and *P. oleae* (as adults and as predated

eggs). The results showed that there is a high rate of predation in the studied olive

orchards, 81% of the eggs were predated, 12.2% hatched, and 6.9% were live eggs.

However, mowed orchards were more effective for controlling *P. oleae* by means of

egg predation rather than tilled orchards, i.e., in mowed orchards, whilst the potential

damage of the pest was higher, egg hatching was rather low. The structure of the adult

arthropod community, i.e., the composition and abundance of families of natural

enemies did not differ between the orchards, but the abundance of the families

Anthocoridae, Miridae and Scelionidae was significantly higher in the mowed orchards.

Finally, the interaction amongst natural enemies and *P. oleae* showed that the families

that better explained the effects on egg predation were Aeolothripidae, Anthocoridae,

Miridae, Chrysopidae (predators), and Formicidae (omnivore). We discuss the results in

terms of ecological interactions of trophic guilds and we conclude that the establishment

and maintenance of ground cover in organic olive orchards, at least in June and July, is

of great significance because it positively affects the egg predation of *P. oleae*. This

effect is especially significant when there is a low abundance of natural enemies in the

olive orchards.

Keywords:

Biological control

Ecological infrastructures

Inter-row management

Trophic guilds

Trophic interactions

1. Introduction

The organic management in olive orchards has been increasing in the Mediterranean

region in recent years (Alonso-Mielgo et al., 2001; Torres-Millares et al., 2017). This

type of management frequently involves the establishment of ground cover within the

orchard, and when possible, the conservation of adjacent semi-natural habitats (Boller et

al., 2004; Landis et al., 2005; Malavolta and Perdikis, 2018). In this region, one of the

most common insects that damages olives is the olive moth, *Prays oleae* Bern

(Lepidoptera: Praydidae) (Tzanakakis, 2006; RAIF, 2018). *Prays oleae*, produces three

generations per year: (1) the phyllophagous generation (feeds on olive leaves from

November to April, and overwinters in the canopy); (2) the anthophagous generation

(feeds on floral buttons from April to June and is the one that lays eggs mainly on the

chalice of the olive fruits); and (3) the carpophagous generation (larvae penetrate the

fruit and feed on the stone from June to October). All three generations can cause

damage to olive orchards and each generation plays an important role in configuring the

size of the next generation. However, the carpophagous larvae can generate significant

damage to olives, which potentially reduces the yield production. Thus, much of the

efforts of pest control are focused on the anthophagous and carpophagous generations

(Ramos et al., 1998; Bento et al., 2001).

Recently, it has been observed that olive orchards have great potential to boost

populations of natural enemies within the orchard, especially when ground cover is

present rather than orchards with bare ground (Herz et al., 2005; Lousão et al., 2007;

Cárdenas et al., 2012; Rodríguez et al., 2012; Paredes et al., 2013a). From a

"biodiversity-ecosystem function" point of view, semi-natural vegetation interspersed

within the growing area or located at their margins can reinforce microclimate

conditions in crops and orchards, and thus provide food and shelter to natural enemies

of insect pests (Tscharntke et al., 2012; Wan et al., 2018). Accordingly, in olive

orchards ground cover plays a major role in modulating such a tendency. For example,

it has been suggested the existence of synergistic effects between ground cover and

natural adjacent vegetation, which jointly promote a high abundance of some (but not

all) predator arthropods of *P. oleae* and *Euphyllura olivina* Costa (Hemiptera: Psyllidae)

within the olive-tree canopy (Paredes et al., 2013a). Recently, Álvarez et al. (2019a)

demonstrated such a synergistic relationship by describing the abundance and

movement of the natural enemies which are boosted by the ground cover. Moreover,

Villa et al. (2016a) observed that ground cover favoured the parasitism of *P. oleae*

larvae by *Ageniaspis fuscicollis* (Dalman) (Hymenoptera: Encyrtidae), whereas

herbicide applications had negative effects.

Nonetheless, a higher abundance of natural enemies does not always suppress pest

abundance and pest damage, which is a problem that has arisen in conservational

biological control (Rusch et al., 2010; Karp et al., 2018). In addition, and unfortunately,

it has been recognized that the positive effects generated by a higher biodiversity on

ecosystem function, i.e., the control of pests, are conditioned by a myriad of factors

(Bianchi et al., 2006; Rusch et al., 2010; Tscharntke et al., 2016; Karp et al., 2018).

Significant efforts have been made by various authors to describe the effects of semi-

natural habitats on the abundance of natural enemies and olive pests (Ruano et al., 2004;

Paredes et al., 2013a; 2013b; Gkisakis et al., 2016; Villa et al., 2016a; 2016b; Porcel et

al., 2017; Álvarez et al., 2019a). However, to the best of our knowledge, there is no

study that has focussed on jointly assessing the effects of ground cover on both the

abundance of natural enemies and the egg predation of *P. oleae*. The aim of this study

was to assess the effectiveness of predation in organic olive orchards with both tilled

and mowed management of the ground cover. Specific goals of the study were: (1) to

assess the relationship between the abundance of natural enemies and egg predation in

both managements, (2) to compare the effectiveness of egg predation between both

managements, and (3) to explore the interaction amongst families of natural enemies

and *P. oleae* adults and predated eggs using unconstrained ordination. We have

hypothesised that when a ground cover is mowed (1) key taxa of natural enemies would

be positively affected; therefore (2) the biological control of *P. oleae*, by means of egg

predation, would increase.

2. Material and methods

2.1. Study area and sampling design

The study was conducted in three consecutive years from 2011 to 2013 in southern

Spain, in the province of Granada. We selected eight organic olive orchards based on

(1) the absence of ground cover (tillage in late spring: tilled) and (2) the use of mowing

techniques during late spring to maintain the ground cover (mowed) (see Table A.1 in

supplementary data). All the orchards were located in areas surrounded by extended

semi-natural habitats interspersed in an olive-orchard matrix including different

management systems (Fig. 1). Agricultural management in these organic orchards was

based on a system of natural regulation (*sensu* Pajarón Sotomayor, 2006), and thus, pest

management did not differ amongst the orchards in the years of study. The distance of

planting was 10×10 m, and two varieties of olive trees were grown: Picual (location

Deifontes) and Lucio (location Granada). The climatic and topographic conditions were

typical of the olive orchards in the study area (see Paredes et al., 2013a; Álvarez et al.,

2019a).

June and July are the months when the anthophagous generation of *P. oleae* laid their

eggs on newly growing olives. We carried out three different types of sampling in both

months per year. Firstly, adult arthropods were collected twice a month by batting four

branches per tree over an entomological net (a sample per tree). Olive trees were

sampled in randomly selected plots formed by four parallel transects with a separation

of 100 m between the transects. Each transect consisted of ten trees of which only five

trees were sampled, following a discontinuous sequence, i.e., 20 samples per plot. After

being collected, the samples were transported to the Department of Zoology, University

of Granada and the Zaidin Experimental Station. The samples were stored individually

and maintained at $-20°C$ until the specimens were identified. The arthropods were

identified to family level, otherwise specified, and the natural enemies were separated

and used for this study. Identification of the natural enemies was based on literature data

(see Table A.4 in supplementary data). Secondly, 200 olives were collected from four

trees (total of 50 olives per tree) randomly selected in each orchard per month (June and

July) and the same tree was never re-sampled. The olives were collected to examine the

eggs laid by *P. oleae*. Thirdly, *P. oleae* adults were collected using pheromone traps (2

traps per orchard), which were randomly distributed in each orchard and changed every

10 days in June and July. Adults and eggs of *P. oleae* were stored and identified at the

Zaidin Experimental Station (CSIC).

2.2. Identification of egg damage

The olives were observed with the help of a microscope-stereoscope to record the

number of olives with laid eggs of *P. oleae*, and to characterize the appearance of the

eggs (Fig. 1). Then, the number of (1) eggs that had hatched and give place to a larva

inside the olive was recorded (hatched eggs); (2) eggs that had not hatched and were

still alive (live eggs) and showed a white to yellowish colour; and (3) eggs that had been

damaged by predators (predated eggs) and of which only the translucent chorion

adhering to the fruit remained.

2.3. Data analysis

For comparison purposes the site (orchard) was used as our experimental unit. Thus, we

pooled together samples by (1) orchard and (2) months for the three years to avoid

pseudo-replication. The orchards were not always the same throughout years, of the

eight orchards, four were sampled in 2011 (2 tilled and 2 mowed), five in 2012 (3 tilled

and 2 mowed, of which 2 were new), and seven in 2013 (2 tilled and 5 mowed, of which

2 were new) (see Table A.1 in supplementary data). Monthly samples were considered

independently. Therefore, for each site we obtained a representative measure of

arthropod abundance and *P. oleae* egg counts.

Raw data of the abundance of natural enemies and *P. oleae* adults was subjected to a

logistic regression approach in order to detect differences between managements. We

used this method instead of mean or median comparison because it is a more suitable

method to detect statistical differences due to the nature of our experimental unit (see

Peng et al., 2002).

We used two approaches to assess the differences in egg predation between the tilled

and mowed managements. Firstly, we fitted a generalized linear mixed model (GLMM)

using a Poisson tendency to test whether or not the relationship between the abundance

of natural enemies and the amount of the predation changed in the two types of

managements. We used a GLMM approach because our experimental unit (orchard)

changed throughout the years of study, in some of the re-sampled orchards farmers

passed from a tilled to a mowed management. Then, to control for such inter-annual

variation and site-management variation we included in the model year and site as

nested random effects. The number of predated eggs was included as the dependent

variable and the type of management and the total abundance of natural enemies were

included as fixed effects (see Table A.2 in supplementary data). Secondly, we assessed

the effectiveness of predation by analysing the potential damage of the pest (number of

olives with any kind of eggs laid $\times$ 100 / total of observed olives) and the rate of egg

hatching (number of hatched eggs $\times$ 100 / all observed eggs minus the predated ones)

(for more detail on these parameters see Ramos et al., 1987; Ramos and Ramos, 1990).

Then, we subjected the data after this transformation to a logistic regression approach in

order to compare both parameters and detect differences between managements.

Finally, non-metric multidimensional scaling (NMDS) was used to assess the overall

pattern of species composition of the natural enemies. Data used for the NMDS were

square-root transformed and subjected to Wisconsin double standardization (Legendre

and Gallagher, 2001). The Bray-Curtis dissimilarity distance was used to compute the

resemblance matrix amongst sites. Species scores, representing the different natural

enemy taxa were added to the final NMDS plot as weighted averages. Based on the

NMDS, smooth surfaces were generated with the data of *P. oleae* adult abundance and

predated eggs to explore associations between families of natural enemies and *P. oleae*.

Smooth surfaces result from fitting thin plate splines in two dimensions using

generalized additive models. The function selects the degree of smoothing by

generalized cross-validation and interpolates the fitted values on the NMDS plot

represented by lines ranking in a gradient (Oksanen et al., 2018) (see Table A.3 in

supplementary data). This method allowed us to indirectly relate different levels of the

abundance of predated eggs and adults of *P. oleae* with the abundance and

correspondence of different families of natural enemies.

Analyses were computed in the R software v.3.6.2 (R Developmental Core Team,

2019). Accordingly, the "lme4" package (Bates et al., 2020) was used to fit GLMM and

the "vegan" package (Oksanen et al., 2018) was used to compute NMDS and smooth

surfaces. Package lme4 was used because whilst other packages are more mature and

better documented, lme4 is fastest, offers built-in facilities for likelihood profiling and

parametric bootstrapping and especially it offers tools for crossed designs (Bates, 2010;

Bates et al., 2018).

3. Results

Overall, 6,400 olives were observed with a total of 15,412 laid eggs of *P. oleae*. 81% of

the eggs were predated, 12.2% hatched, and 6.9% were live eggs. We collected a total

of 62,008 adults of *P. oleae* and a total of 4,001 natural enemy arthropods, of which 36

families were identified (Table 1). 70.7% of the natural enemy specimens were

predators, 26.8% were omnivores, and 2.5% were parasitoids. The most abundant

families of predators were: Anthocoridae, Miridae (order Hemiptera); Chrysopidae

(order Neuroptera); and Thomisidae (order Araneae). Amongst hymenopterans the most

abundant family of parasitoids was the Scelionidae, and Formicidae was the most

abundant family of omnivores and of all the natural enemies.

The structure of the arthropod community, i.e., the composition and abundance of

arthropod families, is mostly the same between tilled and mowed orchards (Fig. 2).

However, six families of natural enemies were present only in one of the two

managements, i.e., mowed: Liocranidae, Uloboridae (order Araneae), Nabidae (order

Hemiptera), Aeolothripidae (order Thysanoptera); and tilled: Malachiidae (order

Coleoptera), Ichneumonidae (order Hymenoptera). Moreover, the abundance of three

families of natural enemies was significantly higher in the mowed orchards,

Anthocoridae (Wald χ^2 = 3.928, df = 1, p = 0.047), Miridae (order Hemiptera) (Wald χ^2

= 5.247, df = 1, p = 0.021), and Scelionidae (order Hymenoptera) (Wald χ^2 = 5.071, df

= 1, p = 0.024) (Fig. 2), as well as the abundance of *P. oleae* adults (Wald χ^2 = 4.624, df

= 1, p = 0.031).

GLMM analysis showed that there is a positive relationship between the abundance of

natural enemies and the amount of predated eggs in both managements (Fig. 3). Both

effects, natural enemy abundance and management, were statistically significant (Table

2). The relationship tended to be high in the mowed orchards and this pattern appeared

at the lowest abundance of natural enemies (Fig. 3). Furthermore, there are differences

in the effectiveness of predation of *P. oleae* eggs between managements, i.e., in mowed

orchards the potential damage of the pest was significantly higher (Wald χ^2 = 8.996, df

= 1, *p* = 0.002) but egg hatching was significantly lower (Wald χ^2 = 5.295, df = 1, *p* =

0.021) than tilled orchards.

The results of the NMDS, which represents the relationship and structure of the

communities of natural enemies and their association with egg predation and the

abundance of adults of *P. oleae*, are shown in Figure 4. Accordingly, the families

Araneidae, Linyphiidae, Liocranidae, Oxiopidae, Salticidae, Theridiidae, Thomisidae,

Uloboridae (order Araneae); Coccinelidae, Malachiidae, Staphylinidae (order

Coleoptera); Anthocoridae, Lygaeidae, Miridae, Nabidae, Reduviidae (order

Hemiptera); Braconidae, Elasmidae, Formicidae, Pteromalidae, Scelionidae (order

Hymenoptera); Chrysopidae (order Neuroptera); Mantidae (order Mantodea);

Raphidiidae (order Raphidioptera); and Aeolothripidae (order Thysanoptera) were

associated with elevated egg predation (Fig. 4a). However, the families Corinidae,

Oxiopidae, Salticidae (order Araneae); Cantharidae, Coccinelidae (order Coleoptera);

Lygaeidae (order Hemiptera); Encyrtidae, Eulophidae, Ichneumonidae (order

Hymenoptera); and Coniopterygidae (order Neuroptera) were associated with a low

abundance of *P. oleae* adults (Fig. 4b). In this type of analysis, an association of a

family of natural enemies with a high number of predated eggs implies that these taxa

could be involved in egg predation (or egg damage), increasing predation rates.

Conversely, an association with a low or intermediate abundance of adults of *P. oleae*

means that such taxa could be feeding on adults, decreasing their abundance to a lower

rate.

Based on the NMDS, trophic status, size, and the morphological features of each family,

the families that are more likely to damage the eggs of *P. oleae* are: Coccinelidae,

Staphylinidae (order Coleoptera); Anthocoridae, Miridae, Nabidae, Reduviidae (order

Hemiptera); Braconidae, Formicidae (order Hymenoptera); Chrysopidae (order

Neuroptera) and Aeolothripidae (order Thysanoptera). However, we have ruled out the

families that had (1) a low abundance in both managements (Coccinelidae,

Staphylinidae, Nabidae, Reduviidae), and (2) no predatory form of feeding (parasitoids:

Braconidae). Hence, only Aeolothripidae, Anthocoridae, Chrysopidae, Formicidae, and

Miridae are the families that could explain the differences in egg predation between the

tilled and mowed orchards.

4. Discussion

In this study, we have assessed the effectiveness of the biological control of *P. oleae* in

organic olive orchards in terms of egg predation by natural enemies. As expected, the

presence of ground cover within the orchards positively affected the predation of eggs

laid by the antophagous generation of *P. oleae*, and thus, egg predation was more

effective in mowed orchards than in tilled orchards.

Organic orchards are very balanced and stable systems (Vossen, 2007). The orchards

that we measured were very similar in their agricultural practices and in their landscape

structure (with the exception of the ground cover management). This was reflected in

the composition and abundance of families of natural enemies. Both local and large-

scale factors can affect the abundance of natural enemies and pests in olive orchards,

such as less pesticide application or microclimate conditions and landscape diversity or

patch size, respectively (Boccaccio and Petacchi, 2009; Rodríguez et al., 2009; Ortega

and Pascual, 2014; Villa et al., 2016a; 2020; Morente et al., 2018; Álvarez et al., 2019a;

2019b; 2021). This may explain why the structure of the arthropod community tended

not to differ in our study. Nonetheless, it has been shown that the abundance of natural

enemies is positively affected by ground cover (Lousão et al., 2007; Cárdenas et al.,

2012; Rodríguez et al., 2012; Álvarez et al., 2019a; 2019b). When analysing the

relationship between egg predation and natural enemy abundance in both managements,

mowed orchards tended to have higher predation as the abundance of natural enemies

increased. Interestingly, the highest differences in predated eggs between mowed and

tilled orchards appeared especially when the levels of natural enemies were low. This

implies that the differences in predation were caused by a subtle, but still higher

abundance of natural enemies.

On the other hand, previous studies on olive orchards showed that the presence of

ground cover had no effect on the abundance of *P. oleae* adults when tilled and mowed

orchards were compared (Paredes et al., 2013b; Paredes et al., 2015a). Nonetheless, it

has been found that certain plant species in the ground cover could promote an increase

in the abundance of *P. oleae* (Villa et al., 2016c). Our results follow such a tendency,

ground cover may increase the abundance of adults of *P. oleae*. Based on the former

results one could assume that ground cover does not promote biological control by

itself. However, we showed that egg hatching of *P. oleae* is lower in orchards with a

ground cover, and thus, less hatching implies that there are potentially less larvae of *P.*

oleae that could damage olives. Therefore, we can assume that the effects of the ground

cover to control *P. oleae* lead towards the predation of eggs rather than attacking the

adults.

Regarding the predators, our results are in agreement with previous studies that have

recorded the role of predator heteropterans, such as Anthocoridae and several species of

Miridae as major predators of olive pests (Mazomenos et al., 1994; Cantero, 1997;

López-Villalta, 1999; Morris et al., 1999; Alvarado et al., 2004; Paredes et al., 2013a;

2015b). The fact that these groups are positively affected by the presence of ground

cover suggests their sensitivity to perturbation. However, it has been shown that

predator heteropterans are more sensitive to the presence of native adjacent vegetation

rather than ground cover, although some species such as *Deraeocoris punctum*

(Rambur), have shown the opposite (Paredes et al., 2013a) implying that differences at

species level are important. These inconsistencies may be the result of the movement of

the natural enemies across habitats within and outside olive orchards. For example,

Álvarez et al. (2019a) showed that predators and parasitoids move from ground cover to

adjacent vegetation and olive trees, respectively, but omnivores move from adjacent

vegetation to ground cover and olive trees, specifically when the ground cover withers.

In addition, there is evidence that *Anthocoris nemoralis* (Fabricius), (Plata et al., 2017)

and some lacewings (Chrysopidae) (Porcel et al., 2017) move from ground cover to

adjacent vegetation.

Lacewing larvae, for example, have been described as one of the main natural enemies

that attack *P. oleae* (Bento et al., 1999; Torres, 2006; Villa et al., 2016b). Lacewing

larvae are positively affected by ground cover (Villa et al., 2016b). Interestingly, in our

study this group did not differ between the tilled and mowed orchards, although we

mainly sampled adults which are very stable geographically (Alcalá-Herrera et al.,

2019). This could be explained by the fact that winged adults have high ranges of

movement (Rusch et al., 2010) and it is possible that they move across the region from

one orchard to another to lay their eggs. Nevertheless, in our study, lacewings were

related with high levels of egg predation, which supports the hypothesis that this group

is of great importance in the biological control of *P. oleae*.

Another family that showed interesting patterns was Aeolothripidae. In our study, it

showed an important interaction with egg predation, but this family was present only in

the mowed orchards. The role of Aeolotrhipidae as a natural enemy of olive pests has

been poorly documented (reviewed by Torres, 2006). It is known that Aeolothripidae

attack other Thysanoptera, however, it also has been suggested that some genera of

Aeolothripidae can feed on the eggs of lepidopterans (Lewis, 1973). Moreover, some

species of European *Aeolothrips* sp. can feed on mites, larvae, and eggs of psyllids and

whiteflies, as well as on aphids (Trdan et al., 2005). Thus, the role of Aeolothripidae in

the predation of the eggs of *P. oleae* should be investigated more thoroughly.

In the case of the omnivores, it is known that ants are important predators of *P. oleae*

(Morris et al., 1999; 2002). Our results showed that ants had the highest abundance

within olive orchards, however, we did not found differences in the abundance of ants

between mowed and tilled orchards. This is of great importance, because a predator that

is not affected by management and has high abundances could be used to enhance local

biological control strategies. Several studies have shown *Tapinoma* ants as the most

abundant type of ant within olive orchards, sometimes representing more than 50 % of

the relative abundance amongst omnivores within olive orchards (Morris et al., 1998a;

1998b;1999; 2002; Morris and Campos, 1999; Redolfi et al., 1999; Pereira et al., 2004;

Rodriguez et al., 2005; Santos et al., 2007; Campos et al., 2011), which makes it one of

the strongest candidates for controlling *P. oleae*. Indeed, some species of the *T.*

nigerrimum complex are beneficial in olive orchards in the southern area of the Iberian

Peninsula (Seifert et al., 2017). Furthermore, it has been found that it is possible to

boost the abundance and trophic interactions of *Tapinoma* ants within the canopy of

olive trees with mature ground cover (Álvarez et al., 2019b) and less pesticide use

(Morente et al., 2018).

In addition, we found that the family Braconidae shows an important association with

predators upon egg predation. Only one species of Braconidae is known to parasitize the

eggs of *P. oleae*: *Chelonus eleaphilus* Silv (Arambourg, 1986). This species is a

poliembrionic parasite, i.e., females oviposit inside the eggs of their prey (Grbic et al.,

1998; Segoli et al., 2010). This species is one of the most important and specific

parasitoids of *P. oleae* in the Mediterranean region. The fact that Braconidae

populations respond to ground cover management may be due to their need for flowers

to feed on (Nave et al., 2016). On the other hand, in our analysis Elasmidae and

Scelionidae showed a similar pattern to Braconidae, but Scelionidae had higher

abundances in mowed orchards. Some species of the family Scelionidae may attack

other natural enemies causing intra-guild predation, such as *Telenomus acrobater* Giard,

which has been described as parasitizing the eggs of lacewings (Alrouedchi, 1981;

Campos, 1986; Rodríguez et al., 2012).

Finally, it is important to point out that the species composition of natural enemies

showed interesting patterns (Fig. 4). The fact that the tendency of the variable predated

eggs depends on the tendency of *P. oleae* adult abundance should be taken into account,

i.e., the more *P. oleae* adults there are, the more the eggs they can lay, and thus, be

predated. This is why the panels in Figure 4, are very similar. However, the families that

are related with the predation on adults are well defined. The spiders are the arthropods

that are most likely to predate adults, which is in agreement with previous studies

(Paredes et al., 2015b). Interestingly, and according to what we have mentioned, in the

NMDS natural enemies assembled in different groups that correspond to their trophic

status. Consequently, several assemblages might fulfil complementary functional roles

determined by the way they catch prey (Uetz et al., 1999; Straub et al., 2008). For

example, it has been found that a single assemblage of natural enemies, such as *A.*

nemoralis, *Brachynotocoris* sp., and *Pseudoloxops coccineus* (Meyer Dur), is better

correlated with the control of *P. oleae* (Paredes et al., 2015b). The assemblage of these

species has been explained as the result of the (complex) life cycle of *P. oleae* (Wilby et

al., 2005) and their preference for eggs (Paredes et al., 2015b). In addition, the arachnid

families Araneidae and Linyphiidae, which are orb-weaving and sheet-weaving spiders

respectively, are most likely to play a role in reducing the adults of *P. oleae* (Paredes et

al., 2015b).

5. Conclusions

A mowing management of the ground cover within olive orchards positively affected

the key natural enemies that play an important role predating *P. oleae*, even though

arthropod communities were similar between tilled and mowed orchards. Hence, the

establishment and maintenance of ground cover in organic olive orchards is of great

significance due to its potential to promote the biological control of *P. oleae* by means

of egg predation, especially when there is a low abundance of natural enemies. The

hypothesis is that an olive orchard with ground cover produces more active and

voracious natural enemies, and it may allow the establishment of more and efficient key

predators. To the best of our knowledge this is the first time that this type of empirical

data has been recorded for olive orchards. The fact that in our study the differences in

the biological control of pests were shown by eggs rather than the abundance of adults

suggests that the studies on biological control should focus on specific instars of the

development of pests where the biological control is more likely to occur, which is a

concern that has already been pointed out for conservation biological control (Karp et

al., 2018). Thus, the effect of landscape structure on egg predation of *P. oleae*

specifically in olive orchards needs to be investigated more thoroughly.

Authors contributions

H.A.A. and F.R. conceived the ideas. F.R. and M.C. designed fieldwork and R.J.M. and

M.M. conducted fieldwork. H.A.A. performed data analysis and write the manuscript.

All authors revised the final version and gave their approval for submission.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors want to thank the owners of the orchards, especially Norberto Recio and

Manuel Recio; Carlos Martínez for his help in the laboratory (UGR collaboration

grant); and all the taxonomist who helped us identifying arthropods. Thanks to Angela

Tate for editing the English language of the manuscript. H. A. Álvarez is grateful to

Gemma Clemente Orta for her help conceptualizing graphic representations and

CONACyT for providing him with an international PhD student grant (registry

332659). This study was financed by The Alhambra and Generalife Governing Board

(contract: 3548-00 and 3548-01) and the Spanish Ministry of Science, Innovation, and

Universities, General Sub-direction of Projects (project AGL2009-09878).

References

Alcalá-Herrera, R., Campos, M., Ruano, F., 2019. Late summer oviposition of green

lacewings (Neuroptera: Chrysopidae) on olive groves and adjacent trees.

Environm. Entomol. 48, 506-513.

Alonso Mielgo, A.M., Sevilla Guzmán, E., Jiménez Romera, M., Guzmán Casado, G.,

2001. Rural development and ecological management of endogenous resources:

the case of mountain olive groves in Los Pedroches comarca (Spain). J. Environ.

Pol. Plan. 3, 163-175.

Alrouechdi, K., Panis. A., 1981. Les parasites de *Chrysoperla carnea* Steph.

(Neuroptera, Chrysopidae) sur Olivier en Provence. Agronomie 1, 139-141.

Alvarado, M., Civantos, M., Duran, J.M., 2004. Plagas. In: Barranco, D., Fernández-

Escobar, R., Rallo, L. (Eds.), El cultivo del olivo. Ediciones Mundi-Prensa,

Madrid, pp. 433-493.

Álvarez, H.A., Morente, M., Oi, F.S., Rodríguez, E., Campos, M., Ruano, F., 2019a.

Semi-natural habitat complexity affects abundance and movement of natural

enemies in organic olive orchards. Agric. Ecosyst. Environ. 285, 106618.

Álvarez, H. A., Morente, M., Campos, M., Ruano, F., 2019b. La madurez de las

cubiertas vegetales aumenta la presencia de enemigos naturales y la resiliencia

de la red trófica de la copa del olivo. Ecosistemas 28, 92-106.

Álvarez, H.A., Morente, M., Ruano, F., 2021. Habitat complexity in organic olive

orchards modulates the abundance of natural enemies but not the attraction to

plant species. BioRxiv 2021.02.04.429588.

Arambourg, Y., 1986. Traité d'Entomologie Oléicole. Consej Oleícola Internacional,

Madrid, Spain, 360 p.

Bates, D., 2010. Lme4: Mixed-effects modeling with R. http://lme4.r-forge.r-

project.org/book/front.pdf

Bates, D., et al. 2018. Linear Mixed-Effects Models using 'Eigen' and S4.

https://CRAN.R-project.org/package=lme4

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R.H.B., Singmann, H.,

Dai, B., Scheipl, F., Grothendieck, G., Green, P., Fox, J., 2015. Package 'lme4'.

Convergence 12, 2.

Bento, A., 1999. Contribuicção para o estabelecimento de um programa de

proteccção integrada contra a tracça da oliveira, *Prays oleae* (Bern.) em Trás-

os-Montes. PhD thesis. Universidade de Trás-os-Montes e Alto Douro, Vila

Real, Portugal.

Bento, A., Torres, L., Lopes, J., Pereira, J.A., 2001. Avaliacção de prejuízos causados

pela tracça da oliveira, Prays oleae (Bern.) em Trás-os-Montes. Rev. Cie. Tec.

Agr. 24, 89-96.

Bianchi, F., Booij, C.J.H., Tscharntke, T., 2006. Sustainable pest regulation in

agricultural landscapes: a review on landscape composition, biodiversity and

natural pest control. Proc. R. Soc. B: Biol. Sci. 273, 1715-1727.

Boccaccio, L., Petacchi, R., 2009. Landscape effects on the complex of *Bactrocera*

oleae parasitoids and implications for conservation biological control.

BioControl 54, 607-616.

Boller, E., Ha□ni, F., Poehling, H.M., 2004. Ecological Infrastructures: Ideabook on

Functional Biodiversity at the Farm Level. IOBC, Lindau.

Campos, M., 1986. Influencia del complejo parasitario sobre las poblaciones de

Chrysoperla carnea (Neuroptera, Chrysopidae) en olivares del sur de España.

Neuroptera Int. 4, 97-105.

Campos, M., Fernández, L., Ruano, F., Cotes, B., Cárdenas, M., Castro, J., 2011. Short

term response of ants to the removal of ground cover in organic olive orchards.

Eur. J. Entomol. 108, 417-423.

Cantero, F.A., 1997. Enfermedades y plagas del olivo. Riquelme y Vargas Ediciones,

S.L. Jaen, Spain, 646 p.

Cárdenas, M., Castro, J., Campos, M., 2012. Short-term response of soil spiders to

cover-crop removal in an organic olive orchard in a Mediterranean setting. J.

Insect Sci. 12, 1-18.

Gkisakis, V., Volakakis, N., Kollaros, D., Bàrberi, P., Kabourakis, E.M., 2016. Soil

arthropod community in the olive agroecosystem: determined by environment

and farming practices in different management systems and agroecological

zones. Agric. Ecosyst. Environ. 218, 178-189.

Grbic, M., Strand, M.R., 1998. Shifts in the life history of parasitic wasps correlate with

pronounced alterations in early development. Proc. Natl. Acad. Sci. 95, 1097-

1101.

Herz, A., et al., 2005. Towards sustainable control of lepidopterous pests in olive

cultivation. Gesunde Pflanz. 57, 117-128.

Malavolta, C., Perdikis, D., 2018. Crop Specific Technical Guidelines for Integrated

Production of Olives. IOBC-WPRS Commission IP Guidelines. 4[th] edition.

https://www.iobc-

wprs.org/members/shop_en.cfm?mod_Shop_detail_produkte=193

Karp, D.S., et al., 2018. Crop pests and predators exhibit inconsistent responses to sur-

rounding landscape composition. Proc. Nat. Acad. Sci. 115, E7863-E7870.

Landis, D.A., Menalled, F.D., Costamagna, A.C., Wilkinson, T.K., 2005. Manipulating

plant resources to enhance beneficial arthropods in agricultural landscapes.

Weed Sci. 53, 902-908.

Legendre, P., Gallagher, E.D., 2001. Ecologically meaningful transformations for ordi-

nation of species data. Oecologia 129, 271-280.

Lewis, T., 1973. Thrips. Their Biology, Ecology and Economic Importance. Academic

Press, New York, 350 p.

López-Villalta, M.C., 1999. Olive pest and disease management. International Olive Oil

Council, Madrid, 207 p.

Loumou, A., Giourga, C., 2003. Olive groves: The life and identity of the

Mediterranean. Agr. Hum. Val. 20, 87-95.

Lousão, C., Bento, A., Campos, M., Ruano, F., Pereira, J.A., 2007. Effects of no tillage

on the abundance and diversity of soil and olive tree canopy arthropods. In:

Kalaitzaki, A. (Ed.), 3rd European Meeting of the IOBC/WPRS study Group

''Integrated Protection of Olive Crops'', Bragança, 131 p.

Mazomenos, B.E. Petrakis, P.V. Roussis, V., Leukidou, I., Ortiz, A., Stefanou, D.,

Pantazis, A., 1994. Natural enemies of major olive pests: community level

model of action. The development of environmentally safe pest controlsystems

for the european olive. AGRE 0013: Éclair Final Tech. Prog. Report, 1990-1994.

Morente, M., Campos, M., Ruano, F., 2018. Evaluation of two different methods to

measure the effects of the management regime on the olive-canopy arthropod

com- munity. Agric. Ecosyst. Environ. 259, 111-118.

Morris, T., Campos, M., 1999. Entomofauna depredadora del suelo del olivar. Zool.

baetica, 10, 149-160.

Morris, T.I., Symondson, W.O.C., Kidd, N.A.C., Jervis, M.A., Campos, M., 1998a. Are

ants significant predators of the olive moth, *Prays oleae*? . Crop Prot. 17, 365-

366.

Morris, T.I., Campos, M., Jervis, M.A., McEwen, P.K., Kidd, N.A.C., 1998b. Potential

effects of various ants species on green lacewing, *Chrysoperla carnea*

(Stephens) (Neuroptera, Chrysopidae) egg number. J. Appl. Ent. 122, 401-403.

Morris, T.I., Campos, M., Kidd, N.A.C., Symondson, W.O.C., 1999. What is

consuming *Prays oleae* (Bernard) (Lep. Yponomeutidae) and when: a

serological solution? . Crop Prot. 18, 17-22.

Morris, T.I., Symondson, W.O.C., Kidd, N.A., Campos, M., 2002. The effect of

different ant species on the olive moth, *Prays oleae* (Bern.), in Spanish olive

orchard. J. Appl. Entomol. 126, 224-230.

Oksanen, J., et al., 2018. Vegan: Community Ecology Package. R package version 2.5-

2. Available at: http://CRAN.R- project.org/package=vegan

Ortega, M., Pascual, S., 2014. Spatio-temporal analysis of the relationship between

landscape structure and the olive fruit fly *Bactrocera oleae* (Diptera:

Tephritidae). Agric. For. Entomol. 16, 14-23.

Pajarón Sotomayor, M., 2006. El cultivo del olivar en producción ecológica. Asociación

para el Desarrollo Sostenible del Poniente Granadino. Proyecto Columela. Loja,

Granada, España. 15 p.

https://www.juntadeandalucia.es/export/drupaljda/olivar.pdf

Paredes, D., Cayuela, L., Campos, M., 2013a. Synergistic effects of ground cover and

adjacent vegetation on natural enemies of olive insect pests. Agric. Ecosyst.

Environ. 173, 72-80.

Paredes, D., Cayuela, L., Gurr, G.M., Campos, M., 2013b. Effect of non-crop

vegetation types on conservation biological control of pests in olive groves.

PeerJ 1, e116.

Paredes, D., Cayuela, L., Gurr, G.M., Campos, M., 2015a. Is ground cover vegetation

an effective biological control enhancement strategy against olive pests?. PLoS

One 10, e0117265.

Paredes, D., Cayuela, L., Gurr, G.M., Campos, M., 2015b. Single best species or natural

enemy assemblages? A correlational approach to investigating ecosystem

function. BioControl 60, 37-45.

Peng, C.Y.J., Lee, K.L., Ingersoll, G.M., 2002. An introduction to logistic regression

analysis and reporting. J. Educ. Res. 96, 3-14.

Pereira, J.A., Bento, A., Cabanas, J.E., Torres, L.M., Herz, A., Hassan, S.A., 2004. Ants

as predators of the egg parasitoid *Trichogramma cacoeciae* (Hymenoptera:

Trichogrammatidae) applied for biological control of the olive moth, *Prays*

oleae (Lepidoptera: Plutellidae) in Portugal. Biocontrol Sci. Techn. 14, 653-664.

Plata, A., Paredes, D., Moreno-Chocano, J., Manjón-Cabezas, J., Campos, M., 2017.

Efecto de la vegetación adyacente en el control biológico de plagas de olivar: el

papel del depredador *Anthocoris nemoralis*. XXXIII Jornadas de la Asociación

Española de Entomología, p. 125.

Porcel, M., Cotes, B., Castro, J., Campos, M., 2017. The effect of resident vegetation

cover on abundance and diversity of green lacewings (Neuroptera: Chrysopidae)

on olive trees. J. Pest Sci. 90, 195-206.

R Development Core Team, 2018. R: a Language and Environment for Statistical

Computing. Available at Vienna: R Foundation for Statistical Computing. http://

www.rproject.org/.

Ramos, P., Campos, M., Ramos, J.M., 1987. Evolución del ataque de *Prays oleae* Bern,

al fruto del olivo. I. Estudio de parámetros y sus relaciones. Bol. San. Veg.

Plagas 13, 129-142.

Ramos, P., Ramos J.M., 1990. Veinte años de observaciones sobre la depredación

oófaga en *Prays oleae* Bern. Granada (España), 1970-1989. Bol. San. Veg.

Plagas 16, 119-127.

Ramos, P., Campos, M., Ramos, J.M., 1998. Long-term study on the evaluation of yield

and economic losses caused by *Prays oleae* Bern. in the olive crop of Granada

(southern Spain). Crop Prot. 17, 645-647.

Red de alerta e Información fitosanitaria de Andalucía (RAIF), 2018. http://www.junta-

deandalucia.es/agriculturapescaydesarrollorural/raif/fichas-fitopatologicas.

Redolfi, I., Tinaut, A., Pascual, F., Campos, M., 1999. Qualitative aspects of

myrmecocenosis (Hym., Formicidae) in olive orchards with different

agricultural management in Spain. J. Appl. Entomol. 123, 621-627.

Rodríguez, E., Lozano, C., Campos, M., 2005. Detection of predation on *Euzophera*

pingüiis (Lepidoptera: Pyralidae) using an enzyme-linked immunosorbent assay.

Eur. J. Entomol. 102: 793-796.

Rodríguez, E., González, B., Campos, M., 2009. Effects of cereal cover crops on the

main insect pests in Spanish olive groves. J. Pest Sci. 82, 179-185.

Rodríguez, E., González, B., Campos, M., 2012. Natural enemies associated with cereal

cover crops in olive groves. Bull. Insectol. 65, 43-49.

Rusch, A., Valantin-Morison, M., Sarthou, J.P., Roger-Estrade, J., 2010. Biological

control of insect pests in agroecosystems: effects of crop management, farming

systems, and seminatural habitats at the landscape scale: a review. Adv. Agron.

109, 219-259.

Santos, S.A., Cabanas, J.E., Pereira, J.A., 2007. Abundance and diversity of soil

arthropods in olive grove ecosystem (Portugal): Effect of pitfall trap type. Eur. J.

Soil Biol. 43, 77-83.

Segoli, M., Harari, A.R., Rosenheim, J.A., Bouskila, A., Keasar, T., 2010. The

evolution of polyembryony in parasitoid wasps. J Evol. Biol. 23, 1807-1819.

Seifert, B., D'Eustacchio, D., Kaufmann, B., Centorame, M., Lorite, P., Modica, M.V.,

2017. Four species within the supercolonial ants of the *Tapinoma nigerrimum*

complex revealed by integrative taxonomy (Hymenoptera: Formicidae).

Myrmecol. News 24, 123-144.

Straub, C.S., Finke, D.L., Snyder, W.E., 2008. Are the conservation of natural enemy

biodiversity and biological control compatible goals? Biol. Control 45, 225-237.

Torres, L., 2006. A fauna auxiliar do olival ea sua conservação. Joao Azevedo. 92 p.

Torres-Miralles, M., Grammatikopoulou, I., Rescia, A.J., 2017. Employing contingent

and inferred valuation methods to evaluate the conservation of olive groves and

associated ecosystem services in Andalusia (Spain). Ecosys. Serv. 26, 258-269.

Trdan, S., Andjus, L., Raspudić, E., Kač, M., 2005. Distribution of *Aeolothrips*

intermedius Bagnall (Thysanoptera: Aeolothripidae) and its potential prey

Thysanoptera species on different cultivated host plants. J. Pest Sci. 78, 217-226.

Tscharntke, T., et al., 2012. Landscape moderation of bio- diversity patterns and

processes – eight hypotheses. Biol. Rev. 87, 661-685.

Tscharntke, T., et al., 2016. When natural habitat fails to enhance biological pest control

– Five hypotheses. Biol. Control 204, 449-458.

Tzanakakis, M., 2006. Insects and Mites Feeding in the Olive. Distribution, Importance,

Habits, Seasonal Development and Dominancy. Brill Academic Publisher,

Leiden.

Uetz, G.W., Halaj, J., Cady, A.B., 1999. Guild structure of spider in major crops. J.

Arachnol. 27, 270-280.

Villa, M., Santos, S.A., Mexia, A., Bento, A., Pereira, J.A., 2016a. Ground cover

management affects parasitism of *Prays oleae* (Bernard). Biol. Control 96, 72-

77.

Villa, M., Santos, S.A., Benhadi-Marín, J., Mexia, A., Bento, A., Pereira, J.A., 2016b.

Life-history parameters of *Chrysoperla carnea* sl fed on spontaneous plant

species and insect honeydews: importance for conservation biological control.

BioControl 61, 533-543.

Villa, M., Marrão, R., Mexia, A., Bento, A., Pereira, J.A., 2016c. Are wild flowers and

insect honeydews potential food resources for adults of the olive moth, *Prays

oleae*?. J. Pest Sci. 90, 185-194.

Villa, M., Santos, S.A., Sousa, J.P., Ferreira, A., da Silva, P.M., Patanita, I., Ortega, M.,

Pascual, S., Pereira, J.A., 2020. Landscape composition and configuration affect

the abundance of the olive moth (*Prays oleae*, Bernard) in olive groves. Agric.

Ecosyst. Environ. 294, 106854.

Vossen, P.M., 2007. Organic olive production manual (Vol. 3505). UCANR

Publications. 105 p.

Wan, N.F., et al., 2018. Increasing plant diversity with border crops reduces insecticide

use and increases crop yield in urban agriculture. eLife 7, e35103.

Wilby, A., Villareal, S.C., Lan, L.P., Heong, K.L., Thomas, M.B., 2005. Functional

benefits on predator species diversity depend on prey identity. Ecol. Entomol.

30, 497-501.

Table 1. Relative abundance (RA), acronyms, presence in orchards: tilled (T) and

mowed (M), and trophic guilds of the families of natural enemies ($n = 36$) identified in

organic olive orchards. Numbers represent presence (1) and absence (0).

Guild	Order	Family	Acronym	T	M	RA (%)
Omnivores	Dermaptera	Forficulidae	Fo	1	1	1,51
	Hymenoptera	Formicidae	For	1	1	98,49
Parasitoids	Hymenoptera	Braconidae	Bra	1	1	8,49
		Elasmidae	El	1	1	12,26
		Encirtidae	En	1	1	5,66
		Eulophidae	Eu	1	1	3,77
		Ichneumonidae	Ich	1	0	0,94
		Pteromalidae	Pt	1	1	26,42
		Scelionidae	Sc	1	1	42,45
Predators	Araneae	Araneidae	Ar	1	1	1,48
		Corinidae	Co	1	1	0,74
		Linyphiidae	Li	1	1	0,60
		Liochranidae	Lio	0	1	0,07
		Oxyopidae	Ox	1	1	2,41
		Philodromidae	Ph	1	1	2,35
		Salticidae	Sa	1	1	6,61
		Theridiidae	The	1	1	1,17
		Thomisidae	Tho	1	1	19,95
		Uloboridae	Ul	0	1	0,03
	Coleoptera	Cantharidae	Ca	1	1	0,74
		Carabidae	Car	1	1	0,07
		Coccinelidae	Coc	1	1	6,17
		Malachiidae	Ma	1	0	0,07
		Staphylinidae	St	1	1	0,10
	Diptera	Asilidae	As	1	1	0,13
		Empididae	Em	1	1	0,64
	Hemiptera	Anthocoridae	An	1	1	11,64
		Lygaeidae	Ly	1	1	0,37
		Miridae	Mi	1	1	31,05
		Nabidae	Na	0	1	0,07
		Reduviidae	Re	1	1	1,11
	Mantodea	Mantidae	Man	1	1	0,47
	Neuroptera	Chrysopidae	Chr	1	1	10,43
		Coniopterygidae	Con	1	1	0,54

Raphidioptera	Raphidiidae	Ra	1	1	0,13
Thysanoptera	Aeolothripidae	Ae	0	1	0,87

671   Table 2. Fixed effects of the fitted model (GLMM) explaining egg predation: type of

672   management and natural enemy abundance.

Variable	df	Wald χ^2	p
Management (tilled or mowed)	1	4.524	0.033
Natural enemy abundance	1	5.841	0.015

Figure 1

Fig. 1. Panoramic view of tilled (left) and mowed (right) organic olive orchards (a).

Oviposition site of *Prays oleae* on an olive (b). Appearance of laid eggs of *P. oleae* on

an olive: live egg (c), hatched egg (d), and predated egg (e). Site and eggs are indicated

by triangles.

Figure 2

Fig. 2. Natural enemy abundance classified by management: tilled (white bars) and

mowed (grey bars). Families are grouped by orders and trophic information. An asterisk

indicates that a family of natural enemy showed significantly differences between

managements. Black points indicate the families that were present in only one of the

managements.

Figure 3

Fig. 3. Relation amongst variables according to GLMM analysis: egg predation, the

natural enemy (NE) abundance, and management.

Figure 4

Figure 4: Non metric multi-dimensional scaling (NMDS) of the total abundance of

natural enemies. Proximity amongst families of natural enemies within the ordination

plot indicates that their abundances are positively related. Lines (smooth surfaces)

represent different levels, in the form of a gradient, of predated egg counts (A) and adult

abundance (B) of *P. oleae*, according to generalised additive models. See family

acronyms in Table 1.