In vivo identification and validation of novel potential predictors for human

cardiovascular diseases

Omar T. Hammouda[1,2], Meng Yue Wu[1], Verena Kaul[1], Jakob Gierten[1,2,3], Thomas

Thumberger[1] and Joachim Wittbrodt[1,*]

[1]Centre for Organismal Studies Heidelberg, Heidelberg University, Im Neuenheimer Feld 230,

69120 Heidelberg, Germany

[2]Heidelberg Biosciences International Graduate School, Heidelberg University, Im

Neuenheimer Feld 501, 69120 Heidelberg, Germany

[3]Department of Pediatric Cardiology and Congenital Heart Diseases, University Hospital

Heidelberg, Im Neuenheimer Feld 430, 69120 Heidelberg, Germany

[*]Corresponding Author

Running title (40 char): Validating understudied GWAS heart genes

Final character count (with spaces): 21,783

Hammouda et al., 2021 Validating understudied GWAS heart genes

Abstract

Genetics crucially contributes to cardiovascular diseases (CVDs), the global leading cause of death. Since the majority of CVDs can be prevented by early intervention there is a high demand for predictive markers. While genome wide association studies (GWAS) correlate genes and CVDs after diagnosis and provide a valuable resource for such markers, preferentially those with previously known or suspected function are addressed further. To tackle the unaddressed blind spot of understudied genes, we particularly focused on the validation of heart GWAS candidates with little or no apparent connection to cardiac function. Building on the high conservation of basic heart function and underlying genetics from fish to human we combined CRISPR/Cas9 genome editing of the orthologs of human GWAS candidates in isogenic medaka with automated high-throughput heart rate analysis. Our functional analyses of understudied human candidates uncovered a prominent fraction of heart rate associated genes from adult human patients displaying a heart rate effect in embryonic medaka already in the injected generation. Following this pipeline, we identified 16 GWAS candidates with potential diagnostic and predictive power for human CVDs.

Keywords: CRISPR / gene validation / GWAS / heart rate / high throughput analysis

Hammouda et al., 2021 Validating understudied GWAS heart genes

Introduction

Genetics crucially contributes to the development and progression of cardiovascular diseases (CVDs), the global leading cause of death (Kathiresan & Srivastava 2012; Cambien & Tiret 2007). Elevated resting heart rate in humans has been widely considered as a potential, modifiable risk factor of cardiovascular and all-cause mortality (Beere et al. 1984; DYER et al. 1980; MD et al. 2010; Gillum et al. 1991). Since the majority of CVDs can be prevented by early intervention (McGill et al. 2008) there is a high demand for diagnostic and predictive CVD markers. Genome wide association studies (GWAS) on human patients correlate genes and CVDs after diagnosis and provide a valuable resource for those putative markers (Eicher et al. 2015). However, genes with pre-assigned cardiac functions are usually more likely to be addressed further, while uncharacterized genes or those with no pre-existing evidence to the heart are often neglected. This is likely due to the lack of experimental pipelines for the rapid and robust validation of such markers with implications for heart function. Recently, we demonstrated the power of targeted genome editing in the small animal model system medaka (*Oryzias latipes*) to validate trabeculation-associated genes (Meyer et al. 2020). The ease of manipulation combined with robust acquisition and analysis pipelines highlight the power of using fish embryos in high-throughput applications (Gierten et al. 2020; Shankaran et al. 2018; Lessman 2011; Oxendine et al. 2006). Embryos of fish model systems undergo extrauterine development in a transparent egg. This allows to monitor heart development and heart rate non-invasively in live undisturbed embryos for an extended period of time. Heart development, function and physiology in fish, though simpler, is comparable to mammals (Nemtsas et al. 2010; Yonekura et al. 2018; Gut et al. 2017). Here we combined targeted genome editing via CRISPR/Cas9 (Stemmer et al. 2015; Thumberger et al. 2021) with automated high-throughput imaging and heart rate analysis in isogenic medaka embryos (Gierten et al. 2020) to enable functional analyses directly in the injected generation (F0). We tested the performance our assay with a positive control (*nkx2-5*), evaluated the random discovery rate and analyzed 40 heart associated genes identified from human GWAS. Our assay uncovered that 57% of candidates assigned to human heart rate in GWAS also affected heart rate in fish embryos. We

Hammouda et al., 2021 Validating understudied GWAS heart genes

have thus experimentally validated understudied human GWAS candidates, identifying 16

genes with potential diagnostic and predictive power for human CVDs.

Results

For the straight forward functional validation of GWAS candidates we aimed at combining

CRISPR/Cas9-mediated targeted gene inactivation in the injected generation of medaka

embryos with high content screening approaches to validate the impact of the loss-of-function

on the heart rate (Fig 1A). We first assessed the efficiency of the CRISPR/Cas9 system by

targeting the *green fluorescent protein* (*gfp*) gene in a transgenic medaka reporter line

expressing GFP and mCherry fluorescent proteins exclusively in the heart via the *cardiac*

myosin light chain 2 (*cmlc2*) promoter. Injection of the *heiCas9* mRNA, i.e. a Cas9 equipped

with an early active nuclear localization signal (Thumberger et al. 2021), together with a guide

RNA targeting *gfp* into medaka embryos at the 1-cell stage resulted in a complete loss of GFP

expression in the heart (n=8/8) (Fig 1B and Fig S1). Only when injected into a single blastomere

of the four-cell stage or later, a mosaic pattern was observable. This demonstrates the high

efficiency of our approach already in the injected generation.

For our functional validation assay, we used the cardiac-specific homeobox-containing

transcription factor NKX2-5 as a positive control. In human patients, a single amino acid

mutation in the homeodomain (R141C) was previously associated with atrial septal defect

(ASD) and shown to cause delayed heart morphogenesis in adult mice (Zakariyah et al. 2017).

To test our high-throughput imaging and heart rate analysis pipeline for functional *in vivo* gene

validation in the injected generation, we targeted the region orthologous to R141C in medaka

embryos using the CRISPR/Cas9 system (Stemmer et al. 2015, Thumberger et al. 2021). As a

negative control, we targeted the *oculocutaneous albinism* 2 locus (*oca2*) (Fukamachi et al.

2004) as a heart-unrelated pigmentation gene. The apparent loss of the eye pigmentation

phenotype (and low degree of mosaicism) (Lischik et al. 2019; Hammouda et al. 2019)

underscores the bi-allelic knock-out efficacy of our system. Injections into wild-type medaka

embryos were performed at the 1-cell stage, and hereafter resulting embryos are referred to as

crispants. To address the impact on the heart rate we raised the medaka crispants until cardiac

Figure 1. Functional gene validation pipeline confirms heart rate phenotype in *nkx2-5* crispants
A Schematic overview of our functional gene validation pipeline: position of human coding SNP mapped to medaka orthologous gene to define region of interest for CRISPR/Cas9 gene targeting (double strand break; DSB). 96-well plate layout of crispant embryos (Target Gene 1 and 2) separated by *GFP* mRNA mock-injected siblings. Embryos are subjected to high-throughput imaging followed by automated heart detection (blue area) and heart rate quantification (graphical output; *HeartBeat* software (Gierten et al. 2020). **B** Confocal images of GFP expression in mock-injected and *gfp* crispant embryo hearts of the *cmlc2::eGFP cmlc2::H2A-mCherry* reporter line (7 dpf). Embryos were injected either at the 1-cell or 4-cell stage. Note: complete loss of GFP expression when injected at the 1-cell stage (n = 8/8), while mosaic expression when injected at 4-cell stage (n = 4/4). Genotyping of *gfp* crispants display

Hammouda et al., 2021 Validating understudied GWAS heart genes

the genetic mosaicism resulting from Cas9-based targeting. Scale bars: 50 µm. For full image refer to Fig S1B. **C** Comparison of the atrium (A, dotted red line) and ventricle (V, dotted yellow line) in *GFP*-injected (Mock) and *nkx2-5* and *oca2* crispant embryos (9 dpf). Note: *nkx2-5* crispant shows dilated heart chambers while mock injected and *oca2* crispant embryos are indistinguishable. Loss of eye pigmentation in *oca2* crispants reflects high efficiency of knock-out rate in the injected generation. **D** Heart rate measurements (beats per minute, bpm) of *GFP*-injected (Mock; dark grey), *nkx2-5* and *oca2* crispant embryos (4 dpf) at 21 and 28°C, before (left) and after (right) exclusion of severely affected embryos (developmental focusing) reveal elevation of mean heart rates in *nkx2-5* targeted embryos, significant at 21°C (red). Significance was determined by two-tailed Student's t-test; *$p < 0.05$, **$p < 0.01$, ns (not significant; light grey). For biological replicates see Source Data Fig 1D.

function was fully developed and the heart rate had reached a plateau at 4 days post fertilization

(Gierten et al. 2020) (4 dpf; developmental stage ~31-32 (Iwamatsu 2004)).

To assess changes in mean heart rate with statistical significance, we took advantage of a 96-

well plate format, and imaged multiple biological replicates of crispants (3 rows; n=36 per

condition) as well as of *GFP mRNA* mock-injected siblings as internal plate control (2 rows;

n=24) (Fig S2A). To assess heart function under different environmental conditions, embryos

were imaged at two different temperatures (21 and 28°C, respectively). Heart rates of all

embryos were quantitatively determined from the imaging data using the *HeartBeat* software

(Gierten et al. 2020), and randomly selected embryos were genotyped to correlate

CRISPR/Cas9 targeting (Hammouda et al. 2019).

While mock injected embryos did not show phenotypes, crispants of the positive control *nkx2-*

5 displayed a variety thereof. These ranged from global severe developmental delays to local

cardiac malformations morphologically resembling the phenotypes previously observed in

zebrafish *nkx2-5* mutants such as enlarged heart chambers (Fig 1C) (Targoff et al. 2013).

Notably, in the negative control (*oca2* crispants) neither cardiac nor developmental phenotypes

were observed (Fig 1C), indicating that genome targeting of *oca2* as well as injection and

handling of the embryos did not impact on heart and general development per se. Quantitative

comparison of cardiac function revealed an overall elevation in the mean heart rate of *nkx2-5*

crispants (21°C 99.7 bpm, 28°C 166 bpm) compared to mock control siblings (21°C 96.1 bpm,

28°C 164 bpm) with a significant (p = 0.0074) difference at 21°C (Fig 1D; left panel). Notably,

independent experimental replicates targeting the same *nkx2-5* exon with two different sgRNAs

robustly yielded a significant heart rate phenotype at 21°C (Fig S2B-C). In contrast, the mean

heart rate in *oca2* crispants was indifferent from mock control at either temperature (21°C 96.4

bpm, 28°C 165 bpm), validating *oca2* as bona fide negative control.

To avoid severe developmental delays in *nkx2-5* crispants to potentially skew heart rate
comparisons, we further applied a developmental focusing filter. Only embryos having
developed beyond stage 28 (Iwamatsu 2004), at which cardiac function was previously shown
to have reached a functional plateau (Gierten et al. 2020), were chosen for statistical analysis.
Developmental focusing, only excluded three embryos from the *nkx2-5* group which did not
impact on the results (Fig 1D; right panel). These results underline the robustness of our
pipeline and demonstrate its sensitivity to detect mild heart rate phenotypes reflecting cardiac
function already at embryonic stages of medaka development.

Next, we determined the baseline probability of heart rate phenotypes by targeting a set of
randomly selected genes with CRISPR/Cas9. From a total of 23622 annotated medaka coding
genes in Ensembl (Yates et al. 2020), we used a random number generator to select 10 genes
(Table 1). For each gene, a random exon was chosen for targeting via CRISPR/Cas9. Heart
rates of 36 crispants were assessed per locus. To control for potential heart rate fluctuations in
embryos within and across different experiments (i.e. 96-well plates), we included mock-
injected siblings as internal plate control. Heart rates of target gene crispants and control
siblings were scored and the means were compared before and after developmental focusing.

Table 1. List of randomly selected genes.
Medaka Ensembl gene names and codes, as well as orthologous human genes as annotated in the 95[th]
Ensembl release.
144

Ensembl ID	Medaka Gene	Orthologous Human Gene
ENSORLG00000006335	novel gene - *cdc42*	na (*CDC42* by name)
ENSORLG00000000979	novel gene - *ogdh*	*OGDH*
ENSORLG00000005268	*duox*	*DUOX1*
ENSORLG00000007310	*git2*	*GIT2*
ENSORLG00000003492	*mus81*	*MUS81*
ENSORLG00000020766	*or124-2*	na
ENSORLG00000007400	*plekha8*	*PLEKHA8*
ENSORLG00000022757	*ttl*	*TTL*
ENSORLG00000005922	*cabp4*	*CABP2*
ENSORLG00000023106	novel gene - *eml6*	*EML6*

Hammouda et al., 2021 Validating understudied GWAS heart genes

Comparative heart rate analysis revealed a heart rate phenotype in two out of the ten randomly selected genes at both temperatures measured (Fig 2, and Fig S3). Remarkably, both genes of the random set, the *oxoglutarate dehydrogenase* (*ogdh*) and the *cell division control protein 42 homolog* (*cdc42*) have been previously associated with heart phenotypes in human GWAS or were shown to play a role in heart function, respectively (Thanassoulis et al. 2013; Qian et al. 2011; Li et al. 2017). These results confirm the reliability of our assay to identify genes of a given set that affect cardiac function. Of note, repeated rounds of random gene selection revealed a similar mean baseline of 10-20% of selected genes being related to the heart in GWAS or previous experimental reports.

Figure 2. Baseline probability of heart rate phenotype assessed via *in vivo* targeting of randomly selected genes
A Heart rate measurements (beats per minute, bpm) of *GFP*-injected (Mock) and corresponding sibling crispant embryos (4 dpf) at 21 and 28°C after developmental focusing. Different experimental plates are represented by breaks on the x-axis. Significant differences in mean heart rates were determined between each crispant embryo group and its corresponding sibling control group by two-tailed Student's t-test; *$p < 0.05$, ***$p < 0.001$, ns (not significant). Crispants showing significant heart rate phenotype (red), *GFP*-injected controls (Mock; dark grey), crispants showing no significant heart rate phenotype (light grey). **B** Heatmap quantitative representation of the data shown in (A); for each measured temperature, the percent change in mean heart rate (HR % Change) between crispants and their corresponding control sibling, flanked by the statistical significance (*p*-value) of the observed change calculated by two-tailed Student's t-test on the full distribution in (A). Genes showing significant heart rate phenotypes are indicated in **bold**. For biological replicates see Source Data Fig S3.

We next applied our pipeline to interrogate a larger, targeted selection of genes associated with

cardiovascular diseases in human GWAS. We used GRASP (Eicher et al. 2015), the genome-

wide repository of associations between single nucleotide polymorphisms (SNPs) and

phenotypes, to compile a list of 40 candidate genes from human GWAS with a coding

association to heart phenotypes (hGWAS genes; Table 2). We focused on genes with no prior

experimental link to heart function, while including few known heart genes as additional

positive controls. To address the specificity of our approach, the selected candidate genes were

categorized according to their association into general heart related (n=17) or related

specifically to heart rate (n= 23; Table2).

Heart rates of candidate gene crispants and control injected siblings were scored and compared

before (Fig S5) and after developmental focusing (Fig 3A and Figure S4). Across the hGWAS

set of 40 genes, comparative heart rate analysis showed statistically significant heart rate

phenotypes in a total of 16 genes (Fig 3B). The five positive controls, known to play key roles

in heart functions such as cardiac contraction (*TTN* (Itoh-Satoh et al. 2002) and *NACA* (Park et

al. 2010)) and heart rate regulation (*CASQ2* (Faggioni & Knollmann 2012), *KCNH2* (Gianulis

& Trudeau 2011) and *SCN5A* (Zaklyazminskaya & Dzemeshkevich 2016; Huttner et al. 2013))

clearly responded in the assay. Beyond known cardiac genes, we revealed new genes linked to

various biological functions (*CCDC141*, *GIGYF1*, *HOMEZ*, *MYRF*, *SMG6*, *CMYA5*, *CNOT1*,

SLC17A3, *TRAPPC12*, *SSPO* and *PADI4*) which up to now, had little to no experimental

evidence in cardiac function (Rathjens et al. 2020; Benson et al. 2017; Yamaguchi et al. 2018;

Elmen et al. 2020).

When analyzing the candidates according to their GWAS association ("heart rate" and "non-

heart rate" phenotypes), we observed a strong positive correlation between the respective

phenotypes observed in medaka crispants already at the embryonic stages used and the

associated phenotype in adult human GWAS. The proportion of heart rate-associated genes in

hGWAS that yield a heart rate phenotype in medaka embryos (13/23) was elevated compared

to the proportion of non-heart rate-associated genes yielding a heart rate phenotype (3/17). Even

when considering the entire group, we observed a higher proportion of genes with an effect on

heart rate in our targeted hGWAS gene set (16/40) compared to our randomly selected gene set

(2/10) (Fig 3C). Taken together, phenotypes in early medaka embryos likely reflect risk factors

Table 2. List of candidate genes extracted from Human GWAS using GRASP 2.0 Database
Human genes are categorized according to their association into "heart rate" (**bold**) and "non-heart rate" (non-bold) related phenotypes in human GWAS.
189

Human Gene	Coding SNP ID	Associated heart phenotype	Association Reference	Medaka orthologue gene (Ensembl release)	Ensembl Gene Code
ATP8B4	rs2452524	Pulse rate	(Hiura et al. 2010)	*atp8b4* (98)	ENSORLG00000005106
CASQ2	rs4074536	QRS interval	(Sotoodehnia et al. 2010)	*casq2* (89)	ENSORLG00000017885
CCDC141	rs17362588	**Heart rate**	(Hoed et al. 2013)	**na (89) (TBLASTN=*ccdc141*)**	**TBLASTN = ENSORLG00000030409**
CEP85L	rs3734381	QRS interval	(Sotoodehnia et al. 2010)	*cep85l* (91)	ENSORLG00000015455
CMYA5	rs10942901	**Heart rate**	(Hoed et al. 2013)	**cmya5 (91)**	**ENSORLG00000008983**
COL9A1	rs592121	Pulse rate	(Hiura et al. 2010)	*col9a1b* (98)	ENSORLG00000010431
GIGYF1	rs221794	**Heart rate**	(Hoed et al. 2013)	**gigyf1 (89)**	**ENSORLG00000003655**
GRID2	rs1385405	Pulse rate	(Hiura et al. 2010)	*grid2* (98)	ENSORLG00000024663
HOMEZ	rs1055061	Sick sinus syndrome	(Holm et al. 2011)	*homeza* (89)	ENSORLG00000012220
KCNH2	rs1805123	QT interval	(Pfeufer et al. 2009)	*kcnh2* (98)	ENSORLG00000004137
MINAR1	rs2297773	Pulse rate	(Hiura et al. 2010)	*minar1* (98)	ENSORLG00000016707

Hammouda et al., 2021 Validating understudied GWAS heart genes

Human Gene	Coding SNP ID	Associated heart phenotype	Association Reference	Medaka orthologue gene (Ensembl release)	Ensembl Gene Code
MYRF	rs174535	RR interval	(Eijgelsheim et al. 2010)	*myrf* (91)	ENSORLG00000006459
NACA	rs2926743	Heart rate	(Hoed et al. 2013)	*naca* (98)	ENSORLG00000012246
OR5AU1	rs4982419	Pulse rate	(Hiura et al. 2010)	na (98) (TBLASTN=no name)	TBLASTN = ENSORLG00000024679
PADI4	rs2240335	Pulse rate	(Hiura et al. 2010)	na (98) (TBLASTN=*padi2*)	TBLASTN = ENSORLG00000007539
PPP1R9A	rs854524	Pulse rate	(Hiura et al. 2010)	*ppp1r9a* (98)	ENSORLG00000004418
RNF207	rs846111	QT interval	(Newton-Cheh et al. 2009)	*rnf207b* (91)	ENSORLG00000017207
SCN5A	rs1805126	QRS interval	(Ritchie et al. 2013)	na (89) (TBLASTN=*scn4ab*)	TBLASTN = ENSORLG00000003273
SSPO	rs10261977	Pulse rate	(Hiura et al. 2010)	*sspo* (98)	ENSORLG00000004121
TRAPPC12	rs6767	Pulse rate	(Hiura et al. 2010)	*trappc12* (98)	ENSORLG00000017859
TTN	rs12476289	QT interval	(Marroni et al. 2009)	*ttn.2* (91)	ENSORLG00000018144
UFSP1	rs12666989	RR interval	(Eijgelsheim et al. 2010)	na (89) (TBLASTN=*ufsp1*)	TBLASTN = ENSORLG00000022928
XYLB	rs17118	PR interval	(Smith et al. 2009)	*xylb* (91)	ENSORLG00000003755
ABCB1	rs1128503	Drug response CVD	(Paré et al. 2013)	*abcb4* (91)	ENSORLG00000009269

Hammouda et al., 2021 Validating understudied GWAS heart genes

Human Gene	Coding SNP ID	Associated heart phenotype	Association Reference	Medaka orthologue gene (Ensembl release)	Ensembl Gene Code
BAG3	rs3858340	Sporadic dilated cardiomyopathy	(Villard et al. 2011)	*bag3* (89)	ENSORLG00000013813
CLCNKA	rs1805152	Sporadic dilated cardiomyopathy	(Villard et al. 2011)	*clcnk* (89)	ENSORLG00000018693
CNOT1	rs11866002	Aortic valve calcium	(Thanassoulis et al. 2013)	*cnot1* (91)	ENSORLG00000013734
EDN1	rs150035515	Aortic valve calcium	(Matsa et al. 2014)	*edn1* (89)	ENSORLG00000009276
HCN4	rs529004	Aortic valve calcium	(Thanassoulis et al. 2013)	*hcn4* (91)	ENSORLG00000013180
MAML3	rs11729794	Congenital heart malformations	(Hu et al. 2013)	na (91) (TBLASTN=*maml3*)	TBLASTN = NCBI Ref Seq: XM_023954746.1
NUBP2	rs344359	LV systolic dysfunction	(Vasan et al. 2009)	*nubp2* (91)	ENSORLG00000007228
PIEZO1	rs2290902	Bicuspid aortic valve	(Wooten et al. 2010)	*piezo1* (91)	ENSORLG00000000402
PLG	rs13231	Aortic valve calcium	(Thanassoulis et al. 2013)	*plg* (91)	ENSORLG00000020532
RGS3	rs12341266	Hypertrophic cardiomyopathy	(Wooten et al. 2013)	*rgs3a* (91)	ENSORLG00000006823
SCMH1	rs10489520	Ischemic stroke	(Ikram et al. 2009)	*scmh1* (91)	ENSORLG00000014207
SH2B3	rs3184504	Tetrology of fallot	(Cordell et al. 2013)	*sh2b3* (91)	ENSORLG00000003569
SLC17A3	rs942379	Bicuspid aortic valve	(Wooten et al. 2010)	*si:ch1073-513e17.1* (91)	ENSORLG00000007671

Hammouda et al., 2021 Validating understudied GWAS heart genes

Human Gene	Coding SNP ID	Associated heart phenotype	Association Reference	Medaka orthologue gene (Ensembl release)	Ensembl Gene Code
SMG6	rs216193	Aortic root size	(Vasan et al. 2009)	*smg6* (91)	ENSORLG00000003317
VEPH1	rs1378796	Sporadic dilated cardiomyopathy	(Villard et al. 2011)	*veph1* (89)	ENSORLG00000012452
ZFHX3	rs2228200	Aortic valve calcium	(Thanassoulis et al. 2013)	*zfhx3* (91)	ENSORLG00000007874

Hammouda et al., 2021 Validating understudied GWAS heart genes

Figure 3 Targeted human heart-GWAS validations reveal new genes affecting heart rate
A Heatmap quantitative representation of the comparative heart rate analysis between each crispant embryo group and its corresponding control sibling group after developmental focusing (also see plots in Fig S4); for each measured temperature, the percent change in mean heart rate (HR % Change) between crispants and their corresponding control sibling, flanked by the statistical significance (*p*-value) of the observed change calculated by two-tailed Student's t-test on the full distribution (Fig S4). Genes showing significant heart rate phenotypes are indicated in **bold**. For biological replicates see Source Data Fig S4. **B** Venn diagram summarizing the genes with significantly different heart rate (HR) phenotypes only at 21°C, only at 28°C or at both temperatures (dark grey). **C** Stacked plots representing percentage of genes showing a significant heart rate phenotype (dark grey) in each group. Number of genes for each group is denoted (n). hGWAS corresponds to the selection of genes associated to heart phenotypes in human GWAS.

193 in human adults, thus we uncovered functionally relevant heart rate phenotypes in previously

194 uncharacterized genes.

Hammouda et al., 2021 Validating understudied GWAS heart genes

In addition to the observed heart rate phenotypes in *trappc12* crispants, we also uncovered
morphological heart phenotypes such as heart looping defects. Where in wild-type medaka,
heart looping usually starts at around stage 27, when the atrium shifts to the right and lies
adjacent to the ventricle (Iwamatsu 2004), *trappc12* crispants revealed strong heart looping
phenotypes (12/46) not observed in *oca2* crispants (0/22) or mock-injected embryos (0/26) (Fig
4A).
In crispants of *scn4ab,* heart rate analysis interestingly uncovered a bimodal distribution, with
a population displaying roughly half the average heart rate at both recorded temperatures (Fig
4B). Visual inspection of the *scn4ab* crispant embryos revealed an arrhythmic heart anomaly
previously reported in zebrafish *scn5a* mutants (Huttner et al. 2013), i.e. known as an atrio-
ventricular block (AV-block), characterized by a delay or disruption of impulse transmission
from the atrium to the ventricle. Scoring the beat frequency of both heart chambers separately
in individual embryos exposed the impaired rhythm of atrial to ventricular contractions, which
resulted in a delay or even skipping of ventricular beats in the *scn4ab* crispants but not in control
siblings (Fig 4C). *scn4ab* crispants displayed various severities of the AV-block from mild
(regular heart beats with occasional beat skipping), to moderate (consistent 2:1 atrial to
ventricular contraction; Movie S1), to severe (3:1 or more; Movie S2). Still heavily affected
scn4ab crispants survived until hatching. Impressively, the prevalence of the arrhythmia
phenotype in *scn4ab* crispants was markedly high, exceeding 90% of the injected embryos,
reflecting the high efficiency of the *heiCas9* and the high penetrance of the mutations
introduced. Notably, the arrhythmia phenotype of *scn4ab* mutants bred to homozygosity did
not differ from the phenotype observed in the injected generation F0 (Movie S1), verifying the
specificity of the phenotype. These results further underscore the efficacy and reliability of
medaka F0 crispant analysis as a rapid validation tool to identify genes with a functional link
to human cardiac diseases.
220
221 **Discussion**
222
223 Most cardiovascular diseases can be prevented if diagnosed and treated early. Previous studies
224 have shown the importance of the resting heart rate as a vital risk factor both in terms of
225 prediction and prevention of CVDs (DYER et al. 1980; MD et al. 2010; Eppinga et al. 2016).

Figure 4. Heart looping defects and AV-block type arrhythmia in medaka crispant embryos
A Confocal images of hearts of mock-injected, *oca2* and *trappc12* crispants of the *cmlc2::eGFP cmlc2::H2A-mCherry* reporter line (7 dpf); note the heart looping defect observed in *trappc12* crispants. Scale bars: 100 µm (First panel on left) and 50 µm (blow-up images). **B** Heart rate measurements (beats per minute, bpm) of *GFP*-injected (Mock; dark grey; n = 22) and *scn4ab* crispant (red; n = 32) embryos (4 dpf) at 21 and 28°C; note the bimodal distribution in *scn4ab* crispants. **C** Paired plots showing heart rate scores for each chamber separately (atrium in blue; ventricle in red) of individual embryos from B at both temperatures.

Hammouda et al., 2021 Validating understudied GWAS heart genes

An increase of 5 beats per minute correlates with a 20% increase in risk of mortality (Eppinga
et al. 2016), and reducing the resting heart rate has proven to improve the clinical outcomes of
various CVDs (Beere et al. 1984; MD et al. 2010).

Human GWAS have been performed in search of genetic determinants of CVDs, and although
a wide array of candidate genes with various functions are being associated to heart phenotypes
in human GWAS, further focus is usually turned to those few genes with pre-existing indication
of cardiac function. Other associated genes with an unknown function or without a pre-existing
functional link to the heart are often neglected and not pursued further, pushing all those into a
blind spot, resulting in a negative loop of discovery. Thus, it is important to address the role of
such genes in heart function through experimental validation in model organisms, in pursuit of
novel markers for CVD diagnosis. We address this blind spot of discovery by applying a high-
throughput heart rate imaging and analysis pipeline coupled to a reverse genetic validation
approach via CRISPR/Cas9 mediated mutagenesis in a genetically suited vertebrate model (Fig
1A).

F0 mutagenesis screens are becoming more and more popular (Teboul 2017; Shankaran et al.
2018; Wu et al. 2018; Heyde et al. 2020), largely due to the improvements in CRISPR/Cas9
gene targeting efficiency by modification of the enzyme or promoting nuclear localization
(Thumberger et al. 2021). Overall, gene targeting in medaka using CRISPR/Cas9 has proven
to be highly efficient, as shown by the prominent loss of GFP expression in *gfp* crispants when
injected at the 1-cell stage (Fig 1B and Fig S1) and the prominent bi-allelic mutagenesis as
apparent by the loss of eye-pigmentation in the *oca2* crispants (Fig 1C and Fig S2A). A
similarly high penetrance was also observed in the *scn4ab* crispants, where we detected and
quantified severe arrhythmia phenotypes such as AV-block with our assay (Fig 4B-C, Movies
S1 and S2). A 90 % prevalence of the arrhythmia phenotype and an absence of global
phenotypes further reflect the specificity of this phenotype.

A subset of the target genes in our assay (e.g. *nkx2-5, smg6, naca, ttn.2, abcb4*) however,
yielded a rather broad range of global developmental phenotypes, potentially reflecting their
important roles in embryonic development. To address this un-avoidable outcome when
tackling genes with broader function (e.g. transcription factors or essential genes), we applied
a developmental focusing filter in the analysis phase. Doing so, we avoid a biased assessment
of the heart rate by ensuring the comparability of the crispant embryos on a global

developmental scale, which in turn allows emphasizing cardiac-specific effects. Interestingly
however, developmental focusing, although deemed important, in only few cases significantly
altered the outcome of the analysis (Fig. S5). This reflects the robustness of the assay and the
homogeneity of CRISPR/Cas9-induced phenotypes in the isogenic background of the medaka
line used.

In a set of ten randomly chosen genes we observed a baseline occurrence of heart-affecting
genes of about 20 % (Fig 2, and Fig S3). Relevantly, both genes were implicated in heart
functions, further reflecting the reliability of our model and approach. For *cdc42*, there is *a*
priori evidence of its human orthologue in heart development as well as in regulating heart
function across species (Qian et al. 2011; Li et al. 2017). Surprisingly, we did not find any
associations (coding or non-coding) of *CDC42* to heart phenotypes in human GWAS according
to the GRASP database (Eicher et al. 2015). As for *ogdh*, no experimental evidence in cardiac
function has been previously reported, but a polymorphism located on one of its exons has been
associated to heart phenotypes in human GWAS (Thanassoulis et al. 2013). Except for *duox*,
which has been reported as having an indirect role in cardiac regeneration in zebrafish (Han et
al. 2014), none of the other randomly selected genes were, to our knowledge, ever connected
with cardiac function. In summary, from our random gene set, the only two affecting the heart
beat (*ogdh* and *cdc42*) are connected to cardiac function by prior evidence (Thanassoulis et al.
2013; Qian et al. 2011; Li et al. 2017).

All but one positive (*HCN4*) control among the hGWAS candidates resulted in a pronounced
heart rate phenotype in our assay, reflecting their role in cardiac contraction (*TTN* (Itoh-Satoh
et al. 2002) and *NACA* (Park et al. 2010)), cardiac conduction and heart rate regulation (*CASQ2*
(Faggioni & Knollmann 2012), *KCNH2* (Gianulis & Trudeau 2011) and *SCN5A*
(Zaklyazminskaya & Dzemeshkevich 2016; Huttner et al. 2013)). In case of the missing heart
rate phenotypes anticipated in *hcn4* crispants, we suspect compensation by its paralog *hcn4l*.

For eleven hGWAS candidate genes, our analysis provided the first experimental evidence
validating a cardiac function, and accordingly put these identified genes under the spotlight as
new targets for future in-depth characterization and as candidates for the prediction of heart
diseases prior to their onset. This is impressively substantiated by the emerging studies on the
CCR4-NOT (*CNOT1*) complex in heart structure and function (Yamaguchi et al. 2018; Elmen
et al. 2020).

Throughout our study, we primarily focused on the heart rate as a measure of cardiac function
due to its ease of quantification and interpretation in high-throughput. However, we did also
notice heart morphological phenotypes in some crispants across our hGWAS candidates list.
Among others, *trappc12* crispants showed heart looping defects, resulting in the improper
placement of the atrium compared to the ventricle (Fig 4A). With advances in AI-based image
analysis, important morphological features could be additionally extracted in high throughput,
leading to a better understanding of factors involved in heart development and function.

Grouping the candidate genes according to their heart GWAS association into heart rate and
non-heart rate-related phenotypes further exposed the prominent positive correlation between
the associated human phenotype and the observed phenotype in medaka (Fig 3C). Medaka's
isogenic background, a product of inbreeding over multiple generations (Wittbrodt et al. 2002),
enabled the detection of subtle changes in heart rate immediately in F0 crispants. This
accelerated the analysis and avoided the necessity to analyze homozygous offspring in the
second and third generation after CRISPR targeting.

It is noteworthy that despite the evolutionary distance from fish to humans the medaka
phenotypes match the class of hGWAS effects. This is even more relevant since the roles of the
genes in medaka were validated in embryos, suggesting that the validated marker genes have
predictive power in humans. This deep functional conservation emphasizes the potential of our
approach for the identification and validation of novel predictive genetic markers for
cardiovascular diseases in humans. We have showcased a highly versatile, sensitive and robust
high-throughput reverse genetic validation assay to address the pool of understudied, neglected
putative candidates. In the future, the combination of genetic validation and drug screening in
a single platform building on our assay will facilitate the simultaneous identification of novel
genetic players and interacting small molecules with rescuing power.

313
314   **Materials and Methods**

315
Ethics Statement
All fish are maintained in closed stocks at Heidelberg University. Medaka (*Oryzias latipes*)
husbandry (permit number 35–9185.64/BH Wittbrodt, Regierungspräsidium Karlsruhe) was
performed according to local animal welfare standards (Tierschutzgesetz §11, Abs. 1, Nr. 1) in

accordance with European Union animal welfare guidelines (Bert et al. 2016). The fish facility

is under the supervision of the local representative of the animal welfare agency. The following

medaka stocks and transgenic lines were used: wild-type Cabs and *cmlc2::eGFP cmlc2::H2A-*

mCherry transgenic HdrR-II strain. Medaka embryos were used at stages prior to stage 42.

Medaka were raised and maintained as described previously (Köster et al. 1997).

Generation of the transgenic cmlc2 dual-reporter line

For dual-color cardiac imaging, *cmlc2::eGFP* and *cmlc2::H2A-mCherry* transgenic medaka

lines were generated in the wild-type HdrR-II background. A modified version of pDestTol2CG

(http://tol2kit.genetics.utah.edu/index.php/PDestTol2CG) was used containing a *cmlc2::eGFP*

reporter cassette. For the second, nuclear reporter, the *eGFP* was replaced by an *H2A-mCherry*

insert. Both plasmids were each co-injected at 10 ng/µl with 10 ng/µl Tol2 transposase mRNA

into HdrR-II one-cell stage embryos using the microinjection technique as previously described

(Rembold et al. 2006) to generate separate reporter lines. A double transgenic line was derived

from a cross of the *cmlc2::eGFP* line to *cmlc2::H2A-mCherry* line and maintained for CRISPR-

Cas9 injections.

Candidate gene selection

For the unbiased gene targeting, an online random number generator was used to generate 10

numbers between 1 and 23622, corresponding to the number of annotated medaka coding genes

in Ensembl (Yates et al. 2020) (Table 1). The number of exons for each gene was counted and

a random number was generated to select the exon for CRISPR/Cas9 targeting. For the targeted

human heart-GWAS (hGWAS) gene selection, the genome-wide repository of associations

between SNPs and phenotypes (GRASP v2.0) was used (Eicher et al. 2015). In the search field,

"Heart" and "Heart rate" were chosen as the respective categories for all heart- and heart rate-

related phenotypes associated in human GWAS, only coding SNPs (i.e. SNP functional class =

exons) were searched for. List of resulting genes was extracted (Table 2), and candidate genes

for the functional validation assay were chosen. The focus was on uncharacterized genes, or

genes with no prior experimental link to heart function, yet some known heart genes were

included as proof of concept. For each hGWAS candidate gene, the corresponding medaka

ortholog was extracted using Ensembl (Yates et al. 2020). For the few genes which did not have

an annotated medaka ortholog, the human protein sequence was BLASTed using the "tblastn"

function of the NCBI BLAST (https://blast.ncbi.nlm.nih.gov/Blast.cgi) and Ensemble

(http://www.ensembl.org/Multi/Tools/Blast) online tools to obtain a target medaka locus. Using

Geneious 8.1.9 (https://www.geneious.com), regions of interest (ROI) on medaka orthologous

genes for CRISPR targeting were primarily chosen based on the corresponding location of

human SNP when aligning the medaka and human protein sequences.

sgRNA target sites selection and in vitro transcription

All sgRNA target sites used in this study are listed in Table S1. sgRNAs were designed with

CCTop as described in Stemmer et al. (Stemmer et al. 2015). sgRNA target sites were selected

based on number of potential off-target sites and their corresponding mismatches. Preferably,

sgRNAs selected had no off-target site or at least 3 nucleotide mismatches. sgRNA for *oca2*

was the same as in Lischik et al. (Lischik et al. 2019). Cloning of sgRNA templates and *in vitro*

transcription was performed as detailed in Stemmer, et al. (Stemmer et al. 2015). All sgRNAs

were initially tested after synthesis for *in vivo* targeting via injections into medaka embryos,

followed by genotyping using our filter-in-tips protocol (Hammouda et al. 2019), in brief terms,

by PCR amplification of target locus followed by T7 Endonuclease I assay (New England

Biolabs).

Microinjection

Medaka one-cell or four-cell stage embryos were injected into the cytoplasm as previously

described (Stemmer et al. 2015). Injection solutions for CRISPR targeting comprised: 150 ng/µl

heiCas9 mRNA (Thumberger et al. 2021), 15 ng/µl respective sgRNA and 10 ng/µl *GFP*

mRNA as injection tracer. Control siblings were injected with 10 ng/µl *GFP* mRNA only.

Injected embryos were incubated at 28 °C in embryo rearing medium (ERM), screened for GFP

expression at 1 dpf and transferred to methylene blue-containing ERM (or plain ERM for

reporter lines) and incubated at 28 °C until heart rate analysis (4 dpf) or confocal microscopy

(7 dpf).

 Sample preparation and Imaging

For the heart rate assay, one day prior to imaging (3 dpf), medaka embryos were transferred

from methylene blue-containing ERM into plain ERM and incubated at 28 °C. On day of

imaging (4 dpf), individual medaka embryos (36 per sgRNA and 24 control injected) were

administered to a 96 U-well microtiter plate (Nunc, Thermofisher #268152) containing 200 µl

ERM per well and sealed using gas-permeable adhesive foil (4titude, Wotton, UK, 4ti-

0516/96). Plates were automatically imaged using an ACQUIFER Imaging Machine (DITABIS

AG, Pforzheim, Germany) at 21 and 28 °C with a 30-minute equilibration period before each

measurement. Images were acquired in brightfield using 130 z-slices (dz = 0 µm) and a 2x Plan

UW N.A. 0.06 objective (Nikon, Düsseldorf, Germany) to capture the centered embryo.

Integration times were fixed with 80 % relative white LED intensity and 10 ms exposure time.

Therefore, the whole 96-well plate was captured, with image sequences (videos) of entire

microwells of approx. 10 seconds with 13 frames per second (fps). More details can be found

in Gierten, et al. (Gierten et al. 2020).

For the live confocal microscopy of the reporter lines, injected embryos were treated from 4

dpf onwards with 5x phenylthiourea (PTU) in ERM solution to prevent pigmentation. On the

day of imaging (7 dpf), PTU solution was washed away with ERM, embryos were rolled on

fine sand paper and de-chorionated by incubation in hatching enzyme. Following de-

chorionation, embryos were treated with 50 mM 2,3-butanedione monoxime (BDM) in 1x

Tricaine solution until de-coupling of heart beat (~40 mins), which resulted in fully dilated heart

chambers. Embryos were mounted on Matek dishes in 1.5% low-melting agarose with 85 mM

BDM in 3x Tricaine solution. To avoid dehydration, mounted samples were covered with 30

mM BDM in 1x Tricaine solution throughout the imaging session. All confocal microscopy

images were acquired at a Leica TCS SP8 with 10x dry or 20× oil objective, z-stacks of 200-

300 µm were acquired with a z-step of 5 µm or 1 µm for 10x and 20x acquired images,

respectively.

HeartBeat **detection and data analysis**

Image optimizations prior to analysis, as well as heart rate analysis using the *HeartBeat*

software were performed as previously described (Gierten et al. 2020). In some instances, heart

rates could not be scored due to inconvenient embryo orientations shielding the view of the

Hammouda et al., 2021 Validating understudied GWAS heart genes

heart. For *scn4ab* crispants with cardiac arrhythmias, atrium and ventricle for individual embryos were separately segmented, and the respective beating frequency for each chamber was measured. Data plots were generated using ggplot2 package (Wickham 2016) in R 3.6.1 (R Core Team, 2019) and R-studio 1.2.1335 (RStudio Team, 2018). Statistical analysis for heart rate comparisons were computed in R. Significant differences were determined by two-tailed Student's t-test. Significant *p*-values are indicated with asterisks (*) with $*p < 0.05$, $**p < 0.01$, $***p < 0.001$ and ns (not significant). Maximum intensity projections of confocal microscopy images were processed via Fiji image processing software.

Embryo genotyping

Nucleic acid extraction and genotyping of embryos was done as previously described (Hammouda et al. 2019). Briefly, after imaging, embryos in 96-well plate were lysed in 50 µl Milli-Q water + 50 µl Fin-Clip lysis buffer each (0.4 M Tris-HCl pH 8.0, 5 mM EDTA pH 8.0, 0.15 M NaCl, 0.1 % SDS in Milli-Q water) using a custom 96-well mortar. The mortar was pre-cleaned by incubation in hypochlorite solution (1:10 dilution of commercial bleach reagent) for at least 15 minutes followed by 5 minutes incubation in Milli-Q water. Plates containing lysed embryos were stored at 4 °C until genotyping. To confirm CRISPR on-target activity, per experimental plate, 2 embryos per condition were chosen at random for genotyping by PCR amplification of target locus using our filter-in-tips approach (Hammouda et al. 2019), followed by T7 Endonuclease I Assay (New England Biolabs). 30 PCR cycles were run in all samples, all primers used for PCR are listed in Table S2. Annealing temperatures were calculated using the online NEB Tm calculator (https://tmcalculator.neb.com/).

Acknowledgments

We thank A. Cornean for providing the *gfp*_T1 guide RNA and for his help in acquiring the confocal images. We thank V. Weinhardt, E. Tsingos and all members of the Wittbrodt lab for their critical, constructive feedback on the procedure and the manuscript, F. Loosli and S. Lemke for their constructive feedback towards the project as well as J. Backs and E. Furlong for an outside perspective. We thank T. Kellner for excellent technical support. We acknowledge the excellent fish husbandry of E. Leist, M. Majewsky and A. Saraceno. We thank

J. Gehrig (ACQUIFER Imaging GmbH) for supporting us with the Imaging machine. O.T.H. is and J.G. was a member of the Heidelberg Biosciences International Graduate School (HBIGS). This work was supported by a fellowship of the Deutsches Zentrum für Herz-Kreislauf-Forschung (DZHK) to O.T.H and through DFG CRC-1324 TP B4 and NIH 5R01ES029917 – 03, KiyosuTox. J.G. was supported by the Deutsche Herzstiftung e.V. (S/02/17) and by an Add-On Fellowship for Interdisciplinary Science of the Joachim Herz Stiftung.

Author contributions

O.T.H., T.T. and J.W. designed the study and implemented the methodology. J.G. generated the transgenic *cmlc2* dual-reporter line. O.T.H., V.K. and M.Y.W. synthesized the guide RNAs. O.T.H. performed experiments. O.T.H. analyzed the data with contributions by M.Y.W. and V.K.. O.T.H. visualized data. O.T.H. and J.W. wrote the first draft of the manuscript. O.T.H., T.T., and J.W. finalized the manuscript. J.W. provided resources and supervised this work.

Conflict of interest

All authors declare no competing interests.

References

Beere, P.A., Glagov, S. & Zarins, C.K., 1984. Retarding effect of lowered heart rate on coronary atherosclerosis. *Science*, 226(4671), p.180.

Benson, M.A. et al., 2017. Ryanodine receptors are part of the myospryn complex in cardiac muscle. *Scientific Reports*, pp.1–12.

Bert, B. et al., 2016. Considerations for a European animal welfare standard to evaluate adverse phenotypes in teleost fish. *The EMBO Journal*, 35(11), pp.1151–1154.

Cambien, F. & Tiret, L., 2007. Genetics of Cardiovascular Diseases. *Circulation*, 116(15), pp.1714–1724.

Cordell, H.J. et al., 2013. Genome-wide association study identifies loci on 12q24 and 13q32 associated with Tetralogy of Fallot. *Human Molecular Genetics*, 22(7), pp.1473–1481.

DYER, A.R. et al., 1980. HEART RATE AS A PROGNOSTIC FACTOR FOR
CORONARY HEART DISEASE AND MORTALITY: FINDINGS IN THREE
CHICAGO EPIDEMIOLOGIC STUDIES. *American Journal of Epidemiology*, 112(6),
pp.736–749.

Eicher, J.D. et al., 2015. GRASP v2.0: an update on the Genome-Wide Repository of
Associations between SNPs and phenotypes. *Nucleic Acids Research*, 43(D1), pp.D799–
D804.

Eijgelsheim, M. et al., 2010. Genome-wide association analysis identifies multiple loci related
to resting heart rate. *Human Molecular Genetics*, 19(19), pp.3885–3894.

Elmen, L. et al., 2020. Silencing of CCR4-NOT complex subunits affect heart structure and
function. *Disease Models & Mechanisms*, pp.dmm.044727–33.

Eppinga, R.N. et al., 2016. Identification of genomic loci associated with resting heart rate
and shared genetic predictors with all-cause mortality. *Nature Genetics*, 48(12), pp.1557–
1563.

Faggioni, M. & Knollmann, B.C., 2012. Calsequestrin 2 and arrhythmias. *American Journal
of Physiology-Heart and Circulatory Physiology*, 302(6), pp.H1250–H1260.

Fukamachi, S. et al., 2004. Conserved function of medaka pink-eyed dilution in melanin
synthesis and its divergent transcriptional regulation in gonads among vertebrates.
Genetics, 168(3), pp.1519–1527.

Grabher, C. & Wittbrodt, J., 2007. Meganuclease and transposon mediated transgenesis in
medaka. *Genome biology*, 8 Suppl 1(Suppl 1), p.S10.

Gianulis, E.C. & Trudeau, M.C., 2011. Rescue of Aberrant Gating by a Genetically Encoded
PAS (Per-Arnt-Sim) Domain in Several Long QT Syndrome Mutant Human Ether-á-go-
go-related Gene Potassium Channels. *Journal of Biological Chemistry*, 286(25),
pp.22160–22169.

Gierten, J. et al., 2020. Automated high-throughput heartbeat quantification in medaka and
zebrafish embryos under physiological conditions. *Scientific Reports*, pp.1–12.

Gillum, R.F., Makuc, D.M. & Feldman, J.J., 1991. Pulse rate, coronary heart disease, and
death: The NHANES I Epidemiologic Follow-up Study. *American Heart Journal*, 121(1),
pp.172–177.

Gut, P. et al., 2017. Little Fish, Big Data: Zebrafish as a Model for Cardiovascular and
Metabolic Disease. *Physiological Reviews*, 97(3), pp.889–938.

Hammouda, O.T. et al., 2019. Swift Large-scale Examination of Directed Genome Editing S.
C. F. Neuhauss, ed. *PLoS ONE*, 14(3), pp.e0213317–11.

Han, P. et al., 2014. Hydrogen peroxide primes heart regeneration with a derepression
mechanism. *Nature Publishing Group*, pp.1–17.

Heyde, von der, B. et al., 2020. Translating GWAS-identified loci for cardiac rhythm and rate
using an in vivo image- and CRISPR/Cas9-based approach. *Scientific Reports*, pp.1–18.

Hiura, Y. et al., 2010. A Genome-Wide Association Study of Hypertension-Related
Phenotypes in a Japanese Population. *Circulation Journal*, 74(11), pp.2353–2359.

Hoed, den, M. et al., 2013. Identification of heart rate–associated loci and their effects on
cardiac conduction and rhythm disorders. *Nature Genetics*, pp.1–14.

Holm, H. et al., 2011. A rare variant in MYH6 is associated with high risk of sick sinus
syndrome. *Nature Genetics*, 43(4), pp.316–320.

Hu, Z. et al., 2013. A genome-wide association study identifies two risk loci for congenital
heart malformations in Han Chinese populations. *Nature Genetics*, pp.1–5.

Huttner, I.G. et al., 2013. A transgenic zebrafish model of a human cardiac sodium channel
mutation exhibits bradycardia, conduction-system abnormalities and early death. *Journal
of Molecular and Cellular Cardiology*, 61(C), pp.123–132.

Ikram, M.A. et al., 2009. Genomewide Association Studies of Stroke. *New England Journal
of Medicine*, 360(17), pp.1718–1728.

Itoh-Satoh, M. et al., 2002. Titin Mutations as the Molecular Basis for Dilated
Cardiomyopathy. *Biochemical and Biophysical Research Communications*, 291(2),
pp.385–393.

Iwamatsu, T., 2004. Stages of normal development in the medaka Oryzias latipes.
Mechanisms of Development, 121(7-8), pp.605–618.

Kathiresan, S. & Srivastava, D., 2012. Genetics of Human Cardiovascular Disease. *Cell*,
148(6), pp.1242–1257.

Koster, R. et al., 1997. Medaka spalt acts as a target gene of hedgehog signaling.
Development, 124(16), p.3147.

Lessman, C.A., 2011. The developing zebrafish (Danio rerio): A vertebrate model for high-
throughput screening of chemical libraries R. S. Tuan, ed. *Birth Defects Research Part C:
Embryo Today: Reviews*, 93(3), pp.268–280.

Li, J. et al., 2017. Essential role of Cdc42 in cardiomyocyte proliferation and cell-cell
adhesion during heart development. *Developmental Biology*, 421(2), pp.271–283.

Lischik, C.Q., Adelmann, L. & Wittbrodt, J., 2019. Enhanced in vivo-imaging in medaka by
optimized anaesthesia, fluorescent protein selection and removal of pigmentation C.
Winkler, ed. *PLoS ONE*, 14(3), pp.e0212956–19.

Marroni, F. et al., 2009. A Genome-Wide Association Scan of RR and QT Interval Duration
in 3 European Genetically Isolated Populations. *Circulation: Cardiovascular Genetics*,
2(4), pp.322–328.

Matsa, L.S. et al., 2014. Endothelin 1 gene as a modifier in dilated cardiomyopathy. *Gene*,
548(2), pp.256–262.

McGill, H.C., Jr, McMahan, C.A. & Gidding, S.S., 2008. Preventing Heart Disease in the 21st
Century. *Circulation*, 117(9), pp.1216–1227.

MD, P.M.B. et al., 2010. Heart rate as a risk factor in chronic heart failure (SHIFT): the
association between heart rate and outcomes in a randomised placebo-controlled trial. *The
Lancet*, 376(9744), pp.886–894.

Meyer, H.V. et al., 2020. Genetic and functional insights into the fractal structure of the heart.
Nature, pp.1–25.

Nemtsas, P. et al., 2010. Adult zebrafish heart as a model for human heart? An
electrophysiological study. *Journal of Molecular and Cellular Cardiology*, 48(1),
pp.161–171.

Newton-Cheh, C. et al., 2009. Common variants at ten loci influence QT interval duration in
the QTGEN Study. *Nature Genetics*, 41(4), pp.399–406.

Oxendine, S.L. et al., 2006. Adapting the medaka embryo assay to a high-throughput
approach for developmental toxicity testing. *NeuroToxicology*, 27(5), pp.840–845.

Paré, G. et al., 2013. Genetic Determinants of Dabigatran Plasma Levels and Their Relation
to Bleeding. *Circulation*, 127(13), pp.1404–1412.

Park, C.Y. et al., 2010. skNAC, a Smyd1-interacting transcription factor, is involved in
cardiac development and skeletal muscle growth and regeneration. *Proc Natl Acad Sci
USA*, 107(48), p.20750.

Pfeufer, A. et al., 2009. Common variants at ten loci modulate the QT interval duration in the
QTSCD Study. *Nature Genetics*, 41(4), pp.407–414.

Qian, L. et al., 2011. Tinman/Nkx2-5 acts via miR-1 and upstream of Cdc42 to regulate heart
function across species. *The Journal of Cell Biology*, 193(7), pp.1181–1196.

R Core Team (2019). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

Rathjens, F.S. et al., 2020. Preclinical evidence for the therapeutic value of TBX5
normalization in arrhythmia control. *Cardiovascular Research*.

Rembold, M. et al., 2006. Transgenesis in fish: efficient selection of transgenic fish by co-
injection with a fluorescent reporter construct. *Nature Protocols*, 1(3), pp.1133–1139.

Ritchie, M.D. et al., 2013. Genome- and Phenome-Wide Analyses of Cardiac Conduction
Identifies Markers of Arrhythmia Risk. *Circulation*, 127(13), pp.1377–1385.

RStudio Team (2018). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA
URL http://www.rstudio.com/

Shankaran, S.S. et al., 2018. CRISPR/Cas9-Directed Gene Editing for the Generation of Loss-
of-Function Mutants in High-Throughput Zebrafish F 0Screens. *Current Protocols in*
Molecular Biology, 119(1), pp.4257–22.

Smith, J.G. et al., 2009. Genome-wide association study of electrocardiographic conduction
measures in an isolated founder population: Kosrae. *HRTHM*, 6(5), pp.634–641.

Sotoodehnia, N. et al., 2010. Common variants in 22 loci are associated with QRS duration
and cardiac ventricular conduction. *Nature Genetics*, pp.1–11.

Stemmer, M. et al., 2015. CCTop: An Intuitive, Flexible and Reliable CRISPR/Cas9 Target
Prediction Tool S. Maas, ed. *PLoS ONE*, 10(4), pp.e0124633–11.

Targoff, K.L. et al., 2013. Nkx genes are essential for maintenance of ventricular identity.
Development, 140(20), pp.4203–4213.

Teboul, L., 2017. Phenotyping first-generation genome editing mutants: a new standard?
Mammalian Genome, 28(7), pp.377–382.

Thanassoulis, G. et al., 2013. Genetic Associations with Valvular Calcification and Aortic
Stenosis. *New England Journal of Medicine*, 368(6), pp.503–512.

Thumberger, T. et al., 2021. hei-tag: a highly efficient tag to boost targeted genome editing.
bioRxiv, p.2021.05.27.445956.

Vasan, R.S. et al., 2009. Genetic Variants Associated With Cardiac Structure and Function: A
Meta-analysis and Replication of Genome-wide Association Data. *JAMA*, 302(2),
pp.168–178.

Villard, E. et al., 2011. A genome-wide association study identifies two loci associated with
heart failure due to dilated cardiomyopathy. *European Heart Journal*, 32(9), pp.1065–
1076.

Wickham, H., 2016. *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag New
York. Available at: https://ggplot2.tidyverse.org.

Wittbrodt, J., Shima, A. & Schartl, M., 2002. MEDAKA — A MODEL ORGANISM FROM
THE FAR EAST. *Nature Reviews Genetics*, 3(1), pp.53–64.

Wooten, E.C. et al., 2010. Application of Gene Network Analysis Techniques Identifies
AXIN1/PDIA2 and Endoglin Haplotypes Associated with Bicuspid Aortic Valve A. E.
Toland, ed. *PLoS ONE*, 5(1), pp.e8830–10.

Wu, R.S. et al., 2018. A Rapid Method for Directed Gene Knockout for Screening in G0
Zebrafish. *Developmental Cell*, 46(1), pp.112–125.e4.

Yamaguchi, T. et al., 2018. The CCR4-NOT deadenylase complex controls Atg7-dependent
cell death and heart function. *Sci. Signal.*, 11(516), p.eaan3638.

Yates, A.D. et al., 2020. Ensembl 2020. *Nucleic Acids Research*, 48(D1), pp.D682–D688.

Yonekura, M. et al., 2018. Medaka as a model for ECG analysis and the effect of verapamil.
Journal of Pharmacological Science, 137(1), pp.55–60.

Zakariyah, A.F. et al., 2017. Congenital heart defect causing mutation in Nkx2.5 displays in
vivo functional deficit. *Journal of Molecular and Cellular Cardiology*, 105(C), pp.89–98.

Zaklyazminskaya, E. & Dzemeshkevich, S., 2016. The role of mutations in the SCN5A gene
in cardiomyopathies. *BBA - Molecular Cell Research*, 1863(Part B), pp.1799–1805.

Hammouda et al., 2021 Validating understudied GWAS heart genes

Figure S1. Highly efficient CRISPR/Cas9-mediated editing in injected (F0) generation
A Distribution of reporter expression in mock injected and *gfp_T1* crispants of the *cmlc2::eGFP cmlc2::H2A-mCherry* reporter line (4 dpf). Embryos were injected either at the 1-cell or 4-cell stage. Note: complete lack of GFP-expressing embryos when injected at the 1-cell stage. Biological replicates for each group is denoted (n). **B** Confocal images of GFP expression in mock-injected and *gfp* crispant embryo hearts of the *cmlc2::eGFP cmlc2::H2A-mCherry* reporter line (7 dpf). Embryos were injected either at the 1-cell or 4-cell stage. Note: complete loss of GFP expression when injected at the 1-cell stage (n = 8/8), while mosaic expression when injected at 4-cell stage (n = 4/4). Scale bars: 100 µm (First panel on left) and 50 µm (blow-up images).

Hammouda et al., 2021 Validating understudied GWAS heart genes

Figure S2. Consistent heart rate phenotype observed in medaka *nkx2-5* crispants
A Overview of 96-well plate with embryos (4 dpf) injected with sgRNA against *nkx2-5* or *oca2*, as well as embryos mock injected with *GFP mRNA* (Fig 1D). Note the loss of eye pigmentation in *oca2* crispant embryos. **B-C** Heart rate measurements of *GFP*-injected (Mock; dark grey) and *nkx2-5* crispant embryos (4 dpf) (**B** second replicate of *nxk2-5_T4*; **C** different sgRNA *nkx2-5_T5* targeting same region of interest) at 21 and 28°C, before and after exclusion of severely affected embryos (< stage 28; developmental focusing). Significant differences are shown in red and were determined by two-tailed Student's t-test; *$p < 0.05$, **$p < 0.01$, ns (not significant; light grey). For biological replicates see Source Data Fig S2.

Hammouda et al., 2021 Validating understudied GWAS heart genes

	21°C				28°C			
	Developmental Focusing				Developmental Focusing			
	Before		After		Before		After	
	HR % Change	*p*-value	HR % Change	*p*-value	HR % Change	*p*-value	HR % Change	*p*-value
cdc42	6.0	2.7e−04	6.4	5.9e−05	5.8	7.7e−05	6.3	2.0e−05
ogdh	3.2	3.4e−02	3.2	3.4e−02	2.8	4.0e−02	2.8	4.0e−02
duox	−1.2	4.2e−01	−1.6	2.9e−01	−2.3	1.3e−01	−2.2	1.7e−01
git2	0.1	9.5e−01	0.5	7.2e−01	0.3	8.4e−01	0.9	5.8e−01
mus81	2.7	1.4e−01	2.7	1.1e−01	1.4	3.5e−01	2.3	8.3e−02
or124−2	−0.3	8.5e−01	−1.3	3.9e−01	−0.1	9.4e−01	0.0	9.9e−01
plekha8	−1.4	5.5e−01	1.3	4.9e−01	−2.6	3.2e−01	0.8	6.7e−01
ttl	0.9	6.5e−01	3.2	8.3e−02	−1.8	4.1e−01	1.7	3.1e−01
cabp4	2.0	2.3e−01	2.0	2.3e−01	1.9	1.9e−01	1.9	1.9e−01
eml6	2.1	2.2e−01	1.6	3.2e−01	2.9	6.0e−02	2.7	7.4e−02

Figure S3. Developmental focusing does not alter analysis outcome of random gene selection
Heatmap quantitative representation of the comparative heart rate analysis between each crispant embryo group and its corresponding control sibling group before and after developmental focusing; for each measured temperature, the percent change in mean heart rate (HR % Change) between crispants and their corresponding control sibling, flanked by the statistical significance (*p*-value) of the observed change calculated by two-tailed Student's t-test on the full distribution. Genes showing significant heart rate phenotypes are indicated in **bold**. For biological replicates see Source Data Fig S3.

Hammouda et al., 2021 Validating understudied GWAS heart genes

Figure S4. Comparative analysis of mean heart rates in targeted hGWAS gene selection
Heart rate measurements (beats per minute, bpm) of *GFP*-injected (Mock; dark grey) and corresponding sibling crispant embryos (4 dpf) at 21 and 28°C after developmental focusing (also see heatmap representation of the data in Fig 3A). Different experimental plates are represented by breaks on the x-axis. Significant differences in mean heart rates were determined between each crispant embryo group and its corresponding sibling control group by two-tailed Student's t-test; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ns (not significant). Red groups correspond to crispants showing significant heart rate phenotypes, and light grey groups correspond to crispants showing no significant heart rate phenotype. For biological replicates see Source Data Fig S4.

Hammouda et al., 2021 Validating understudied GWAS heart genes

Fig S5. Developmental focusing does not alter analysis outcome of targeted hGWAS genes
Heatmap quantitative representation of the comparative heart rate analysis between each crispant embryo group and its corresponding control sibling group before and after developmental focusing; for each measured temperature, the percent change in mean heart rate (HR % Change) between crispants and their corresponding control siblings, flanked by the statistical significance (p-value) of the observed change, calculated by two-tailed Student's t-test on the full distribution. Genes showing significantly different heart rate phenotypes are indicated in **bold**. For biological replicates see Source Data Fig S4.

Hammouda et al., 2021 Validating understudied GWAS heart genes

Movie S1. Moderate AV-block arrhythmia observed in medaka F0 *scn4ab* crispants and
homozygous F2 mutants
Side by side comparison of rhythmic heartbeat of *GFP*-injected (Mock; left) and arrhythmic *scn4ab*
crispants (F0; middle) as well as homozygous mutants (F2; right) displaying 2:1 AV-block phenotype.
Videos of medaka embryos (5 dpf) were acquired using a stereomicroscope under bright field
illumination.
636

637 **Movie S2. Severe AV-block arrhythmia observed in medaka *scn4ab* crispants**
638 Side by side comparison of rhythmic heartbeat of *GFP*-injected (Mock; left) and arrhythmic *scn4ab*
639 crispants displaying severe AV-block phenotype (right). Videos of medaka embryos (9 dpf) were
640 acquired using a stereomicroscope under bright field illumination.
641

642 **Table S1. List of sgRNAs used**
643 sgRNA target sites given in 5'-3' direction, protospacer adjacent motif (PAM) in square brackets.
644

sgRNA	Target sequence [PAM]
abcb4_T7	ATGTCCCTGAGTGTGAAGAG[TGG]
atp8b4_T2	TAAGTTCTACGATAACACCC[TGG]
bag3_T1	GAGCAGGGCGCCGTTCACCC[GGG]
cabp4_T9	GAATTTGACTACGATGCAGA[TGG]
casq2_T1	TGTTGTCTGGGATGGGTTCG[TGG]
ccdc141_T1	CCAGGAACCTGAAGGTTGTC[AGG]
cdc42_T2	AGAGCGGAGAGAAACTGGCT[CGG]
cep85l _T5	AACCCAGGACTTCTCAGATA[GGG]
clcnk_T10	TTCTCCACTGGAGTAGTTTT[TGG]
cmya5_T1	GTAGAACAGGTAATTCTCGT[TGG]
cnot1_T5	GCCAAGTGTTGACAATACTG[AGG]
col9a1b_T2	GACGGACGCGTAGGCATTCC[AGG]
duox_T3	CATTCGGCACGCTTTCTCTA[AGG]
edn1_T4	GCCGACGGAGTCTGCGCGGA[GGG]
eml6_T2	CTGCGCTGTTCGCACGCTAA[AGG]
gfp_T1	AGCACTGCACGCCGTAGGTC[AGG]
gigyf1_T3	GAATGAACCGGCATGAACGC[CGG]
git2_T3	TAAACGCCTTCGAAACACGG[AGG]
grid2_T8	AAAGGCTACGGCTAAGGGTC[TGG]
hcn4_T5	AAACTCCCTTCGAACTTGTG[AGG]
homeza_T1	GCTACCAGCAGGTGCGAGAT[TGG]
kcnh2_T1	CATCACTGCTGGGAGAACCG[GGG]
maml3_T19	TCATGTAAGGTGTCATCATA[GGG]

Hammouda et al., 2021 Validating understudied GWAS heart genes

sgRNA	Target sequence [PAM]
minar1_T1	GTTGCCGTCGGCGACGCGTA[GGG]
mus81_T7	AAGAGGATGGACGACCTCTG[TGG]
myrf_T1	CCTTATTGGAGTCCATATTG[TGG]
naca_T1	TTGGTCTTAGGCAAGTAACG[GGG]
nkx2-5_T4	GCCGCGGGTCCTCTTCTCCC[AGG]
nkx2-5_T5	CGGACAGACCCAAGCCCCGG[AGG]
nubp2_T4	GAACTGAATGTGGCACTGTT[AGG]
oca2_T3	TTGCAGGAATCATTCTGTGT[GGG]
ogdh_T4	CAAAGCTGGACTTGGCCTCA[GGG]
or124-2_T1	GCCCACGTGGTCTCGTACGC[CGG]
or5au1_T2	GTCATGGACTCTTGCCTGTA[TGG]
padi2_T4	AGACTGGTATTACGGTTAAG[AGG]
piezo1_T1	TTGGGAGCCCGGATGTAGTT[AGG]
plekha8_T9	GCGGTGAAGCTTTCATAACA[CGG]
plg_T1	GGCAACGGGGCCAATTATCG[AGG]
ppp1r9a_T7	TGGTTCTTAGGAGATTCGGG[TGG]
rgs3a_T6	GATCAAGTCACAGTCCAAGA[TGG]
rnf207b_T2	TAGACTTGTTTGTACTGATC[TGG]
scmh1_T2	TCTGACCTGCGCTGTACGAG[TGG]
scn4ab_T2	TTAGGCTTAGCAATCCGCAA[CGG]
sh2b3_T1	AAGAAGTCCGTCGCTGCAAT[CGG]
slc17a3_T5	GTCTTGGCGCCGATTGTGAC[AGG]
smg6_T2	AAAGATAAAACCAAGGGCGT[AGG]
sspo_T4	GTGCACCAAGTCATGTGGTT[GGG]
trappc12_T2	CAACACGCAGTGCCTCAAGC[TGG]
ttl_T1	GCTGGTAAATTACTACAGAG[GGG]
ttn.2_T1	GAGCTAGCTGTCAAAGCCAT[GGG]
ufsp1_T1	GCTGGAGAGCATCGCAGTCC[AGG]
veph1_T1	GGCTCTGGTGGAGGTGTCCC[AGG]
xylb_T1	TTTCCAGTGTCAGTACCTAC[AGG]
zfhx3_T1	AGCTGAGCGCACCCTGCCTG[AGG]

Hammouda et al., 2021 Validating understudied GWAS heart genes

650   **Table S2. List of primers used for genotyping by PCR**
651

Target Gene	Primers (5' – 3' direction)
abcb4 Lf	CCTCTGCAGAAGCTGGATGG
abcb4 Lr	TTTGCGATGGCGTGTGAGCG
atp8b4 Lf	ACCTACAGTCAAAGGGAAATGAGC
atp8b4 Lr	ACGCTGCCTCGTCTGGTGAC
bag3 Lf	CTTCGCTTAGGAGCCAGTCC
bag3 Lr	CAGTATGGTCTGAACCCGCC
cabp4 Lf	GTGCCATCTTGCCATGCTGC
cabp4 Lr	ATCCTGCTGCTCCTGGTCCC
casq2 Lf	CGAACAGGTCAAACCGTGTG
casq2 Lr	GGAACTTGCACAAACGGACC
ccdc141 Lf	GGCTACGGATGAAGGAGCTC
ccdc141 Lr	TCCTGGCTAAGTGAAGCAGC
cdc42 Lf	TCCAGAGCTGCGAGGATGGC
cdc42 Lr	CCAGCCGCTCTGTTCTGGGC
cep85l Lf	AGGAGTGAGCTGAGGATGGG
cep85l Lr	GCATCCTCACTCTAGCCGGG
clcnk Lf	GATTCATGCTGTGTTCCCGC
clcnk Lr	CCACTGAGTTGGTCGCTTTG
cmya5 Lf	TTTGTGGGAGCTCGTCTGGG
cmya5 Lr	GCCACGCTGAGATGGAGCCC
cnot1 Lf	CCGGTTGTGGCGCCTTGAGG
cnot1 Lr	GCTTTGGGTCTGGCACTGCG
col9a1b Lf	AGGGACTCACTTACCGGAAGCCC
col9a1b Lr	CTGGTCCTCAGGGAGTGGCAGG
duox Lf	GACGCTGTCAGCTCGCACTG
duox Lr	GGTCCACTCATCGCTCCCTC
edn1 Lf	TCTCCGTGCTGTCAGTGTTC
edn1 Lr	ATGCTGGTTGCCATGGAGTC
eml6 Lf	ACGCCACGGACAGCATGGTC
eml6 Lr	TCACTGACAGCTTTGAAGGTTGACG
gfp Lf	GTGAGCAAGGGCGAGGAGCT
gfp Lr	CTTGTACAGCTCGTCCATGC
gigyf1 Lf	ACAGGAGATAGCAGCAGCAG

Hammouda et al., 2021 Validating understudied GWAS heart genes

Target Gene	Primers (5' – 3' direction)
gigyf1 Lr	CGTCCCTGATTGTAGATTTGCG
git2 Lf	TTGCTGTCGCCGCTGCTCTG
git2 Lr	GGGCTCATGGGACCTTATGTTGG
grid2 Lf	ACCTACAGACCTGTAACTCCTTCGC
grid2 Lr	ACCTGGAGCACCTACACCACAGG
hcn4 Lf	GACCCGTCAAGTGCCACAGG
hcn4 Lr	CACACCCTCCTTGTCCCTTG
homeza Lf	GCCAAGGAAGACCAAAGAGC
homeza Lr	TCACAAACCTCGCTCTAGGC
kcnh2 Lf	TCGCCCAGGCAAGTCCAACG
kcnh2 Lr	ACCTCATCGCCACTTGAGTGTG
maml3 Lf	CGGCAGCCATGTTGTCGTAC
maml3 Lr	AAGCGGGTTATGTGTGCTGC
minar1 Lf	CGGTCAGCATGAACTAAGCTTCGG
minar1 Lr	TGGAGACGGACAGTGATTCCAGTGG
mus81 Lf	GAAGACGACGAGACGGCCGG
mus81 Lr	TGACCCGAACAGTACCTTTGTG
myrf Lf	GGTCCACCAACCGCCTGCC
myrf Lr	AGCATCCTAGCATTGCAGCC
naca Lf	TCGCTCTCCTCCTGTACTGTAGGG
naca Lr	TCTGAGTCTACTGAAGCCAGCC
nkx2-5 lf	TAGCTTTGAGGCCTGCAGAC
nkx2-5 lr	CGCATAGTTCGTGTTGCAGG
nubp2 Lf	ACTGATGAGCCAGCCAACCC
nubp2 Lr	TGGTCTGTGGCCACATGGTG
oca2 Lf	GTTAAAACAGTTTCTTAAAAAGAACAGGA
oca2 Lr	AGCAGAAGAAATGACTCAACATTTTG
ogdh Lf	GAACACTCGTCTGGCCGGCC
ogdh Lr	TACAGCAAGTCCCGCCCACG
or124-2 Lf	TGAGGCTCAGACCCTCCTGG
or124-2 Lr	ATGCGGTCCGGATGAGCTGG
or5au1 Lf	GCATTCAGCAGTTCTTCTGTCTTC
or5au1 Lr	GGCTTTGATAGATCTGTGTGGAAGC
padi2 Lf	GCATGACACTACCTGAGAATGAAAC
padi2 Lr	AGCCTCAGTCATCTCAGTAAAGC

Hammouda et al., 2021 Validating understudied GWAS heart genes

Target Gene	Primers (5' – 3' direction)
piezo1 Lf	GCAGCAGCAGGAAGGTCAGG
piezo1 Lr	CCGCTGCATTAGACCACTGC
plekha8 Lf	CTGGCACCTTCCTGTCCAGC
plekha8 Lr	ACGGAAGTCGCCCAAGTCCG
plg Lf	CACAAACACAGCCGCACGCC
plg Lr	GTGGGCGTGTCCGAGTCGAC
ppp1r9a Lf	CGAGGTCCAACCGAGGCAGC
ppp1r9a Lr	TCTCTGTCAGATCTGAGCCGGG
rgs3a Lf	TGAGCTGTGTTCGCCCACTC
rgs3a Lr	GGAGGGCTTCAACTGTGAGG
rnf207b Lf	GCAGAAGCCTCCCATTGACG
rnf207b Lr	CCTCTGGCTGACCGCTCCTG
scmh1 Lf	ACTCTGTTCGAGCGCCTCCC
scmh1 Lr	AGGCGCATCCTGCAGCTTCC
scn4ab Lf	TCACCACTTCTGCAGGATGG
scn4ab Lr	GAGGCGTAGTGATCTGACGG
sh2b3 Lf	TTTCTGCCGGCATCTGCTCC
sh2b3 Lr	CCTGCTGGTCGTTGTCCGTC
slc17a3 Lf	CCGCTGTGCCCAGGAGCAAG
slc17a3 Lr	AGGAAACGACCCGTGACCAC
smg6 Lf	CCCAGCATCAGTGGAGGTGC
smg6 Lr	CCCTGAGCCTCAGTCCCAGC
sspo Lf	TGCAGTCGAGGGTCAGTGGTCG
sspo Lr	CACAGGATCGCGGGCAAGAAGG
trappc12 Lf	GTGCCTGCTGTACCTGGGCC
trappc12 Lr	ACCACCACCCTCTCTCCCAGC
ttl Lf	CCGGGCGGGACCAACAACTG
ttl Lr	GGAGTCAGACAGCTCTGGGC
ttn.2 Lf	CAGAGCTGGACGGACATCGG
ttn.2 Lr	GGAGCGCTGACGTCTGCCTC
ufsp1 Lf	TGGGAAAGGACTAGAGGAGGG
ufsp1 Lr	GACACTCGCTTCCAGGACAC
veph1 Lf	TCTGTCAGAGCGGATGCAAG
veph1 Lr	AGTCTCTGATCGGGATGCAAC
xylb Lf	CCTGCAGCAACATGGCAGTGTG

Target Gene	Primers (5' – 3' direction)
xylb Lr	ACCATCACTGTGGTCAATGGCTGC
zfhx3 Lf	GGCTGGGTGGTGGGTGTCAG
zfhx3 Lr	TGTGCGGCTAGGTGGTGGAC

Source Data Fig 1D. Biological replicates for Fig 1D

Ns correspond to number of scored embryos per condition. before DF = before developmental focusing (i.e. all scored embryos). after DF = after developmental focusing (i.e. only scored embryos looking older than stage 28)

Gene_sgRNA	Ns before DF		Ns after DF		Ns mock	
	21C	**28C**	**21C**	**28C**	**21C**	**28C**
nkx2-5_T4	34	34	31	31	22	23
oca2_T1	35	35	35	35		

Source Data Fig S2. Biological replicates for Fig S2 B-C

Ns correspond to number of scored embryos per condition. before DF = before developmental focusing (i.e. all scored embryos). after DF = after developmental focusing (i.e. only scored embryos looking older than stage 28)

Gene_sgRNA	Ns before DF		Ns after DF		Ns mock	
	21C	**28C**	**21C**	**28C**	**21C**	**28C**
nkx2-5_T4	34	36	31	33	19	23
nkx2-5_T5	33	33	31	31	22	21

Source Data Fig S3. Biological replicates for Fig 2 and S3

Ns correspond to number of scored embryos per condition. before DF = before developmental focusing (i.e. all scored embryos). after DF = after developmental focusing (i.e. only scored embryos looking older than stage 28)

Gene	Ns before DF		Ns after DF		Ns mock before -> after DF	
	21C	**28C**	**21C**	**28C**	**21C**	**28C**
cdc42	32	32	29	30	22	20
ogdh	31	33	31	33		
duox	32	33	27	28	19	21
git2	35	33	33	31		
mus81	27	33	25	31	20 -> 18	21 -> 20
or124-2	34	34	30	30		
plekha8	35	31	29	26	21	23
ttl	33	30	28	24		
cabp4	34	34	34	34	21	21
eml6	36	34	34	32		

Source Data Fig S4. Biological replicates for Fig 3A, S4 and S5

Ns correspond to number of scored embryos per condition. before DF = before developmental focusing (i.e. all scored embryos). after DF = after developmental focusing (i.e. only scored embryos looking older than stage 28)

Gene	Ns before DF		Ns after DF		Ns mock before -> after DF	
	21C	**28C**	**21C**	**28C**	**21C**	**28C**
bag3	32	33	32	33	22	24
edn1	35	34	35	34		
zfhx3	33	35	33	35	23	22
nubp2	36	34	36	34		
minar1	33	33	30	30	24	22
ppp1r9a	32	34	32	34		
atp8b4	35	34	32	31	24	23
padi2	31	36	31	36		
sspo	33	32	33	32	22	22
scn4ab	35	32	35	32		
ufsp1	33	32	33	32	23	24
veph1	34	35	34	35		
clcnk	34	31	34	31	22	23
rgs3a	26	28	21	23		
myrf	34	34	33	33	24 -> 23	23 -> 22
cnot1	36	35	22	22		
trappc12	35	35	34	34	23	24
or5au1	34	33	34	33	22	22
naca	33	32	27	27		
abcb4	36	35	36	35	23	22
smg6	36	35	30	29		
maml3	35	35	34	34	23	24
plg	35	36	33	34		
sh2b3	35	36	35	36	24	24
rnf207b	35	36	33	34		
slc17a3	36	35	36	35	22	24
scmh1	33	33	33	33		
hcn4	35	36	33	34	23	24
grid2	30	34	30	34	23	20
casq2	59	56	53	50	25 -> 24	24 -> 23
ccdc141	36	34	36	34	22	23
gigyf1	34	35	34	35		

Hammouda et al., 2021 Validating understudied GWAS heart genes

Gene	Ns before DF		Ns after DF		Ns mock before -> after DF	
	21C	**28C**	**21C**	**28C**	**21C**	**28C**
homeza	32	35	32	35	19	23
cep85l	35	36	35	36	22	24
piezo1	35	33	35	33		
ttn.2	34	35	34	35	23	24
cmya5	33	34	33	34		
xylb	35	33	35	33	23	22
col9a1b	35	35	34	34		
kcnh2	34	36	34	36	21 -> 20	21