1 **Touch-triggered bZIP translocation regulates elongation and secondary wall biosynthesis**
2

3 Joshua H. Coomey[1,2], Kirk J.-M. MacKinnon[1,3], Pubudu P. Handakumbura[1,2], Ian W.
4 McCahill[1,2], Gina M. Trabucco[1,3], Jessica Mazzola[1], Nicole A. Leblanc[1], Rithany Kheam[1],
5 Miriam Hernandez-Romero[1,2], Kerrie Barry[4], Lifeng Liu[4], Ji E. Lee[4], John P. Vogel[4], Ronan C.
6 O'Malley[4], James J. Chambers[5], Samuel P. Hazen[1,2,3]
7
8

[1]Biology Department, University of Massachusetts, Amherst, MA 01003, USA,

[2]Plant Biology Graduate Program, University of Massachusetts, Amherst, MA 01003, USA,

[3]Molecular and Cellular Biology Graduate Program, University of Massachusetts, Amherst, MA
01003, USA,

[4]DOE Joint Genome Institute, Berkeley, CA 94720, USA,

[5]Institute for Applied Life Science, University of Massachusetts, Amherst, MA 01003, USA,

Author for correspondence:
Samuel P. Hazen, hazen@bio.umass.edu
17
ORCID
Joshua H. Coomey, https://orcid.org/0000-0001-5866-6671
Pubudu P. Handakumbura, https://orcid.org/0000-0002-1708-8315
Kerrie Barry, https://orcid.org/0000-0002-8999-6785
Ji E. Lee, https://orcid.org/0000-0001-8325-1689
John P. Vogel, https://orcid.org/0000-0003-1786-2689
James J. Chambers, https://orcid.org/0000-0003-3883-8215
Samuel P. Hazen, https://orcid.org/0000-0001-9131-8994
26

27 **AUTHOR CONTRIBUTIONS**
28 JHC and SH conceived and designed the study. JHC, PH, MH-R, JJC developed techniques and
29 reagents. JHC, PH, GT, JM, NL, RK, KB, LL, JL, ZV, and RO acquired the data. JHC, KM, and
30 SH analyzed and interpreted the data. JHC, KM, IM, SH drafted the manuscript. SH agrees to
31 serve as the author responsible for contact and ensures communication.

ABSTRACT

Plant growth is mediated by the integration of internal and external cues, perceived by cells and transduced into a developmental program that gives rise to cell division, elongation, and wall thickening. Extra-, inter-, and intra- physical cellular forces contribute to this regulation. Across the plant kingdom thigmomorphogenesis is widely observed to alter plant morphology by reducing stem height and increasing stem diameter. The transcriptome is highly responsive to touch, including components of calcium signalling pathways and cell wall modification. One aspect of this cascade involves gibberellins and bZIP family transcription factors. Here, we present data connecting thigmomorphogenesis with secondary cell wall synthesis by gibberellin (GA) inactivation and bZIP translocation into the nucleus. *Brachypodium distachyon* SECONDARY WALL INTERACTING bZIP (SWIZ) protein translocated into the nucleus in response to mechanical stimulation. This translocation was mitigated by supplement with exogenous bioactive GA, and induced by chemical inhibition of GA synthesis. Misregulation of *SWIZ* expression resulted in plants hypersensitive to mechanical stimulation, with reduced height and increased secondary wall thickness. Classical touch responsive genes were upregulated in *B. distachyon* following touch, and we observe significant induction of the glycoside hydrolase 17 family, which may be indicative of a grass specific facet of thigmomorphogenesis. SWIZ binding to an E-box variant in exons and introns was associated with immediate activation followed by repression of gene expression. Thus, SWIZ is a transcriptional regulatory component of thigmomorphogenesis, which includes the thickening of secondary cell walls.

INTRODUCTION

Forces both internal and external to a cell influence growth. Turgor pressure in conjunction with anisotropic cell wall dynamics determine direction of growth and thus cell shape. Force perception between neighboring cells plays a critical role in the development and maintenance of tissue form and function, such as the lobed, interlocking, pavement cells on the leaf epidermis, or the developmental hotpots in the apical meristem (Hamant et al., 2008; Uyttewaal et al., 2012; Bidhendi et al., 2019). Specific inter-cell forces result in dynamic remodelling of the cortical cytoskeleton, with subsequent changes in cellulose microfibril alignment and alterations to other cell wall components such as pectin methyl esterification (Hamant et al., 2008; Uyttewaal et al., 2012; Bidhendi and Geitmann, 2018; Altartouri et al., 2019; Bidhendi et al., 2019). The classic hallmarks of touch responsive growth, or thigmomorphogenesis, include reduced plant height, increased radial growth in plants with a cambial meristem, increased branching, and delayed flowering time (Jaffe, 1973; Biro et al., 1980; Jaffe et al., 1980; Braam, 2004). These attributes have been leveraged by farmers for hundreds of years. As early as 1680 records show Japanese farmers tread on young wheat and barley seedlings to elicit increased branching, spikes per plant, and grain weight per plant, along with stronger roots (Iida, 2014). This practice, known as mugifumi, continues today with mechanized rollers.

Calcium is a critical cellular signalling molecule, and cytosolic Ca^{2+} fluctuations are induced by
mechanical stimuli and implicated in mechanosignaling (Monshausen et al., 2009; Szczegielniak
et al., 2012). Mechanical stimulus can significantly remodel gene expression (Braam and Davis,
1990; Braam, 2004; Lee et al., 2005). The so-called *TOUCH* (*TCH*) genes in *Arabidopsis*
thaliana, encode calmodulin (*AtTCH1/AtCaM2*), calmodulin-like proteins (*AtTCH2/AtCML24*,
AtTCH3/CML12), and a xyloglucan endotransglucosylase/hydrolase (*AtTCH4/AtXTH22*) (Braam
and Davis, 1990). Touch responsive gene expression patterns often overlap with other stimuli
such as dark, cold, and hormone treatment (Polisensky and Braam, 1996; Lee et al., 2005). In
addition to calcium binding and signaling, genes related to cell wall modification and a variety of
transcription factors and kinases are regulated by mechanical stimulus, as well as genes involved
in hormone homeostasis and signalling.

The dynamic control of bioactive cellular gibberellic acid (GA) is linked to
thigmomorphogenesis. GA is a growth promoting hormone derived through multiple enzymatic
steps from the diterpene pathway, with interconvertible bioactive and inactive species. The
bioactive forms of GA interact with protein receptors that regulate transcription factors
controlling growth. Mechanical force application results in the inactivation of cellular GA by
GA20 oxidases (Lange and Lange, 2015). The addition of exogenous bioactive GA to
mechanically stressed plants remediates their height phenotype, and mutations in *Atga20ox* do
not show thigmomorphogenic phenotypes under mechanical stress treatment (Lange and Lange,
2015). Recent work studying the translocation dynamics of several bZIP family transcription
factors has provided new links for calcium and GA signalling in mechanoperception signalling.

Group I bZIPs are also implicated in mechanosensing. AtVIP1 and related Group I bZIP proteins
translocate from the cytoplasm to the nucleus in response to a variety of biotic and abiotic
stimuli, including hypo-osmotic conditions (Tsugama et al., 2012; Tsugama et al., 2014;
Tsugama et al., 2016). This translocation appears to be dependent on protein phosphorylation,
either from MITOGEN ACTIVATED PROTEIN KINASE 3 during pathogen invasion, or via
calcium dependent protein kinases (CDPK). AtVIP1 interacts with calmodulins, the calcium
binding proteins involved in many calcium signalling events. The Group I bZIP *Nt REPRESSOR*
OF SHOOT GROWTH (*NtRSG*) in tobacco plays a role in maintaining GA homeostasis, wherein
it translocates from the cytoplasm to the nucleus in response to cellular bioactive GA levels
(Igarashi et al., 2001; Ishida et al., 2008; Fukazawa et al., 2010). Under standard conditions,
NtRSG is largely localized in the cytoplasm, and to some extent in the nucleus, where it directly
activates expression of *NtEnt-kaurene oxidase*, an enzyme at an early step in the GA
biosynthesis pathway. After cellular GAs were chemically inhibited, NtRSG translocated to the
nucleus and also activated *NtGA20oxidase1* (*NtGA20ox1*), an enzyme further downstream in the
GA biosynthesis pathway that is in part responsible for converting GA species to their bioactive
form (Fukazawa et al., 2010; Fukazawa et al., 2011). This suggests that nuclear-localized NtRSG
acts to promote bioactive GA synthesis in response to low GA conditions. As with AtVIP1,
NtRSG translocation relies on CDPK mediated phosphorylation (Ishida et al., 2008; Ito et al.,

2017). Both AtVIP1 and NtRSG associate with 14-3-3 proteins in the cytoplasm while
phosphorylated (Ishida et al., 2004; Ito et al., 2014; Ito et al., 2017; Tsugama et al., 2018).
Phosphatase activity causes NtRSG and the related AtbZIP29 to dissociate from the 14-3-3
protein and enter the nucleus (Van Leene et al., 2016). *NtRSG* is named for the dwarf phenotype
observed when a dominant negative form of the protein is overexpressed. In these dwarf plants,
there were reduced levels of bioactive GA and reduced elongation of stem internode cells
(Fukazawa et al., 2000).

The development of secondary cell walls is a distinct and critical developmental process in plant
growth. Following cellular elongation, some cells, particularly in vascular and structural tissues,
increase their mechanical strength by depositing a thick layer of rigid material between the
plasma membrane and primary cell wall. These secondary walls are made of crystalline cellulose
interwoven with hemicelluloses and phenolic lignin polymers. Although functionally similar,
secondary walls in monocotyledonous plants have key differences from those in eudicots,
including distinct hemicellulose chemistry and differences in lignin biosynthesis. Grasses also
produce mixed-linkage glucans (MLGs), a wall polysaccharide that is rarely found outside the
commelinid monocots (Coomey et al., 2020).The synthesis and deposition of secondary wall
polymers is tightly controlled spatially and temporally by a well defined transcriptional
regulatory network. Evidence in both eudicots and grasses suggest a system of feed-forward
loops regulating transcription of wall synthesizing enzymes, with NAC family transcription
factors activating wall synthesis genes as well as MYB family and other transcription factors that
further promote secondary wall synthesis (McCahill and Hazen, 2019). These networks are very
similar between grasses and eudicots, with some components in each that have yet to be
described in the other. There are currently no bZIP family members in any secondary wall
regulatory model.

Beyond differences at the chemical and regulatory levels, grasses employ a fundamentally
different growth mechanism than eudicots. Grasses do not have a cambium layer, and thus no
lateral meristem. Stem elongation comes from cellular division and elongation in a discrete series
of intercalary meristems, called nodes, with one internode region elongating and pushing up
subsequent nodes. Detailed studies of thigmomorphogenesis have been conducted almost
exclusively in eudicots However, recent work in the model cereal grass *Brachypodium*
distachyon shows general overlap with conventional dicot thigmomorphogenesis, albeit with
some notable differences, such as no change in stem diameter and increased flowering time
(Gladala-Kostarz et al., 2020).

Thigmomorphogenesis is a widely observed phenomenon that results in reduced height,
increased radial growth, and increased branching. The mechanisms behind this form of growth
are not yet fully understood, but involve aspects of hormone regulation, Ca^{2+} signalling, Group I
bZIP intracellular translocation, and changes in gene expression. Here we describe the function
of a *B. distachyon* bZIP transcription factor, SECONDARY WALL ASSOCIATED bZIP
(Bradi1g17700) and its role in touch response and cell wall biosynthesis.

RESULTS

SWIZ **is a Group I bZIP transcription factor and candidate cell wall regulator.**

To identify genes involved in the regulation of secondary cell wall thickening, transcript
abundance was measured in *B. distachyon* leaf, root, and stem tissue (Trabucco et al., 2013). A
gene annotated as a bZIP transcription factor, Bradi1g17700, was highly expressed in root and
stem relative to leaf (**Fig. 1A**). Bradi1g17700 is also a member of a 112 gene coexpression
network (**Supplemental Table S1**) that includes genes highly expressed in the peduncle (Sibout
et al., 2017). Phylogenetic analysis of Bradi1g17700, hereinafter referred to as *SECONDARY*
WALL INTERACTING bZIP (*SWIZ*), amino acid sequence shows it to be an ortholog of the *A.*
thaliana Group I bZIPs (Jakoby et al., 2002; Dröge-Laser et al., 2018), and closely related to
AtbZIP18 and *AtbZIP52* (**Fig. 1B**).

SWIZ translocates into the nucleus in response to mechanical stimulus

Given the translocation dynamics described for other Group I bZIPs, we hypothesized that SWIZ
protein may similarly translocate within the cell in response to mechanical force. To test this, the
roots of transgenic plants overexpressing either *SWIZ* fused to *GFP* (*SWIZ:GFP-OE*) or *GFP*
alone (*GFP-OE*) were observed following a mechanical stimulus (**Fig 2A**). In control plants,
GFP signal was present in both the cytosol and nucleus, which remained constant over the
imaging period (**Fig 2B**). GFP signal was mostly observed in the cytosol in *SWIZ:GFP-OE*
plants, but following mechanical stimulus, nuclear GFP signal increased substantially, reaching a
peak around 30 min post stimulus and returned to near basal levels by about 60 min (**Fig 2B**).

The dynamics and repeatability of SWIZ nuclear translocation were further investigated by
sequential stimulus treatments. Touch response to stimulus can saturate at a certain number of
treatments (Martin et al., 2010; Leblanc-Fournier et al., 2014; Moulia et al., 2015). To test if
SWIZ translocation dynamics varied after repeated treatments, we applied mechanical force to
SWIZ:GFP-OE roots at regular intervals. A second stimulus was given 90 min after the first, and
a third at 180 min. Following each mechanical stimulation, SWIZ consistently translocated from
cytoplasm to nucleus (**Fig 2D**). This suggests that SWIZ translocation dynamics are not impacted
by repeated stimulus events 90 min apart.

When conducting these translocation assays, root tissue in the field of view was treated with
mechanical force and nuclei in that region were tracked and quantified for fluorescence. To
determine if the SWIZ translocation signal is transmitted outside the specific region of stimulus,
two regions of the same *SWIZ:GFP-OE* root separated by 3 cm were simultaneously observed.
The stimulated region showed typical SWIZ:GFP nuclear signal accumulation and in the region
below no translocation was observed (**Fig 2C**). At 120 min, the treatments were reversed, with
the lower root region receiving a stimulus while the upper region was unperturbed. The lower
region showed SWIZ:GFP nuclear translocation while the upper region did not. Thus, the touch
stimulus perceived by SWIZ is localized to directly perturbed cells.

***B. distachyon* shows conventional thigmomorphogenic phenotypes of reduced height and**
increased branching.

Given the touch responsive nature of SWIZ protein translocation and the evidence suggesting a
role as a cell wall regulator, we hypothesized that SWIZ may play a role in mediating
thigmomorphogenesis. To investigate this, we first established a touch response assay and
assessed the thigmomorphogenic response in *B. distachyon*. Wildtype plants were perturbed with
a metal bar once every 90 min for two or three weeks (**Supplemental Fig 1**). After the treatment
period, all plants were allowed to recover and grow undisturbed until senescence (**Fig 3A**). Two
week stressed plants were significantly shorter than controls, and three week stressed plants were
shorter still (**Fig 3B**). Despite this difference in height, there was not a significant difference
between the groups in terms of aboveground biomass weight (**Fig 3C**). Three week stressed
plants had significantly more branches ($p < 0.05$), with a non-significant increase in two week
stressed plants (**Fig 3D**).

The effect of repeated mechanical stress on stem biology and interfascicular fiber cell walls was
assessed by making transverse sections of the peduncle and the third internode (**Fig 3E**). The
mechanical stress assay contacted the growing plant at ~5 cm above the soil, such that the third
internode region had already elongated prior to stress treatment, while the developing peduncle
was directly contacted by the metal bar during elongation. In this way, the third internode served
as an internal control tissue that did not experience direct mechanical perturbation. Sections were
stained with phloroglucinol-HCl to detect lignin in secondary cell walls. The degree of staining
and stem morphology in the third internode did not appear to be impacted by mechanical
perturbation, but the peduncle showed distinct morphological changes in the three week stressed
plants (**Fig 3E**). Interfascicular fiber cell walls were more lightly stained, and large, non-lignified
pockets containing starch granules were evident around the stem periphery, between the minor
bundles (**Supplemental Fig 2**). There was no significant difference in fiber wall thickness
between the internode regions or from differing touch treatments (**Fig 3F**).

Thigmomorphogenesis in *SWIZ* genetic reagents

To test the role of SWIZ in regulating thigmomorphogenesis and secondary cell wall
development we tested wildtype, *SWIZ-OE*, and *swiz-amiRNA* under two weeks of mechanical
perturbation as described above (**Supplemental Fig 3**). In control conditions, there were no
differences among genotypes . In mechanically stressed conditions, both *SWIZ-OE* and *swiz-*
amiRNA were significantly shorter than wildtype, and *swiz-amiRNA* had significantly more
branches (**Fig 4A-C**).

Transverse sections of the stem were made in the third elongated internode and the peduncle and
stained with phloroglucinol-HCl (**Fig 4D**). The third internode was not directly contacted during
mechanical perturbation, and as expected these sections did not show a difference in staining or
fiber wall thickness compared to control conditions. However, both *SWIZ-OE* and *swiz-amiRNA*
stained lighter than wildtype, independent of mechanical perturbation. The peduncle was directly

contacted during mechanical perturbation, and exhibited comparable lignin staining in both
SWIZ-OE and *swiz-amiRNA* to wildtype under touch perturbation, while remaining lighter
staining than wildtype in control conditions. *SWIZ-OE* touched peduncles showed significantly
thicker interfascicular fiber cell walls (**Fig 4E**). Thus, positive or negative modulation of *SWIZ*
transcript interferes with lignin accumulation in control conditions, and *SWIZ-OE* wall
thickening was more responsive to touch relative to no treatment.

Transcriptional response to touch
To assess the impact of *SWIZ* on gene expression in combination with touch treatment, we
measured transcript abundance by sequencing mRNA in the roots of wildtype, *SWIZ-OE*, and
swiz-amiRNA, just prior to touch and at 10, 30, and 60 min following touch treatment (**Fig 5A**).
Principal component analysis of transcript abundance shows the greatest amount of variance is
attributed to the first principal component, 53%, where samples cluster based on genotype (**Fig
5B**). The second principal component, which accounts for 15% of the variance, distinguished
between pre-touch and 10 min following touch, with the last two time points clustering similarly.
238
The wildtype transcriptome was massively remodelled in response to touch, with 8,902
transcripts differentially expressed ($q < 0.1$) at 10 min post touch, 5,682 transcripts at 30 min,
and 7,672 transcripts at 60 min (**Supplemental Tables 2-4**). Touch response studies in *A.*
thaliana describe similar changes that include numerous calmodulin (CAM), calmodulin-like
(CmL), and xyloglucan endo-transglycosylase/hydrolase (XTH) activity (**Fig 5C**)(Braam and
Davis, 1990; Lee et al., 2005). Based on homology with *TOUCH1* (*TCH1*), *TCH2*, and *TCH3*, 79
CAM and CmL genes were identified, the majority of which showed upregulation in wildtype
plants following touch treatment (**Supplemental Table 5**). Similarly, homology with *TCH4*
identified 37 *B. distachyon XTH* and *XTH-like* genes (Eklöf and Brumer, 2010). Of these, 8
genes were upregulated by touch in wildtype plants, including the two with the highest similarity
to *TCH4* (**Fig 5D**) (**Supplemental Table 6**). Since xyloglucans are in relatively low abundance
in grass cell walls compared to dicots, we also investigated the effect of touch on other families
of the glycosyl hydrolase superfamily (Tyler et al., 2010). In particular, GH17 genes were
differentially regulated following touch. The GH17 family is broadly described as modifying β-
1,3-glucans. We measured transcript abundance for 35 of the 53 annotated GH17 family
members, with 8, 11, and 16 members showing differential expression at 10, 30 and 60 min
respectively. Fisher's exact test determined significant enrichment of GH17 expression at time
60 ($p = 9.307e\text{-}3$) (**Fig 5D**, **Supplemental Table 7**). Online resources are available to explore the
RNA-seq data at https://hazenlab.shinyapps.io/swiztc/.
258
Cell wall genes are downregulated immediately following touch, then induced concurrent
with SWIZ nuclear translocation and more strongly in *SWIZ-OE*
Genes related to secondary cell wall synthesis were immediately downregulated in wildtype
plants following touch, but were subsequently upregulated (**Fig 6**). Xylan, cellulose, MLG, and
callose synthesis associated genes were upregulated 30 and 60 min following touch, while lignin

biosynthesis and BAHD acyltransferase genes were upregulated at 60 min. Network analysis of
gene expression patterns over time using the iDREM software showed that in *SWIZ-OE* plants,
cell wall genes were more rapidly induced (**Supplemental Fig 4A-C**). Prior to touch, nearly all
of the cell wall associated genes investigated were significantly downregulated relative to
wildtype, then significantly upregulated following touch (**Fig 6**). In *SWIZ-OE*, cell wall related
terms were enriched in paths that had stronger upregulation earlier than the same terms in
wildtype (**Supplemental Fig 4A-C**). Notably, aspects of cell wall polysaccharide biosynthesis,
particularly glucan biosynthesis, were significantly enriched in *SWIZ-OE* path D, while not
significantly enriched in wildtype or *swiz-amiRNA* (**Supplemental Table 8**). This is mirrored
when looking at specific gene expression, with primary wall associated CESAs, MLG, and
callose synthesis genes all significantly upregulated following touch in *SWIZ-OE*. In *swiz-*
amiRNA, cell wall related genes were generally not differentially expressed compared to
wildtype (**Fig 6**). Thus, cell wall genes were immediately repressed by touch, followed by
significant activation. That activation occurs more rapidly in *SWIZ-OE* plants and is associated
with the timing of SWIZ protein nuclear translocation.
279
SWIZ protein binding in the gene body is associated with dynamic changes in gene
expression
Next we investigated the direct binding targets of SWIZ protein by performing DNA affinity
purification sequencing (DAP-seq) with whole genomic DNA. In total, SWIZ interacted with
2,455 distinct positions in the genome. Those regions were significantly enriched for two
sequence motifs, (A/C)CAGNCTG(T/G) and (A/C)CAGCTG(T/G) that are variants of the E-box
motif (**Fig 7A**). A plurality of the binding sites were between the translational start and stop sites
with 21% of the peaks in introns and 21% in exons (**Fig 7B**). Fewer than 10% of the peaks
occurred in UTRs and the promoter (defined as 5 kb upstream of the 5 'UTR) and intergenic
regions each accounted for 25% of the binding sites. Thus, SWIZ protein preferentially bound to
an E-box like motif and often in the gene body (**Fig 7C**).
291
To investigate the effects of possible SWIZ binding on gene expression, we compared the genes
differentially expressed in the *SWIZ-OE* and *swiz-amiRNA* touch response RNA-seq time courses
with the promoter and gene body binding targets identified by the DAP-seq assay. Prior to touch,
genes with promoter-SWIZ protein interactions were most often downregulated in *SWIZ-OE* and
upregulated in *swiz-amiRNA* (**Table 1**). That same trend was observed for gene body targets in
untouched plants. However, some SWIZ targets behaved differently following touch. SWIZ
promoter binding targets were most often upregulated in *SWIZ-OE* in all three post-touch time
points. Thus, SWIZ promoter binding was coincident with repression, but following a touch
stimulus, activation was more prominent. However, this difference between untreated and
touched was not as pronounced in gene body targets, with more upregulated genes 10 min
following touch and more downregulated 30 and 60 min following touch. To further explore
these trends we conducted iDREM cluster analysis of the differentially expressed genes that

were also SWIZ protein targets (**Fig 7D**). For all genotypes, the network analysis revealed a pathway to increased gene expression by 60 min following touch. The network pathway trend for both promoter and gene binding targets in wildtype was immediate repression at 10 min followed by increased expression at 30 and 60 min post-stimulus. A unique pathway was observed among gene body binding targets in *SWIZ-OE*; transcript abundance was reduced following touch. Thus, SWIZ binding to a promoter or gene body was more strongly associated with increased expression with the exception of gene body targets in touched *SWIZ-OE* plants, which were more strongly repressed.

Among the genes differentially expressed in response to both touch, *SWIZ-OE* transgene, and bound by SWIZ in the first intron is the mixed glucan synthase *CSLF6* (**Fig 8A**). The transcript abundance of *GA2ox3*, which inactivates bioactive GA, was bound by SWIZ protein in both the gene body and 3'UTR and expression significantly increased 30 min after touch in *SWIZ-OE* (**Fig 8B**). An example of gene body binding repression in Bradi1g30252, the closest ortholog to a wheat transcription factor induced by hypoxia, *TaMYB1* (Lee 2007). This gene was touch induced in wildtype and repressed in *SWIZ-OE* with a binding site in the 5'UTR (**Fig 8C**). The NAC35 transcription factor Bradi2g28020 exhibited a similar expression to *MYB44* and SWIZ bound the promoter region (**Fig 8D**).

SWIZ translocation is sensitive to cellular GA levels, and may act in maintaining GA homeostasis following mechanical stimuli

Mechanical forces have been shown to inactivate cellular GA, and that cellular GA levels dictate cytosolic versus nuclear localizations of *N. benthamiana* Group I bZIPs similar to SWIZ (Fukazawa et al., 2010; Lange and Lange, 2015). To investigate this in relationship to SWIZ, *SWIZ:GFP-OE* roots were treated with bioactive GA or paclobutrazol, an inhibitor of GA synthesis. SWIZ:GFP signal amplitude was reduced in roots treated with bioactive GA, particularly at the 50 and 100 mM level. After 60 min the nuclear signal had returned to near pre-stimulus levels, and a second stimulus was applied (**Fig 9A**). This time all GA treated roots showed a similar reduction in nuclear signal, nearly half of the untreated control. These results suggest that the addition of exogenous GA reduces SWIZ translocation to the nucleus following mechanical force application.

Roots treated with 10 mM paclobutrazol did not show altered translocation dynamics compared to the untreated control, but roots treated with 50 or 100 mM paclobutrazol initially showed greater SWIZ:GFP nuclear localization, even prior to mechanical stimulus (**Fig 9B**). SWIZ:GFP signal in the 100 mM paclobutrazol treated roots was high prior to treatment and remained high even after the other nuclear signals diminished. This suggests that pharmacological reduction of bioactive GA levels triggers SWIZ nuclear translocation even without the application of mechanical force.

Given the role of GA in touch responsive bZIP translocation dynamics, we measured how GA
related gene expression changed in response touch. In wildtype following touch, GA synthesis
and GA activating genes *KS*, *KAO4*, and *GA20ox4* were downregulated, while the genes
encoding GA inactivating enzymes GA2ox4 and GA2ox9 were significantly upregulated (**Fig**
9C). In *SWIZ-OE*, *KS* was upregulated prior to touch and 10 min post touch. *GA20ox4*, a gene
responsible for synthesizing bioactive GA, was downregulated prior to touch in *SWIZ-OE* plants,
then significantly upregulated following touch. At 30 and 60 min *GA2ox4* and *GA2ox9* were
upregulated, along with the DELLA protein Rht-1 (**Fig 9C**). Here we show consistent patterns of
gene expression associated with the touch inactivation of GA and that SWIZ protein promotes
expression of GA synthesis and GA activation genes following mechanical perturbation.
352
353   **DISCUSSION**
354
Touch stimulus is generally an inhibitor of plant elongation growth, but promotes branching and
radial expansion (Jaffe, 1973; Braam, 2004; Chehab et al., 2009). The majority of this work has
been done in dicots, where increased radial growth has been associated with greater deposition
of secondary wall forming cells, particularly in woody species such as poplar (Biro et al., 1980;
Coutand et al., 2009; Börnke and Rocksch, 2018; Roignant et al., 2018; Niez et al., 2019). Our
understanding of thigmomorphogenesis in grasses is limited, and mostly in the agricultural
context of lodging (Shah et al., 2019). While these studies highlight the importance of stem
strength and associated cell wall defects, they do relatively little to elucidate the
mechanosensitive response. Recent work by Gladala-Kostarz et al (2020) describes grass touch
response to both wind and direct mechanical treatment, with an emphasis on cell walls and stem
anatomy. Touch treatment significantly increased lignin content, and also altered the relative
abundance of wall bound hydroxycinnamates. Wall polysaccharide content, particularly glucose
and galactose, also increased. Consistent with increased secondary wall lignification, touched
plants had increased wall stiffness and decreased saccharification. Our study of wildtype *B.*
distachyon plants with two or three weeks of regular touch revealed a decrease in plant height.
We also observed an increase in branching with increasing touch treatment. However, we did not
observe a change in total biomass at senescence, perhaps due to compensation by increased
branching.
373   Specific gene expression patterns have become molecular hallmarks of plant touch
response, most notably induction of the *TCH* and similar genes. Orthologs of wall modifying
XTHs and CAM/CML signalling genes are upregulated in response to mechanical stimulation
across species, and as we present here, in *B. distachyon*. Touch elicits major global changes in
gene expression, with 2.5% to 10% of the genes assayed in *A. thaliana* and poplar differentially
expressed (Lee et al., 2005; Pomiès et al., 2017; Van Moerkercke et al., 2019). In sorghum
(*Sorghum bicolor*), leaf sheath imposes mechanical constraints on emerging buds, and removing
this dramatically altered gene expression, with 42% of genes differentially expressed over a 9 h
period (Liu and Finlayson, 2019). Our observations of short term (<2 h) touch responses within
stress related pathways align with observations from poplar and *A. thaliana*, with rapid induction

of *TCH*, biotic, and abiotic stress related genes (Pomiès et al., 2017). Canonical touch responsive
genes such as the orthologs of *TCH1-4* and related *XTH* and *CML* genes were upregulated
immediately following stimulus. However, we also note the previously unreported touch-
induction of GH17 family genes, as well as *CSLF6*, suggesting a role for MLG and β-1,3 glucan
modification in grass touch response.
In both poplar and in our analysis of *B. distachyon* touch responses, touch-regulated
expression of secondary cell wall related transcripts differed between early and late timepoints.
Upon touch, we observed immediate repression of many key cell wall biosynthetic enzymes,
followed by significant upregulation one hour after stimulus. In poplar, no significant repression
of these transcripts was reported, but terms related to wood or cell wall synthesis were also not
enriched among touch-regulated transcripts until 24 or 72 hours after touch (Pomiès et al., 2017).
While differences in the method used to quantify gene expression, and in the temporal sampling
scheme complicate direct comparisons of plant touch response experiments, this may suggest
that delayed induction of secondary cell wall transcripts is a common feature of plant touch
responses.
398
Thigmomorphogenesis, GA physiology, and secondary cell walls
Our understanding of touch responsive gene expression is further informed by the dynamics of
SWIZ nuclear translocation. The timing of SWIZ nuclear accumulation following mechanical
stimulus was consistent with observations of *A. thaliana* Group I bZIPs (Tsugama et al., 2014;
Tsugama et al., 2016). However, by adopting a finer temporal sampling scheme, and quantifying
nuclear signal sooner after touch (2 vs. 30 min), our results clarify the rapid and immediate
nature of this translocation. Furthermore, we show that the speed and magnitude of SWIZ
nuclear accumulation is not diminished over three successive touch treatments. Notably, poplar
transcriptional responses to a second touch stimulus were significantly attenuated relative to a
single treatment, and in separate experiments, trees subjected to a daily touch treatment over four
days no longer showed touch-induced increases in their rates of radial growth (Martin et al.,
2010; Pomiès et al., 2017). Thus, if *B. distachyon* is likewise desensitized to repeated touch, the
high repeatability of SWIZ translocation might implicate a mechanism downstream of touch
perception and bZIP translocation in mediating that shift. Like the touch-responsive transcription
factor AtVIP1, SWIZ translocation occurred in the specific area of touch treatment, suggesting a
cellular rather than tissue level response. SWIZ also displays a highly similar pattern of GA
responsive translocation to the well-studied bZIP NtRSG (Ishida et al., 2004; Fukazawa et al.,
2010; Fukazawa et al., 2011). Exogenous bioactive GA dampened SWIZ nuclear translocation,
while inhibition of GA synthesis with paclobutrazol induced translocation, even prior to touch.
Furthermore, like NtRSG, SWIZ appears to both respond to and regulate GA bioactivity. In
wildtype, GA inactivating enzymes GA2ox4 and GA2ox9 were upregulated following touch,
consistent with previous reports (Lange and Lange, 2015). In *SWIZ-OE* we also observe touch
induced expression of *GA20ox4*, a GA activating enzyme, and as with NtRSG, this response can
restore bioactive GA following touch induced degradation (Ishida et al., 2004; Fukazawa et al.,

2010; Fukazawa et al., 2011). Altered GA pathways are a well described mechanism for reduced
stem elongation (Lange and Lange, 2006; Nagai et al., 2020), and dwarfing has been described
for dominant negative mutants of Group I bZIPs *NtRSG* and *OsRF2A* (Fukazawa et al., 2000;
Dai et al., 2003). *SWIZ-OE* and *swiz-amiRNA* were both significantly shorter following touch
treatment, suggesting that GA misregulation may contribute to this thigmomorphogenic
phenotype. Another component of this reduced elongation may be increased secondary cell wall
deposition, as this could terminate elongation. Prior to touch, almost the entire monolignol
biosynthetic pathway, as well as laccase genes that radicalize monolignols, and BAHD
acyltransferases that add hydroxycinnamic acids to wall polymers, are significantly
downregulated in *SWIZ-OE* plants. Following touch, most of these genes become significantly
upregulated. This activity is consistent with our observations of increased lignin staining in
touched *SWIZ-OE* peduncles, and aligns with the compositional data presented by Gladala-
Kostarz et al (2020). This activation may be a result of direct binding of SWIZ to modulate
expression, or an indirect effect from another transcriptional regulator such as the NAC
transcription factor *SWN5*, capable of activating the full developmental program for secondary
cell wall synthesis (Valdivia et al., 2013), which is significantly upregulated in *SWIZ-OE*
following touch.
440
While the E-box (CANNTG) was initially described as a bHLH binding motif, many bZIP
groups in *A. thaliana* have been shown to bind similar sequences. The two closets *A. thaliana*
orthologs of SWIZ , bZIP18 and 52, both bind (A/C)CAG(G/C)(T/C), while VIP1 binds
(A/C)CAGCT(G/A) (O'Malley et al., 2016). Using the same technique, we identified the SWIZ
binding motifs as (A/C)CAGNCTG(T/G) and (A/C)CAGCTG(T/G). Both of these sequences are
similar to those bound by *A. thaliana* orthologs, although the presence of an ambiguous
nucleotide in the core of one variant is not previously reported for other Group I bZIPs. Among
genes bound by SWIZ *in vitro,* we observed both activation and repression following touch,
suggesting a complex regulatory function for SWIZ. Furthermore, SWIZ direct binding sites
were found in both promoter regions and in the body of genes and most often in genes activated
by touch. SWIZ gene body binding targets tended to be repressed in *SWIZ-OE* plants following
touch, without either being clearly associated with up or down regulation. AtVIP1 and AtbZIP29
have both been described as activators, particularly of genes related to biotic and abiotic stress
response, cell cycle control, vascular development, and cell wall biogenesis (Yin et al., 1997;
Ringli and Keller, 1998; Pitzschke et al., 2009; Van Leene et al., 2016). AtbZIP18 is suggested
to have repressive roles (Tsugama et al., 2012; Gibalová et al., 2017).

bZIPs are known to homo- and heterodimerize through their leucine zipper domains (Schütze et
al., 2008), and in *A. thaliana* the combinatorial interactions of different bZIP groups have been
fairly well described (Deppmann et al., 2004; Ehlert et al., 2006; Grigoryan and Keating, 2006;
Weltmeier et al., 2006; Schütze et al., 2008). These interactions can have a synergistic effect on
transcriptional activity, and can result in unique binding interactions (Schütze et al., 2008; Van
Leene et al., 2016). It is conceivable that this heterodimerization may explain some aspects of

our results, such as why so many of the cell wall genes that are upregulated in *SWIZ-OE* after

touch are significantly downregulated prior to touch. One hypothesis is that SWIZ, as with many

other bZIPs, can regulate targets as a heterodimer. The abundance of SWIZ protein in the

cytoplasm in *SWIZ-OE* plants prior to touch may act as a sink, sequestering interacting partners

that would otherwise be regulating targets independent of a touch stimulus. Similarly, redundant

or compensatory function among similar bZIPs may explain why the *swiz-amiRNA* plants did not

show significant and opposite phenotypes from the *SWIZ-OE*. Further experimentation is needed

to elucidate this bZIP regulatory network in response to touch and other stimuli.

Conclusions

Proteins orthologous to SWIZ in other species have been implicated in cell wall development

and remodelling, but not cell wall thickening. SWIZ appears to control stem height and cell wall

thickening in response to mechanical stimuli, which regulates SWIZ translocation into the

nucleus by acting on the cellular bioactive GA pool. Touch has been shown to decrease cellular

bioactive GA levels in *A. thaliana*, and Group I bZIPs are known to translocate to the nucleus in

response to touch and to low bioactive GA. Direct mechanical stimulation and chemical

inhibition of GA biosynthesis both caused SWIZ to translocate from the cytoplasm into the

nucleus. This translocation can be reduced by supplementing with exogenous bioactive GA. GA

biosynthesis and signalling genes were upregulated in *SWIZ-OE* lines following touch, indicating

that SWIZ may act in managing GA homeostasis following mechanical perturbation. Altering

SWIZ function may impact this GA homeostasis and result in reduced elongation following

touch. Touch significantly remodeled the *B. distachyon* transcriptome, with notable changes in

wall polysaccharide biosynthetic gene expression not previously reported. Enhanced SWIZ

function through overexpression amplified the expression of cell wall related genes and synthesis

of interfascicular fiber walls in touched tissues. Greater secondary wall synthesis in response to

touch may also explain reduced stem elongation in *SWIZ-OE*. Together, the evidence presented

here connects mechanosensitive bZIP translocation with dynamic changes in cellular GA levels

that have been observed separately in other systems and shows a link between touch response

and secondary cell wall transcriptional regulation (**Fig 10**).

MATERIALS AND METHODS

Phylogenetic analysis

Protein sequences described for *A. thaliana*, *B. distachyon*, and *O. sativa* (Liu and Chu, 2015) as

Group II bZIPs were selected and searched against the most recent genome annotations in

Phytozome v12.1 (https://phytozome.jgi.doe.gov). The *Nicotiana tabacum* homologs NtRSGa

and NtRSGb were also added to the analysis. Protein sequences were aligned using the MAFFT

service for multiple sequence alignment with the iterative refinement method L-INS-I (Katoh et

al., 2019). The alignment was used to construct the phylogenetic tree using a maximum

likelihood analysis with a bootstrap resampling value of 1000 in W-IQ-TREE (Trifinopoulos et

al., 2016). All of the proteins included in the phylogenetic analysis are described in
Supplemental File 1.

Plant transformation

Overexpression and artificial microRNA transgenes and transgenic plants were constructed as
previously described (Handakumbura et al., 2013). The full length coding sequence of
Bradi1g17700 was amplified from cDNA and cloned into the pOL001 ubigate ori1 binary
expression vector (Handakumbura et al., 2018) to make the *SWIZ-OE* trangene. The coding
sequence lacking the stop codon was amplified and fused in frame with the coding sequence of
Aequorea victoria enhanced GFP in the pOL001 vector to generate *SWIZ:GFP-OE*. Artificial
microRNA constructs were generated as previously described (Handakumbura et al., 2013) to
include a targeting sequence specific to the Bradi1g17700 transcript,
CAGCCTGAACCTTCCCAGTCC. This sequence was identified using the WMD3 webtool
(Schwab et al., 2006; Warthmann et al., 2008). The modified amiRNA sequence was cloned into
pOL001 for overexpression.

Translocation assay

Seeds were surface sterilized and grown vertically on 1X MS media,, pH 5.7, without sucrose for
6d at 28°C in the dark. After 6 d, seedlings were moved to treatment plates containing 1X MS
media, pH 5.7, plus varying concentrations (0, 10, 50, or 100mM) of GA4, paclobutrazol. Roots
were positioned such that they were completely submerged in media and relatively flat along the
imaging plane After a 6h incubation, plants were stimulated and imaged.

All observations were made on a Nikon A1R scanning confocal microscope using a Plan Apo
10x 0.5NA objective. Root areas to be observed were located by eye using transmitted light and
then confirmed under confocal conditions using a 488 nm laser and green filtered PMT detector.
The X, Y, and Z coordinate stage locations of each region were programmed into the Nikon NIS
Elements 5.2 Advanced Research V6 software package for automated imaging. After all target
regions were programmed, 30 min of imaging began pre-treatment, with images captured every 2
min.

To elicit the touch response, the observed root region was gently probed for 5 sec with a blunt
metal probe while observing through the eyepiece. Images were captured for 60-90 min post
treatment. For experiments with multiple stimulus events, the timelapse sequence was paused
and roots were probed as described for the relevant stimulus events.

Analysis of GFP signal was done using the Nikon NIS Elements Advanced Research V5
software package. For each time series, the frame showing maximal nuclear signal was selected
and used as a reference point. Using the General Analysis tool, a channel for GFP signal was
established and thresholded for intensity and particle size to identify the nuclear regions. These
regions were added to the timelapse image series as a binary layer and then converted to static
regions of interest for quantification. The GFP intensity under each nuclear region of interest was
calculated for the course of the timelapse and the average signal from each nucleus was plotted
for intensity over time.

Plant growth

Accession Bd21-3 was used for all experiments. Seeds were stratified on wet paper towel
wrapped in foil to exclude light for 10 days at 4°C before being planted in Promix BX potting
mix in SC10 Ray Leach Cone-tainers (Stuewe & Sons Inc). Plants were grown in a Percival
PGC-15 growth chamber with day/night conditions of 20h light at 24°C and 4h dark at 18°C,
respectively.

Thigmomatic construction and operation

The Thigmomatic is a basic robotic device that sweeps plants with a metal bar at regular
intervals to elicit a touch response. The device was constructed from aluminum V-Slot linear rail
(Openbuilders Partstore, Monreoville, NJ) and bracket joints for the upright supports (20x20
mm), cross bars (20x20 mm), and tracks (20x40 mm). Two gantry carts ride along the 20x40 mm
V-Slot linear rails, connected by a 6.35 mm diameter metal rod bolted to the carts. Their
movement is powered by a belt driven linear actuator system using a NEMA 17 stepper motor
with a 12V 18W AC/DC power supply. The stepper motor provides fine spatial control over the
gantry cart position with bi-directional motion. Motor function is controlled by a Raspberry Pi
3B microcomputer equipped with a stepper motor HAT (Adafruit Industries, New York). The
Thigmomatic was programmed to cover a specified distance in one direction once every 90 min.

Transverse stem sections and histology

The main stem of senesced plants was taken and the internode of interest removed and embedded
in 8% agarose. A Leica VT1000 Vibratome was used to make 55 μm thick transverse sections.
Histochemical staining was carried out using phloroglucinol-HCl as previously described (Matos
et al., 2013). Images were obtained at 4, 10, and 20X using a Nikon Eclipse E200MV R light
microscope and PixeLINK 3 MP camera. Transverse sections imaged at 20x were used for cell
wall thickness measurements. Interfascicular fiber cells separated by one cell layer from the
mestome cells on the phloem side of major vascular bundles were targeted for measurement.
Using ImageJ, lines were drawn across two walls of adjoining cells. The resulting line length
was divided by two to give one cell wall width. Approximately 15 measurements were made for
each plant.

RNA extraction and quantification and analysis

Seedlings were grown for 6 d on vertical 1X MS agar plates. Touch treatment was performed as described above using a metal probe along the entire length of the root. Untouched samples were collected immediately before touch treatment began, and touched samples were collected after the designated time. Three roots were pooled per biological replicate, and RNA was extracted using the Qiagen RNeasy Plant Mini Kit with on-column DNA digestion with RNase-free DNase I (Qiagen). Strand-specific libraries were prepared using the Illumina TruSeq kit. Libraries were sequenced using Illumina technology. Sequences were processed as previously described (MacKinnon et al., 2020) and then checked for differential expression using the likelihood ratio test. Specific contrasts (i.e., wildtype vs *SWIZ-OE*) were identified and compared using the Wald test. Statistical enrichment of gene families was assessed using Fisher's exact test. Raw read data were deposited in the European Nucleotide Archive for public access (Accession no.: XYZ).

DNA affinity purification sequencing

DNA affinity purification was carried out as previously described (Handakumbura et al., 2018). In brief, transcription factor coding sequences were HALO tagged and mixed with Bd21 genomic DNA for *in vitro* binding. Protein-DNA was crosslinked, fragmented, immunoprecipitated using the HALO antibody, barcoded, and sequenced. Reads were mapped back to the Bd21 genome using HiSat2 (Kim et al., 2015) to identify binding target loci. Peak calling and motif analysis was done using HOMER v4.10 (Hypergeometric Optimization of Motif EnRichment) suite (Heinz et al., 2010). Motif enrichment was calculated against the hypergeometric distribution; the significance threshold was set to $p < 0.05$. The nearest annotated gene to a bound peak was used for GO analysis.

Gene Ontology analysis

Phytozome was used to find orthologs for all *B. distachyon* v3.1 genes as the reciprocal best match to *A. thaliana* TAIRv10 protein sequences. Arabidopsis gene identifiers were submitted to g:Profiler (Raudvere et al., 2019) for KEGG and Wiki pathway enrichment analysis.

iDREM network analysis

Probabilistic graphical models that predict diverging gene expression paths and the points at which regulators may influence those paths were generated using iDREM (Ding et al., 2018). Briefly, this software applies an input-output hidden Markov model to time course gene expression data overlaid with static regulatory information, in this case SWIZ protein-DNA interactions identified from DAP-seq. GO analysis, described above, was also applied to the gene sets in each path identified by iDREM.

SUPPLEMENTAL MATERIALS

Supplemental Table S1. Cluster 56 from network analysis of gene expression atlas (Sibout et al. 2017) includes Bradi1g17700.

Supplemental Table S2. Differentially expressed genes in wildtype Bd21-3 root tissue 10 min after touch treatment.

Supplemental Table S3. Differentially expressed genes in wildtype Bd21-3 root tissue 30 min after touch treatment.

Supplemental Table S4. Differentially expressed genes in wildtype Bd21-3 root tissue 60 min after touch treatment.

Supplemental Table S5. Calmodulin and calmodulin-like touch responsive gene expression in wildtype Bd21-3 root tissue.

Supplemental Table S6. XTH and XTH-like touch responsive gene expression in wildtype Bd21-3 root tissue.

Supplemental Table S7. Glycoside hydrolase touch responsive gene expression in wildtype Bd21-3 root tissue.

Supplemental Table S8. GO analysis of genes in iDREM identified pathways derived from differential gene expression following touch in wildtype, swiz-amiRNA, and SWIZ-OE root tissue.

Supplemental Figure 1. Thigmomatic, an automated system to administer touch stimulus. (A) Overview of Thigmomatic inside a Percival PGC-15 growth chamber showing linear rail based frame (1), gantry carts (2), NEMA17 stepper motor (3), 12V 18W AC/DC power supply (4), and Raspberry Pi 3b microcomputer (5). (B) Thigmomatic making contact with a *Brachypodium distachyon* plant.

Supplemental Figure 2. Regions in the periphery of the stem without secondary cell walls containing starch granules. Lugol's iodine stained starch granules of peduncle transverse sections of *Brachypodium distachyon* treated with three weeks of mechanical stress. Arrow indicates an example starch granule. Scale bar = 100 μm.

Supplemental Figure 3. Description of *SWIZ* transgenic reagents. (A) Diagram of *SWIZ* overexpression transgenes. The *ZmUbi* promoter was used to drive expression of the *SWIZ* coding sequence, either alone or fused in frame with *eGFP*. (B) Artificial microRNA construct for targeted *SWIZ* transcript knockdown. The maize ubiquitin promoter was used to drive expression of an amiRNA that has homology to the first *SWIZ* exon and forms a hairpin after transcription. The 21-mer nucleotide sequence used for target specificity is listed, with red

postions showing hairpin mismatches that are predicted to interact with the DICER complex.
Relative level of *SWIZ* transcript measured by RT-qPCR in whole stem tissue was collected 1 d
after inflorescence emergence in *SWIZ:GFP-OE* (C) and *swiz-amiRNA* (D) lines. LB, left border;
ZmUbi prom, maize ubiquitin promoter; Hyg, hygromycin phosphotransferase gene; NOS,
nopaline synthase terminator; RB, right border. ns: $p > 0.05$, : *: $p < 0.05$.

Supplemental Figure 4. Network analysis of differential gene expression paths and enriched GO
terms. (A) Path determinations from iDREM time course analysis. Each line represents a set of
genes with similar expression level patterns over the time course relative to time 0, pre-touch.
(B) Heatmap of selected GO term enrichment of genes from iDREM path analysis. Color
indicates qvalue score, with a cutoff of 0.1. Terms without a colored block are not significantly
enriched in that path.

Supplemental File 1. Gene names and amino acid sequences used in the SWIZ bZIP Group I
phylogenetic analysis.

ACCESSION NUMBERS

AtbZIP18 (At2g40620), AtbZIP29 (At4g38900), AtbZIP52 (At1g06850), AtTCH1/AtCaM2
(At2g41110), AtTCH2/AtCML24 (At5g37770), AtTCH3/CML12 (At2g41100),
AtTCH4/AtXTH22 (At5g57560), CAD1 (Bradi3g17920), CESA4 (Bradi4g28350), COMT6
(Bradi3g16530), CSLF6 (Bradi3g16307), GA2ox3 (Bradi2g50280), MYB44 (Bradi1g30252),
NAC35 (Bradi2g28020), NtRSGa (Niben101Scf01150g00005), NtRSGb
(Niben101Scf02191g00014), SWIZ (Bradi1g17700), VIP1 (At1g43700).

ACKNOWLEDGEMENTS

This work was supported by the Office of Science, Biological and Environmental Research,
Department of Energy (DE-FG02-08ER64700DE and DE-SC0006641) and The National
Science Foundation Division of Integrative Organismal Systems (NSF IOS-1558072) to S.P.H..
The work conducted by the US DOE Joint Genome Institute is supported by the Office of
Science of the US Department of Energy under Contract no. DE-AC02-05CH11231. The
microscopy data was gathered in the Light Microscopy Facility and Nikon Center of Excellence
at the Institute for Applied Life Sciences, UMass Amherst with support from the Massachusetts
Life Sciences Center.

Table 1. The count of differentially expressed genes relative to untouched wildtype plants that were SWIZ DAP-seq targets in either the promoter or gene body regions.

Genotype	Time after touch	Promoter		Gene body	
		Up	Down	Up	Down
	(min)	percent (count)			
SWIZ-OE	0	42.0 (103)	58.0 (142)	48.0 (286)	52.0 (310)
	10	57.0 (73)	43.0 (55)	53.0 (167)	47.0 (148)
	30	59.4 (79)	40.6 (54)	41.2 (136)	58.8 (194)
	60	64.2 (77)	35.8 (43)	45.7 (128)	54.3 (152)
swiz-amiRNA	0	67.1 (47)	32.9 (23)	60.0 (78)	40.0 (52)
	10	56.1 (37)	43.9 (29)	55.9 (76)	44.1 (60)
	30	65.0 (13)	35.0 (7)	29.4 (5)	70.6 (12)
	60	53.8 (7)	46.2 (6)	27.8 (5)	72.2 (13)

FIGURE LEGENDS

Figure 1. SWIZ is a Group I bZIP highly expressed in maturing stem and root. (A) *SWIZ* transcript abundance in *Brachypodium distachyon* leaf, root, and stem tissue measured by microarray. Mean +/- standard deviation of three biological replicates. (B) Phylogeny analysis of amino acid sequences from *B. distachyon* (green), *Oryza sativa* (blue), *Arabidopsis thaliana* (red), *Nicotiana tabacum* (orange) shows *SWIZ* (black dot). The phylogeny was reconstructed using maximum likelihood analysis with a bootstrap resampling value of 1000. The numbers labeled on the branches are the posterior probability to support each clade. The protein sequences, full locus IDs, and plant species with genome source are provided in Supplemental File 1.

Figure 2. SWIZ translocates to the nucleus in response to mechanical stimulus, specifically in regions directly stimulated. (A) Image of *SWIZ:GFP-OE* and *GFP-OE* roots prior to stimulus and 30 min post stimulus. Roots were observed immediately following mechanical perturbation. (B) Quantification of nuclear signal. n=14-20 nuclei. (C) SWIZ translocation occured in the local area of the stimulus. At 30 min, stimulus was applied to an upper region of the root, while at 120 min it was applied to a lower region approximately 3 cm below. n=109,

184 nuclei respectively for upper and lower regions. (D) *SWIZ:GFP-OE* roots were imaged by
confocal microscopy with stimulus applied in the field of view at 0, 90, and 180 min. n=126
nuclei. (B-D) Images were taken every 2 min. Nuclear GFP signal was quantified in selected
nuclei at each time point. The average nuclear GFP signal is represented by the line with error
bars indicating standard error of the mean.

Figure 3. *Brachypodium distachyon* displays classic thigmomorphogenic phenotypes. One-
week-old wildtype plants were treated with touch stimulus every 90 min. (A) Left to right, plants
that experienced no stress, two weeks stress, three weeks stress and were then allowed to
recover, imaged one week after the end of treatment (B) height, (C) aboveground non-grain
biomass weight, and (D) branch number were measured at senescence. ns: $p > 0.05$, *: $p < 0.05$,
n = 5 plants per treatment. (E) Transverse sections of the peduncle or third internode were take
from control, 2 week stressed, and 3 week stressed plants and stained with phloroglucinol-HCl.
(F) Quantification of interfascicular fiber wall thickness. Scale bar = 100 µm. n = 3 plants per
treatment.

Figure 4. *SWIZ* transgenes enhance thigmomorphogenic response. Wildtype, *SWIZ-OE*, and
swiz-amiRNA plants were placed under control conditions or received two weeks of mechanical
stress every 90 min. After senescence, plants were phenotyped for (A) height, (B) branch
number, and (C) aboveground biomass weight. ns: $p > 0.05$, *: $p < 0.05$. n = 5 plants per
treatment. (D) The main stem was collected and transverse sections made of the peduncle and
third elongated internode. Sections were stained with phloroglucinol-HCl. Scale bar = 100 µm.
(E) Quantification of interfascicular fiber wall thickness. n = 4 to 6 plants per genotype, per
treatment.

Figure 5. Transcriptome analysis of touch response in *Brachypodium distachyon* roots. (A)
Root tissue was sampled just prior to touch (t=0), and at 10, 30, and 60 min following touch
treatment in wildtype, *SWIZ-OE*, and *swiz-amiRNA*. (B) Principal component analysis of
variance in gene expression across samples shows the greatest difference corresponding to
genotype, with the second greatest corresponding to time after touch. (C) Canonical, as well as
novel touch responsive genes are upregulated in *B. distachyon* following touch. Closest orthologs
of the *Arabidopsis thaliana TOUCH* genes encoding calmodulin-like (CML) and xyloglucan
endo-transglycosylase/hydrolases (XTH) are upregulated following touch, as are previously
unreported members of the glycosyl hydrolase 17 (GH17) family. Full list of gene expression for
TOUCH and glycoside hydrolases defined by Tyler et al. (2010) in Supplemental Tables 5,6,and
7.

Figure 6. Gene expression analysis of cell wall related genes. Log fold-change of gene
expression in wildtype, *SWIZ-OE*, and *swiz-amiRNA* relative to wildtype expression at time 0,
pre-touch. Bar color indicates class of cell wall gene. Error bars indicate standard deviation of

three biological replicates. Asterisks represent significantly differentially expressed genes, q-
value < 0.1 for Wald test of each time and genotype compared to wild type at time 0.
728
Figure 7. DNA affinity purification sequencing to determine SWIZ binding sites. (A) Top
two most statistically enriched sequence motifs in SWIZ binding sites. (B) Distribution of
binding sites across genomic features, relative to primary transcripts of the *Brachypodium*
distachyon annotation v 3.1. (C) Relative distribution of binding sites centered on the
transcriptional start site (TSS, blue dashed line), transcriptional termination site (TTS, red dashed
line) represents the average length of all annotated transcripts, approximately 4.5 kb away from
the TSS. (D) Path determinations from iDREM time course analysis of differentially expressed
genes that also have DAP-seq binding sites. Each line represents a set of genes with similar
expression level patterns over the time course relative to time 0, pre-touch.
738
Figure. 8 SWIZ binding targets differentially expressed in response to touch and SWIZ.
Gene expression over time of selected genes with SWIZ binding sites. Line graphs are the
average transcript abundance of three biological replicates for each time point. Binding site
determined as peaks of sequence alignment. Scale bar unit is bases. Direction of transcription is
shown with arrows on the gene model, 5' and 3' UTRs are depicted by narrowed rectangles on
the gene model.
745
Figure 9. SWIZ translocation is dependent on bioactive GA levels and impacts GA related
gene expression. (A) SWIZ translocation to the nucleus is dampened by the addition of
exogenous GA4. (B) SWIZ translocation to the nucleus is enhanced by the addition of
paclobutrazol (PBZ), a GA synthesis inhibitor. The average nuclear GFP signal is represented by
the trend line, with error bars indicating standard deviation of the mean. n= 85-125 (A) and n=
64-142 (B) nuclei per treatment. Gene expression analysis of GA related genes. (C) Log fold
change of gene expression in wildtype samples, relative to their expression at time 0, pre touch.
(D) Log fold change of gene expression in SWIZ OE and swiz amiRNA samples, relative to
their expression in wildtype at the same time points. Bar color indicates relevant function in GA
synthesis and signalling.
756
757   **Figure 10. SWIZ regulates cell wall synthesis and GA homeostasis in response to**
758   **mechanical stimulus.** Prior to touch, SWIZ is localized to the cytosol and partially in the
759   nucleus. The bioactive GA pool is depleted in response to touch, triggering SWIZ to translocate
760   to the nucleus where it positively regulates cell wall synthesis and GA activating enzymes.
761
REFERENCES

Altartouri B, Bidhendi AJ, Tani T, Suzuki J, Conrad C, Chebli Y, Liu N, Karunakaran C,
Scarcelli G, Geitmann A (2019) Pectin chemistry and cellulose crystallinity govern

pavement cell morphogenesis in a multi-step mechanism. Plant Physiol **181**: 127–141

Bidhendi AJ, Altartouri B, Gosselin FP, Geitmann A (2019) Mechanical stress initiates and
sustains the morphogenesis of wavy leaf epidermal cells. Cell Rep **28**: 1237–1250.e6

Bidhendi AJ, Geitmann A (2018) Finite element modeling of shape changes in plant cells.
Plant Physiol **176**: 41–56

Biro RL, Hunt ER, Erner Y, Jaffe MJ (1980) Thigmomorphogenesis: changes in cell division
and elongation in the internodes of mechanically-perturbed or ethrel-treated bean plants.
Ann Bot **45**: 655–664

Börnke F, Rocksch T (2018) Thigmomorphogenesis – Control of plant growth by mechanical
stimulation. Sci Hortic **234**: 344–353

Braam J (2004) In touch: plant responses to mechanical stimuli. New Phytol **165**: 373–389

Braam J, Davis RW (1990) Rain-, wind-, and touch-induced expression of calmodulin and
calmodulin-related genes in Arabidopsis. Cell **60**: 357–364

Chehab EW, Eich E, Braam J (2009) Thigmomorphogenesis: a complex plant response to
mechano-stimulation. J Exp Bot **60**: 43–56

Coomey JH, Sibout R, Hazen SP (2020) Grass secondary cell walls, *Brachypodium distachyon*
as a model for discovery. New Phytol **227**: 1709–1724

Coutand C, Martin L, Leblanc-Fournier N, Decourteix M, Julien J-L, Moulia B (2009)
Strain mechanosensing quantitatively controls diameter growth and PtaZFP2 gene
expression in poplar. Plant Physiol **151**: 223–232

Dai S, Petruccelli S, Ordiz MI, Zhang Z, Chen S, Beachy RN (2003) Functional analysis of
RF2a, a rice transcription factor. J Biol Chem **278**: 36396–36402

Deppmann CD, Acharya A, Rishi V, Wobbes B, Smeekens S, Taparowsky EJ, Vinson C
(2004) Dimerization specificity of all 67 B-ZIP motifs in *Arabidopsis thaliana*: a
comparison to *Homo sapiens* B-ZIP motifs. Nucleic Acids Res **32**: 3435–3445

Ding J, Hagood JS, Ambalavanan N, Kaminski N, Bar-Joseph Z (2018) iDREM: Interactive
visualization of dynamic regulatory networks. PLoS Comput Biol **14**: e1006019

Dröge-Laser W, Snoek BL, Snel B, Weiste C (2018) The Arabidopsis bZIP transcription factor
family-an update. Curr Opin Plant Biol **45**: 36–49

Ehlert A, Weltmeier F, Wang X, Mayer CS, Smeekens S, Vicente-Carbajosa J, Dröge-

Laser W (2006) Two-hybrid protein-protein interaction analysis in Arabidopsis protoplasts: establishment of a heterodimerization map of group C and group S bZIP transcription factors. Plant J **46**: 890–900

Eklöf JM, Brumer H (2010) The XTH gene family: an update on enzyme structure, function, and phylogeny in xyloglucan remodeling. Plant Physiol **153**: 456–466

Fukazawa J, Nakata M, Ito T, Matsushita A, Yamaguchi S, Takahashi Y (2011) bZIP transcription factor RSG controls the feedback regulation of NtGA20ox1 via intracellular localization and epigenetic mechanism. Plant Signal Behav **6**: 26–28

Fukazawa J, Nakata M, Ito T, Yamaguchi S, Takahashi Y (2010) The transcription factor RSG regulates negative feedback of NtGA20ox1 encoding GA 20-oxidase. Plant J **62**: 1035–1045

Fukazawa J, Sakai T, Ishida S, Yamaguchi I, Kamiya Y, Takahashi Y (2000) Repression of shoot growth, a bZIP transcriptional activator, regulates cell elongation by controlling the level of gibberellins. Plant Cell **12**: 901–915

Gibalová A, Steinbachová L, Hafidh S, Bláhová V, Gadiou Z, Michailidis C, Műller K, Pleskot R, Dupľáková N, Honys D (2017) Characterization of pollen-expressed bZIP protein interactions and the role of ATbZIP18 in the male gametophyte. Plant Reprod **30**: 1–17

Gladala-Kostarz A, Doonan JH, Bosch M (2020) Mechanical stimulation in *Brachypodium distachyon*: implications for fitness, productivity and cell wall properties. Plant Cell Environ **43**: 1314–1330

Grigoryan G, Keating AE (2006) Structure-based prediction of bZIP partnering specificity. J Mol Biol **355**: 1125–1142

Hamant O, Heisler MG, Jönsson H, Krupinski P, Uyttewaal M, Bokov P, Corson F, Sahlin P, Boudaoud A, Meyerowitz EM, et al (2008) Developmental patterning by mechanical signals in Arabidopsis. Science **322**: 1650–1655

Handakumbura P, Matos D, Osmont K, Harrington M, Heo K, Kafle K, Kim S, Baskin T, Hazen S (2013) Perturbation of *Brachypodium distachyon CELLULOSE SYNTHASE A4* or *7* results in abnormal cell walls. BMC Plant Biol **13**: 131

Handakumbura PP, Brow K, Whitney IP, Zhao K, Sanguinet KA, Lee SJ, Olins J, Romero-Gamboa SP, Harrington MJ, Bascom CJ, et al (2018) *SECONDARY WALL ASSOCIATED MYB1* is a positive regulator of secondary cell wall thickening in *Brachypodium distachyon* and is not found in the Brassicaceae. Plant J **96**: 485–699

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H,
Glass CK (2010) Simple combinations of lineage-determining transcription factors prime
cis-regulatory elements required for macrophage and B cell identities. Mol Cell **38**: 576–
589

Igarashi D, Ishida S, Fukazawa J, Takahashi Y (2001) 14-3-3 proteins regulate intracellular
localization of the bZIP transcriptional activator RSG. Plant Cell **13**: 2483–2497

Iida H (2014) Mugifumi, a beneficial farm work of adding mechanical stress by treading to
wheat and barley seedlings. Front Plant Sci **5**: 453

Ishida S, Fukazawa J, Yuasa T, Takahashi Y (2004) Involvement of 14-3-3 signaling protein
binding in the functional regulation of the transcriptional activator REPRESSION OF
SHOOT GROWTH by gibberellins. Plant Cell **16**: 2641–2651

Ishida S, Yuasa T, Nakata M, Takahashi Y (2008) A tobacco calcium-dependent protein
kinase, CDPK1, regulates the transcription factor REPRESSION OF SHOOT GROWTH in
response to gibberellins. Plant Cell **20**: 3273–3288

Ito T, Ishida S, Oe S, Fukazawa J, Takahashi Y (2017) Autophosphorylation affects substrate-
binding affinity of tobacco Ca2+-dependent protein kinase1. Plant Physiol **174**: 2457–2468

Ito T, Nakata M, Fukazawa J, Ishida S, Takahashi Y (2014) Scaffold function of Ca2+-
dependent protein kinase: tobacco Ca2+-DEPENDENT PROTEIN KINASE1 transfers 14-
3-3 to the substrate REPRESSION OF SHOOT GROWTH after phosphorylation. Plant
Physiol **165**: 1737–1750

Jaffe MJ (1973) Thigmomorphogenesis: The response of plant growth and development to
mechanical stimulation : With special reference to Bryonia dioica. Planta **114**: 143–157

Jaffe MJ, Biro R, Bridle K (1980) Thigmomorphogenesis: Calibration of the parameters of the
sensory function in beans. Physiol Plant **49**: 410–416

Jakoby M, Weisshaar B, Dröge-Laser W, Vicente-Carbajosa J, Tiedemann J, Kroj T,
Parcy F, bZIP Research Group (2002) bZIP transcription factors in Arabidopsis. Trends
Plant Sci **7**: 106–111

Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Brief Bioinform **20**: 1160–1166

Kim D, Langmead B, Salzberg SL (2015) HISAT: a fast spliced aligner with low memory
requirements. Nat Methods **12**: 357

Lange MJ, Lange T (2006) Gibberellin biosynthesis and the regulation of plant development.
Plant Biol **8**: 281–290

Lange MJP, Lange T (2015) Touch-induced changes in Arabidopsis morphology dependent on
gibberellin breakdown. Nat Plants **1**: 14025

Leblanc-Fournier N, Martin L, Lenne C, Decourteix M (2014) To respond or not to respond,
the recurring question in plant mechanosensitivity. Front Plant Sci **5**: 401

Lee D, Polisensky DH, Braam J (2005) Genome-wide identification of touch- and darkness-
regulated Arabidopsis genes: a focus on calmodulin-like and XTH genes. New Phytol **165**:
429–444

Liu R, Finlayson SA (2019) Sorghum tiller bud growth is repressed by contact with the
overlying leaf. Plant Cell Environ **42**: 2120–2132

Liu X, Chu Z (2015) Genome-wide evolutionary characterization and analysis of bZIP
transcription factors and their expression profiles in response to multiple abiotic stresses in
Brachypodium distachyon. BMC Genomics **16**: 227

MacKinnon KJ-M, Cole BJ, Yu C, Coomey JH, Hartwick NT, Remigereau M-S, Duffy T,
Michael TP, Kay SA, Hazen SP (2020) Changes in ambient temperature are the prevailing
cue in determining *Brachypodium distachyon* diurnal gene regulation. New Phytol **227**:
1649–1667

Martin L, Leblanc-Fournier N, Julien J-L, Moulia B, Coutand C (2010) Acclimation
kinetics of physiological and molecular responses of plants to multiple mechanical loadings.
J Exp Bot **61**: 2403–2412

Matos DA, Whitney IP, Harrington MJ, Hazen SP (2013) Cell walls and the developmental
anatomy of the *Brachypodium distachyon* stem internode. PLoS One **8**: e80640

McCahill IW, Hazen SP (2019) Regulation of cell wall thickening by a medley of mechanisms.
Trends Plant Sci **24**: 853–866

Monshausen GB, Bibikova TN, Weisenseel MH, Gilroy S (2009) Ca2+ regulates reactive
oxygen species production and pH during mechanosensing in Arabidopsis roots. Plant Cell
21: 2341–2356

Moulia B, Coutand C, Julien J-L (2015) Mechanosensitive control of plant growth: bearing the
load, sensing, transducing, and responding. Front Plant Sci **6**: 52

Nagai K, Mori Y, Ishikawa S, Furuta T, Gamuyao R, Niimi Y, Hobo T, Fukuda M, Kojima

M, Takebayashi Y, et al (2020) Antagonistic regulation of the gibberellic acid response
during stem growth in rice. Nature **584**: 109–114

Niez B, Dlouha J, Moulia B, Badel E (2019) Water-stressed or not, the mechanical acclimation
is a priority requirement for trees. Trees **33**: 279–291

O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti
A, Ecker JR (2016) Cistrome and epicistrome features shape the regulatory DNA
landscape. Cell **165**: 1280–1292

Pitzschke A, Djamei A, Teige M, Hirt H (2009) VIP1 response elements mediate mitogen-
activated protein kinase 3-induced stress gene expression. Proc Natl Acad Sci U S A **106**:
18414–18419

Polisensky DH, Braam J (1996) Cold-shock regulation of the Arabidopsis TCH genes and the
effects of modulating intracellular calcium levels. Plant Physiol **111**: 1271–1279

Pomiès L, Decourteix M, Franchel J, Moulia B, Leblanc-Fournier N (2017) Poplar stem
transcriptome is massively remodelled in response to single or repeated mechanical stimuli.
BMC Genomics **18**: 300

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) g:Profiler: a
web server for functional enrichment analysis and conversions of gene lists. Nucleic Acids
Res **47**: W191–W198

Ringli C, Keller B (1998) Specific interaction of the tomato bZIP transcription factor VSF-1
with a non-palindromic DNA sequence that controls vascular gene expression. Plant Mol
Biol **37**: 977–988

Roignant J, Badel É, Leblanc-Fournier N, Brunel-Michac N, Ruelle J, Moulia B,
Decourteix M (2018) Feeling stretched or compressed? The multiple mechanosensitive
responses of wood formation to bending. Ann Bot **121**: 1151–1161

Schütze K, Harter K, Chaban C (2008) Post-translational regulation of plant bZIP factors.
Trends Plant Sci **13**: 247–255

Schwab R, Ossowski S, Riester M, Warthmann N, Weigel D (2006) Highly specific gene
silencing by artificial microRNAs in Arabidopsis. Plant Cell **18**: 1121–1133

Shah L, Yahya M, Shah SMA, Nadeem M, Ali A, Ali A, Wang J, Riaz MW, Rehman S, Wu
W, et al (2019) Improving Lodging Resistance: Using Wheat and Rice as Classical
Examples. Int J Mol Sci. doi: 10.3390/ijms20174211

Sibout R, Proost S, Hansen BO, Vaid N, Giorgi FM, Ho-Yue-Kuang S, Legée F, Cézart L,
Bouchabké-Coussa O, Soulhat C, et al (2017) Expression atlas and comparative
coexpression network analyses reveal important genes involved in the formation of lignified
cell wall in Brachypodium distachyon. New Phytol **215**: 1009–1025

Szczegielniak J, Borkiewicz L, Szurmak B, Lewandowska-Gnatowska E, Statkiewicz M,
Klimecka M, Cieśla J, Muszyńska G (2012) Maize calcium-dependent protein kinase
(ZmCPK11): local and systemic response to wounding, regulation by touch and components
of jasmonate signaling. Physiol Plant **146**: 1–14

Trabucco GM, Matos DA, Lee SJ, Saathoff A, Priest H, Mockler TC, Sarath G, Hazen SP
(2013) Functional characterization of cinnamyl alcohol dehydrogenase and caffeic acid *O*-
methyltransferase in *Brachypodium distachyon*. BMC Biotechnol **13**: 61

Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ (2016) W-IQ-TREE: a fast online
phylogenetic tool for maximum likelihood analysis. Nucleic Acids Res **44**: W232–W235

Tsugama D, Liu S, Fujino K, Takano T (2018) Calcium signalling regulates the functions of
the bZIP protein VIP1 in touch responses in *Arabidopsis thaliana*. Ann Bot **122**: 1219–1229

Tsugama D, Liu S, Takano T (2012) A bZIP protein, VIP1, is a regulator of osmosensory
signaling in Arabidopsis. Plant Physiol **159**: 144–155

Tsugama D, Liu S, Takano T (2016) The bZIP protein VIP1 Is involved in touch responses in
Arabidopsis roots. Plant Physiol **171**: 1355–1365

Tsugama D, Liu S, Takano T (2014) Analysis of functions of VIP1 and its close homologs in
osmosensory responses of *Arabidopsis thaliana*. PLoS One **9**: e103930

Tyler L, Bragg JN, Wu J, Yang X, Tuskan GA, Vogel JP (2010) Annotation and comparative
analysis of the glycoside hydrolase genes in *Brachypodium distachyon*. BMC Genomics **11**:
600

Uyttewaal M, Burian A, Alim K, Landrein B, Borowska-Wykręt D, Dedieu A, Peaucelle A,
Ludynia M, Traas J, Boudaoud A, et al (2012) Mechanical stress acts via katanin to
amplify differences in growth rate between adjacent cells in Arabidopsis. Cell **149**: 439–451

Valdivia ER, Herrera MT, Gianzo C, Fidalgo J, Revilla G, Zarra I, Sampedro J (2013)
Regulation of secondary wall synthesis and cell death by NAC transcription factors in the
monocot *Brachypodium distachyon*. J Exp Bot **64**: 1333–1343

Van Leene J, Blomme J, Kulkarni SR, Cannoot B, De Winne N, Eeckhout D, Persiau G,
Van De Slijke E, Vercruysse L, Vanden Bossche R, et al (2016) Functional

characterization of the Arabidopsis transcription factor bZIP29 reveals its role in leaf and
root development. J Exp Bot **67**: 5825–5840

Van Moerkercke A, Duncan O, Zander M, Šimura J, Broda M, Vanden Bossche R, Lewsey
MG, Lama S, Singh KB, Ljung K, et al (2019) A MYC2/MYC3/MYC4-dependent
transcription factor network regulates water spray-responsive gene expression and
jasmonate levels. Proc Natl Acad Sci U S A **116**: 23345–23356

Warthmann N, Chen H, Ossowski S, Weigel D, Hervre P (2008) Highly specific gene
silencing by artificial miRNAs in rice. PLoS One **3**: e1829

Weltmeier F, Ehlert A, Mayer CS, Dietrich K, Wang X, Schütze K, Alonso R, Harter K,
Vicente-Carbajosa J, Dröge-Laser W (2006) Combinatorial control of Arabidopsis
proline dehydrogenase transcription by specific heterodimerisation of bZIP transcription
factors. EMBO J **25**: 3133–3143

Yin Y, Zhu Q, Dai S, Lamb C, Beachy RN (1997) RF2a, a bZIP transcriptional activator of the
phloem-specific rice tungro bacilliform virus promoter, functions in vascular development.
EMBO J **16**: 5247–5259

Figure 1. SWIZ is a Group I bZIP highly expressed in maturing stem and root. (A) *SWIZ* transcript abundance in *Brachypodium distachyon* leaf, root, and stem tissue measured by microarray. Mean +/- standard deviation of three biological replicates. (B) Phylogeny analysis of amino acid sequences from *B. distachyon* (green), *Oryza sativa* (blue), *Arabidopsis thaliana* (red), *Nicotiana tabacum* (orange) shows *SWIZ* (black dot). The phylogeny was reconstructed using maximum likelihood analysis with a bootstrap resampling value of 1000. The numbers labeled on the branches are the posterior probability to support each clade. The protein sequences, full locus IDs, and plant species with genome source are provided in Supplemental File 1.

Figure 2. SWIZ translocates to the nucleus in response to mechanical stimulus, specifically in regions directly stimulated. (A) Image of *SWIZ:GFP-OE* and *GFP-OE* roots prior to stimulus and 30 min post stimulus. Roots were observed immediately following mechanical perturbation. (B) Quantification of nuclear signal. n=14-20 nuclei. (C) SWIZ translocation occured in the local area of the stimulus. At 30 min, stimulus was applied to an upper region of the root, while at 120 min it was applied to a lower region approximately 3 cm below. n=109, 184 nuclei respectively for upper and lower regions. (D) *SWIZ:GFP-OE* roots were imaged by confocal microscopy with stimulus applied in the field of view at 0, 90, and 180 min. n=126 nuclei. (B-D) Images were taken every 2 min. Nuclear GFP signal was quantified in selected nuclei at each time point. The average nuclear GFP signal is represented by the line with error bars indicating standard error of the mean.

Figure 3. *Brachypodium distachyon* displays classic thigmomorphogenic phenotypes. One-week-old wildtype plants were treated with touch stimulus every 90 min. (A) Left to right, plants that experienced no stress, two weeks stress, three weeks stress and were then allowed to recover, imaged one week after the end of treatment (B) height, (C) aboveground non-grain biomass weight, and (D) branch number were measured at senescence. ns: $p > 0.05$, *: $p < 0.05$, n = 5 plants per treatment. (E) Transverse sections of the peduncle or third internode were take from control, 2 week stressed, and 3 week stressed plants and stained with phloroglucinol-HCl. (F) Quantification of interfascicular fiber wall thickness. Scale bar = 100 μm. n = 3 plants per treatment.

Figure 4. *SWIZ* transgenes enhance thigmomorphogenic response. Wildtype, *SWIZ-OE*, and *swiz-amiRNA* plants were placed under control conditions or received two weeks of mechanical stress every 90 min. After senescence, plants were phenotyped for (A) height, (B) branch number, and (C) aboveground biomass weight. ns: *p* > 0.05, *: *p* < 0.05. n = 5 plants per treatment. (D) The main stem was collected and transverse sections made of the peduncle and third elongated internode. Sections were stained with phloroglucinol-HCl. Scale bar = 100 μm. (E) Quantification of interfascicular fiber wall thickness. n = 4 to 6 plants per genotype, per treatment.

Figure 5. Transcriptome analysis of touch response in *Brachypodium distachyon* roots. (A) Root tissue was sampled just prior to touch (t=0), and at 10, 30, and 60 min following touch treatment in wildtype, *SWIZ-OE*, and *swiz-amiRNA*. (B) Principal component analysis of variance in gene expression across samples shows the greatest difference corresponding to genotype, with the second greatest corresponding to time after touch. (C) Canonical, as well as novel touch responsive genes are upregulated in *B. distachyon* following touch. Closest orthologs of the *Arabidopsis thaliana TOUCH* genes encoding calmodulin-like (CML) and xyloglucan endo-transglycosylase/hydrolases (XTH) are upregulated following touch, as are previously unreported members of the glycosyl hydrolase 17 (GH17) family. Full list of gene expression for *TOUCH* and glycoside hydrolases defined by Tyler et al. (2010) in Supplemental Tables 5,6,and 7.

Figure 6. Gene expression analysis of cell wall related genes. Log fold-change of gene expression measured by RNA-seq in wildtype, *SWIZ-OE*, and *swiz-amiRNA* relative to wildtype expression at time 0, pre-touch. Bar color indicates class of cell wall gene. Error bars indicate standard deviation of three biological replicates. Asterisks represent significantly differentially expressed genes, q-value < 0.1 for Wald test of each time and genotype compared to wild type at time 0.

Figure 7. DNA affinity purification sequencing to determine SWIZ binding sites. (A) Top two most statistically enriched sequence motifs in SWIZ binding sites. (B) Distribution of binding sites across genomic features, relative to primary transcripts of the *Brachypodium distachyon* annotation v 3.1. (C) Relative distribution of binding sites centered on the transcriptional start site (TSS, blue dashed line), transcriptional termination site (TTS, red dashed line) represents the average length of all annotated transcripts, approximately 4.5 kb away from the TSS. (D) Path determinations from iDREM time course analysis of differentially expressed genes that also have DAP-seq binding sites. Each line represents a set of genes with similar expression level patterns over the time course relative to time 0, pre-touch.

Figure. 8 SWIZ binding targets differentially expressed in response to touch and SWIZ. Gene expression over time of selected genes with SWIZ binding sites. Line graphs are the average transcript abundance of three biological replicates for each time point. Binding site determined as peaks of sequence alignment. Scale bar unit is bases. Direction of transcription is shown with arrows on the gene model, 5' and 3' UTRs are depicted by narrowed rectangles on the gene model.

Figure 9. SWIZ translocation is dependent on bioactive GA levels and impacts GA related gene expression. (A) SWIZ translocation to the nucleus is dampened by the addition of exogenous GA4. (B) SWIZ translocation to the nucleus is enhanced by the addition of paclobutrazol (PBZ), a GA synthesis inhibitor. The average nuclear GFP signal is represented by the trend line, with error bars indicating standard deviation of the mean. n= 85-125 (A) and n= 64-142 (B) nuclei per treatment. Gene expression analysis of GA related genes. (C) Log fold change of gene expression relative to their expression at time 0, pre touch. Bar color indicates relevant function in GA synthesis and signalling.

Figure 10. SWIZ regulates cell wall synthesis and GA homeostasis in response to mechanical stimulus. Prior to touch, SWIZ is localized to the cytosol and partially in the nucleus. The bioactive GA pool is depleted in response to touch, triggering SWIZ to translocate to the nucleus where it positively regulates cell wall synthesis and GA activating enzymes.

Parsed Citations

Altartouri B, Bidhendi AJ, Tani T, Suzuki J, Conrad C, Chebli Y, Liu N, Karunakaran C, Scarcelli G, Geitmann A (2019) Pectin chemistry and cellulose crystallinity govern pavement cell morphogenesis in a multi-step mechanism. Plant Physiol 181: 127–141
  Google Scholar: Author Only Title Only Author and Title

Bidhendi AJ, Altartouri B, Gosselin FP, Geitmann A (2019) Mechanical stress initiates and sustains the morphogenesis of wavy leaf epidermal cells. Cell Rep 28: 1237–1250.e6
  Google Scholar: Author Only Title Only Author and Title

Bidhendi AJ, Geitmann A (2018) Finite element modeling of shape changes in plant cells. Plant Physiol 176: 41–56
  Google Scholar: Author Only Title Only Author and Title

Biro RL, Hunt ER, Erner Y, Jaffe MJ (1980) Thigmomorphogenesis: changes in cell division and elongation in the internodes of mechanically-perturbed or ethrel-treated bean plants. Ann Bot 45: 655–664
  Google Scholar: Author Only Title Only Author and Title

Börnke F, Rocksch T (2018) Thigmomorphogenesis – Control of plant growth by mechanical stimulation. Sci Hortic 234: 344–353
  Google Scholar: Author Only Title Only Author and Title

Braam J (2004) In touch: plant responses to mechanical stimuli. New Phytol 165: 373–389
  Google Scholar: Author Only Title Only Author and Title

Braam J, Davis RW (1990) Rain-, wind-, and touch-induced expression of calmodulin and calmodulin-related genes in Arabidopsis. Cell 60: 357–364
  Google Scholar: Author Only Title Only Author and Title

Chehab EW, Eich E, Braam J (2009) Thigmomorphogenesis: a complex plant response to mechano-stimulation. J Exp Bot 60: 43–56
  Google Scholar: Author Only Title Only Author and Title

Coomey JH, Sibout R, Hazen SP (2020) Grass secondary cell walls, Brachypodium distachyon as a model for discovery. New Phytol 227: 1709–1724
  Google Scholar: Author Only Title Only Author and Title

Coutand C, Martin L, Leblanc-Fournier N, Decourteix M, Julien J-L, Moulia B (2009) Strain mechanosensing quantitatively controls diameter growth and PtaZFP2 gene expression in poplar. Plant Physiol 151: 223–232
  Google Scholar: Author Only Title Only Author and Title

Dai S, Petruccelli S, Ordiz MI, Zhang Z, Chen S, Beachy RN (2003) Functional analysis of RF2a, a rice transcription factor. J Biol Chem 278: 36396–36402
  Google Scholar: Author Only Title Only Author and Title

Deppmann CD, Acharya A, Rishi V, Wobbes B, Smeekens S, Taparowsky EJ, Vinson C (2004) Dimerization specificity of all 67 B-ZIP motifs in Arabidopsis thaliana: a comparison to Homo sapiens B-ZIP motifs. Nucleic Acids Res 32: 3435–3445
  Google Scholar: Author Only Title Only Author and Title

Ding J, Hagood JS, Ambalavanan N, Kaminski N, Bar-Joseph Z (2018) iDREM: Interactive visualization of dynamic regulatory networks. PLoS Comput Biol 14: e1006019
  Google Scholar: Author Only Title Only Author and Title

Dröge-Laser W, Snoek BL, Snel B, Weiste C (2018) The Arabidopsis bZIP transcription factor family-an update. Curr Opin Plant Biol 45: 36–49
  Google Scholar: Author Only Title Only Author and Title

Ehlert A, Weltmeier F, Wang X, Mayer CS, Smeekens S, Vicente-Carbajosa J, Dröge-Laser W (2006) Two-hybrid protein-protein interaction analysis in Arabidopsis protoplasts: establishment of a heterodimerization map of group C and group S bZIP transcription factors. Plant J 46: 890–900
  Google Scholar: Author Only Title Only Author and Title

Eklöf JM, Brumer H (2010) The XTH gene family: an update on enzyme structure, function, and phylogeny in xyloglucan remodeling. Plant Physiol 153: 456–466
  Google Scholar: Author Only Title Only Author and Title

Fukazawa J, Nakata M, Ito T, Matsushita A, Yamaguchi S, Takahashi Y (2011) bZIP transcription factor RSG controls the feedback regulation of NtGA20ox1 via intracellular localization and epigenetic mechanism. Plant Signal Behav 6: 26–28
  Google Scholar: Author Only Title Only Author and Title

Fukazawa J, Nakata M, Ito T, Yamaguchi S, Takahashi Y (2010) The transcription factor RSG regulates negative feedback of NtGA20ox1 encoding GA 20-oxidase. Plant J 62: 1035–1045
  Google Scholar: Author Only Title Only Author and Title

Fukazawa J, Sakai T, Ishida S, Yamaguchi I, Kamiya Y, Takahashi Y (2000) Repression of shoot growth, a bZIP transcriptional activator, regulates cell elongation by controlling the level of gibberellins. Plant Cell 12: 901–915

Google Scholar: Author Only Title Only Author and Title

Gibalová A, Steinbachová L, Hafidh S, Bláhová V, Gadiou Z, Michailidis C, Műller K, Pleskot R, Dupľáková N, Honys D (2017) Characterization of pollen-expressed bZIP protein interactions and the role of ATbZIP18 in the male gametophyte. Plant Reprod 30: 1–17
  Google Scholar: Author Only Title Only Author and Title

Gladala-Kostarz A, Doonan JH, Bosch M (2020) Mechanical stimulation in Brachypodium distachyon: implications for fitness, productivity and cell wall properties. Plant Cell Environ 43: 1314–1330
  Google Scholar: Author Only Title Only Author and Title

Grigoryan G, Keating AE (2006) Structure-based prediction of bZIP partnering specificity. J Mol Biol 355: 1125–1142
  Google Scholar: Author Only Title Only Author and Title

Hamant O, Heisler MG, Jönsson H, Krupinski P, Uyttewaal M, Bokov P, Corson F, Sahlin P, Boudaoud A, Meyerowitz EM, et al (2008) Developmental patterning by mechanical signals in Arabidopsis. Science 322: 1650–1655
  Google Scholar: Author Only Title Only Author and Title

Handakumbura P, Matos D, Osmont K, Harrington M, Heo K, Kafle K, Kim S, Baskin T, Hazen S (2013) Perturbation of Brachypodium distachyon CELLULOSE SYNTHASE A4 or 7 results in abnormal cell walls. BMC Plant Biol 13: 131
  Google Scholar: Author Only Title Only Author and Title

Handakumbura PP, Brow K, Whitney IP, Zhao K, Sanguinet KA, Lee SJ, Olins J, Romero-Gamboa SP, Harrington MJ, Bascom CJ, et al (2018) SECONDARY WALL ASSOCIATED MYB1 is a positive regulator of secondary cell wall thickening in Brachypodium distachyon and is not found in the Brassicaceae. Plant J 96: 485–699
  Google Scholar: Author Only Title Only Author and Title

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK (2010) Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. Mol Cell 38: 576–589
  Google Scholar: Author Only Title Only Author and Title

Igarashi D, Ishida S, Fukazawa J, Takahashi Y (2001) 14-3-3 proteins regulate intracellular localization of the bZIP transcriptional activator RSG. Plant Cell 13: 2483–2497
  Google Scholar: Author Only Title Only Author and Title

Iida H (2014) Mugifumi, a beneficial farm work of adding mechanical stress by treading to wheat and barley seedlings. Front Plant Sci 5: 453
  Google Scholar: Author Only Title Only Author and Title

Ishida S, Fukazawa J, Yuasa T, Takahashi Y (2004) Involvement of 14-3-3 signaling protein binding in the functional regulation of the transcriptional activator REPRESSION OF SHOOT GROWTH by gibberellins. Plant Cell 16: 2641–2651
  Google Scholar: Author Only Title Only Author and Title

Ishida S, Yuasa T, Nakata M, Takahashi Y (2008) A tobacco calcium-dependent protein kinase, CDPK1, regulates the transcription factor REPRESSION OF SHOOT GROWTH in response to gibberellins. Plant Cell 20: 3273–3288
  Google Scholar: Author Only Title Only Author and Title

Ito T, Ishida S, Oe S, Fukazawa J, Takahashi Y (2017) Autophosphorylation affects substrate-binding affinity of tobacco Ca2+-dependent protein kinase1. Plant Physiol 174: 2457–2468
  Google Scholar: Author Only Title Only Author and Title

Ito T, Nakata M, Fukazawa J, Ishida S, Takahashi Y (2014) Scaffold function of Ca2+-dependent protein kinase: tobacco Ca2+-DEPENDENT PROTEIN KINASE1 transfers 14-3-3 to the substrate REPRESSION OF SHOOT GROWTH after phosphorylation. Plant Physiol 165: 1737–1750
  Google Scholar: Author Only Title Only Author and Title

Jaffe MJ (1973) Thigmomorphogenesis: The response of plant growth and development to mechanical stimulation : With special reference to Bryonia dioica. Planta 114: 143–157
  Google Scholar: Author Only Title Only Author and Title

Jaffe MJ, Biro R, Bridle K (1980) Thigmomorphogenesis: Calibration of the parameters of the sensory function in beans. Physiol Plant 49: 410–416
  Google Scholar: Author Only Title Only Author and Title

Jakoby M, Weisshaar B, Dröge-Laser W, Vicente-Carbajosa J, Tiedemann J, Kroj T, Parcy F, bZIP Research Group (2002) bZIP transcription factors in Arabidopsis. Trends Plant Sci 7: 106–111
  Google Scholar: Author Only Title Only Author and Title

Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. Brief Bioinform 20: 1160–1166
  Google Scholar: Author Only Title Only Author and Title

Kim D, Langmead B, Salzberg SL (2015) HISAT: a fast spliced aligner with low memory requirements. Nat Methods 12: 357

Google Scholar: Author Only Title Only Author and Title

Lange MJ, Lange T (2006) Gibberellin biosynthesis and the regulation of plant development. Plant Biol 8: 281–290

Lange MJP, Lange T (2015) Touch-induced changes in Arabidopsis morphology dependent on gibberellin breakdown. Nat Plants 1: 14025

Leblanc-Fournier N, Martin L, Lenne C, Decourteix M (2014) To respond or not to respond, the recurring question in plant mechanosensitivity. Front Plant Sci 5: 401

Lee D, Polisensky DH, Braam J (2005) Genome-wide identification of touch- and darkness-regulated Arabidopsis genes: a focus on calmodulin-like and XTH genes. New Phytol 165: 429–444

Liu R, Finlayson SA (2019) Sorghum tiller bud growth is repressed by contact with the overlying leaf. Plant Cell Environ 42: 2120–2132

Liu X, Chu Z (2015) Genome-wide evolutionary characterization and analysis of bZIP transcription factors and their expression profiles in response to multiple abiotic stresses in Brachypodium distachyon. BMC Genomics 16: 227

MacKinnon KJ-M, Cole BJ, Yu C, Coomey JH, Hartwick NT, Remigereau M-S, Duffy T, Michael TP, Kay SA, Hazen SP (2020) Changes in ambient temperature are the prevailing cue in determining Brachypodium distachyon diurnal gene regulation. New Phytol 227: 1649–1667

Martin L, Leblanc-Fournier N, Julien J-L, Moulia B, Coutand C (2010) Acclimation kinetics of physiological and molecular responses of plants to multiple mechanical loadings. J Exp Bot 61: 2403–2412

Matos DA, Whitney IP, Harrington MJ, Hazen SP (2013) Cell walls and the developmental anatomy of the Brachypodium distachyon stem internode. PLoS One 8: e80640

McCahill IW, Hazen SP (2019) Regulation of cell wall thickening by a medley of mechanisms. Trends Plant Sci 24: 853–866

Monshausen GB, Bibikova TN, Weisenseel MH, Gilroy S (2009) Ca2+ regulates reactive oxygen species production and pH during mechanosensing in Arabidopsis roots. Plant Cell 21: 2341–2356

Moulia B, Coutand C, Julien J-L (2015) Mechanosensitive control of plant growth: bearing the load, sensing, transducing, and responding. Front Plant Sci 6: 52

Nagai K, Mori Y, Ishikawa S, Furuta T, Gamuyao R, Niimi Y, Hobo T, Fukuda M, Kojima M, Takebayashi Y, et al (2020) Antagonistic regulation of the gibberellic acid response during stem growth in rice. Nature 584: 109–114

Niez B, Dlouha J, Moulia B, Badel E (2019) Water-stressed or not, the mechanical acclimation is a priority requirement for trees. Trees 33: 279–291

O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A, Ecker JR (2016) Cistrome and epicistrome features shape the regulatory DNA landscape. Cell 165: 1280–1292

Pitzschke A, Djamei A, Teige M, Hirt H (2009) VIP1 response elements mediate mitogen-activated protein kinase 3-induced stress gene expression. Proc Natl Acad Sci U S A 106: 18414–18419

Polisensky DH, Braam J (1996) Cold-shock regulation of the Arabidopsis TCH genes and the effects of modulating intracellular calcium levels. Plant Physiol 111: 1271–1279

Pomiès L, Decourteix M, Franchel J, Moulia B, Leblanc-Fournier N (2017) Poplar stem transcriptome is massively remodelled in response to single or repeated mechanical stimuli. BMC Genomics 18: 300

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) g:Profiler: a web server for functional enrichment analysis and conversions of gene lists. Nucleic Acids Res 47: W191–W198

Google Scholar: Author Only Title Only Author and Title

Ringli C, Keller B (1998) Specific interaction of the tomato bZIP transcription factor VSF-1 with a non-palindromic DNA sequence that controls vascular gene expression. Plant Mol Biol 37: 977–988
  Google Scholar: Author Only Title Only Author and Title

Roignant J, Badel É, Leblanc-Fournier N, Brunel-Michac N, Ruelle J, Moulia B, Decourteix M (2018) Feeling stretched or compressed? The multiple mechanosensitive responses of wood formation to bending. Ann Bot 121: 1151–1161
  Google Scholar: Author Only Title Only Author and Title

Schütze K, Harter K, Chaban C (2008) Post-translational regulation of plant bZIP factors. Trends Plant Sci 13: 247–255
  Google Scholar: Author Only Title Only Author and Title

Schwab R, Ossowski S, Riester M, Warthmann N, Weigel D (2006) Highly specific gene silencing by artificial microRNAs in Arabidopsis. Plant Cell 18: 1121–1133
  Google Scholar: Author Only Title Only Author and Title

Shah L, Yahya M, Shah SMA, Nadeem M, Ali A, Ali A, Wang J, Riaz MW, Rehman S, Wu W, et al (2019) Improving Lodging Resistance: Using Wheat and Rice as Classical Examples. Int J Mol Sci. doi: 10.3390/ijms20174211
  Google Scholar: Author Only Title Only Author and Title

Sibout R, Proost S, Hansen BO, Vaid N, Giorgi FM, Ho-Yue-Kuang S, Legée F, Cézart L, Bouchabké-Coussa O, Soulhat C, et al (2017) Expression atlas and comparative coexpression network analyses reveal important genes involved in the formation of lignified cell wall in Brachypodium distachyon. New Phytol 215: 1009–1025
  Google Scholar: Author Only Title Only Author and Title

Szczegielniak J, Borkiewicz L, Szurmak B, Lewandowska-Gnatowska E, Statkiewicz M, Klimecka M, Cieśla J, Muszyńska G (2012) Maize calcium-dependent protein kinase (ZmCPK11): local and systemic response to wounding, regulation by touch and components of jasmonate signaling. Physiol Plant 146: 1–14
  Google Scholar: Author Only Title Only Author and Title

Trabucco GM, Matos DA, Lee SJ, Saathoff A, Priest H, Mockler TC, Sarath G, Hazen SP (2013) Functional characterization of cinnamyl alcohol dehydrogenase and caffeic acid O-methyltransferase in Brachypodium distachyon. BMC Biotechnol 13: 61
  Google Scholar: Author Only Title Only Author and Title

Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ (2016) W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids Res 44: W232–W235
  Google Scholar: Author Only Title Only Author and Title

Tsugama D, Liu S, Fujino K, Takano T (2018) Calcium signalling regulates the functions of the bZIP protein VIP1 in touch responses in Arabidopsis thaliana. Ann Bot 122: 1219–1229
  Google Scholar: Author Only Title Only Author and Title

Tsugama D, Liu S, Takano T (2012) A bZIP protein, VIP1, is a regulator of osmosensory signaling in Arabidopsis. Plant Physiol 159: 144–155
  Google Scholar: Author Only Title Only Author and Title

Tsugama D, Liu S, Takano T (2016) The bZIP protein VIP1 Is involved in touch responses in Arabidopsis roots. Plant Physiol 171: 1355–1365
  Google Scholar: Author Only Title Only Author and Title

Tsugama D, Liu S, Takano T (2014) Analysis of functions of VIP1 and its close homologs in osmosensory responses of Arabidopsis thaliana. PLoS One 9: e103930
  Google Scholar: Author Only Title Only Author and Title

Tyler L, Bragg JN, Wu J, Yang X, Tuskan GA, Vogel JP (2010) Annotation and comparative analysis of the glycoside hydrolase genes in Brachypodium distachyon. BMC Genomics 11: 600
  Google Scholar: Author Only Title Only Author and Title

Uyttewaal M, Burian A, Alim K, Landrein B, Borowska-Wykręt D, Dedieu A, Peaucelle A, Ludynia M, Traas J, Boudaoud A, et al (2012) Mechanical stress acts via katanin to amplify differences in growth rate between adjacent cells in Arabidopsis. Cell 149: 439–451
  Google Scholar: Author Only Title Only Author and Title

Valdivia ER, Herrera MT, Gianzo C, Fidalgo J, Revilla G, Zarra I, Sampedro J (2013) Regulation of secondary wall synthesis and cell death by NAC transcription factors in the monocot Brachypodium distachyon. J Exp Bot 64: 1333–1343
  Google Scholar: Author Only Title Only Author and Title

Van Leene J, Blomme J, Kulkarni SR, Cannoot B, De Winne N, Eeckhout D, Persiau G, Van De Slijke E, Vercruysse L, Vanden Bossche R, et al (2016) Functional characterization of the Arabidopsis transcription factor bZIP29 reveals its role in leaf and root development. J Exp Bot 67: 5825–5840
  Google Scholar: Author Only Title Only Author and Title

Van Moerkercke A, Duncan O, Zander M, Šimura J, Broda M, Vanden Bossche R, Lewsey MG, Lama S, Singh KB, Ljung K, et al (2019) A MYC2/MYC3/MYC4-dependent transcription factor network regulates water spray-responsive gene expression and jasmonate levels.

Proc Natl Acad Sci U S A 116: 23345–23356
  Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Warthmann N, Chen H, Ossowski S, Weigel D, Hervre P (2008) Highly specific gene silencing by artificial miRNAs in rice. PLoS One 3: e1829
  Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Weltmeier F, Ehlert A, Mayer CS, Dietrich K, Wang X, Schütze K, Alonso R, Harter K, Vicente-Carbajosa J, Dröge-Laser W (2006) Combinatorial control of Arabidopsis proline dehydrogenase transcription by specific heterodimerisation of bZIP transcription factors. EMBO J 25: 3133–3143
  Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Yin Y, Zhu Q, Dai S, Lamb C, Beachy RN (1997) RF2a, a bZIP transcriptional activator of the phloem-specific rice tungro bacilliform virus promoter, functions in vascular development. EMBO J 16: 5247–5259
  Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>