Mechanically induced localization of SECONDARY WALL INTERACTING bZIP is associated with thigmomorphogenic and secondary cell wall gene expression

Joshua H. Coomey[1,2], Kirk J.-M. MacKinnon[1,3], Ian W. McCahill[1,2], Bahman Khahani[1,2], Pubudu P. Handakumbura[1,2], Gina M. Trabucco[1,3], Jessica Mazzola[1], Nicole A. Leblanc[1], Rithany Kheam[1], Miriam Hernandez-Romero[1,2], Kerrie Barry[4], Lifeng Liu[4], Ji E. Lee[4], John P. Vogel[4], Ronan C. O'Malley[4], James J. Chambers[5], Samuel P. Hazen[1,2,3]

[1]Biology Department, University of Massachusetts, Amherst, MA 01003, USA,

[2]Plant Biology Graduate Program, University of Massachusetts, Amherst, MA 01003, USA,

[3]Molecular and Cellular Biology Graduate Program, University of Massachusetts, Amherst, MA 01003, USA,

[4]DOE Joint Genome Institute, Berkeley, CA 94720, USA,

[5]Institute for Applied Life Science, University of Massachusetts, Amherst, MA 01003, USA,

Author for correspondence:
Samuel P. Hazen, hazensam@umass.edu

ORCID
Joshua H. Coomey, https://orcid.org/0000-0001-5866-6671
Kirk J.-M. MacKinnon, https://orcid.org/0000-0001-9698-3985
Bahman Khahani, https://orcid.org/0000-0003-4499-2825
Pubudu P. Handakumbura, https://orcid.org/0000-0002-1708-8315
Kerrie Barry, https://orcid.org/0000-0002-8999-6785
Ji E. Lee, https://orcid.org/0000-0001-8325-1689
John P. Vogel, https://orcid.org/0000-0003-1786-2689
James J. Chambers, https://orcid.org/0000-0003-3883-8215
Samuel P. Hazen, https://orcid.org/0000-0001-9131-8994

AUTHOR CONTRIBUTIONS
JC, IM, and SH conceived and designed the study. JC, IM, PH, GT, MH-R, JV, RO developed techniques and reagents. JC, IM, JM, NL, RK, KB, LL, JL, JV. and RO acquired the data. JC, KM, IM, BK, and SH analyzed and interpreted the data. JC, KM, IM, and SH drafted the manuscript. SH agrees to serve as the author responsible for contact and ensures communication.

Abstract

Plant growth requires the integration of internal and external cues, perceived and transduced into a developmental program of cell division, elongation, and wall thickening. Mechanical forces contribute to this regulation, and thigmomorphogenesis typically includes reducing stem height, increasing stem diameter, and a canonical transcriptomic response. We present data on a bZIP transcription factor involved in this process in grasses. *Brachypodium distachyon* SECONDARY WALL INTERACTING bZIP (SWIZ) protein translocated into the nucleus following mechanostimulation. Classical touch responsive genes were upregulated in *B. distachyon* roots following touch, including significant induction of the glycoside hydrolase 17 family, which may be unique to grass thigmomorphogenesis. SWIZ protein binding to an E-box variant in exons and introns was associated with immediate activation followed by repression of gene expression. *SWIZ* overexpression resulted in plants with reduced stem and root elongation. These data further define plant touch-responsive transcriptomics and physiology, offering insights to grass mechanotranduction dynamics.

INTRODUCTION

Forces both internal and external to a cell influence growth. Turgor pressure in conjunction with anisotropic cell wall dynamics direct plant cell shape and expansion. Force perception between neighboring cells is critical in the development and maintenance of tissue form and function, such as the interlocking pavement cells on the leaf epidermis, or the developmental hotspots in the apical meristem (Hamant et al., 2008; Uyttewaal et al., 2012; Bidhendi et al., 2019). Specific inter-cell forces result in dynamic remodeling of the cortical cytoskeleton, with subsequent changes in cellulose microfibril alignment and alterations to other cell wall components such as pectin methyl esterification (Hamant et al., 2008; Uyttewaal et al., 2012; Bidhendi and Geitmann, 2018; Altartouri et al., 2019; Bidhendi et al., 2019). The classic hallmarks of touch responsive growth, or thigmomorphogenesis, include reduced plant height, increased radial growth in plants with a cambial meristem, increased branching, and delayed flowering time (Jaffe, 1973; Biro et al., 1980; Jaffe et al., 1980; Braam, 2004). These attributes have been leveraged by farmers for hundreds of years. As early as 1680, records show Japanese farmers tread on young wheat and barley seedlings to elicit increased branching, spikes per plant, and grain weight per plant, along with stronger roots (Iida, 2014). This practice, known as mugifumi, continues today with mechanized rollers. Thigmomorphogenesis in belowground tissues have also been studied to some extent, with the impact of stiffer substrates eliciting changes in root length and straightness, with the implication of hormonal signaling pathways in mediating this response (Lourenço et al., 2015; Lee et al., 2019; Nam et al., 2020).

Mechanical stimulus can significantly remodel gene expression (Braam and Davis, 1990; Braam, 2004; Lee et al., 2005). The so-called *TOUCH* (*TCH*) genes in *Arabidopsis thaliana*, encode calmodulin (*AtTCH1/AtCaM2*), calmodulin-like proteins (*AtTCH2/AtCML24*, *AtTCH3/CML12*), and a xyloglucan endotransglucosylase/hydrolase (*AtTCH4/AtXTH22*) (Braam and Davis, 1990). Touch responsive gene expression patterns often overlap with other stimuli such as dark, cold, and hormone treatment (Polisensky and Braam, 1996; Lee et al., 2005). In addition to calcium binding and signaling, genes related to cell wall modification and a variety of transcription

factors and kinases are regulated by mechanical stimulus, as well as genes involved in hormone
homeostasis and signaling.

Group I bZIPs are also implicated in mechanosensing. VIRE2-INTERACTING PROTEIN 1
(AtVIP1) and related Group I bZIP proteins translocate from the cytoplasm to the nucleus in
response to a variety of biotic and abiotic stimuli, including hypo-osmotic conditions (Tsugama
et al., 2012; Tsugama et al., 2014; Tsugama et al., 2016). The Group I bZIP *Nt REPRESSOR OF*
SHOOT GROWTH (*NtRSG*) in tobacco plays a role in maintaining GA homeostasis, wherein it
translocates to the nucleus in response to cellular bioactive GA levels (Igarashi et al., 2001;
Ishida et al., 2008; Fukazawa et al., 2010; Ito et al., 2017). Translocation appears to be dependent
on protein phosphorylation, either from MITOGEN ACTIVATED PROTEIN KINASE 3 during
pathogen invasion, or via calcium dependent protein kinases. When phosphorylated, Group I
bZIPs associate with 14-3-3 proteins in the cytoplasm until phosphatase activity releases them
for nuclear translocation (Ishida et al., 2004; Ito et al., 2014; Ito et al., 2017; Tsugama et al.,
2018; Van Leene et al., 2016).

Secondary cell walls deposited between the plasma membrane and primary cell wall provide
mechanical strength in vascular and structural tissues. Secondary walls are made of crystalline
cellulose, hemicelluloses, and phenolic lignin polymers. Although functionally similar,
secondary walls in monocotyledonous plants have key differences from eudicots, including
distinct hemicellulose chemistry and differences in lignin biosynthesis. Grasses also produce
mixed-linkage glucans (MLGs), a wall polysaccharide that is rarely found outside the
commelinid monocots (Coomey et al., 2020). A tightly controlled network of feed-forward loops
regulates the transcription of wall synthesizing enzymes, with NAC family transcription factors
activating wall synthesis genes as well as MYB family and other transcription factors that further
promote secondary wall synthesis (McCahill and Hazen, 2019). These networks are similar
between grasses and eudicots, with some components in each that have yet to be described in the
other (Coomey et al., 2020). There are currently no bZIP family members in any secondary wall
regulatory model.

Grasses employ a fundamentally different growth mechanism than eudicots, as they lack a
cambium layer, and thus no lateral meristem. Stem elongation comes from division and
elongation in a discrete series of intercalary meristems, called nodes, with one internode region
elongating and pushing up subsequent nodes. Detailed studies of thigmomorphogenesis have
been conducted almost exclusively in eudicots. However, recent work in the model cereal grass
Brachypodium distachyon shows general overlap with conventional eudicot
thigmomorphogenesis but with no change in stem diameter and increased time to flower
(Gladala-Kostarz et al., 2020).

Thigmomorphogenesis is a widely observed phenomenon that results in reduced height,
increased radial growth, and increased branching. The mechanisms behind this form of growth
are not yet fully understood, but involve aspects of hormone regulation, Ca^{2+} signaling, Group I
bZIP intracellular translocation, and changes in gene expression. Here we describe the
transcriptional response to mechanical stimulation and the function of a *B. distachyon* bZIP
transcription factor, SECONDARY WALL ASSOCIATED bZIP (Bradi1g17700) and its role in
touch response and cell wall biosynthesis.

RESULTS

SWIZ is a Group I bZIP transcription factor and candidate cell wall regulator

To identify genes involved in the regulation of secondary cell wall thickening, Trabucco et al., (2013) measured transcript abundance in *B. distachyon* leaf, root, and stem tissue. A gene annotated as a bZIP transcription factor, Bradi1g17700, was highly expressed in root and stem relative to leaf (**Supplemental Figure S1A**). Bradi1g17700 is also a member of a 112 gene coexpression network (**Supplemental Table S1**) that includes genes highly expressed in the peduncle (Sibout et al., 2017). Phylogenetic analysis of Bradi1g17700, hereinafter referred to as SECONDARY WALL INTERACTING bZIP (SWIZ), amino acid sequence shows it to be an ortholog of the *A. thaliana* Group I bZIPs (Jakoby et al., 2002; Dröge-Laser et al., 2018), and closely related to *AtbZIP18* and *AtbZIP52* (**Supplemental Figure S1B, Supplemental File S1**).

SWIZ translocates into the nucleus in response to mechanical stimulus

Group I bZIPs in *A. thaliana* translocate between the cytosol and nucleus in response to external cellular force. As an ortholog of these proteins, we hypothesized that SWIZ protein may similarly translocate within the cell in response to mechanical force. To test this, the roots of plants overexpressing either *SWIZ* fused to *GFP* (*SWIZ:GFP-OE*) or *GFP* alone (*GFP-OE*) were observed following a mechanical stimulus (**Fig 1A, Supplemental Files 2 and 3**). In control plants, GFP signal was both cytosolic and nuclear, which remained constant over the imaging period (**Fig 1B**). GFP signal was mostly observed in the cytosol in *SWIZ:GFP-OE* plants, but following mechanical stimulus, nuclear GFP signal increased substantially, peaking around 30 min post-stimulus and returned to near basal levels by ~60 min, while untouched plants showed no change in signal localization (**Fig 1B**).

Touch response to stimulus can saturate at a certain number of treatments (Martin et al., 2010; Leblanc-Fournier et al., 2014; Moulia et al., 2015). To test if SWIZ translocation dynamics varied after repeated treatments, we applied mechanical force to *SWIZ:GFP-OE* roots at regular intervals. A second stimulus was given 90 min after the first, and a third at 180 min. Following each mechanical stimulation, SWIZ consistently translocated from cytoplasm to nucleus (**Fig 1D**). This suggests that SWIZ translocation dynamics are not impacted by repeated stimulus events 90 min apart.

To determine if the signal triggering SWIZ translocation is spread beyond the specifically stimulated region, two regions of the same *SWIZ:GFP-OE* root separated by 3 cm were simultaneously observed. The stimulated region showed typical SWIZ:GFP nuclear signal accumulation and in the region below no translocation was observed (**Fig 1C**). At 120 min, the treatments were reversed, with the lower root region receiving a stimulus while the upper region was unperturbed. The lower region showed SWIZ:GFP nuclear translocation while the upper region did not. Thus, the touch stimulus is localized directly to the perturbed region.

Figure 1. SWIZ translocates to the nucleus in response to mechanical stimulus, specifically in regions directly stimulated. (A) Image of *SWIZ:GFP-OE* and *GFP-OE* roots prior to stimulus and 30 min post stimulus. Roots were observed immediately following mechanical perturbation. (B) Quantification of nuclear signal in control (purple) and touched (teal) conditions for *GFP-OE* (left) and *SWIZ:GFP-OE* (right). n = 14-20 nuclei. (C) SWIZ translocation occurred in the local area of the stimulus. At 30 min, stimulus was applied to an upper region of the root, while at 120 min it was applied to a lower region approximately 3 cm below. n = 109, 184 nuclei respectively for upper and lower regions. Scale bar = 100 μm, (D) *SWIZ:GFP-OE* roots were imaged by confocal microscopy with stimulus applied in the field of view at 0, 90, and 180 min. n = 126 nuclei. (B-D) Images were taken every 2 min. Nuclear GFP signal was quantified in selected nuclei at each time point. The average nuclear GFP signal is represented by the line with error bars indicating standard error of the mean. Scale bar = 100 μm. n = 4-6 plants per treatment.

126
Transcriptional response to touch

Having established the nuclear translocation of SWIZ in response to touch, we then investigated
what effect touch and *SWIZ* overexpression during touch may have on gene expression. We
measured transcript abundance by sequencing mRNA from wildtype and *SWIZ-OE* root tissue
just prior to touch (0 min) and at 10, 30, and 60 min following touch treatment along the entire
length of the root (**Fig 2A, Supplemental Fig S2**). Principal component analysis of transcript
abundance shows the greatest amount of variance is attributed to the first principal component,
60%, where samples cluster based on genotype (**Fig 2B**). The second principal component,
which accounts for 17% of the variance, distinguished between pre-touch and 10 min following
touch, with the last two time points clustering similarly.
137
The wildtype transcriptome was massively remodeled in response to touch, with 8,902
transcripts differentially expressed ($q < 0.1$) at 10 min post touch, 5,682 transcripts at 30 min,
and 7,672 transcripts at 60 min (**Supplemental Tables S2-4**). Canonical touch-responsive genes
(TCH) include calmodulins (CAM), calmodulin-like (CmL), and xyloglucan endo-
transglycosylase/hydrolases (XTH) (Braam and Davis, 1990; Lee et al., 2005). Based on
homology with *TCH1*, *TCH2*, and *TCH3*, 79 CAM and CmL genes were identified, the majority
of which showed upregulation in wildtype plants following touch treatment (**Fig 2C**,
Supplemental Table S11). Similarly, homology with *TCH4* identified 37 *B. distachyon XTH*
and *XTH-like* genes. Of these, 8 genes were upregulated by touch in wildtype plants, including
the two with the highest similarity to *TCH4* (Bradi1g33810 and Bradi33840) (**Fig 2C**,
Supplemental Table S12). We also investigated the effect of touch on other glycosyl hydrolase
families (Tyler et al., 2010). GH17 genes were differentially regulated following touch. The
GH17 family is broadly described as modifying β-1,3-glucans. Thirty-five of the 53 annotated
GH17 members were measured in our data, with 8, 11, and 16 members differentially expressed
at 10, 30 and 60 min respectively. Fisher's exact test determined significant enrichment of GH17
expression at time 60 ($p = 9.307e-3$) (**Fig 2C, Supplemental Table S12**). An interactive
platform for exploring this *B. distachyon* touched-root transcriptome is available online at
(https://hazenlab.shinyapps.io/swiztc/).
156
Cell wall genes are downregulated immediately following touch, then induced concurrent
with SWIZ nuclear translocation and more strongly in *SWIZ-OE*
Genes related to secondary cell wall synthesis were immediately downregulated in wildtype
plants following touch, but were subsequently upregulated (**Fig 3, Supplemental Table S5-10**).
Xylan, cellulose, MLG, and callose synthesis associated genes were upregulated 30 and 60 min
following touch, while lignin biosynthesis and BAHD acyltransferase genes were upregulated at
60 min. Prior to touch, nearly all cell wall associated genes investigated were significantly
downregulated relative to wildtype, then significantly upregulated following touch (**Fig 3**). In
SWIZ-OE, cell wall-related terms were enriched in paths that had stronger upregulation earlier
than the same terms in wildtype (**Supplemental Fig S3A-B**). Network analysis of gene
expression patterns over time using the iDREM software showed that in *SWIZ-OE* plants, cell
wall genes were more rapidly induced. Notably, cell wall polysaccharide biosynthesis,
particularly glucan biosynthesis, were significantly enriched in *SWIZ-OE* path D, while not
significantly enriched in wildtype (**Supplemental Table S14**). This is mirrored when looking at
specific gene expression, with primary wall CESAs, MLG, and callose synthesis genes all
significantly upregulated following touch in *SWIZ-OE*. Thus, cell wall genes were immediately

repressed by touch, followed by significant activation, which occurs more rapidly in *SWIZ-OE*
plants and is associated with the timing of SWIZ nuclear translocation.

Figure 2. Transcriptome analysis of touch response in *Brachypodium distachyon* roots.
(A) Root tissue was sampled just prior to touch (t = 0), and at 10, 30, and 60 min following
touch treatment in wildtype and *SWIZ-OE*. (B) Principal component analysis of gene
expression across samples shows the greatest difference corresponding to genotype and the
second greatest corresponding to time after touch. (C) Canonical, as well as novel touch
responsive genes are upregulated in *B. distachyon* following touch. Closest orthologs of the
Arabidopsis thaliana TOUCH genes encoding calmodulin-like (CML) and xyloglucan endo-
transglycosylase/hydrolases (XTH) are upregulated following touch, as are previously
unreported members of the glycosyl hydrolase 17 (GH17) family. Full list of gene expression
for *TOUCH* and glycoside hydrolases defined by Tyler et al. (2010) in Supplemental Tables
S10-12. Significance denoted by * reflecting $q < 0.1$ compared to expression at t = 0, with q-
values representing Wald test p-values adjusted for false discovery rate.

SWIZ protein binding in the gene body is associated with dynamic changes in gene expression

Next, we investigated the direct binding targets of SWIZ protein by performing DNA affinity purification sequencing (DAP-seq) with whole genomic DNA. SWIZ interacted with 2,455 distinct positions in the genome (**Supplemental Table S15**). Those regions were significantly enriched for two E-box motif variants, (A/C)CAGNCTG(T/G) and (A/C)CAGCTG(T/G) (**Fig 4A**). Numerous binding sites were between the translational start and stop sites, with 21% of peaks in introns and 21% in exons (**Fig 4B**). Fewer than 10% of the peaks occurred in UTRs and the promoter (5 kb upstream of the 5 'UTR) and intergenic regions each accounted for 25%. Thus, SWIZ protein is preferentially bound to an E-box like motif and often in the gene body (**Fig 4C**).

Figure 3. Gene expression analysis of cell wall related genes. Log fold-change of gene expression measured by RNA-seq in wildtype and *SWIZ-OE*, presented as relative to wildtype expression at time 0, pre-touch. Bar color indicates class of cell wall gene. Error bars indicate standard deviation of three biological replicates. Significance denoted by * reflecting $q < 0.1$ compared to wildtype expression at t = 0, with q-values representing Wald test p-values adjusted for false discovery rate. Legend abbreviations: BAHD, BAHD (BEAT, AHCT, HCBT, and DAT) acyltransferases; CESA, cellulose synthase; MLG, mixed-linkage glucans.

We next compared genes differentially expressed in the *SWIZ-OE* touch response RNA-seq time courses with the binding targets identified by DAP-seq (**Supplemental Table S5-9,15**). Prior to touch, genes with promoter-SWIZ interactions were most often downregulated in *SWIZ-OE* relative to wildtype (**Table 1**). Following touch, SWIZ promoter binding targets were most often upregulated in *SWIZ-OE* in all three post-touch time points. Thus, SWIZ promoter binding was coincident with repression, but following a touch stimulus, activation was more prominent. This difference between untreated and touched was not as pronounced in gene body targets, with more upregulated genes 10 min following touch and more downregulated 30 and 60 min

following touch. To further explore these trends, we conducted iDREM cluster analysis of the differentially expressed genes that were also SWIZ protein targets (**Fig 4D**). For both wildtype and *SWIZ-OE*, the network analysis revealed a pathway to increased gene expression by 60 min following touch. The network pathway trend for both promoter and gene binding targets in wildtype was immediate repression at 10 min followed by increased expression at 30 and 60 min post-stimulus. A unique pathway was observed among gene body binding targets in *SWIZ-OE*; transcript abundance was reduced following touch. Thus, SWIZ binding to a promoter or gene body was more strongly associated with increased expression except for gene body targets in touched *SWIZ-OE* plants, which were more strongly repressed.

Several vignettes stood out as examples of genes differentially expressed by touch, *SWIZ-OE* transgene, and being bound by SWIZ at various locations. The mixed glucan synthase *CSLF6* was bound in the first intron and upregulated by touch and *SWIZ-OE* (**Fig 4E**). *GA2ox3*, which inactivates bioactive GA, was bound by SWIZ protein in both the gene body and 3'UTR and was upregulated 30 min after touch in *SWIZ-OE* (**Fig 4F**). An example of gene body binding repression is *SWAM3* (Bradi1g30252), the closest ortholog to a wheat transcription factor induced by hypoxia, *TaMYB1* (Lee et al., 2006; Handakumbura et al., 2018) (**Fig 4G**). A membrane-associated transcription factor, NAC35, was bound by SWIZ in the promoter region and downregulated in SWIZ-OE (**Fig 4H**).

Cis-regulatory sequences associated with mechanical stress, wounding, and cell wall synthesis are enriched among touch responsive genes

Touch-responsive genes in wildtype were analyzed for enrichment of putative *cis*-regulatory elements (CREs). We identified several sequences significantly enriched among touch-responsive transcripts (**Fig 5, Supplemental Fig S4, Supplemental Table S16-17**). Several of those have been previously described as touch responsive, including the Rapid Stress Response Element, the GCC boxes AGCCGCC and GCCGCC, a sequence referred to as both the E-box and G-box (CACGTG), CM2, AP2-like, GGNCCCAC site II element, P-box, and the GRF and FAR1 binding sites (Rushton et al., 2002; Walley et al., 2007; Doherty et al., 2009; Fernández-Calvo et al., 2011; Moore et al., 2022). Other putative CREs have not previously been identified as touch responsive. One CRE was exclusively enriched in touch-repressed genes at all time points, the TCP site II element TGGGC. A GATA-like binding site was enriched among 10 min repressed transcripts. The homeobox binding site motif was enriched among both induced and repressed genes. The CGCG-box was enriched among induced genes at all timepoints like RSRE, CM2, and FAR1. The two CREs associated with secondary cell wall thickening were also significantly enriched (Coomey et al., 2020); the VNS element among induced genes, and the AC element ACC(A/T)ACC with a unique profile. AC elements were enriched among repressed genes at 10 min, in both induced and repressed genes at 30 min, and among induced genes only at 60 min.

Figure 4. DNA affinity purification sequencing to determine SWIZ binding sites and differential expression in response to touch and *SWIZ-OE*. (A) Top two most statistically enriched sequence motifs in SWIZ binding sites. (B) Distribution of binding sites across genomic features, relative to primary transcripts of the *Brachypodium distachyon* annotation v 3.1. (C) Relative distribution of binding sites centered on the transcriptional start site (TSS, blue dashed line), transcriptional termination site (TTS, red dashed line) represents the average length of all annotated transcripts, approximately 4.5 kb away from the TSS. (D) Path determinations from iDREM time course analysis of differentially expressed genes that also have DAP-seq binding sites. Each line represents a set of genes with similar expression level patterns over the time course relative to time 0, pre-touch. Wildtype: promoter and gene body = 170 and 146 genes, respectively. *SWIZ-OE*: promoter = 163 genes, gene body A and B = 141 and 60, respectively. (E-H) Gene expression over time of selected genes with SWIZ binding sites. Line graphs are the average transcript abundance of three biological replicates for each time point. Binding site determined as peaks of sequence alignment. Scale bar unit is bases. Direction of transcription is shown with arrows on the gene model, 5' and 3' UTRs are depicted by narrowed rectangles on the gene model.

240
Root morphology is altered by *SWIZ-OE* and mechanical treatment
Given the SWIZ protein mechanical response and regulatory influence, we then investigated how
SWIZ might impact plant growth in response to touch. We challenged wildtype and *SWIZ-OE*
roots with the same style of touch treatment used in the translocation and gene expression
experiments, but repeated twice daily for a period of five days. In both touch and control
conditions, *SWIZ-OE* roots were significantly shorter than wildtype, suggesting a dwarfing effect
from *SWIZ* overabundance (**Supplemental Fig S5A-C**). Mechanical challenges to roots have
been reported to impact root straightness, a trait that has also been described as being impacted
in other bZIP studies (Van Leene et al., 2016; Zha et al., 2016). In control conditions, we
observed that *SWIZ-OE* roots were significantly less straight than wildtype, while this was not
observed in in response to touch (**Supplemental Fig S5D**). We further tested the
mechanoresponse of *SWIZ-OE* roots by growing seedlings on plates with increasing degrees of
plate angle at 10°, 20°, 30°, and 40° from vertical as a form of mechanostimulation (Oliva and
Dunand, 2007; Zha et al., 2016) (**Supplemental Fig S6**). Wildtype *B. distachyon* roots displayed
decreasing root length with increasing plate angle. *SWIZ-OE* roots were shorter than wildtype at
all plate angles tested (**Supplemental Fig S6**). Root straightness did not show any significant
differences.
258
B. distachyon **displays classic thigmomorphogenic phenotypes**
We next investigated the effect of touch treatment on aboveground tissue. Wildtype plants were
perturbed with a metal bar once every 90 min for either two or three weeks (**Supplemental Fig
S7 and S8**). After the treatment period, all plants were allowed to recover and grow to
senescence (**Supplemental Fig S8**). Two-week stressed plants were significantly shorter than
control plants, and three-week stressed plants were shorter still (**Supplemental Fig S8A-B**).
Despite this difference in height, there was no significant difference in aboveground biomass
(**Supplemental Fig S8C**). Three-week stressed plants had significantly more branches,
(**Supplemental Fig S8D**). Transverse stem cross sections in the third elongated internode and
peduncle did not show a significant difference in cell wall thickness or phloroglucinol staining in
response to touch (**Supplemental Fig S8E-G**).

Figure 5. Sequence motifs enriched in the *cis*-regulatory regions of touch responsive *Brachypodium distachyon* genes. Negative log *p*-values for *cis*-elements, known and not known to be touch responsive. RSRE - Rapid Stress Response Element, FAR1 - FAR-RED impaired response1, GRF - Growth Regulating Factor, VNS - VND, NST/SND, SMB.

272
Stem height and cell wall thickening are affected by *SWIZ-OE* and touch treatment
To test the role of *SWIZ* in regulating thigmomorphogenesis and secondary cell wall
development, we tested wildtype and *SWIZ-OE* under two weeks of mechanical perturbation as
described above. In control conditions, there were no differences among genotypes in height,
weight, or branching. Touch significantly shortened both wildtype and *SWIZ-OE* stems relative
to control conditions, but not relative to each other (**Supplemental Fig S9-10**). Transverse
sections of the stem were made in the peduncle, the last elongated stem internode where the
touch treatment occurred during stem elongation, and stained with phloroglucinol-HCl. *SWIZ-*
OE touched peduncles showed significantly thicker interfascicular fiber cell walls compared to
untouched, but not significantly different from touched wildtype (**Supplemental Fig S10B-C**).
283
284 **DISCUSSION**
285
Touch stimulus is generally an inhibitor of plant elongation growth, but promotes branching and
radial expansion (Jaffe, 1973; Braam, 2004; Chehab et al., 2009). The majority of this work has
been done in dicots, where increased radial growth has been associated with greater deposition of
secondary wall forming cells, particularly in woody species such as poplar (Biro et al., 1980;
Coutand et al., 2009; Börnke and Rocksch, 2018; Roignant et al., 2018; Niez et al., 2019). Our

understanding of thigmomorphogenesis in grasses is limited, and mostly in the agricultural
context of lodging (Shah et al., 2019). While these studies highlight the importance of stem
strength and associated cell wall defects, they do relatively little to elucidate the
mechanosensitive response. Recent work by Gladala-Kostarz et al (2020) describes grass touch
response to both wind and direct mechanical treatment, with an emphasis on cell walls and stem
anatomy. Touch treatment significantly increased lignin content and wall stiffness.
297
Specific gene expression patterns have become molecular hallmarks of plant touch response,
most notably induction of the *TCH* and similar genes. Orthologs of wall modifying XTHs and
CAM/CML signaling genes are upregulated in response to mechanical stimulation across
species, and as we present here, in *B. distachyon*. Touch elicits major global changes in gene
expression, with 2.5% to 10% of the genes assayed in *A. thaliana* and poplar differentially
expressed (Lee et al., 2005; Pomiès et al., 2017; Van Moerkercke et al., 2019). In sorghum
(*Sorghum bicolor*), leaf sheath imposes mechanical constraints on emerging buds, and removing
this dramatically altered gene expression, with 42% of genes differentially expressed over a 9 h
period (Liu and Finlayson, 2019). Recent work in cereal species, including oat, wheat, and
barley, showed 2-5% differential gene expression within 2h of mechanostimulation of young
leaves with a soft brush (Darwish et al., 2023). In our analysis we applied a *q*-value significance
cutoff of 0.1, which accounts to some extent for the larger number of genes we find differentially
expressed compared to other reports. An interactive application
(https://hazenlab.shinyapps.io/swiztc) allows users to sort gene expression results by *q*-value, *p*-
value, or log-2 fold change, and may be utilized by readers to make their own comparisons. Our
results are similar to studies in other species in the strong induction of canonical TCH genes,
along with biotic and abiotic stress related genes (Pomiès et al., 2017). Canonical touch
responsive genes such as the orthologs of *TCH1-4* and related *XTH* and *CML* genes were
upregulated immediately following stimulus. We also note the previously unreported touch-
induction of GH17 family genes, as well as *CSLF6*, suggesting a role for MLG and β-1,3 glucan
modification in grass touch response.
319
Touch-regulated expression of secondary cell wall related transcripts appears to differ between
early and late timepoints. Upon touch, we observed immediate repression of many key cell wall
biosynthetic enzymes, followed by significant upregulation one hour after stimulus. In poplar, no
significant repression of these transcripts was reported, but terms related to wood or cell wall
synthesis were also not enriched among touch-regulated transcripts until 24 or 72 hours after
touch (Pomiès et al., 2017). In other cereals, genes related to cell wall polysaccharides were
upregulated 25-60 min following touch while lignin biosynthetic genes down regulated (Darwish
et al., 2023). While differences in mechanostimulation, tissue type, gene expression
quantification, and temporal sampling schemes complicate direct comparisons of plant touch
response experiments, together these data suggest that delayed induction of secondary cell wall
transcripts is a common feature of plant touch responses.
331
Thigmomorphogenesis and secondary cell walls
The timing of SWIZ nuclear accumulation following mechanostimulation was consistent with *A.*
thaliana Group I bZIPs (Tsugama et al., 2014; Tsugama et al., 2016). However, by adopting a
finer temporal sampling scheme, and quantifying nuclear signal sooner after touch (2 vs. 30
min), our results clarify the rapidity of this translocation. Furthermore, we show that the speed

and magnitude of SWIZ nuclear accumulation is not diminished over successive touch
treatments. Poplar transcriptional response to a second touch stimulus was significantly
attenuated relative to a single treatment and trees given daily touch treatment over four days no
longer showed touch-induced increases in their rates of radial growth (Martin et al., 2010;
Pomiès et al., 2017). If *B. distachyon* is likewise desensitized to repeated touch, the repeatability
of SWIZ translocation implies a mechanism downstream of touch perception and bZIP
translocation in mediating that shift.

Although we did not observe a robust secondary wall response in touched stems, touch-
responsive wall synthesis is still implicated by our transcriptomic data. Two CREs not previously
associated with touch response, the AC and VNS elements, were identified in our touch-
responsive and SWIZ-targeted datasets. These sites are bound by NAC and MYB proteins that
regulate the thickening of secondary cell walls (Ohtani et al., 2011; Zhong et al., 2011; Kim et
al., 2012; Zhong and Ye, 2012; Handakumbura et al., 2018; Olins et al., 2018; Tamura et al.,
2019).

Prior to touch, almost the entire monolignol biosynthetic pathway was significantly
downregulated in *SWIZ-OE* plants. Following touch, most of these genes were significantly
upregulated. This activity is consistent with the compositional data presented by Gladala-Kostarz
et al. (2020). This activation may be a result of direct binding of SWIZ, or an indirect effect from
other transcriptional regulators like the NAC transcription factor *SWN5* that is significantly
upregulated in *SWIZ-OE* following touch and capable of activating the full developmental
program for secondary cell wall synthesis (Valdivia et al., 2013). The E-box (CANNTG) was
initially described as a bHLH binding motif, many bZIP phylogenetic groups in *A. thaliana* have
been shown to bind similar sequences (O'Malley et al., 2016). Using the same technique, we
identified the SWIZ binding motifs as (A/C)CAGNCTG(T/G) and (A/C)CAGCTG(T/G), very
similar to those bound by *A. thaliana* orthologs, although an ambiguous nucleotide in the core of
one variant is not reported for other Group I bZIPs. Genes bound by SWIZ *in vitro* show both
activation and repression following touch, suggesting a complex regulatory function for SWIZ.
Furthermore, SWIZ direct binding sites were found in both promoter regions and gene bodies,
and most often in genes activated by touch. SWIZ gene body binding targets tended to be
repressed in *SWIZ-OE* plants following touch, without either being clearly associated with up or
down regulation. AtVIP1 and AtbZIP29 have both been described as activators (Yin et al., 1997;
Ringli and Keller, 1998; Pitzschke et al., 2009; Van Leene et al., 2016) while AtbZIP18 is
described as a repressor (Tsugama et al., 2012; Gibalová et al., 2017). The directionality of
SWIZ transcriptional control remains to be determined.

CONCLUSIONS

Touch significantly remodeled the *B. distachyon* transcriptome, with notable changes in wall
polysaccharide biosynthetic gene expression not previously reported and revealed an enrichment
of secondary cell wall associated CREs, the AC and VNS elements. SWIZ mechanotropism
shows similar dynamics to other bZIP proteins and the timing of this translocation is consistent
with the enhanced touch-responsive gene expression in *SWIZ-OE*. Some of the data presented
here relies on overexpression of *SWIZ*, which comes with certain limitations as to how directly
the phenotypic and molecular results relate to SWIZ-specific function in wildtype plants. Further

experimentation is needed to elucidate the bZIP regulatory network in response to touch and
other stimuli.

MATERIALS AND METHODS

Phylogenetic analysis
Protein sequences described for *A. thaliana*, *B. distachyon*, and *O. sativa* (Liu and Chu, 2015) as
Group II bZIPs were selected and searched against the most recent genome annotations in
Phytozome v12.1 (https://phytozome.jgi.doe.gov). The *Nicotiana tabacum* homologs NtRSGa
and NtRSGb were also added to the analysis. Protein sequences were aligned by MAFFT using
the L-INS-I model (Katoh et al., 2019). A maximum-likelihood phylogenetic tree bootstrap
resampling value of 1000 was generated in W-IQ-TREE (Trifinopoulos et al., 2016). All proteins
in the phylogenetic analysis are described in **Supplemental File S1**.
392
Plant transformation
Overexpression constructs and plant transformation carried out as previously described using
accession Bd21-3 (Handakumbura et al., 2013). Bradi1g17700 coding sequence was amplified
from cDNA and cloned into the pOL001 ubigate ori1 binary expression vector (Handakumbura
et al., 2018) to make the *SWIZ-OE* trangene. The coding sequence without stop codon was
amplified and cloned in frame with the enhanced GFP coding sequence in the pOL001 vector to
generate *SWIZ:GFP-OE*. For both constructs, three independent events were analyzed with no
phenotypic difference between them for height or translocation dynamics among the *SWIZ:GFP-*
OE lines.
402
Translocation assay
Bd21-3 seeds were surface sterilized and grown vertically on 1X MS media, pH 5.7, without
sucrose for 6d at 28°C in the dark. After 6 d, seedlings were moved to treatment plates
containing 1X MS media, pH 5.7.

All observations were made on a Nikon A1R scanning confocal microscope using a Plan Apo
10x 0.5NA objective and PMT detector. Root areas were located by eye using transmitted light
and then imaged with excitation at 488 nm and emission captured at 510-530 nm. Roots were
imaged for 30 min pre-treatment, with images captured every 2 min.

To elicit the touch response, the observed root region was gently probed 5 times in ~5 sec with a
blunt probe while observing through the eyepiece (**Supplemental Fig S7**). Images were captured
for 60-90 min post treatment. For experiments with multiple stimulus events, the timelapse
sequence was paused and roots were probed as described for the relevant stimulus events.

Analysis of GFP signal was done using the Nikon NIS Elements Advanced Research V5
software package. Nuclear regions were thresholded for intensity and particle size to identify
regions of interest. Fluorescence intensity was calculated for each ROI and averaged at each
timepoint.
419
Thigmomatic construction and operation
The Thigmomatic is a basic robotic device that sweeps plants with a metal bar at regular
intervals to elicit a touch response. The device was constructed from aluminum V-Slot linear rail
(Openbuilders Partstore, Monreoville, NJ) and bracket joints for the upright supports (20x20

mm), cross bars (20x20 mm), and tracks (20x40 mm). Two gantry carts ride along the 20x40 mm
V-Slot linear rails, connected by a 6.35 mm diameter metal rod bolted to the carts. Their
movement is powered by a belt driven linear actuator system using a NEMA 17 stepper motor
with a 12V 18W AC/DC power supply. Motor function is controlled by a Raspberry Pi 3B
microcomputer with stepper motor HAT (Adafruit Industries, New York). The Thigmomatic was
programmed to cover a specified distance in one direction once every 90 min.
430
Transverse stem sections and histology

Internode segments of the main stem were embedded in 8% agarose. A Leica VT1000
Vibratome was used to make 55 µm thick transverse sections. Histochemical staining with
phloroglucinol-HCl was done as previously described (Matos et al., 2013). Images were obtained
at 4, 10, and 20X using a Nikon Eclipse E200MV R light microscope and PixeLINK 3 MP
camera. Cell wall thickness was quantified for interfascicular fiber cells separated by one cell
layer from the mestome cells on the phloem side of major vascular bundles. Using ImageJ, lines
were drawn across two adjoining walls divided by two yielding one cell wall width.

RNA extraction and quantification and analysis

Seedlings were grown for 6 d on vertical 1X MS agar plates. Touch treatment was performed as
described above using a metal probe along the entire length of the root. Untouched samples were
collected immediately before touch treatment and touched samples collected 10, 30, and 60 min
post-treatment. Three roots were pooled per biological replicate, and RNA extracted by Qiagen
RNeasy Plant Mini Kit with on-column DNA digestion with RNase-free DNase I (Qiagen).
Strand-specific libraries were prepared using the Illumina TruSeq kit. Libraries were sequenced
using Illumina technology and processed as previously described (MacKinnon et al., 2020).
Briefly, quality was checked with FastQC (Andrews, 2010), aligned to the Bd21 reference
genome (v3.1) using HiSat2 (Kim et al., 2015), then assembled and quantified using StringTie
(Pertea et al., 2015). Transcripts were normalized and differential expression tested using the
likelihood ratio test from the R (v3.6.0) package DESeq2 (Love et al., 2014). Benjamini-
Hochberg p-value adjustments were applied to account for multiple testing with a significance
cutoff of 0.1 and 30,380 of 34,310 reference genes had non-zero read counts after normalization,
with an average mapping percentage of 97.8% for all libraries, as determined by SAMtools (Li et
al., 2009). Specific treatment contrasts (i.e., wildtype vs *SWIZ-OE*) were compared by Wald test
from DESeq2 (Love et al., 2014). Statistical enrichment of gene families was assessed using
Fisher's exact test. Raw read data was deposited in the European Nucleotide Archive for public
access (Accession no.: E-MTAB-10084).

DNA affinity purification sequencing

DNA affinity purification was carried out as previously described (Handakumbura et al., 2018).
In brief, transcription factor coding sequences were HALO tagged and mixed with Bd21
genomic DNA for *in vitro* binding. Protein-DNA was crosslinked, fragmented,
immunoprecipitated using the HALO antibody, barcoded, and sequenced. Reads were mapped to
the Bd21 genome using HiSat2 (Kim et al., 2015) to identify binding target loci. Peak calling and
motif analysis was done using HOMER v4.10 (Hypergeometric Optimization of Motif
EnRichment) suite (Heinz et al., 2010). Motif enrichment was calculated against the
hypergeometric distribution; the significance threshold was set to $p < 0.05$. The nearest annotated
gene to a bound peak was used for GO analysis. Raw read data were deposited in the European
Nucleotide Archive for public access (Accession no.: E-MTAB-10066).

469
Gene Ontology analysis
Phytozome was used to find orthologs for all *B. distachyon* v3.1 genes as the reciprocal best
match to *A. thaliana* TAIRv10 protein sequences. Arabidopsis gene identifiers were submitted to
g:Profiler (Raudvere et al., 2019) for KEGG and Wiki pathway enrichment analysis.

iDREM network analysis
Probabilistic graphical models that predict diverging gene expression paths were generated using
iDREM (Ding et al., 2018). Briefly, this software applies an input-output hidden Markov model
to time course gene expression data overlaid with static regulatory information, in this case
SWIZ DAP-seq protein-DNA interactions. GO analysis, described above, was performed for
each path identified by iDREM.

Cis-regulatory sequence analysis
Differentially expressed genes following touch were categorized as increasing or decreasing in
transcript abundance at each time point. Homer v.4.10 identified regulatory sequences in the
1000 bp upstream of the transcriptional start site of differentially expressed genes previously
identified in *A. thaliana* DAP-seq analysis (Heinz et al., 2010; O'Malley et al., 2016). We also
applied the growing k-mer approach to identify CREs (Moore et al., 2022). In brief, touch-
responsive genes were divided into six groups: up or down regulated each timepoint. The 1000
bp upstream of differential expressed genes were searched for all possible 6-mers, plus one
additional base (A,T,C,G), and then tested for *p*-value shift. If the *p*-value was lower, the 7-
mer(s) was kept and grown further up to 12-mer length. The Tomtom tool in MEME Suite 5.4.1
found similarities between significant CREs and the *A. thaliana* DAP-seq database, with a false
discovery rate cutoff of < 0.01.

Root touch experiment
Bd21-3 seeds were surface sterilized and plated on 1x MS, pH 5.7, containing 0.05 % MES as a
buffering agent and 1% plant agar (Gold Bio). Seeds were stratified on plates in the dark at 4 ℃
for 2 days and then transferred to a Percival PGC-15 growth chamber with day/night conditions
of 16h light at 24 ℃ and 8h dark at 18 ℃, respectively and grown at a ~10° angle from vertical.
After 2 days, a touch location was designated by selecting the lower of 1 cm up from the root tip
or 1 cm down from the seed and marked on the plate. For 5 days, this marked spot was treated
twice daily, two hours before and two hours after chamber midday (ZT6 and ZT10). Touch
treatment consisted of five firm presses with the side of a sterile pipette tip.

Root length and straightness measurement
Plates were photographed at the conclusion of the experiment. Semi-automated measurements of
root length were performed using the Smart Roots plugin for ImageJ (Lobet et al., 2011).
Straightness was quantified as described in (Swanson et al., 2015); the straight line distance
between root tip and the base of the seed was measured, and this value was divided by the traced
length of the root.

SUPPLEMENTAL MATERIALS

Supplemental Table S1. Cluster 56 from network analysis of gene expression atlas (Sibout et al. 2017) includes Bradi1g17700.

Supplemental Table S2. Differentially expressed genes in wildtype root tissue 10 min after touch treatment.

Supplemental Table S3. Differentially expressed genes in wildtype root tissue 30 min after touch treatment.

Supplemental Table S4. Differentially expressed genes in wildtype root tissue 60 min after touch treatment.

Supplemental Table S5. Differentially expressed genes in *SWIZ-OE* root tissue prior to touch treatment (0 min).

Supplemental Table S6. Differentially expressed genes in *SWIZ-OE* root tissue 10 min after touch treatment.

Supplemental Table S7. Differentially expressed genes in *SWIZ-OE* root tissue 30 min after touch treatment.

Supplemental Table S8. Differentially expressed genes in *SWIZ-OE* root tissue 60 min after touch treatment.

Supplemental Table S9. Overlap of differentially expressed genes in wildtype and *SWIZ-OE* root tissue before touch (0 min) and 10, 30, and 60 min after touch treatment.

Supplemental Table S10. Count of differentially expressed genes in wildtype and *SWIZ-OE* root tissue 10, 30, and 60 min after touch treatment and the count of differentially expressed genes shared between genotypes at each timepoint.

Supplemental Table S11. Calmodulin and calmodulin-like touch responsive gene expression in wildtype Bd21-3 root tissue.

Supplemental Table S12. XTH and XTH-like touch responsive gene expression in wildtype root tissue.

Supplemental Table S13. Glycoside hydrolase touch responsive gene expression in wildtype root tissue.

Supplemental Table S14. GO analysis of genes in iDREM identified pathways derived from differential gene expression following touch in wildtype and *SWIZ-OE* root tissue.

Supplemental Table S15. DAP-seq results of SWIZ genome binding locations.

Supplemental Table S16. All putative CREs enriched in differentially expressed genes at each timepoint identified using the k-mer approach.

Supplemental Table S17. CREs described for Arabidopsis thaliana in O'Malley et al. 2016 that match CREs identified using the k-mer approach.

557
Supplemental Figure S1. SWIZ is a Group I bZIP highly expressed in maturing stem and
root. (A) *SWIZ* transcript abundance in *Brachypodium distachyon* leaf, root, and stem tissue
measured by microarray. Mean +/- standard deviation of three biological replicates. (B)
Phylogeny analysis of amino acid sequences from *B. distachyon* (green), *Oryza sativa* (blue),
Arabidopsis thaliana (red), *Nicotiana tabacum* (orange) shows *SWIZ* (black dot). The phylogeny
was reconstructed using maximum likelihood analysis with a bootstrap resampling value of
1000. The numbers labeled on the branches are the posterior probability to support each clade.
The protein sequences, full locus IDs, and plant species with genome source are provided in
Supplemental File S1.

Supplemental Figure S2. Description of *SWIZ* transgenic reagents. (A) Diagram of *SWIZ*
overexpression transgenes. The *ZmUbi* promoter was used to drive expression of the *SWIZ*
coding sequence, either alone or fused in frame with *eGFP*. (B) Transcript abundance of *SWIZ*
measured by RNA-sequencing. Root tissue was sampled just prior to touch (t = 0), and at 10, 30,
and 60 min following touch treatment in wildtype and *SWIZ-OE*. LB, left border; *ZmUbi* prom,
maize ubiquitin promoter; Hyg, hygromycin phosphotransferase gene; NOS, nopaline synthase
terminator; RB, right border.

Supplemental Figure S3. Network analysis of differential gene expression paths and enriched GO terms. (A) Path determinations from iDREM time course analysis. Each line represents a set of genes with similar expression level patterns over the time course relative to time 0, pre-touch. (B) Heatmap of selected GO term enrichment of genes from iDREM path analysis. Color indicates *q*-value score, with a cutoff of 0.1. Terms without a colored block are not significantly enriched in that path.

Supplemental Figure S4. Brachypodium distachyon cis-regulatory sequences most overrepresented at each time course after touching. Each nucleotide sequence is a position probability matrix motif derived from DNA-affinity purification sequencing and identified as enriched under different time courses. The height of the letter at each position is proportional to the probability of a given nucleotide.

Supplemental Figure S5. *SWIZ-OE* roots are shorter than wildtype with no significant difference in straightness in response to direct touch. Wildtype and *SWIZ-OE* seedlings grown on plates under control (A) or touched (B) conditions for five days. Touched plants were probed with a pipette tip twice a day. (C) Quantification of root length in control and touched conditions. Significance denoted by compact letter display reflecting Tukey HSD adjusted *p*-values < 0.05. (D) Quantification of root straightness in control and touch conditions. Significance denoted by * reflecting Wilcoxon sign-ranked test for non-parametric data with *p*-value < 0.05, ns = not significant. Scale bar = 1 cm.

599
Supplemental Figure S6. Wildtype roots shorten with increasing plate angle, while *SWIZ-OE* roots are consistently short, with no significant difference in straightness. Wildtype and *SWIZ-OE* seedlings grown on plates at (A) 10°, 20°, 30°, and (B) 40° incline from vertical. (C) Quantification of root length. (D) Quantification of primary root straightness. Significance denoted by compact letter display reflecting Tukey HSD adjusted *p*-values < 0.05. Scale bar = 1 cm.

Supplemental Figure S7. Thigmomatic, an automated system to administer touch stimulus.
(A) Overview of Thigmomatic inside a Percival PGC-15 growth chamber showing linear rail
based frame (1), gantry carts (2), NEMA17 stepper motor (3), 12V 18W AC/DC power supply
(4), and Raspberry Pi 3b microcomputer (5). (B) Thigmomatic making contact with a
Brachypodium distachyon plant.

Supplemental Figure S8. Brachypodium distachyon displays classic thigmomorphogenic phenotypes. One-week-old wildtype plants were treated with touch stimulus every 90 min. (A) Left to right, plants that experienced no stress, two weeks stress, three weeks stress and were then allowed to recover, imaged one week after the end of treatment (B) height, (C) aboveground non-grain biomass weight, and (D) branch number were measured at senescence. n = 4-5 plants per treatment. Significance denoted by compact letter display reflecting Tukey HSD adjusted *p*-values < 0.05. (E) Transverse sections of the peduncle or third internode were taken from control, 2 week stressed, and 3 week stressed plants and stained with phloroglucinol-HCl. (F) Quantification of interfascicular fiber wall thickness. Scale bar = 100 µm. n = 3 plants per

622   treatment.

Supplemental Figure S9. Biomass and branching in SWIZ-OE under touch treatment
shows no difference from wildtype. Wildtype and SWIZ-OE plants were placed under control
conditions or received two weeks of mechanical stress every 90 min. After senescence, branch
number (A) and aboveground biomass weight (B) were quantified. Significant differences were
not observed following ANOVA and Tukey HSD testing.

Supplemental Figure S10. *SWIZ-OE* **marginally enhanced thigmomorphogenic response in
stems.** Wildtype and *SWIZ-OE* plants were placed under control conditions or received two
weeks of mechanical stress every 90 min. (A) Quantification of main stem height at senescence.
(B) Quantification of interfascicular fiber wall thickness in transverse cross sections of the
peduncle. n = 4 to 6 plants per genotype, per treatment. (C) Representative cross sections stained
with phloroglucinol-HCl. Scale bar = 100 µm. Significance denoted by compact letter display
reflecting Tukey HSD adjusted *p*-values < 0.05.

Supplemental Figure S11. Method of root touch treatment. A blunt probe formed from a
glass Pasteur pipette was used to gently tap on the root five times in areas like those indicated by
the orange arrows.

**Supplemental File S1. Gene names and amino acid sequences used in the SWIZ bZIP
Group I phylogenetic analysis.**

**Supplemental File S2. Movie of *GFP-OE* root images captured for 60-90 min post touch
treatment.**

**Supplemental File S3. Movie file of a *SWIZ:GFP-OE* root images captured for 60-90 min
post touch treatment.**

ACCESSION NUMBERS

AtbZIP18 (At2g40620), AtbZIP29 (At4g38900), AtbZIP52 (At1g06850), AtTCH1/AtCaM2
(At2g41110), AtTCH2/AtCML24 (At5g37770), AtTCH3/CML12 (At2g41100),
AtTCH4/AtXTH22 (At5g57560), CAD1 (Bradi3g17920), CESA4 (Bradi4g28350), COMT6
(Bradi3g16530), CSLF6 (Bradi3g16307), GA2ox3 (Bradi2g50280), SWAM3/MYB44
(Bradi1g30252), NAC35 (Bradi2g28020), NtRSGa (Niben101Scf01150g00005), NtRSGb
(Niben101Scf02191g00014), SWIZ (Bradi1g17700), VIP1 (At1g43700).

FINANCIAL SUPPORT

This work was supported by the Office of Science, Biological and Environmental Research,
Department of Energy (DE-FG02-08ER64700DE and DE-SC0006641), the National Science
Foundation Division of Integrative Organismal Systems (NSF IOS-1558072), and the United
States Department of Agriculture's National Institute of Food and Agriculture and Massachusetts
Agriculture Experiment Station (MAS00534) to S.P.H., the Constantine J. Gilgut Fellowship to
J.H.C, I.W.M., and P.P.H., and the Lotta M. Crabtree Fellowship to J.H.C., I.W.M., and K.J-
M.M. The work conducted by the U.S. Department of Energy Joint Genome Institute, a DOE
Office of Science User Facility, is supported by the Office of Science of the U.S. Department of
Energy operated under Contract No. DE-AC02-05CH11231. The microscopy data was gathered
in the Light Microscopy Facility and Nikon Center of Excellence at the Institute for Applied Life
Sciences, UMass Amherst with support from the Massachusetts Life Sciences Center.
670

671 **CONFLICT OF INTEREST**
672 None
673
DATA AVAILABILITY STATEMENT
The data that support the findings of this study are openly available at E-MTAB-10066 and E-
MTAB-10084 in the European Nucleotide Archive and https://hazenlab.shinyapps.io/swiztc.

Table 1. The number of SWIZ Dap-seq targets differentially expressed in wildtype plants after
touch treatment relative to untouched wildtype The location of SWIZ DAP-seq binding, in either
the promoter or gene body regions, is noted as is the direction of differential expression, up or
down.

Time after touch	Promoter		Gene body	
	Up	Down	Up	Down
(min)	percent (count)			
0	42.0 (103)	58.0 (142)	48.0 (286)	52.0 (310)
10	57.0 (73)	43.0 (55)	53.0 (167)	47.0 (148)
30	59.4 (79)	40.6 (54)	41.2 (136)	58.8 (194)
60	64.2 (77)	35.8 (43)	45.7 (128)	54.3 (152)

681
FIGURE LEGENDS

Figure 1. SWIZ translocates to the nucleus in response to mechanical stimulus, specifically
in regions directly stimulated. (A) Image of *SWIZ:GFP-OE* and *GFP-OE* roots prior to
stimulus and 30 min post stimulus. Roots were observed immediately following mechanical
perturbation. (B) Quantification of nuclear signal in control (purple) and touched (teal)
conditions for *GFP-OE* (left) and *SWIZ:GFP-OE* (right). n = 14-20 nuclei. (C) SWIZ

translocation occurred in the local area of the stimulus. At 30 min, stimulus was applied to an
upper region of the root, while at 120 min it was applied to a lower region approximately 3 cm
below. n = 109, 184 nuclei respectively for upper and lower regions. Scale bar = 100 µm, (D)
SWIZ:GFP-OE roots were imaged by confocal microscopy with stimulus applied in the field of
view at 0, 90, and 180 min. n = 126 nuclei. (B-D) Images were taken every 2 min. Nuclear GFP
signal was quantified in selected nuclei at each time point. The average nuclear GFP signal is
represented by the line with error bars indicating standard error of the mean. Scale bar = 100 µm.
n = 4-6 plants per treatment.

696
Figure 2. Transcriptome analysis of touch response in *Brachypodium distachyon* roots. (A)
Root tissue was sampled just prior to touch (t = 0), and at 10, 30, and 60 min following touch
treatment in wildtype and *SWIZ-OE*. (B) Principal component analysis of gene expression across
samples shows the greatest difference corresponding to genotype and the second greatest
corresponding to time after touch. (C) Canonical, as well as novel touch responsive genes are
upregulated in *B. distachyon* following touch. Closest orthologs of the *Arabidopsis thaliana*
TOUCH genes encoding calmodulin-like (CML) and xyloglucan endo-
transglycosylase/hydrolases (XTH) are upregulated following touch, as are previously
unreported members of the glycosyl hydrolase 17 (GH17) family. Full list of gene expression for
TOUCH and glycoside hydrolases defined by Tyler et al. (2010) in Supplemental Tables S10-12.
Significance denoted by * reflecting $q < 0.1$ compared to expression at t = 0, with q-values
representing Wald test p-values adjusted for false discovery rate.

709
Figure 3. Gene expression analysis of cell wall related genes. Log fold-change of gene
expression measured by RNA-seq in wildtype and *SWIZ-OE*, presented as relative to wildtype
expression at time 0, pre-touch. Bar color indicates class of cell wall gene. Error bars indicate
standard deviation of three biological replicates. Significance denoted by * reflecting $q < 0.1$
compared to wildtype expression at t = 0, with q-values representing Wald test p-values adjusted
for false discovery rate. Legend abbreviations: BAHD, BAHD (BEAT, AHCT, HCBT, and
DAT) acyltransferases; CESA, cellulose synthase; MLG, mixed-linkage glucans.

717
Figure 4. DNA affinity purification sequencing to determine SWIZ binding sites and
differential expression in response to touch and *SWIZ-OE*. (A) Top two most statistically
enriched sequence motifs in SWIZ binding sites. (B) Distribution of binding sites across genomic
features, relative to primary transcripts of the *Brachypodium distachyon* annotation v 3.1. (C)
Relative distribution of binding sites centered on the transcriptional start site (TSS, blue dashed
line), transcriptional termination site (TTS, red dashed line) represents the average length of all
annotated transcripts, approximately 4.5 kb away from the TSS. (D) Path determinations from
iDREM time course analysis of differentially expressed genes that also have DAP-seq binding
sites. Each line represents a set of genes with similar expression level patterns over the time
course relative to time 0, pre-touch. Wildtype: promoter and gene body = 170 and 146 genes,
respectively. *SWIZ-OE*: promoter = 163 genes, gene body A and B = 141 and 60, respectively.
(E-H) Gene expression over time of selected genes with SWIZ binding sites. Line graphs are the
average transcript abundance of three biological replicates for each time point. Binding site
determined as peaks of sequence alignment. Scale bar unit is bases. Direction of transcription is
shown with arrows on the gene model, 5' and 3' UTRs are depicted by narrowed rectangles on
the gene model.

734
735 **Figure 5. Sequence motifs enriched in the *cis*-regulatory regions of touch responsive**
736 ***Brachypodium distachyon* genes.** Negative log *p*-values for *cis*-elements, known and not known
737 to be touch responsive. RSRE - Rapid Stress Response Element, FAR1 - FAR-RED impaired
738 response1, GRF - Growth Regulating Factor, VNS - VND, NST/SND, SMB.
739
740

REFERENCES

Altartouri B, Bidhendi AJ, Tani T, Suzuki J, Conrad C, Chebli Y, Liu N, Karunakaran C, Scarcelli G, Geitmann A (2019) Pectin chemistry and cellulose crystallinity govern pavement cell morphogenesis in a multi-step mechanism. Plant Physiol **181**: 127–141

Andrews S (2010) FastQC: a quality control tool for high throuput sequence data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc:

Bidhendi AJ, Altartouri B, Gosselin FP, Geitmann A (2019) Mechanical stress initiates and sustains the morphogenesis of wavy leaf epidermal cells. Cell Rep **28**: 1237–1250.e6

Bidhendi AJ, Geitmann A (2018) Finite element modeling of shape changes in plant cells. Plant Physiol **176**: 41–56

Biro RL, Hunt ER, Erner Y, Jaffe MJ (1980) Thigmomorphogenesis: changes in cell division and elongation in the internodes of mechanically-perturbed or ethrel-treated bean plants. Ann Bot **45**: 655–664

Börnke F, Rocksch T (2018) Thigmomorphogenesis – Control of plant growth by mechanical stimulation. Sci Hortic **234**: 344–353

Braam J (2004) In touch: plant responses to mechanical stimuli. New Phytol **165**: 373–389

Braam J, Davis RW (1990) Rain-, wind-, and touch-induced expression of calmodulin and calmodulin-related genes in Arabidopsis. Cell **60**: 357–364

Chehab EW, Eich E, Braam J (2009) Thigmomorphogenesis: a complex plant response to mechano-stimulation. J Exp Bot **60**: 43–56

Coomey JH, Sibout R, Hazen SP (2020) Grass secondary cell walls, *Brachypodium distachyon* as a model for discovery. New Phytol **227**: 1709–1724

Coutand C, Martin L, Leblanc-Fournier N, Decourteix M, Julien J-L, Moulia B (2009) Strain mechanosensing quantitatively controls diameter growth and PtaZFP2 gene expression in poplar. Plant Physiol **151**: 223–232

Deppmann CD, Acharya A, Rishi V, Wobbes B, Smeekens S, Taparowsky EJ, Vinson C (2004) Dimerization specificity of all 67 B-ZIP motifs in *Arabidopsis thaliana*: a comparison to *Homo sapiens* B-ZIP motifs. Nucleic Acids Res **32**: 3435–3445

Ding J, Hagood JS, Ambalavanan N, Kaminski N, Bar-Joseph Z (2018) iDREM: Interactive visualization of dynamic regulatory networks. PLoS Comput Biol **14**: e1006019

Doherty CJ, Van Buskirk HA, Myers SJ, Thomashow MF (2009) Roles for Arabidopsis CAMTA transcription factors in cold-regulated gene expression and freezing tolerance. Plant Cell **21**: 972–984

Dröge-Laser W, Snoek BL, Snel B, Weiste C (2018) The Arabidopsis bZIP transcription factor family-an update. Curr Opin Plant Biol **45**: 36–49

Ehlert A, Weltmeier F, Wang X, Mayer CS, Smeekens S, Vicente-Carbajosa J, Dröge-Laser W (2006) Two-hybrid protein-protein interaction analysis in Arabidopsis protoplasts: establishment of a heterodimerization map of group C and group S bZIP transcription factors. Plant J **46**: 890–900

Fernández-Calvo P, Chini A, Fernández-Barbero G, Chico J-M, Gimenez-Ibanez S, Geerinck J, Eeckhout D, Schweizer F, Godoy M, Franco-Zorrilla JM, et al (2011) The Arabidopsis bHLH transcription factors MYC3 and MYC4 are targets of JAZ repressors and act additively with MYC2 in the activation of jasmonate responses. Plant Cell **23**: 701–715

Fukazawa J, Nakata M, Ito T, Matsushita A, Yamaguchi S, Takahashi Y (2011) bZIP transcription factor RSG controls the feedback regulation of NtGA20ox1 via intracellular localization and epigenetic mechanism. Plant Signal Behav **6**: 26–28

Fukazawa J, Nakata M, Ito T, Yamaguchi S, Takahashi Y (2010) The transcription factor RSG regulates negative feedback of NtGA20ox1 encoding GA 20-oxidase. Plant J **62**: 1035–1045

Fukazawa J, Sakai T, Ishida S, Yamaguchi I, Kamiya Y, Takahashi Y (2000) Repression of shoot growth, a bZIP transcriptional activator, regulates cell elongation by controlling the level of gibberellins. Plant Cell **12**: 901–915

Gibalová A, Steinbachová L, Hafidh S, Bláhová V, Gadiou Z, Michailidis C, Műller K, Pleskot R, Dupľáková N, Honys D (2017) Characterization of pollen-expressed bZIP protein interactions and the role of ATbZIP18 in the male gametophyte. Plant Reprod **30**: 1–17

Gladala-Kostarz A, Doonan JH, Bosch M (2020) Mechanical stimulation in *Brachypodium distachyon*: implications for fitness, productivity and cell wall properties. Plant Cell Environ **43**: 1314–1330

Grigoryan G, Keating AE (2006) Structure-based prediction of bZIP partnering specificity. J Mol Biol **355**: 1125–1142

Hamant O, Heisler MG, Jönsson H, Krupinski P, Uyttewaal M, Bokov P, Corson F, Sahlin P, Boudaoud A, Meyerowitz EM, et al (2008) Developmental patterning by mechanical signals in Arabidopsis. Science **322**: 1650–1655

Handakumbura P, Matos D, Osmont K, Harrington M, Heo K, Kafle K, Kim S, Baskin T, Hazen S (2013) Perturbation of *Brachypodium distachyon CELLULOSE SYNTHASE A4* or *7* results in abnormal cell walls. BMC Plant Biol **13**: 131

Handakumbura PP, Brow K, Whitney IP, Zhao K, Sanguinet KA, Lee SJ, Olins J, Romero-Gamboa SP, Harrington MJ, Bascom CJ, et al (2018) *SECONDARY WALL*

ASSOCIATED MYB1 is a positive regulator of secondary cell wall thickening in *Brachypodium distachyon* and is not found in the Brassicaceae. Plant J **96**: 485–699

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK (2010) Simple combinations of lineage-determining transcription factors prime *cis*-regulatory elements required for macrophage and B cell identities. Mol Cell **38**: 576–589

Igarashi D, Ishida S, Fukazawa J, Takahashi Y (2001) 14-3-3 proteins regulate intracellular localization of the bZIP transcriptional activator RSG. Plant Cell **13**: 2483–2497

Iida H (2014) Mugifumi, a beneficial farm work of adding mechanical stress by treading to wheat and barley seedlings. Front Plant Sci **5**: 453

Ishida S, Fukazawa J, Yuasa T, Takahashi Y (2004) Involvement of 14-3-3 signaling protein binding in the functional regulation of the transcriptional activator REPRESSION OF SHOOT GROWTH by gibberellins. Plant Cell **16**: 2641–2651

Ishida S, Yuasa T, Nakata M, Takahashi Y (2008) A tobacco calcium-dependent protein kinase, CDPK1, regulates the transcription factor REPRESSION OF SHOOT GROWTH in response to gibberellins. Plant Cell **20**: 3273–3288

Ito T, Ishida S, Oe S, Fukazawa J, Takahashi Y (2017) Autophosphorylation affects substrate-binding affinity of tobacco Ca2+-dependent protein kinase1. Plant Physiol **174**: 2457–2468

Ito T, Nakata M, Fukazawa J, Ishida S, Takahashi Y (2014) Scaffold function of Ca2+-dependent protein kinase: tobacco Ca2+-DEPENDENT PROTEIN KINASE1 transfers 14-3-3 to the substrate REPRESSION OF SHOOT GROWTH after phosphorylation. Plant Physiol **165**: 1737–1750

Jaffe MJ (1973) Thigmomorphogenesis: The response of plant growth and development to mechanical stimulation : With special reference to Bryonia dioica. Planta **114**: 143–157

Jaffe MJ, Biro R, Bridle K (1980) Thigmomorphogenesis: Calibration of the parameters of the sensory function in beans. Physiol Plant **49**: 410–416

Jakoby M, Weisshaar B, Dröge-Laser W, Vicente-Carbajosa J, Tiedemann J, Kroj T, Parcy F, bZIP Research Group (2002) bZIP transcription factors in Arabidopsis. Trends Plant Sci **7**: 106–111

Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. Brief Bioinform **20**: 1160–1166

Kim D, Langmead B, Salzberg SL (2015) HISAT: a fast spliced aligner with low memory requirements. Nat Methods **12**: 357

Kim W-C, Ko J-H, Han K-H (2012) Identification of a *cis*-acting regulatory motif recognized by MYB46, a master transcriptional regulator of secondary wall biosynthesis. Plant Mol

Biol **78**: 489–501

Leblanc-Fournier N, Martin L, Lenne C, Decourteix M (2014) To respond or not to respond, the recurring question in plant mechanosensitivity. Front Plant Sci **5**: 401

Lee D, Polisensky DH, Braam J (2005) Genome-wide identification of touch- and darkness-regulated Arabidopsis genes: a focus on calmodulin-like and XTH genes. New Phytol **165**: 429–444

Lee H-J, Kim H-S, Park JM, Cho HS, Jeon JH (2019) PIN-mediated polar auxin transport facilitates root obstacle avoidance. New Phytol. doi: 10.1111/nph.16076

Lee TG, Jang CS, Kim JY, Kim DS, Park JH, Kim DY, Seo YW (2006) A Myb transcription factor (TaMyb1) from wheat roots is expressed during hypoxia: roles in response to the oxygen concentration in root environment and abiotic stresses. Physiol Plant **129**: 375–385

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup (2009) The Sequence Alignment/Map format and SAMtools. Bioinformatics **25**: 2078–2079

Liu R, Finlayson SA (2019) Sorghum tiller bud growth is repressed by contact with the overlying leaf. Plant Cell Environ **42**: 2120–2132

Liu X, Chu Z (2015) Genome-wide evolutionary characterization and analysis of bZIP transcription factors and their expression profiles in response to multiple abiotic stresses in Brachypodium distachyon. BMC Genomics **16**: 227

Lobet G, Pagès L, Draye X (2011) A novel image-analysis toolbox enabling quantitative analysis of root system architecture. Plant Physiol **157**: 29–39

Lourenço TF, Serra TS, Cordeiro AM, Swanson SJ, Gilroy S, Saibo NJM, Oliveira MM (2015) The Rice E3-Ubiquitin Ligase HIGH EXPRESSION OF OSMOTICALLY RESPONSIVE GENE1 Modulates the Expression of ROOT MEANDER CURLING, a Gene Involved in Root Mechanosensing, through the Interaction with Two ETHYLENE-RESPONSE FACTOR Transcription Factors. Plant Physiol **169**: 2275–2287

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol **15**: 550

MacKinnon KJ-M, Cole BJ, Yu C, Coomey JH, Hartwick NT, Remigereau M-S, Duffy T, Michael TP, Kay SA, Hazen SP (2020) Changes in ambient temperature are the prevailing cue in determining *Brachypodium distachyon* diurnal gene regulation. New Phytol **227**: 1649–1667

Martin L, Leblanc-Fournier N, Julien J-L, Moulia B, Coutand C (2010) Acclimation kinetics of physiological and molecular responses of plants to multiple mechanical loadings. J Exp Bot **61**: 2403–2412

Matos DA, Whitney IP, Harrington MJ, Hazen SP (2013) Cell walls and the developmental
anatomy of the *Brachypodium distachyon* stem internode. PLoS One **8**: e80640

McCahill IW, Hazen SP (2019) Regulation of cell wall thickening by a medley of mechanisms.
Trends Plant Sci **24**: 853–866

Monshausen GB, Bibikova TN, Weisenseel MH, Gilroy S (2009) Ca2+ regulates reactive
oxygen species production and pH during mechanosensing in Arabidopsis roots. Plant Cell
21: 2341–2356

Moore BM, Lee YS, Wang P, Azodi C, Grotewold E, Shiu S-H (2022) Modeling temporal
and hormonal regulation of plant transcriptional response to wounding. Plant Cell **34**: 867–
888

Moulia B, Coutand C, Julien J-L (2015) Mechanosensitive control of plant growth: bearing the
load, sensing, transducing, and responding. Front Plant Sci **6**: 52

Nam BE, Park Y-J, Gil K-E, Kim J-H, Kim JG, Park C-M (2020) Auxin mediates the touch-
induced mechanical stimulation of adventitious root formation under windy conditions in
Brachypodium distachyon. BMC Plant Biol **20**: 335

Niez B, Dlouha J, Moulia B, Badel E (2019) Water-stressed or not, the mechanical acclimation
is a priority requirement for trees. Trees **33**: 279–291

Ohtani M, Nishikubo N, Xu B, Yamaguchi M, Mitsuda N, Goué N, Shi F, Ohme-Takagi M,
Demura T (2011) A NAC domain protein family contributing to the regulation of wood
formation in poplar. Plant J **67**: 499–512

Olins JR, Lin L, Lee SJ, Trabucco GM, MacKinnon KJM, Hazen SP (2018) Secondary wall
regulating NACs differentially bind at the promoter at a *CELLULOSE SYNTHASE A4* cis-
eQTL. Front Plant Sci **9**: 1895

Oliva M, Dunand C (2007) Waving and skewing: how gravity and the surface of growth media
affect root development in Arabidopsis. New Phytol **176**: 37–43

O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti
A, Ecker JR (2016) Cistrome and epicistrome features shape the regulatory DNA
landscape. Cell **165**: 1280–1292

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL (2015)
StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat
Biotechnol **33**: 290–295

Pitzschke A, Djamei A, Teige M, Hirt H (2009) VIP1 response elements mediate mitogen-
activated protein kinase 3-induced stress gene expression. Proc Natl Acad Sci U S A **106**:
18414–18419

Polisensky DH, Braam J (1996) Cold-shock regulation of the Arabidopsis TCH genes and the

effects of modulating intracellular calcium levels. Plant Physiol **111**: 1271–1279

Pomiès L, Decourteix M, Franchel J, Moulia B, Leblanc-Fournier N (2017) Poplar stem
transcriptome is massively remodelled in response to single or repeated mechanical stimuli.
BMC Genomics **18**: 300

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) g:Profiler: a
web server for functional enrichment analysis and conversions of gene lists. Nucleic Acids
Res **47**: W191–W198

Ringli C, Keller B (1998) Specific interaction of the tomato bZIP transcription factor VSF-1
with a non-palindromic DNA sequence that controls vascular gene expression. Plant Mol
Biol **37**: 977–988

Roignant J, Badel É, Leblanc-Fournier N, Brunel-Michac N, Ruelle J, Moulia B,
Decourteix M (2018) Feeling stretched or compressed? The multiple mechanosensitive
responses of wood formation to bending. Ann Bot **121**: 1151–1161

Rushton PJ, Reinstädler A, Lipka V, Lippok B, Somssich IE (2002) Synthetic plant
promoters containing defined regulatory elements provide novel insights into pathogen- and
wound-induced signaling. Plant Cell **14**: 749–762

Schütze K, Harter K, Chaban C (2008) Post-translational regulation of plant bZIP factors.
Trends Plant Sci **13**: 247–255

Shah L, Yahya M, Shah SMA, Nadeem M, Ali A, Ali A, Wang J, Riaz MW, Rehman S, Wu
W, et al (2019) Improving lodging resistance: using wheat and rice as classical examples.
Int J Mol Sci. doi: 10.3390/ijms20174211

Sibout R, Proost S, Hansen BO, Vaid N, Giorgi FM, Ho-Yue-Kuang S, Legée F, Cézart L,
Bouchabké-Coussa O, Soulhat C, et al (2017) Expression atlas and comparative
coexpression network analyses reveal important genes involved in the formation of lignified
cell wall in *Brachypodium distachyon*. New Phytol **215**: 1009–1025

Swanson SJ, Barker R, Ye Y, Gilroy S (2015) Evaluating mechano-transduction and touch
responses in plant roots. Methods Mol Biol **1309**: 143–150

Szczegielniak J, Borkiewicz L, Szurmak B, Lewandowska-Gnatowska E, Statkiewicz M,
Klimecka M, Cieśla J, Muszyńska G (2012) Maize calcium-dependent protein kinase
(ZmCPK11): local and systemic response to wounding, regulation by touch and components
of jasmonate signaling. Physiol Plant **146**: 1–14

Tamura T, Endo H, Suzuki A, Sato Y, Kato K, Ohtani M, Yamaguchi M, Demura T (2019),
Affinity-based high-resolution analysis of DNA binding by VASCULAR-RELATED NAC-
DOMAIN7 via fluorescence correlation spectroscopy. Plant J **100**: 298-313

Trabucco GM, Matos DA, Lee SJ, Saathoff A, Priest H, Mockler TC, Sarath G, Hazen SP
(2013) Functional characterization of cinnamyl alcohol dehydrogenase and caffeic acid *O-*

methyltransferase in *Brachypodium distachyon*. BMC Biotechnol **13**: 61

Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ (2016) W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids Res **44**: W232–W235

Tsugama D, Liu S, Fujino K, Takano T (2018) Calcium signalling regulates the functions of the bZIP protein VIP1 in touch responses in *Arabidopsis thaliana*. Ann Bot **122**: 1219–1229

Tsugama D, Liu S, Takano T (2012) A bZIP protein, VIP1, is a regulator of osmosensory signaling in Arabidopsis. Plant Physiol **159**: 144–155

Tsugama D, Liu S, Takano T (2016) The bZIP protein VIP1 Is involved in touch responses in Arabidopsis roots. Plant Physiol **171**: 1355–1365

Tsugama D, Liu S, Takano T (2014) Analysis of functions of VIP1 and its close homologs in osmosensory responses of *Arabidopsis thaliana*. PLoS One **9**: e103930

Tyler L, Bragg JN, Wu J, Yang X, Tuskan GA, Vogel JP (2010) Annotation and comparative analysis of the glycoside hydrolase genes in *Brachypodium distachyon*. BMC Genomics **11**: 600

Uyttewaal M, Burian A, Alim K, Landrein B, Borowska-Wykręt D, Dedieu A, Peaucelle A, Ludynia M, Traas J, Boudaoud A, et al (2012) Mechanical stress acts via katanin to amplify differences in growth rate between adjacent cells in Arabidopsis. Cell **149**: 439–451

Valdivia ER, Herrera MT, Gianzo C, Fidalgo J, Revilla G, Zarra I, Sampedro J (2013) Regulation of secondary wall synthesis and cell death by NAC transcription factors in the monocot *Brachypodium distachyon*. J Exp Bot **64**: 1333–1343

Van Leene J, Blomme J, Kulkarni SR, Cannoot B, De Winne N, Eeckhout D, Persiau G, Van De Slijke E, Vercruysse L, Vanden Bossche R, et al (2016) Functional characterization of the Arabidopsis transcription factor bZIP29 reveals its role in leaf and root development. J Exp Bot **67**: 5825–5840

Van Moerkercke A, Duncan O, Zander M, Šimura J, Broda M, Vanden Bossche R, Lewsey MG, Lama S, Singh KB, Ljung K, et al (2019) A MYC2/MYC3/MYC4-dependent transcription factor network regulates water spray-responsive gene expression and jasmonate levels. Proc Natl Acad Sci U S A **116**: 23345–23356

Walley J, Coughlan S, Hudson ME, Covington MF, Kaspi R, Banu G, Harmer SL, Dehesh K (2007) Mechanical stress induces biotic and abiotic stress responses via a novel cis-element. PLoS Genet **3**: 1800–1812

Weltmeier F, Ehlert A, Mayer CS, Dietrich K, Wang X, Schütze K, Alonso R, Harter K, Vicente-Carbajosa J, Dröge-Laser W (2006) Combinatorial control of Arabidopsis proline dehydrogenase transcription by specific heterodimerisation of bZIP transcription factors. EMBO J **25**: 3133–3143

987   **Yin Y, Zhu Q, Dai S, Lamb C, Beachy RN** (1997) RF2a, a bZIP transcriptional activator of the
988   phloem-specific rice tungro bacilliform virus promoter, functions in vascular development.
989   EMBO J **16**: 5247–5259

990   **Zha G, Wang B, Liu J, Yan J, Zhu L, Yang X** (2016) Mechanical touch responses of
991   Arabidopsis TCH1-3 mutant roots on inclined hard-agar surface. International Agrophysics
992   **30**: 65

993   **Zhong R, Lee C, McCarthy RL, Reeves CK, Jones EG, Ye Z-H** (2011) Transcriptional
994   activation of secondary wall biosynthesis by rice and maize NAC and MYB transcription
995   factors. Plant Cell Physiol **52**: 1856–1871

996   **Zhong R, Ye Z-H** (2012) MYB46 and MYB83 bind to the SMRE sites and directly activate a
997   suite of transcription factors and secondary wall biosynthetic genes. Plant Cell Physiol **53**:
998   368–380