Touch-triggered bZIP translocation regulates elongation and secondary wall biosynthesis

Joshua H. Coomey[1,2], Kirk J.-M. MacKinnon[1,3], Ian W. McCahill[1,2], Pubudu P. Handakumbura[1,2], Gina M. Trabucco[1,3], Jessica Mazzola[1], Nicole A. Leblanc[1], Rithany Kheam[1], Miriam Hernandez-Romero[1,2], Kerrie Barry[4], Lifeng Liu[4], Ji E. Lee[4], John P. Vogel[4], Ronan C. O'Malley[4], James J. Chambers[5], Samuel P. Hazen[1,2,3]

[1]Biology Department, University of Massachusetts, Amherst, MA 01003, USA,

[2]Plant Biology Graduate Program, University of Massachusetts, Amherst, MA 01003, USA,

[3]Molecular and Cellular Biology Graduate Program, University of Massachusetts, Amherst, MA 01003, USA,

[4]DOE Joint Genome Institute, Berkeley, CA 94720, USA,

[5]Institute for Applied Life Science, University of Massachusetts, Amherst, MA 01003, USA,

Author for correspondence:
Samuel P. Hazen, hazen@bio.umass.edu

ORCID
Joshua H. Coomey, https://orcid.org/0000-0001-5866-6671
Pubudu P. Handakumbura, https://orcid.org/0000-0002-1708-8315
Kerrie Barry, https://orcid.org/0000-0002-8999-6785
Ji E. Lee, https://orcid.org/0000-0001-8325-1689
John P. Vogel, https://orcid.org/0000-0003-1786-2689
James J. Chambers, https://orcid.org/0000-0003-3883-8215
Samuel P. Hazen, https://orcid.org/0000-0001-9131-8994

AUTHOR CONTRIBUTIONS
JHC and SH conceived and designed the study. JHC, IM, JJC, MH-R, PH, developed techniques and reagents. GT, IM, JHC, JL, JM, JV, KB, LL, NL, PH, RK, , and RO acquired the data. JHC, KM, and SH analyzed and interpreted the data. JHC, KM, IM, SH drafted the manuscript. SH agrees to serve as the author responsible for contact and ensures communication.

Abstract

Plant growth is mediated by the integration of internal and external cues, perceived by cells and transduced into a developmental program that gives rise to cell division, elongation, and wall thickening. Extra-, inter-, and intra- physical cellular forces contribute to this regulation. Across the plant kingdom thigmomorphogenesis is widely observed to alter plant morphology by reducing stem height and increasing stem diameter. The transcriptome is highly responsive to touch, including components of calcium signaling pathways and cell wall modification. One aspect of this cascade involves gibberellins (GA) and bZIP family transcription factors. Here, we present data connecting thigmomorphogenesis with secondary cell wall synthesis by GA inactivation and bZIP translocation into the nucleus. *Brachypodium distachyon* SECONDARY WALL INTERACTING bZIP (SWIZ) protein translocated into the nucleus in response to mechanical stimulation. This translocation was mitigated by supplementing with exogenous bioactive GA, and induced by chemical inhibition of GA synthesis. Misregulation of *SWIZ* expression resulted in plants with reduced stem and root elongation, and following mechanical stimulation, increased secondary wall thickness. Classical touch responsive genes were upregulated in *B. distachyon* following touch, and we observe significant induction of the glycoside hydrolase 17 family, which may be indicative of a facet of grass thigmomorphogenesis. SWIZ protein binding to an E-box variant in exons and introns was associated with immediate activation followed by repression of gene expression. Thus, SWIZ is a transcriptional regulatory component of thigmomorphogenesis, which includes the thickening of secondary cell walls.

INTRODUCTION

Forces both internal and external to a cell influence growth. Turgor pressure in conjunction with anisotropic cell wall dynamics determine direction of growth and thus cell shape. Force perception between neighboring cells plays a critical role in the development and maintenance of tissue form and function, such as the lobed, interlocking, pavement cells on the leaf epidermis, or the developmental hotspots in the apical meristem (Hamant et al., 2008; Uyttewaal et al., 2012; Bidhendi et al., 2019). Specific inter-cell forces result in dynamic remodelling of the cortical cytoskeleton, with subsequent changes in cellulose microfibril alignment and alterations to other cell wall components such as pectin methyl esterification (Hamant et al., 2008; Uyttewaal et al., 2012; Bidhendi and Geitmann, 2018; Altartouri et al., 2019; Bidhendi et al., 2019). The classic hallmarks of touch responsive growth, or thigmomorphogenesis, include reduced plant height, increased radial growth in plants with a cambial meristem, increased branching, and delayed flowering time (Jaffe, 1973; Biro et al., 1980; Jaffe et al., 1980; Braam, 2004). These attributes have been leveraged by farmers for hundreds of years. As early as 1680 records show Japanese farmers tread on young wheat and barley seedlings to elicit increased branching, spikes per plant,

and grain weight per plant, along with stronger roots (Iida, 2014). This practice, known as
mugifumi, continues today with mechanized rollers.

Calcium is a critical cellular signaling molecule, and cytosolic Ca^{2+} fluctuations are induced by
mechanical stimuli and implicated in mechanosignaling (Monshausen et al., 2009; Szczegielniak
et al., 2012). Mechanical stimulus can significantly remodel gene expression (Braam and Davis,
1990; Braam, 2004; Lee et al., 2005). The so-called *TOUCH* (*TCH*) genes in *Arabidopsis*
thaliana, encode calmodulin (*AtTCH1*/*AtCaM2*), calmodulin-like proteins (*AtTCH2*/*AtCML24*,
AtTCH3/*CML12*), and a xyloglucan endotransglucosylase/hydrolase (*AtTCH4*/*AtXTH22*) (Braam
and Davis, 1990). Touch responsive gene expression patterns often overlap with other stimuli
such as dark, cold, and hormone treatment (Polisensky and Braam, 1996; Lee et al., 2005). In
addition to calcium binding and signaling, genes related to cell wall modification and a variety of
transcription factors and kinases are regulated by mechanical stimulus, as well as genes involved
in hormone homeostasis and signaling.

The dynamic control of bioactive cellular gibberellic acid (GA) is linked to
thigmomorphogenesis. GA is a growth promoting hormone derived through multiple enzymatic
steps from the diterpene pathway, with interconvertible bioactive and inactive species. The
bioactive forms of GA interact with protein receptors that regulate transcription factors
controlling growth. Mechanical force application results in the inactivation of cellular GA by
GA20 oxidases (Lange and Lange, 2015). The addition of exogenous bioactive GA to
mechanically stressed plants remediates their height phenotype, and mutations in *Atga20ox* do
not show thigmomorphogenic phenotypes under mechanical stress treatment (Lange and Lange,
2015). Recent work studying the translocation dynamics of several bZIP family transcription
factors has provided new links for calcium and GA signaling in mechanoperception signaling.

Group I bZIPs are also implicated in mechanosensing. VIRE2-INTERACTING PROTEIN 1
(AtVIP1) and related Group I bZIP proteins translocate from the cytoplasm to the nucleus in
response to a variety of biotic and abiotic stimuli, including hypo-osmotic conditions (Tsugama
et al., 2012; Tsugama et al., 2014; Tsugama et al., 2016). This translocation appears to be
dependent on protein phosphorylation, either from MITOGEN ACTIVATED PROTEIN
KINASE 3 during pathogen invasion, or via calcium dependent protein kinases (CDPK). AtVIP1
interacts with calmodulins, the calcium binding proteins involved in many calcium signaling
events. The Group I bZIP *Nt REPRESSOR OF SHOOT GROWTH* (*NtRSG*) in tobacco plays a
role in maintaining GA homeostasis, wherein it translocates from the cytoplasm to the nucleus in
response to cellular bioactive GA levels (Igarashi et al., 2001; Ishida et al., 2008; Fukazawa et
al., 2010). Under standard conditions, NtRSG is largely localized in the cytoplasm, and to some
extent in the nucleus, where it directly activates expression of *NtEnt-kaurene oxidase*, an enzyme
at an early step in the GA biosynthesis pathway. After cellular GAs were chemically inhibited,
NtRSG translocated to the nucleus and also activated *NtGA20oxidase1* (*NtGA20ox1*), an enzyme
further downstream in the GA biosynthesis pathway that is in part responsible for converting GA
species to their bioactive form (Fukazawa et al., 2010; Fukazawa et al., 2011). This suggests that

nuclear-localized NtRSG acts to promote bioactive GA synthesis in response to low GA
conditions. As with AtVIP1, NtRSG translocation relies on CDPK mediated phosphorylation
(Ishida et al., 2008; Ito et al., 2017). Both AtVIP1 and NtRSG associate with 14-3-3 proteins in
the cytoplasm while phosphorylated (Ishida et al., 2004; Ito et al., 2014; Ito et al., 2017;
Tsugama et al., 2018). Phosphatase activity causes NtRSG and the related AtbZIP29 to
dissociate from the 14-3-3 protein and enter the nucleus (Van Leene et al., 2016). *NtRSG* is
named for the dwarf phenotype observed when a dominant negative form of the protein is
overexpressed. In these dwarf plants, there were reduced levels of bioactive GA and reduced
elongation of stem internode cells (Fukazawa et al., 2000).

The development of secondary cell walls is a distinct and critical developmental process in plant
growth. Following cellular elongation, some cells, particularly in vascular and structural tissues,
increase their mechanical strength by depositing a thick layer of rigid material between the
plasma membrane and primary cell wall. These secondary walls are made of crystalline cellulose
interwoven with hemicelluloses and phenolic lignin polymers. Although functionally similar,
secondary walls in monocotyledonous plants have key differences from those in eudicots,
including distinct hemicellulose chemistry and differences in lignin biosynthesis. Grasses also
produce mixed-linkage glucans (MLGs), a wall polysaccharide that is rarely found outside the
commelinid monocots (Coomey et al., 2020). The synthesis and deposition of secondary wall
polymers is tightly controlled spatially and temporally by a well defined transcriptional
regulatory network. Evidence in both eudicots and grasses suggest a system of feed-forward
loops regulating transcription of wall synthesizing enzymes, with NAC family transcription
factors activating wall synthesis genes as well as MYB family and other transcription factors that
further promote secondary wall synthesis (McCahill and Hazen, 2019). These networks are very
similar between grasses and eudicots, with some components in each that have yet to be
described in the other. There are currently no bZIP family members in any secondary wall
regulatory model.

Beyond differences at the chemical and regulatory levels, grasses employ a fundamentally
different growth mechanism than eudicots. Grasses do not have a cambium layer, and thus no
lateral meristem. Stem elongation comes from cellular division and elongation in a discrete series
of intercalary meristems, called nodes, with one internode region elongating and pushing up
subsequent nodes. Detailed studies of thigmomorphogenesis have been conducted almost
exclusively in eudicots However, recent work in the model cereal grass *Brachypodium*
distachyon shows general overlap with conventional dicot thigmomorphogenesis, albeit with
some notable differences, such as no change in stem diameter and increased time to flower
(Gladala-Kostarz et al., 2020).

Thigmomorphogenesis is a widely observed phenomenon that results in reduced height,
increased radial growth, and increased branching. The mechanisms behind this form of growth
are not yet fully understood, but involve aspects of hormone regulation, Ca^{2+} signaling, Group I
bZIP intracellular translocation, and changes in gene expression. Here we describe the

transcriptional response to mechanical stimulation and the function of a *B. distachyon* bZIP

transcription factor, SECONDARY WALL ASSOCIATED bZIP (Bradi1g17700) and its role in

touch response and cell wall biosynthesis.

RESULTS

SWIZ is a Group I bZIP transcription factor and candidate cell wall regulator

To identify genes involved in the regulation of secondary cell wall thickening, transcript abundance was measured in *B. distachyon* leaf, root, and stem tissue (Trabucco et al., 2013). A gene annotated as a bZIP transcription factor, Bradi1g17700, was highly expressed in root and stem relative to leaf (**Fig 1A**). Bradi1g17700 is also a member of a 112 gene coexpression network (**Supplemental Table S1**) that includes genes highly expressed in the peduncle (Sibout et al., 2017). Phylogenetic analysis of Bradi1g17700, hereinafter referred to as *SECONDARY WALL INTERACTING bZIP* (*SWIZ*), amino acid sequence shows it to be an ortholog of the *A. thaliana* Group I bZIPs (Jakoby et al., 2002; Dröge-Laser et al., 2018), and closely related to *AtbZIP18* and *AtbZIP52* (**Fig 1B**).

SWIZ translocates into the nucleus in response to mechanical stimulus

Group I bZIPs in *A. thaliana* have been described as playing intersectional roles in regulating growth and development, and in doing so display dynamic translocation between the cytosol and the nucleus in response to external cellular force. As an ortholog of these mechanotropic proteins, we hypothesized that SWIZ protein may similarly translocate within the cell in response to mechanical force and contribute to plant growth regulation. To test this, the roots of transgenic plants overexpressing either *SWIZ* fused to *GFP* (*SWIZ:GFP-OE*) or *GFP* alone (*GFP-OE*) were observed following a mechanical stimulus (**Fig 2A**). In control plants, GFP signal was present in both the cytosol and nucleus, which remained constant over the imaging period (**Fig 2B**). GFP signal was mostly observed in the cytosol in *SWIZ:GFP-OE* plants, but following mechanical stimulus, nuclear GFP signal increased substantially, reaching a peak around 30 min post stimulus and returned to near basal levels by about 60 min (**Fig 2B**).

The dynamics and repeatability of SWIZ nuclear translocation were further investigated by sequential stimulus treatments. Touch response to stimulus can saturate at a certain number of treatments (Martin et al., 2010; Leblanc-Fournier et al., 2014; Moulia et al., 2015). To test if SWIZ translocation dynamics varied after repeated treatments, we applied mechanical force to *SWIZ:GFP-OE* roots at regular intervals. A second stimulus was given 90 min after the first, and a third at 180 min. Following each mechanical stimulation, SWIZ consistently translocated from cytoplasm to nucleus (**Fig 2D**). This suggests that SWIZ translocation dynamics are not impacted by repeated stimulus events 90 min apart.

When conducting these translocation assays, root tissue in the field of view was treated with mechanical force and nuclei in that region were tracked and quantified for fluorescence. To determine if the signal triggering SWIZ translocation is spread beyond the specifically stimulated region, two regions of the same *SWIZ:GFP-OE* root separated by 3 cm were simultaneously observed. The stimulated region showed typical SWIZ:GFP nuclear signal accumulation and in the region below no translocation was observed (**Fig 2C**). At 120 min, the treatments were

Figure 1. SWIZ is a Group I bZIP highly expressed in maturing stem and root. (A) *SWIZ* transcript abundance in *Brachypodium distachyon* leaf, root, and stem tissue measured by microarray. Mean +/- standard deviation of three biological replicates. (B) Phylogeny analysis of amino acid sequences from *B. distachyon* (green), *Oryza sativa* (blue), *Arabidopsis thaliana* (red), *Nicotiana tabacum* (orange) shows *SWIZ* (black dot). The phylogeny was reconstructed using maximum likelihood analysis with a bootstrap resampling value of 1000. The numbers labeled on the branches are the posterior probability to support each clade. The protein sequences, full locus IDs, and plant species with genome source are provided in Supplemental File 1.

192   reversed, with the lower root region receiving a stimulus while the upper region was

193   unperturbed. The lower region showed SWIZ:GFP nuclear translocation while the upper region

194   did not. Thus, the touch stimulus perceived by SWIZ is localized directly to the perturbed region.

195   **Transcriptional response to touch**

Figure 2. SWIZ translocates to the nucleus in response to mechanical stimulus, specifically in regions directly stimulated. (A) Image of *SWIZ:GFP-OE* and *GFP-OE* roots prior to stimulus and 30 min post stimulus. Roots were observed immediately following mechanical perturbation. (B) Quantification of nuclear signal. n = 14-20 nuclei. (C) SWIZ translocation occured in the local area of the stimulus. At 30 min, stimulus was applied to an upper region of the root, while at 120 min it was applied to a lower region approximately 3 cm below. n = 109, 184 nuclei respectively for upper and lower regions. (D) *SWIZ:GFP-OE* roots were imaged by confocal microscopy with stimulus applied in the field of view at 0, 90, and 180 min. n = 126 nuclei. (B-D) Images were taken every 2 min. Nuclear GFP signal was quantified in selected nuclei at each time point. The average nuclear GFP signal is represented by the line with error bars indicating standard error of the mean.

196   Having established the nuclear translocation of SWIZ in response to touch, we then investigated
197   what effect touch and SWIZ overabundance during touch may have on gene expression. To
198   assess this, we measured transcript abundance by sequencing mRNA from wildtype and *SWIZ-*

Figure 3. Transcriptome analysis of touch response in *Brachypodium distachyon* roots. (A) Root tissue was sampled just prior to touch (t = 0), and at 10, 30, and 60 min following touch treatment in wildtype and *SWIZ-OE.* (B) Principal component analysis of gene expression across samples shows the greatest difference corresponding to genotype and the second greatest corresponding to time after touch. (C) Canonical, as well as novel touch responsive genes are upregulated in *B. distachyon* following touch. Closest orthologs of the *Arabidopsis thaliana TOUCH* genes encoding calmodulin-like (CML) and xyloglucan endo-transglycosylase/hydrolases (XTH) are upregulated following touch, as are previously unreported members of the glycosyl hydrolase 17 (GH17) family. Full list of gene expression for *TOUCH* and glycoside hydrolases defined by Tyler et al. (2010) in Supplemental Tables 5, 6,and 7. Significance denoted by * reflecting q < 0.1 compared to expression at t = 0, with q-values representing Wald test p-values adjusted for false discovery rate.

OE root tissue just prior to touch (0 min) and at 10, 30, and 60 min following touch treatment (**Fig 3A**). Principal component analysis of transcript abundance shows the greatest amount of variance is attributed to the first principal component, 60%, where samples cluster based on

genotype (**Fig 3B**). The second principal component, which accounts for 17% of the variance, distinguished between pre-touch and 10 min following touch, with the last two time points clustering similarly.

The wildtype transcriptome was massively remodeled in response to touch, with 8,902 transcripts differentially expressed (q < 0.1) at 10 min post touch, 5,682 transcripts at 30 min, and 7,672 transcripts at 60 min (**Supplemental Tables 2-4**). Touch response studies in *A. thaliana* describe similar changes that include numerous calmodulin (CAM), calmodulin-like (CmL), and xyloglucan endo-transglycosylase/hydrolase (XTH) activity (**Fig 3C**)(Braam and Davis, 1990; Lee et al., 2005). Based on homology with *TCH1*, *TCH2*, and *TCH3*, 79 CAM and CmL genes were identified, the majority of which showed upregulation in wildtype plants following touch treatment (**Supplemental Table 5**). Similarly, homology with *TCH4* identified 37 *B. distachyon XTH* and *XTH-like* genes. Of these, 8 genes were upregulated by touch in wildtype plants, including the two with the highest similarity to *TCH4* (Bradi1g33810 and Bradi33840) (**Fig 3C**, **Supplemental Table 6**). Since xyloglucans are in relatively low abundance in grass cell walls compared to eudicots, we also investigated the effect of touch on other families of the glycosyl hydrolase superfamily (Tyler et al., 2010). In particular, GH17 genes were differentially regulated following touch. The GH17 family is broadly described as modifying β-1,3-glucans. We observed measurable transcript abundance for 35 of the 53 annotated GH17 family members, with 8, 11, and 16 members showing differential expression at 10, 30 and 60 min respectively. Fisher's exact test determined significant enrichment of GH17 expression at time 60 (*p* = 9.307e-3) (**Fig 3C**, **Supplemental Table 7**). Online resources are available to explore the RNA-seq data at (https://hazenlab.shinyapps.io/swiztc/), which provides an interactive platform for interrogating the *B. distachyon* touched-root transcriptome.

Cell wall genes are downregulated immediately following touch, then induced concurrent with SWIZ nuclear translocation and more strongly in *SWIZ-OE*
Genes related to secondary cell wall synthesis were immediately downregulated in wildtype plants following touch, but were subsequently upregulated (**Fig 4**). Xylan, cellulose, MLG, and callose synthesis associated genes were upregulated 30 and 60 min following touch, while lignin biosynthesis and BAHD acyltransferase genes were upregulated at 60 min. Network analysis of gene expression patterns over time using the iDREM software showed that in *SWIZ-OE* plants, cell wall genes were more rapidly induced (**Supplemental Fig 2A-B**). Prior to touch, nearly all of the cell wall associated genes investigated were significantly downregulated relative to wildtype, then significantly upregulated following touch (**Fig 4**). In *SWIZ-OE*, cell wall related terms were enriched in paths that had stronger upregulation earlier than the same terms in wildtype (**Supplemental Fig 2A-B**). Notably, aspects of cell wall polysaccharide biosynthesis, particularly glucan biosynthesis, were significantly enriched in *SWIZ-OE* path D, while not significantly enriched in wildtype (**Supplemental Table 8**). This is mirrored when looking at specific gene expression, with primary wall associated CESAs, MLG, and callose synthesis

Figure 4. Gene expression analysis of cell wall related genes. Log fold-change of gene expression measured by RNA-seq in wildtype and *SWIZ-OE*, presented as relative to wildtype expression at time 0, pre-touch. Bar color indicates class of cell wall gene. Error bars indicate standard deviation of three biological replicates. Significance denoted by * reflecting q < 0.1 compared to wildtype expression at t = 0, with q-values representing Wald test p-values adjusted for false discovery rate.

242 genes all significantly upregulated following touch in *SWIZ-OE*. Thus, cell wall genes were
243 immediately repressed by touch, followed by significant activation. That activation occurs more
244 rapidly in *SWIZ-OE* plants and is associated with the timing of SWIZ protein nuclear
245 translocation.
246

SWIZ protein binding in the gene body is associated with dynamic changes in gene
expression
Next we investigated the direct binding targets of SWIZ protein by performing DNA affinity
purification sequencing (DAP-seq) with whole genomic DNA. In total, SWIZ interacted with
2,455 distinct positions in the genome. Those regions were significantly enriched for two
sequence motifs, (A/C)CAGNCTG(T/G) and (A/C)CAGCTG(T/G) that are variants of the E-box
motif (**Fig 5A**). A plurality of the binding sites were between the translational start and stop sites
with 21% of the peaks in introns and 21% in exons (**Fig 5B**). Fewer than 10% of the peaks
occurred in UTRs and the promoter (defined as 5 kb upstream of the 5 'UTR) and intergenic
regions each accounted for 25% of the binding sites. Thus, SWIZ protein is preferentially bound
to an E-box like motif and often in the gene body (**Fig 5C**).
258

259 To investigate the effects of possible SWIZ binding on gene expression, we compared the genes
260 differentially expressed in the *SWIZ-OE* touch response RNA-seq time courses with the promoter
261 and gene body binding targets identified by the DAP-seq assay. Prior to touch, genes with

Figure 5. DNA affinity purification sequencing to determine SWIZ binding sites. (A) Top two most statistically enriched sequence motifs in SWIZ binding sites. (B) Distribution of binding sites across genomic features, relative to primary transcripts of the *Brachypodium distachyon* annotation v 3.1. (C) Relative distribution of binding sites centered on the transcriptional start site (TSS, blue dashed line), transcriptional termination site (TTS, red dashed line) represents the average length of all annotated transcripts, approximately 4.5 kb away from the TSS. (D) Path determinations from iDREM time course analysis of differentially expressed genes that also have DAP-seq binding sites. Each line represents a set of genes with similar expression level patterns over the time course relative to time 0, pre-touch.

promoter-SWIZ protein interactions were most often downregulated in *SWIZ-OE* relative to
wildtype (**Table 1**). However, some SWIZ targets behaved differently following touch. SWIZ
promoter binding targets were most often upregulated in *SWIZ-OE* in all three post-touch time
points. Thus, SWIZ promoter binding was coincident with repression, but following a touch
stimulus, activation was more prominent. However, this difference between untreated and
touched was not as pronounced in gene body targets, with more upregulated genes 10 min
following touch and more downregulated 30 and 60 min following touch. To further explore
these trends we conducted iDREM cluster analysis of the differentially expressed genes that
were also SWIZ protein targets (**Fig 5D**). For both wildtype and *SWIZ-OE*, the network analysis

revealed a pathway to increased gene expression by 60 min following touch. The network
pathway trend for both promoter and gene binding targets in wildtype was immediate repression
at 10 min followed by increased expression at 30 and 60 min post-stimulus. A unique pathway
was observed among gene body binding targets in *SWIZ-OE*; transcript abundance was reduced
following touch. Thus, SWIZ binding to a promoter or gene body was more strongly associated
with increased expression with the exception of gene body targets in touched *SWIZ-OE* plants,
which were more strongly repressed.
278
Among the genes differentially expressed in response to both touch, *SWIZ-OE* transgene, and
bound by SWIZ in the first intron is the mixed glucan synthase *CSLF6* (**Fig 6A**). The transcript
abundance of *GA2ox3*, which inactivates bioactive GA, was bound by SWIZ protein in both the
gene body and 3'UTR and expression significantly increased 30 min after touch in *SWIZ-OE*
(**Fig 6B**). An example of gene body binding repression is Bradi1g30252, the closest ortholog to a
wheat transcription factor induced by hypoxia, *TaMYB1* (Lee et al., 2006). This gene was touch
induced in wildtype and repressed in *SWIZ-OE* with a binding site in the 5'UTR (**Fig 6C**). A
membrane-associated transcription factor, NAC35, exhibited a similar expression to *SWAM3*
(Handakumbura et al., 2018) and SWIZ bound the promoter region (**Fig 6D**).
288
B. distachyon displays classic thigmomorphogenic phenotypes
We next investigated the effect of touch treatment on aboveground tissue. Wildtype plants were
perturbed with a metal bar once every 90 min for either two or three weeks (**Supplemental Fig
3, 4**). After the treatment period, all plants were allowed to recover and grow undisturbed until
senescence (**Supplemental Fig 4**). Two week stressed plants were significantly shorter than
control plants, and three week stressed plants were shorter still (**Supplemental Fig 4A-B**).
Despite this difference in height, there was not a significant difference between the groups in
terms of aboveground plant weight (**Supplemental Fig 4C**). Three week stressed plants had
significantly more branches ($p < 0.05$), with a non-significant increase in two week stressed
plants (**Supplemental Fig 4D**). Transverse stem cross sections in the third elongated internode
and pedncule did not show a significant difference in cell wall thickness or phloroglucinol
staining in response to touch (**Supplemental Fig 4E-G**).
301
Root morphology is altered by *SWIZ-OE* and touch treatment
Given the SWIZ protein mechanotropic response and regulatory influence, we then investigated
how SWIZ might impact plant growth in response to touch. We challenged wildtype and *SWIZ-*
OE roots with the same style of touch treatment used in the translocation and gene expression
experiments, but repeated twice daily for a period of five days. In both touch and control
conditions, *SWIZ-OE* roots were significantly shorter than wildtype, suggesting a dwarfing effect
from *SWIZ* overabundance (**Fig 7**). Mechanical challenges to roots have been reported to impact
root straightness, a trait that has also been described as being impacted in other bZIP studies
(Van Leene et al., 2016; Zha et al., 2016). In response to touch treatment, we observed that

Figure 6. SWIZ binding targets differentially expressed in response to touch and *SWIZ-OE*. Gene expression over time of selected genes with SWIZ binding sites. Line graphs are the average transcript abundance of three biological replicates for each time point. Binding site determined as peaks of sequence alignment. Scale bar unit is bases. Direction of transcription is shown with arrows on the gene model, 5' and 3' UTRs are depicted by narrowed rectangles on the gene model.

311   *SWIZ-OE* roots were significantly less straight than wildtype, while this was not observed in
312   control conditions (**Fig 7**). We further tested the mechano-response of *SWIZ-OE* roots by
313   growing seedlings on plates with increasing degrees of plate angle at 10°, 20°, 30°, and 40° from
314   vertical (**Fig 8**). Greater angles result in more direct contact between the root tip and media and
315   thus increased mechanical stimulation in *A .thaliana (Oliva and Dunand, 2007; Zha et al., 2016)*.

Figure 7. *SWIZ-OE* roots are shorter than wildtype and less straight in response to direct touch. Wildtype and *SWIZ-OE* seedlings grown on plates under control (A) or touched (B) conditions for five days. Touched plants were probed with a pipette tip twice a day. (C) Quantification of root length in control and touched conditions. Significance denoted by * reflecting t-test p-value < 0.05, ns = not significant. (D) Quantification of root straightness in control and touch conditions. Significance denoted by * reflecting Wilcoxon sign-ranked test for non-parametric data with p-value < 0.05, ns = not significant.. Scale bar = 1 cm.

316   Wildtype *B. distachyon* roots displayed decreasing root length with increasing plate angle. *SWIZ-*
317   *OE* roots were shorter than wildtype at all plate angles tested (**Fig 8**). Root straightness did not
318   show any significant differences. Thus, *SWIZ-OE* roots were shorter than wildtype regardless of
319   direct touch stimulus or increasing plate angle, and significantly less straight under direct touch.
320

Figure 8. Wildtype roots shorten with increasing plate angle, while _SWIZ-OE_ roots are consistently short, with no significant difference in straightness. Wildtype and _SWIZ-OE_ seedlings grown on plates at (A) 10°, 20°, 30°, and (B) 40° incline from vertical. (C) Quantification of root length. (D) Quantification of primary root straightness. Significance denoted by * reflecting t-test p-value < 0.05, ns = not significant. Scale bar = 1 cm.

321   **Stem height and cell wall thickening are affected by _SWIZ-OE_ and touch treatment**

To test the role of *SWIZ* in regulating thigmomorphogenesis and secondary cell wall development, we tested wildtype and *SWIZ-OE* under two weeks of mechanical perturbation as described above. In control conditions, there were no differences among genotypes in height, weight, or branching. Touch significantly shortened both widltype and *SWIZ-OE* stems relative to control conditions, but not relative to each other. (**Fig 9A, Supplemental Fig 5**). Thus, *SWIZ-OE* stems are inherently shorter than wildtype.

Transverse sections of the stem were made in the peduncle, the last elongated stem internode where the touch treatment occurred during stem elongation, and stained with phloroglucinol-HCl. *SWIZ-OE* touched peduncles showed significantly thicker interfascicular fiber cell walls (**Fig 9B-C**). Thus, overabundance of *SWIZ* transcript sensitizes wall thickening in response to touch treatment.

SWIZ translocation is sensitive to cellular GA levels, and may act in maintaining GA homeostasis following mechanical stimuli

Mechanical forces have been shown to inactivate cellular GA, and that cellular GA levels dictate cytosolic versus nuclear localizations of *N. benthamiana* Group I bZIPs similar to SWIZ (Fukazawa et al., 2010; Lange and Lange, 2015). To investigate this in relation to SWIZ, *SWIZ:GFP-OE* roots were treated with bioactive GA or paclobutrazol, an inhibitor of GA synthesis. SWIZ:GFP signal amplitude was reduced in roots treated with bioactive GA, particularly at the 50 and 100 mM level. After 60 min the nuclear signal had returned to near pre-stimulus levels, and a second stimulus was applied (**Fig 10A**). This time all GA treated roots showed a similar reduction in nuclear signal, nearly half of the untreated control. These results suggest that the addition of exogenous GA reduces SWIZ translocation to the nucleus following mechanical force application.

Roots treated with 10 mM paclobutrazol did not show altered translocation dynamics compared to the untreated control, but roots treated with 50 or 100 mM paclobutrazol initially showed greater SWIZ:GFP nuclear localization, even prior to mechanical stimulus (**Fig 10B**). SWIZ:GFP signal in the 100 mM paclobutrazol treated roots was high prior to treatment and remained high even after the other nuclear signals diminished. This suggests that pharmacological reduction of bioactive GA levels triggers SWIZ nuclear translocation even without the application of mechanical force.

Given the role of GA in touch responsive bZIP translocation dynamics, we measured how GA related gene expression changed in response touch. In wildtype following touch, GA synthesis and GA activating genes *KS*, *KAO4*, and *GA20ox4* were downregulated, while the genes encoding GA inactivating enzymes GA2ox4 and GA2ox9 were significantly upregulated (**Fig 10C**). In *SWIZ-OE*, *KS* was upregulated prior to touch and 10 min post touch. *GA20ox4*, a gene responsible for synthesizing bioactive GA, was downregulated prior to touch in *SWIZ-OE* plants, then significantly upregulated following touch. At 30 and 60 min *GA2ox4* and *GA2ox9* were upregulated, along with the DELLA protein Rht-1 (**Fig 10C**). Here we show consistent patterns

Figure 9. *SWIZ-OE* enhanced thigmomorphogenic response in stems. Wildtype and *SWIZ-OE* plants were placed under control conditions or received two weeks of mechanical stress every 90 min. (A) Quantification of main stem height at senescence. (B) Quantification of interfascicular fiber wall thickness in transverse cross sections of the peduncle. n = 4 to 6 plants per genotype, per treatment. (C) Representative cross sections stained with phloroglucinol-HCl. Scale bar = 100 µm. Significance denoted by * reflecting t-test p-value < 0.05, ns = not significant.

360   of gene expression associated with the touch inactivation of GA and that SWIZ protein promotes
361   expression of GA synthesis and GA activation genes following mechanical perturbation.

Figure 10. SWIZ translocation is dependent on bioactive GA levels and impacts GA related gene expression. (A) SWIZ translocation to the nucleus is dampened by the addition of exogenous GA4. (B) SWIZ translocation to the nucleus is enhanced by the addition of paclobutrazol (PBZ), a GA synthesis inhibitor. The average nuclear GFP signal is represented by the trend line, with error bars indicating standard deviation of the mean. Imaging was initiated at t = 0 min, while touch treatment was initiated at t = 10 min. n = 85-125 (A) and n = 64-142 (B) nuclei per treatment. Gene expression analysis of GA related genes. (C) Log fold change of gene expression relative to their expression at time 0, pre-touch. Bar color indicates relevant function in GA synthesis and signalling. Significance denoted by * reflecting q < 0.1 compared to wildtype expression at t = 0, with q-values representing Wald test p-values adjusted for false discovery rate.

363
364

DISCUSSION

Touch stimulus is generally an inhibitor of plant elongation growth, but promotes branching and radial expansion (Jaffe, 1973; Braam, 2004; Chehab et al., 2009). The majority of this work has been done in dicots, where increased radial growth has been associated with greater deposition of secondary wall forming cells, particularly in woody species such as poplar (Biro et al., 1980; Coutand et al., 2009; Börnke and Rocksch, 2018; Roignant et al., 2018; Niez et al., 2019). Our understanding of thigmomorphogenesis in grasses is limited, and mostly in the agricultural context of lodging (Shah et al., 2019). While these studies highlight the importance of stem strength and associated cell wall defects, they do relatively little to elucidate the mechanosensitive response. Recent work by Gladala-Kostarz et al (2020) describes grass touch response to both wind and direct mechanical treatment, with an emphasis on cell walls and stem anatomy. Touch treatment significantly increased lignin content, and also altered the relative abundance of wall bound hydroxycinnamates. Wall polysaccharide content, particularly glucose and galactose, also increased. Consistent with increased secondary wall lignification, touched plants had increased wall stiffness and decreased saccharification.

Specific gene expression patterns have become molecular hallmarks of plant touch response, most notably induction of the *TCH* and similar genes. Orthologs of wall modifying XTHs and CAM/CML signaling genes are upregulated in response to mechanical stimulation across species, and as we present here, in *B. distachyon*. Touch elicits major global changes in gene expression, with 2.5% to 10% of the genes assayed in *A. thaliana* and poplar differentially expressed (Lee et al., 2005; Pomiès et al., 2017; Van Moerkercke et al., 2019). In sorghum (*Sorghum bicolor*), leaf sheath imposes mechanical constraints on emerging buds, and removing this dramatically altered gene expression, with 42% of genes differentially expressed over a 9 h period (Liu and Finlayson, 2019). Our observations of short term (<2 h) touch responses within stress related pathways align with observations from poplar and *A. thaliana*, with rapid induction of *TCH*, biotic, and abiotic stress related genes (Pomiès et al., 2017). Canonical touch responsive genes such as the orthologs of *TCH1-4* and related *XTH* and *CML* genes were upregulated immediately following stimulus. However, we also note the previously unreported touch-induction of GH17 family genes, as well as *CSLF6*, suggesting a role for MLG and β-1,3 glucan modification in grass touch response.

In both poplar and in our analysis of *B. distachyon* touch responses, touch-regulated expression of secondary cell wall related transcripts differed between early and late timepoints. Upon touch, we observed immediate repression of many key cell wall biosynthetic enzymes, followed by significant upregulation one hour after stimulus. In poplar, no significant repression of these transcripts was reported, but terms related to wood or cell wall synthesis were also not enriched among touch-regulated transcripts until 24 or 72 hours after touch (Pomiès et al., 2017). While differences in the method used to quantify gene expression, and in the temporal sampling scheme

405 complicate direct comparisons of plant touch response experiments, this may suggest that
406 delayed induction of secondary cell wall transcripts is a common feature of plant touch
407 responses.
408
Thigmomorphogenesis, GA physiology, and secondary cell walls
Our understanding of touch responsive gene expression is further informed by the dynamics of
SWIZ nuclear translocation. The timing of SWIZ nuclear accumulation following mechanical
stimulus was consistent with observations of *A. thaliana* Group I bZIPs (Tsugama et al., 2014;
Tsugama et al., 2016). However, by adopting a finer temporal sampling scheme, and quantifying
nuclear signal sooner after touch (2 vs. 30 min), our results clarify the rapid and immediate
nature of this translocation. Furthermore, we show that the speed and magnitude of SWIZ
nuclear accumulation is not diminished over three successive touch treatments. Notably, poplar
transcriptional responses to a second touch stimulus were significantly attenuated relative to a
single treatment, and in separate experiments, trees subjected to a daily touch treatment over four
days no longer showed touch-induced increases in their rates of radial growth (Martin et al.,
2010; Pomiès et al., 2017). Thus, if *B. distachyon* is likewise desensitized to repeated touch, the
high repeatability of SWIZ translocation might implicate a mechanism downstream of touch
perception and bZIP translocation in mediating that shift. Like the touch-responsive transcription
factor AtVIP1, SWIZ translocation occurred in the specific area of touch treatment, suggesting a
local response to stimulus rather than a tissue or whole organism response. SWIZ also displays a
highly similar pattern of GA responsive translocation to the well-studied bZIP NtRSG (Ishida et
al., 2004; Fukazawa et al., 2010; Fukazawa et al., 2011). Exogenous bioactive GA dampened
SWIZ nuclear translocation, while inhibition of GA synthesis with paclobutrazol induced
translocation, even prior to touch. Furthermore, like NtRSG, SWIZ appears to both respond to
and regulate GA bioactivity. In wildtype, GA inactivating enzymes GA2ox4 and GA2ox9 were
upregulated following touch, consistent with previous reports (Lange and Lange, 2015). In
SWIZ-OE we also observe touch induced expression of *GA20ox4*, a GA activating enzyme, and
as with NtRSG, this response can restore bioactive GA following touch induced degradation
(Ishida et al., 2004; Fukazawa et al., 2010; Fukazawa et al., 2011). Altered GA pathways are a
well described mechanism for reduced stem elongation (Lange and Lange, 2006; Nagai et al.,
2020), and dwarfing has been described for dominant negative mutants of Group I bZIPs *NtRSG*
and *OsRF2A* (Fukazawa et al., 2000; Dai et al., 2003). *SWIZ-OE* stems were significantly shorter
following touch treatment, suggesting that GA misregulation may contribute to this
thigmomorphogenic phenotype. *SWIZ-OE* roots were shorter than wildtype under control, direct
touch, and multiple plate growth angles. This may suggest that *SWIZ-OE* roots grown in agar
media experience a sufficient degree of mechanical input to reduce elongation. It is also
conceivable that misregulation of GA related genes by *SWIZ* overabundance inherently results in
shorter tissues.
443

Another component of this reduced elongation may be increased secondary cell wall deposition, as this could terminate elongation. Prior to touch, almost the entire monolignol biosynthetic pathway, including laccase genes that radicalize monolignols, and BAHD acyltransferases that add hydroxycinnamic acids to wall polymers, are significantly downregulated in *SWIZ-OE* plants. Following touch, most of these genes become significantly upregulated. This activity is consistent with our observations of increase in wall thickness in touched *SWIZ-OE* peduncles, and aligns with the compositional data presented by Gladala-Kostarz et al. (2020). This activation may be a result of direct binding of SWIZ to modulate expression, or an indirect effect from another transcriptional regulator such as the NAC transcription factor *SWN5* that is significantly upregulated in *SWIZ-OE* following touch and capable of activating the full developmental program for secondary cell wall synthesis (Valdivia et al., 2013). GA has also been implicated in regulating cell wall thickening, and the interactions between SWIZ and touch mediation of GA levels may contribute to this aspect of thigmomorphogenesis (Huang et al., 2015; Falcioni et al., 2018; Wang et al., 2021).

While the E-box (CANNTG) was initially described as a bHLH binding motif, many bZIP groups in *A. thaliana* have been shown to bind similar sequences. The two closets *A. thaliana* orthologs of SWIZ , bZIP18 and 52, both bind (A/C)CAG(G/C)(T/C), while VIP1 binds (A/C)CAGCT(G/A) (O'Malley et al., 2016). Using the same technique, we identified the SWIZ binding motifs as (A/C)CAGNCTG(T/G) and (A/C)CAGCTG(T/G). Both of these sequences are similar to those bound by *A. thaliana* orthologs, although the presence of an ambiguous nucleotide in the core of one variant is not previously reported for other Group I bZIPs. Among genes bound by SWIZ *in vitro,* we observed both activation and repression following touch, suggesting a complex regulatory function for SWIZ. Furthermore, SWIZ direct binding sites were found in both promoter regions and in the body of genes and most often in genes activated by touch. SWIZ gene body binding targets tended to be repressed in *SWIZ-OE* plants following touch, without either being clearly associated with up or down regulation. AtVIP1 and AtbZIP29 have both been described as activators, particularly of genes related to biotic and abiotic stress response, cell cycle control, and development. (Yin et al., 1997; Ringli and Keller, 1998; Pitzschke et al., 2009; Van Leene et al., 2016). AtbZIP18 is suggested to have repressive roles (Tsugama et al., 2012; Gibalová et al., 2017).

bZIPs are known to homo- and heterodimerize through their leucine zipper domains (Schütze et al., 2008), and in *A. thaliana* the combinatorial interactions of different bZIP groups have been fairly well described (Deppmann et al., 2004; Ehlert et al., 2006; Grigoryan and Keating, 2006; Weltmeier et al., 2006; Schütze et al., 2008). These interactions can have a synergistic effect on transcriptional activity, and can result in unique binding interactions (Schütze et al., 2008; Van Leene et al., 2016). It is conceivable that this heterodimerization may explain some aspects of our results, such as why so many of the cell wall genes that are upregulated in *SWIZ-OE* after touch are significantly downregulated prior to touch. One hypothesis is that SWIZ, as with many other bZIPs, can regulate targets as a heterodimer. The abundance of SWIZ protein in the

cytoplasm in *SWIZ-OE* plants prior to touch may act as a sink, sequestering interacting partners
that would otherwise be regulating targets independent of a touch stimulus. Further
experimentation is needed to elucidate this bZIP regulatory network in response to touch and
other stimuli.

CONCLUSIONS

Proteins orthologous to SWIZ in other species have been implicated in cell wall development
and remodeling, but not cell wall thickening. SWIZ may influence root length inherently, and
stem height and cell wall thickening in response to mechanical stimuli, which regulates SWIZ
translocation into the nucleus by acting on the cellular bioactive GA pool. Touch has been shown
to decrease cellular bioactive GA levels in *A. thaliana*, and Group I bZIPs are known to
translocate to the nucleus in response to touch and to low levels of bioactive GA. Direct
mechanical stimulation and chemical inhibition of GA biosynthesis both caused SWIZ to
translocate from the cytoplasm into the nucleus. This translocation can be reduced with
exogenous bioactive GA. GA biosynthesis and signaling genes were upregulated in *SWIZ-OE*
lines following touch, indicating that SWIZ may act in managing GA homeostasis following
mechanical perturbation. Altering *SWIZ* function may impact GA homeostasis and result in
reduced elongation. Touch significantly remodeled the *B. distachyon* transcriptome, with notable
changes in wall polysaccharide biosynthetic gene expression not previously reported. Enhanced
SWIZ function through overexpression amplified the expression of cell wall related genes and
synthesis of interfascicular fiber walls in touched tissues. Greater secondary wall synthesis in
response to touch may also explain reduced stem elongation in *SWIZ-OE*. Together, the evidence
presented here connects mechanotropic bZIP dynamics with changes in cellular GA levels that
have been observed separately in other systems, and shows a link between touch response and
secondary cell wall transcriptional regulation (**Fig 11**).

MATERIALS AND METHODS

Phylogenetic analysis
Protein sequences described for *A. thaliana*, *B. distachyon*, and *O. sativa* (Liu and Chu, 2015) as
Group II bZIPs were selected and searched against the most recent genome annotations in
Phytozome v12.1 (https://phytozome.jgi.doe.gov). The *Nicotiana tabacum* homologs NtRSGa
and NtRSGb were also added to the analysis. Protein sequences were aligned using the MAFFT
service for multiple sequence alignment with the iterative refinement method L-INS-I (Katoh et
al., 2019). The alignment was used to construct the phylogenetic tree using a maximum
likelihood analysis with a bootstrap resampling value of 1000 in W-IQ-TREE (Trifinopoulos et
al., 2016). All of the proteins included in the phylogenetic analysis are described in
Supplemental File 1.
520

Figure 11. SWIZ regulates cell wall synthesis and GA homeostasis in response to mechanical stimulus. Prior to touch, SWIZ is localized to the cytosol and partially in the nucleus. The bioactive GA pool is depleted in response to touch, triggering SWIZ to translocate to the nucleus where it positively regulates cell wall synthesis and GA activating enzymes.

Plant transformation

Overexpression and artificial microRNA transgenes and transgenic plants were constructed as previously described (Handakumbura et al., 2013). The full length coding sequence of Bradi1g17700 was amplified from cDNA and cloned into the pOL001 ubigate ori1 binary expression vector (Handakumbura et al., 2018) to make the *SWIZ-OE* trangene. The coding sequence lacking the stop codon was amplified and fused in frame with the coding sequence of *Aequorea victoria* enhanced GFP in the pOL001 vector to generate *SWIZ:GFP-OE*.

Translocation assay

Seeds were surface sterilized and grown vertically on 1X MS media, pH 5.7, without sucrose for 6d at 28°C in the dark. After 6 d, seedlings were moved to treatment plates containing 1X MS media, pH 5.7, plus varying concentrations (0, 10, 50, or 100 mM) of GA4, paclobutrazol. Roots were positioned such that they were completely submerged in media and relatively flat along the imaging plane After a 6 h incubation, plants were stimulated and imaged.

All observations were made on a Nikon A1R scanning confocal microscope using a Plan Apo 10x 0.5NA objective. Root areas to be observed were located by eye using transmitted light and then confirmed under confocal conditions using a 488 nm laser and green filtered PMT detector. The X, Y, and Z coordinate stage locations of each region were programmed into the Nikon NIS

Elements 5.2 Advanced Research V6 software package for automated imaging. After all target regions were programmed, 30 min of imaging began pre-treatment, with images captured every 2 min.

To elicit the touch response, the observed root region was gently probed for 5 sec with a blunt metal probe while observing through the eyepiece. Images were captured for 60-90 min post treatment. For experiments with multiple stimulus events, the timelapse sequence was paused and roots were probed as described for the relevant stimulus events.

Analysis of GFP signal was done using the Nikon NIS Elements Advanced Research V5 software package. For each time series, the frame showing maximal nuclear signal was selected and used as a reference point. Using the General Analysis tool, a channel for GFP signal was established and thresholded for intensity and particle size to identify the nuclear regions. These regions were added to the timelapse image series as a binary layer and then converted to static regions of interest for quantification. The GFP intensity under each nuclear region of interest was calculated for the course of the timelapse and the average signal from each nucleus was plotted for intensity over time.

Thigmomatic construction and operation

The Thigmomatic is a basic robotic device that sweeps plants with a metal bar at regular intervals to elicit a touch response. The device was constructed from aluminum V-Slot linear rail (Openbuilders Partstore, Monreoville, NJ) and bracket joints for the upright supports (20x20 mm), cross bars (20x20 mm), and tracks (20x40 mm). Two gantry carts ride along the 20x40 mm V-Slot linear rails, connected by a 6.35 mm diameter metal rod bolted to the carts. Their movement is powered by a belt driven linear actuator system using a NEMA 17 stepper motor with a 12V 18W AC/DC power supply. The stepper motor provides fine spatial control over the gantry cart position with bi-directional motion. Motor function is controlled by a Raspberry Pi 3B microcomputer equipped with a stepper motor HAT (Adafruit Industries, New York). The Thigmomatic was programmed to cover a specified distance in one direction once every 90 min. Accession Bd21-3 was used for these experiments. Seeds were stratified on wet paper towel wrapped in foil to exclude light for 10 days at 4°C before being planted in Promix BX potting mix in SC10 Ray Leach Cone-tainers (Stuewe & Sons Inc). Plants were grown in a Percival PGC-15 growth chamber with day/night conditions of 20 h light at 24°C and 4 h dark at 18°C, respectively.

Transverse stem sections and histology

The main stem of senesced plants was taken and the internode of interest removed and embedded in 8% agarose. A Leica VT1000 Vibratome was used to make 55 μm thick transverse sections. Histochemical staining was carried out using phloroglucinol-HCl as previously described (Matos et al., 2013). Images were obtained at 4, 10, and 20X using a Nikon Eclipse E200MV R light microscope and PixeLINK 3 MP camera. Transverse sections imaged at 20x were used for cell

578 wall thickness measurements. Interfascicular fiber cells separated by one cell layer from the
579 mestome cells on the phloem side of major vascular bundles were targeted for measurement.
580 Using ImageJ, lines were drawn across two walls of adjoining cells. The resulting line length
581 was divided by two to give one cell wall width. Approximately 15 measurements were made for
582 each plant.

RNA extraction and quantification and analysis
Seedlings were grown for 6 d on vertical 1X MS agar plates. Touch treatment was performed as
described above using a metal probe along the entire length of the root. Untouched samples were
collected immediately before touch treatment began, and touched samples were collected after
the designated time. Three roots were pooled per biological replicate, and RNA was extracted
using the Qiagen RNeasy Plant Mini Kit with on-column DNA digestion with RNase-free DNase
I (Qiagen). Strand-specific libraries were prepared using the Illumina TruSeq kit. Libraries were
sequenced using Illumina technology. Sequences were processed as previously described
(MacKinnon et al., 2020) and then checked for differential expression using the likelihood ratio
test. Specific contrasts (i.e., wildtype vs *SWIZ-OE*) were identified and compared using the Wald
test. Statistical enrichment of gene families was assessed using Fisher's exact test. Raw read data
were deposited in the European Nucleotide Archive for public access (Accession no.: E-MTAB-
10084).

DNA affinity purification sequencing
DNA affinity purification was carried out as previously described (Handakumbura et al., 2018).
In brief, transcription factor coding sequences were HALO tagged and mixed with Bd21
genomic DNA for *in vitro* binding. Protein-DNA was crosslinked, fragmented,
immunoprecipitated using the HALO antibody, barcoded, and sequenced. Reads were mapped
back to the Bd21 genome using HiSat2 (Kim et al., 2015) to identify binding target loci. Peak
calling and motif analysis was done using HOMER v4.10 (Hypergeometric Optimization of
Motif EnRichment) suite (Heinz et al., 2010). Motif enrichment was calculated against the
hypergeometric distribution; the significance threshold was set to $p < 0.05$. The nearest annotated
gene to a bound peak was used for GO analysis. Raw read data were deposited in the European
Nucleotide Archive for public access (Accession no.: E-MTAB-10066).

Gene Ontology analysis
609 Phytozome was used to find orthologs for all *B. distachyon* v3.1 genes as the reciprocal best
610 match to *A. thaliana* TAIRv10 protein sequences. Arabidopsis gene identifiers were submitted to
611 g:Profiler (Raudvere et al., 2019) for KEGG and Wiki pathway enrichment analysis.

iDREM network analysis
Probabilistic graphical models that predict diverging gene expression paths and the points at
which regulators may influence those paths were generated using iDREM (Ding et al., 2018).
Briefly, this software applies an input-output hidden Markov model to time course gene

expression data overlaid with static regulatory information, in this case SWIZ protein-DNA
interactions identified from DAP-seq. GO analysis, described above, was also applied to the gene
sets in each path identified by iDREM.
619

Root Touch Experiment
Seeds were surface sterilized and plated on 1x MS, pH 5.7, containing 0.05 % MES as a
buffering agent and 1% plant agar (Gold Bio). Seeds were stratified on plates in the dark at 4 ℃
for 2 days and then transferred to a Percival PGC-15 growth chamber with day/night conditions
of 16h light at 24 ℃ and 8h dark at 18 ℃, respectively and grown at a ~10° angle from vertical.
After 2 days of preliminary growth, a touch location was designated by selecting the lower of 1
cm up from the root tip or 1 cm down from the seed and marked on the plate. Each day for 5
days, this marked spot was treated twice a day, two hours before and two hours after chamber
midday (ZT6 and ZT10). Touch treatment consisted of 5 firm presses with the side of a sterile
pipette tip, just hard enough to cause elastic deformation of the media underlying the root when
viewed through a dissecting microscope. To ensure unbiased application of touch treatment and
control for any blocking effects, each plate contained one group of four *SWIZ-OE* seeds and one
group of four Bd21-3 wildtype seeds, with the experimenter blinded to their respective positions
for the duration of the experiment.
634

Root length and straightness measurement
Plates were photographed at the conclusion of the experiment. To generate high contrast images
suitable for root tracing, plate images were separated into hue, saturation and brightness
channels, and the saturation channel was selected and inverted. Semi-automated measurements
of root length were performed using the Smart Roots plugin for ImageJ (Lobet et al., 2011).
Straightness was quantified as described in (Swanson et al., 2015); the straight line distance
between root tip and the base of the seed was measured, and this value was divided by the traced
length of the root.

SUPPLEMENTAL MATERIALS

Supplemental Table S1. Cluster 56 from network analysis of gene expression atlas (Sibout et al. 2017) includes Bradi1g17700.

Supplemental Table S2. Differentially expressed genes in wildtype Bd21-3 root tissue 10 min after touch treatment.

Supplemental Table S3. Differentially expressed genes in wildtype Bd21-3 root tissue 30 min after touch treatment.

Supplemental Table S4. Differentially expressed genes in wildtype Bd21-3 root tissue 60 min after touch treatment.

Supplemental Table S5. Calmodulin and calmodulin-like touch responsive gene expression in wildtype Bd21-3 root tissue.

Supplemental Table S6. XTH and XTH-like touch responsive gene expression in wildtype Bd21-3 root tissue.

Supplemental Table S7. Glycoside hydrolase touch responsive gene expression in wildtype Bd21-3 root tissue.

Supplemental Table S8. GO analysis of genes in iDREM identified pathways derived from differential gene expression following touch in wildtype, swiz-amiRNA, and SWIZ-OE root tissue.

Supplemental Figure 1. Description of *SWIZ* transgenic reagents. (A) Diagram of *SWIZ* overexpression transgenes. The *ZmUbi* promoter was used to drive expression of the *SWIZ* coding sequence, either alone or fused in frame with *eGFP*. (B) Relative level of *SWIZ* transcript measured by RT-qPCR in whole stem tissue was collected 1 d after inflorescence emergence in *SWIZ:GFP-OE*. LB, left border; *ZmUbi* prom, maize ubiquitin promoter; Hyg, hygromycin phosphotransferase gene; NOS, nopaline synthase terminator; RB, right border. ns: $p > 0.05$, : *: $p < 0.05$.

Supplemental Figure 2. Network analysis of differential gene expression paths and enriched GO terms. (A) Path determinations from iDREM time course analysis. Each line represents a set of genes with similar expression level patterns over the time course relative to time 0, pre-touch. (B) Heatmap of selected GO term enrichment of genes from iDREM path analysis. Color indicates qvalue score, with a cutoff of 0.1. Terms without a colored block are not significantly enriched in that path.

Supplemental Figure 3. Thigmomatic, an automated system to administer touch stimulus.
(A) Overview of Thigmomatic inside a Percival PGC-15 growth chamber showing linear rail based frame (1), gantry carts (2), NEMA17 stepper motor (3), 12V 18W AC/DC power supply (4), and Raspberry Pi 3b microcomputer (5). (B) Thigmomatic making contact with a *Brachypodium distachyon* plant.

Supplemental Figure 4. *Brachypodium distachyon* displays classic thigmomorphogenic phenotypes. One-week-old wildtype plants were treated with touch stimulus every 90 min. (A) Left to right, plants that experienced no stress, two weeks stress, three weeks stress and were then allowed to recover, imaged one week after the end of treatment (B) height, (C) aboveground non-grain biomass weight, and (D) branch number were measured at senescence. ns: $p > 0.05$, *: $p < 0.05$, n = 5 plants per treatment. (E) Transverse sections of the peduncle or third internode were take from control, 2 week stressed, and 3 week stressed plants and stained with phloroglucinol-HCl. (F) Quantification of interfascicular fiber wall thickness. Scale bar = 100 μm. n = 3 plants per treatment.

Supplemental Figure 5. Biomass and branching in SWIZ-OE under touch treatment shows no difference from wildtype. Wildtype and *SWIZ-OE* plants were placed under control conditions or received two weeks of mechanical stress every 90 min. After senescence, aboveground biomass weight (A) and branch number (B) were quantified. p value calculated by student t-test.

Supplemental File 1. Gene names and amino acid sequences used in the SWIZ bZIP Group I phylogenetic analysis.

ACCESSION NUMBERS

AtbZIP18 (At2g40620), AtbZIP29 (At4g38900), AtbZIP52 (At1g06850), AtTCH1/AtCaM2 (At2g41110), AtTCH2/AtCML24 (At5g37770), AtTCH3/CML12 (At2g41100), AtTCH4/AtXTH22 (At5g57560), CAD1 (Bradi3g17920), CESA4 (Bradi4g28350), COMT6 (Bradi3g16530), CSLF6 (Bradi3g16307), GA2ox3 (Bradi2g50280), MYB44 (Bradi1g30252), NAC35 (Bradi2g28020), NtRSGa (Niben101Scf01150g00005), NtRSGb (Niben101Scf02191g00014), SWIZ (Bradi1g17700), VIP1 (At1g43700).

ACKNOWLEDGEMENTS

This work was supported by the Office of Science, Biological and Environmental Research, Department of Energy (DE-FG02-08ER64700DE and DE-SC0006641) and The National Science Foundation Division of Integrative Organismal Systems (NSF IOS-1558072) to S.P.H.. The work conducted by the US DOE Joint Genome Institute is supported by the Office of Science of the US Department of Energy under Contract no. DE-AC02-05CH11231. The microscopy data was gathered in the Light Microscopy Facility and Nikon Center of Excellence

714   at the Institute for Applied Life Sciences, UMass Amherst with support from the Massachusetts
715   Life Sciences Center.

717 **Table 1.** The count of differentially expressed genes relative to untouched wildtype plants that
718 were SWIZ DAP-seq targets in either the promoter or gene body regions.

Time after touch	Promoter		Gene body	
	Up	Down	Up	Down
(min)	percent (count)			
0	42.0 (103)	58.0 (142)	48.0 (286)	52.0 (310)
10	57.0 (73)	43.0 (55)	53.0 (167)	47.0 (148)
30	59.4 (79)	40.6 (54)	41.2 (136)	58.8 (194)
60	64.2 (77)	35.8 (43)	45.7 (128)	54.3 (152)

719
FIGURE LEGENDS

Figure 1. SWIZ is a Group I bZIP highly expressed in maturing stem and root. (A) *SWIZ*
transcript abundance in *Brachypodium distachyon* leaf, root, and stem tissue measured by
microarray. Mean +/- standard deviation of three biological replicates. (B) Phylogeny analysis of
amino acid sequences from *B. distachyon* (green), *Oryza sativa* (blue), *Arabidopsis thaliana*
(red), *Nicotiana tabacum* (orange) shows *SWIZ* (black dot). The phylogeny was reconstructed
using maximum likelihood analysis with a bootstrap resampling value of 1000. The numbers
labeled on the branches are the posterior probability to support each clade. The protein
sequences, full locus IDs, and plant species with genome source are provided in Supplemental
File 1.

Figure 2. SWIZ translocates to the nucleus in response to mechanical stimulus, specifically
in regions directly stimulated. (A) Image of *SWIZ:GFP-OE* and *GFP-OE* roots prior to
stimulus and 30 min post stimulus. Roots were observed immediately following mechanical
perturbation. (B) Quantification of nuclear signal. n=14-20 nuclei. (C) SWIZ translocation
occured in the local area of the stimulus. At 30 min, stimulus was applied to an upper region of
the root, while at 120 min it was applied to a lower region approximately 3 cm below. n=109,
184 nuclei respectively for upper and lower regions. (D) *SWIZ:GFP-OE* roots were imaged by
confocal microscopy with stimulus applied in the field of view at 0, 90, and 180 min. n=126
nuclei. (B-D) Images were taken every 2 min. Nuclear GFP signal was quantified in selected
nuclei at each time point. The average nuclear GFP signal is represented by the line with error
bars indicating standard error of the mean.

Figure 3. Transcriptome analysis of touch response in *Brachypodium distachyon* roots. (A)
Root tissue was sampled just prior to touch (t = 0), and at 10, 30, and 60 min following touch
treatment in wildtype and *SWIZ-OE*. (B) Principal component analysis of gene expression across
samples shows the greatest difference corresponding to genotype and the second greatest
corresponding to time after touch. (C) Canonical, as well as novel touch responsive genes are
upregulated in *B. distachyon* following touch. Closest orthologs of the *Arabidopsis thaliana*
TOUCH genes encoding calmodulin-like (CML) and xyloglucan endo-
transglycosylase/hydrolases (XTH) are upregulated following touch, as are previously
unreported members of the glycosyl hydrolase 17 (GH17) family. Full list of gene expression for
TOUCH and glycoside hydrolases defined by Tyler et al. (2010) in Supplemental Tables 5, 6,
and 7. Significance denoted by * reflecting q < 0.1 compared to expression at t = 0, with q-values
representing Wald test p-values adjusted for false discovery rate.

Figure 4. Gene expression analysis of cell wall related genes. Log fold-change of gene
expression in wildtype, *SWIZ-OE*, and *swiz-amiRNA* relative to wildtype expression at time 0,
pre-touch. Bar color indicates the class of cell wall gene. Error bars indicate standard deviation
of three biological replicates. Significance denoted by * reflecting q < 0.1 compared to wildtype
expression at t = 0, with q-values representing Wald test p-values adjusted for false discovery
rate.

Figure 5. DNA affinity purification sequencing to determine SWIZ binding sites. (A) Top
two most statistically enriched sequence motifs in SWIZ binding sites. (B) Distribution of
binding sites across genomic features, relative to primary transcripts of the *Brachypodium*
distachyon annotation v 3.1. (C) Relative distribution of binding sites centered on the
transcriptional start site (TSS, blue dashed line), transcriptional termination site (TTS, red dashed
line) represents the average length of all annotated transcripts, approximately 4.5 kb away from
the TSS. (D) Path determinations from iDREM time course analysis of differentially expressed
genes that also have DAP-seq binding sites. Each line represents a set of genes with similar
expression level patterns over the time course relative to time 0, pre-touch.

**Figure 6. SWIZ binding targets are differentially expressed in response to touch and
SWIZ.** Gene expression over time of selected genes with SWIZ binding sites. Line graphs are
the average transcript abundance of three biological replicates for each time point. Binding site
determined as peaks of sequence alignment. Scale bar unit is bases. Direction of transcription is
shown with arrows on the gene model, 5' and 3' UTRs are depicted by narrowed rectangles on
the gene model.

**Figure 7. *SWIZ-OE* roots are shorter than wildtype and less straight in response to direct
touch.** Wildtype and *SWIZ-OE* seedlings grown on plates under control (A) or touched (B)

conditions for five days. Touched plants were probed with a pipette tip twice a day. (C)
Quantification of root length in control and touched conditions. Significance denoted by *
reflecting t-test p-value < 0.05, ns = not significant. (D) Quantification of root straightness in
control and touch conditions. Significance denoted by * reflecting Wilcoxon sign-ranked test for
non-parametric data with p-value < 0.05, ns = not significant.. Scale bar = 1 cm.

Figure 8. Wildtype roots shorten with increasing plate angle, while *SWIZ-OE* roots are
consistently short, with no significant difference in straightness. Wildtype and *SWIZ-OE*
seedlings grown on plates at (A) 10°, 20°, 30°, and (B) 40° incline from vertical. (C)
Quantification of root length. (D) Quantification of primary root straightness. Significance
denoted by * reflecting t-test p-value < 0.05, ns = not significant.Scale bar = 1 cm.

Figure 9. *SWIZ-OE* enhanced thigmomorphogenic response in stems. Wildtype and *SWIZ-*
OE plants were placed under control conditions or received two weeks of mechanical stress
every 90 min. (A) Quantification of main stem height at senescence. (B) Quantification of
interfascicular fiber wall thickness in transverse cross sections of the peduncle. n = 4 to 6 plants
per genotype, per treatment. (C) Representative cross sections stained with phloroglucinol-HCl.
Scale bar = 100 μm. Significance denoted by * reflecting t-test p-value < 0.05, ns = not
significant.

Figure 10. SWIZ translocation is dependent on bioactive GA levels and impacts GA related
gene expression. (A) SWIZ translocation to the nucleus is dampened by the addition of
exogenous GA4. (B) SWIZ translocation to the nucleus is enhanced by the addition of
paclobutrazol (PBZ), a GA synthesis inhibitor. The average nuclear GFP signal is represented by
the trend line, with error bars indicating standard deviation of the mean. n= 85-125 (A) and n=
64-142 (B) nuclei per treatment. Gene expression analysis of GA related genes. (C) Log fold
change of gene expression in wildtype samples, relative to their expression at time 0, pre touch.
(D) Log fold change of gene expression in SWIZ OE and swiz amiRNA samples, relative to
their expression in wildtype at the same time points. Bar color indicates relevant function in GA
synthesis and signaling. Significance denoted by * reflecting q < 0.1 compared to wildtype
expression at t = 0, with q-values representing Wald test p-values adjusted for false discovery
rate.

Figure 11. SWIZ regulates cell wall synthesis and GA homeostasis in response to
mechanical stimulus. Prior to touch, SWIZ is localized to the cytosol and partially in the
nucleus. The bioactive GA pool is depleted in response to touch, triggering SWIZ to translocate
to the nucleus where it positively regulates cell wall synthesis and GA activating enzymes.

Parsed Citations

Altartouri B, Bidhendi AJ, Tani T, Suzuki J, Conrad C, Chebli Y, Liu N, Karunakaran C, Scarcelli G, Geitmann A (2019) Pectin chemistry and cellulose crystallinity govern pavement cell morphogenesis in a multi-step mechanism. Plant Physiol 181: 127–141
  Google Scholar: Author Only Title Only Author and Title

Bidhendi AJ, Altartouri B, Gosselin FP, Geitmann A (2019) Mechanical stress initiates and sustains the morphogenesis of wavy leaf epidermal cells. Cell Rep 28: 1237–1250.e6
  Google Scholar: Author Only Title Only Author and Title

Bidhendi AJ, Geitmann A (2018) Finite element modeling of shape changes in plant cells. Plant Physiol 176: 41–56
  Google Scholar: Author Only Title Only Author and Title

Biro RL, Hunt ER, Erner Y, Jaffe MJ (1980) Thigmomorphogenesis: changes in cell division and elongation in the internodes of mechanically-perturbed or ethrel-treated bean plants. Ann Bot 45: 655–664
  Google Scholar: Author Only Title Only Author and Title

Börnke F, Rocksch T (2018) Thigmomorphogenesis – Control of plant growth by mechanical stimulation. Sci Hortic 234: 344–353
  Google Scholar: Author Only Title Only Author and Title

Braam J (2004) In touch: plant responses to mechanical stimuli. New Phytol 165: 373–389
  Google Scholar: Author Only Title Only Author and Title

Braam J, Davis RW (1990) Rain-, wind-, and touch-induced expression of calmodulin and calmodulin-related genes in Arabidopsis. Cell 60: 357–364
  Google Scholar: Author Only Title Only Author and Title

Chehab EW, Eich E, Braam J (2009) Thigmomorphogenesis: a complex plant response to mechano-stimulation. J Exp Bot 60: 43–56
  Google Scholar: Author Only Title Only Author and Title

Coomey JH, Sibout R, Hazen SP (2020) Grass secondary cell walls, Brachypodium distachyon as a model for discovery. New Phytol 227: 1709–1724
  Google Scholar: Author Only Title Only Author and Title

Coutand C, Martin L, Leblanc-Fournier N, Decourteix M, Julien J-L, Moulia B (2009) Strain mechanosensing quantitatively controls diameter growth and PtaZFP2 gene expression in poplar. Plant Physiol 151: 223–232
  Google Scholar: Author Only Title Only Author and Title

Dai S, Petruccelli S, Ordiz MI, Zhang Z, Chen S, Beachy RN (2003) Functional analysis of RF2a, a rice transcription factor. J Biol Chem 278: 36396–36402
  Google Scholar: Author Only Title Only Author and Title

Deppmann CD, Acharya A, Rishi V, Wobbes B, Smeekens S, Taparowsky EJ, Vinson C (2004) Dimerization specificity of all 67 B-ZIP motifs in Arabidopsis thaliana: a comparison to Homo sapiens B-ZIP motifs. Nucleic Acids Res 32: 3435–3445
  Google Scholar: Author Only Title Only Author and Title

Ding J, Hagood JS, Ambalavanan N, Kaminski N, Bar-Joseph Z (2018) iDREM: Interactive visualization of dynamic regulatory networks. PLoS Comput Biol 14: e1006019
  Google Scholar: Author Only Title Only Author and Title

Dröge-Laser W, Snoek BL, Snel B, Weiste C (2018) The Arabidopsis bZIP transcription factor family-an update. Curr Opin Plant Biol 45: 36–49
  Google Scholar: Author Only Title Only Author and Title

Ehlert A, Weltmeier F, Wang X, Mayer CS, Smeekens S, Vicente-Carbajosa J, Dröge-Laser W (2006) Two-hybrid protein-protein interaction analysis in Arabidopsis protoplasts: establishment of a heterodimerization map of group C and group S bZIP transcription factors. Plant J 46: 890–900
  Google Scholar: Author Only Title Only Author and Title

Falcioni R, Moriwaki T, de Oliveira DM, Andreotti GC, de Souza LA, Dos Santos WD, Bonato CM, Antunes WC (2018) Increased gibberellins and light levels promotes cell wall thickness and enhance lignin deposition in xylem fibers. Front Plant Sci 9: 1391
  Google Scholar: Author Only Title Only Author and Title

Fukazawa J, Nakata M, Ito T, Matsushita A, Yamaguchi S, Takahashi Y (2011) bZIP transcription factor RSG controls the feedback regulation of NtGA20ox1 via intracellular localization and epigenetic mechanism. Plant Signal Behav 6: 26–28
  Google Scholar: Author Only Title Only Author and Title

Fukazawa J, Nakata M, Ito T, Yamaguchi S, Takahashi Y (2010) The transcription factor RSG regulates negative feedback of NtGA20ox1 encoding GA 20-oxidase. Plant J 62: 1035–1045

Google Scholar: Author Only Title Only Author and Title

Fukazawa J, Sakai T, Ishida S, Yamaguchi I, Kamiya Y, Takahashi Y (2000) Repression of shoot growth, a bZIP transcriptional activator, regulates cell elongation by controlling the level of gibberellins. Plant Cell 12: 901–915
  Google Scholar: Author Only Title Only Author and Title

Gibalová A, Steinbachová L, Hafidh S, Bláhová V, Gadiou Z, Michailidis C, Müller K, Pleskot R, Dupľáková N, Honys D (2017) Characterization of pollen-expressed bZIP protein interactions and the role of ATbZIP18 in the male gametophyte. Plant Reprod 30: 1–17
  Google Scholar: Author Only Title Only Author and Title

Gladala-Kostarz A, Doonan JH, Bosch M (2020) Mechanical stimulation in Brachypodium distachyon: implications for fitness, productivity and cell wall properties. Plant Cell Environ 43: 1314–1330
  Google Scholar: Author Only Title Only Author and Title

Grigoryan G, Keating AE (2006) Structure-based prediction of bZIP partnering specificity. J Mol Biol 355: 1125–1142
  Google Scholar: Author Only Title Only Author and Title

Hamant O, Heisler MG, Jönsson H, Krupinski P, Uyttewaal M, Bokov P, Corson F, Sahlin P, Boudaoud A, Meyerowitz EM, et al (2008) Developmental patterning by mechanical signals in Arabidopsis. Science 322: 1650–1655
  Google Scholar: Author Only Title Only Author and Title

Handakumbura P, Matos D, Osmont K, Harrington M, Heo K, Kafle K, Kim S, Baskin T, Hazen S (2013) Perturbation of Brachypodium distachyon CELLULOSE SYNTHASE A4 or 7 results in abnormal cell walls. BMC Plant Biol 13: 131
  Google Scholar: Author Only Title Only Author and Title

Handakumbura PP, Brow K, Whitney IP, Zhao K, Sanguinet KA, Lee SJ, Olins J, Romero-Gamboa SP, Harrington MJ, Bascom CJ, et al (2018) SECONDARY WALL ASSOCIATED MYB1 is a positive regulator of secondary cell wall thickening in Brachypodium distachyon and is not found in the Brassicaceae. Plant J 96: 485–699
  Google Scholar: Author Only Title Only Author and Title

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK (2010) Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. Mol Cell 38: 576–589
  Google Scholar: Author Only Title Only Author and Title

Huang D, Wang S, Zhang B, Shang-Guan K, Shi Y, Zhang D, Liu X, Wu K, Xu Z, Fu X, et al (2015) A gibberellin-mediated DELLA-NAC signaling cascade regulates cellulose synthesis in rice. Plant Cell 27: 1681–1696
  Google Scholar: Author Only Title Only Author and Title

Igarashi D, Ishida S, Fukazawa J, Takahashi Y (2001) 14-3-3 proteins regulate intracellular localization of the bZIP transcriptional activator RSG. Plant Cell 13: 2483–2497
  Google Scholar: Author Only Title Only Author and Title

Iida H (2014) Mugifumi, a beneficial farm work of adding mechanical stress by treading to wheat and barley seedlings. Front Plant Sci 5: 453
  Google Scholar: Author Only Title Only Author and Title

Ishida S, Fukazawa J, Yuasa T, Takahashi Y (2004) Involvement of 14-3-3 signaling protein binding in the functional regulation of the transcriptional activator REPRESSION OF SHOOT GROWTH by gibberellins. Plant Cell 16: 2641–2651
  Google Scholar: Author Only Title Only Author and Title

Ishida S, Yuasa T, Nakata M, Takahashi Y (2008) A tobacco calcium-dependent protein kinase, CDPK1, regulates the transcription factor REPRESSION OF SHOOT GROWTH in response to gibberellins. Plant Cell 20: 3273–3288
  Google Scholar: Author Only Title Only Author and Title

Ito T, Ishida S, Oe S, Fukazawa J, Takahashi Y (2017) Autophosphorylation affects substrate-binding affinity of tobacco Ca2+-dependent protein kinase1. Plant Physiol 174: 2457–2468
  Google Scholar: Author Only Title Only Author and Title

Ito T, Nakata M, Fukazawa J, Ishida S, Takahashi Y (2014) Scaffold function of Ca2+-dependent protein kinase: tobacco Ca2+-DEPENDENT PROTEIN KINASE1 transfers 14-3-3 to the substrate REPRESSION OF SHOOT GROWTH after phosphorylation. Plant Physiol 165: 1737–1750
  Google Scholar: Author Only Title Only Author and Title

Jaffe MJ (1973) Thigmomorphogenesis: The response of plant growth and development to mechanical stimulation : With special reference to Bryonia dioica. Planta 114: 143–157
  Google Scholar: Author Only Title Only Author and Title

Jaffe MJ, Biro R, Bridle K (1980) Thigmomorphogenesis: Calibration of the parameters of the sensory function in beans. Physiol Plant 49: 410–416

Google Scholar: Author Only Title Only Author and Title

Jakoby M, Weisshaar B, Dröge-Laser W, Vicente-Carbajosa J, Tiedemann J, Kroj T, Parcy F, bZIP Research Group (2002) bZIP transcription factors in Arabidopsis. Trends Plant Sci 7: 106–111
  Google Scholar: Author Only Title Only Author and Title

Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. Brief Bioinform 20: 1160–1166
  Google Scholar: Author Only Title Only Author and Title

Kim D, Langmead B, Salzberg SL (2015) HISAT: a fast spliced aligner with low memory requirements. Nat Methods 12: 357
  Google Scholar: Author Only Title Only Author and Title

Lange MJ, Lange T (2006) Gibberellin biosynthesis and the regulation of plant development. Plant Biol 8: 281–290
  Google Scholar: Author Only Title Only Author and Title

Lange MJP, Lange T (2015) Touch-induced changes in Arabidopsis morphology dependent on gibberellin breakdown. Nat Plants 1: 14025
  Google Scholar: Author Only Title Only Author and Title

Leblanc-Fournier N, Martin L, Lenne C, Decourteix M (2014) To respond or not to respond, the recurring question in plant mechanosensitivity. Front Plant Sci 5: 401
  Google Scholar: Author Only Title Only Author and Title

Lee D, Polisensky DH, Braam J (2005) Genome-wide identification of touch- and darkness-regulated Arabidopsis genes: a focus on calmodulin-like and XTH genes. New Phytol 165: 429–444
  Google Scholar: Author Only Title Only Author and Title

Lee TG, Jang CS, Kim JY, Kim DS, Park JH, Kim DY, Seo YW (2006) A Myb transcription factor (TaMyb1) from wheat roots is expressed during hypoxia: roles in response to the oxygen concentration in root environment and abiotic stresses. Physiol Plant 129: 375–385
  Google Scholar: Author Only Title Only Author and Title

Liu R, Finlayson SA (2019) Sorghum tiller bud growth is repressed by contact with the overlying leaf. Plant Cell Environ 42: 2120–2132
  Google Scholar: Author Only Title Only Author and Title

Liu X, Chu Z (2015) Genome-wide evolutionary characterization and analysis of bZIP transcription factors and their expression profiles in response to multiple abiotic stresses in Brachypodium distachyon. BMC Genomics 16: 227
  Google Scholar: Author Only Title Only Author and Title

Lobet G, Pagès L, Draye X (2011) A novel image-analysis toolbox enabling quantitative analysis of root system architecture. Plant Physiol 157: 29–39
  Google Scholar: Author Only Title Only Author and Title

MacKinnon KJ-M, Cole BJ, Yu C, Coomey JH, Hartwick NT, Remigereau M-S, Duffy T, Michael TP, Kay SA, Hazen SP (2020) Changes in ambient temperature are the prevailing cue in determining Brachypodium distachyon diurnal gene regulation. New Phytol 227: 1649–1667
  Google Scholar: Author Only Title Only Author and Title

Martin L, Leblanc-Fournier N, Julien J-L, Moulia B, Coutand C (2010) Acclimation kinetics of physiological and molecular responses of plants to multiple mechanical loadings. J Exp Bot 61: 2403–2412
  Google Scholar: Author Only Title Only Author and Title

Matos DA, Whitney IP, Harrington MJ, Hazen SP (2013) Cell walls and the developmental anatomy of the Brachypodium distachyon stem internode. PLoS One 8: e80640
  Google Scholar: Author Only Title Only Author and Title

McCahill IW, Hazen SP (2019) Regulation of cell wall thickening by a medley of mechanisms. Trends Plant Sci 24: 853–866
  Google Scholar: Author Only Title Only Author and Title

Monshausen GB, Bibikova TN, Weisenseel MH, Gilroy S (2009) Ca2+ regulates reactive oxygen species production and pH during mechanosensing in Arabidopsis roots. Plant Cell 21: 2341–2356
  Google Scholar: Author Only Title Only Author and Title

Moulia B, Coutand C, Julien J-L (2015) Mechanosensitive control of plant growth: bearing the load, sensing, transducing, and responding. Front Plant Sci 6: 52
  Google Scholar: Author Only Title Only Author and Title

Nagai K, Mori Y, Ishikawa S, Furuta T, Gamuyao R, Niimi Y, Hobo T, Fukuda M, Kojima M, Takebayashi Y, et al (2020) Antagonistic regulation of the gibberellic acid response during stem growth in rice. Nature 584: 109–114

Google Scholar: Author Only Title Only Author and Title

Niez B, Dlouha J, Moulia B, Badel E (2019) Water-stressed or not, the mechanical acclimation is a priority requirement for trees. Trees 33: 279–291
Google Scholar: Author Only Title Only Author and Title

Oliva M, Dunand C (2007) Waving and skewing: how gravity and the surface of growth media affect root development in Arabidopsis. New Phytol 176: 37–43
Google Scholar: Author Only Title Only Author and Title

O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A, Ecker JR (2016) Cistrome and epicistrome features shape the regulatory DNA landscape. Cell 165: 1280–1292
Google Scholar: Author Only Title Only Author and Title

Pitzschke A, Djamei A, Teige M, Hirt H (2009) VIP1 response elements mediate mitogen-activated protein kinase 3-induced stress gene expression. Proc Natl Acad Sci U S A 106: 18414–18419
Google Scholar: Author Only Title Only Author and Title

Polisensky DH, Braam J (1996) Cold-shock regulation of the Arabidopsis TCH genes and the effects of modulating intracellular calcium levels. Plant Physiol 111: 1271–1279
Google Scholar: Author Only Title Only Author and Title

Pomiès L, Decourteix M, Franchel J, Moulia B, Leblanc-Fournier N (2017) Poplar stem transcriptome is massively remodelled in response to single or repeated mechanical stimuli. BMC Genomics 18: 300
Google Scholar: Author Only Title Only Author and Title

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) g:Profiler: a web server for functional enrichment analysis and conversions of gene lists. Nucleic Acids Res 47: W191–W198
Google Scholar: Author Only Title Only Author and Title

Ringli C, Keller B (1998) Specific interaction of the tomato bZIP transcription factor VSF-1 with a non-palindromic DNA sequence that controls vascular gene expression. Plant Mol Biol 37: 977–988
Google Scholar: Author Only Title Only Author and Title

Roignant J, Badel É, Leblanc-Fournier N, Brunel-Michac N, Ruelle J, Moulia B, Decourteix M (2018) Feeling stretched or compressed? The multiple mechanosensitive responses of wood formation to bending. Ann Bot 121: 1151–1161
Google Scholar: Author Only Title Only Author and Title

Schütze K, Harter K, Chaban C (2008) Post-translational regulation of plant bZIP factors. Trends Plant Sci 13: 247–255
Google Scholar: Author Only Title Only Author and Title

Shah L, Yahya M, Shah SMA, Nadeem M, Ali A, Ali A, Wang J, Riaz MW, Rehman S, Wu W, et al (2019) Improving lodging resistance: using wheat and rice as classical examples. Int J Mol Sci. doi: 10.3390/ijms20174211
Google Scholar: Author Only Title Only Author and Title

Sibout R, Proost S, Hansen BO, Vaid N, Giorgi FM, Ho-Yue-Kuang S, Legée F, Cézart L, Bouchabké-Coussa O, Soulhat C, et al (2017) Expression atlas and comparative coexpression network analyses reveal important genes involved in the formation of lignified cell wall in Brachypodium distachyon. New Phytol 215: 1009–1025
Google Scholar: Author Only Title Only Author and Title

Swanson SJ, Barker R, Ye Y, Gilroy S (2015) Evaluating mechano-transduction and touch responses in plant roots. Methods Mol Biol 1309: 143–150
Google Scholar: Author Only Title Only Author and Title

Szczegielniak J, Borkiewicz L, Szurmak B, Lewandowska-Gnatowska E, Statkiewicz M, Klimecka M, Cieśla J, Muszyńska G (2012) Maize calcium-dependent protein kinase (ZmCPK11): local and systemic response to wounding, regulation by touch and components of jasmonate signaling. Physiol Plant 146: 1–14
Google Scholar: Author Only Title Only Author and Title

Trabucco GM, Matos DA, Lee SJ, Saathoff A, Priest H, Mockler TC, Sarath G, Hazen SP (2013) Functional characterization of cinnamyl alcohol dehydrogenase and caffeic acid O-methyltransferase in Brachypodium distachyon. BMC Biotechnol 13: 61
Google Scholar: Author Only Title Only Author and Title

Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ (2016) W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids Res 44: W232–W235
Google Scholar: Author Only Title Only Author and Title

Tsugama D, Liu S, Fujino K, Takano T (2018) Calcium signalling regulates the functions of the bZIP protein VIP1 in touch responses in Arabidopsis thaliana. Ann Bot 122: 1219–1229
Google Scholar: Author Only Title Only Author and Title

Tsugama D, Liu S, Takano T (2012) A bZIP protein, VIP1, is a regulator of osmosensory signaling in Arabidopsis. Plant Physiol 159: 144–155

Google Scholar: Author Only Title Only Author and Title

Tsugama D, Liu S, Takano T (2016) The bZIP protein VIP1 Is involved in touch responses in Arabidopsis roots. Plant Physiol 171: 1355–1365

Google Scholar: Author Only Title Only Author and Title

Tsugama D, Liu S, Takano T (2014) Analysis of functions of VIP1 and its close homologs in osmosensory responses of Arabidopsis thaliana. PLoS One 9: e103930

Google Scholar: Author Only Title Only Author and Title

Tyler L, Bragg JN, Wu J, Yang X, Tuskan GA, Vogel JP (2010) Annotation and comparative analysis of the glycoside hydrolase genes in Brachypodium distachyon. BMC Genomics 11: 600

Google Scholar: Author Only Title Only Author and Title

Uyttewaal M, Burian A, Alim K, Landrein B, Borowska-Wykręt D, Dedieu A, Peaucelle A, Ludynia M, Traas J, Boudaoud A, et al (2012) Mechanical stress acts via katanin to amplify differences in growth rate between adjacent cells in Arabidopsis. Cell 149: 439–451

Google Scholar: Author Only Title Only Author and Title

Valdivia ER, Herrera MT, Gianzo C, Fidalgo J, Revilla G, Zarra I, Sampedro J (2013) Regulation of secondary wall synthesis and cell death by NAC transcription factors in the monocot Brachypodium distachyon. J Exp Bot 64: 1333–1343

Google Scholar: Author Only Title Only Author and Title

Van Leene J, Blomme J, Kulkarni SR, Cannoot B, De Winne N, Eeckhout D, Persiau G, Van De Slijke E, Vercruysse L, Vanden Bossche R, et al (2016) Functional characterization of the Arabidopsis transcription factor bZIP29 reveals its role in leaf and root development. J Exp Bot 67: 5825–5840

Google Scholar: Author Only Title Only Author and Title

Van Moerkercke A, Duncan O, Zander M, Šimura J, Broda M, Vanden Bossche R, Lewsey MG, Lama S, Singh KB, Ljung K, et al (2019) A MYC2/MYC3/MYC4-dependent transcription factor network regulates water spray-responsive gene expression and jasmonate levels. Proc Natl Acad Sci U S A 116: 23345–23356

Google Scholar: Author Only Title Only Author and Title

Wang Y, Yu W, Ran L, Chen Z, Wang C, Dou Y, Qin Y, Suo Q, Li Y, Zeng J, et al (2021) DELLA-NAC interactions mediate GA signaling to promote secondary cell wall formation in cotton stem. Front Plant Sci 12: 655127

Google Scholar: Author Only Title Only Author and Title

Weltmeier F, Ehlert A, Mayer CS, Dietrich K, Wang X, Schütze K, Alonso R, Harter K, Vicente-Carbajosa J, Dröge-Laser W (2006) Combinatorial control of Arabidopsis proline dehydrogenase transcription by specific heterodimerisation of bZIP transcription factors. EMBO J 25: 3133–3143

Google Scholar: Author Only Title Only Author and Title

Yin Y, Zhu Q, Dai S, Lamb C, Beachy RN (1997) RF2a, a bZIP transcriptional activator of the phloem-specific rice tungro bacilliform virus promoter, functions in vascular development. EMBO J 16: 5247–5259

Google Scholar: Author Only Title Only Author and Title

Zha G, Wang B, Liu J, Yan J, Zhu L, Yang X (2016) Mechanical touch responses of Arabidopsis TCH1-3 mutant roots on inclined hard-agar surface. International Agrophysics 30: 65

Google Scholar: Author Only Title Only Author and Title