1

DNA damage and somatic mutations engraved on the genomes of

mammalian cells after irradiation with a nail polish dryer

Maria Zhivagui[1-3], Areebah Hoda[1], Noelia Valenzuela[3], Yi-Yu Yeh[2], Jason Dai[2], Yudou He[1-3],

Bennett Van Houten[4], and Ludmil B. Alexandrov[1-3*]

<u>Affiliations</u>

[1]Department of Cellular and Molecular Medicine, UC San Diego, La Jolla, CA, 92093, USA

[2]Department of Bioengineering, UC San Diego, La Jolla, CA, 92093, USA

[3]Moores Cancer Center, UC San Diego, La Jolla, CA, 92037, USA

[4]UPMC-Hillman Cancer Center, University of Pittsburgh, PA 15213

[*]Correspondence should be addressed to L2alexandrov@health.ucsd.edu.

ABSTRACT

Ultraviolet A light (UVA) is commonly emitted by nail polish dryers with recent reports

suggesting that long-term use of UV-nail polish dryers may increase the risk for developing skin

cancer. However, no experimental evaluation has been conducted to reveal the effect of radiation

emitted by UV-nail polish dryers on mammalian cells. Here, we examine the DNA damage and

somatic mutations found on the genomes of human and murine cells due to irradiation by a UV-

nail dryer. High levels of reactive oxygen species were observed in all UVA-irradiated samples,

consistent with elevated 8-oxo-7,8-dihydroguanine damage and mitochondrial dysfunction.

Analysis of somatic mutations revealed a dose-dependent increase of C:G>A:T substitutions in

irradiated samples with a pattern similar to the ones of COSMIC signatures SBS18/36, mutational

signatures attributed to reactive oxygen species. Re-examination of previously generated skin

cancer genomics data revealed that SBS18/36 are ubiquitously present in melanoma and account

for ~12% of the observed driver mutations. In summary, this study demonstrates that radiation

emitted by UV-nail polish dryers can both damage DNA and permanently engrave somatic

mutations on the genomes of mammalian cells. These results have far-reaching implications in

regard to public health and to preventing skin cancer due to occupational- or consumer-based

exposure to ultraviolet light from artificial sources.

INTRODUCTION

Ultraviolet (UV) light is a type of electromagnetic radiation with a wavelength ranging between 10nm and 400nm. Since wavelengths below 280nm are generally blocked by the Earth's stratospheric ozone, the UV light that reaches the Earth's surface is between 280nm and 400nm[1]. The UV spectrum can be further categorized based on its effect on human skin as well as on its ability to induce DNA damage[2]. Ultraviolet B light (UVB; 280-315nm) accounts for about 10% of the UV found on Earth, penetrates the outer layer of the skin, and it induces a plethora of DNA lesions including cyclobutane-pyrimidine dimers and 6-4 photoproducts[3,4]. In contrast, ultraviolet A light (UVA; 315-400nm) constitutes approximately 90% of the ultraviolet radiation that reaches the surface of the Earth, it can penetrate the skin more deeply, and it causes little direct DNA damage as UVA is poorly absorbed by DNA[2,5-12].

The International Agency for Research on Cancer has classified broadband UVA (315-400nm) as a Group 1 carcinogen, based on sufficient evidence for carcinogenicity in both humans and experimental models combined with strong mechanistic considerations[13]. While UVA is found in sunlight, most of UVA environmental toxicity has been attributed to the use of commercial products, such as tanning beds[14]. Consistently, meta-analyses have shown a causal relation between skin cancer and irradiation with UV-emitting tanning devices[13,15]. Further, it has been demonstrated that skin squamous-cell carcinoma will develop in mice after long-term exposure to broadband UVA[16-19]. Prior experimental studies have also suggested that UVA irradiation leads to indirect DNA damage mostly through the accumulation of 8-oxo-7,8-dihydro-2′-deoxyguanosine (8-oxo-dG) derived from reactive oxygen species[20-24]. Studies using reporter single-gene assays have identified an enrichment of C:G>A:T mutations in UVA irradiated samples consistent with

damage due to 8-oxo-dG[25-28]. Despite prior evidence for carcinogenicity of broadband UVA (315-

400nm), UVA radiation in subsets of this spectrum is widely used in a surfeit of consumer products

without extensive evaluation of the potential carcinogenic and mutagenic effects of these products.

One prominent example is UV-nail polish dryers, which have become increasingly popular in the

last decade[29,30].

UV-nail lamps are used to cure and dry nail polish formulas, known as gels, which are oligomers

requiring exposure to UV radiation to harden into polymers. These UV-gel nail devices release

UVA radiation with either very little or, in most cases, no UVB radiation[31]. UV-gel nail devices

contain multiple bulbs, emitting UV wavelengths between 340 and 395nm that can react and

activate the photo-initiators in a gel[29]. Concerns have been raised regarding the magnitude of DNA

damage that can be posed from exposure to UV-nail machines and their potential role in skin

carcinogenesis[30,32]. Notably, in most cases, both nails and hands are irradiated up to 10 minutes

with a UV-nail dryer per session[32]. The number of nail salon clients is estimated to reach 8 clients

a day per nail technician, accounting for approximately 3 million daily clients in the United

States[33]. Typically, regular users change their gel manicures every 2 weeks[32]. Recently, a small

number of melanoma and non-melanoma cases, reported either on the nail or on the dorsum of the

hand, have also been putatively attributed to exposure to UV radiation emitted by nail polish

dryers[32,34].

To evaluate the DNA damage and mutagenic effects of ultraviolet radiation emitted by nail polish

UV-dryers on mammalian cells, we performed an *in vitro* irradiation of human and murine primary

cells using distinct acute and chronic exposure protocols. Irradiated and control cells were

subjected to multiple assays for measuring DNA damage as well as to DNA sequencing either by

bulk whole-genome sequencing after clonal expansion or by single-molecule duplex sequencing

without clonal expansion (**Figure 1**).

RESULTS

Experimental Design and Examination of Cytotoxicity

To study the cytotoxic effect of irradiation by a UV-nail polish dryer, mouse embryonic fibroblasts

(MEFs), human foreskin fibroblasts (HFFs), and adult human epidermal keratinocytes (HEKa)

were exposed under several distinct conditions (**Figure 1**). Each primary cell line was irradiated

one, two, or three times, with the duration of each exposure lasting between 0 and 20 minutes. Cell

viability was measured 48 hours after the final irradiation with each condition repeated at least

three times. Analysis of cell viability revealed that UV radiation induced cytotoxicity with higher

number of exposures causing a lower cell viability (**Figure 2*a***). For example, in all cell line

models, a single 20-minute irradiation resulted in 20% to 30% cell death, while three consecutive

20-minute exposures caused between 65% and 70% cell death (q-values < 0.05; Mann-Whitney U

tests; **Figure 2*a***).

Assessment of DNA damage was performed on MEFs and HFFs after either two 20-minute

irradiations taking place within 2 hours in a single day (termed, acute exposure; **Figure 1*a&c***) or

after three 20-minute irradiations each occurring in 3 consecutive days (termed, chronic exposure;

Figure 1*a&c*). Assessment of DNA damage was also performed at the same timepoint for

unexposed MEF and HFF cells. Moreover, to assess the presence of somatic mutations, both

irradiated and control cells were subjected to barrier bypass-clonal expansion[35], grown for the

same number of doubling populations, and subjected to bulk whole-genome sequencing (**Figure

1*c***; **Supplemental Figure 1**). Specifically, irradiated and control MEF cells were grown for ~28

doubling populations prior to bulk whole-genome sequencing, while exposed and unexposed HFF

cells were grown for ~35 rounds of cell division (**Figure 1*c***; **Supplemental Figure 1**).

In the case of HEKa, cells were only chronically exposed with three 20-minute irradiations each

occurring in 3 consecutive days (**Figure 1b**). Assessment of DNA damage in HFFs was conducted

similarly to the assessments perform for MEFs and HFFs (**Figure 1c**). Nevertheless, somatic

mutations in HEKa cells were detected using an orthogonal approach to the one applied for MEFs

and HFFs by avoiding barrier bypass-clonal expansion. Specifically, irradiated and control HEKa

cells were grown for exactly 14 days, corresponding to approximately 8 doubling populations, and

were subsequently subjected to targeted single-molecule duplex sequencing[36].

Quantification of Genotoxicity

Genotoxicity of mammalian cells irradiated with a UV-nail polish dryer was evaluated 4 hours

after exposure using γH2Ax immunofluorescence[37]. Statistically significant increases in the

number of γH2Ax foci were observed when comparing irradiated MEF, HFF, and HEKa to control

cells for all cell lines as well as for all types of exposure (**Figure 2b-e**). For MEFs, the number of

γH2Ax foci increased 186-fold for acute exposure when compared to control samples (q-value:

3.6×10^{-4}; Mann-Whitney U test) and 183.3-fold for chronic exposure (q-value: 2.5×10^{-4}; **Figure**

2c). Acutely exposed HFFs had 154.7-fold elevation of the number of γH2Ax foci when compared

to control samples (q-value: 3.6×10^{-4}), while chronically exposed HFFs exhibited 137-fold

elevation (q-value: 1.2×10^{-5}; **Figure 2d**). Similarly, irradiated HEKa cells showed a significant

increase of γH2Ax foci when compared to control samples with 398-fold elevation for acute

exposure (q-value: 3×10^{-9}) and 260-fold for chronic exposure (q-value: 2×10^{-9}; **Figure 2e**).

Evaluation of Photoproducts' Formation

Immunofluorescence was used to evaluate whether irradiation from a UV-nail dryer resulted in the

formation of cyclobutane pyrimidine dimers (CPDs) or pyrimidine-pyrimidone (6-4) photoproduct

(6-4PPs) in all cell lines models after either acute or chronic exposure (**Supplementary Figure 2**).

As a positive control, each cell line model was also exposed to ultraviolet C light (UVC), which is

known to form these photoproducts[38]. Our results demonstrated that, in contrast to UVC radiation,

UV emitted by a nail-polish dryer induced neither CPDs nor 6-4PPs in any of the exposed cell

lines (**Supplementary Figure 2**).

Quantification of Oxidative Damage

To evaluate oxidative damage after irradiation, we utilized three independent *in vitro* assays,

including: *(i)* CellROX for detection of oxidative stress[39]; *(ii)* enzyme-linked immunosorbent assay

(ELISA) for quantification of 8-oxo-dG[40,41]; and *(iii)* OxiSelect™ for detection of cytosolic and

extra-cellular reactive oxygen species (ROS)[42].

CellROX green is a fluorogenic probe used for measuring oxidative stress in live cells, namely

superoxide radical anion ($O_2^•$) and hydroxyl radical ($•OH$)[39]. The reduced state of the CellROX

dye is weakly fluorescent inside cells but once oxidized by ROS, the dye binds to nucleic acids in

cells and exhibits bright green photostable fluorescence[39]. The three cell models (MEF, HFF, and

HEKa) were tested using CellROX green dye immediately after irradiation (**Figure 1c**). Live cell

imaging results unveiled a clear increase of CellROX green fluorescence in irradiated cells

compared to controls in all cases (**Figure 3a-b**). To ensure that the CellROX fluorescence is due

to ROS formation in the cells, we further challenged the HEKa cells with a ROS scavenger, namely

N-acetyl-L-cysteine (NAC), prior to irradiation[43]. Quantification of the number of CellROX foci

without NAC treatment revealed a significant increase of oxidative stress in irradiated HEKa cells

when compared to the controls (5.5-fold increase for acute, q-value $< 2.2 \times 10^{-16}$; 4.4-fold increase

for chronic, q-value $< 2.2 \times 10^{-16}$; Mann-Whitney U tests). Importantly, the number of CellROX

foci significantly decreased after NAC treatment when compared to untreated cells (acute: 5.2-

fold decrease, q-value$= 2 \times 10^{-3}$; chronic: 4.2-fold decrease, q-value$= 1.4 \times 10^{-3}$), confirming that

irradiation by a nail polish dryer is responsible for oxidative stress induction in mammalian cells.

ELISA quantifications of 8-oxo-dG were performed for all cell line modes in 3 distinct timepoints:

(i) immediately after exposure; *(ii)* 4 hours post-irradiation; and *(iii)* 24 hours post-irradiation

(**Figure 3c-e**). Additionally, three positive controls, namely, hydrogen peroxide (H_2O_2), ultraviolet

C radiation (UVC), and potassium bromate ($KBrO_3$), were also evaluated for each of the cell lines

immediately after exposure (**Figure 3c-e**). Overall, statistically significant enrichments of 8-oxo-

dG concentrations, ranging from 5- to 20-fold (q-values < 0.05), were observed in irradiated cells

when compared to the control cells for all cell line models and for all exposure types (**Figure 3c-**

e). In all cases, this level of enrichment was comparable or higher to the one exhibited by the

exposure to positive controls. Moreover, the amount of 8-oxo-dG remained stable even 24 hours

after irradiation (**Figure 3c-e**). These results confirm that irradiation by a UV-nail polish dryer

causes high levels of oxidative stress mediated by ROS formation and subsequent oxidation of

DNA bases via 8-oxo-dG formation.

To further evaluate the total free radicals in the irradiated MEF and HFF cells, we utilized

OxiSelect™ *In Vitro* ROS/RNS Assay Kit. This assay employs a proprietary quenched fluorogenic

probe, dichlorodihydrofluorescin DiOxyQ (DCFH-DiOxyQ), which is newly developed derivative

of DCFDA and has been shown to be more specific for the detection of ROS/RNS[42,44,45]. Cytosolic

and extra-cellular ROS signals were evaluated in MEF and HFF samples, in three different

timepoints: *(i)* immediately after exposure, *(ii)* 20 minutes post-irradiation, and *(iii)* 24 hours post-

irradiation. (**Supplementary Figure 3**). ROS-induced florescence was elevated in both acutely

and chronically exposed mammalian cells compared to the untreated cells. For both MEFs and

HFFs, acute exposure resulted in 2- to 8-fold increase of both cytosolic and extra-cellular ROS

compared to their respective controls (q-values: 2×10^{-5} and 1.7×10^{-6} for MEF cytosolic and

extracellular, respectively; q-values: 2×10^{-3} and 3×10^{-5} for HFF cytosolic and extracellular,

respectively; Mann-Whitney U tests; **Supplementary Figure 3**).

Examining Mitochondrial Membrane Potential and ROS Generation

As mitochondria are particularly sensitive to excessive ROS generated by UV radiation and they

play key role in oxidants production[46,47], we also examined mitochondrial potential membrane and

evaluated the mitochondrial ROS production after irradiation with a UV-nail polish dryer.

Mitochondrial activity and functionality were conducted in all three cellular models 24 hours post-

irradiation by using a tetramethylrhodamine dye (TMRM), which is taken up by mitochondria with

a high proton gradient and is a measure of well coupled mitochondria (**Figure 4*a***; **Supplementary**

Figure 4). In all three cellular models, irradiation resulted in reduced TMRM fluorescence

indicating a likely mitochondrial loss of mitochondrial membrane potential (**Figure 4*a***;

Supplementary Figure 4). Moreover, MitoSOX red mitochondrial superoxide indicator dye was

applied to HEKa cells 24 hours post-radiation. MitoSOX selectively targets the mitochondria and,

after being oxidized by superoxide, it produces red fluorescence upon binding to nucleic acid in

the cells. Live cell imaging of acutely and chronically irradiated HEKa cells revealed MitoSOX

red fluorescence when compared to unexposed HEKa controls (**Figure 4*b***). Quantification of the

number of MitoSOX foci per cell revealed a significant increase of oxidative stress in mitochondria

of irradiated HEKa cells when compared to the controls (8.3-fold increase for acute, q-value: 7.4

x 10^{-12}; 17-fold increase for chronic, q-value < 3.6 x 10^{-10}; Mann-Whitney U tests; **Figure 4c**).

This increase in MitoSOX fluorescence putatively suggests an elevation of oxidative stress in

mitochondria caused by irradiation. Taken altogether, these data suggest that UVA light can cause

mitochondrially generated ROS and persistent nuclear DNA damage.

Somatic Mutations After Barrier Bypass-clonal Expansion

To evaluate the somatic mutations engraved by UV radiation from a nail dryer, we utilized two

orthogonal sequencing approach: *(i)* bulk whole-genome sequencing of immortalized HFF and

MEF clones, grown for the same number of doubling populations after barrier bypass-clonal

expansion (**Supplementary Figure 1**); and *(ii)* single molecule duplex sequencing for HEKa cells

without clonal expansion (**Figure 1b**). A total of 15 HFF and 15 MEF immortalized clones were

subjected to bulk whole-genome sequencing (WGS) at 30x coverage, including: *(i)* 5 HFF and 5

MEF clones after acute irradiation; *(ii)* 5 HFF and 5 MEF clones after chronic irradiation; *(iii)* 5

HFF and 5 MEF unirradiated control clones. Somatic mutations were derived for immortalized

clones by comparing the WGS data of each immortalized clone to the WGS data from the

respective primary cell line using previously established state-of-the-art bioinformatics pipelines

(**Figure 1**; **Supplementary Figure 5**). Examining the variant allele frequencies (VAFs) for single

base substitutions revealed that the majority of MEF mutations are subclonal with a mean VAF of

approximately 0.25, while most HFF mutations are clonal with a mean VAF of approximately 0.50

(**Supplementary Figure 6a-b**). These results are consistent with previous observations in these

model systems[35,48] and they reflect experimental differences between the murine and human cell

line models after clonal expansion. Specifically, MEFs undergoing barrier-bypass clonal

expansion are most likely to yield two subclonal populations[35,48], while HFFs undergoing single-

cell subcloning have a high propensity for selecting a single clone[35].

Single base substitutions (SBSs) were 2- and 2.5-fold elevated in acutely and chronically irradiate

MEF cells, respectively, when compared to control cells (q-values<0.01; Mann-Whitney U tests;

Figure 5a). Stratification of SBSs based on their simplest six channel classification (*i.e.*, C>A,

C>G, C>T, T>A, T>C, T>G; each mutation referred to by the pyrimidine base of the Watson-

Crick DNA base-pair) revealed a significant increase of C>A transversions in acute and chronic

MEFs exposed to UV emitted from a nail dryer: 2.8- and 4.2-fold increase for acute and chronic,

respectively (q-values: 0.02 and 0.016; Mann-Whitney U tests; **Figure 5b**). Importantly, the

number of C>A single base substitutions found in MEF clones was positively correlated with the

number of UV exposures (**Figure 5c**; Spearman correlation: 0.64; p-value*:* 0.0098).

Immortalized HFF clones exhibited similar behaviors to the ones of MEF clones, albeit, in most

cases, with smaller effect sizes. Specifically, substitutions were 1.2- and 1.4-fold enriched in

acutely and chronically irradiated HFF cells compared to control cells (q-values<0.01; Mann-

Whitney U tests; **Figure 5a**). An elevation of C>A mutations was also observed in acutely

irradiated HFF clones (1.22-fold increase with q-value: 0.015; Mann-Whitney U tests; **Figure 5b**)

as well as in chronically irradiated HFF clones (1.63-fold increase with q-value: 7.9×10^{-3}; Mann-

Whitney U tests; **Figure 5b**). Similarly, the number of C>A transversions exhibited a positive

correlation with the number of UV exposures in HFFs (**Figure 5c**; Spearman correlation: 0.85; p-

value*:* 5.9×10^{-5}). No differences in the numbers of small insertions or deletions were observed

between the irradiated and the control cells for any of the used cell line models (**Figure 5d**).

Somatic Mutations After Duplex Sequencing

In principle, detection of somatic mutations in experimental systems requires clonal expansion[35] and, for MEF and HFF cell lines, we utilized barrier bypass-clonal expansions prior to bulk whole-genome sequencing. Recently, single-molecule duplex sequencing approaches have provided an alternative high-fidelity protocol for directly measuring the mutagenic impact of genotoxic agents without the need for clonal expansion, whilst improving the accuracy of NGS by more than 10,000-fold and enabling sensitive detection of extremely rare mutations, such as those induced by mutagens[36,49,50]. Importantly, duplex sequencing does not sequence a single cell or even a small number of cells, rather, the method generally interrogates millions of double-stranded DNA coming from many thousands of cells[51]. Thus, the mutational profile from duplex sequencing provides an average somatic mutational profile across the complete population of exposed cells[36,49,50].

We used a commercial targeted duplex sequencing protocol for assessing mutagenesis without clonal expansion[36] both to provide an orthogonal validation of the previous results and to account for any potential artifacts due to clonal expansion in MEF and HFF cell lines. Specifically, we detected somatic mutations using duplex sequencing data from 3 chronically irradiated and 3 unirradiated control HEKa cells (**Method**). Similar to irradiated HFFs and MEFs, chronically exposed HEKa exhibited 1.3-fold increase of substitutions when compared to control cells (**Figure 6a**; q-value: 0.02; Mann-Whitney test). Importantly, C>A transversions were 2.2-fold enriched in chronically irradiate HEKa cells compared to control cells (q-values: 0.04; Mann-Whitney tests; **Figure 6b**). Moreover, a positive correlation between the number of C>A transversion and the

number of UV exposures was also evident in HEKa cells (**Figure 6c**; Spearman correlation: 0.88;

p-value*:* 0.021). Consistent with prior results, no changes were observed for small insertions or

deletions between irradiated and control HEKa cells (**Figure 6d**).

Mutational Signatures Engraved by Irradiating with a UV-nail Polish Dryer

Previously, we have shown that different endogenous and exogenous mutational processes engrave

characteristic patterns of somatic mutations, termed, mutational signatures[52-54]. To evaluate the

mutational signatures engraved by irradiation with a UV-nail polish dryer, we performed separate

analysis of mutational signatures for MEFs, HFFs, and HEKa data. To explain the SBS patterns

of mutations observed in the irradiated cells, any known COSMIC signature[55] that improved the

reconstruction above the background models was allowed. For the analyses of MEF data, in

addition to known COSMIC signatures, we also allowed mouses specific mutational signatures

previously observed either *in vitro*[48] or *in vivo*[56] murine models (**Supplementary Figure 7a**).

Our analyses revealed that only COSMIC signatures SBS18 and SBS36 (SBS18/36), both

previously attributed to reactive oxygen species[57,58], were found to be operative in all irradiated

MEF and HFF cells. It should be noted that ROS signatures SBS18 and SBS36 have cosine

similarity of 0.914 (**Figure 7a**), which, in many cases, makes them hard to distinguish from one

another and these signatures are commonly reported together as SBS18/36[59-61]. Our analyses

revealed that SBS18/36 were the only signatures enriched in irradiated HFF and MEF samples

when compared to controls (**Figure 7b-c**; q-values<0.05; Mann-Whitney tests). No other

mutational signature found in these samples was elevated in irradiated samples when compared to

control samples (**Supplementary Figure 7b&d**). Moreover, the identified sets of operative

mutational signatures allowed accurately reconstructing the mutational patterns in both MEFs and

HFFs (**Supplementary Figure** 7*c&e*).

Observing signatures SBS18/36, which are almost exclusively characterized by C>A substitutions

(**Figure 7***a*), in irradiated clones confirms that radiation emitted by a UV-nail polish dryer not only

oxidizes DNA but also results in permanently engraved patterns of C>A somatic mutations on the

genomes of irradiated cells. Similarly, an almost 3-fold enrichment of signatures SBS18/36 (p-

value < 0.0001; Fisher's exact test) was observed in keratinocytes irradiated with a UV-nail polish

dryer when compared to controls. Specifically, 65% of the mutations in UV-irradiated HEKa cells

were attributed to SBS18/36 (**Figure 7***d-e*), accounting for a total of 151 mutations

(**Supplementary Figure 7***f*). In contrast, in unirradiated control samples, SBS18/36 accounted for

23% of mutations (*i.e.*, 21 mutations; **Figure 7***d-e***; Supplementary Figure 7***f-g*).

Re-examination of the recently published set of 144 whole-genome sequenced skin cancers from

the Pan-Cancer Analysis of Whole Genomes (PCAWG) project[62] revealed a positive correlation

between C>A and C>T mutations (Pearson's correlation: 0.87; p-value < 2.2 x 10^{-16}; **Figure 7***f*).

Moreover, an average skin cancer appears to harbor approximately 2,500 C>A substitutions (95%

CI: 2,015 – 2,868) in their genome with 80% of their pattern attributed to signatures SBS18/36.

However, these C>A substitutions account for only 2.3% of the mutations observed in the 144

melanomas; 85% of mutations in these melanomas are generated by signature 7 – mutational

signature attributed to cyclobutane-pyrimidine dimers and 6-4 photoproducts from ultraviolet

light[62]. Importantly, 20 of the 169 substitutions (11.83%) identified as driver mutations[62] in these

samples can be attributed to signatures SBS18/36. This indicates that, while signatures SBS18/36

generate low numbers of mutations in skin cancer, they play an important role in generating

somatic mutations that contribute towards tumor development and cancer evolution.

DISCUSSION

In this report, we employed well-controlled experimental models for UV irradiation using a UV-nail polish dryer and evaluated the DNA damage and mutagenic changes engraved on the genomes of mammalian cells (**Figure 1**). The utilized *in vitro* clonal expansion models, MEFs and HFFs, manifest the specific features required for mimicking human carcinogenesis and for recapitulating the activity of characteristic mutational processes[35]. Both MEFs and HFFs underwent clonal expansion after a barrier bypass step (senescence for MEFs and single-cell bottleneck for HFFs) leading to the formation of exposure-derived clones, which were grown for the same number of doubling populations (**Supplementary Figure 1**). As an orthogonal approach, we also utilized HEKa primary cells, which cannot go through clonal expansion in culture. Cytotoxicity and genotoxicity were observed in all irradiated cells; importantly, we observed an elevation of ROS in irradiated cells leading to increased levels of 8-oxo-dG, which was confirmed by multiple *in vitro* assays (**Figure 3**). Moreover, mitochondrial dysfunction was evident upon UVA radiation (**Figure 4** and **Supplementary Figure 4**). Genomic profiling revealed higher levels of somatic mutations in irradiated cells with C>A mutations being highly elevated in exposed samples (**Figures 5** and **6**). Consistent with the increase of ROS, COSMIC signatures SBS18/36, two mutational signatures previously attributed to reactive oxygen species[57,58], were engraved on the genomes of irradiated samples and enriched in irradiated samples when compared to controls (**Figure 7**).

Broadband UVA (315-420nm) has been extensively studied in the context of tanning devices and classified as carcinogen by the International Agency for Research on Cancer[13]. Prior experimental studies with broadband UVA have shown that it causes an accumulation of 8-oxo-dG[20-24],

generates C>A mutations in single-gene assays[25-28], and even induces tumors in mice[63]. Indeed,

prior studies have shown that UVA can generate low level of C>T somatic mutations consistent

with pyrimidine-pyrimidine photodimers[64]. Intriguingly, in this study, we demonstrate that UVA

with wavelengths between 365 and 395nm, which is generally considered to be safe and is

commonly used in a plethora of consumer products, causes DNA oxidative damage leading to

C>A mutations; no evidence was found that radiation generated by the UV-nail polish machine

generates any pyrimidine-pyrimidine photodimers (**Supplementary Figure 2**).

Our findings are also consistent with prior cancer epidemiological observations. In principle, males

have significantly higher risk than females to develop melanoma[65,66]. Surprisingly, this trend is

reversed when one examines cancer risk in melanomas of the hand in younger women.

Specifically, young females (ages 15 to 39) have approximately 2-fold higher risk for developing

melanoma of the upper extremities compared to young males[67]. This higher risk is possibly due to

lifestyle choices, one of which may be radiation from consumer products such as UV-nail polish

dryers[67]. It should be noted that exposure to UV-light from the sun cannot reasonably explain the

higher risk in young females as there is no observed differences in melanoma risk between young

males and females for other parts of the body commonly exposed to the sunlight, *viz.*, face, head,

and neck[66,67].

While this report demonstrates that radiation from UV-nail polish dryers is cytotoxic, genotoxic,

and mutagenic, it does not provide direct evidence for increased cancer risk in human beings. Prior

studies have shown that an increase in mutagenesis will likely lead to an increase in cancer risk[68-

70]. Further, several anecdotal cases have demonstrated that cancers of the hand are likely due to

radiation from UV-nail polish dryers[32,34]. Taken together, our experimental results and the prior

evidence strongly suggest that radiation emitted by UV-nail polish dryers may cause cancers of

the hand and that UV-nail polish dryers, similar to tanning beds[71], may increase the risk of early-

onset skin cancer[72]. Nevertheless, future large-scale epidemiological studies are warranted to

accurately quantify the risk for skin cancer of the hand in people regularly using UV-nail polish

dryers. It is likely that such studies will take at least a decade to complete and to subsequently

inform the general public. In the meantime, this report aims to raise awareness of the likely cancer-

causing role of artificial UV lamps and caution people who regularly use UV-nail polish dryers.

MATERIALS AND METHODS

UV-nail polish machine characteristics

A 54-Watt UV nail drying machine was purchased (model: MelodySusie), harboring 6 bulbs that

emit UV photons for curing gel nail polish. Based on the manufacturer's specifications, the UV

nail drying machine emits ultraviolet A light in wavelengths between 365 and 395nm. The UV

power density was evaluated using a UV513AB Digital UVA/UVB Meter that can measure UV

radiation between 280 and 400 nm. The machine stabilizes at $\sim$7.5 mW/cm^2, within minutes

(**Supplementary Figure 8*a-b*),** putting it on the lower end of the power density spectrum

calculated for commonly used UV-nail machines[31] (median of 10.6 mW/cm^2). In this study, in

most cases, we performed a 20-minute UV exposure session which is equivalent to a total amount

of energy delivered per unit area of 9 J/cm^2:

*7.5 mW/cm^2 * 20 minutes = 7.5 mJ/(sec*cm^2) * 1200 sec = 9000 mJ/cm^2 = 9 J/cm^2*

Cell culture, irradiation, and immortalization of mouse embryonic fibroblasts (MEFs)

Primary mouse embryonic fibroblasts (MEFs) were expanded in Advanced DMEM supplemented

with 15% fetal bovine serum, 1% penicillin/streptomycin, 1% sodium pyruvate and 1% glutamine,

and incubated in 20% O_2 and 5% CO_2. At passage 2, the primary cells were seeded in six-well

plates for 24 hours, until adherence, then exposed for 20 minutes to the UV-nail machine in pre-

warmed sterile PBS. Control cells were kept in PBS for 20 minutes. Cells were irradiated with a

UV nail polish dryer, acutely (*i.e.*, twice a day, with 60 minutes break between each of the 2

sessions) as well as chronically (*i.e.*, once every day for up to 3 consecutive days). At the end of

every irradiation, the cells were washed, and complete pre-warmed medium was replenished.

Exposed and control primary cells were cultivated until bypassed a barrier step through

senescence. Upon barrier bypass, the cells reached clonal expansion allowing the isolation of *bona*

fide cell clones (**Supplementary Figure 1a**). In all cases, irradiated and control MEF cells were

grown for ~28 doubling populations prior to bulk whole-genome sequencing. Fifteen clonally

expanded populations were successfully isolated and subjected to bulk whole-genome sequencing,

including: 5 replicates acute exposure, 5 replicates of chronic exposure, and 5 replicates for

unirradiated control cells. All cell cultures were routinely tested for the absence of mycoplasma.

Cell culture, irradiation, and immortalization of human foreskin fibroblasts (HFFs)

Primary human cells derived from human foreskin fibroblasts (HFFs) were provided to us by Dr.

John Murray (Indiana University Bloomington). Early passage cells were expanded in Advanced

DMEM supplemented with 10% fetal bovine serum and 1% glutamine in 5% CO_2 incubator.

Irradiation with a UV-nail polish dryer followed the same protocol as the one for MEFs (**Figure**

1a). Similarly, control HFF cells were kept in PBS for 20 minutes during all irradiations. Clonal

expansion was carried out using serial dilutions procedure in 96-well plates, assuming 30%

probability of a single-cell clone formation per well (**Supplementary Figure 1b**). Wells were

washed weekly, until clones reached confluency and were transferred progressively to T-75 flasks.

In all cases, irradiated and control HFF cells were grown for ~35 rounds of cell division. Fifteen

clonally expanded populations were successfully isolated and subjected to bulk whole-genome

sequencing, including: 5 replicates acute exposure, 5 replicates of chronic exposure, and 5

replicates for unirradiated control cells. All cell cultures were routinely tested for the absence of

mycoplasma.

Cell culture and irradiation of adult human epidermal keratinocytes (HEKa)

Primary human keratinocytes, derived from normal adult human epidermal keratinocytes (HEKa), were purchased from ATCC (PCS-200-011; Lot number: 70033063). Early passage cells were expanded in Dermal Cell Basal Medium (PCS-200-030) supplemented with Keratinocyte Growth Kit (PCS-200-040) and antibiotics (Penicillin: 10 Units/mL, Streptomycin: 10 µg/mL, Amphotericin B: 25 ng/mL) in 5% CO_2 incubator. At passage 2, the primary cells were seeded at a density of 2.5 x 10^3 cells/cm^2 in 10-cm dishes for 24 hours. Subsequently, cells were only chronically irradiated with a UV-nail polish dryer following the same protocol as the for MEFs and HFFs (**Figure 1b**). Similarly, control HEKa cells were kept in PBS for 20 minutes during all irradiations. In all cases, irradiated and control HEKa cells were grown for exactly 14 days, corresponding to approximately 8 rounds of cell division, which allows sufficient time for DNA adduct/damage to be fixed as somatic mutation(s) but not enough time to result in a clonal expansion[36]. Unexposed cells were grown for the same duration as irradiated cells. Six cell populations were successfully isolated and subjected to subjected to duplex sequencing, including: 3 replicates for chronic exposure and 3 replicates for unirradiated control cells. All cell cultures were routinely tested for the absence of mycoplasma.

Cell viability and cytotoxicity assays

Primary cells were seeded in 24-well plates and exposed to the UV drying device as indicated. Cell viability was measured 48 hours after treatment cessation using the Cell-Counting Kit 8 (CCK-8) from Dojindo. Plates were incubated for 3 hours at 37°C and absorbance was measured at 450 nm using the Infinite 200 Tecan *i*-control plate reader machine. The CCK-8 assay was performed in at least 4 replicates for each experimental condition. Trypan Blue exclusion assay

was also performed for assessing cell viability upon exposure, validating the choice for selecting

the irradiation condition that causes around 50% cell death (**Supplementary Figure 8c**).

Assessment of genotoxicity using γH2Ax immunofluorescence

Immunofluorescence staining was carried out using a monoclonal antibody specific for Ser139-

phosphorylated H2Ax (γH2Ax) (9718, Cell Signaling Technology). Briefly, primary cells were

seeded on coverslips in 24-well plates and, the following day, irradiated in triplicates as previously

described. Four hours after treatment cessation, the cells were fixed with 4% formaldehyde at room

temperature for 15 minutes, followed by blocking in 5% normal goat serum (5425, Life

Technologies) for 60 minutes. Subsequently, the cells were incubated with γH2Ax-antibody (1:400

in 1% BSA) at 4°C, overnight. A fluorochrome-conjugated anti-rabbit secondary antibody (4412,

Cell Signaling Technology) was then incubated for 1 hour at room temperature. ß-actin staining

was followed using phalloidin (8953, Cell Signaling Technology) incubated for 15 minutes at

room temperature. Coverslips were mounted in ProLong Gold Antifade Reagent with DAPI (8961,

Cell Signaling Technology), overnight. Immunofluorescence images were captured using a

Confocal Laser Scanning Biological Microscope Olympus FV1000 Fluoview. Quantification of

γH2Ax fluorescent cells was computed using Fiji software.

Investigation of photoproducts using CPD and 6-4PP immunofluorescence

Immunofluorescence staining was carried out using anti-cyclobutane pyrimidine dimers (CPDs)

monoclonal antibodies (clone KTM53, Kamiya Biomedical) and anti-6-4 photoproducts

monoclonal antibodies (Clone 64M-2, Cosmo Bio). Briefly, primary cells were seeded on

coverslips in 24-well plates and, the following day, irradiated in triplicates as previously described.

Subsequently, the cells were washed with ice-cold CSK buffer, containing 100mM NaCl, 300mM

Glucose, 10mM PIPES, 3mM MgCl2 and 0.5% Triton, then fixed with 4% formaldehyde at room

temperature (RT) for 15 minutes, followed by permeabilization using 0.5% Triton for 5 minutes

on ice. DNase I treatment was performed to denaturalize the genomic DNA in the cells using 20U

DNase I for 40s per coverslip (NEB; DNase I (RNase-free), M0303S). For blocking, we used 1%

BSA (A3059, Sigma) and 5% normal goat serum (5425, Life Technologies) for 60 minutes.

Afterwards, the cells were incubated with the monoclonal antibodies (1:50 in 1% BSA) at 4°C,

overnight. A fluorochrome-conjugated anti-mouse Alexa Fluor 594 secondary antibody (8890S,

Cell Signaling Technology) was then incubated for 1 hour at room temperature. Coverslips were

mounted in ProLong Gold Antifade Reagent with DAPI (8961, Cell Signaling Technology),

overnight. Immunofluorescence images were captured using Nikon A1R-STORM Super

Resolution Microscope.

Quantification of reactive oxygen species via CellROX

Oxidative stress induces the production of reactive oxygen species (ROS) and reactive nitrogen

species (RNS) in cells. We employed the CellROX™ Green Reagent for detection of oxidative

stress (ThermoFisher Scientific; catalog number: C10444), immediately after irradiation. MEF,

HFF, and HEKa primary cells were seeded on Mattek 35-mm dishes from (P35G-1.5-14-C)

appropriate for high quality live cell imaging. CellROX green was added in pre-warmed complete

media to a final concentration of 5μM and incubated for 30 minutes at 37 ºC. CellROX solution

was washed carefully 3 times with pre-warmed PBS and normal media was replenished. Live cell

imaging was conducted using Nikon A1R-STORM Super Resolution Microscope. To ensure

CellROX selectivity towards ROS components, cells were challenged with 2mM N-acetyl-L-

cysteine (NAC) for 1 hour prior to UVA exposure. NAC is a known ROS scavenger. After

CellROX addition, live cell images were used for quantification of CellROX foci number, with

and without NAC pre-treatment, using General Analysis pipeline on Nikon WorkStation 5317.

Quantification of reactive oxygen species via OxiSelect

Additionally, we also utilized the OxiSelect™ *In Vitro* ROS/RNS Assay Kit (Green Fluorescence),

from Cell Biolabs, to evaluate the level of oxidative damage induced after irradiation, intra- and

extra-cellularly, immediately after exposure as well as 20 minutes and 1 day after irradiation.

Primary MEFs and HFFs were irradiated in triplicates in a 6-well plate, twice a day (acute

exposure) and once every day for 3 consecutive days (chronic exposure). PBS solutions were

collected immediately after the last UV treatments. After every treatment, the cells were washed

with pre-warmed PBS and complete media was replenished for the accounted waiting timepoints.

Thereafter, media solutions were collected 20 minutes and 24 hours after treatment cessation.

These solutions were used to assess extracellular ROS signals. Cytosolic ROS production was

evaluated after trypsinization of the cells, lysis of cellular membrane with 0.5% TritonX-100 and

centrifugation for 5 minutes at 10,000 rpm. Samples were loaded into a 96 black well plate,

together with varying concentrations of hydrogen peroxide for generating the standard curve, and

fluorescence signals were recorded using the Infinite 200 Tecan *i*-control plate reader machine at

480 nm excitation / 530 nm emission.

Quantification of 8-hydroxy-2'-deoxyguanosine via ELISA

We quantified 8-hydroxy-2'-deoxyguanosine (8-oxo-dG) for oxidative DNA damage in DNA

samples using the EpiQuik™ 8-OHdG DNA Damage Quantification Direct Kit (Colorimetric)

from Epigentek following manufacturer instructions. Briefly, primary MEF, HFF, and HEKa

primary cells were seeded on 10-cm dishes overnight. Three positive control exposures were

employed, after concentration optimization, namely 400μM H_2O_2 for 1 hour, 20J/m^2 UVC, and

4mM $KBrO_3$ for 1 hour. Following irradiation with nail polish dryer, cells were collected at the

selected timepoints (immediately, 4 hours, and 24 hours after irradiation) and subjected to genomic

DNA extraction. Following Qubit DNA quantification, 300ng of DNA was loaded into strip wells

that are specifically treated to have a high DNA affinity. 8-oxo-dG was detected using capture and

detection antibodies. The detected signal is enhanced, and the readout was measured at 450 nm

absorbance using the Infinite 200 Tecan *i*-control plate reader machine.

Evaluation of mitochondrial ROS production and membrane potential

Image-iT™ TMRM Reagent, and MitoSOX™ Red Mitochondrial Superoxide Indicator

(ThermoFisher Scientific; catalog number: I34361 and M36008, respectively), are two dyes that

target mitochondria in live cells, the former localizing active mitochondria with normal

mitochondria membrane potential, and the latter with affinity to ROS production. MitoTracker™

Green FM (ThermoFisher Scientific, catalog number: M7514) was utilized to localize

mitochondria regardless of mitochondrial membrane potential. Briefly, primary cells were seeded

on Mattek 35-mm dishes (P35G-1.5-14-C) overnight and incubated for 30 minutes in pre-warmed

live imaging staining solution (ThermoFisher Scientific, catalog number: A14291DJ) containing

100μM MitoTracker reagent. TMRM dye was added in pre-warmed Live Imagining Solution to a

final concentration of 100nM and incubated for 30 minutes at 37 ºC. MitoSOX solution was

prepared in pre-warmed Live Imagining Solution to a final concentration of 200nM and incubated

for 10 minutes at 37 ºC. All solutions were washed carefully 3 times with pre-warmed PBS and

normal media was replenished. Live cell imaging was conducted using Nikon A1R-STORM Super

Resolution Microscope. Fluorescence data quantification was carried out using General Analysis

pipeline on Nikon WorkStation 5317.

DNA extraction and bulk whole-genome sequencing

Genomic DNA from MEF and HFF primary cells and immortalized clones were extracted using

Qiagen DNeasy Blood & Tissue kit, following the manufacturer instructions. Quality and quantity

of DNA were checked using NanoDrop and Qubit instruments. Around 2ug of DNA was thereby

extracted per each sample. High-quality DNA for 32 samples were sent to Novogene for whole-

genome library preparation and whole-genome sequencing at 30x coverage using paired-end 150

base-pair run mode with Illumina's HiSeq-XTen. The 32 samples included: *(i)* one primary HFF

and one primary MEF used as normal samples in the mutation calling; *(ii)* 5 HFF and 5 MEF

immortalized clones after acute irradiation; *(iii)* 5 HFF and 5 MEF immortalized clones after

chronic irradiation; and *(iv)* 5 HFF and 5 MEF unirradiated immortalized control clones.

Identification of somatic mutations from whole-genome bulk sequencing

FASTQ files were subjected to BWA-MEM alignment using GRCm38 and GRCh38 as reference

genomes for MEF and HFF, respectively. Our methodology for identification of somatic mutations

from bulk sequencing data follows established approaches from large genomics consortia[73,74].

Briefly, ensemble variant calling of somatic mutations was performed using three independent

variant callers Mutect2[75], VarScan2[76], Strelka2[77], and MuSe[78]. Any mutation identified by at least

two out of the four variant callers was considered a *bona fide* mutation. *Bona fide* mutations were

subsequently filtered to remove any residual SNPs based on dbSNP annotation by variant effect

predictor[79]. Further, any mutations shared between two or more samples and clustered mutations

were removed as these reflect either residual germline mutations or mutations under positive

selection (**Supplementary Figure 5**). Overall, a total of 118,429 unique somatic mutations were

detected across all sequenced samples prior to filtering. Consistent with prior datasets[73,74], the

germline filtering removed 18.7% of these mutations with another 5.2% removed by the clustered

filter. The remaining set of somatic mutations were used in the subsequent analyses and the

evaluation for mutational signatures.

DNA extraction and duplex sequencing

Genomic DNA from HEKa cells were extracted using Qiagen DNeasy Blood & Tissue kit,

following manufacturer instructions, with one exception during the initial proteinaseK digestion

for which the samples were incubated at 37℃ rather than 54℃ for 1 hour, and DNA yield was

eluted in IDTE buffer (10mM Tris HCl pH 8.0 + 0.1mM EDTA). Quality and quantity of DNA

were checked using NanoDrop and Qubit instruments. High-quality DNA for 7 samples were sent

to TwinStrand Biosciences for targeted library preparation, targeted duplex sequencing, and data

analysis. The seven samples included: *(i)* one primary HEKa sample used in the mutation calling;

(ii) 3 HEKa samples after chronic irradiation; and *(iii)* 3 HEKa unirradiated immortalized control

clones. Briefly, duplex library preparation was performed using a human mutagenesis panel (~50

kb distributed across the whole genome in regions not predicted to be under positive or negative

selection) on DNA input sufficient to generate between 500 million to 1 billion informative Duplex

base pairs per sample. Each sample was prepared in two batches: one with the standard

mutagenesis assay and one with a prototype rapid workflow version of the assay. All libraries were

sequenced on an Illumina NovaSeq 6000.

Identification of somatic mutations from duplex sequencing

All raw NovaSeq sequencing FASTQ data was processed through the TwinStrand cloud-based

human mutagenesis pipeline to generate error-corrected BAM files and variant call files for each

sample. Mutagenesis analysis was carried out using the *Min* assumption for mutant frequency

calculation as standardly done[36]. This method counts each variant only once, regardless of the

number of reads that contain the non-reference allele.

$$Mutant\ Frequency = \frac{\text{Number of Unique Variants}}{\text{Total non N Duplex Bases}}$$

The remaining set of somatic mutations were used in the subsequent analyses and evaluation for

mutational signatures.

Duplex sequencing yield results

Average raw reads per sample	**~239 million**
Average duplex bases per sample	**~982 million**
Average mean on target duplex depth	**15,204x**
Interspecies contamination	**Negative**
Inter-sample contamination	**Negative**
Hybrid selection efficacy	**~99.81%**
Variants VAF used	**≤ 1%**

Analysis of mutational signatures and additional examinations

Analysis of mutational signatures was performed using our previously derived set of reference

mutational signatures[55] as well as our previously established methodology with the SigProfiler

suite of tools used for summarization, simulation, visualization, and assignment of mutational

signatures[62,80,81]. Variant allele frequencies (VAF) were calculated using integrative genomics

600   viewer[82]. R version 3.6.1 was used to plot data (*i.e.,* ggplot2, easyGgplot2, ComplexHeatmap[83],

601   and circlize[84] packages), to compute p-values (ggpubr package), to perform correlation analyses

602   (corrr package), and to compute the cosine similarity (lsa[85] package).

 FIGURE LEGENDS

Figure 1. Overview of the overall study design. *(a)* Primary mammalian cells, MEFs and HFFs,

were expanded into 6-well plates and treated with ultraviolet light (UV) emitted from a UV-nail

polish dryer for 20 minutes, twice a day within one single day, termed *acute UV exposure*. For

chronic UV exposure, primary cells were exposed consecutively in three different days with each

exposure lasting 20 minutes. Control samples were maintained in the dark in pre-warmed PBS for

20 minutes during each exposure session. After recovery and cellular selection, whether through

senescence bypass or single-cell subcloning, control and irradiated cell were grown for the same

number of doubling populations and subjected to bulk whole-genome sequencing. Analysis of

whole-genome sequenced samples was performed using our established pipelines for mutation

calling. Analysis of mutational signatures was performed using the SigProfiler suite of tools. *(b)*

Primary HEKa cells were irradiated in 10-cm dishes using UV-nail polish dryer for 20 minutes for

three consecutive days. Control samples were maintained in the dark in pre-warmed PBS for 20

minutes during each exposure session. HEKa cells were kept in culture for recovery and

replication, allowing around 8 doubling-populations after which the cells were subjected to high

coverage duplex Sequencing. Analysis of duplex sequencing samples was performed using

commercially established pipelines. Analysis of mutational signatures was performed using the

SigProfiler suite of tools. *(c)* Asterisks in panels *(a)* and *(b)* denote timepoints when DNA damage

and other assessments were performed for each condition, including interrogation of cytotoxicity,

genotoxicity, oxidative damage, and mitochondrial damage.

Figure 2. Cytotoxicity and genotoxicity in mammalian cells after irradiation with a UV-nail

dryer. *(a)* Cytotoxicity assessment following exposure of primary MEFs (top panel), HFFs

(middle panel), and HEKa (bottom panel) to UV radiation emitted from a UV-nail polish dryer for

different timepoints, ranging from 0 to 20 minutes. Multiple UV-exposure sessions were tested

with one hour difference between each consecutive exposure, including: grey – one exposure in a

day; yellow – two exposures in a day; red – three exposures in a day. Formazan dye absorbance

was measured 48 hours after treatment cessation and was normalized to the number of unirradiated

control cells at timepoint 0. The results are presented as mean percentage ± standard deviation

from at least four replicates. *(b)* DNA damage evaluation by immunofluorescence of Ser139-

pjosphorylated histone H2Ax (γH2Ax). Primary MEFs (top panel), HFFs (middle panel) and

HEKa (bottom panel), were exposed to UV radiation either acutely or chronically. DAPI is shown

in blue, Phalloidin in red, and γH2Ax in green. *(c-e)* Quantification of the number of γH2Ax foci

was performed by analyzing at least 100 cells per condition, collected from at least triplicates, in

MEF *(c)*, HFF *(d)* and HEKa *(e)* cells. Statically significant results from FDR corrected Mann-

Whitney U tests are denoted as: ***q-value<0.001 and ****q-value<0.0001.

Figure 3. Oxidative damage in mammalian cells after irradiation with a UV-nail dryer. (*a*)

Assessment of oxidative DNA damage by live imaging of CellROX green reagent in control and

irradiated MEF (top panel), HFF (middle panel), and HEKa (bottom panel) cells. DAPI is shown

in blue while CellROX is shown in green. *(b)* HEKa cells were challenged with 2mM N-acetyl-L-

cysteine (NAC) for 1 hour prior to irradiation. Quantification of the number of CellROX foci was

performed by analyzing at least 100 cells per condition and comparing cells with and without NAC

pre-treatment, collected from triplicates. Statically significant results from FDR corrected Mann-

Whitney U tests are denoted as: ***q-value<0.001 and ****q-value<0.0001. *(c-e)* Oxidative

DNA damage measurement of 8-oxo-dG by ELISA in MEF *(c)*, HFF *(d)*, and HEKa *(e)* cells.

Measurements of 8-oxo-dG were performed at different timepoints (0, 4, 24 hours post-treatment).

Measurements of 8-oxo-dG were also performed on a negative control (NC), consisting of water

only, and on three positive controls, namely, hydrogen peroxide (H_2O_2), ultraviolet C radiation

(UVC), and potassium bromate ($KBrO_3$). Statically significant results from FDR corrected Student

t-tests are denoted as: *q-value<0.05; *q-value<0.01; ***q-value<0.001; and ****q-

value<0.0001.

Figure 4. Mitochondrial disruption after irradiation with a UV-nail dryer in HEKa cells. *(a)*

Evaluating the mitochondrial membrane potential after irradiation of HEKa cells with a UV-nail

dryer. DAPI is shown in blue, MitoTracker in green, and tetramethylrhodamine dye (TMRM) in

red. Yellow corresponds to overlaps between MitoTracker and TMRM. *(b)* Examining

mitochondrial ROS production in HEKa cells. DAPI is shown in blue, MitoTracker in green, and

MitoSOX in red. Yellow corresponds to overlaps between MitoTracker and MitoSOX. In both

panels *(a)* and *(b)*, MitoTracker green reagent is used to localize mitochondria regardless of

mitochondria membrane potential. *(c)* Quantification of the number of MitoSOX foci in at least

100 cells per condition in HEKa, collected from at least triplicates. Statically significant results

from FDR corrected Mann-Whitney U tests are denoted as: ***q-value<0.001 and ****q-

value<0.0001.

Figure 5. Mutations found in the genomes of MEFs and HFFs irradiated by a UV-nail dryer.

(a) Mutation count per megabase (Mb) detected in the different conditions, represented in colors,

in MEFs and HFFs. *(b)* Fold-increase of single base substitutions in UV-treated clones compared

to controls. Fold increase is expressed as mean fold-change ± SE (standard error). *(c)* Spearman's

correlations between the number of C>A substitutions and the number UV exposures in the five independent UV-treated clones. Acute and chronic exposures correspond to 2 and 3 exposures, respectively. *(d)* Number of small insertions and deletions (indels) in MEF and HFF clones for each irradiation condition. *: q-value<0.05; **: q-value<0.01; ***: q-value<0.001; ****: q-value<0.0001.

Figure 6. Mutation analysis of HEKa cells irradiated by a UV-nail dryer. *(a)* Mutation count per megabase (Mb) detected in the different conditions, represented in colors, in MEFs and HFFs. *(b)* Fold-increase of single base substitutions in UV-treated clones compared to controls. Fold increase is expressed as mean fold-change ± SE (standard error). *(c)* Spearman's correlations between the number of C>A substitutions and the number UV exposures in each of the 3 biological replicates. *(d)* Number of small insertions and deletions (indels) in HEKa cells.

Figure 7. Somatic Mutations due to SBS18 in cells and PCAWG samples. *(a)* Mutational signatures COSMIC SBS18 and COSMIC SBS36 observed in UV irradiated cells. *(b)* Number of mutations per megabase attributed to COSMIC SBS18/36 in UV exposed MEF clones, per condition, *i.e.*, control (n=3/5), chronic (n=5/5), and acute (n=5/5) exposure. *(c)* Number of mutation per megabase attributed to COSMIC SBS18/36 in UV exposed HFF clones, per condition, *i.e.*, control (n=3/5), chronic (n=5/5), and acute (n=5/5) exposure. *(d)* Mutational patterns of irradiated (upper track) and control (lower track) primary HEKa cells. *(e)* COSMIC signatures assignment of UV-irradiated HEKa cells (upper track) and control (lower track). *(f)* Correlation analysis of C>A and C>T mutations in 144 PCAWG skin cancer samples. *(g)* Percentage contribution of the C>A mutations to SBS18/36 in PCAWGS skin cancer samples.

 SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1. Exposure and barrier bypass-clonal expansion assays. *(a)* MEFs

were treated at early passages and led through senescence manifested by a plateau-like curve where

cells lost the ability to efficiently duplicate and grow. Cells went through senescence bypass and

eventually emerge into immortalized clones, able to multiply indefinitely, a process called barrier-

bypass clonal expansion, emulating oncogenesis. Irradiated and control MEF cells were grown for

~28 doubling populations prior to bulk whole-genome sequencing (WGS). Exemplars are provided

for one control, one acutely irradiated, and one chronically irradiated MEF. Y-axes reflect the

number of doubling populations, while X-axes correspond to the number of days in culture. *(b)*

HFFs were treated at early passages and led through recovery phase before initiating a single cell

subcloning assay using a serial dilutions technic. *Bona fide* single-cell clones were collected at the

end of experiment. Irradiated and control HEF cells were grown for ~35 doubling populations

prior to bulk whole-genome sequencing (WGS). Exemplars are provided for one control, one

acutely irradiated, and one chronically irradiated HFF. Y-axes reflect the number of doubling

populations, while X-axes correspond to the number of days in culture.

Supplementary Figure 2. Evaluation of cyclobutane pyrimidine dimers and 6-4

photoproducts irradiated by a UV-nail dryer. *(a-c)* Immunofluorescence of cyclobutane

pyrimidine dimers (CPDs) and 6-4 photoproducts (6-4PPs) in MEF *(a)*, HFF *(b)*, and HEKa *(c)*

cells after acute or chronic irradiation with a UV-nail dryer. DAPI is shown in blue while CPDs

and 6-4PPs are shown in red. A positive control, namely, cells irradiated with ultraviolet C (UVC)

light (1x 10J/m^2), is also provided for each cell line.

Supplementary Figure 3. Cytosolic and extra-cellular measurement of reactive oxygen species in MEFs and HFFs. Accumulation of reactive oxygen species (ROS) through time in primary MEFs (left panel) and HFFs (right panel) after acute or chronic irradiation with a UV-nail dryer. Each circle corresponds to a single experiment for a particular exposure. All y-axes reflect, either cytosolic or extracellular, relative ROS fluorescence by normalizing the fluorescence of a particular exposure to the average fluorescence of the appropriate controls. Statically significant results from FDR corrected Mann-Whitney U tests are denoted as: ***q-value<0.001 and ****q-value<0.0001.

Supplementary Figure 4. Mitochondrial membrane potential after irradiation with a UV-nail dryer. Evaluating the mitochondrial membrane potential after irradiation of MEF cells *(a)* and HFF cells *(b)* with a UV-nail dryer. DAPI is shown in blue, MitoTracker in green, and tetramethylrhodamine dye (TMRM) in red. Yellow corresponds to overlaps between MitoTracker and TMRM.

Supplementary Figure 5. Schematic of whole-genome data analysis. Primary cells were sequenced and used as a germline reference. We employed three variant callers (Mutect2, VarScan2, Strelka2, and MuSe) in matched tumor-normal mode, with the primary cells' BAM files as normal for each cell line. We selected variants called in at least 2 of the variant callers in order to increase variant calling quality. Germline single nucleotide polymorphisms (SNP) were removed using the dbSNP database. Any clustered mutation as well as any mutation observed in 2 or more samples were also removed. The final set of somatic mutations were analyzed by the SigProfiler suite of tools.

Supplementary Figure 6. Analysis of variant allelic frequency analysis. *(a)* Variant allelic

frequency of *bona fide* mutations in MEF clones for each irradiation condition. *(b)* Variant allelic

frequency of *bona fide* mutations in HFF clones for each irradiation condition.

Supplementary Figure 7. Mutation analysis of UV-treated MEF and HFF clones and

PCAWG samples. *(a)* Mutational signature mSBS_N3 identified *in vivo* in mouse models. *(b)*

COSMIC signatures assigned to MEF clones per condition (control, acute and chronic). *(c)*

Mutational signatures reconstruction of MEF mutational patterns. Accuracy is measured in cosine

similarity between the original pattern of a sample and the pattern of a sample reconstructed using

the assigned mutational signatures. *(d)* COSMIC signatures assigned to HFF clones per condition

(control, acute and chronic). *(e)* Mutational signatures reconstruction of HFF mutational patterns.

(f) COSMIC signatures assigned to UV-irradiated HEKa cells and control, grouped by SBS18/36

and other signatures. Numbers reflect the number of somatic mutations assigned to the signatures

(upper track). Percentages reflect the number of somatic mutations assigned to the signatures

divided by the total number of mutations (lower track). *(g)* Mutational signatures reconstruction

of HEKa mutational patterns. *(h)* Correlation analysis of C>A and C>T mutations using PCAWG

(whole-genome) and TCGA (whole-exome) data of skin cancer samples. *(i)* Percentage

contribution of the C>A mutations to SBS18/36 in PCAWG and TCGA skin cancer samples.

Supplementary Figure 8. Toxicity and characteristics of the UV-nail polish dryer machine.

(a) Intensity of UV radiation (mW/cm^2) at different position points (vertical and horizontal) in the

employed UV nail polish dryer. *(b)* Stability assessment of the UV intensity emitted from the UV-

nail polish dryer over time.

ACKNOWLEDGEMENTS

The authors would like to thank Dr. John Murray (Indiana University Bloomington) for

providing human foreskin fibroblasts. This work was supported by Alfred P. Sloan Research

Fellowship. LBA is also an Abeloff V scholar and he is personally supported by a Packard

Fellowship for Science and Engineering. This work was also supported by the US National

Institute of Health grants R01ES030993-01A1, R01ES032547-01, and R01CA269919-01 to

LBA as well as R35ES031638-03 to BVH. The funders had no roles in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

AUTHOR CONTRIBUTIONS

LBA and MZ designed all experiments with help from BVH. MZ performed all experiments

with assistance from AH, NV and YY. MZ performed all data analysis with help from CY, JD,

and YH. BVH assisted in the writing of the manuscript and helped interpreting the experimental

results. The manuscript was written by LBA and MZ with input from all other authors. LBA

supervised the overall project and writing of the manuscript. All authors read and approved the

final manuscript.

COMPETING INTERESTS

LBA is a compensated consultant and has equity interest in io9, LLC. His spouse is an employee

of Biotheranostics, Inc. LBA is also an inventor of a US Patent 10,776,718 for source identification

by non-negative matrix factorization. LBA declares provisional patent applications for "Clustered

mutations for the treatment of cancer" (U.S. provisional application serial number 63/289,601),

"Artificial intelligence architecture for predicting cancer biomarker" (serial number 63/269,033),

"Methods and systems for detecting homologous recombination deficiency in cancer therapies"

(serial number 63/366,392), and "Drugs for treating head and neck cancers with chromosome 9p

loss" (serial number 63/367,846). All other authors declare no competing interests.

DATA AVAILABILITY

All whole-genome and duplex sequencing data have been deposited to Sequence Read Archive

(SRA) and can be downloaded using accession number: PRJNA667106.

REFERENCES

Khan, A. Q., Travers, J. B. & Kemp, M. G. Roles of UVA radiation and DNA damage responses in melanoma pathogenesis. *Environ Mol Mutagen* **59**, 438-460, doi:10.1002/em.22176 (2018).

Pfeifer, G. P. Mechanisms of UV-induced mutations and skin cancer. *Genome Instab Dis* **1**, 99-113, doi:10.1007/s42764-020-00009-8 (2020).

You, Y. H., Szabó, P. E. & Pfeifer, G. P. Cyclobutane pyrimidine dimers form preferentially at the major p53 mutational hotspot in UVB-induced mouse skin tumors. *Carcinogenesis* **21**, 2113-2117, doi:10.1093/carcin/21.11.2113 (2000).

Pfeifer, G. P. Formation and processing of UV photoproducts: effects of DNA sequence and chromatin environment. *Photochem Photobiol* **65**, 270-283, doi:10.1111/j.1751-1097.1997.tb08560.x (1997).

Cadet, J. *et al.* Effects of UV and visible radiation on DNA-final base damage. *Biol Chem* **378**, 1275-1286 (1997).

Cadet, J., Douki, T. & Ravanat, J. L. Oxidatively generated damage to cellular DNA by UVB and UVA radiation. *Photochem Photobiol* **91**, 140-155, doi:10.1111/php.12368 (2015).

Zhang, X. *et al.* Induction of 8-oxo-7,8-dihydro-2'-deoxyguanosine by ultraviolet radiation in calf thymus DNA and HeLa cells. *Photochem Photobiol* **65**, 119-124, doi:10.1111/j.1751-1097.1997.tb01886.x (1997).

Besaratinia, A. *et al.* DNA lesions induced by UV A1 and B radiation in human cells: comparative analyses in the overall genome and in the p53 tumor suppressor gene. *Proc Natl Acad Sci U S A* **102**, 10058-10063, doi:10.1073/pnas.0502311102 (2005).

Schuch, A. P., Moreno, N. C., Schuch, N. J., Menck, C. F. M. & Garcia, C. C. M. Sunlight damage to cellular DNA: Focus on oxidatively generated lesions. *Free Radic Biol Med* **107**, 110-124, doi:10.1016/j.freeradbiomed.2017.01.029 (2017).

Brem, R., Li, F. & Karran, P. Reactive oxygen species generated by thiopurine/UVA cause irreparable transcription-blocking DNA lesions. *Nucleic Acids Res* **37**, 1951-1961, doi:10.1093/nar/gkp070 (2009).

Douki, T. & Sage, E. Dewar valence isomers, the third type of environmentally relevant DNA photoproducts induced by solar radiation. *Photochem Photobiol Sci* **15**, 24-30, doi:10.1039/c5pp00382b (2016).

Rochette, P. J. *et al.* UVA-induced cyclobutane pyrimidine dimers form predominantly at thymine-thymine dipyrimidines and correlate with the mutation spectrum in rodent cells. *Nucleic Acids Res* **31**, 2786-2794, doi:10.1093/nar/gkg402 (2003).

El Ghissassi, F. *et al.* A review of human carcinogens--part D: radiation. *Lancet Oncol* **10**, 751-752, doi:10.1016/s1470-2045(09)70213-x (2009).

Sample, A. & He, Y. Y. Mechanisms and prevention of UV-induced melanoma. *Photodermatol Photoimmunol Photomed* **34**, 13-24, doi:10.1111/phpp.12329 (2018).

cancer, I. A. f. R. o. C. W. G. o. a. u. U. l. a. s. The association of use of sunbeds with cutaneous malignant melanoma and other skin cancers: A systematic review. *Int J Cancer* **120**, 1116-1122, doi:10.1002/ijc.22453 (2007).

van Weelden, H., de Gruijl, F. R., van der Putte, S. C., Toonstra, J. & van der Leun, J. C. The carcinogenic risks of modern tanning equipment: is UV-A safer than UV-B? *Arch Dermatol Res* **280**, 300-307, doi:10.1007/bf00440604 (1988).

843   17   van Weelden, H., van der Putte, S. C., Toonstra, J. & van der Leun, J. C. UVA-induced
844   tumours in pigmented hairless mice and the carcinogenic risks of tanning with UVA. *Arch*
845   *Dermatol Res* **282**, 289-294, doi:10.1007/bf00375721 (1990).
846   18   Sterenborg, H. J. & van der Leun, J. C. Tumorigenesis by a long wavelength UV-A source.
847   *Photochem Photobiol* **51**, 325-330, doi:10.1111/j.1751-1097.1990.tb01718.x (1990).
848   19   Kelfkens, G., de Gruijl, F. R. & van der Leun, J. C. Tumorigenesis by short-wave
849   ultraviolet A: papillomas versus squamous cell carcinomas. *Carcinogenesis* **12**, 1377-
850   1382, doi:10.1093/carcin/12.8.1377 (1991).
851   20   Greinert, R. *et al.* UVA-induced DNA double-strand breaks result from the repair of
852   clustered oxidative DNA damages. *Nucleic Acids Res* **40**, 10263-10273,
853   doi:10.1093/nar/gks824 (2012).
854   21   Didier, C., Emonet-Piccardi, N., Béani, J. C., Cadet, J. & Richard, M. J. L-arginine
855   increases UVA cytotoxicity in irradiated human keratinocyte cell line: potential role of
856   nitric oxide. *FASEB J* **13**, 1817-1824, doi:10.1096/fasebj.13.13.1817 (1999).
857   22   Francis, A. J. & Giannelli, F. Cooperation between human cells sensitive to UVA
858   radiations: a clue to the mechanism of cellular hypersensitivity associated with different
859   clinical conditions. *Exp Cell Res* **195**, 47-52, doi:10.1016/0014-4827(91)90498-j (1991).
860   23   Eisenstark, A. Mutagenic and lethal effects of near-ultraviolet radiation (290-400 nm) on
861   bacteria and phage. *Environ Mol Mutagen* **10**, 317-337, doi:10.1002/em.2850100311
862   (1987).
863   24   Ridley, A. J., Whiteside, J. R., McMillan, T. J. & Allinson, S. L. Cellular and sub-cellular
864   responses to UVA in relation to carcinogenesis. *Int J Radiat Biol* **85**, 177-195,
865   doi:10.1080/09553000902740150 (2009).
866   25   Besaratinia, A., Bates, S. E., Synold, T. W. & Pfeifer, G. P. Similar mutagenicity of
867   photoactivated porphyrins and ultraviolet A radiation in mouse embryonic fibroblasts:
868   involvement of oxidative DNA lesions in mutagenesis. *Biochemistry* **43**, 15557-15566,
869   doi:10.1021/bi048717c (2004).
870   26   Besaratinia, A., Kim, S. I., Bates, S. E. & Pfeifer, G. P. Riboflavin activated by ultraviolet
871   A1 irradiation induces oxidative DNA damage-mediated mutations inhibited by vitamin C.
872   *Proc Natl Acad Sci U S A* **104**, 5953-5958, doi:10.1073/pnas.0610534104 (2007).
873   27   Kim, S. I., Pfeifer, G. P. & Besaratinia, A. Mutagenicity of ultraviolet A radiation in the
874   lacI transgene in Big Blue mouse embryonic fibroblasts. *Mutat Res* **617**, 71-78,
875   doi:10.1016/j.mrfmmm.2006.12.003 (2007).
876   28   Pfeifer, G. P., You, Y. H. & Besaratinia, A. Mutations induced by ultraviolet light. *Mutat*
877   *Res* **571**, 19-31, doi:10.1016/j.mrfmmm.2004.06.057 (2005).
878   29   Rieder, E. A. & Tosti, A. Cosmetically Induced Disorders of the Nail with Update on
879   Contemporary Nail Manicures. *J Clin Aesthet Dermatol* **9**, 39-44 (2016).
880   30   Bollard, S. M. *et al.* Skin cancer risk and the use of UV nail lamps. *Australas J Dermatol*
881   **59**, 348-349, doi:10.1111/ajd.12806 (2018).
882   31   Shipp, L. R., Warner, C. A., Rueggeberg, F. A. & Davis, L. S. Further investigation into
883   the risk of skin cancer associated with the use of UV nail lamps. *JAMA Dermatol* **150**, 775-
884   776, doi:10.1001/jamadermatol.2013.8740 (2014).
885   32   Ratycz, M. C., Lender, J. A. & Gottwald, L. D. Multiple Dorsal Hand Actinic Keratoses
886   and Squamous Cell Carcinomas: A Unique Presentation following Extensive UV Nail
887   Lamp Use. *Case Rep Dermatol* **11**, 286-291, doi:10.1159/000503273 (2019).

888 33 Ceballos, D. M. *et al.* Biological and environmental exposure monitoring of volatile
889   organic compounds among nail technicians in the Greater Boston area. *Indoor Air* **29**, 539-
890   550, doi:10.1111/ina.12564 (2019).

891 34 MacFarlane, D. F. & Alonso, C. A. Occurrence of nonmelanoma skin cancers on the hands
892   after UV nail light exposure. *Arch Dermatol* **145**, 447-449,
893   doi:10.1001/archdermatol.2008.622 (2009).

894 35 Zhivagui, M., Korenjak, M. & Zavadil, J. Modelling Mutation Spectra of Human
895   Carcinogens Using Experimental Systems. *Basic Clin Pharmacol Toxicol* **121 Suppl 3**,
896   16-22, doi:10.1111/bcpt.12690 (2017).

897 36 Schmitt, M. W. *et al.* Detection of ultra-rare mutations by next-generation sequencing.
898   *Proc Natl Acad Sci U S A* **109**, 14508-14513, doi:10.1073/pnas.1208715109 (2012).

899 37 Sharma, A., Singh, K. & Almasan, A. Histone H2AX phosphorylation: a marker for DNA
900   damage. *Methods Mol Biol* **920**, 613-626, doi:10.1007/978-1-61779-998-3_40 (2012).

901 38 Budden, T. & Bowden, N. A. The role of altered nucleotide excision repair and UVB-
902   induced DNA damage in melanomagenesis. *Int J Mol Sci* **14**, 1132-1151,
903   doi:10.3390/ijms14011132 (2013).

904 39 Yang, Z. & Choi, H. Single-Cell, Time-Lapse Reactive Oxygen Species Detection in E.
905   coli. *Curr Protoc Cell Biol* **80**, e60, doi:10.1002/cpcb.60 (2018).

906 40 Suzuki, T. & Kamiya, H. Mutations induced by 8-hydroxyguanine (8-oxo-7,8-
907   dihydroguanine), a representative oxidized base, in mammalian cells. *Genes Environ* **39**,
908   2, doi:10.1186/s41021-016-0051-y (2017).

909 41 Douki, T. *et al.* Oxidation of guanine in cellular DNA by solar UV radiation: biological
910   role. *Photochem Photobiol* **70**, 184-190 (1999).

911 42 KESTON, A. S. & BRANDT, R. THE FLUOROMETRIC ANALYSIS OF
912   ULTRAMICRO QUANTITIES OF HYDROGEN PEROXIDE. *Anal Biochem* **11**, 1-5,
913   doi:10.1016/0003-2697(65)90034-5 (1965).

914 43 Ezeriņa, D., Takano, Y., Hanaoka, K., Urano, Y. & Dick, T. P. N-Acetyl Cysteine
915   Functions as a Fast-Acting Antioxidant by Triggering Intracellular H. *Cell Chem Biol* **25**,
916   447-459.e444, doi:10.1016/j.chembiol.2018.01.011 (2018).

917 44 BRANDT, R. & KESTON, A. S. SYNTHESIS OF
918   DIACETYLDICHLOROFLUORESCIN: A STABLE REAGENT FOR
919   FLUOROMETRIC ANALYSIS. *Anal Biochem* **11**, 6-9, doi:10.1016/0003-
920   2697(65)90035-7 (1965).

921 45 Bass, D. A. *et al.* Flow cytometric studies of oxidative product formation by neutrophils: a
922   graded response to membrane stimulation. *J Immunol* **130**, 1910-1917 (1983).

923 46 Murphy, M. P. How mitochondria produce reactive oxygen species. *Biochem J* **417**, 1-13,
924   doi:10.1042/BJ20081386 (2009).

925 47 Brand, R. M. *et al.* Targeting Mitochondrial Oxidative Stress to Mitigate UV-Induced Skin
926   Damage. *Front Pharmacol* **9**, 920, doi:10.3389/fphar.2018.00920 (2018).

927 48 Zhivagui, M. *et al.* Experimental and pan-cancer genome analyses reveal widespread
928   contribution of acrylamide exposure to carcinogenesis in humans. *Genome Res* **29**, 521-
929   531, doi:10.1101/gr.242453.118 (2019).

930 49 Salk, J. J., Schmitt, M. W. & Loeb, L. A. Enhancing the accuracy of next-generation
931   sequencing for detecting rare and subclonal mutations. *Nat Rev Genet* **19**, 269-285,
932   doi:10.1038/nrg.2017.117 (2018).

50   Salk, J. J. & Kennedy, S. R. Next-Generation Genotoxicology: Using Modern Sequencing
Technologies to Assess Somatic Mutagenesis and Cancer Risk. *Environ Mol Mutagen* **61**,
135-151, doi:10.1002/em.22342 (2020).

51   Abbasi, A. & Alexandrov, L. B. Significance and limitations of the use of next-generation
sequencing technologies for detecting mutational signatures. *DNA Repair (Amst)* **107**,
103200, doi:10.1016/j.dnarep.2021.103200 (2021).

52   Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**,
415-421, doi:10.1038/nature12477 (2013).

53   Alexandrov, L. B. & Stratton, M. R. Mutational signatures: the patterns of somatic
mutations hidden in cancer genomes. *Curr Opin Genet Dev* **24**, 52-60,
doi:10.1016/j.gde.2013.11.014 (2014).

54   Alexandrov, L. B. & Zhivagui, M. in *Encyclopedia of Cancer (Third Edition)*   499-510
(2019).

55   Alexandrov, L. B., Nik-Zainal, S., Wedge, D. C., Campbell, P. J. & Stratton, M. R.
Deciphering signatures of mutational processes operative in human cancer. *Cell Rep* **3**,
246-259, doi:10.1016/j.celrep.2012.12.008 (2013).

56   Riva, L. *et al.* The mutational signature profile of known and suspected human carcinogens
in mice. *Nat Genet* **52**, 1189-1197, doi:10.1038/s41588-020-0692-4 (2020).

57   Pilati, C. *et al.* Mutational signature analysis identifies MUTYH deficiency in colorectal
cancers and adrenocortical carcinomas. *J Pathol* **242**, 10-15, doi:10.1002/path.4880
(2017).

58   Viel, A. *et al.* A Specific Mutational Signature Associated with DNA 8-Oxoguanine
Persistence in MUTYH-defective Colorectal Cancer. *EBioMedicine* **20**, 39-49,
doi:10.1016/j.ebiom.2017.04.022 (2017).

59   Georgeson, P. *et al.* Identifying colorectal cancer caused by biallelic MUTYH pathogenic
variants using tumor mutational signatures. *Nat Commun* **13**, 3254, doi:10.1038/s41467-
022-30916-1 (2022).

60   Levatić, J., Salvadores, M., Fuster-Tormo, F. & Supek, F. Mutational signatures are
markers of drug sensitivity of cancer cells. *Nat Commun* **13**, 2926, doi:10.1038/s41467-
022-30582-3 (2022).

61   Georgeson, P. *et al.* Evaluating the utility of tumour mutational signatures for identifying
hereditary colorectal cancer and polyposis syndrome carriers. *Gut* **70**, 2138-2149,
doi:10.1136/gutjnl-2019-320462 (2021).

62   Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer. *Nature*
578, 94-101, doi:10.1038/s41586-020-1943-3 (2020).

63   Trucco, L. D. *et al.* Ultraviolet radiation-induced DNA damage is prognostic for outcome
in melanoma. *Nat Med* **25**, 221-224, doi:10.1038/s41591-018-0265-6 (2019).

64   Moreno, N. C. *et al.* Whole-exome sequencing reveals the impact of UVA light
mutagenesis in xeroderma pigmentosum variant human cells. *Nucleic Acids Res* **48**, 1941-
1953, doi:10.1093/nar/gkz1182 (2020).

65   Feigelson, H. S. *et al.* Melanoma incidence, recurrence, and mortality in an integrated
healthcare system: A retrospective cohort study. *Cancer Med* **8**, 4508-4516,
doi:10.1002/cam4.2252 (2019).

66   Ferlay, J. *et al.* Estimating the global cancer incidence and mortality in 2018: GLOBOCAN
sources and methods. *Int J Cancer* **144**, 1941-1953, doi:10.1002/ijc.31937 (2019).

978	67	Weir, H. K. *et al.* Melanoma in adolescents and young adults (ages 15-39 years): United
979		States, 1999-2006. *J Am Acad Dermatol* **65**, S38-49, doi:10.1016/j.jaad.2011.04.038
980		(2011).

67 Weir, H. K. *et al.* Melanoma in adolescents and young adults (ages 15-39 years): United
States, 1999-2006. *J Am Acad Dermatol* **65**, S38-49, doi:10.1016/j.jaad.2011.04.038
(2011).
68 Tomasetti, C., Marchionni, L., Nowak, M. A., Parmigiani, G. & Vogelstein, B. Only three
driver gene mutations are required for the development of lung and colorectal cancers. *Proc*
Natl Acad Sci U S A **112**, 118-123, doi:10.1073/pnas.1421839112 (2015).
69 Tomasetti, C., Li, L. & Vogelstein, B. Stem cell divisions, somatic mutations, cancer
etiology, and cancer prevention. *Science* **355**, 1330-1334, doi:10.1126/science.aaf9011
(2017).
70 Balmain, A. The critical roles of somatic mutations and environmental tumor-promoting
agents in cancer risk. *Nat Genet* **52**, 1139-1143, doi:10.1038/s41588-020-00727-5 (2020).
71 Zhang, M. *et al.* Use of tanning beds and incidence of skin cancer. *J Clin Oncol* **30**, 1588-
1593, doi:10.1200/JCO.2011.39.3652 (2012).
72 O'Sullivan, N. A. & Tait, C. P. Tanning bed and nail lamp use and the risk of cutaneous
malignancy: a review of the literature. *Australas J Dermatol* **55**, 99-106,
doi:10.1111/ajd.12145 (2014).
73 Consortium, I. T. P.-C. A. o. W. G. Pan-cancer analysis of whole genomes. *Nature* **578**,
82-93, doi:10.1038/s41586-020-1969-6 (2020).
74 Ellrott, K. *et al.* Scalable Open Science Approach for Mutation Calling of Tumor Exomes
Using Multiple Genomic Pipelines. *Cell Syst* **6**, 271-281.e277,
doi:10.1016/j.cels.2018.03.002 (2018).
75 David Benjamin *et al.* (bioRxiv 2019).
76 Koboldt, D. C., Larson, D. E. & Wilson, R. K. Using VarScan 2 for Germline Variant
Calling and Somatic Mutation Detection. *Curr Protoc Bioinformatics* **44**, 15.14.11-17,
doi:10.1002/0471250953.bi1504s44 (2013).
77 Kim, S. *et al.* Strelka2: fast and accurate calling of germline and somatic variants. *Nat*
Methods **15**, 591-594, doi:10.1038/s41592-018-0051-x (2018).
78 Fan, Y. *et al.* MuSE: accounting for tumor heterogeneity using a sample-specific error
model improves sensitivity and specificity in mutation calling from sequencing data.
Genome Biol **17**, 178, doi:10.1186/s13059-016-1029-6 (2016).
79 McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol* **17**, 122,
doi:10.1186/s13059-016-0974-4 (2016).
80 Bergstrom, E. N. *et al.* SigProfilerMatrixGenerator: a tool for visualizing and exploring
patterns of small mutational events. *BMC Genomics* **20**, 685, doi:10.1186/s12864-019-
6041-2 (2019).
81 Bergstrom, E. N., Barnes, M., Martincorena, I. & Alexandrov, L. B. Generating realistic
null hypothesis of cancer mutational landscapes using SigProfilerSimulator. *BMC*
Bioinformatics **21**, 438, doi:10.1186/s12859-020-03772-3 (2020).
82 Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer (IGV):
high-performance genomics data visualization and exploration. *Brief Bioinform* **14**, 178-
192, doi:10.1093/bib/bbs017 (2013).
83 Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847-2849,
doi:10.1093/bioinformatics/btw313 (2016).

84   Gu, Z., Gu, L., Eils, R., Schlesner, M. & Brors, B. circlize Implements and enhances
1023   circular visualization in R. *Bioinformatics* **30**, 2811-2812,
1024   doi:10.1093/bioinformatics/btu393 (2014).
1025   85   Günther, F., Dudschig, C. & Kaup, B. LSAfun--An R package for computations based on
1026   Latent Semantic Analysis. *Behav Res Methods* **47**, 930-944, doi:10.3758/s13428-014-
1027   0529-0 (2015).
1028

Assessment	Assays	Timepoint
Cytotoxicity	Cell Counting kit (CCK-8)	48h
	Trypan Blue	
Genotoxicity	γH2Ax immunofluorescence	4h
	CPD/6-4PP immunofluorescence	Immediately
Oxidative damage	CellROX green reagent	Immediately
	OxiSelect In vitro ROS/RNS assay kit	Immediately
		20min
		24h
	EpiQuik 8-OHdG DNA damage quantification kit	Immediately
		4h
		24h
Mitochondrial damage	MitoSOX mitochondrial superoxide indicator	24h
	TMRM mitochondrial membrane potential indicator	

Figure 1

Figure 3

Figure 4

a

MEF   MEF   HEF
       HEKa   HEKa
              HFF

Single Base Substitutions per Megabase (SBS/Mb)

Control (grey)
Acute (orange)
Chronic (red)

b

HEKa   MEF Acute   MEF Chronic
        HFF

C>A, C>G, C>T, T>A, T>C, T>G, All

Fold change to the control

HFF Acute   HFF Chronic

c

MEF

$R = 0.64$, $p = 0.0098$

C>A substitutions per Megabase

HFF

$R = 0.85$, $p = 5.9e{-}05$

C>A substitutions per Megabase

Number of UVA exposure

d

MEF   HFF

Control   Acute   Chronic

Co   Ch

INDELs
deletions
insertions

Figure 5

Figure 6

Figure 7