Systematic monitoring of photosynthetically-derived oxidative signals by a peroxiredoxin-based biosensor

Nardy Lampl[1], Idan Nissan[1], Raz Lev[1], Gal Gilad[1], Matanel Hipsch[1] and Shilo Rosenwasser[1]

Affiliations:

[1]The Robert H. Smith Institute of Plant Sciences and Genetics in Agriculture, The Hebrew University, Rehovot, Israel

 *Corresponding author. email: shilo.rosenwaser@mail.huji.ac.il (SR)

Abstract: Transmission of reductive cues from the photosynthetic electron transport chain to redox-regulated proteins plays a crucial role in activating chloroplast metabolism. However, deciphering the role of their counterbalanced oxidative signals is challenging due to monitoring difficulties. Here, we demonstrate the light-depended redox modification of chloroplast targeted 2-Cys peroxiredoxins and introduced a peroxiredoxin-based biosensor to monitor the dynamic changes in photosynthetically-derived oxidative signals. Systematic monitoring of probe oxidation demonstrated the induction of oxidative signals under habitual light intensities and their inverse relationship with chloroplastic NADPH levels, unraveling the combined activity of reducing and oxidizing signals in fine-tuning chloroplast activity. Considering that peroxiredoxins mediate oxidative inhibition of carbon assimilation, the measured light-dependent oxidative signals reflect the limitation to photosynthesis imposed by the redox regulatory system.

One-Sentence Summary: A genetically encoded biosensor unmasked the dominant role of photosynthetically-derived oxidative signals under habitual conditions.

Introduction

Sophisticated mechanisms enabling adjustments of light-capturing reactions and downstream metabolic processes have evolved in sessile plants, allowing them to reach optimal photosynthetic performance. Under low-light conditions, efficient photosynthetic activity and tight regulation of the reducing power distribution between essential metabolic pathways are required for optimal utilization of the available energy input. On the other hand, energy dissipation under high-light (HL) conditions is critical to avoid over-production of harmful reactive oxygen species (ROS) (Asada, 2006). Reductive and oxidative signals, both emanating from the photosynthetic electron transport chain (PETC), and their transfer to regulated thiol proteins, play a significant role in linking the performance of photosynthesis to chloroplast transcription, translation, and metabolic processes, thereby enabling rapid acclimation to instantaneous changes in daytime photon fluxes (Buchanan and Balmer, 2005; Schürmann and Buchanan, 2008b; Foyer and Noctor, 2009; Dietz and Pfannschmidt, 2011; Scheibe and Dietz, 2012).

Reductive signals in chloroplasts are mainly generated by the flux of electrons derived from ferredoxin to target proteins, via ferredoxin-dependent thioredoxin reductase (FTR), to thioredoxins (Trxs) (Schürmann and Buchanan, 2008) and via NADPH-dependent thioredoxin reductases C (NTRC) (Serrato et al., 2004), thereby linking chloroplast metabolism to light availability and photosynthetic activity. Counteracting oxidative signals are mainly generated from electron flux through the water-water cycle (WWC). During the WWC, electrons are donated from photosystem I (PSI) to molecular oxygen in the Mehler reaction, yielding superoxide radicals ($O_2^{.-}$), which are then dismutated to molecular oxygen (O_2) and H_2O_2 in a reaction catalyzed by superoxide dismutase (SOD) (Mehler, 1951; Rizhsky et al., 2003; Mittler et al., 2004). The production of H_2O_2 through the WWC allows communication between the photosynthetic light reactions and downstream metabolic processes by transmitting oxidative signals to redox-regulated proteins (Apel and Hirt, 2004; Mullineaux et al., 2018; Exposito-Rodriguez et al., 2017; Smirnoff and Arnaud, 2019).

The detoxification of photosynthetically produced H_2O_2 is mediated by chloroplast-targeted ascorbate peroxidases (APXs) and 2-Cys peroxiredoxins (2-Cys Prxs) (Awad et al., 2015; Dietz, 2016). The latter are highly abundant and remarkably efficient thiol peroxidases, displaying high catalytic efficiency of up to 10^8 M^{-2} s^{-1} (König et al., 2002; Dietz et al., 2006). Electrons delivered from the PETC, reduce 2-Cys Prx via the FTR/TRX pathways or through NADPH via NTRC activity (Collin et al., 2003; Pérez-Ruiz and Cejudo, 2009; Pulido et al., 2010; Cerveau et al., 2016). Therefore, the 2-Cys Prxs redox state is determined by the balance between photosynthetically produced reducing and oxidizing equivalents. Importantly, recent studies have demonstrated oxidative signal transmission, mediated by 2-Cys Prxs, from H_2O_2 to target proteins, at the onset of the light and dark periods and under low- and fluctuating-light conditions (Dangoor et al., 2012; Eliyahu et al., 2015; Ojeda et al., 2018; Vaseghi et al., 2018; Yoshida et al., 2018). Despite the considerable importance of the regulatory role of 2-Cys Prx, most of our current knowledge is based on measuring its oxidation state using western blot analysis or examination of mutant lines performance, rendering it challenging to detect the transmission of oxidative signals in real-time and with high temporal resolution.

Genetically encoded redox-sensitive green fluorescent protein (roGFP) probes enable, due to their high sensitivity and reversibility, systematic *in vivo* mapping of H_2O_2 production and the glutathione redox state (E_{GSH}), in different sub-cellular compartments (Dooley et al., 2004; Hanson et al., 2004; Jiang et al., 2006; Meyer et al., 2007; Gutscher et al., 2008; Gutscher et al.,

2009; Meyer and Dick, 2010; Nietzel et al., 2019; Müller-Schüssele et al., 2021). In vitro characterization of roGFP showed that its reduction is mediated by Glutaredoxins (Grxs), which catalyze the reversible electron flow between GSH and target proteins (Meyer et al., 2007; Gutscher et al., 2008). Recently, an H_2O_2 sensor based on a redox relay mechanism between Tsa2, a highly efficient *S. cerevisiae* 2-Cys peroxiredoxin, and roGFP2 was developed (Morgan et al., 2016). To increase the probe's sensitivity to low H_2O_2 levels, a mutation in the resolving cysteine (C_R) of Tsa2 (roGFP2-Tsa2ΔC_R) was introduced, thus preventing competition between the Trx-mediated reduction of the Tsa2 moiety and roGFP oxidation. The resulting probe enabled the measurement of endogenous basal H_2O_2 levels in yeast and *Chlamydomonas reinhardtii* cells (Morgan et al., 2016; Niemeyer et al., 2020); however, such a roGFP2 Prx☐based probe has not been yet implemented to measure the dynamic oxidation of 2-Cys Prx in higher plants.

Here, we employed chl-roGFP2- PrxΔC_R, a new 2-Cys Prx biosensor, to establish the role of 2-Cys Prx in light-dependent oxidative signal transmission in *Arabidopsis* plants. By parallel measurements of the endogenous 2-Cys Prx and chl-roGFP2-PrxΔC_R oxidation states, we showed that 2-Cys Prx is directly regulated by light intensity. Furthermore, the dynamics of 2-Cys Prx-dependent oxidative signals, throughout the day, under various light conditions, were resolved by systematic monitoring of the probe's redox state and showed negative correlation with the availability of chloroplast NADPH. Determination of the light intensity range of 2-Cys Prx-mediated redox signaling suggests its role in redox signaling under habitual conditions rather than oxidative stress.

Results

Recent studies have demonstrated oxidative signal transmission mediated by the highly abundant chloroplast 2-Cys peroxiredoxins (2-Cys Prxs), a thoroughly studied scavenger of WWC-derived H_2O_2. Thus, we sought to explore the direct effect of light intensity on the redox state of 2-Cys Prx. In agreement with a previous report (Dangoor et al., 2012), transferring plants from dark to normal growth light (GL, 120 µmol photons m^{-2} s^{-1}) at the beginning of daylight, resulted in a reduction of 2-Cys Prx catalytic sites, as reflected by the dominance of the dimer (oxidized) band under dark (at the end of the night) and by the prolonged decrease in its relative abundance under light conditions (Fig. 1A). Importantly, this reduction was not observed in plants that were pretreated with 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU), an inhibitor of photosynthetic electron transport (Fig. 1A), simulating dark conditions. To distinguish between the fully oxidized (2S-S) and half-oxidized (1S-S) forms, a gel-shift assay was carried out with methoxypolyethylene glycol-maleimide (mPEG). A mixture of three bands was identified, with the higher and middle bands trapped with mPEG and corresponding to the 2S-S and 1S-S forms, respectively (Fig. 1B&C). Under dark conditions, a relatively high level of the fully oxidized form was visible. On transfer of plants from dark to GL, the relative levels of the fully reduced form significantly increased and were paralleled by a decrease in the fully oxidized form. Such a reduction was not detected in plants treated with DCMU under GL, indicating that PETC was the source of the reduction power.

Plant exposure to high light (HL) intensities (600 or 1000 µmol photons m^{-2} s^{-1}) (Fig. 1A) resulted in a similar reduction of 2-Cys Prx upon dark-to-light transition for the two tested HL intensities, which stabilized at a higher oxidation state as compared with GL. A significant amount of fully oxidized 2-Cys Prx arose in plants exposed to 1000 µmol photons m^{-2} s^{-1} (Fig.

1B&C), suggesting that increased illumination enhanced 2-Cys Prx dimer formation, in which both catalytic sites participate in disulfide exchange reactions. Plant exposure to constant HL conditions or DCMU also led to higher levels of hyperoxidized 2-Cys Prx than GL, as measured using SO_3 antibodies (fig. S1).

Fig. 1 Light responsiveness of the 2-Cys Prx redox state. Plants were kept in the dark for 8 h and then exposed for 5.5 h to the indicated light intensities. Total protein was extracted using trichloroacetic acid (TCA), and free thiols were blocked with N-ethylmaleimide (NEM). (**A**) An immunoblot, using anti-Prx, showing the oxidized and reduced states of 2-Cys

5   Prx, as dimers and monomers, respectively. (**B**) Schematic illustration of a redox mobility gel assay (with samples from A). (**C**) Gel-shift assay with methoxypolyethylene glycol-maleimide (mPEG) showing the in vivo redox state of 2-Cys Prx

residues in plants exposed to light intensities as in A. Following NEM treatment, 1,4-Dithiothreitol (DTT) was applied to reduce disulfides before mPEG labeling. The number of oxidized Cys residues (Ox1 and Ox2) was determined by the increased molecular mass of 2-Cys Prx after labeling with mPEG. (**D**) Light intensity over time. (**E**) Plants were exposed to the light regime indicated in D. Immunoblot assay showing the oxidized and reduced states of 2-Cys Prx, as dimers and monomers, respectively (anti-Prx) and 2-Cys Prx hyperoxidation (anti-SO$_3$).

An apparent light-dependent oxidation response of 2-Cys Prx was observed under prolonged, physiologically relevant light conditions (Fig. 1 D&E). Interestingly, the major shift in oxidation was during the non-stressed light regime, in the transition from low (80-130 µmol photons m^{-2} s^{-1}) to moderate (240-280 µmol photons m^{-2} s^{-1}) light intensities, at the initiation and termination of the light phase (Fig.1 D&E), implying its signaling role under habitual light conditions. Notably, accumulation of the Prx hyperoxidation form was not detected (Fig. 1E), suggesting that a gradual increase in light conditions inhibits 2-Cys Prx overoxidation even while reaching HL conditions. These observations demonstrated that the 2-Cys Prx redox state is directly affected by the PETC and that its oxidation state under gradual changes in light intensities that mimic natural conditions in the field, is firmly linked to changes in light intensities.

We recognized that the observed oxidation patterns of 2-Cys Prx in response to changing light intensities reflect the balance between its peroxidase activity and its reduction by TRXs. Therefore, to explicitly inspect the oxidative signals mediated by 2-Cys Prx activity, we generated, based on a previous work in yeast (Morgan et al., 2016), a chl-roGFP2-PrxΔC$_R$ probe by genetically fusing roGFP2 to a mutated version of 2-Cys Prx A (BAS1), in which the resolving cysteine (C$_R$) was changed to alanine, impeding its TRX-dependent reduction (Fig. 2A-C). BAS1 was fused at its N terminus to roGFP2, since initial attempts to fuse it at its C terminus resulted in probe cleavage (fig. S2). Interestingly, the cleaved roGFP obtained from the fusion of BAS1 at its C-terminus was localized in the chloroplast (fig. S2), implying that the C terminus of BAS1 is prone to cleavage by specific chloroplast proteases. A dynamic range of 5.6 was calculated between the fully oxidized (R$_{oxi}$=0.45) and fully reduced (R$_{red}$=0.08) forms, ensuring a sufficient signal to noise ratio (Fig. 2 D). Whole-plant ratiometric analysis (Hipsch et al., 2021) showed a higher oxidation state of the chl-roGFP2-PrxΔC$_R$ probe compared to chl-roGFP2 under steady-state conditions (GL) (~ 60% and ~ 40%, respectively, Fig. 2 E-G), a phenomenon also observed with the roGFP2-Tsa2ΔC$_R$ probe (Morgan et al., 2016; Niemeyer et al., 2020), indicating a direct effect of 2-Cys Prx on the roGFP moiety. Shifting plants from dark to GL induced a slight change in the redox state of chl-roGFP2-PrxΔC$_R$ and significant changes were visualized in its oxidation degree (OxD) upon shifting plants from dark to different HL intensities (450, 750, or 1000 µmol photons m^{-2} s^{-1}), reaching its almost fully oxidized state within 5 min of transition to light (Fig. 2 E&G). In contrast, the unfused chl-roGFP2 probe was only moderately oxidized under HL conditions (Fig. 2 E&F). These findings demonstrate the enhanced light sensitivity of chl-roGFP2-PrxΔC$_R$ compared to chl-roGFP2, achieved by positioning BAS1 in close proximity to roGFP2. As both probes are subjected to reduction in a GSH/GRX-dependent manner, the comparison between their dynamic oxidation enables resolution of the effect on H$_2$O$_2$-dependent thiol oxidases mediated by BAS1.

Fig. 2 Light responsiveness of chloroplast-targeted peroxiredoxin-based redox probes. (**A**) Schematic diagram of the gene cassette used to transform *Arabidopsis* plants. (**B**) Subcellular localization of chl-roGFP2-PrxΔC_R in mesophyll cells, as detected by confocal microscopy. (**C**) Detection of the fused sensor protein by western blot analysis using either anti-Prx or anti-GFP antibodies. (**D**) Fluorescence ratios (405/465) in plants expressing chl-roGFP2-PrxΔC_R under steady-state conditions and following treatments with 100 mM DTT or 500 mM H_2O_2. Fluorescence emitted from three-week-old plants grown in soil was recorded using a plate reader. (**E**) Whole-plant ratiometric analysis of chl-roGFP2 and chl-

roGFP2-PrxΔC$_R$ fluorescence under dark and 5 min following transfer from dark to different light intensities. Ratiometric analysis of plants treated with 100 mM DTT or 500 mM H$_2$O$_2$ is also presented. The roGFP2 fluorescence images were captured at 510 nm, following excitation at 400 nm and 465 nm. (**F, G**) Quantification of ratiometric images, presented as box-plots.

We further investigated the daily dynamics of the chl-roGFP2-PrxΔC$_R$ oxidation state in comparison to chl-roGFP2 using an automated system that allows for continuous measurements of fluorescence signals emanating from living plants expressing genetically encoded biosensors (Fig. 3A-C, Methods). Monitoring probe oxidation throughout the day further demonstrated the higher sensitivity of chl-roGFP2-PrxΔC$_R$ to light compared to chl-roGFP2. While both probes were oxidized upon transition from dark to light under all examined light conditions, significantly higher oxidation values were recorded under HL in chl-roGFP2-PrxΔC$_R$ (~85-95%) as compared to chl-roGFP2 plants (~65-70 %; Fig. 3B-C).

To characterize chl-roGFP2-PrxΔC$_R$ oxidation dynamics under a range of light intensities, we monitored their OxD in plants exposed to gradual changes in light intensities that mimic natural conditions, as described in Fig.1E. For comparison, the oxidation dynamics of the chl-roGFP2 and chl-roGFP2-Prx probes (without the mutation in the C$_R$, fig. S4), were also monitored, as was NADPH availability, using the recently developed *Arabidopsis* NADPH sensor (TKTP-iNAP4) (Tao et al., 2017; Lim et al., 2020). As shown in Fig. 3D, the four probes exhibited entirely different oxidation patterns. The chl-roGFP2-PrxΔC$_R$ OxD sharply increased from the light onset, reaching its full oxidation state (~100%) after approximately two hours, when the light intensity was ∼300 µmol photons m^{-2} s^{-1}. A gradual reduction in OxD was only detected 14 h after light onset, when light intensities were dimmed again. A lower magnitude of oxidation of chl-roGFP2-Prx when compared to chl-roGFP2-PrxΔC$_R$ was recorded, supporting the higher sensitivity of the latter to oxidative signals, achieved by the C$_R$ mutation. Notably, the oxidative patterns of chl-roGFP2-PrxΔC$_R$ and chl-roGFP2-Prx probes in response to changes in light intensities were similar to those of endogenous 2-Cys Prx (Fig. 3D and Fig.1E), reinforcing the probe's capabilities to detect 2-Cys Prx-mediated redox signals. We reasoned that the differences between roGFP2-PrxΔC$_R$ and chl-roGFP2-Prx OxD reflected reducing equivalents transferred to 2-Cys Prx by TRXs or NTRC activity. In contrast to the oxidation patterns of chl-roGFP2-Prx and chl-roGFP2-PrxΔC$_R$, the unfused chl-roGFP2 showed a gradual increase in OxD, reaching its highest level (65%) after 3 h, when the light intensity was ∼600 µmol photons m^{-2} s^{-1}. From this point onward, a gradual decrease in chl-roGFP2 oxidation was observed, reaching its lowest OxD (42%) at the beginning of the dark period.

Strikingly, the redox state of chl-roGFP2-PrxΔC$_R$ negatively correlated with the dynamic changes in stromal NADPH availability. While the NADPH level was high and stable at night, a sharp decrease was detected under light, starting after light onset and reaching its lowest level at ~300 µmol photons m^{-2} s^{-1}. A gradual increase in NADPH level was observed ~14 h after light onset, when light intensities were lowered, until it reached its highest level at the beginning of the night. These results suggest that the availability of stromal NADPH governs the induction of photosynthetically-derived oxidative signals.

Fig. 3 Systematic monitoring of light-induced oxidative signals. (**A**) Schematic illustration of an automated system used to monitor fluorescence signals in living plants. (**B, C**) Dynamic changes in chl-roGFP2 and chl-roGFP2-PrxΔC_R oxidation degree (OxD) in plants exposed to growth light (GL) (120 µmol photons m^{-2} s^{-1}) or high light (HL) (600 or 1000 µmol photons m m^{-2} s^{-1}) after 8 h in the dark. (**D**) Dynamic changes in OxD of chl-roGFP2-PrxΔC_R, chl-roGFP2-Prx and chl-roGFP2 as well as TKTP-iNAP4 ratio (400/485nm) throughout the day, are presented. The applied light intensities are depicted as a dashed yellow line. For each line, between 24 and 40 plants divided into 3-5 independent plates were analyzed and consolidated in a "sliding window" (n=3-5) display. Values represent means ± SE. The color bar denotes the light conditions: black-dark, yellow-light.

The increasing recognition that oxidative signals play a dominant role in fine-tuning of metabolic activity under low-light conditions (Dangoor et al., 2012; Eliyahu et al., 2015; Vaseghi et al., 2018; Wolf et al., 2020) drove us to further investigate the sensitivity of the chl-roGFP2-PrxΔC$_R$ redox state to low-light intensity. To this end, plants were exposed to a gradual increase in light intensities, starting with ~22 µmol photons m^{-2} s^{-1} and increasing up to 330µmol photons m–2 s (Fig. 4A). An immediate reduction of chl-roGFP2 was observed at ~22 µmol photons m^{-2} s^{-1}, followed by oxidation at a light intensity of ~44 µmol photons m^{-2} s^{-1}, with oxidation reaching a maximum under 330 µmol photons m^{-2} s^{-1}. In contrast, gradual oxidation of chl-roGFP-PrxΔC$_R$ was observed with progression through the low-light gradient. Stromal NADPH availability negatively correlated with chl-roGFP2-PrxΔC$_R$ as in Fig. 3D. Interestingly, the chl-roGFP2-PrxΔC$_R$ and chl-roGFP2-Prx probes showed contrasting responses upon transition from dark to low light (Fig. 4B). While chl-roGFP2-PrxΔC$_R$ OxD slightly increased when plants were transferred to low light, chl-roGFP2-Prx underwent a significant reduction. This reduction continued for at least two hours from light onset until stabilization at a lower OxD (35%). The differences between the OxD of roGFP2-PrxΔC$_R$, chl-roGFP2-Prx, and the unfused chl-roGFP2 after the onset of low light demonstrate the high magnitude of reductive signals under low-light intensities, emanating from TRXs and GSH/GRXs activity and which are attenuated by induced oxidative signals (Fig. 4C).

Discussion

Tracking the in vivo dynamics of oxidative thiol modulations is central to understating the role of redox regulation in adjusting metabolic activity. Exploring the diurnal redox dynamics of the CBC enzymes, FBPase and SBPase, showed a change from fully oxidized to an almost fully reduced state of FBPase during gradual increases in light intensity from dark to 350 µmol photons m^{-2} s^{-1} (Yoshida et al., 2014). Intriguingly, the observed light range in which FBPase reduction occurred matched the range of light-induced changes in 2-Cys redox signaling, as observed either by direct measurements of the 2-Cys Prx redox state or by using the roGFP2-PrxΔC$_R$ (Fig. 1E&3D&4A). Considering the sensitivity of FBPase to 2-Cys Prx-mediated oxidation (Yoshida et al., 2018), these results suggest that the reductive and oxidative pathways are interconnected in regulating carbon assimilation rates under low- and moderate-light conditions. Furthermore, the inverse relationship between NADPH level and the induction of oxidative signals suggests a feedback mechanism in which NADPH consumption may trigger oxidative signals, leading to inhibition of CBC enzyme activities and to a consequent decrease in carbon assimilation rates.

Due to the heterogeneous organization and spatial arrangement of crop canopies, leaves lower in the canopy experience sporadic HL, scattered with low-light levels of variable durations. Given the dominant role of redox signaling at low to moderate light intensities, the level of activation and deactivation of CBC enzymes in lower leaves will vary throughout the day, with both the reductive and oxidizing signal shaping the rate of carbon assimilation rate, defining the contribution of these leaves to the total carbon assimilation rates (Sassenrath-Cole and Pearcy, 1994). The inherent redox changes that stimulate the activation and deactivation of CBC enzymes link carbon fixation to electron transport activity and energy flux but also govern limitation to optimal canopy assimilation under fluctuating light intensities. Given the role of 2-Cys Prx in deactivating CBC enzyme activity, our data suggest that the roGFP2-PrxΔC$_R$ probe estimates the degree of limitation of photosynthesis imposed by oxidative signals in plants, laying the foundation for increasing photosynthesis efficiency through redox modulations.

Fig. 4 chl-roGFP2-PrxΔC$_R$ reports on the induction of oxidative signals under low light. (**A**) Daily dynamic changes in the oxidation degree (OxD) of chl-roGFP2-PrxΔC$_R$ and chl-roGFP2, and in TKTP-iNAP4 ratios, in plants exposed to a gradual increase (low to moderate) light intensities followed by an equivalent decrease, are presented. Monitoring was performed in 3-week-old plants grown in 12-well plates. For each line, between 24 and 40 plants divided into 3-5 independent plates were analyzed and consolidated in a "sliding window" (n=3-5) display. Values represent means ± SE. The color bar denotes the light conditions: black-dark, yellow-light. (**B**) Dynamic changes in chl-roGFP2-PrxΔC$_R$, chl-roGFP2-Prx and chl-roGFP2 OxD at the onset of low light (22 µmol photons m^{-2} s^{-1}). The applied light intensities are depicted as a dashed yellow line. (**C**) A proposed model of the oxidation and reduction powers that shape the oxidation state of the chl-roGFP2-PrxΔC$_R$, chl-roGFP2-Prx and chl-roGFP2 sensors.

Materials and Methods

Plant growth conditions and treatment

Arabidopsis thaliana ecotype Col-0 plants were grown in a greenhouse under a 16/8 h light/darkness cycle. Immunoassay, redox imaging and thiol labelling experiments were performed on whole, three-week-old *Arabidopsis* plants. For the 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) treatment, three-week-old plants were sprayed with 150μM DCMU (D2425-100G, Sigma) in the darkness phase, 1 hour before the onset of the light period.

Construction of roGFP2-PrxΔCR and PrxΔCR-roGFP2 probes

Both fusion probes (roGFP2-PrxΔC_R and PrxΔC_R-roGFP2) were synthesized with codons optimized for *Arabidopsis* expression (Genewiz). Chloroplast targeting was achieved by using the 2-Cys peroxiredoxin A signal peptide. The genetically fused sequences were cloned into the plant cloning vector pART7, using the XhoI and HindIII restriction enzymes. The whole construct, including the CaMV 35S promoter and ocs terminator, was then cloned into the binary vector pART27, using the NotI restriction enzyme. The pART27 plasmids containing the roGFP2-PrxΔC_R and PrxΔC_R-roGFP2 probe constructs, were transformed into GV3101 Agrobacterium tumefaciens. Transformation of *Arabidopsis* thaliana (Columbia, Col-0) was then performed by floral dip (Clough and Bent, 1998). Transformant lines were selected based on kanamycin resistance and the chl-roGFP2 fluorescence signal.

Screening of chl-roGFP2-Prx and chl-roGFP2-PrxΔC_R DNA insertion in transformed plants

The substitution of the resolving cysteine (C_R) by alanine in chl-roGFP2-PrxΔC_R plants was verified by DNA insertion screening. DNA extraction and PCR amplification were carried out using REDExtract-N-AmpTM Plant PCR kit (sigma-aldrich). The sequence of both primers, forward and reverse are as follows:

Forward primer: CATCCAAGAGAACCCAGATGAAGCT

Reverse primer: TTGCAGAGAAATATTCTTTAGACAAC

The Forward primer was designed to form HindIII restriction site sequence in the PCR products of Prx thus allowing us to distinguish between Prx and PrxΔC_R.

PCR product for Prx (80bp):

CATCCAAGAGAACCCAGATGAAGCTTGCCCAGCTGGTTGGAAACCCGGTGAGAAGT
CTATGAAGCCCGATCCTAAGTTGTCTAAAGAATATTTCTCTGCAA

PCR product for PrxΔC_R (101bp):

CATCCAAGAGAACCCAGATGAAGCTGCTCCAGCTGGTTGGAAACCCGGTGAGAAGT
CTATGAAGCCCGATCCTAAGTTGTCTAAAGAATATTTCTCTGCAA

The PCR products were digested with HindIII and results in 80bp product for Prx and 101bp for PrxΔC_R.

Determination of 2-Cys Prx redox state by immunolabeling

NEM labeling.

Protein extracts were prepared in the presence of 1 ml cold 10% trichloroacetic acid (TCA) (dissolved in water) to retain the in vivo thiol oxidation status of proteins. Proteins were precipitated for 30 min on ice, in the darkness, and centrifuged at 19,000 xg for 20 min, at 4°C. The pellet was then washed four times with 100% cold acetone. After removal of the residual acetone, the pellet was resuspended in urea buffer (8M urea, 100mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (pH 7.2), 1mM EDTA, 2% (w/v) SDS, protease inhibitors cocktail (PI) (Calbiochem) and 100mM N-ethylmaleimide (NEM) (E3876, Sigma)) dissolved in ethanol. Samples were then incubated for 30 min, at room temperature, followed by centrifugation (19,000 xg, 20 min, 4°C) and washed (4 times) with 100% cold acetone. The dry pellets were resuspended in urea buffer without NEM. The sample buffer (x3) contained 150mM Tris-HCl, pH 6.8, 6% (w/v) SDS, 30% glycerol and 0.3% pyronin Y. Gel fractionation was carried out without a reducing agent, on a precast 4–15% polyacrylamide gel. Fractionated proteins were transferred to a polyvinylidene fluoride membrane (Bio-Rad), using the Trans-Blot Turbo Transfer System (Bio-Rad) with Trans-Blot Turbo Midi Transfer Packs. The membrane was incubated with anti-PRX antibodies (1:1,000) (kindly provided by Prof. Avichai Danon) or an anti-peroxiredoxin-SO$_3$ antibody (Abcam, ab16830, 1:2000), both followed by anti-rabbit horseradish peroxidase (HRP)-conjugated IgG (1:20,000) (Agrisera). Chemiluminescence was detected using the Advanced Molecular Imager HT (Spectral Ami-HT, Spectral Instruments Imaging, LLC., USA).

mPEG labeling

Protein extract was prepared, using a hand homogenizer, in 1ml cold 10% trichloroacetic acid (TCA) (dissolved in water) to retain the in vivo thiol oxidation status of proteins. Proteins were precipitated for 30 min, on ice, in the darkness, and then centrifuged at 19,000 xg for 20 min at 4°C. The pellet was washed four times with 100% cold acetone, after which, residual acetone was removed and the pellet was resuspended in urea buffer with NEM. The samples were then incubated for 30 min at room temperature. Samples were then reduced by addition of 100mM DTT (60 min, room temperature). TCA (10%) was then added and samples were precipitated for 30 min, on ice, in the darkness. The TCA-treated extracts were then centrifuged at 19,000 xg, 4°C for 20 min, supernatants were removed and the pellets were washed three times with 100% acetone. The dry pellets were resuspended in urea buffer containing 10mM methoxy-polyethylene glycol maleimide (mPEG, MW 5000 g/mol, Sigma), and incubated at 28 °C for 2 h. The reaction was stopped by adding an equal volume of sample buffer containing 50 mM DTT. Samples were then separated by 10% Tricine–SDS–PAGE and then transferred to a polyvinylidene fluoride membrane (Bio-Rad) under the same conditions described above. The membrane was incubated with anti-PRX antibodies (1:1,000) and then with anti-rabbit HRP-conjugated IgG (1:20,000).

Probe detection using western analysis

Experiments were performed on 10□day□old seedlings. Whole-plant contents were extracted with extraction buffer (20 mM Tris, pH 8.0, 1 mM EDTA, and 50 mM NaCl), and then centrifuged at 19,000 xg, for 20 min, at 4°C. Gel fractionation of 100 μg extracted protein was carried out by reducing 10% SDS□PAGE. Fractionated proteins were then transferred onto a polyvinylidene fluoride membrane, which was then incubated with anti-PRX antibodies

(1:1,000) or anti-GFP (abcam) antibodies (1:5,000), and then with anti-rabbit HRP-conjugated IgG (1:20,000).

Confocal microscopy

Confocal microscopy analysis was carried out on 2□week□old transgenic roGFP2, roGFP2-Prx, roGFP2-PrxΔC_R and PrxΔC_R-roGFP2-expressing plants. Images were captured with a Leica TCS SP8 confocal system (Leica Microsystems) and the LAS X Life Science Software, while using a HC PL APO ×40/1.10 objective. To capture chl-roGFP2 fluorescence, samples were excited at 488 nm and emission was measured at 500–520 nm. For chlorophyll fluorescence, excitation was at 488 nm and emission was at 670 nm. Merged images were generated using Fiji (Image 1.A) software.

roGFP2 fluorescence measurements and analysis

Redox imaging was performed on three-week-old transgenic chl-roGFP2 and roGFP2-PrxΔC_R-expressing plants. Whole-plant roGFP2 fluorescence was detected using an Advanced Molecular Imager HT, and images were taken using the AMIview software. For roGFP fluorescence images, excitation was at 465nm (reduction) and 405nm (oxidation), followed by emission at 510nm. Chlorophyll fluorescence, which was used to select plant tissue pixels, was measured following excitation at 405 and emission at 670. For each probe, images were taken under the same settings of LED intensities and exposure time. Ratiometric images were created by dividing, pixel by pixel, the 405 nm image by the 465 nm image, and displaying in false colors. Images were processed using a custom-written Matlab script. For calibration of the probe response, detached plants were immersed in 0.5M H_2O_2 or 100 mM DTT, and ratiometric images for fully oxidized and fully reduced states, respectively, were obtained.

Daily roGFP2 fluorescence was measured using a Tecan Spark® multimode microplate reader (Haber et al., 2021). For both probes (roGFP2, roGFP2-Prx and roGFP2-PrxΔC_R), roGFP fluorescence was measured using excitation at 485nm/20 (reduction) and 400nm/20 (oxidation), followed by emission at 520nm/10; gain values were adjusted to avoid signal saturation. For chlorophyll detection, 400nm/20 was applied for excitation, followed by 670nm/40 for emission. For automatic detection of chl-roGFP2 signals, 10-day-old plants expressing the roGFP2-based probes or wild type plants were transferred to solid peat plugs in 12-well plates. The plates were placed in a growth chamber (Fytoscope FS-SI-4600, Photon Systems Instruments, Czech Republic) under normal growth conditions for another 5 days before the start of the experiment. An automated robot (KiNEDx KX.01467, paa automations, Hampshire, UK) controlled by a self-compiled program (Overlord™, paa automations), inserted the plates into the plate reader to measure chl-roGFP2 fluorescence. For the plate-reader analysis, a 9-by-9-pixel matrix was formed for each well. Chlorophyll fluorescence was detected to create a chlorophyll mask, which was then used to choose pixels that returned a positive chlorophyll fluorescence signal, and only those pixels were considered for the roGFP analysis. The average signals of non-fluorescent plants (WT) was calculated, and the values were deducted from the values detected in the chl-roGFP2 fluorescence analysis. A similar experimental setup was used to measure the fluorescence emitted from the TKTP-iNAP4 and TKTP-iNAPc expressing lines. The pH corrected ratio for TKTP-iNAP4 was calculated by dividing the fluorescence ratio values obtained from the TKTP-iNAP4 line by those obtained from the TKTP-iNAPc line, according to Lim et al., (2020) . To determine sensor OxD, detached roGFP-based probe-expressing and WT

plants were immersed in 0.5M H_2O_2 or 100 mM DTT and fluorescence was measured. roGFP2 OxD was calculated based on the fluorescence signal according to Equation (Meyer et al., 2007).

Equation 1:

$$xD_{roGFP2} = \frac{R - R_{red}}{(I_{485min}/I_{485max})(R_{ox} - R) + (R - R_{red})}$$

where R represents the ratio (400/485) at each point in the experiment, Rred represents the ratio under fully reduced conditions, Rox represents the ratio under fully oxidized conditions, I485min represents the fluorescence emitted at 520 nm when excited at 485 nm under fully oxidized conditions and I485max represents the fluorescence emitted at 520 nm when excited at 485 nm under fully reduced conditions.

A self-written Matlab script was used to analyze the many files that contained the measurements of each experiment. In addition, experimental metadata such as plant lines, light irradiance, temperature, relative humidity and CO_2 concentration were also collected and added to the output file.

Author contributions: N.L. and S.R. conceived and designed the research. N.L., I.N., R.L., G.G., M.H., and S.R. performed the research. N.L., R.L., M.H., and S.R. analyzed the data. N.L., and S.R. wrote the manuscript. Competing interests: The authors declare no competing interests.

Acknowledgments: We thank Avihai Danon for his critical comments on the manuscript and for kindly providing us the anti-2-Cys Prx antibody. We thank Bruce Morgan for his kindly help in initiating this project.

Funding: This research was supported by the Israel Science Foundation (grant No.826/17 and No. 827/17).

References

Apel K, Hirt H (2004) Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. Annu Rev Plant Biol **55**: 373–399

Asada K (2006) Production and scavenging of reactive oxygen species in chloroplasts and their functions. Plant Physiol **141**: 391 LP – 396

Awad J, Stotz HU, Fekete A, Krischke M, Engert C, Havaux M, Berger S, Mueller MJ (2015) 2-Cysteine peroxiredoxins and thylakoid ascorbate peroxidase create a water-water cycle that is essential to protect the photosynthetic apparatus under high light stress conditions. Plant Physiol **167**: 1592 LP – 1603

Buchanan BB, Balmer Y (2005) REDOX REGULATION: A Broadening horizon. Annu Rev Plant Biol **56**: 187–220

Cerveau D, Kraut A, Stotz HU, Mueller MJ, Cout Y, Rey P (2016) Characterization of the

Arabidopsis thaliana 2-Cys peroxiredoxin interactome. Plant Sci **252**: 30–41

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium -mediated transformation of Arabidopsis thaliana. Plant J **16**: 735–743

Collin V, Issakidis-Bourguet E, Marchand C, Hirasawa M, Lancelin J-M, Knaff DB, Miginiac-Maslow M (2003) The arabidopsis plastidial thioredoxins: new functions and new insights into specificity . J Biol Chem **278**: 23747–23752

Dangoor I, Peled-Zehavi H, Wittenberg G, Danon A (2012) A chloroplast light-regulated oxidative sensor for moderate light intensity in Arabidopsis. Plant Cell **24**: 1894–906

Dietz K-J (2016) Thiol-based peroxidases and ascorbate peroxidases: why plants rely on multiple peroxidase systems in the photosynthesizing chloroplast? Mol Cells **39**: 20–5

Dietz K-J, Jacob S, Oelze M-L, Laxa M, Tognetti V, de Miranda SMN, Baier M, Finkemeier I (2006) The function of peroxiredoxins in plant organelle redox metabolism. J Exp Bot **57**: 1697–1709

Dietz K-J, Pfannschmidt T (2011) Novel regulators in photosynthetic redox control of plant metabolism and gene expression. Plant Physiol **155**: 1477 LP – 1485

Dooley CT, Dore TM, Hanson GT, Jackson WC, Remington SJ, Tsien RY (2004) Imaging dynamic redox changes in mammalian cells with green fluorescent protein indicators. J Biol Chem **279**: 22284–22293

Exposito-Rodriguez M, Laissue PP, Yvon-Durocher G, Smirnoff N, Mullineaux PM (2017) Photosynthesis-dependent H_2O_2 transfer from chloroplasts to nuclei provides a high-light signalling mechanism. Nat Commun 8: 49

Eliyahu E, Rog I, Inbal D, Danon A (2015) ACHT4-driven oxidation of APS1 attenuates starch synthesis under low light intensity in *Arabidopsis* plants. Proc Natl Acad Sci **112**: 12876 LP – 12881

Foyer CH, Noctor G (2009) Redox regulation in photosynthetic organisms. Regulation **11**: 861–905

Gutscher M, Pauleau A-L, Marty L, Brach T, Wabnitz GH, Samstag Y, Meyer AJ, Dick TP (2008) Real-time imaging of the intracellular glutathione redox potential. Nat Methods **5**: 553–559

Gutscher M, Sobotta MC, Wabnitz GH, Ballikaya S, Meyer AJ, Samstag Y, Dick TP (2009) Proximity-based protein thiol oxidation by H_2O_2-scavenging peroxidases. J Biol Chem **284**: 31532–31540

Haber Z, Lampl N, Meyer AJ, Zelinger E, Hipsch M, Rosenwasser S (2021) Resolving diurnal dynamics of the chloroplastic glutathione redox state in Arabidopsis reveals its photosynthetically-derived oxidation. Plant Cell. doi: 10.1093/plcell/koab068

Hanson GT, Aggeler R, Oglesbee D, Cannon M, Capaldi RA, Tsien RY, Remington SJ (2004) Investigating mitochondrial redox potential with redox-sensitive green fluorescent protein indicators. J Biol Chem **279**: 13044–13053

Hipsch M, Lampl N, Zelinger E, Barda O, Waiger D, Rosenwasser S (2021) Sensing stress responses in potato with whole-plant redox imaging. Plant Physiol. doi: 10.1093/plphys/kiab159

Jiang K, Schwarzer C, Lally E, Zhang S, Ruzin S, Machen T, Remington SJ, Feldman L (2006) Expression and characterization of a redox-sensing green fluorescent protein (reduction-oxidation-sensitive green fluorescent protein) in Arabidopsis. Plant Physiol **141**: 397 LP – 403

König J, Baier M, Horling F, Kahmann U, Harris G, Schürmann P, Dietz K-J (2002) The plant-specific function of 2-Cys peroxiredoxin-mediated detoxification of peroxides in the redox-hierarchy of photosynthetic electron flux. Proc Natl Acad Sci **99**: 5738 LP – 5743

Lim SL, Voon CP, Guan X, Yang Y, Gardeström P, Lim BL (2020) In planta study of photosynthesis and photorespiration using NADPH and NADH/NAD+ fluorescent protein sensors. Nat Commun. doi: 10.1038/s41467-020-17056-0

Mehler AH (1951) Studies on reactions of illuminated chloroplasts: I. Mechanism of the reduction of oxygen and other hill reagents. Arch Biochem Biophys **33**: 65–77

Meyer a J, Dick TP (2010) Fluorescent protein-based redox probes. Antioxid Redox Signal **13**: 621–650

Meyer AJ, Brach T, Marty L, Kreye S, Rouhier N, Jacquot J-P, Hell R (2007) Redox-sensitive GFP in Arabidopsis thaliana is a quantitative biosensor for the redox potential of the cellular glutathione redox buffer. Plant J **52**: 973–986

Mittler R, Vanderauwera S, Gollery M, Van Breusegem F (2004) Reactive oxygen gene network of plants. Trends Plant Sci **9**: 490–498

Morgan B, Van Laer K, Owusu TNE, Ezeriņa D, Pastor-Flores D, Amponsah PS, Tursch A, Dick TP (2016) Real-time monitoring of basal H_2O_2 levels with peroxiredoxin-based probes. Nat Chem Biol **12**: 437–443

Müller-Schüssele SJ, Schwarzländer M, Meyer AJ (2021) Live monitoring of plant redox and energy physiology with genetically encoded biosensors. Plant Physiol **186**: 93–109

Mullineaux PM, Exposito-Rodriguez M, Laissue PP, Smirnoff N (2018) ROS-dependent signalling pathways in plants and algae exposed to high light: Comparisons with other eukaryotes. Free Radic Biol Med **122**: 52–64

Niemeyer J, Scheuring D, Oestreicher J, Morgan B, Schroda M (2020) Real-time monitoring of subcellular H_2O_2 distribution in *Chlamydomonas reinhardtii*; bioRxiv 2020.11.13.382085

Nietzel T, Elsässer M, Ruberti C, Steinbeck J, Ugalde JM, Fuchs P, Wagner S, Ostermann L, Moseler A, Lemke P, et al (2019) The fluorescent protein sensor roGFP2-Orp1 monitors in vivo H_2O_2 and thiol redox integration and elucidates intracellular H_2O_2 dynamics during elicitor-induced oxidative burst in Arabidopsis. New Phytol **221**: 1649–1664

Ojeda V, Pérez-Ruiz JM, Cejudo FJ (2018) 2-Cys Peroxiredoxins participate in the oxidation of chloroplast enzymes in the dark. Mol Plant **11**: 1377–1388

Pérez-Ruiz JM, Cejudo FJ (2009) A proposed reaction mechanism for rice NADPH thioredoxin reductase C, an enzyme with protein disulfide reductase activity. FEBS Lett **583**: 1399–1402

Pulido P, Spínola MC, Kirchsteiger K, Guinea M, Pascual MB, Sahrawy M, Sandalio LM, Dietz K-J, González M, Cejudo FJ (2010) Functional analysis of the pathways for 2-Cys peroxiredoxin reduction in Arabidopsis thaliana chloroplasts. J Exp Bot **61**: 4043–4054

Rizhsky L, Liang H, Mittler R (2003) The water-water cycle is essential for chloroplast protection in the absence of stress. J Biol Chem **278**: 38921–38925

Sassenrath-Cole GF, Pearcy RW (1994) Regulation of photosynthetic induction state by the magnitude and duration of low light exposure. Plant Physiol **105**: 1115 LP – 1123

Scheibe R, Dietz KJ (2012) Reduction-oxidation network for flexible adjustment of cellular metabolism in photoautotrophic cells. Plant, Cell Environ **35**: 202–216

Schürmann P, Buchanan BB (2008) The ferredoxin/thioredoxin system of oxygenic photosynthesis. Antioxid Redox Signal **10**: 1235–74

Serrato AJ, Pérez-Ruiz JM, Spínola MC, Cejudo FJ (2004) A Novel NADPH thioredoxin reductase, localized in the chloroplast, which deficiency causes hypersensitivity to abiotic stress in Arabidopsis thaliana. J Biol Chem **279**: 43821–43827

Smirnoff N, Arnaud D (2019) Hydrogen peroxide metabolism and functions in plants. New

Phytol **221**: 1197–1214

Tao R, Zhao Y, Chu H, Wang A, Zhu J, Chen X, Zou Y, Shi M, Liu R, Su N, Du J, Zhou HM, Zhu L, Qian X, Liu H, Loscalzo J, Yang Y (2017) Genetically encoded fluorescent sensors reveal dynamic regulation of NADPH metabolism. Nat Methods 14: 720–728

Vaseghi M-J, Chibani K, Telman W, Liebthal MF, Gerken M, Schnitzer H, Mueller SM, Dietz K-J (2018) The chloroplast 2-cysteine peroxiredoxin functions as thioredoxin oxidase in redox regulation of chloroplast metabolism. Elife **7**: e38194

Wolf B-C, Isaacson T, Tiwari V, Dangoor I, Mufkadi S, Danon A (2020) Redox regulation of PGRL1 at the onset of low light intensity. Plant J **103**: 715–725

Yoshida K, Hara A, Sugiura K, Fukaya Y, Hisabori T (2018) Thioredoxin-like2/2-Cys peroxiredoxin redox cascade supports oxidative thiol modulation in chloroplasts. Proc Natl Acad Sci **115**: E8296 LP-E8304

Yoshida K, Matsuoka Y, Hara S, Konno H, Hisabori T (2014) Distinct redox behaviors of chloroplast thiol enzymes and their relationships with photosynthetic electron transport in arabidopsis thaliana. Plant Cell Physiol **55**: 1415–1425