Mutation of lipoprotein processing pathway gene *lspA* or inhibition of LspA activity by

globomycin increases MRSA resistance to β-lactam antibiotics

Claire Fingleton[1], Merve S. Zeden[1], Emilio Bueno[2], Felipe Cava[2] and James P. O'Gara[1]*

[1]Microbiology, School of Natural Sciences, National University of Ireland, Galway, Ireland.

[2]Department of Molecular Biology, Umeå University, MIMS - Laboratory for Molecular

Infection Medicine Sweden, Umeå, Sweden.

*For correspondence: jamesp.ogara@nuigalway.ie

Running title: Lipoprotein processing and β-lactam susceptibility in MRSA

Abstract

The *Staphylococcus aureus* cell envelope comprises numerous components, including

peptidoglycan (PG), wall teichoic acids (WTA), lipoteichoic acids (LTA), targeted by

antimicrobial drugs. MRSA resistance to methicillin is mediated by the *mecA*-encoded β-

lactam-resistant transpeptidase, penicillin binding protein 2a (PBP2a). However, PBP2a-

dependent β-lactam resistance is also modulated by the activity of pathways involved in the

regulation or biosynthesis of PG, WTA or LTA. Here, we report that mutation of the lipoprotein

signal peptidase II gene, *lspA*, from the lipoprotein processing pathway, significantly increased

β-lactam resistance in MRSA. Mutation of *lgt*, which encodes diacylglycerol transferase (Lgt)

responsible for synthesis of the LspA substrate did not impact β-lactam susceptibility.

Consistent with previous reports, *lgt* and *lspA* mutations impaired growth in chemically

defined media, but not in complex broth. MRSA exposure to the LspA inhibitor globomycin

also increased β-lactam resistance. Mutation of *lgt* in an *lspA* background restored β-lactam

resistance to wild type. The *lspA* mutation had no effect on PBP2a expression, PG composition

or autolytic activity indicating a potential role for WTA or LTA. The *lspA* and *lgt* mutants

exhibited marginally increased resistance to the D-alanine pathway inhibitor D-cycloserine. In

addition, mutation of *lgt* and multicopy *lspA* expression, but not mutation of *lspA,* significantly

increased susceptibility to the lipoteichoic acid synthase inhibitor Congo red revealing

complex interplay between lipoprotein processing mutations and the expression/stability of

cell surface glycopolymers. These findings indicate that accumulation of the LspA substrate,

diacylglyceryl lipoprotein, increases MRSA resistance to β-lactam antibiotics through impacts

on cell envelope components other than PG.

Introduction

The cell envelope of *Staphylococcus aureus* comprises a cytoplasmic membrane surrounded

by a thick peptidoglycan layer, cell wall-anchored proteins, lipoteichoic acids (LTA), wall

teichoic acids (WTA) and cell surface proteins. Accurate biosynthesis, assembly and stability

of these cell envelope components is essential for the growth and pathogenesis of *S. aureus*,

and is the target of numerous antimicrobial agents (1). The peptidoglycan layer determines

cell shape and protects the cell from osmotic lysis, cell surface proteins have important roles

in adhesion, biofilm formation, and immune evasion, and teichoic acids are involved in

protecting the cell from the activity of cationic antimicrobial peptides.

Methicillin resistance in MRSA is mediated by the *mecA*-encoded, low-affinity penicillin-

binding protein 2a (PBP2a) carried on the mobile staphylococcal cassette chromosome *mec*

resistance (HeR) under laboratory growth conditions. HeR strains can become highly or

homogeneously resistant (HoR) after selection on elevated β-lactam concentrations via

poorly understood mechanisms, which require accessory mutations at other chromosomal

loci frequently associated with activation of the stringent response, cyclic-di-adenosine

monophosphate (c-di-AMP) signalling pathway (2-5), the activity of RNA polymerase (6) and

the ClpXP chaperone-protease complex (7,8). In addition, methicillin susceptible *S. aureus*

strains (MSSA) lacking *mecA* can also acquire low-level resistance through adaptive mutations

impacting the c-di-AMP signalling pathway and ClpXP activity (9).

Bacterial lipoproteins are a class of lipid-modified membrane proteins, involved in a range of

diverse functions such as; nutrient acquisition (10), signal transduction (11), respiration (12),

protein folding (13), virulence (14), antibiotic resistance (15) and host invasion (16). Mature

lipoproteins are composed of lipid moieties, specifically acyl groups, linked to the N-terminal

of a protein. The hydrophobic nature of the acyl groups serves as a membrane anchor for the

lipoprotein (17). In Gram-negative bacteria, lipoproteins reside in both the cytoplasmic and

outer membranes, while in Gram-positive bacteria they are anchored in the outer leaflet of

the cytoplasmic membrane and the protein portion may extend into the cell wall and beyond

(18).

Lipoprotein genes are estimated to comprise 1-3% to all genes in bacterial genomes (18).

While many lipoproteins have been identified and experimentally validated, others are

putatively identified using predictive software, thus their functions remain unknown (19). A

recent bioinformatic evaluation of Staphylococcal lipoproteins in the MRSA strain USA300

identified 67 lipoproteins, comprising 2.57% of all genes (18). When grouped by function, 25

of the 67 lipoproteins were implicated in ion (notably iron) and nutrient transport, 8 were

ascribed miscellaneous functions including sex pheromone biosynthesis, respiration,

chaperone-folding, and protein translocation and 15 were classified as tandem lipoproteins,

of which 9 are "lipoprotein-like" lipoproteins, known to play a role in host cell invasion (16).

The remaining 19 lipoproteins were not assigned any known function.

Lipoproteins are synthesised in a precursor form called preprolipoproteins. The N-terminal

domain includes a type II signal peptide, approximately 20 amino acids in length (20), which

enables translocation of preprolipoproteins to the cytoplasmic membrane, predominantly via

the general secretory (Sec) pathway (21). The signal peptide has 3 distinct domains; a

positively charged N domain, a hydrophobic H domain and a C-terminal lipobox. The lipobox

is comprised of a conserved 3-amino acid sequence $[LVI]_{-3}$ $[ASTVI]_{-2}$ $[GAS]_{-1}$ in front of an

invariant cysteine residue $[C]_{+1}$ (22,23). This lipobox serves as a recognition site for enzymes

of the lipoprotein processing pathway, enabling lipid modification of the cysteine residue, and

cleavage of the signal peptide between the amino acid at position -1 and the +1 cysteine (22).

The first enzyme in the lipoprotein processing pathway is diacylglycerol transferase (Lgt)

which covalently attaches a diacylglycerol molecule from phosphatidyl glycerol onto the

sulfhydryl group of the invariant cysteine, resulting in a prolipoprotein (24). This diacylglycerol

serves as a membrane anchor. Next, the type II lipoprotein signal peptidase (Lsp) cleaves the

signal peptide between the amino acid at position -1 and +1, leaving the invariant cysteine

residue as the new terminal amino acid (25). Lgt and Lsp are conserved in all bacterial species.

In Gram-negative bacteria, a third step is catalysed by the enzyme N-acyl transferase (Lnt),

which transfers an N-acyl group onto the invariant cysteine residue at the N-terminal of the

protein (26). Lnt homologs have been identified in high-GC Gram-positive bacteria (27) but

not in low-GC Firmicutes. Despite the lack of an apparent Lnt homolog, N-acylated

lipoproteins have been identified in *S. aureus* (28) and recent work has identified two novel

non-contiguous genes *lnsA* and *lnsA* which catalyse the N-terminal acylation of lipoproteins

in *S. aureus* (29). The membrane metalloprotease Eep and the EcsAB transporter were shown

to be involved in the processing and export of linear peptides, including the signal peptide

cleaved by LspA in the lipoprotein processing pathway (30-32). Lgt and Lsp are essential for

the viability of Gram-negative bacteria (33). In contrast, *lgt* and *lspA* mutations do not impact

viability in Gram-positive bacteria (34), but are associated with changes in growth,

immunogenicity (35) and virulence (14) phenotypes.

In this study, we characterised the impact of *lspA and lgt* mutations, alone and in

combination, on susceptibility to β-lactams, D-cycloserine and Congo red, growth, PBP2a

expression, peptidoglycan structure, and autolytic activity in MRSA. The impact of

globomycin, which is known to inhibit LspA activity, on β-lactam susceptibility was also

characterised. Our data suggest that accumulation of the LspA substrate, diacylglyceryl-

prolipoprotein, modulates resistance to β-lactam antibiotics in MRSA.

Results

Mutation of *lspA* in MRSA increases resistance to β-lactam antibiotics. The Nebraska Transposon Mutant Library (NTML) (36) was screened to identify mutants exhibiting altered susceptibility to cefoxitin, which is recommended as a surrogate for measuring *mecA*-mediated oxacillin resistance in clinical laboratories, in accordance with Clinical and Laboratory Standards Institute (CLSI) guidelines for disk diffusion susceptibility assays. Mutants identified by this screen included NE869 (*yjbH)* (37), NE1909 (*sagA*) (38) and NE810 (*cycA*)(39), all of which have previously been implicated in β-lactam resistance. A new mutant identified in this screen was NE1757 (*lspA*::Tn), which exhibited increased resistance to cefoxitin (Fig 1A, B). PCR was used to confirm the presence of *lspA*::Tn allele in NE1757 (data not shown). Using E-test strips, as described in the methods, oxacillin MIC of the *lspA* mutant NE1757 was found to be 128 - 256 µg/ml, compared to 32 - 64 µg/ml for JE2 (Fig. 1C). Two JE2 transductants carrying the *lspA*::Tn allele from NE1757 also exhibited increased resistance to oxacillin (Fig. 1C).

The increased oxacillin resistance phenotype of NE1757 was also complemented by the introduction of a plasmid (pLI50)-borne copy of the wild type *lspA* gene (p*lspA*) into the mutant. Growth of JE2, NE1757, NE1757 pLI50 and NE1757 p*lspA* on MHA 2% NaCl supplemented with oxacillin 32 µg/ml visually demonstrated that carriage of the complementation plasmid reversed the increased oxacillin resistance phenotype of NE1757 (Fig. S1). Measurement of oxacillin MICs by agar dilution showed that NE1757 and NE1757 pLI50 had MICs of 256 µg/ml, while JE2 and the complemented strain NE1757 p*lspA* had MICs of 64 µg/ml (Table 1).

Comparison of JE2 and NE1757 growth in MHB, MHB 2% NaCl, TSB and TSB 0.5 mg/ml oxacillin revealed no significant differences (Fig. S2). Similarly, population analysis profiling revealed that the heterogeneous pattern of oxacillin resistance expressed by JE2 was unchanged in NE1757 (Fig. S3). These observations indicate that the increased β-lactam resistance phenotype of NE1757 was not attributable to any growth advantage or change in the heterogeneous/homogeneous oxacillin resistance profile.

Comparative WGS analysis confirmed that the only change in the NE1757 genome was the insertion of the *Bursa aurealis* transposon in the *lspA* gene at position 1192002 (Table S1) and there were no SNPs present. The NE1757 genome was also checked manually for zero

147   coverage regions to confirm the absence of any large deletions and insertions. Taken together
148   these data indicate that mutation of *lspA*, which encodes lipoprotein signal peptidase II in the
149   lipoprotein processing pathway (Fig. 2), increases resistance to oxacillin and cefoxitin in JE2.
150
Mutation of *lspA* does not affect PBP2a expression or peptidoglycan structure and
crosslinking. One of the numerous lipoproteins processed by LspA is PrsA, a chaperone and
foldase protein, reported to play a role in PBP2a folding and β-lactam resistance (40). To
investigate if improper processing of PrsA, or another lipoprotein, in the *lspA* mutant
impacted on PBP2a, Western blotting was used to compare PBP2a expression in JE2, NE1757
and NE1757 p*lspA.* The MSSA strain 8325-4 was included as a *mecA*-negative control. Growth
of JE2 and NE1757 in MHB without 2% NaCl at 37°C (data not shown) or MHB with 2% NaCl
at 35°C supplemented with 0.5 μg/ml oxacillin (Fig. 3A) revealed similar levels of PBP2a
expression in all strains.
Quantitative peptidoglycan compositional analysis was performed using UPLC analysis of
muramidase-digested muropeptide fragments extracted from exponential or stationary
phase cultures of JE2 and NE1757 grown in MHB or MHB supplemented with oxacillin 3 mg/ml
or 32 μg/ml. The PG profile of JE2 and the *lspA* mutant NE1757 were similar under all growth
conditions tested (Fig. 3B). Thus, supplementation of MHB with oxacillin was associated with
a significant changes in muropeptide oligomerization and reduced crosslinking, but these
effects were the same in both JE2 and NE1757 (Fig. S4). The total PG concentrations extracted
from JE2 and NE1757 cell pellets was also the same (data not shown). Comparison of Triton
X-100-induced autolysis in JE2 and NE1757 also revealed identical autolytic profiles (Fig. S5).
Finally, the NaCl tolerance phenotypes of JE2 and NE1757 were also similar (Fig. S6), indicating
that c-di-AMP signalling, which has previously been implicated in the control of β-lactam
resistance, autolytic activity and NaCl tolerance (4,5) was unaffected by the *lspA* mutation.
These data indicate that increased β-lactam resistance in the *lspA* mutant was not associated
with significance changes in PG abundance, structure, crosslinking, c-di-AMP signalling or
autolytic activity.
175
Exposure to the LspA inhibitor globomycin also increases β-lactam resistance. Globomycin
is a natural peptide antibiotic, first discovered in 1978, produced by 4 different strains of
actinomycetes (41,42). It is an inhibitor of LspA and works by sterically blocking the active site

of the enzyme (43). Globomycin has moderate to strong antibacterial activity against many

Gram-negative species and has been proposed to cause disruption of cell surface integrity

(44). However, despite its ability to inhibit LspA, globomycin does not have significant

antimicrobial activity against Gram-positive bacteria including *S. aureus*, with MICs >100

μg/ml (41,42,45).

Because disruption of *lspA* increased resistance to β-lactams, we hypothesized that chemical

inhibition of LspA by globomycin may also be associated with increased β-lactam resistance.

To test this hypothesis, the susceptibility of JE2 and NE1757 to oxacillin was determined in

the presence or absence of globomycin. A series of JE2 and NE1757 cultures grown in MHB

2% NaCl were used to determine that oxacillin 40 μg/ml inhibited growth of JE2 but not

NE1757 (Fig. 4A) as predicted. Next, JE2 MHB 2% NaCl 40 μg/ml oxacillin cultures were further

supplemented with globomycin 10, 20, 30, 40 or 50 μg/ml (Fig. 4B). Oxacillin-induced

inhibition of JE2 growth was rescued globomycin, optimally at 10, 20 and 30 μg/ml (Fig. 4B).

Growth of JE2 in oxacillin 40 μg/ml and globomycin 50 μg/ml was substantially impacted

compared to lower globomycin concentration (Fig. 4B) indicating that the antagonism of

oxacillin by globomycin was dose dependent.

These experiments were extended to USA300_FPR3757, from which JE2 is derived and

ATCC43300, a SCC*mec* type II MRSA clinical isolate. Oxacillin concentrations of 50 μg/ml and

30 μg/ml inhibited growth of USA300 and ATCC4330, respectively (Fig. 5). Globomycin

concentrations (determined empirically for each strain) of 20 μg/ml for USA300 (Fig. 4C) and

30 μg/ml for ATCC43300 (Fig. 4D) rescued growth in the presence of oxacillin. These data

demonstrate that the increased oxacillin resistance phenotype observed in the *lspA* mutant

NE1757, can be replicated by globomycin-induced inhibition of LspA activity in wild type JE2

and other MRSA strains.

In contrast to the observation that globomycin increased β-lactam resistance in wild type JE2,

USA300 and ATCC43300, the growth of NE1757 in a range of globomycin concentrations from

10 – 50 μg/ml had a dose-dependent and negative effect on growth in the presence of

oxacillin 40μg/ml (Fig. S7). Taken together, these data suggest that globomycin antagonises

β-lactam antibiotics, increasing MRSA resistance to oxacillin and cefotaxime in LspA-

dependent manner. However, in the absence of *lspA*, the combination of globomycin and

oxacillin interferes with growth of JE2, particularly at higher concentrations of globomycin,

perhaps due to off-target effects.

To determine if globomycin could also increase resistance to other classes of β-lactam

antibiotics, its effect on cefotaxime resistance in JE2, USA300 and ATCC43300 was evaluated.

Cefotaxime was chosen because it is a 3[rd] generation cephalosporin with broad spectrum

activity against Gram-positive and Gram-negative bacteria commonly used in the clinic,

whereas oxacillin is a narrow-spectrum penicillin antibiotic. Cefotaxime 40 µg/ml inhibited

growth of JE2 and USA300 (Fig. 5A and B), while 30 µg/ml inhibited growth of ATCC43300 (Fig.

5C). Globomycin (30 µg/ml) rescued growth of all three strains in oxacillin (Fig. 5A, B and C)

suggesting that globomycin-mediated inhibition of LspA may antagonise the activity of many

or all β-lactam antibiotics.

Mutation of *lgt* increases susceptibility to the lipoteichoic acid synthase inhibitor Congo

red. Congo red was recently identified as a selective inhibitor of lipoteichoic acid synthase

(LtaS) activity (46). To investigate the possible involvement of LTA expression or stability in

the increased resistance of the *lspA* mutant to β-lactams serial dilutions of overnight *lspA*, *lgt*

and *lspA*/*lgt* mutant cultures were spotted onto TSA 0.125% Congo Red. Our data showed

that mutation of *lgt* dramatically increased susceptibility to the selective lipoteichoic acid

synthase (LtaS) inhibitor Congo red (Fig. 6) suggesting that impaired lipoprotein processing

affects the expression or stability of LTA. However, while the *lgt*/*lspA* double mutant was even

more susceptible to Congo red than the single *lgt* mutant, the single *lspA* mutations in NE1757

and NE1757 MM did not significantly alter Congo red susceptibility and complementation of

NE1757 was also associated with increased susceptibility (Fig. 6). These data show that

mutation of *lgt* or multicopy expression of *lspA* both increase susceptibility to this LTA

inhibitor indicating that the lipoprotein processing pathway impacts LTA synthesis/stability in

a complex manner. The mutations in the *lgt* and *lspA* genes alone, and in particular when

combined, significantly increased resistance to the alanylation pathway inhibitor D-

cycloserine (DCS)(Table 1). However lack of correlation between the effects of *lgt* and *lspA*

mutations on susceptibility to Congo red, DCS and β-lactams indicates that further analysis is

needed to better understand the interactions between lipoprotein processing pathway

intermediates and the expression/stability of LTA and WTA.

Mutation of *lgt* in the *lspA* background restores wild type levels of β-lactam resistance. LspA catalyses the second major step in the lipoprotein processing pathway (Fig. 2). To probe the contribution of lipoprotein processing to LspA-controlled oxacillin resistance, we compared the impact of *lgt, lspA* and *lgt/lspA* mutants on growth and resistance to oxacillin, as well as cefotaxime, nafcillin and vancomycin. Lgt catalyses the addition of a diacylglycerol moiety onto preprolipoproteins, from which the signal peptide is then cleaved by LspA (Fig. 2). To construct a *lspA/lgt* double mutant the erythromycin resistance marker of the *lspA*::Tn allele in NE1757 was first exchanged for a markerless transposon to generate a strain designated NE1757 MM into which the erythromycin-marked *lgt*::Tn allele from NE1905 was transduced. Comparison of growth of JE2, NE1757, NE1757 MM, NE1905 and the *lgt/lspA* double mutant NE1757/NE1905 revealed no significant differences MHB, MHB with 2% NaCl or TSB (Fig. 7A, B, C). However, consistent with previous analysis of a *lgt* mutant (35), the *lspA, lgt* and in particular the *lgt/lspA* mutants exhibited impaired growth in CDM (Fig. 7D), suggesting that aberrant lipoprotein processing may affect nutrient acquisition under substrate-limiting conditions.

The *lgt* mutant NE1905 exhibited no changes in susceptibility to oxacillin, cefotaxime, nafcillin or vancomycin (Table 1). As observed for oxacillin, the *lspA* mutant NE1757 was more resistant to cefotaxime and nafcillin and these phenotypes were complemented by the p*lspA* plasmid (Table 1). Neither the *lspA* nor *lgt* mutations increased resistance to vancomycin (Table 1). Oxacillin and nafcillin MICs were restored to wild type levels in the *lspA/lgt* double mutant, and the cefotaxime MIC was significantly reduced (Table 1).

Taken together, these data indicate that while mutation of *lspA* or *lgt* or both impact growth in CDM, only mutation of *lspA* alone is associated with increased β-lactam resistance. The *lgt* mutation and possible accumulation of unprocessed prolipoproteins (Fig. 8A,B) does not increase β-lactam resistance, whereas the possible accumulation of diacylglyceryl-lipoprotein in a LspA mutant (Fig. 8C, D) is associated with this phenotype. The oxacillin, cefotaxime and nafcillin MICs of the *ecsB* mutant from the NTML were the same as wild type (Table 1) indicating that downstream processing of the LspA-cleaved signal peptide is not associated with altered β-lactam resistance.

Discussion

Advances in our understanding of the accessory factors that control levels of *mecA*/PBP2a-dependent resistance to methicillin has the potential to reveal new therapeutic targets and drugs that may facilitate the reintroduction of other β-lactam antibiotics for the treatment of MRSA infections. In this study, we demonstrated that mutation of the lipoprotein processing pathway gene *lspA*, or inhibition of LspA with globomycin increased resistance to β-lactam antibiotics. Although numerous mutations impacting the stringent response (ppGPpp) and c-di-AMP signalling are associated with the transition from a heterogeneous to homogeneous pattern of resistance and elevated PBP2a expression (2,3), our data show that the *lspA* mutation was not associated with a HoR phenotype, increased PBP2a expression, or altered NaCl tolerance (which is controlled by c-di-AMP) (4,5). On the other hand, changes in β-lactam resistance independent of altered PBP2a regulation has long been known (47-49), and several auxiliary factors known to influence β-lactam resistance in MRSA have been described (49-54). In addition to unchanged PBP2a expression, no evidence of peptidoglycan remodelling was observed in NE1757 after growth in the presence or absence of oxacillin, potentially implicating wall teichoic acid (WTA) or lipoteichoic acid (LTA) synthesis or stability in the *lspA* mutant phenotype. Inhibition of WTA synthesis was previously shown to decrease β-lactam resistance in a PBP2a-independent manner (50). Reduced LTA stability, as evidenced by Western blotting and increased susceptibility to the selective lipoteichoic acid synthase inhibitor Congo red, was recently correlated with a PBP2a-independent reduction in β-lactam resistance in auxiliary factor *auxA* and *auxB* mutants (49). Interestingly AuxA is structurally similar to SecDF (49) and may interact with Sec pathway and lipoprotein processing (Fig. 8). The *lgt* mutant was significantly more susceptible to Congo red than the *lspA* mutant and multicopy expression of *lspA* also increased Congo red susceptibility suggesting that although the lipoprotein processing pathway modulates LTA synthesis/stability, the relationship between these two pathways appears to be complex.

Consistent with previous studies of lipoprotein pathway processing mutants in *S. aureus* (35), *Listeria monocytogenes* (55) and *Streptococcus agalactiae* (56), our analysis also showed that the *lgt, lspA* and *lgt/lspA* double mutants all exhibited impaired growth in CDM but not in TSB or MHB indicating that the impact of lipoprotein processing pathway mutations on nutrient acquisition and growth under nutrient limiting conditions can be compensated in rich media.

Mutation of the *lgt* gene from the lipoprotein processing pathway (Figs. 2, 7) did not affect β-

lactam resistance and introduction of the *lgt* mutation into a *lspA* mutant restored wild type

levels of resistance. These data implicate accumulation of diacylglycerol lipoprotein in

elevated β-lactam resistance. Consistent with this possibility, lipoproteins were retained in

the membrane of a *S. agalactiae lspA* mutant, but were released into the supernatant in large

concentrations by *lgt* and *lgt/lspA* mutants (56). Lipoproteins synthesised by the *S. agalactiae*

lspA mutant retained signal peptide, which was absent in a *lgt* mutant with LspA activity.

Importantly, signal peptide processing also occurred in the *lgt/lspA* double mutant, albeit

with cleavage occurring between different amino acids, implicating the involvement of an

alternative signal peptidase. Our analysis revealed no change in oxacillin susceptibility in the

MRSA *lgt/lspA* double mutant, indicating that even if an alternative peptidase can cleave the

signal peptide, this may not impact β-lactam resistance. Furthermore, mutation of *ecsB*,

which has recently been implicated in export of linear peptides from the lipoprotein

processing pathway in *S. aureus* (32), did not change the oxacillin MIC in JE2 (Table 1) also

indicating that downstream processing of signal peptides cleaved from lipoproteins by LspA

is not associated with altered β-lactam resistance. In *L. monocytogen*es, deletion of *lgt* also

led to significant release of lipoproteins into the supernatant. However, treatment of the *L.*

monocytogenes lgt mutant with globomycin (inhibiting LspA activity) resulted in enhanced

lipoprotein retention in the membrane (55), suggesting that the impact of globomycin and

lspA mutation on lipoprotein processing is not necessarily the same. In a *S. aureus lgt* mutant,

the Götz group reported that the majority of lipoprotein (lacking signal peptide) was released

into the supernatant (35). Taken together, the data suggest that accumulation of membrane-

anchored diacylglycerol lipoprotein with uncleaved signal peptide, or lipoprotein that is mis

localised or released due to aberrant signal peptide processing by an alternative peptidase, is

accompanied by increased β-lactam resistance in MRSA.

Experimental procedures

Bacterial strains and culture conditions. Bacterial strains and plasmids used in this study are listed in Table S2. *Escherichia coli* strains were grown in Luria-Bertani (LB) broth or agar (LBA). *S. aureus* strains were grown in Tryptic Soy Broth (TSB), Tryptic Soy Agar (TSA) or chemically defined medium (CDM) (57). Muller Hinton Broth (MHB) or Muller Hinton Agar (MHA) (Oxoid) supplemented with 2% NaCl where indicated, were used for antimicrobial susceptibility testing (AST). Antibiotic concentrations used were 10 µg/ml erythromycin, 10 µg/ml chloramphenicol, 75 µg/ml kanamycin, 100 µg/ml ampicillin.

Two hundred and fifty ml flasks were filled with 25 ml growth media, and overnight cultures were used to inoculate the media at a starting OD_{600} of 0.05. Overnight cultures were grown in TSB, and washed once in 5 ml PBS before being used to inoculate the CDM cultures. Flasks were incubated at 37°C shaking at 200 rpm. OD_{600} readings were measured at 1 - 2 h intervals. Colony forming units (CFU) were enumerated in serially diluted 20 µl aliquots removed from flask cultures. Three independent biological replicates were performed for each strain and the resulting data plotted using GraphPad Prism software.

Data from growth experiments in a Tecan Sunrise microplate instrument were recorded by Magellan software. Overnight cultures were adjusted to an OD_{600} of 1 in fresh media and 10 µl inoculated into 200 µl media per well before being incubated at 37°C for 24 h with shaking. OD_{600} was recorded every 15 min. For CDM, overnight TSB cultures were first washed in 5 ml PBS and adjusted to OD_{600} of 1 in PBS.

Genetic manipulation of *S. aureus*. Phage 80α transduction was used to verify the association between antibiotic resistance phenotypes and transposon insertion-marked mutations from the NTML as described previously (39). Transductants were verified by PCR amplification of the target locus using primers listed in Table S3. The plasmid pTnT, which contains a truncated, markerless transposon was used to construct a markerless *lspA* mutant designated NE1757 MM, as described previously (58). A double *lgt/lspA* double mutant was subsequently constructed using phage 80α to transduce the *lgt*::Tn allele from NE1905 into NE1757 MM. A 1324 bp fragment encompassing the *lspA* gene was PCR amplified from JE2 genomic DNA using primers NE1757_INF#3_Fwd and NE1757_INF#3_Rev (Table 2), purified with GenElute™ PCR Clean-Up Kit and cloned into the *E. coli - Staphylococcus* shuttle vector pLI50 digested with *Eco*RI (New England Biolabs) using In-Fusion® HD Cloning Kit (Clontech) to generate

p*lspA.* Using electroporation, p*lspA* was transformed sequentially into Stellar™ (*E. coli* HST08) (Clontech), *S. aureus* RN4220 and finally NE1757.

Disk diffusion susceptibility assays. Cefoxitin disk diffusion susceptibility assays were performed in accordance with CLSI guidelines. Briefly, isolates were grown at 37°C on MHA for 24 h and 5 - 6 colonies were resuspended in 0.85% saline to OD_{600} of 0.08 - 0.1 (0.5 McFarland; 1×10^8 CFU/ml) and swabbed onto MHA plates with a uniform agar depth of 4 mm. A 30 μg cefoxitin disk (Oxoid) was applied, the plate incubated at 35°C for 16 - 18 h and the zone of inhibition diameter measured. Strains were classified as sensitive, intermediate, or resistant, according to CLSI criteria.

Minimum inhibitory concentration (MIC) measurements. For oxacillin M.I.C.Evaluators (Oxoid) several colonies from 24 h MHA plates were resuspended in 0.85% saline to OD_{600} of 0.08 - 0.1, (0.5 McFarland standard) and evenly swabbed onto MHA 2% NaCl (4 mm agar depth). An M.I.C.Evaluator strip was applied and the plate incubated at 35°C for 24 h. Three biological repeats were performed for each strain.

For broth microdilution MIC measurements using 96-well plates, each plate row was used to prepare two-fold dilutions of antibiotic in MHB, typically ranging from 256 - 0.5 μg/ml across 10 wells. For oxacillin and nafcillin MIC measurements, MHB 2% NaCl was used. Several colonies from 24 h MHA plates were resuspended in 0.85% saline to OD_{600} of 0.08 - 0.1 (0.5 McFarland standard), diluted 1:20 in 0.85% saline and 10 μl of this cell suspension used to inoculate each well (approximately 5×10^4 CFU/well) in a final volume of 100 μl. The plates were incubated at 35°C for 16 - 20 h, or 24 h incubation for oxacillin and nafcillin. The MIC was the lowest concentration of antimicrobial agent that completely inhibited growth.

Freshly prepared MHA plates (with 2% NaCl when using oxacillin and nafcillin) were supplemented with antimicrobial agents at 0.5, 1, 2, 4, 8, 16, 32, 64, 128 and 256 μg/ml to perform agar dilution MIC measurements. The inoculum was prepared as described for the broth microdilution method above before being further diluted 1:20 in 0.85% saline and 4 μl spot-inoculated onto each plate (approximately 10^4 CFU per 5-8 mm diameter spot). The MIC was the lowest concentration of antimicrobial agent that completely inhibited growth after 16-18 h (24 h for oxacillin and nafcillin) at 35°C, disregarding a single colony or a faint haze associated with the inoculum.

Congo red susceptibility assays were performed as described previously (59). Briefly,
overnight cultures grown in TSB were adjusted to an OD_{600} of 1 in PBS, serially diluted and
spotted onto TSA plates supplemented with 1% aqueous Congo red solution (VWR) to a final
concentration of 0.125%. Plates were incubated for 24 h at 37°C. These assays were
performed with at least 3 biological repeats and representative image is shown for TSA
0.125% Congo red.

Autolytic activity assays. Overnight cultures (20 µl) were inoculated into 20 ml TSB, grown at
37°C (200 rpm) to OD_{600} of 0.5, washed with 20 ml cold PBS, resuspended in 1 ml cold PBS
and finally adjusted to OD_{600} of 1. Triton X-100 was added at a final concentration of 0.1% v/v
and the cell suspension incubated at 37°C with shaking (200 rpm). OD_{600} was recorded every
30 min for 4 h. Autolytic activity was expressed as a percentage of the initial OD_{600}. NE406
(*atl*::Tn) was used as a control and at least 3 biological replicates performed for each strain.

Globomycin and β-lactam antibiotic synergy/antagonism assays. One hundred µl of MHB
cefotaxime or MHB 2% NaCl oxacillin was added to the individual wells of 96-well plates. The
oxacillin or cefotaxime concentration chosen for each strain was based on approximate MICs
i.e. the lowest antibiotic concentration that inhibited growth. Overnight MHB cultures were
resuspended in PBS at OD_{600} of 0.1 (0.5 McFarland standard) and then further diluted 1:20
before 10 µl (approximately 5×10^5 CFU/ml) was added to each well and the plates were
incubated at 35°C with shaking on a Tecan Sunrise microplate instrument for 20 h
(cefotaxime) or 24 h (oxacillin). Globomycin ranging from 10 - 100 µg/ml was then added to
the cefotaxime or oxacillin cultures to measure potential synergism or antagonism. Three
independent biological replicates were performed for each strain and antibiotic combination.

PBP2a western blot analysis. Overnight MHB cultures were used to inoculate 25 ml of MHB
2% NaCl, with or without 0.5 µg/ml oxacillin to a starting OD_{600} of 0.05, incubated at 35°C (200
rpm shaking) until an OD_{600} of 0.8 was reached before the cells were pelleted and
resuspended in PBS to an OD_{600} of 10. Six µl of lysostaphin (10 µg/ml) and 1 µl of DNase (10
µg/ml) was added to 500 µl of this concentrated cell suspension before being incubated at
37°C for 40 min. Next, 50 µl of 10% SDS was added and the incubation continued for a further
20 min. The lysed cells were then pelleted in a microcentrifuge for 15 min, following which
the protein-containing supernatant was collected and total protein concentration

determined using the Pierce® BCA Protein Assay Kit. Samples containing 8 µg total protein

were mixed 1:1 with protein loading buffer (2x) (National Diagnostics) and incubated at 95°C

for 5 min and loaded onto a 7.5% Tris-Glycine gel and separated at 120 V for 60 mins.

Electrophoretic transfer to a PVDF membrane was carried out at 30 V for 30 min on the Trans-

Blot Turbo Transfer System (Biorad). The PVDF membrane was blocked overnight in 5% skim

milk powder in PBS at 4°C. The following day, the membrane was washed in fresh PBS. Anti-

PBP2a (Abnova) was diluted 1:1000 in PBS-Tween 20 (0.1%) and incubated with the

membrane for 1 h at room temperature. The membrane was washed in PBS to remove

unbound antibody. The secondary antibody, HRP-rec-Protein G (Invitrogen) was diluted

1:2000 in PBS-Tween 20 (0.1%) and incubated with the membrane at room temperature for

1 h. Visualisation of the membrane was performed with the Opti-4CN Substrate kit (Biorad).

Three independent experiments were performed and representative images of the

developed PVDF membrane were recorded.

Population level antibiotic resistance profile analysis. Characterisation of the population

resistance profile was performed as described previously (60). Overnight cultures were grown

in TSB, adjusted to an OD_{600} of 1, 10-fold serially diluted from 10^{-1} to 10^{-7}, and a 20 µl aliquot

of each dilution plated onto a series of TSA agar plates supplemented with oxacillin 0.25, 0.5,

1, 2, 4, 8, 16, 32, 64 and 128 µg/ml. CFUs were enumerated after overnight incubation at 37°C

and the results were expressed as CFU/ml at each oxacillin concentration. Three independent

experiments were performed for each strain.

Antibiotic tolerance assay. Tolerance assays were performed as described previously (61).

Briefly overnight TSB cultures were sub-cultured into 25 ml of fresh TSB in 250 ml flasks at a

starting OD_{600} of 0.05 and grown to an OD_{600} of 0.5 at 37°C with 200 rpm shaking. At this time

(T_0) an aliquot was removed for CFU enumeration and 12.5 µg/ml oxacillin promptly added

before the cultures were re-incubated. Antibiotic tolerance was expressed as the % CFU/ml

after 2, 4, 6, 8 and 24 h growth in the antibiotic compared to the CFU/ml at T_0. The results

represent 3 biological replicates of each strain.

Salt tolerance assay. Overnight TSB cultures were adjusted to an OD_{600} of 1 in fresh TSB and

serially diluted from 10^{-1} to 10^{-7}. Four µl of each dilution was spot-inoculated onto TSA 2.2M

NaCl plates and incubated at 37°C overnight.

Genomic DNA extraction and Whole Genome Sequencing (WGS). Genomic DNA (gDNA)

extractions were performed using the Wizard® Genomic DNA Purification Kit (Promega)

following pre-treatment of *S. aureus* cells with 10 µg/ml lysostaphin (Ambi Products LLC) at

37°C for 30 min. WGS was performed by MicrobesNG (http://www.microbesng.uk) using an

Illumina sequencing platform with 2x250 bp paired-end reads. CLC Genomics Workbench

software (Qiagen) was used for genome sequencing analysis of strains. As a reference

genome, a contig was produced for wild type JE2 by mapping Illumina reads onto the closely

related USA300 FPR3757 genome sequence (RefSeq accession number NC_07793.1). The

Illumina short read sequences from NE1757 were then mapped onto the assembled JE2

sequence and the presence of the transposon insertion confirmed. Single Nucleotide

Polymorphisms (SNPs), deletions or insertions were mapped in the NE1757 genome and

presence of large deletions ruled out by manually searching for zero coverage regions using

the CLC Genomics Workbench software.

Peptidoglycan analysis. For each strain and growth condition tested, independent

quadruplicate 50 ml cultures were grown to an OD_{600} of 0.5, harvested and resuspended in 5

ml PBS before peptidoglycan was extracted as described previously (39,62). Mass

spectrometry was performed on a Waters XevoG2-XS QTof mass spectrometer. Structural

characterization of muropeptides was determined based on their MS data and MS/MS

fragmentation pattern, matched with PG composition and structure reported previously (63-

66).

Acknowledgements

This study was funded by grants from the Health Research Board (HRA-POR-2015-1158 and

ILP-POR-2019-102) (www.hrb.ie), the Irish Research Council (GOIPG/2016/36)

(www.research.ie), Science Foundation Ireland (19/FFP/6441) (www.sfi.ie) to J.P.O'G, and

Svenska Forskningsrådet Formas to F.C. We are grateful to Christopher Campbell and Laura

Gallagher for assistance and advice throughout this study. The illustrations in Figures 2 and 8

were created with Biorender.com.

498   **Table and Figures**
499

Table 1. Antibacterial activity of oxacillin, cefotaxime, nafcillin vancomycin and D-cycloserine (MIC measurements; μg/ml[1]) against ATCC 29213 (control), JE2, NE1757 (*lspA*::Tn), NE1757 p*lspA*, NE1757 pLI50, NE1757 MM (markerless *lspA* mutant), NE1757 MM p*lspA*, JE2 *lspA*::Tn #2 (transductant), NE1905 (*lgt*::Tn) and NE1757/NE1905 double mutant *lspA*/*lgt*.

Strain	Oxacillin	Cefotaxime	Nafcillin	Vancomycin	D-Cycloserine
ATCC 29213	≥1	2	0.5	1	32
JE2	32 - 64	64	32	1	32
NE1757 (*lspA*)	128 - 256	256	64	1	16 - 32
NE1757 p*lspA*	64	64	32	1	16 - 32
NE1757 pLI50	256	256	-	1	16 - 32
NE1757 MM	128 - 256	256	64	-	-
NE1757 MM p*lspA*	64	64	32	-	-
JE2 *lspA*::Tn #2	128 - 256	256	-	1	16 - 32
NE1905 (*lgt*)	64	64	32	1	16 - 32
NE1757/NE1905 (*lspA*/*lgt*)	64	128	32	1	16
NE107 (*ecsB*)	64	64	32	1	16 - 32

[1] MIC measurements (μg/ml) were performed by MH agar dilution in accordance with CLSI standards.

Fig. 1. Mutation of *lspA* increases resistance to cefoxitin and oxacillin. (A) Average diameters of the cefoxitin disk zones of inhibition for JE2 and NE1757 (*lspA*::Tn) from 4 independent experiments, plotted using Prism software (GraphPad). (B) Representative image of JE2 (left) and NE1757 (right) grown on MH agar with a cefoxitin 30 µg disk. (C) M.I.C.Evaluator measurement of oxacillin minimum inhibitory concentrations (MICs) in JE2, NE1757 (*lspA*::Tn), and two independent JE2 transductants (#2 and #4) carrying the *lspA*::Tn allele grown on MHB 2% NaCl agar. This assay was repeated 3 independent times for each strain and a representative image is shown.

528
529
Fig. 2. Role of LspA in the proposed model of lipoprotein processing in Gram-positive
bacteria. Lipoproteins are synthesized in the cytoplasm as precursors. **1.** Prolipoprotein is
translocated usually via the Sec machinery through recognition of its signal peptide (SP). This
sequence also contains the lipobox specific for bacterial lipoproteins. **2.** Diacylglyceryl
transferase (Lgt) catalyzes the transfer of a diacylglyceryl group from phosphatidylglycerol
onto the prolipoprotein resulting in diacylglyceryl-prolipoprotein. **3.** Lipoprotein signal
peptidase II (LspA) recognizes the diacylglyceryl-modified signal peptide and cleaves between
the amino acid at position-1 and the lipid-modified cysteine residue at +1. **4.** The lipoprotein
N-acylation transferase system (Lns) converts diacyl lipoproteins to triacylated lipoprotein
(29). **5.** Eep and EcsAB play roles in the processing and secretion of linear peptides (32).
540

Fig. 3. Mutation of *lspA* does not affect PBP2a expression levels or peptidoglycan structure and crosslinking. (A) Western blot of PBP2a protein in JE2, NE1757 (*lspA*), NE1757 p*lspA* and MSSA strain 8325-4 (negative control). Cells were grown to exponential stage in MHB 2% NaCl

supplemented with 0.5 µg/ml oxacillin, with the exception of 8325-4 which was grown in MHB
2% NaCl. For each sample, 8 µg total protein was run on a 7.5% Tris-Glycine gel, transferred
to a PVDF membrane and probed with anti-PBP2a (1:1000), followed by HRP-conjugated
protein G (1:2000) and colorimetric detection with Opti-4CN Substrate kit. Three independent
experiments were performed and a representative image is shown. (**B**) Representative UV
chromatograms of peptidoglycan extracted from JE2 and NE1757 collected from cultures
grown to exponential or stationary phase in MHB or MHB supplemented with oxacillin 3
µg/ml or 32 µg/ml. Each profile shown is a representative of 3 biological replicates.

Fig. 4. Mutation of *lspA* or exposure to globomycin increases oxacillin resistance. (A) JE2 and NE1757 (*lspA*) grown in MHB 2% NaCl supplemented with oxacillin 40 µg/ml. **(B)** JE2 grown in MHB 2% NaCl supplemented with oxacillin 40 µg/ml and globomycin concentrations ranging from 10-50 µg/ml or MHB 2% NaCl supplemented with oxacillin 40 µg/ml and 0.6% ethanol (control solvent for globomycin). **(C)** USA300 grown in MHB 2% NaCl supplemented with oxacillin 50 µg/ml, MHB 2% NaCl supplemented with oxacillin 50 µg/ml and globomycin 20 µg/ml or MHB 2% NaCl supplemented with oxacillin 50 µg/ml and 0.4% ethanol (control solvent for globomycin). **(D)** ATCC 43300 grown in MHB 2% NaCl supplemented with oxacillin 30 µg/ml, MHB 2% NaCl supplemented with oxacillin 30 µg/ml and globomycin 30 µg/ml or MHB 2% NaCl supplemented with oxacillin 50 µg/ml and 0.6% ethanol (control solvent for globomycin). The oxacillin and globomycin concentrations used in these experiments were determined empirically for each strain. Cultures were grown in a Tecan Sunrise incubated microplate reader for 24 h at 35°C. OD_{600} was recorded at 15 min intervals and growth curves were plotted in Prism software (GraphPad). The data presented are the average of 3 independent biological replicates, and error bars represent standard deviations.

Fig. 5. Globomycin increases cefotaxime resistance in JE2, USA300 and ATCC43300. (A) JE2 was grown in MHB 2% NaCl supplemented with cefotaxime (CTX) 40 µg/ml, MHB 2% NaCl supplemented with CTX 40 µg/ml and globomycin 30 µg/ml or MHB 2% NaCl supplemented with CTX 40 µg/ml and 0.6% ethanol (control solvent for globomycin). **(B)** USA300 was grown in MHB 2% NaCl supplemented with CTX 40 µg/ml, MHB 2% NaCl supplemented with CTX 40 µg/ml and globomycin 30 µg/ml or MHB 2% NaCl supplemented with CTX 40 µg/ml and 0.6% ethanol (control solvent for globomycin). **(C)** ATCC 43300 was grown in MHB 2% NaCl supplemented with CTX 30 µg/ml, MHB 2% NaCl supplemented with CTX 30 µg/ml and globomycin 30 µg/ml or MHB 2% NaCl supplemented with CTX 30 µg/ml and 0.6% ethanol (control solvent for globomycin). The cefotaxime and globomycin concentrations used in these experiments were determined empirically for each strain. The cultures were grown in a Tecan

583   Sunrise incubated microplate reader for 24 h at 35ºC OD_{600} was recorded at 15 min intervals
584   and growth curves were plotted in Prism software (GraphPad). The data presented are the
585   average of 3 independent biological replicates, and error bars represent standard deviations.
586

587
588
Fig. 6. Mutation of *lgt* increases susceptibility to Congo red. Overnight cultures of JE2,
NE1757 (*lspA*), NE1905 (*lgt*), NE1757/NE1905 (*lgt/lspA*), NE1757 (markerless *lspA* mutation),
NE1757 pLI50, NE1757 p*lspA* and JE2 pLI50 were grown in TSB (without antibiotic selection),
normalised to an OD_{600} of 1 in PBS, serially diluted and spotted onto TSA plates supplemented
with 0.125% Congo red, and the plates were incubated for 24 h at 37°C. A representative
image is shown.
595

Fig. 7. Mutation of *lspA* or *lgt* impacts growth in nutrient limited media but not in complex media. Growth of JE2 wild type, NE1757 (*lspA*::Tn), NE1757 MM (markerless *lspA*::Tn), NE1905 (*lgt*::Tn) and NE1757/NE1905 (markerless *lspA*::Tn / *lgt*::Tn) in Muller Hinton broth **(A)**, Muller Hinton broth with 2% NaCl **(B)**, TSB **(C)**, and chemically-defined medium **(D)**. Growth experiments were performed in 96-well hydrophobic plates in a Tecan Sunrise incubated microplate reader for 24 h at 37°C. OD_{600} was recorded at 15 min intervals and growth curves were plotted in Prism software (GraphPad). The data presented are the average of 3 independent biological replicates, and error bars represent standard deviations.

2. Lgt-mediated addition of diacylgylcerol moiety

A

1. Sec-dependent transport of prolipoprotein to membrane

3. LspA-mediated cleavage of signal peptide

No Lgt-mediated addition of diacylglycerol moiety

B

LspA-mediated cleavage of lipoprotein signal peptide occurs despite lack of diacylglycerol modification

Lgt-mediated addition of diacylgylcerol moiety

C

No LspA-mediated cleavage of the signal peptide; membrane-localised diacylglycerol liporotein is either unprocessed or the signal peptide is cleaved by an alternative signal peptidase

No Lgt-mediated addition of diacylglycerol moiety

D

No LspA-mediated cleavage of the signal peptide; prolipoproteins may be retained in the membrane or released after cleavage of the signal peptidase by an alternative signal paptidase

**Fig. 8. Predicted impacts of *lgt*, *lspA* and *lgt/lspA* mutations on lipoprotein processing in *S.*
***aureus*. (A)** Overview of lipoprotein processing in Gram-positive bacteria. **(B)** The *lgt* mutant
lacks diacylglyceryl transferase activity and is unable to add the diacylglycerol molecule to the
prolipoprotein, thus blocking accumulation of the LspA substrate. The prolipoprotein is likely
to be released from the membrane due to LspA-mediated cleavage of the signal peptide signal
peptide. **(C)** LspA-mediated cleavage of the signal peptide is absent in the *lspA* mutant, but
an alternative signal peptidase may undertake this activity. **(D)** Prolipoproteins are not
processed in mutants lacking Lgt and LspA activity, and may be retained in the membrane or
released after cleavage of the signal peptide by an alternative signal peptidase.

626 **Supplementary Tables and Figures**
627

630 **Supplementary Table S1.** Genome sequence changes in NE1757 (*lspA*::Tn)

Reference Position[1]	Type[2]	Ref.[3]	Allele[4]	Freq.[5]	Annotation
1192002 - 1192003	INS	TA	T-Tn-A	100	*Bursa aurealis* transposon

[1]Reference position: Position in the USA300 FPR3757 genome sequence (NC_007793.1).

[2]Type of mutation: INS, insertion.

[3]Ref: Nucleotide base in the USA300 FPR3757 reference genome.

[4]Allele: Nucleotide base at the same position in NE1757.

[5]Freq.: % Frequency at which INS was found in NE1757 compared to JE2

Table S2. Bacterial strains and plasmids used in this study

Strain/plasmid	Description and resistance	Source/reference
***Escherichia coli* strains**		
HST08	TaKaRa *E. coli* HST08 Premium Electro-Cells	TaKaRa
HST08 pLI50	*E. coli* HST08 pLI50; Amp[R]	This study
HST08 p*lspA*	*E. coli* HST08 p*lspA*; Amp[R]	This study
***Staphylococcus aureus* strains**		
JE2	JE2 (plasmid-cured derivative of strain LAC)	(36)
8325-4	NCTC 8325 derivative cured of prophages, methicillin susceptible, CC8.	(67)
RN4220	RN4220 (Restriction-deficient derivative of 8325-4)	(68)
BHICC	MRSA clinical isolate; SCC*mec* type II; CC8	(39)
ATCC 29213	Quality control strain for susceptibility testing; oxacillin sensitive	*S. aureus* ATCC 29213
ATCC 43300	Quality control strain for susceptibility testing; oxacillin resistant, SCC*mec* Type II	*S. aureus* ATCC 43300
USA300	USA300_FPR3757	(69)
NE1757 (*lspA*)	JE2 *lspA*::Tn; Erm[R]	(36)
NE1905 (*lgt*)	JE2 *lgt*::Tn; Erm[R]	(36)
NE1757 MM	JE2 *lspA*:Tn (truncated, markerless Tn); Erm[S]	This study
NE1757/NE1905	JE2 *lspA* MM *lgt*::Tn; Erm[R]	This study
RN4220 pTnT	RN4220 pTnT	(36)
RN4220 pLI50	RN4220 pLI50; Cam[R]	This study
RN4220 p*lspA*	RN4220 p*lspA*	This study
NE1757 p*lspA*	JE2 *lspA*::Tn p*lspA*; Erm[R] Cam[R]	This study
NE1757 pLI50	JE2 *lspA*::Tn pLI50; Erm[R] Cam[R]	This study
JE2 pLI50	JE2 pLI50; Cam[R]	This study
JE2 *lspA* MM pLI50	JE2 *lspA* MM pLI50; Cam[R]	This study
NE1757 MM p*lspA*	JE2 *lspA* MM p*lspA*; Cam[R]	This study
JE2 *lspA*::Tn #2	JE2 *lspA*::Tn; Erm[R]	This study
JE2 *lspA*::Tn #4	JE2 *lspA*::Tn; Erm[R]	This study
NE788	JE2 *trkA*::Tn; Erm[R]	(36)
NE460	JE2 *atl*::Tn; Erm[R]	(36)
Plasmids		
pTnT	Plasmid to replace Erm[R] marker from transposon mutants with a markerless mutation. Cam[R]	(58)
pLI50	*S. aureus-E. coli* shuttle vector	(70)
p*lspA*	pLI50 carrying the *lspA* gene from JE2	This study

641   **Table S3**. Oligonucleotides used in this study

Target	Name	Primer Sequence (5' – 3')
lspA (infusion primers)	NE1757_INF#3_Fwd	TCGTCTTCAAGAATTTTATGAAGGAGGCTGGGACA
	NE1757_INF#3_Rev	TACCGAGCTCGAATTCAGGCAGCAACTTATCTACACG
Tn-check primers	**Name**	**Primer Sequence (5' – 3')**
lspA	NE1757_fwd	GTTCCAGCCTGCTTTCCTAATT
	NE1757_rev	ACACGCATACCTGTTTGTTCT
lgt	NE1905_fwd	GCATTAACACGGCCGAAGAA
	NE1905_rev	CAACCGTACCAGCTGCAAC

642
643

Supplementary Fig. S1. Complementation of the NE1757 mutant with the _lspA_ gene is
accompanied by a wild type oxacillin resistance phenotype. JE2, NE1757 (_lspA_::Tn), NE1757
pLI50 and NE1757 p_lspA_ were grown on MHA 2% NaCl supplemented with oxacillin 32 µg/ml.
The plate shows 5 biological replicates each for NE1757 pLI50 and NE1757 p_lspA_. ATCC 29213
was included as an oxacillin susceptible control).

Supplementary Fig. S2. Mutation of *lspA* does not impact growth in MHB or TSB media.
Growth of JE2 and NE1757 cultures in **(A)** MHB, **(B)** MHB 2% NaCl, **(C)** TSB and **(D)** TSB
0.5mg/ml oxacillin. MHB cultures were grown in 96-well plates in a Tecan Sunrise incubated
microplate reader for 24 h at 37°C. The OD_{600} was recorded at 15 min intervals and growth
curves were plotted in Prism software (GraphPad). TSB cultures were grown in flasks and
the OD_{600} monitored every 2 h. All data presented are the average of 3 independent biological
replicates, and error bars represent standard deviations.

Supplementary Fig. S3. Population analysis profiling shows that the *lspA* mutant NE1757
exhibits heterogenous resistance to oxacillin. Overnight cultures of JE2, NE1757 and BH1CC
were grown in TSB, adjusted to OD_{600} of 1, serially diluted and plated onto TSA and TSA
supplemented with 0.25, 0.5, 1, 2, 4, 8, 16, 32, 64 and 128 µg/ml oxacillin. CFUs were
enumerated after overnight incubation at 37°C. The data are expressed as CFU/ml at each
oxacillin concentration, plotted using Prism software (GraphPad). Three independent
experiments were performed, and error bars represent standard deviations. BH1CC, which
exhibits homogenous oxacillin resistance, was included as a positive control.
670

671
672
Supplementary Fig. S4. Relative proportions of cell wall muropeptide fractions based on
oligomerization and relative cross-linking efficiency of cell wall muropeptide fractions in
peptidoglycan extracted from JE2 and NE1757. Cells were collected from cultures grown to
exponential phase in MHB or MHB supplemented with oxacillin 3 μg/ml or 32 μg/ml. Each
profile shown is a representative of 3 biological replicates. Significant differences determined
using Students t-test (**P < .01; ***P < .001; ****P<.0001).
679

680
681
Supplementary Fig. S5. Autolytic activity is unaffected by the *lspA* mutation. Triton X-100-
induced autolysis of JE2, NE1757 (*lspA::*Tn) and NE406 (*atl*::Tn, negative control). The strains
were grown to OD_{600} = 0.5 in MHB medium at 37°C, before being washed in cold PBS and
resuspended in 0.1% Triton X-100. The OD_{600} was monitored, and autolysis was expressed as
a percentage of the initial OD_{600}. The experiments were repeated 3 independent times,
plotted using Prism software (GraphPad) and standard deviations are shown.
688

Supplementary Fig. S6. Mutation of *lspA* has no significant impact on salt tolerance in JE2.
Overnight cultures of JE2, NE1757, NE1757 pLI50, NE1757 p*lspA*, JE2 *lspA*::Tn #2 transductant
and NE788 (JE2 *ktr*::Tn, NaCl-sensitive control) were grown in TSB and cell density was
standardised to OD$_{600}$ of 1. Four ml aliquots from 10-fold serial dilutions were spotted onto
TSA supplemented with 2.2 M NaCl and the plates incubated overnight at 37°C. Three
independent experiments were carried out and a representative image of a plate is shown

Supplementary Fig. S7. Globomycin does not increase oxacillin resistance in the *lspA* mutant NE1757. NE1757 was grown in MHB 2% NaCl supplemented with a sub-inhibitory concentration of oxacillin (40 µg/ml) and a range (10 – 50 µg/ml) of globomycin concentrations. The solvent for globomycin, 0.6% ethanol, was included as a control. The cultures were grown in a Tecan Sunrise incubated microplate reader for 24 h at 35°C. OD_{600} was recorded at 15 min intervals and growth curves were plotted in Prism software (GraphPad). The data presented are the average of 3 independent biological replicates, and error bars represent standard deviations.

References

1.   Gründling, A. (2016) Cell Wall Assembly and Physiology. in *Staphylococcus: Genetics and Physiology* (Somerville, G. A. ed.), Caister Academic Press, UK. pp 133-170

2.   Dordel, J., Kim, C., Chung, M., Pardos de la Gandara, M., Holden, M. T., Parkhill, J., de Lencastre, H., Bentley, S. D., and Tomasz, A. (2014) Novel determinants of antibiotic resistance: identification of mutated loci in highly methicillin-resistant subpopulations of methicillin-resistant Staphylococcus aureus. *MBio* **5**, e01000

3.   Pozzi, C., Waters, E. M., Rudkin, J. K., Schaeffer, C. R., Lohan, A. J., Tong, P., Loftus, B. J., Pier, G. B., Fey, P. D., Massey, R. C., and O'Gara, J. P. (2012) Methicillin resistance alters the biofilm phenotype and attenuates virulence in Staphylococcus aureus device-associated infections. *PLoS Pathog* **8**, e1002626

4.   Corrigan, R. M., Abbott, J. C., Burhenne, H., Kaever, V., and Grundling, A. (2011) c-di-AMP is a new second messenger in Staphylococcus aureus with a role in controlling cell size and envelope stress. *PLoS Pathog* **7**, e1002217

5.   Dengler, V., McCallum, N., Kiefer, P., Christen, P., Patrignani, A., Vorholt, J. A., Berger-Bachi, B., and Senn, M. M. (2013) Mutation in the C-di-AMP cyclase dacA affects fitness and resistance of methicillin resistant Staphylococcus aureus. *PLoS One* **8**, e73512

6.   Panchal, V. V., Griffiths, C., Mosaei, H., Bilyk, B., Sutton, J. A. F., Carnell, O. T., Hornby, D. P., Green, J., Hobbs, J. K., Kelley, W. L., Zenkin, N., and Foster, S. J. (2020) Evolving MRSA: High-level beta-lactam resistance in Staphylococcus aureus is associated with RNA Polymerase alterations and fine tuning of gene expression. *PLoS Pathog* **16**, e1008672

7.   Baek, K. T., Grundling, A., Mogensen, R. G., Thogersen, L., Petersen, A., Paulander, W., and Frees, D. (2014) beta-Lactam resistance in methicillin-resistant Staphylococcus aureus USA300 is increased by inactivation of the ClpXP protease. *Antimicrob Agents Chemother* **58**, 4593-4603

8.   Jensen, C., Baek, K. T., Gallay, C., Thalso-Madsen, I., Xu, L., Jousselin, A., Ruiz Torrubia, F., Paulander, W., Pereira, A. R., Veening, J. W., Pinho, M. G., and Frees, D. (2019) The ClpX chaperone controls autolytic splitting of Staphylococcus aureus daughter cells, but is bypassed by beta-lactam antibiotics or inhibitors of WTA biosynthesis. *PLoS Pathog* **15**, e1008044

9.   Giulieri, S. G., Guerillot, R., Kwong, J. C., Monk, I. R., Hayes, A. S., Daniel, D., Baines, S., Sherry, N. L., Holmes, N. E., Ward, P., Gao, W., Seemann, T., Stinear, T. P., and Howden, B. P. (2020) Comprehensive Genomic Investigation of Adaptive Mutations Driving the Low-Level Oxacillin Resistance Phenotype in Staphylococcus aureus. *mBio* **11**

10.   Shahmirzadi, S. V., Nguyen, M. T., and Gotz, F. (2016) Evaluation of Staphylococcus aureus Lipoproteins: Role in Nutritional Acquisition and Pathogenicity. *Front Microbiol* **7**, 1404

11.   Dartois, V., Djavakhishvili, T., and Hoch, J. A. (1997) KapB is a lipoprotein required for KinB signal transduction and activation of the phosphorelay to sporulation in Bacillus subtilis. *Mol Microbiol* **26**, 1097-1108

12.   Bengtsson, J., Tjalsma, H., Rivolta, C., and Hederstedt, L. (1999) Subunit II of Bacillus subtilis cytochrome c oxidase is a lipoprotein. *J Bacteriol* **181**, 685-688

13.   Jacobs, M., Andersen, J. B., Kontinen, V., and Sarvas, M. (1993) Bacillus subtilis PrsA is required in vivo as an extracytoplasmic chaperone for secretion of active enzymes synthesized either with or without pro-sequences. *Mol Microbiol* **8**, 957-966

14. Mei, J. M., Nourbakhsh, F., Ford, C. W., and Holden, D. W. (1997) Identification of
Staphylococcus aureus virulence genes in a murine model of bacteraemia using
signature-tagged mutagenesis. *Mol Microbiol* **26**, 399-407

15. Nielsen, J. B., and Lampen, J. O. (1982) Membrane-bound penicillinases in Gram-
positive bacteria. *J Biol Chem* **257**, 4490-4495

16. Nguyen, M. T., Kraft, B., Yu, W., Demircioglu, D. D., Hertlein, T., Burian, M., Schmaler,
M., Boller, K., Bekeredjian-Ding, I., Ohlsen, K., Schittek, B., and Gotz, F. (2015) The
nuSaalpha Specific Lipoprotein Like Cluster (lpl) of S. aureus USA300 Contributes to
Immune Stimulation and Invasion in Human Cells. *PLoS Pathog* **11**, e1004984

17. Kovacs-Simon, A., Titball, R. W., and Michell, S. L. (2011) Lipoproteins of bacterial
pathogens. *Infect Immun* **79**, 548-561

18. Nguyen, M. T., Matsuo, M., Niemann, S., Herrmann, M., and Gotz, F. (2020)
Lipoproteins in Gram-Positive Bacteria: Abundance, Function, Fitness. *Front Microbiol*
11, 582582

19. Schmollinger, M., Fischer, I., Nerz, C., Pinkenburg, S., Gotz, F., Kaufmann, M., Lange,
K. J., Reuter, R., Rosenstiel, W., and Zell, A. (2004) ParSeq: searching motifs with
structural and biochemical properties. *Bioinformatics* **20**, 1459-1461

20. Inouye, S., Wang, S., Sekizawa, J., Halegoua, S., and Inouye, M. (1977) Amino acid
sequence for the peptide extension on the prolipoprotein of the Escherichia coli outer
membrane. *Proc Natl Acad Sci U S A* **74**, 1004-1008

21. Buddelmeijer, N. (2015) The molecular mechanism of bacterial lipoprotein
modification--how, when and why? *FEMS Microbiol Rev* **39**, 246-261

22. Hayashi, S., and Wu, H. C. (1990) Lipoproteins in bacteria. *J Bioenerg Biomembr* **22**,
451-471

23. Braun, V., and Wu, H. C. (1994) Lipoproteins, structure, function, biosynthesis and
model for protein export. in *New Comprehensive Biochemistry* (Ghuysen, J.-M., and
Hakenbeck, R. eds.), Elsevier. pp 319-341

24. Sankaran, K., and Wu, H. C. (1994) Lipid modification of bacterial prolipoprotein.
Transfer of diacylglyceryl moiety from phosphatidylglycerol. *J Biol Chem* **269**, 19701-
19706

25. Ichihara, S., Hussain, M., and Mizushima, S. (1982) Mechanism of export of outer
membrane lipoproteins through the cytoplasmic membrane in Escherichia coli.
Binding of lipoprotein precursors to the peptidoglycan layer. *J Biol Chem* **257**, 495-500

26. Sankaran, K., Gupta, S. D., and Wu, H. C. (1995) Modification of bacterial lipoproteins.
Methods Enzymol **250**, 683-697

27. Tschumi, A., Nai, C., Auchli, Y., Hunziker, P., Gehrig, P., Keller, P., Grau, T., and Sander,
P. (2009) Identification of apolipoprotein N-acyltransferase (Lnt) in mycobacteria. *J
Biol Chem* **284**, 27146-27156

28. Asanuma, M., Kurokawa, K., Ichikawa, R., Ryu, K. H., Chae, J. H., Dohmae, N., Lee, B.
L., and Nakayama, H. (2011) Structural evidence of alpha-aminoacylated lipoproteins
of Staphylococcus aureus. *Febs J* **278**, 716-728

29. Gardiner, J. H. t., Komazin, G., Matsuo, M., Cole, K., Gotz, F., and Meredith, T. C. (2020)
Lipoprotein N-Acylation in Staphylococcus aureus Is Catalyzed by a Two-Component
Acyl Transferase System. *mBio* **11**

30. An, F. Y., Sulavik, M. C., and Clewell, D. B. (1999) Identification and characterization of
a determinant (eep) on the Enterococcus faecalis chromosome that is involved in
production of the peptide sex pheromone cAD1. *J Bacteriol* **181**, 5915-5921

31.   Varahan, S., Harms, N., Gilmore, M. S., Tomich, J. M., and Hancock, L. E. (2014) An ABC
805   transporter is required for secretion of peptide sex pheromones in Enterococcus
806   faecalis. *mBio* **5**, e01726-01714
807   32.   Schilcher, K., Caesar, L. K., Cech, N. B., and Horswill, A. R. (2020) Processing, Export,
808   and Identification of Novel Linear Peptides from Staphylococcus aureus. *mBio* **11**
809   33.   Gan, K., Gupta, S. D., Sankaran, K., Schmid, M. B., and Wu, H. C. (1993) Isolation and
810   characterization of a temperature-sensitive mutant of Salmonella typhimurium
811   defective in prolipoprotein modification. *J Biol Chem* **268**, 16544-16550
812   34.   Leskela, S., Wahlstrom, E., Kontinen, V. P., and Sarvas, M. (1999) Lipid modification of
813   prelipoproteins is dispensable for growth but essential for efficient protein secretion
814   in Bacillus subtilis: characterization of the Lgt gene. *Mol Microbiol* **31**, 1075-1085
815   35.   Stoll, H., Dengjel, J., Nerz, C., and Gotz, F. (2005) Staphylococcus aureus deficient in
816   lipidation of prelipoproteins is attenuated in growth and immune activation. *Infect*
817   *Immun* **73**, 2411-2423
818   36.   Fey, P. D., Endres, J. L., Yajjala, V. K., Widhelm, T. J., Boissy, R. J., Bose, J. L., and Bayles,
819   K. W. (2013) A genetic resource for rapid and comprehensive phenotype screening of
820   nonessential Staphylococcus aureus genes. *MBio* **4**, e00537-00512
821   37.   Gohring, N., Fedtke, I., Xia, G., Jorge, A. M., Pinho, M. G., Bertsche, U., and Peschel, A.
822   (2011) New role of the disulfide stress effector YjbH in beta-lactam susceptibility of
823   Staphylococcus aureus. *Antimicrob Agents Chemother* **55**, 5452-5458
824   38.   Chan, Y. G., Frankel, M. B., Missiakas, D., and Schneewind, O. (2016) SagB
825   Glucosaminidase Is a Determinant of Staphylococcus aureus Glycan Chain Length,
826   Antibiotic Susceptibility, and Protein Secretion. *J Bacteriol* **198**, 1123-1136
827   39.   Gallagher, L. A., Shears, R. K., Fingleton, C., Alvarez, L., Waters, E. M., Clarke, J., Bricio-
828   Moreno, L., Campbell, C., Yadav, A. K., Razvi, F., O'Neill, E., O'Neill, A. J., Cava, F., Fey,
829   P. D., Kadioglu, A., and O'Gara, J. P. (2020) Impaired Alanine Transport or Exposure to
830   d-Cycloserine Increases the Susceptibility of MRSA to beta-lactam Antibiotics. *J Infect*
831   *Dis* **221**, 1000-1016
832   40.   Roch, M., Lelong, E., Panasenko, O. O., Sierra, R., Renzoni, A., and Kelley, W. L. (2019)
833   Thermosensitive PBP2a requires extracellular folding factors PrsA and HtrA1 for
834   Staphylococcus aureus MRSA beta-lactam resistance. *Commun Biol* **2**, 417
835   41.   Inukai, M., Takeuchi, M., Shimizu, K., and Arai, M. (1978) Mechanism of action of
836   globomycin. *J Antibiot (Tokyo)* **31**, 1203-1205
837   42.   Inukai, M., Nakajima, M., Osawa, M., Haneishi, T., and Arai, M. (1978) Globomycin, a
838   new peptide antibiotic with spheroplast-forming activity. II. Isolation and physico-
839   chemical and biological characterization. *J Antibiot (Tokyo)* **31**, 421-425
840   43.   Vogeley, L., El Arnaout, T., Bailey, J., Stansfeld, P. J., Boland, C., and Caffrey, M. (2016)
841   Structural basis of lipoprotein signal peptidase II action and inhibition by the antibiotic
842   globomycin. *Science* **351**, 876-880
843   44.   Yakushi, T., Tajima, T., Matsuyama, S., and Tokuda, H. (1997) Lethality of the covalent
844   linkage between mislocalized major outer membrane lipoprotein and the
845   peptidoglycan of Escherichia coli. *J Bacteriol* **179**, 2857-2862
846   45.   Garland, K., Pantua, H., Braun, M. G., Burdick, D. J., Castanedo, G. M., Chen, Y. C.,
847   Cheng, Y. X., Cheong, J., Daniels, B., Deshmukh, G., Fu, Y., Gibbons, P., Gloor, S. L., Hua,
848   R., Labadie, S., Liu, X., Pastor, R., Stivala, C., Xu, M., Xu, Y., Zheng, H., Kapadia, S. B.,
849   and Hanan, E. J. (2020) Optimization of globomycin analogs as novel gram-negative
850   antibiotics. *Bioorg Med Chem Lett* **30**, 127419

46.   Vickery, C. R., Wood, B. M., Morris, H. G., Losick, R., and Walker, S. (2018)
Reconstitution of Staphylococcus aureus Lipoteichoic Acid Synthase Activity Identifies
Congo Red as a Selective Inhibitor. *J Am Chem Soc* **140**, 876-879

47.   Parvez, M. A., Shibata, H., Nakano, T., Niimi, S., Fujii, N., Arakaki, N., and Higuti, T.
(2008) No relationship exists between PBP 2a amounts expressed in different MRSA
strains obtained clinically and their beta-lactam MIC values. *J Med Invest* **55**, 246-253

48.   Hartman, B. J., and Tomasz, A. (1986) Expression of methicillin resistance in
heterogeneous strains of Staphylococcus aureus. *Antimicrob Agents Chemother* **29**,
85-92

49.   Mikkelsen, K., Sirisarn, W., Alharbi, O., Alharbi, M., Liu, H., Nohr-Meldgaard, K., Mayer,
K., Vestergaard, M., Gallagher, L. M., Derrick, J. P., McBain, A. J., Biboy, J., Vollmer, W.,
O'Gara, J. P., Grunert, T., Ingmer, H., and Xia, G. (2021) The Novel Membrane-
Associated Auxiliary Factors AuxA and AuxB Modulate beta-lactam Resistance in
MRSA by stabilizing Lipoteichoic Acids. *Int J Antimicrob Agents*, 106283

50.   Campbell, J., Singh, A. K., Santa Maria, J. P., Jr., Kim, Y., Brown, S., Swoboda, J. G.,
Mylonakis, E., Wilkinson, B. J., and Walker, S. (2011) Synthetic lethal compound
combinations reveal a fundamental connection between wall teichoic acid and
peptidoglycan biosyntheses in Staphylococcus aureus. *ACS Chem Biol* **6**, 106-116

51.   Brown, S., Xia, G., Luhachack, L. G., Campbell, J., Meredith, T. C., Chen, C., Winstel, V.,
Gekeler, C., Irazoqui, J. E., Peschel, A., and Walker, S. (2012) Methicillin resistance in
Staphylococcus aureus requires glycosylated wall teichoic acids. *Proc Natl Acad Sci U*
S A **109**, 18909-18914

52.   Memmi, G., Filipe, S. R., Pinho, M. G., Fu, Z., and Cheung, A. (2008) Staphylococcus
aureus PBP4 is essential for beta-lactam resistance in community-acquired methicillin-
resistant strains. *Antimicrob Agents Chemother* **52**, 3955-3966

53.   Lee, S. H., Jarantow, L. W., Wang, H., Sillaots, S., Cheng, H., Meredith, T. C., Thompson,
J., and Roemer, T. (2011) Antagonism of chemical genetic interaction networks
resensitize MRSA to beta-lactam antibiotics. *Chem Biol* **18**, 1379-1389

54.   Gardete, S., Wu, S. W., Gill, S., and Tomasz, A. (2006) Role of VraSR in antibiotic
resistance and antibiotic-induced stress response in Staphylococcus aureus.
Antimicrob Agents Chemother **50**, 3424-3434

55.   Baumgartner, M., Karst, U., Gerstel, B., Loessner, M., Wehland, J., and Jansch, L. (2007)
Inactivation of Lgt allows systematic characterization of lipoproteins from Listeria
monocytogenes. *J Bacteriol* **189**, 313-324

56.   Henneke, P., Dramsi, S., Mancuso, G., Chraibi, K., Pellegrini, E., Theilacker, C., Hubner,
J., Santos-Sierra, S., Teti, G., Golenbock, D. T., Poyart, C., and Trieu-Cuot, P. (2008)
Lipoproteins are critical TLR2 activating toxins in group B streptococcal sepsis. *J*
Immunol **180**, 6149-6158

57.   Halsey, C. R., Lei, S., Wax, J. K., Lehman, M. K., Nuxoll, A. S., Steinke, L., Sadykov, M.,
Powers, R., and Fey, P. D. (2017) Amino Acid Catabolism in Staphylococcus aureus and
the Function of Carbon Catabolite Repression. *mBio* **8**

58.   Bose, J. L., Fey, P. D., and Bayles, K. W. (2013) Genetic tools to enhance the study of
gene function and regulation in Staphylococcus aureus. *Appl Environ Microbiol* **79**,
2218-2224

59.   Karinou, E., Schuster, C. F., Pazos, M., Vollmer, W., and Grundling, A. (2019)
Inactivation of the Monofunctional Peptidoglycan Glycosyltransferase SgtB Allows
Staphylococcus aureus To Survive in the Absence of Lipoteichoic Acid. *J Bacteriol* **201**

60. Tomasz, A., Nachman, S., and Leaf, H. (1991) Stable classes of phenotypic expression
in methicillin-resistant clinical isolates of staphylococci. *Antimicrob Agents Chemother*
35, 124-129
61. Griffiths, J. M., and O'Neill, A. J. (2012) Loss of function of the gdpP protein leads to
joint beta-lactam/glycopeptide tolerance in Staphylococcus aureus. *Antimicrob*
Agents Chemother **56**, 579-581
62. Alvarez, L., Hernandez, S. B., de Pedro, M. A., and Cava, F. (2016) Ultra-Sensitive, High-
Resolution Liquid Chromatography Methods for the High-Throughput Quantitative
Analysis of Bacterial Cell Wall Chemistry and Structure. *Methods Mol Biol* **1440**, 11-27
63. de Jonge, B. L., Chang, Y. S., Gage, D., and Tomasz, A. (1992) Peptidoglycan
composition of a highly methicillin-resistant Staphylococcus aureus strain. The role of
penicillin binding protein 2A. *J Biol Chem* **267**, 11248-11254
64. De Jonge, B. L., Gage, D., and Xu, N. (2002) The carboxyl terminus of peptidoglycan
stem peptides is a determinant for methicillin resistance in Staphylococcus aureus.
Antimicrob Agents Chemother **46**, 3151-3155
65. Kuhner, D., Stahl, M., Demircioglu, D. D., and Bertsche, U. (2014) From cells to
muropeptide structures in 24 h: peptidoglycan mapping by UPLC-MS. *Sci Rep* **4**, 7494
66. Boneca, I. G., Xu, N., Gage, D. A., de Jonge, B. L., and Tomasz, A. (1997) Structural
characterization of an abnormally cross-linked muropeptide dimer that is
accumulated in the peptidoglycan of methicillin- and cefotaxime-resistant mutants of
Staphylococcus aureus. *J Biol Chem* **272**, 29053-29059
67. Horsburgh, M. J., Aish, J. L., White, I. J., Shaw, L., Lithgow, J. K., and Foster, S. J. (2002)
sigmaB modulates virulence determinant expression and stress resistance:
characterization of a functional rsbU strain derived from Staphylococcus aureus 8325-
4. *J Bacteriol* **184**, 5457-5467
68. Kreiswirth, B. N., Lofdahl, S., Betley, M. J., O'Reilly, M., Schlievert, P. M., Bergdoll, M.
S., and Novick, R. P. (1983) The toxic shock syndrome exotoxin structural gene is not
detectably transmitted by a prophage. *Nature* **305**, 709-712
69. Diep, B. A., Gill, S. R., Chang, R. F., Phan, T. H., Chen, J. H., Davidson, M. G., Lin, F., Lin,
J., Carleton, H. A., Mongodin, E. F., Sensabaugh, G. F., and Perdreau-Remington, F.
(2006) Complete genome sequence of USA300, an epidemic clone of community-
acquired meticillin-resistant Staphylococcus aureus. *Lancet* **367**, 731-739
70. Lee, C. Y., Buranen, S. L., and Ye, Z. H. (1991) Construction of single-copy integration
vectors for Staphylococcus aureus. *Gene* **103**, 101-105
932