Sensitive visualization of SARS-CoV-2 RNA with

CoronaFISH

Elena Rensen[1,$], Stefano Pietropaoli[2,$], Florian Mueller[1,$,#], Christian Weber[1], Sylvie Souquere[3],

Pierre Isnard[4], Marion Rabant[4], Jean-Baptiste Gibier[5], Etienne Simon-Loriere[6], Marie-Anne

Rameix-Welti[7,8], Gérard Pierron[9,#], Giovanna Barba-Spaeth[2,#], Christophe Zimmer[1,#]

[1]Imaging and Modeling Unit, Institut Pasteur, 25 rue du Docteur Roux, 75015 Paris, France

[2]Structural Virology Unit, Institut Pasteur, 25 rue du Docteur Roux, 75015 Paris, France

[3]Gustave Roussy, AMMICA UMS-3655, 94805 Villejuif, France

[4]Anatomie et Cytologie Pathologiques, CHU Necker-Enfants Malades, 149 rue de Sèvres,

75015 Paris, France

[5]Service d'Anatomo- pathologie, Centre de Biologie Pathologie, CHU Lille, Lille, France

[6]Evolutionary genomics of RNA viruses, Institut Pasteur, 25 rue du Docteur Roux, 75015 Paris,

France

[7]Université Paris-Saclay, INSERM, Université de Versailles St. Quentin, UMR 1173 (2I),

Montigny-le-Bretonneux 78180, France.

[8]AP-HP, Université Paris Saclay, Hôpital Ambroise Paré, Laboratoire de Microbiologie,

Boulogne-Billancourt 92104, France

[9]Gustave Roussy, CNRS-UMR 9196, 94805 Villejuif, France

[$] Equal contribution, [#] Corresponding authors

Abstract

The current COVID-19 pandemic is caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The positive-sense single-stranded RNA virus contains a single linear RNA segment that serves as a template for transcription and replication, leading to the synthesis of positive and negative-stranded viral RNA (vRNA) in infected cells. Tools to visualize viral RNA directly in infected cells are critical to analyze its replication cycle, screen for therapeutic molecules or study infections in human tissue. Here, we report the design, validation and initial application of fluorescence *in situ* hybridization (FISH) probes to visualize positive or negative RNA of SARS-CoV-2 (CoronaFISH). We demonstrate sensitive visualization of vRNA in African green monkey and several human cell lines, in patient samples and human tissue. We further demonstrate the adaptation of CoronaFISH probes to electron microscopy (EM). We provide all required oligonucleotide sequences, source code to design the probes, and a detailed protocol. We hope that CoronaFISH will complement existing techniques for research on SARS-CoV-2 biology and COVID-19 pathophysiology, drug screening and diagnostics.

Introduction

Coronavirus disease (COVID-19) emerged by the end of 2019 in Wuhan, China, and led to more than 100 million infections and over 2.1 million deaths (Johns Hopkins University Dashboard). Its causative agent, Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), is an enveloped, positive-sense, single-stranded RNA virus. Upon infection, viral replication occurs in the host cell's cytoplasm, which is massively reorganized (V'kovski *et al*, 2020). The genomic positive-strand viral RNA serves as a template for transcription and replication. The virus synthesizes its own RNA-dependent RNA polymerase (RdRP) to generate negative-sense RNA replication intermediates. This negative strand acts as template for replication of new full-length positive-stranded RNA genomes and for transcription of several smaller, subgenomic positive-stranded RNAs (sgRNAs). These sgRNAs are then used to synthesize all other viral proteins in spatially confined replication complexes. Mature virions are exocytosed and released from the infected host cell. Despite recent progress, many aspects of the SARS-CoV-2 viral replication cycle, including the subcellular location of viral RNA synthesis, are still not fully understood and under active investigation (V'kovski *et al*, 2020).

Several established techniques allow studying SARS-CoV-2 and its interaction with its host. Immunofluorescence (IF) permits the visualization of viral and host proteins in the spatial context of a single cell. However, the development of specific antibodies against novel viruses is time- and cost-intensive, especially if specificity over other closely related viruses is required. Further, the presence in cells of structural viral proteins, such as the Spike protein, does not necessarily imply active viral replication (Cheung *et al*, 2005; Tang *et al*, 2020) and their subcellular localization may not reflect that of the vRNA strands. Other molecular methods, such as RT-PCR, provide an accurate, quantitative readout of viral load and replication dynamics, but are bulk measurements over large cell populations that mask variability between cells and

provide no information about the subcellular localization of the virus. RNAseq permits a

complete view of the transcriptome of both the host and virus, including in single cells, albeit

again without spatial information (Kim *et al*, 2020; Bost *et al*, 2020).

Unlike immunostaining, PCR or sequencing methods, fluorescence *in situ* hybridization (FISH)

offers the capacity to directly and specifically visualize viral RNA in single cells (Raj *et al*, 2008;

Itzkovitz & van Oudenaarden, 2011; Mueller *et al*, 2013). In RNA-FISH, single RNA molecules

are typically targeted with 10–50 fluorescently labeled probes consisting of short (20-30

nucleotides), custom synthesized oligonucleotides with bioinformatically designed sequences.

Individual RNAs are subsequently visible as bright, diffraction-limited spots under a microscope,

and can be detected with appropriate image analysis methods (Raj *et al*, 2008; Mueller *et al*,

2013). We recently introduced smiFISH (Tsanov *et al*, 2016), an inexpensive variant of this

approach that has been used in biological samples ranging from single-cell organisms such as

bacteria and yeast to whole tissue sections and organs (Trcek *et al*, 2012; Wu *et al*, 2018;

Wang, 2019; Chen *et al*, 2016; Skinner *et al*, 2013), and is ideally suited to visualize RNA

viruses and study their subcellular localization and kinetics in host cells (King *et al*, 2018;

Rensen *et al*, 2020).

In this study, we designed and validated smiFISH probes against the positive and negative RNA

strands of SARS-CoV-2 (CoronaFISH). We demonstrate highly specific viral detection in cell

culture, in patient isolates, and in tissue samples. We further demonstrate the flexibility of these

probes by adapting them for electron microscopy *in situ* hybridization (EM-ISH). CoronaFISH

provides a flexible, cost-efficient and versatile platform for studying SARS-CoV-2 replication at

the level of single cells in culture or in tissue, and can potentially be employed for drug

screening and diagnosis.

Results

Design of probes specific for SARS-CoV-2

Our RNA-FISH approach employs two types of bioinformatically designed DNA oligonucleotides (oligos) (Tsanov *et al*, 2016): (i) unlabeled primary oligos consisting of two parts: a specific sequence complementary to a selected subregion of the target RNA and a readout sequence that is identical among all primary oligos (FLAP sequence), (ii) a fluorescently labeled secondary oligo complementary to the FLAP sequence, allowing visualization by light microscopy. These oligos are hybridized to each other *in vitro* before their use for cellular imaging (**Fig 1a**).

A cell infected by SARS-CoV-2 can contain the incoming positive strand, the negative-strand replication intermediate, as well as replicated full-length and sub-genomic positive-strand RNA molecules (**Fig 1b**)**.** We designed two sets of 96 probes, one against the positive strand, and one against the negative strand of SARS-CoV-2 (**Fig 1c**). For more details on the probe design workflow, see the Methods section and the source code (https://github.com/muellerflorian/corona-fish). In brief, we identified an initial list of more than 600 potential probe sequences with our previously described method Oligostan (Tsanov *et al*, 2016). We then further screened these probes to be robust to known genomic variations of SARS-CoV-2 (as of April 2020), while removing probes with affinity to other known β-coronaviruses or viruses frequently causing similar respiratory diseases in human such as Influenza. Lastly, we removed probes overlapping with the transcriptome of several relevant host organisms (human, mouse, African green monkey, hamster and ferret). To establish the final list of 96 probes (**Fig 1c**), we chose probes targeting regions with the highest NGS coverage. The complete list of probe sequences is provided in **Supplementary Table S1**.

Visualization of SARS-CoV-2 in Vero cells with CoronaFISH

To test our probes, we first used Vero cells (African green monkey), which are known to be permissive for SARS-CoV-2 replication (Takayama, 2020) (see Methods). We processed the samples for FISH following the protocol detailed in **Supplementary Note 1** and imaged cells 18h post infection (p.i.) with a multiplicity of infection (MOI) of 0.2, as well as uninfected control cells. The positive-strand and negative-strand probe-sets were both labeled with the fluorophore Cy3 and imaged separately in distinct experiments.

In uninfected samples, cells displayed only weak and diffuse fluorescent signal when labeled with either probe-set, consistent with unspecific background labeling common in RNA-FISH (Tsanov *et al*, 2016), and occasional bright spots could be observed, mostly located outside cells, presumably due to unspecific probe aggregation (**Fig 1d-e** and **S1a**). By contrast, in infected samples, a large proportion of cells showed very strong and localized signal in large regions of the cytoplasm (**Fig 1d-e** and **S1a**). Quantification of the fluorescent intensities per cell (see Methods) indicated that 26% of infected cells labeled for the positive strand (n=242) had intensities exceeding a threshold that excluded >99% of uninfected cells (n=83) (**Fig 1e**). The fluorescence intensity in these cells was on average 23 times higher than this threshold (s.d. 27) (**Fig 1e**). For the negative strands, we counted 5% of Vero cells with intensities above the similarly defined threshold in infected samples (n=307), vs. <1% for uninfected cells (n=103), with on average 2.5 times higher intensities (s.d. 2.2).

The fluorescent signal for the positive-strand was remarkably strong compared to the uninfected control images (**Fig 1e**). This is consistent with an extremely high per-cell viral yield, which has previously been reported for Vero cells (Ogando *et al*, 2020). The RNA signal was perinuclear and restricted to the cytoplasm, consistent with cytoplasmic replication, as previously reported

for other *Coronaviridae* and recently for SARS-CoV-2 (Snijder *et al*, 2020; Fung & Liu, 2019;

Stertz *et al*, 2007; Klein *et al*, 2020). Interestingly, we observed bright foci of different sizes and

intensities, some of which displayed hollow structures reminiscent of the replication organelles

(RO) or double-membrane vesicle (DMV) structures described previously (Ogando *et al*, 2020;

Klein *et al*, 2020). The signal of the negative strand was substantially dimmer than for the

positive-strand (**Fig 1e**), in agreement with previous reports that the replication intermediate

negative strand is less abundant (Wolff *et al*, 2020), but potentially also reflecting diminished

labeling efficiency of this probe set. The negative-strand RNA likewise forms clusters of different

sizes and intensities in the proximity of the nucleus (**Fig 1d, Fig EV1a**).

We next wanted to demonstrate the ability of CoronaFISH to simultaneously visualize two RNA

species with dual-color imaging (Tsanov *et al*, 2016). For this purpose, we used different FLAP

sequences on the primary probes set (FLAP-X for positive-strand, and FLAP-Y for negative-

strand probes), and labeled them with spectrally different fluophores (Atto488 and Cy5). We

then imaged infected Vero cells at 18 h p.i. These images clearly show the presence of both

positive and negative RNA strands in the same cells and the same subcellular regions, although

the relative abundance of both strands appeared to vary from cell to cell and colocalization was

only partial (**Fig 1f, Fig EV1b**). The ability to visualize positive and negative RNA together

opens the door to analyzing the intracellular organization of viral transcription and replication

dynamics.

Our data thus show that CoronaFISH probes can sensitively and specifically detect the positive

and negative strands of SARS-CoV-2 in infected Vero cells.

Applicability of CoronaFISH to drug screening

We next asked if CoronaFISH can be used to test pharmacological treatments against SARS-CoV-2. To this end, we treated SARS-CoV-2 infected Vero cells with Remdesivir, an adenosine nucleoside triphosphate analog that reduces viral replication *in vitro* by inhibiting the RdRP (Gordon *et al*, 2020). We first established an inhibition curve that showed, in our system, an IC50 concentration of 2.8 µM for Remdesivir (**Fig EV1c**). We then performed FISH against the positive strand using Atto-488 labeled probes in infected cells left untreated or treated with 10 µM of Remdesivir. In untreated samples, 26% of cells (n=143) displayed a strong fluorescence signal (above the 99% threshold defined for n= 273 uninfected cells), as above, whereas Remdesivir-treated cells only showed background signal similar to uninfected cells (0% of n=179 cells above threshold) (**Fig 1g,e**). These data suggest that our CoronaFISH probes can be used to test molecules for their ability to inhibit SARS-CoV-2 replication.

Detection of SARS-CoV-2 RNA in human cell lines

We next tested the probes in several human cell lines known to be permissive to SARS-CoV-2 (Takayama, 2020): Caco-2 (human intestinal epithelial cells), Huh7 (hepatocyte-derived carcinoma cells), and Calu-3 (human lung cancer cells). Each cell line was infected with MOI 0.2 and fixed at 36 h p.i. Titration on the supernatants of the infected cells revealed vastly different replication efficiencies as tested by focus forming assay (**Table 1**). Caco2 and Huh7 cells showed rather low virus levels in the same order of magnitude as Vero cells ($2\text{-}5\text{x}10^3$ FFU/ml), while Calu3 showed two orders of magnitude higher levels of viral RNA ($2\text{x}10^5$ FFU/ml).

We then performed FISH against the positive RNA strand. As for the Vero cells above, the uninfected controls of all human cell types showed only a weak background signal, while a

strong signal could be detected in the infected cells (**Fig 2a-c**). However, depending on the cell-

type, the number of infected cells, as well as the amount and cellular localization of vRNA

detected by CoronaFISH were very different, consistent with different replication dynamics of

SARS-CoV-2 in these cell lines. In Caco-2 cells, only a minority of cells appeared infected (19%

of n=1,752 cells above intensity threshold defined from n=229 uninfected cells as above),

displaying rather low-intensity values, indicating a low abundance of positive-strand viral RNA

(6.2-fold above threshold, s.d. 6.8). Huh7 cells were more permissive for viral infection

manifesting in more cells displaying vRNA (74% of n=546 above threshold defined on n=246

uninfected cells) and higher RNA signal intensity (11-fold above threshold, s.d. 10). Lastly, all

Calu cells were infected (100% of n=773 cells above threshold defined on n=479 uninfected

cells) and the signal intensity was higher than in Caco-2 and Huh7 cells (28-fold above

threshold, s.d. 14). These data show that CoronaFISH probes can also be used in cell lines of

human origin with similar detection performance as in Vero cells.

Detection of SARS-CoV-2 RNA in human tissue

Next, we aimed to test if our approach can be used to detect SARS-CoV-2 RNA in samples

from patients. The major histopathological finding of the pulmonary system of post mortem

COVID-19 patients with acute respiratory distress syndrome (ARDS) is diffuse alveolar damage

in the acute or organizing phases. Lung tissue examination mainly shows evidence of intra-

alveolar hemorrhage and edema with fibrin and hyaline membranes developed on alveolar walls

at the acute phase and proliferative and fibrotic lesions in the alveolar septal walls at the

organizing phase (Hanley *et al*, 2020; Bradley *et al*, 2020). However, these lesions are common

to multiple forms of ARDS and not specific to COVID-19 and do not shed light on the underlying

etiology. CoronaFISH therefore has the potential to specifically detect SARS-CoV-2 infection

and characterize viral tropism within the tissue.

As a negative control, we imaged a tissue section sample obtained from a deceased adult patient with diffuse alveolar damage from ARDS prior to the COVID-19 pandemic (see Methods). Histological analysis showed diffuse alveolar damage with an important alveolar hemorrhage, an intra-alveolar and interstitial edema with polymorphic inflammatory infiltrate and the presence of early hyaline membrane adjacent to alveolar walls (**Fig EV2a**). When staining this sample with the positive-strand CoronaFISH probes, no strong fluorescent signal was detected, despite the presence of alveolar damage comparable to patients suffering from COVID-19 (**Fig EV2 b-c**).

As a positive control, we used samples from a COVID-19 patient who died 3 days after admission to the intensive care unit. Histological analysis showed diffuse alveolar damage at the organizing phase with intra-alveolar hyaline membranes and fibrin together with interstitial fibrotic lesions with polymorphic inflammatory cell infiltrate of alveolar walls (**Fig EV2d**). Whereas histology by itself is not sufficient to diagnose lung tissue infection, CoronaFISH revealed infected cells with large cytoplasmic RNA aggregates (**Fig 3a**), illustrating that viral presence can also be detected in tissue sections. Because of the extensive destruction of tissue architecture, determining the affected area of the parenchyma is challenging. However, the cell types (e.g. macrophages or type 2 pneumocytes) infected by SARS-CoV-2 could be revealed by combining CoronaFISH with immunostaining.

Detection of SARS-CoV-2 RNA in nasal swabs

Motivated by the previous results, we next attempted to detect viral presence in human samples used for COVID-19 diagnostics. Nasal swabs were collected from patients with respiratory symptoms as part of routine care at the Hospital Ambroise Paré (Paris) (**Fig 3b**). Samples were screened for the presence of SARS-CoV-2 with RT-PCR. The remainder of the sample not used for diagnostics was deposited on coverglass and we performed CoronaFISH against the

positive-strand RNA of SARS-CoV-2. Negative samples (RT-PCR Ct value above detection

limit) gave no specific signal, but some areas showed substantially higher background than in

the cultured cell lines (**Fig EV3**). This background was rather homogenous, and thus distinct

from the RNA signal seen so far in infected cells above. However, in a RT-PCR positive sample

(Ct value = 21), we detected a strong RNA-FISH signal in a subset of cells (**Fig 3c**) when

staining for the positive strand. Therefore, although a systematic analysis on many more

samples will be required to assess specificity and specificity, our CoronaFISH probes may allow

the detection of SARS-CoV-2 in patient-derived samples in a clinical setting.

Electron microscopy visualization of SARS-CoV-2 RNA

Above, we demonstrated how CoronaFISH allows using different fluorescently labeled

secondary detector oligos. We reasoned that this flexibility extends beyond conventional

fluorescence microscopy, and could also allow for other imaging modalities including electron

microscopy (EM). EM is optimally suited to reveal how infection alters the cellular ultrastructure.

Indeed, conventional EM images of glutaraldehyde fixed samples showed a dramatic

reorganization of the cytoplasm of Vero cells upon SARS-CoV-2 infection, characterized by a

loss of Golgi stacks (**Fig EV4a,b**) and prominent new features, including numerous DMVs (**Fig

EV4c**), which have recently been identified as the main replication organelles of SARS-CoV-2

(Snijder *et al*, 2020; Cortese *et al*, 2020; Klein *et al*, 2020; Wolff *et al*, 2020). Budding of viral

particles appeared restricted to the lumen of the endoplasmic reticulum (ER) (data not shown)

and to electron-lucent vesicles derived from the ER-Golgi intermediate compartment (ERGIC)

that were distinct from DMVs (**Fig EV4d**), also in agreement with prior studies (Sicari *et al*,

2020).

Coupling EM with RNA *in situ* hybridization (EM-ISH) would allow for ultrastructural studies of

SARS-CoV-2 infected cells with direct visualization of the viral RNA. We previously used EM-

ISH to detect various cellular RNAs using DNA- or ribo-probes (Hubstenberger *et al*, 2017;

Yamazaki *et al*, 2018). Here, we adapted this labeling approach by using the same 96 primary

oligos against the positive strand of SARS-CoV-2 as before, hybridized to a secondary oligo

with Biotin at its 5' end. These hybrids were detected with an anti-biotin antibody conjugated to

10 nm gold particles (**Fig 4a**). EM imaging was performed on thin sections (80 nm) of Lowicryl

K4M-embedded Vero cells, either uninfected or infected (MOI 0.1, 36 h p.i.).

In uninfected samples, very few gold particles were detected (**Fig 4b**). Manual counting on

random fields yielded a mean of only 0.5 +/- 0.3 gold particles/μm^2 in nuclear and cytoplasmic

areas (n=9 regions, for a total of 94.4 μm^2), indicative of low background labeling. Although a

much larger number of small particles was visible, their size was consistent with ribosomes

rather than gold particles (**Fig 4e,f**). In infected cells, gold particles were visible in large

numbers at several locations, notably at intracytoplasmic (**Fig 4c**) and extracellular viral

particles (**Fig EV4e**), as expected. However, DMVs were the most heavily labelled structures

(**Fig 4d**), with manual counts of 180 +/- 39 gold particles/μm2 in DMV zones, a 360 fold

increase over uninfected cells (n=11 regions, for a total of 11.9 μm2). Strikingly, we observed

gold particles accumulating along DMV internal 10 nm thick fibers and at the periphery of DMVs

(**Fig 4d, Fig EV4f**). Although the nature of these fibers remains to be determined, this

accumulation of gold particles could reflect a slow export of the viral genomes through the

recently described pores spanning the DMV double membrane(Wolff *et al*, 2020). Finally, gold

particles also labeled large lysosomal organelles, shown to play a role in exocytosis of mouse β-

Coronaviruses (Ghosh *et al*, 2020) and containing densely-packed SARS-CoV-2 virions (**Fig

EV4f**).

These data demonstrate the flexibility of our probe sets, permitting their use for both

fluorescence and EM imaging, and their potential for ultrastructural studies of SARS-CoV-2

replication kinetics.

288 Discussion

Here, we presented CoronaFISH, an approach based on FISH (Tsanov *et al*, 2016) permitting

the detection of the positive and negative RNA strands of SARS-CoV-2. We validated sensitive

and specific detection of SARS-CoV-2 RNA by fluorescence microscopy in Vero cells, several

human cell lines (Caco-2, Huh7, and Calu-3), human lung tissue and nasal swabs. Lastly, we

demonstrated the flexibility of our approach, by adapting it for EM imaging of the viral RNA.

Our two probe sets each consist of 96 probes, each conjugated to two fluorescent dyes,

theoretically enabling 192 fluorescent dyes to target each RNA strand spaced along the entire

SARS-CoV-2 RNA. This results in a very bright signal, allowing viral RNA detection in

challenging samples, as demonstrated with the patient samples, where the increased signal

intensity helps to distinguish true signal despite high autofluorescence. Because our probes

span the entire length of the ~30 Kb viral RNA, they should enjoy high robustness against

mutations or partial RNA degradation. Probes are provided individually in a 96-well plate format,

therefore subsets of probes against specific regions (e.g. to target specific viral genes) can be

selected individually. Further, the secondary detector probes can easily be swapped, allowing

the use of different fluorophores and simultaneous imaging of positive and negative strands or

even change the imaging modality, as demonstrated by our EM-ISH experiments. This labeling

flexibility, together with the fact that our probes target the entire genome (as opposed to only the

Spike gene), make CoronaFISH a useful alternative to commercial FISH-based methods for

SARS-Cov-2 RNA detection such as HuluFISH and RNAscope.

Compared to immunofluorescence, our hybridization based technique offers several

advantages. First, CoronaFISH directly visualizes the viral genome (and/or its replication

intermediate) rather than viral proteins. This provides a more specific indication of viral presence

and replication, since viral proteins may be found in cells or subcellular compartments where the

viral genome is absent and/or where it does not replicate. Thus, CoronaFISH could be

instrumental in distinguishing productive from non productive (abortive) infection (Fehr &

Perlman, 2015), as has been reported for example in the context of antibody dependent

enhancement of SARS-CoV-2 infection (Lee *et al*, 2020). Thereby, CoronaFISH offers a

powerful tool to help define the molecular mechanisms of SARS-CoV-2 pathogenesis. In

addition, the ability to distinguish and simultaneously visualize positive and negative RNA

strands permits the study of replication kinetics in single cells and to uncover spatiotemporal

aspects of the infection cycle. Second, the high specificity of these probes owing to their unique

complementarity to the SARS-CoV-2 sequence allows to distinguish it from other related

viruses, which can be a common problem for antibodies against similar epitopes of different

related viral strains. Third, probes can be synthesized within a few days, permitting quick

turnover compared to antibody production. Fourth, the CoronaFISH approach is inexpensive.

Primary probes can be ordered at low cost, and the provided quantities (smallest synthesis

scale provides nanomoles for each oligo) suffice for thousands of experiments. This makes

CoronaFISH attractive for high throughput image-based screening of large libraries of antiviral

compounds, as illustrated by our Remdesivir experiment.

CoronaFISH can also be used in combination with immunofluorescence for the detection of viral

or host proteins (Rensen *et al*, 2020) and is compatible with GFP stains (Tsanov *et al*, 2016).

Furthermore, RNA-FISH and IF combined have also been shown to be suitable for flow

cytometry and fluorescence activated cell sorting (FACS) (Arrigucci *et al*, 2017). More complex

implementations enable multiplexed detection of multiple RNA species (Pichon *et al*, 2018), and

will thus permit to probe the host-pathogen interaction at the single-cell-single-virus level.

Compared to the analysis of viral RNA using (single-cell) RNA-seq, CoronaFISH provides

information on single cells in their spatial context, since experiments do not require cell

dissociation. Our approach can thus deliver insights into the viral life-cycle, including occurrence

and abundance of positive and negative-strand RNA, their subcellular localization, and their

interplay with the host, as well as with structures induced by SARS-CoV-2 infection (DMVs,

replication compartments, ERGIC). CoronaFISH also provides a unique tool to study virus-host

interactions in tissue. Furthermore, studying viral RNA presence in thousands of cells is

possible by using automated image analysis, and can hence allow the detection of rare events.

This will allow examining the effects of infection on aspects such as cell morphology, cell fusion,

cell-to-cell transmission, or tissue (re)organization.

Our data on nasal swabs suggest that CoronaFISH may be used on clinical samples and

potentially as the basis for diagnostic tests. Unlike standard RT-PCR tests, CoronaFISH does

not require RNA extraction or enzymatic reagents, which have at times been in short supply. As

mentioned above, because we chose probes targeting the whole length of the SARS-CoV-2

genomes, CoronaFISH detection is also likely to be more robust to mutations, such as the

recently identified VUI-202012/01 (aka B.1.1.7) variant, which has been reported to yield

negative results in some PCR tests based on the Spike gene (Investigation of novel SARS-

COV-2 variant Variant of Concern 202012/01, 2). The cost of CoronaFISH reagents also

compares favorably to those used in standard PCR assays. Despite these advantages, a

diagnostic test based on CoronaFISH would face two hurdles: speed and a need for a

fluorescence microscope. The duration of the FISH experiment (~2 days) is currently too long

for a rapid test. However, microfluidic devices can be used to reduce this delay to less than 15

minutes comparable to fast antigenic tests (Shaffer *et al*, 2015). The requirement for a

fluorescence microscope may also be alleviated using cheap do-it-yourself imaging systems, for

example smartphones combined with inkjet-printed lenses (Sung *et al*, 2017; Cybulski *et al*,

2014). Such portable and low-cost imaging systems could potentially facilitate point-of-care

diagnostics.

We believe that the probes and complementary labeling approaches described here expand the

toolbox for studying SARS-CoV-2 and hope that the resources provided (sequences, protocol,

and source code) will facilitate their adoption by the community to better understand, diagnose

and fight this virus.

372 Acknowledgments

We would like to thank Edouard Bertrand, who originally developed smiFISH, and Hans

Johansson for insightful discussions. We also thank Felix Rey for having helped set up the

collaboration and Guillaume Dumenil (Ultrastructural Bioimaging UTechS of Institut Pasteur) for

having suggested the application of CoronaFISH to electron microscopy and for follow-up

discussions. We thank Nathalie Jolly and Nathalie Clément (Center for Translational Science,

Institut Pasteur) for their help in obtaining authorization to use patient samples. ESL

acknowledges funding from the French Government's Investissement d'Avenir program,

'INCEPTION' (ANR-16-CONV-0005). GBS acknowledges funding by the Institut Pasteur

Coronavirus task force (don AXA COVID-19 project COVID-SPREAD). This work was funded by

Institut Pasteur.

Materials and Methods

Probe design

Entire code for probe-design is available on GitHub: https://github.com/muellerflorian/corona-fish.

The probe-design involves several steps to ensure high sensitivity for the detection of SAR-CoV-2 RNA, while minimizing false positive detection of other β-coronaviruses, other pathogens provoking similar symptoms, and transcripts of the host organism. In parenthesis, we list how many probes remain after each selection step for +strand and -strand targeting probes.

The initial list of probes for the plus and minus strand was generated with Oligostan (N=615 / 608) (Tsanov *et al*, 2016). We then selected all probes with a GC content between 40-60%, probes satisfying at least 2 out of 5 previously established criteria for efficient oligo design (Xu *et al*, 2009) (N=385 / 362).

To guarantee that the probes are insensitive towards known mutations of SARS-CoV-2, we selected only probes with not more than 2 mismatches with any of 2500 aligned SARS-CoV-2 sequences (N=333 / 311).

We then performed a local blast against other β-coronaviruses (SARS, MERS, HKU1, OC43, NL63, or 229E), other pathogens and viruses causing similar symptoms (Mycobacterium tuberculosis, Human parainfluenza virus type 1-4, Respiratory syncytial virus, Human metapneumovirus, Mycoplasma pneumoniae, Chlamydophila pneumoniae, Influenza A-D, Rhinovirus/enterovirus), as well as the transcriptome of the most common host organisms

(homo sapiens, mus musculus, African Green monkey, hamsters, ferret). We excluded all

probes with more than 22 matches in any of these blast searches (N=115 / 114).

Lastly, we selected the 96 probes with the highest NGS coverage. Probe sequences are

provided in **Supplementary Table 1**.

smiFISH

To visualize vRNA molecules, we used the smiFISH approach (Tsanov *et al*, 2016). Unlabelled

primary probes are designed to target the RNA of interest, and can be pre-hybridized with

fluorescently labeled secondary detector oligonucleotides for visualization. Probes were

designed as described above.

A detailed protocol is available in **Supplementary Note 1**. Cells were fixed in 4% PFA for 30

min, washed twice with PBS++ and stored in nuclease-free 70% ethanol at -20 °C until labeling.

On the day of the labeling, the samples were brought to room temperature, washed twice with

wash buffer A (2x SSC in nuclease-free water) for 5 min, followed by two washing steps with

washing buffer B (2X SSC and 10% formamide in nuclease-free water) for 5 min. The target-

specific primary probes were pre-hybridized with the fluorescently labeled secondary probes via

a complementary binding readout sequence. The reaction mixture contained primary probes at

a final concentration of 40 pm, and secondary probes at a final concentration of 50 pm in 1X

NEBuffer buffer (New England Biolabs). Pre-hybridization was performed in a PCR machine

with the following cycles: 85 °C for 3 min, followed by heating to 65 °C for 3 min, and a further 5

min heating at 25 °C. 2 µL of this FISH-probe stock solution was added to 100 µL of

hybridization buffer (10% (w/v) dextran, 10% formamide, 2X SSC in nuclease-free water).

Samples were placed on Parafilm in a humidified chamber on 100 µL of hybridization solution,

sealed with Parafilm, and incubated overnight at 37°C. The next day, cells were washed in the

dark at 37°C without shaking for >30min twice with pre-warmed washing buffer B. Sample were

washed once with PBS for 5 min, stained with DAPI in PBS (1:10000) for 5 min, and washed

again in PBS for 5 min. Samples were mounted in ProLong Gold antifade mounting medium.

Infection of cell lines

437   The viral stock used originates from BetaCoV/France/IDF0372/2020 and was kindly gifted by

438   the National Reference Centre for Respiratory Viruses at Institut Pasteur, Paris, and originally

439   supplied through the European Virus Archive goes Global platform.

Vero cells

The day before the infection, Vero cells were trypsinized and diluted in DMEM – Glutamax 10%

FBS. They were then seeded, 8*104 / well, in a 12-multiwell with coverslips. The day of the

infection the medium of the cells was discarded and the monolayers were infected with SARS-

CoV-2 virus in DMEM – Glutamax 2% FBS for 1h at 37°C 5% CO_2 at a multiplicity of infection

(MOI) of 0.02. The MOI used were 0.02. After the desired infection duration, the supernatant

was collected for virus titration and the cells were washed with PBS++, fixed with 4% EM-grade

PFA for 30 minutes at RT and processed for smiFISH.

Vero cells for EM-ISH

The day before the infection, Vero cells were trypsinized and diluted in DMEM – Glutamax 10%

FBS. They were then seeded, 7×10^5 cells / well, in a 6 multiwell plate. The day of the infection

the medium of the cells was discarded and cells were infected with SARS-CoV-2 virus in DMEM

– Glutamax 2% FBS at multiplicity of infection (MOI) of 0.1. After 36h, the supernatant was

collected for virus titration and the cells were washed with PBS++. The monolayers were fixed

using 4% EM-grade PFA in 0.1M Sorensen's buffer for 1 hour at 4°C.

Human cell lines

The day before the infection, Huh7, CaCo-2, Calu-3 and Vero cells were trypsinized and diluted

in DMEM – Glutamax 10% FBS (Huh7 and Vero), MEM 20% FBS + NEAA, Sodium Pyruvate

and Glutamax (CaCo-2), RPMI 20% FBS + NEAA (Calu-3). After 6 hours the medium of the

cells was discarded and the monolayers were infected in triplicate with a multiplicity of infection

(MOI) of 0.2 with SARS-CoV-2 virus in DMEM – Glutamax 2% FBS. After 36h the supernatant

was collected for virus titration and the cells were washed with PBS++. The monolayers were

fixed using 4% EM-grade PFA for 30 minutes at RT and processed for smiFISH.

Titration protocol (Focus Forming Assay)

Vero cells were seeded in 96-multi wells at 2×10^4 cells/well in DMEM – Glutamax 10% FBS.

The following day the supernatants to be titered were thawed and serially diluted in tenfold

steps in DMEM – Glutamax 1% FBS. 100 ul of the dilutions were used to infect the Vero

monolayers, and incubated for 2 h at 37˚C 5% CO2. The infection medium was then discarded

and a semi-solid media containing MEM 1X, 1.5% CMC, 10% FBS was added to the

monolayers. The cells were incubated at 37˚C 5% CO_2 for 36 hours. Cells were then fixed with

4% formaldehyde and foci were revealed using a rabbit anti-SARS-CoV N antibody and

matching secondary HRP-conjugated secondary antibodies. Foci were visualized by

diaminobenzidine (DAB) staining and counted using an Immunospot S6 Analyser (Cellular

Technology Limited CTL). Viral titers were expressed as focus forming units (FFU)/ml.

Inhibitors assay

To determine IC50 of Remdesivir in our system, Vero cells were pre-treated with serial dilution

of Remdesivir (100nM-100μM) for 1h at 37C before infection with SARS-CoV-2 at MOI 0.02.

After 2hs infection the virus inoculum was removed, cells were replenished with drug-containing

media and incubated for two days. Supernatant was then collected and titered by focus forming

assay. IC50 values were calculated by non linear regression analysis (log(inhibitor) vs response

– Variable slope (four parameters)) using Prism 6, GraphPad software. For the FISH experiment

Vero cells infected at MOI 0.1 were treated with Remdesivir 10 µM and fixed 24hs p.i.

Image-analysis of infected cell lines

Nuclei were automatically segmented in 2D images with an ImJoy (Ouyang *et al*, 2019) plugin

using the CellPose model (Stringer *et al*, 2020). Source code for segmentation is available here:

https://github.com/fish-quant/segmentation. Equidistant regions with a width of 50 pixels were

calculated around each nucleus. Overlapping regions from different nuclei were removed. 3D

FISH images were transformed into 2D images with a maximum intensity projection along z.

Signal intensity for each cell was determined as the 90% quantile of all pixels in the equidistant

region around its nucleus.

Lung tissue

Lung autopsy material from the COVID-19 patient was provided by the human biological sample

bank of the Lille COVID working group "LICORN" (Lille, France). The use of this autopsy

material for research purposes was approved by local ethics review committees at Lille Hospital

(Lille, France). Lung autopsy material from the control patient with diffuse alveolar damage prior

COVID-19 pandemic was provided by the Pathological department of Necker-Enfants maladies

Hospital (Paris, France).

Lung autopsy material was fixed in 10% neutral buffered formalin and paraffin embedded, 4 µm

sections were stained with haematoxylin and eosin staining for histological analysis using light

microscopy. SARS-CoV-2 RNA was detected as described above.

Nasal swab patient samples

Respiratory specimens (nasal swabs) have been collected from patients with respiratory

symptoms as part of routine care at the Hospital Ambroise Paré (Paris) in late March 2020. No

additional samples were collected in the course of this work. Patients were contacted, informed

about the research project, and given the possibility to oppose the use of their samples for this

project. Lack of opposition to participate in clinical research was verified in the records of all

patients whose samples were used here. This project has been recorded in the French public

register Health-data-Hub (n°F20200717122429). Processing of personal data for this study

complies with the requirements of the "reference methodology MR-004" established by the

French Data Protection Authority (CNIL) regarding data processing in health research.

Samples were screened for the presence of SARS-CoV-2 with RT-PCR as described below.

Thin-layer preparation from respiratory specimens was achieved through the cytocentrifugation

(800rpm, 10min) of 150µL of the remaining respiratory sample with a Thermo Scientific Cytospin

4 cytocentrifuge. Cells were fixed in PBS-PFA 4% for 30 min and then conserved frozen in

100% ethanol until FISH was performed.

RT-PCR

RNA extraction: 400 µL of clinical samples were extracted in 300 µL of elution buffer (Total NA

Lysis/Binding buffer) for 20 min at room temperature with gentle agitation. RNA was extracted

with the MagnaPURE compact (Roche) and the MagNA Pure Compact DNA Isolation Kit I

(Roche) following the protocol "Total_NA_Plasma_external_lysis purification protocol". Final

dilution of RNA was in 50 µL elution buffer.

RT-PCR: Screening for SARS-CoV-2 was performed by RT-PCR following a modified protocol

recommended by the French National Reference Center for Respiratory Viruses, Institut

Pasteur, Paris using Ag Path-ID One-Step RT-PCR kit® (Thermofisher). PCR reaction was run

on the ABI PRISM® 7900 system (Applied Biosystems) with the following cycle settings: 45°C

10'; 95°C 5'; 45 X (95°C 15''; 58°C 45''). Primer sequences and concentration are provided in

Supplementary Table 2.

FISH

Thin-layered samples on a cover-slide suitable for FISH were obtained with a Cytospin protocol.

150 µL of the sample were deposited in a Cytofunnel (Thermo Scientific 1102548). Samples

were then centrifuged (800 rpm, 10 min, room temperature) with a Cytospin 4 cytocentrifuge

(Thermo Scientific) on Cytoslides (Thermo Scientific 5991059). Cells were fixed in PBS-PFA 4%

for 30 min and then conserved frozen in 70% ethanol at -20°C. FISH protocol was performed

with Cy3-labeled plus-strand probes labeled with as described above, with one exception: for

hybridization 400 µL hybridization buffer was used per sample instead of 100 µL (with the same

final probe concentration).

EM-ISH

Fixation and embedding for electron microscopy

For Epon embedding, cell cultures were fixed for 1 hour at 4°C in 2% glutaraldehyde (Taab

Laboratory Equipment, Reading, UK) in 0.1 M phosphate buffer, pH 7.3. During fixation, cells

were scraped off from the plastic substratum and centrifuged at 5000g for 15 min. Cell pellets

were dehydrated in increasing concentrations of ethanol and embedded in Epon.

Polymerization was carried out for 48 hours at 64°C. Ultrathin sections were collected on

Formvar-carbon-coated copper grids (200 mesh) and stained briefly with standard uranyl-

acetate and lead-citrate solutions.

Embedding in Lowicryl K4M (Chemische Werke Lowi, Waldkraiburg, Germany) was carried

out on Vero cells fixed either in 4% formaldehyde (Merck, Darmstadt, Germany) or in 2%

glutaraldehyde at 4°C. Cell pellets were equilibrated in 30% methanol and deposited in a Leica

EM AFS2/FSP automatic reagent handling apparatus (Leica Microsystems). Lowicryl

polymerization under UV was for 40 h at – 20°C followed by 40 h at +20°C Ultrathin-sections of

Lowicryl-embedded material were collected on Formvar-carbon-coated gold grids (200 mesh)

and stored until use.

Electron microscopic *in situ* hybridization (EM-ISH)

At the EM level, the SARS-CoV-2 RNA (+) strand probe was composed of the same set of 96

oligodeoxynucleotides that was used for RNA-FISH. The secondary oligonucleotide, however,

was modified by a custom-added biotin residue at its 5'-end (Qiagen). Hybrids of the CoV-2

RNAs with the probe were detected with a goat anti-biotin antibody conjugated to 10 nm gold

particles (BBI international).

High resolution in situ hybridization was carried out essentially as described previously

(Hubstenberger *et al*, 2017; Yamazaki *et al*, 2018). The hybridization solution contained 50%

deionized formamide, 10% dextran sulfate, 2 x SSC, and a final concentration of 80 ng/ml of a

mix of 1 µg/µl primary oligonucleotides and 1.2 µg/µl of biotinylated secondary oligonucleotide

stored at -20°C. EM-grids, with ultra-thin sections of either formaldehyde- or glutaraldehyde-

fixed cells side down, were floated for 3 h at 37-42°C on a 1.5 µl drop of hybridization solution

deposited on a parafilm in a moist glass chamber. EM-grids were then briefly rinsed over three

drops of PBS and incubated 30 min at RT on a drop of goat anti-biotin antibody (BBI

International) conjugated to 10 nm gold particles diluted 1/25 in PBS. EM-grids were further

rinsed on 2 drops of PBS and finally washed with a brief jet of deionized water at high intensity.

Following a 10 min drying on filter paper with thin-sections on top, the EM grids were stained 1

min on a drop of 4% uranyl acetate in water and dried on filter paper 30 min before observation

under the EM. Standard lead citrate staining was omitted to favor higher contrast of gold

particles over the moderately-stained cellular structures.

Sections were analyzed with a Tecnai Spirit transmission electron microscope (FEI, Hillsboro,

OR), and digital images were taken with an SIS MegaviewIII charge-coupled device camera

(Olympus, Tokyo, Japan). Quantitation was performed by manually counting gold particles on

surfaces that were measured using analySIS software (Olympus Soft Imaging Solutions,

Munster, Germany). Statistics were calculated using Excel (Microsoft, Redmond, WA).

References

Arrigucci R, Bushkin Y, Radford F, Lakehal K, Vir P, Pine R, Martin D, Sugarman J, Zhao Y,
Yap GS, *et al* (2017) FISH-Flow, a protocol for the concurrent detection of mRNA and
protein in single cells using fluorescence in situ hybridization and flow cytometry. *Nat*
Protoc 12: 1245–1260
Bost P, Giladi A, Liu Y, Bendjelal Y, Xu G, David E, Blecher-Gonen R, Cohen M, Medaglia C, Li
H, *et al* (2020) Host-Viral Infection Maps Reveal Signatures of Severe COVID-19
Patients. *Cell* 181: 1475-1488.e12
Bradley BT, Maioli H, Johnston R, Chaudhry I, Fink SL, Xu H, Najafian B, Deutsch G, Lacy JM,
Williams T, *et al* (2020) Histopathology and ultrastructural findings of fatal COVID-19
infections in Washington State: a case series. *The Lancet* 396: 320–332
Chen F, Wassie AT, Cote AJ, Sinha A, Alon S, Asano S, Daugharthy ER, Chang J-B,
Marblestone A, Church GM, *et al* (2016) Nanoscale imaging of RNA with expansion
microscopy. *Nat Methods* 13: 679–684
Cheung CY, Poon LLM, Ng IHY, Luk W, Sia S-F, Wu MHS, Chan K-H, Yuen K-Y, Gordon S,
Guan Y, *et al* (2005) Cytokine Responses in Severe Acute Respiratory Syndrome
Coronavirus-Infected Macrophages In Vitro: Possible Relevance to Pathogenesis. *J Virol*
79: 7819–7826
Cortese M, Lee J-Y, Cerikan B, Neufeldt CJ, Oorschot VMJ, Köhrer S, Hennies J, Schieber NL,
Ronchi P, Mizzon G, *et al* (2020) Integrative Imaging Reveals SARS-CoV-2-Induced
Reshaping of Subcellular Morphologies. *Cell Host Microbe* 28: 853-866.e5
Cybulski JS, Clements J & Prakash M (2014) Foldscope: Origami-Based Paper Microscope.
PLoS ONE 9
Fehr AR & Perlman S (2015) Coronaviruses: an overview of their replication and pathogenesis.
Methods Mol Biol Clifton NJ 1282: 1–23
Fung TS & Liu DX (2019) Human Coronavirus: Host-Pathogen Interaction. *Annu Rev Microbiol*
73: 529–557
Ghosh S, Dellibovi-Ragheb TA, Kerviel A, Pak E, Qiu Q, Fisher M, Takvorian PM, Bleck C, Hsu
VW, Fehr AR, *et al* (2020) β-Coronaviruses Use Lysosomes for Egress Instead of the
Biosynthetic Secretory Pathway. *Cell* 183: 1520-1535.e14
Gordon CJ, Tchesnokov EP, Woolner E, Perry JK, Feng JY, Porter DP & Götte M (2020)
Remdesivir is a direct-acting antiviral that inhibits RNA-dependent RNA polymerase from
severe acute respiratory syndrome coronavirus 2 with high potency. *J Biol Chem* 295:
6785–6797
Hanley B, Naresh KN, Roufosse C, Nicholson AG, Weir J, Cooke GS, Thursz M, Manousou P,
Corbett R, Goldin R, *et al* (2020) Histopathological findings and viral tropism in UK
patients with severe fatal COVID-19: a post-mortem study. *Lancet Microbe* 1: e245–
e253
Hubstenberger A, Courel M, Bénard M, Souquere S, Ernoult-Lange M, Chouaib R, Yi Z, Morlot
J-B, Munier A, Fradet M, *et al* (2017) P-Body Purification Reveals the Condensation of
Repressed mRNA Regulons. *Mol Cell* 68: 144-157.e5
Investigation of novel SARS-COV-2 variant Variant of Concern 202012/01
Itzkovitz S & van Oudenaarden A (2011) Validating transcripts with probes and imaging
technology. *Nat Methods* 8: S12–S19
Johns Hopkins University Dashboard
Kim D, Lee J-Y, Yang J-S, Kim JW, Kim VN & Chang H (2020) The Architecture of SARS-CoV-
2 Transcriptome. *Cell* 181: 914-921.e10
King BR, Samacoits A, Eisenhauer PL, Ziegler CM, Bruce EA, Zenklusen D, Zimmer C, Mueller

F & Botten J (2018) Visualization of Arenavirus RNA Species in Individual Cells by
636   Single-Molecule Fluorescence In Situ Hybridization Suggests a Model of Cyclical
637   Infection and Clearance during Persistence. *J Virol* 92
638   Klein S, Cortese M, Winter SL, Wachsmuth-Melm M, Neufeldt CJ, Cerikan B, Stanifer ML,
639   Boulant S, Bartenschlager R & Chlanda P (2020) SARS-CoV-2 structure and replication
640   characterized by in situ cryo-electron tomography. *Nat Commun* 11: 5885
641   Lee WS, Wheatley AK, Kent SJ & DeKosky BJ (2020) Antibody-dependent enhancement and
642   SARS-CoV-2 vaccines and therapies. *Nat Microbiol* 5: 1185–1191
643   Mueller F, Senecal A, Tantale K, Marie-Nelly H, Ly N, Collin O, Basyuk E, Bertrand E, Darzacq
644   X & Zimmer C (2013) FISH-quant: automatic counting of transcripts in 3D FISH images.
645   *Nat Methods* 10: 277–278
646   Ogando NS, Dalebout TJ, Zevenhoven-Dobbe JC, Limpens RWAL, van der Meer Y, Caly L,
647   Druce J, de Vries JJC, Kikkert M, Bárcena M, *et al* (2020) SARS-coronavirus-2
648   replication in Vero E6 cells: replication kinetics, rapid adaptation and cytopathology. *J
649   Gen Virol*
650   Ouyang W, Mueller F, Hjelmare M, Lundberg E & Zimmer C (2019) ImJoy: an open-source
651   computational platform for the deep learning era. *Nat Methods* 16: 1199–1200
652   Pichon X, Lagha M, Mueller F & Bertrand E (2018) A Growing Toolbox to Image Gene
653   Expression in Single Cells: Sensitive Approaches for Demanding Challenges. *Mol Cell*
654   71: 468–480
655   Raj A, van den Bogaard P, Rifkin SA, van Oudenaarden A & Tyagi S (2008) Imaging individual
656   mRNA molecules using multiple singly labeled probes. *Nat Meth* 5: 877–879
657   Rensen E, Mueller F, Scoca V, Parmar JJ, Souque P, Zimmer C & Di Nunzio F (2020)
658   Clustering and reverse transcription of HIV-1 genomes in nuclear niches of
659   macrophages. *EMBO J*: e105247
660   Shaffer SM, Joshi RP, Chambers BS, Sterken D, Biaesch AG, Gabrieli DJ, Li Y, Feemster KA,
661   Hensley SE, Issadore D, *et al* (2015) Multiplexed detection of viral infections using rapid
662   in situ RNA analysis on a chip. *Lab Chip* 15: 3170–3182
663   Sicari D, Chatziioannou A, Koutsandreas T, Sitia R & Chevet E (2020) Role of the early
664   secretory pathway in SARS-CoV-2 infection. *J Cell Biol* 219
665   Skinner SO, Sepúlveda LA, Xu H & Golding I (2013) Measuring mRNA copy number in
666   individual Escherichia coli cells using single-molecule fluorescent in situ hybridization.
667   *Nat Protoc* 8: 1100–1113
668   Snijder EJ, Limpens RWAL, Wilde AH de, Jong AWM de, Zevenhoven-Dobbe JC, Maier HJ,
669   Faas FFGA, Koster AJ & Bárcena M (2020) A unifying structural and functional model of
670   the coronavirus replication organelle: Tracking down RNA synthesis. *PLOS Biol* 18:
671   e3000715
672   Stertz S, Reichelt M, Spiegel M, Kuri T, Martínez-Sobrido L, García-Sastre A, Weber F & Kochs
673   G (2007) The intracellular sites of early replication and budding of SARS-coronavirus.
674   *Virology* 361: 304–315
675   Stringer C, Wang T, Michaelos M & Pachitariu M (2020) Cellpose: a generalist algorithm for
676   cellular segmentation. *Nat Methods*
677   Sung Y, Campa F & Shih W-C (2017) Open-source do-it-yourself multi-color fluorescence
678   smartphone microscopy. *Biomed Opt Express* 8: 5075–5086
679   Takayama K (2020) In Vitro and Animal Models for SARS-CoV-2 research. *Trends Pharmacol
680   Sci*
681   Tang T, Bidon M, Jaimes JA, Whittaker GR & Daniel S (2020) Coronavirus membrane fusion
682   mechanism offers a potential target for antiviral development. *Antiviral Res* 178: 104792
683   Trcek T, Chao JA, Larson DR, Park HY, Zenklusen D, Shenoy SM & Singer RH (2012) Single-
684   mRNA counting using fluorescent in situ hybridization in budding yeast. *Nat Protoc* 7:
685   408–419

Tsanov N, Samacoits A, Chouaib R, Traboulsi A-M, Gostan T, Weber C, Zimmer C, Zibara K,
Walter T, Peter M, *et al* (2016) smiFISH and FISH-quant - a flexible single RNA
detection approach with super-resolution capability. *Nucleic Acids Res* 44: e165
V'kovski P, Kratzel A, Steiner S, Stalder H & Thiel V (2020) Coronavirus biology and replication:
implications for SARS-CoV-2. *Nat Rev Microbiol*: 1–16
Wang S (2019) Single Molecule RNA FISH (smFISH) in Whole-Mount Mouse Embryonic
Organs. *Curr Protoc Cell Biol* 83: e79
Wolff G, Limpens RWAL, Zevenhoven-Dobbe JC, Laugks U, Zheng S, Jong AWM de, Koning
RI, Agard DA, Grünewald K, Koster AJ, *et al* (2020) A molecular pore spans the double
membrane of the coronavirus replication organelle. *Science* 369: 1395–1398
Wu C, Simonetti M, Rossell C, Mignardi M, Mirzazadeh R, Annaratone L, Marchiò C, Sapino A,
Bienko M, Crosetto N, *et al* (2018) RollFISH achieves robust quantification of single-
molecule RNA biomarkers in paraffin-embedded tumor tissue samples. *Commun Biol* 1:
1–8
Xu Q, Schlabach MR, Hannon GJ & Elledge SJ (2009) Design of 240,000 orthogonal 25mer
DNA barcode probes. *Proc Natl Acad Sci U S A* 106: 2289–2294
Yamazaki T, Souquere S, Chujo T, Kobelke S, Chong YS, Fox AH, Bond CS, Nakagawa S,
Pierron G & Hirose T (2018) Functional Domains of NEAT1 Architectural lncRNA Induce
Paraspeckle Assembly through Phase Separation. *Mol Cell* 70: 1038-1053.e7

Tables

Cell line	Viral titer [FFU/ml]
Caco2	2×10^3
Huh7	4×10^3
Calu3	2×10^5
Vero	5×10^3

709 **Table 1**. Titration on the supernatant of infected mammalian cells.

 Figure Legends

 Figure 1. Visualizing SARS-CoV-2 with CoronaFISH

(a) Principle of CoronaFISH. 96 primary probes are pre-hybridized *in vitro* with dye-carrying secondary probes via the FLAP sequence. Resulting duplexes are subsequently hybridized in cells to target the SARS-CoV-2 positive or negative RNA. **(b)** Replication cycle of SARS-CoV-2. Incoming, genomic positive-strand RNA is used to produce viral polymerase. Polymerase produces a negative-strand replication intermediate, which serves as a template for synthesis of full-length positive-strand and shorter sub-genomic RNAs. The latter are used to produce other viral proteins. **(c)** Genome of SARS-CoV-2 with indicated probe positions targeting the positive and negative strand. **(d)** Images of uninfected and infected Vero cells with either the positive or negative strands detected with Cy3-labeled probes. Shown are zoom-ins on individual cells. Full-size images in **Fig EV1a**. First column shows uninfected control, second and third column infected cells with different intensity scalings as indicated in brackets (the first and second values in brackets indicate the pixel values corresponding to the lowest and brightest intensities in the displayed image, respectively). Scale bars 5 µm. Scale bar in red inset 1 µm. **(e)** Quantification of signal intensities in individual cells. Dashed line is the 99% quantile estimated from uninfected samples. **(f)** Simultaneous imaging of positive and negative strands with dual-color CoronaFISH. Scale bar in full image 10 µm, in inset 2 µm. **(g)** Images of Remdesivir treated (right) or untreated cells (left). Scale bars 30 µm. (**h**) Quantification of Remdesivir treated cells performed as in e).

Figure 2. CoronaFISH in human cell lines

(a-b) FISH against the positive strand of SARS-CoV-2 in three different cell lines. (**a**) Full field of

views, scale bars 30 µm , (b) zoom-ins on indicated green rectangles, scale bars 10 µm. **(c)**

FISH signal intensities of cells in a small region around the nucleus of each cell. Box plots as in

Fig 1e.

Figure 3. CoronaFISH in human tissue and nasal swabs

(a) Detection of positive-strand SARS-CoV-1 in human lung tissue. Scale bar 20 µm. Right

image is a magnified view of the green boxed region of interest. Negative control and

histopathology images in **Fig EV3. (b)** Nasal swabs were used to perform RT-PCR and the

surplus was used for imaging experiments. **(c)** FISH against SARS-CoV-2 RNA in a patient

sample tested positive for SARS-CoV-2. Positive-strand RNA was labeled with Cy3 (white),

nucleus in blue (DAPI). Scale bar 20 µm. Large field of view and negative controls in

Fig EV2b,c

Figure 4. CoronaFISH for electron microscopy

(a) Principle of EM-ISH performed on thin sections of Lowicryl K4M-embedded infected and

uninfected Vero cells. The asterisk indicates biotin at the 5'-end of the secondary oligo that is

recognized by the anti-biotin antibody conjugated to 10 nm gold particles. As sketched, only

virions with a section on the upper face of the ~80 nm ultrathin section will be labelled. **(b)**

Uninfected control samples. Blue oval surrounds an example of sparse background staining by

electron-dense gold particles. The less defined punctate structures such as those lining up the

ER (middle panel) are ribosomes (see panels **e,f**). Nu: nucleus; M: mitochondria; ER:

endoplasmic reticulum. **(c, d)** Overviews of SARS-CoV-2 infected Vero cells showing major

cytoplasmic vacuolization by virus-induced DMVs. Positive strand of SARS-CoV-2 can be

detected over intracytoplasmic aggregates of viral particles (**c**). See **Fig EV4e** for extracellular

aggregates. DMVs were the most heavily labelled structures, and displayed viral RNAs

accumulating especially on peripheral 10 nm fibers (**d,** rightmost frame and **Fig EV4f**). By

contrast, mitochondria or nuclei were not significantly labelled. DMV: double membrane

vesicles. Nu: nucleus; M: mitochondria. (**e**) Averaged image from punctate structures in panels

b-d detected with FISH-quant and aligned to the same center (Mueller *et al*, 2013). **(f)** Line

profiles through the averaged spots in **e**. The punctate structures visible in infected cells have a

size in agreement with the 10 nm nano-gold particles, whereas the punctate structures in

uninfected cells are substantially larger and consistent with 30 nm ribosomes.

Figure 1

Figure 2

a

Histopathology

Lung biopsy

CoronaFISH

Lung tissue

+ strand (Cy3)
DAPI

b

RT-PCR

Nasal swab

CoronaFISH

c Nasal swab (Ct=21)

[300, 3000]

+ strand (Cy3)
DAPI

Figure 3

Figure 4