The epoxyeicosatrienoic pathway is intact in endothelial and smooth muscle cells exposed to aldosterone excess

Short title: Intact EET pathway in hyperaldosteronism

Laura BRUNNENKANT[a*], Yao MENG[a*], Jing SUN[a], Jair GONZALEZ MARQUES[b], Berthold KOLETZKO[b], Michael MEDEROS Y SCHNITZLER[c,d], Thomas GUDERMANN[c,d], Tracy Ann WILLIAMS[a,e], Felix BEUSCHLEIN[a,f], Daniel A HEINRICH[a], Christian ADOLF[a], Martin REINCKE[a,d], and Holger SCHNEIDER[a‡]

[a]Department of Medicine IV, University Hospital, LMU Munich, Munich, Germany

[b]Department of Pediatrics, Dr. von Hauner Children's Hospital, University Hospital, LMU Munich, Munich, Germany

[c]Walther Straub Institute of Pharmacology and Toxicology, Faculty of Medicine, LMU Munich, Munich, Germany

[d]DZHK (German Centre for Cardiovascular Research), Munich Heart Alliance, Munich, Germany

[e]Division of Internal Medicine and Hypertension, Department of Medical Sciences, University of Turin, Turin, Italy

[f]Klinik für Endokrinologie, Diabetologie und Klinische Ernährung, Universitätsspital Zürich, Zurich, Switzerland

*shared first authorship

‡corresponding author

Correspondence should be addressed to:

Dr Holger Schneider

Endocrine Research Unit

Medizinische Klinik und Poliklinik IV

Klinikum der Universität München

Ziemssenstr. 1

80336 München

Germany

Telephone: +49-89-4400-52943

Fax: +49-89-4400-57288

Email: holger.schneider@med.uni-muenchen.de

Word count: 5091

Number of tables: 0

Number of figures: 6

Number of supplementary digital content files: 1

Abstract

Objectives:

Endothelial dysfunction (ED) is considered to be a major driver of the increased incidence of cardiovascular disease in primary aldosteronism (PA). Whether the epoxyeicosatrienoic acid (EET) pathway, involving the release of beneficial endothelium-derived lipid mediators, contributes to ED in PA is unknown. Preclinical evidence suggests this pathway to be relevant in the pathogenesis in various models of experimental hypertension. In addition, an orally available soluble epoxide hydrolase inhibitor, which halts the breakdown of EETs, has already passed a phase 1 clinical trial.

We, therefore, exposed primary human coronary artery endothelial cells to 1 nM aldosterone.

Methods:

We used qPCR to investigate changes in the expression levels of essential genes for the synthesis and degradation of EETs as well as mass spectrometry to determine endothelial synthetic capacity to release EETs upon stimulation. We also assessed primary human coronary artery smooth muscle cells for expression of putative EET receptor ion channels or downstream mediators as well as for the calcium response to EETs using calcium imaging.

Results:

No major aldosterone-related expression changes were detected on the endothelial as well as the smooth muscle side. Stimulated release of endothelial EETs was unaffected. Likewise, the smooth muscle calcium response was unchanged after aldosterone excess.

Conclusions:

61 The EET pathway is not negatively affected by increased aldosterone concentrations

62 as seen in PA. Modulating the EET pathway with therapeutic intent in patients with PA

63 might therefore be assessed in future preclinical and clinical trials to address ED.

64 Keywords: aldosterone, smooth muscle, endothelium, vascular remodeling,

65 hyperaldosteronism, eicosanoids

Introduction

Primary aldosteronism (PA) is the most common cause of endocrine hypertension. In

recent years, it has been appreciated that in up to 12 % of patients arterial hypertension

can be attributed to aldosterone excess[1]. It is of particular importance to address this

condition adequately as high levels of aldosterone cause a significantly greater

prevalence of atherosclerotic sequelae as reflected by blood-pressure-independent

detrimental effects on the cardiovascular system[2]. In particular, renal injury, atrial

fibrillation, coronary artery disease and stroke even when compared to essentially

hypertensive control patients who had been matched for age, sex and blood pressure

are more abundant in patients with PA. One of the earliest steps in the pathogenesis

of atherosclerosis is endothelial dysfunction[3]. In line with previous observations

indicating a high prevalence of atherosclerotic disease, endothelial dysfunction has

been demonstrated in patients with PA[4–6]. Endothelial dysfunction is a descriptive

term which does not allow for a mechanistic attribution. While experimental studies

have linked aldosterone-mediated endothelial dysfunction to a release of constrictive

prostanoids[7] or endothelin-1[8], most lines of explanation focus on impaired

endothelial nitric oxide (NO) generation, decreased NO bioavailability or decreased

smooth muscle response to NO[9–11].

However, the relative importance of NO as vasodilator decreases with ageing and

sedentary life style[12] and, consequently, a shift to other endothelial mediators seems

to occur. Other non-nitric oxide endothelial factors encompass substances like

prostacyclin (PGI$_2$) and a heterogeneous group of mediators summarized as EDHF

(endothelium-derived hyperpolarizing factor). Diverse mechanisms such as localized

increases in extracellular potassium via release from the endothelial cytosol,

hyperpolarizing currents through myoendothelial gap junctions, release of hydrogen

peroxide and arachidonic-acid-derived epoxyeicosatrienoic acids (EETs) have been

proposed to constitute molecular correlates of EDHF[13].

Specifically, the role of EETs in aldosterone-mediated endothelial dysfunction has

barely been investigated so far. EETs comprise 4 bioactive regioisomers which are

named according to the position of the epoxide group (5,6, 8,9, 11,12 and 14,15 EET).

They are degraded into inactive DHETs (dihydroxyeicosatrienoic acids) by epoxide

hydrolases. Clinical data indicate that levels of circulating 14,15 DHET, the breakdown

product of 14,15 EET, are increased in patients with PA, which might point to increased

breakdown of active EETs[14]. Further, experimental hypertension induced by

administration of the mineralocorticoid precursor DOCA (deoxycorticosterone acetate)

was reversed after administration of an inhibitor of soluble epoxide hydrolase[15],

which in turn would increase the bioavailability of endothelium-derived EETs. These

findings suggest a particular role of EETs in the pathogenesis of mineralocorticoid-

induced endothelial dysfunction and warrant further investigation.

Since the impact of aldosterone excess on the EET pathway in both major vascular

cell types is unknown, we set out to explore for putative pathologic effects using

primary human coronary artery cell lines. We addressed both the capacity of

endothelial cells to synthesize and release EETs as well as the calcium response of

smooth muscle cells to investigate which cell type might be specifically perturbed. We

deliberately focussed on pharmacologic strategies to allow for extrapolation of our

results to clinical cohorts.

Materials and Methods

The authors refer to the online-available supplement with a detailed list of materials

and methods used in the experiments.

Statistics: GraphPad Prism 8 (GraphPad Software, San Diego, CA, USA) was used

for construction of graphs and computation of statistical significances. Bar graphs show

mean ± SEM. Statistical tests comprised t-tests or 1-Way ANOVA followed by indicated

post hoc tests, wherever applicable. Statistical significance levels were set to a two-

sided p-value of <0.05.

Results

In endothelial cells (EC), EETS are synthesized by CYP2C isoenzymes[16,17] as well

as CYP2J2[18]. Assessment of the expression status of CYP-epoxygenases revealed

that human coronary artery EC express CYP2C8 and CYP2J2. CYP2C9 mRNA could

not be detected.   Neither CYP2J2 nor CYP2C8 showed significant changes in

transcript levels upon challenge with aldosterone, with or without co-stimulation by

cortisol (Figure 1A, 1B).

There are at least two epoxide hydrolases which have been reported to be capable of

converting EETs to inactive DHETs, microsomal epoxide hydrolase (EPHX1) and

soluble epoxide hydrolase (EPHX2), with the latter reportedly being the most important

one[19].

Microsomal epoxide hydrolase (EPHX1) and soluble epoxide hydrolase (sEH, EPHX2)

were not dysregulated in aldosterone-treated smooth muscle cells (SMC) (Figure 2C,

2E) and EC (Figure 2D, 2F) when compared to their appropriate time-controls. A

striking observation, however, was the drastically increased expression of EPHX1

over EPHX2 within both cell types: In 48 h cortisol-treated SMC normalized expression

levels ($2^{-\Delta CT}$) of EPHX1 versus EPHX2 were 0.24 ± 0.02 versus 0.002 ± 0.0001; 48 h

F-treated EC: 0.83 ± 0.13 versus 0.003 ± 0.001.

Further results showing the treatment response on MR and GR target genes in EC and

SMC as well as expression of HSD11B1 and HSD11B2 are provided in the

supplement.

We next directly assessed the concentrations of endothelial EETs after exposure to 48

h of aldosterone and cortisol by high pressure liquid chromatography tandem mass

spectrometry (HPLC MS/MS). Attention was paid to establishment of the method

according to the criteria of the European Medicines Agency[20] (see Supplemental

Tables 2 and 3). We used two concentrations of acetylcholine (ACh), 1 and 100 µM.

The method was able to quantify all 4 EET regioisomers and their respective DHET

epoxide hydrolase products. 14,15 EET and its respective diol were the most abundant

regioisomers. ACh-stimulated secretion of EETs was unaltered when compared to

cells only treated with cortisol. Likewise, MR antagonism or inhibition of sEH with GSK

2256294 (GSK) parallel to application of aldosterone and cortisol did not resulted in

major changes despite that GSK increased 11,12 EET concentrations in ACh 1 µM-

stimulated supernatants compared to controls.

Higher concentrations of EETs and lower concentrations of DHETs were measured in

the supernatants of cells stimulated with 100 µM *vs.* 1 µM ACh. The sum of EETs and

DHETs as a measure of CYP-epoxygenase function did not differ across the two ACh

concentrations. Epoxide hydrolase activity was inhibited as a function of ACh

concentration, as evidenced by the DHET/EET ratio. In line with this interpretation,

GSK precluded the ACh-mediated decrease in 14,15 DHET/EET. Individual

concentrations of eicosanoids and derived measures are provided in Supplemental

Tables S4 and S5.

EETs reportedly operate indirectly via BK$_{Ca}$-mediated hyperpolarization which leads to

a decrease in smooth muscle calcium, relaxation and – eventually – vasodilation[21].

Activation of BK$_{Ca}$ channels in turn is mediated by calcium transients through TRPV4,

which are direct targets of EETs[22].

In SMC, TRPV4 expression was unrelated to the presence of aldosterone (Figure 3A).

Likewise, no difference in L-type calcium channel expression (CACNA1C coding for

Cav1.2) was detected related to the presence or absence of aldosterone with or without

cortisol (Figure 3B). Aldosterone alone did not increase CACNA1C transcript levels

while cortisol alone reduced Cav1.2 mRNA. Of note, the glucocorticoid receptor

antagonist mifepristone (RU486) did not antagonize the cortisol-mediated reduction in

Cav1.2 mRNA.

The expression of BK$_{Ca}$ alpha subunit (KCNMA1) was reduced in SMC treated with

aldosterone and cortisol as compared to cortisol alone (Figure 4A). Further

examination revealed that the BK$_{Ca}$ message reduction was most pronounced after

addition of cortisol. Addition of RU486 to either cortisol or the combined treatment with

aldosterone and cortisol substantially increased BK$_{Ca}$ message levels without apparent

synergism with eplerenone (Figure 4B). However, despite the reproducible reduction

in KCNMA1 mRNA, no effect on protein level could be detected in Western blot (Figure

4C).

14,15 EET-elicited calcium transients in Fura2-loaded SMC were unchanged in cells

pre-treated with aldosterone and cortisol for 48 hours when compared to cortisol-

treated cells (Figure 5). Application of MRA reduced the calcium transient significantly

when compared to the control group.

SMC were submitted to a concentration response curve for 14,15 EET in a plate reader

format. A fourth group was included which was treated for 48 hours with 1 µM EET +

3 µM GSK in addition to aldosterone and cortisol. EET and GSK were applied together

to correct for SMCs not being able to synthesize EETs themselves. Relative increases

in fluorescence were comparable across all three treatment groups until 1 µM EET

concentration (Figure 6). At 1 µM, Ep- and EET+GSK-treated cells showed a decline

in fluorescence response to EET application. The fluorescence in Aldo-treated cells,

however, did not differ to the cortisol-treated control group at any point of the EET

concentration-response curve.

Discussion

In the present study we investigated whether concentrations of aldosterone which are

typically found in patients with PA impair human coronary artery EC synthetic capacity

to release epoxyeicosatrienoic acids (EET). We also addressed whether human

coronary artery SMC reactivity to EETs would be hampered by excess of aldosterone.

The two cell types were chosen deliberately, since the effects of EETs have been

extensively validated in coronary arteries from a variety of species, including

bovine[23] and human arteries[24].

Effects of excess of aldosterone on endothelial CYP-epoxygenase expression levels

have not been investigated so far. We report that aldosterone essentially does neither

alter the expression of CYP2C8 and CYP2J2, nor of epoxide hydrolases which might

degrade EETs. Importantly, we were able to corroborate these findings in functional

studies: EET release from endothelial cells was not affected by excessive

concentrations of aldosterone. In line with our findings, no change in renal expression

levels of soluble epoxide hydrolase was detected in a mouse model of DOCA-induced

hypertension[25].

We used ACh to trigger endothelial EET release. Two concentrations of ACh yielded

a graded endothelial response which was most obvious in the DHET/EET ratio. We

therefore concluded that the lack of a difference in stimulated EETs between the

treatment groups was not masked by exhaustion of endothelial secretory capacity with

maximum ACh concentrations. Consistent with earlier reports, endothelial cells

released mostly 14,15 EET and 11,12 EET[26,27], a finding compatible with the

preferential synthesis patterns of CYP2C8 and CYP2J2[28,29].

In patients with PA, higher serum levels of 14,15 DHET were detected which was

interpreted as increased breakdown of 14,15 EET; however, no ratio of DHET/EET

was measured. DHET levels positively correlated with the extent of abdominal aortic

calcification[14]. While we also observed numerically higher values of 14,15 DHET in

our aldosterone- and cortisol-treated ECs, the differences were not significant. As a

caveat, circulating 14,15 DHET and its precursor substance are not necessarily

endothelium-derived since other cells such as erythrocytes have been shown to

release EETs[30].

The DHET/EET ratio decreased with the higher concentration of ACh. At the same

time, the sum of all EETs and DHETs as time integral of EET production remained

constant. We therefore conclude that higher concentrations of ACh lead to an inhibition

of epoxide hydrolase. It remains to be determined whether in this context the

predominant epoxide hydrolase is sEH or mEH. Also, it appears that this study is the

first to suggest an ACh concentration-dependent negative regulation of epoxide

hydrolase activity.

We detected a minimum 120-fold higher relative expression level of microsomal over

soluble epoxide hydrolase in both EC as well as SMC. The negligible effect of GSK in

increasing the levels of EETs and decreasing DHET concentrations after stimulation

with acetylcholine might be a direct consequence of the low expression of EPHX2.

In SMC, we could confirm intact expression levels of both TRPV4 and BK_{Ca} channels

as well as Cav1.2 channels, indicating an intact molecular machinery to transduce

endothelium-derived EET signals. In line with these findings, calcium responses to

EETs were comparable between control cells and aldosterone-treated SMC.

We could show synergistic glucocorticoid and mineralocorticoid small repressor effects

on BK_{Ca} transcription. The isolated effect of aldosterone, however, did not translate

into diminished protein levels as also reflected in unchanged calcium responses to

14,15 EET. We concluded that BK_{Ca} channels are mostly regulated by glucocorticoids

and that their transcriptional control is most likely not a primary mechanism of

aldosterone-mediated alterations in vascular smooth muscle function.

Conflicting data from mouse models on the role of A on BK_{Ca} expression do not allow

to integrate our data into a clear picture[31–33].

To the best of our knowledge, this study is the first to suggest glucocorticoid receptor

and MR synergism in transcriptional repression of BK_{Ca} channels. It will be intriguing

to study these effects further in populations with excessive glucocorticoid secretion (i.e.

Cushing disease), co-secretion of gluco- and mineralocorticoids (as often found in

primary aldosteronism[34]) or chronic exogenous glucocorticoid administration.

The smooth muscle calcium transient to 14,15 EET was unaffected by exposure to

aldosterone. MRA antagonism, however, severely blunted the calcium transient to

EET. Conceivably, calcium imaging unmasked an inhibitory effect of eplerenone on

pannexin 1 hemichannels[35,36] which became obvious at 1 μM EET in both the single

cell imaging as well as the plate reader format.

Stimulation with single EET concentrations has previously been shown to increase

global smooth muscle intracellular calcium[37]. True concentration-response curves in

SMC for cytosolic calcium have not been performed to date.

Of note, while most EET concentrations led to similar responses among all treatments,

MR antagonism as well as chronic exposure of aldosterone-treated cells to EET plus

sEH inhibitor drastically reduced the calcium response to 1 and 10 µM 14,15 EET.

Reduced intracellular calcium should also translate into vasodilation[38]. Therefore,

sEH inhibition might constitute a means to increase vasodilation to EETs.

Our study has shown that aldosterone-mediated endothelial dysfunction does not

seem to impact on the EET pathway. Experimental trials have shown that soluble

epoxide hydrolase inhibition was able to reverse DOCA-induced hypertension and

endothelial dysfunction[15]. Thus, although soluble epoxide hydrolase inhibition

probably cannot be simply understood as a reversal of the underlying

pathophysiological processes, its benefits in conditions of mineralocorticoid excess

should be further investigated. Beneficial effects might result from destiffening

endothelial cells[39] by reducing ENaC channels[40] which would allow for increased

NO release[41]. Another favorable mode of action in mineralocorticoid excess might

be EET-mediated attenuation of inflammation[42]. Moreover, EETs have also been

described to ameliorate insulin resistance[43]. Since patients with PA frequently show

impaired insulin secretion[44], an epoxide hydrolase inhibitor therapy might - again -

be of potential additional benefit for patients with PA.

The main limitations of our study consist in its *in vitro* nature precluding any study of

heterocellular communication or interaction with systemic factors. In addition, the time

scale at which the exposure to aldosterone excess can be studied in primary cell

culture systems does not mirror the prolonged exposure in humans.

On the other hand, we believe that a particular point of strength is that we tried to

account for both mineralocorticoid and glucocorticoid effects at both steroid receptors,

a situation akin to that *in vivo*. Further, our EET quantification underwent an extensive

validation protocol according to standards of the European Medicines Agency[20]

(Supplemental Tables S2 and S3).

Perspectives

A substantial proportion of patients with PA still suffers from hypertension even after

unilateral adrenalectomy and cure of aldosterone excess [45]. Moreover, medication

side effects of MRA hamper therapeutic adherence of patients with bilateral disease.

There is, thus, an urgent need for novel pharmacological strategies targeting the

downstream effector cells of mineralocorticoid excess. Based on our results, it is

therefore tempting to propose that the EET-EDHF axis might be intact in patients with

PA. Therapeutic modulation of EET half-lives with epoxide hydrolase inhibitors in

conditions of mineralocorticoid excess should, therefore, be investigated in future

experiments since the structures responsible for signal transduction in arterial cells –

at least in our primary cell culture system – remained intact.

Acknowledgements

299 We are indebted to Michaela Höhne and Laura Danner for excellent technical

300 assistance.

Sources of funding

H. Schneider was supported by the Clinician Scientist PRogram In Vascular MEdicine

(PRIME) funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) - project number MA 2186/14-1. The study received further support

through the European Research Council (ERC) under the European Union's Horizon

2020 research and innovation programme (grant agreement No [694913] to M.

Reincke) and through the Deutsche Forschungsgemeinschaft (DFG) (CRC/TRR 205,

Project No. 314061271-TRR205 "The Adrenal: Central Relay in Health and Disease"

to F. Beuschlein, M. Reincke and T.A. Williams), and the Else Kröner-Fresenius

Stiftung in support of the German Conn Registry-Else-Kröner Hyperaldosteronism

Registry (2013_A182 and 2015_A171 to M. Reincke).

Conflicts of Interest/Disclosures

None.

References

1   Monticone S, Burrello J, Tizzani D, Bertello C, Viola A, Buffolo F, *et al.*
Prevalence and Clinical Manifestations of Primary Aldosteronism Encountered
in Primary Care Practice. *J Am Coll Cardiol* 2017; 69:1811–1820.

2   Ohno Y, Sone M, Inagaki N, Yamasaki T, Ogawa O, Takeda Y, *et al.*
Prevalence of Cardiovascular Disease and Its Risk Factors in Primary
Aldosteronism: A Multicenter Study in Japan. *Hypertension* 2018; 71:530–537.

3   Hopkins P. Molecular Biology of Atherosclerosis. *Physiol Rev* 2013; 93:1317–
1542.

4   Tsuchiya K, Yoshimoto T, Hirata Y. Endothelial dysfunction is related to
aldosterone excess and raised blood pressure. *Endocr J* 2009; 56:553–9.

5   Matsumoto T, Oki K, Kajikawa M, Nakashima A, Maruhashi T, Iwamoto Y, *et al.*
Effect of Aldosterone-Producing Adenoma on Endothelial Function and Rho-
Associated Kinase Activity in Patients With Primary Aldosteronism.
Hypertension 2015; 65:841–848.

6   Kishimoto S, Oki K, Maruhashi T, Kajikawa M, Matsui S, Hashimoto H, *et al.*
Eplerenone improves endothelial function and arterial stiffness and inhibits
Rho-associated kinase activity in patients with idiopathic hyperaldosteronism. *J*
Hypertens 2018; :1.

7   Blanco-Rivero J, Cachofeiro V, Lahera V, Aras-Lopez R, Márquez-Rodas I,
Salaices M, *et al.* Participation of prostacyclin in endothelial dysfunction
induced by aldosterone in normotensive and hypertensive rats. *Hypertension*
2005; 46:107–112.

8   Lariviere R, Thibault G, Schiffrin EL. Increased endothelin-1 content in blood
vessels of deoxycorticosterone acetate-salt hypertensive but not in
spontaneously hypertensive rats. *Hypertension* 1993; 21:294–300.

9   Leopold JA, Dam A, Maron BA, Scribner AW, Liao R, Handy DE, *et al.*
Aldosterone impairs vascular reactivity by decreasing glucose-6-phosphate
dehydrogenase activity. *Nat Med* 2007; 13:189–197.

10   Nagata D, Takahashi M, Sawai K, Tagami T, Usui T, Shimatsu A, *et al.*
Molecular mechanism of the inhibitory effect of aldosterone on endothelial NO
synthase activity. *Hypertension* 2006; 48:165–171.

11   Garnier A, Bendall JK, Fuchs S, Escoubet B, Rochais F, Hoerter J, *et al.*
Cardiac specific increase in aldosterone production induces coronary
dysfunction in aldosterone synthase-transgenic mice. *Circulation* 2004;

350   110:1819–1825.

Taddei S, Galetta F, Virdis A, Ghiadoni L, Salvetti G, Franzoni F, *et al.* Physical activity prevents age-related impairment in nitric oxide availability in elderly athletes. *Circulation* 2000; 101:2896–2901.

Félétou M. Endothelium-dependent hyperpolarization and endothelial dysfunction. *J Cardiovasc Pharmacol* 2016; 67:373–387.

Liu P, Zhang S, Gao J, Lin Y, Shi G, He W, *et al.* Downregulated Serum 14, 15-Epoxyeicosatrienoic Acid Is Associated With Abdominal Aortic Calcification in Patients With Primary Aldosteronism. *Hypertension* 2018; 71:592–598.

Loch D, Hoey A, Morisseau C, Hammock BO, Brown L. Prevention of hypertension in DOCA-salt rats by an inhibitor of soluble epoxide hydrolase. *Cell Biochem Biophys* 2007; 47:87–98.

Bolz S-S, Fisslthaler B, Pieperhoff S, de Wit C, Fleming I, Busse R, *et al.* Antisense oligonucleotides against cytochrome P450 2C8 attenuate EDHF-mediated Ca 2+ changes and dilation in isolated resistance arteries. *FASEB J* 2000; 14:255–260.

Fisslthaler B, Popp R, Kiss L, Potente M, Harder DR, Fleming I, *et al.* Cytochrome P450 2C is an EDHF synthase in coronary arteries. *Nature* 1999; 401:493–7.

Node K, Huo Y, Ruan X, Yang B, Spiecker M, Ley K, *et al.* Anti-inflammatory Properties of Derived Eicosanoids. *Science (80-)* 1999; 285:1276–1280.

Edin ML, Hamedani BG, Gruzdev A, Graves JP, Lih FB, Arbes SJ, *et al.* Epoxide hydrolase 1 (EPHX1) hydrolyzes epoxyeicosanoids and impairs cardiac recovery after ischemia. *J Biol Chem* 2018; 293:3281–3292.

European Medicines Agency. Guideline on bioanalytical method validation. ; 2011. doi:10.4155/bio.12.44

Archer SL, Gragasin FS, Wu X, Wang S, McMurtry S, Kim DH, *et al.* Endothelium-Derived Hyperpolarizing Factor in Human Internal Mammary Artery Is 11,12-Epoxyeicosatrienoic Acid and Causes Relaxation by Activating Smooth Muscle BKCa Channels. *Circulation* 2003; 107:769–776.

Watanabe H, Vriens J, Prenen J, Droogmans G, Voets T, Nilius B. Anandamide and arachidonic acid use epoxyeicosatrienoic acids to activate TRPV4 channels. *Nature* 2003; 424:434–438.

Gauthier KM, Edwards EM, Falck JR, Reddy DS, Campbell WB. 14,15-Epoxyeicosatrienoic acid represents a transferable endothelium-dependent relaxing factor in bovine coronary arteries. *Hypertension* 2005; 45:666–671.

Larsen BT, Miura H, Hatoum OA, Campbell WB, Hammock BD, Zeldin DC, *et al.* Epoxyeicosatrienoic and dihydroxyeicosatrienoic acids dilate human coronary arterioles via BKCa channels: Implications for soluble epoxide hydrolase inhibition. *Am J Physiol - Hear Circ Physiol* 2006; 290:491–499.

Lee DL, Wilson JL, Duan R, Hudson T, El-Marakby A. Peroxisome proliferator-activated receptor-α activation decreases mean arterial pressure, plasma interleukin-6, and COX-2 while increasing renal CYP4A expression in an acute

model of DOCA-salt hypertension. *PPAR Res* 2011; 2011.
doi:10.1155/2011/502631

26   Fisslthaler B, Popp R, Michaelis UR, Kiss L, Fleming I, Busse R. Cyclic stretch
enhances the expression and activity of coronary endothelium-derived
hyperpolarizing factor synthase. *Hypertension* 2001; 38:1427–1432.

27   Nithipatikom K, Pratt P, Campbell WB. Determination of EETs using microbore
liquid chromatography with fluorescence detection. *Am J Physiol - Hear Circ*
Physiol 2000; 279:H857–H862.

28   Wu S, Moomaw CR, Tomer KB, Falck JR, Zeldin DC. Molecular Cloning and
Expression of CYP2J2, a Human Cytochrome P450 Arachidonic Acid
Epoxygenase Highly Expressed in Heart. *J Biol Chem* 1996; 271:3460–3468.

29   Zeldin DC, DuBois RN, Falck JR, Capdevila JH. Molecular cloning, expression
and characterization of an endogenous human cytochrome P450 arachidonic
acid epoxygenase isoform. Arch. Biochem. Biophys. 1995; 322:76–86.

30   Jiang H, Anderson GD, McGiff JC. Red blood cells (RBCs),
epoxyeicosatrienoic acids (EETs) and adenosine triphosphate (ATP).
Pharmacol Reports 2010; 62:468–474.

31   Ambroisine ML, Favre J, Oliviero P, Rodriguez C, Gao J, Thuillez C, *et al.*
Aldosterone-induced coronary dysfunction in transgenic mice involves the
calcium-activated potassium (BKCa) channels of vascular smooth muscle cells.
Circulation 2007; 116:2435–2443.

32   McCurley A, Pires PW, Bender SB, Aronovitz M, Zhao MJ, Metzger D, *et al.*
Direct regulation of blood pressure by smooth muscle cell mineralocorticoid
receptors. *Nat Med* 2012; 18:1429–1433.

33   Khan M, Meuth AI, Brown SM, Chandrasekar B, Bowles DK, Bender SB.
Aldosterone impairs coronary adenosine-mediated vasodilation via reduced
functional expression of Ca 2+ -activated K + channels. *Am J Physiol Circ*
Physiol 2019; 317:H357–H363.

34   Arlt W, Lang K, Sitch AJ, Dietz AS, Rhayem Y, Bancos I, *et al.* Steroid
metabolome analysis reveals prevalent glucocorticoid excess in primary
aldosteronism. *JCI insight* 2017; 2:0–14.

35   Locovei S, Wang J, Dahl G. Activation of pannexin 1 channels by ATP through
P2Y receptors and by cytoplasmic calcium. *FEBS Lett* 2006; 580:239–244.

36   Good ME, Chiu YH, Poon IKH, Medina CB, Butcher JT, Mendu SK, *et al.*
Pannexin 1 channels as an unexpected new target of the anti-hypertensive
drug spironolactone. *Circ Res* 2018; 122:606–615.

37   Fang X, Weintraub NL, Stoll LL, Spector AA. Epoxyeicosatrienoic acids
increase intracellular calcium concentration in vascular smooth muscle cells.
Hypertension 1999; 34:1242–1246.

38   Amberg GC, Navedo MF. Calcium Dynamics in Vascular Smooth Muscle.
Microcirculation 2013; 20:281–289.

39   Drüppel V, Kusche-Vihrog K, Grossmann C, Gekle M, Kasprzak B, Brand E, *et*
al. Long-term application of the aldosterone antagonist spironolactone prevents

stiff endothelial cell syndrome. *FASEB J* 2013; 27:3652–3659.

40   Pavlov TS, Ilatovskaya D V., Levchenko V, Mattson DL, Roman RJ,
Staruschenko A. Effects of cytochrome P-450 metabolites of arachidonic acid
on the epithelial sodium channel (ENaC). *Am J Physiol - Ren Physiol* 2011;
301:672–681.

41   Oberleithner H, Callies C, Kusche-Vihrog K, Schillers H, Shahin V, Riethmüller
C, *et al.* Potassium softens vascular endothelium and increases nitric oxide
release. *Proc Natl Acad Sci U S A* 2009; 106:2829–2834.

42   He Z, Zhang X, Chen C, Wen Z, Hoopes SL, Zeldin DC, *et al.* Cardiomyocyte-
specific expression of CYP2J2 prevents development of cardiac remodelling
induced by angiotensin II. *Cardiovasc Res* 2015; 105:304–317.

43   Luria A, Bettaieb A, Xi Y, Shieh GJ, Liu HC, Inoue H, *et al.* Soluble epoxide
hydrolase deficiency alters pancreatic islet size and improves glucose
homeostasis in a model of insulin resistance. *Proc Natl Acad Sci U S A* 2011;
108:9038–9043.

44   Fischer E, Adolf C, Pallauf A, Then C, Bidlingmaier M, Beuschlein F, *et al.*
Aldosterone excess impairs first phase insulin secretion in primary
aldosteronism. *J Clin Endocrinol Metab* 2013; 98:2513–2520.

45   Williams TA, Lenders JWM, Mulatero P, Burrello J, Rottenkolber M, Adolf C, *et*
al. Outcomes after adrenalectomy for unilateral primary aldosteronism: An
international consensus on outcome measures and analysis of remission rates
in an international cohort. *Lancet Diabetes Endocrinol* 2017; 5:689–699.

Figure legends

Figure 1: **A, B:** Effect of duration of exposure to 1 nM of aldosterone and co-stimulation with
cortisol on CYP2C8 (**A**) and CYP2J2 (**B**) mRNA expression in human coronary artery
endothelial cells. Each individual relative expression value was normalized to the mean of the
48 h cortisol control. Aldo, aldosterone, Ep, eplerenone. **C, D:** Effect of duration of exposure
to 1 nM of aldosterone and co-stimulation with cortisol on EPHX1 (microsomal epoxide
hydrolase) mRNA expression in human coronary artery smooth muscle cells (**C**) and
endothelial cells (**D**). Aldo, aldosterone, Ep, eplerenone.*: $p < 0.05$ vs. Aldo+cortisol 24 h, t-test.
E, F: Effect of duration of exposure to 1 nM of aldosterone and co-stimulation with cortisol on
EPHX2 (soluble epoxide hydrolase) mRNA expression in human coronary artery smooth
muscle cells (**E**) and endothelial cells (**F**). Aldo, aldosterone, Ep, eplerenone.* $p < 0.05$ vs.
Aldo+cortisol 24 h, t-test.

Figure 2: Endothelial cells were synchronized under serum-free conditions, treated with the
indicated pharmacological agents in serum-free medium for 48 hours, loaded with 10 µM
arachidonic acid and then stimulated either with 1 or 100 µM acetylcholine in HEPES buffer to
provoke the release of EETs. Subsequently, supernatants were collected and quantified via
HPLC-MS/MS. Dashed bright red line marks the values of control cells stimulated with 1 µM
ACh, dashed blue line marks the control cells stimulated with 100 µM ACh. Shown are the
sums of all active EETs (5,6; 8,9; 11,12; 14,15 EET, **A**), sum of all inactive EETs (5,6; 8,9;
11,12; 14,15 DHET, **B**), the ratio of the two sums (sum DHET/sumEET, **C**) as a measure of
expoxide hydrolase activity and the sum of EET and DHET as a time integral and cumulative

measure of EET production as time integral and cumulative measure of total EET production
by CYP-epoxigenases (**D**). ACh, acetylcholine; Aldo, aldosterone; Ep, eplerenone; GSK,
GSK2256294. N=3-9 per group, *p<0.05, **p<0.01, 1-way ANOVA, Tukey.

Figure 3: A: Effect of duration of exposure to 1 nM of aldosterone and co-stimulation with
cortisol on TRPV4 mRNA expression in human coronary artery smooth muscle cells. **B:** Effect
of 48 hours exposure to 1 nM of aldosterone and co-stimulation with cortisol on CACNA1C
mRNA expression in human coronary artery smooth muscle cells. N=3 per group. Aldo,
aldosterone, Ep, eplerenone, RU, mifepristone. *, p<0.05 vs. DMSO; **, p<0.01 vs. DMSO; #,
p<0.05 vs. Aldo; ##. P<.0.01 vs. Aldo; 1-Way ANOVA, Tukey.

Figure 4: A: Effect of exposure to 1 nM of aldosterone and co-stimulation with cortisol on BK_{Ca}
alpha subunit mRNA expression in human coronary artery smooth muscle cells. Aldo,
aldosterone, Ep, eplerenone. N=6 experiments per group,*p<0.05, Dunnett. **B:** Relative
contribution of glucocorticoid receptor stimulation on downregulation of BK_{Ca} alpha subunit
mRNA in SMC. Aldo, aldosterone; Ep, eplerenone; RU, RU486 (mifepristone). N=3-6 per
group.*p<0.05 vs. control (w/o cortisol), ****,p<0.001 vs. control (w/o cortisol); ##, p<0.01 vs.
A 1nM; §§, p<0.01 vs. Aldo+cortisol; $, p<0.05 vs. Aldo+cortisol+Ep. 1-Way-ANOVA,
Tukey..**C, D**: Western blot of cortisol, Aldo+cortisol and Aldo+cortisol+Ep-treated SMC for BK_{Ca}
alpha subunit and beta-actin. N=4 experiments per group.

Figure 5: Calcium transient after EET as observed in single smooth muscle cell calcium
imaging. Shown are intracellular baseline calcium and peak calcium responses after
application of 1 µM 14,15 EET. Aldo, aldosterone; EET, 14,15 epoxyeicostrienoic acid; Ep,
eplerenone. ***, p<0.001; ****, p<0.0001;#, p<0.05 vs. peak cortisol (1-Way ANOVA, Tukey),
n (cells/total number of experiments)= 17/3 (cortisol), 18/3 (Aldo+cortisol), 23/3
(Aldo+cortisol+Ep).

Figure 6: Calcium concentration-response curve for 14,15 EET in fluo4-labelled smooth
muscle cells cultured for 48 hours in the presence of the indicated treatments. Aldo,
aldosterone; EET, 14,15 epoxyeicostrienoic acid; Ep, eplerenone; GSK, GSK2256294. N=9-
21 recordings per concentration. *, p<0.05 (Aldo+cortisol vs. Aldo+cortisol+EET+GSK and
Aldo+cortisol vs. Aldo+cortisol+Ep), 2-Way ANOVA, Dunnett.

A

B

C

D

E

F