Pharmacokinetics of Orally Administered GS-441524 in Dogs

Victoria C. Yan[1]*, Sunada Khadka[1,2], Kenisha Arthur[1], Cong-Dat Pham[1], Matthew J.

Yan[3], Alexander J. Yan[3], Jeffrey J. Ackroyd[1], Dimitra K. Georgiou[1], and Florian L.

Muller[1]

[1]Department of Cancer Systems Imaging, University of Texas MD Anderson Cancer

Center, Houston, TX, USA

[2]Department of Cancer Biology, University of Texas MD Anderson Cancer Center,

Houston, TX, USA

[3]Department of Chemistry, Boston College, Chestnut Hill, MA, USA

*Email: victoriacyanide@gmail.com Twitter: @victoriacyanide

Abstract.

Despite being FDA-approved for COVID-19, the clinical efficacy of remdesivir (Veklury[®])

remains contentious. We previously described the pharmacokinetic, pharmacodynamic

and toxicological rationales on the greater suitability of its parent nucleoside, GS-

441524, for COVID-19 treatment. Here, we assess the oral bioavailability of GS-441524

in beagle dogs and show that plasma concentrations approximately 24-fold higher than

the EC_{50} against SARS-CoV-2 are easily and safely sustained. These data support

translation of GS-441524 as an oral agent for COVID-19.

Remdesivir (RDV) is currently the only FDA-approved anti-viral drug for the treatment of

COVID-19 despite exhibiting just modest efficacy in one double-blind, placebo-

controlled randomized clinical trial (RCT) (1); other RCTs have thus far found no

statistically significant improvement in mortality (2) or time to clinical improvement (3).

As a phosphoramidate prodrug of the McGuigan class (4), RDV is structurally

susceptible to conversion to its active nucleoside triphosphate (NTP; GS-443902) form

by enzymes that are abundant in the liver (CES1/CTSA/HINT1) but minimally expressed

in alveolar type 2 cells (AT2) (5), the cell type most susceptible to SARS-CoV-2

infection (6). Preferential liver metabolism of RDV results in on-target dose limiting

toxicity (DLT) that precludes dose escalation despite its modest clinical performance (7,

8). Such shortcomings are exacerbated by the hydrophobic nature of RDV, which

requires complex excipients that could implicate kidney function (9, 10). Another major

drawback with RDV is its requisite intravenous (IV) administration (11), making

outpatient therapy, early intervention, and prophylaxis impractical.

Cognizant of these limitations, we have asserted that the parent nucleoside of RDV,

GS-441524, would be better suited for the treatment for COVID-19 (5). GS-441524 is

the persistent, predominant metabolite in plasma following IV infusion of RDV in

preclinical species (12–14) and in humans with a half-life ($T_{1/2}$) of >24 h (8, 15). The

extent to which GS-441524 contributes to the overall anti-SARS-CoV-2 activity of RDV

when administered to patients remains unclear. In contrast to RDV, which is

preferentially bioconverted to GS-443902 by liver-abundant enzymes (16, 17), GS-

441524 is bioconverted to GS-443902 by nucleoside kinases (likely adenosine kinase,

ADK) that are broadly expressed across all tissues. Due to its demonstrably better

safety profile (7, 18), a significant advantage that GS-441524 possesses over RDV is

the possibility for dose escalation without liver-related DLTs, as this would increase the

concentration of bioactive GS-443902 in AT2 cells. Cell-based studies have shown that

GS-441524 is a potent inhibitor of SARS-CoV-2-infected cells, with EC_{50} values on the

same order of magnitude as that of RDV (EC_{50}= 0.47-1.09 μM) (19). Efficacy studies

conducted in cats with natural presentations feline infectious peritonitis (FIP) as a result

of infection by the closely related feline coronavirus (FCoV) have demonstrated up to 96%

cure rate with subcutaneously (SC) administered GS-441524 (20–22). Efficacy studies

conducted in mice infected with either SARS-CoV-2 or murine hepatitis virus (MHV, a

closely related coronavirus) have shown that GS-441524 is capable of reducing viral

loads in pathologically relevant organs without obvious adverse events (23). Given the

scarcity of simple outpatient treatment options for COVID-19 (24), these especially

encouraging data warrant translation of GS-441524 to the clinic. Here, we provide

pharmacokinetic (PK) evidence in dogs supporting the ability for GS-441524 to be

investigated as an oral agent for COVID-19.

To validate our assertion that McGuigan prodrugs such as RDV are heavily subject to

first-pass metabolism, we first conducted a single-dose, equimolar comparison between

orally (PO) administered GS-441524 (6.5 mg/kg) and RDV (13 mg/kg) in dogs. Male

beagles (N=1 per group) were administered excipient-less capsule formulations of either

GS-441524 or RDV and plasma concentrations of GS-441524 were evaluated at

predetermined timepoints (**Figure 1a**). No adverse events were observed in either

dosing group. As expected, plasma concentrations of GS-441524 following

administration of RDV were poor, with C_{max} values roughly 25-fold lower than that

observed when GS-441524 was administered directly (172 vs. 4580 ng/mL, respectively;

Figure 1a, c, Supplementary Data File 1). Interestingly, there was an observable

difference in T_{max} values, with plasma concentrations of GS-441524 peaking at

approximately 3 h following RDV administration versus 1 h following direct dosing of

GS-441524. This shift in T_{max} values with RDV PO administration suggests a

mechanism of systemic release similar to that observed for the McGuigan prodrug

sofosbuvir, wherein rapid hepatic extraction of intact prodrug forms a reservoir of active

NTP and hydrolyzed nucleoside—the latter of which is then slowly released into

systemic circulation (25). Given that plasma concentrations of GS-441524 following

RDV administration are below the range of reported anti-SARS-CoV-2 EC_{50} values and

that long-term PO dosing of RDV at 13 mg/kg is almost certainly therapeutically

prohibitive in humans due to hepatotoxicity concerns (8), these data allude to the

infeasibility of administering RDV PO for COVID-19. At the same time, we find that

direct PO administration of GS-441524 results in plasma concentration exceeding the

range of reported anti-SARS-CoV-2 EC_{50} values for at least 8 h (**Figure 1a,**

Supplementary Data File 1). Peak concentrations of GS-441524 reached 15.45 µM

and were obtained at approximately 1 h (**Figure 1a, c**), indicating high oral absorption of

GS-441524 in dog even in the absence of excipients.

Prior studies assessing the oral bioavailability (OBV, F%) of GS-441524 in dogs

following both IV (2 mg/kg) and PO (5 mg/kg) administration have found that the drug is

efficiently absorbed, with an OBV of 85% (NCATS OpenData Portal). Such studies

examined PO absorption of GS-441524 using a solution formulation prepared at a final

concentration of 2.5 mg/mL. We sought to determine whether similarly favorable F%

could be achieved using an excipient-less capsule formulation, which would greatly

ease outpatient administration. The wide range of F% observed in other preclinical

species (**Table 1**; NCATS OpenData Portal) and the unusual solubility properties of GS-

441524 are reminiscent of that observed with the FDA-approved nucleoside analogue

acyclovir (**Table 1**), which was ultimately formulated as an excipient-less tablet (26).

Direct comparison of PK parameters between solution and capsule formulations of GS-

441524 indicates a similar pattern of high drug absorption, with T_{max} values of 0.5 and 1

h, respectively (**Figure 1b**). Between solution and capsule formulations, PK parameters

were generally similar; it should be noted that the capsule dose was slightly higher than

the solution dose (5 mg/kg vs. 6.5 mg/kg). Adjusting for sampling timeframes, these

data indicate that the C_{max} value was higher with the solution formulation (5060 vs. 4580

ng/mL) but the AUC value were somewhat higher with the capsule formulation (17916

vs.19151 ng/mL; **Figure 1c**). Such observations appear consistent with the general

observation that liquid formulations tend to be more readily absorbed than their pill

counterparts (27). Nevertheless, the estimated OBV using this capsule formulation

remains high at about 76% (**Figure 1b**). These data hint at the feasibility of using an

excipient-less pill formulation for GS-441524 for outpatient treatment.

There are some limitations associated with this study. First, the sample size in the

capsule study is small, which may not capture possible variability associated with this

formulation. Second, nucleoside analogues generally tend to exhibit higher OBV in dogs

than in other preclinical species perhaps due to the presence of a paracellular

nucleoside transporter that is absent in humans and non-human primates (28). As a

result, the OBV of nucleoside analogues in dogs tends to overestimate that observed in

humans (**Supplementary Data File 2**). While not specifically explored in this study, it

should be noted that—at the other end of the OBV spectrum—OBV of nucleoside

analogues in non-human primates tend to under-predict that observed in humans

(**Supplementary Data File 3**). Nevertheless, these data suggest the feasibility of using

a simple, excipient-less capsule formulation of GS-441524. As a prodrug inhibitor of the

SARS-CoV-2 RNA-dependent RNA polymerase (RdRp), GS-441524 is aptly poised to

demonstrate consistent efficacy among new mutations of SARS-CoV-2, as RdRp is

much less susceptible to efficacy-altering mutations than is the spike protein (29, 30).

Thus, clinical translation of GS-441524 is imperative.

Drug formulation.

For capsule studies, GS-441524 and RDV were purchased at the highest commercially available quality from MedKoo Biosciences; purity was verified by ultra-performance liquid chromatography mass spectrometry (UPLC-MS) and nuclear magnetic resonance (NMR) spectroscopy (^{1}H, ^{13}C) in-house. For capsule studies, gelatin capsules (size 5, XPRS Nutra) were tightly packed with either GS-441524 (65 mg) or RDV (136.74 mg) without additional excipients. For solution studies, GS-441524 was purchased at the highest commercially available from AK Scientific and characterized by NCATS. Formulations for PO and IV studies conducted by NCATS are described on the NCATS OpenData Portal. Briefly, GS-441524 was dissolved in a solution containing 5% ethanol, 30% propylene glycol, 45% PEG-400, 20% water with 1 equivalent HCl for a final concentration of 2.5 mg/mL.

Single dose GS-441524 and RDV in dogs via capsule formulation.

All capsule form studies were performed at Charles River Laboratories (Wilmington, MA) with IACUC approval (#20236536). Fasted male adult beagles (10 kg; N=1 per compound) were administered either GS-441524 (6.5 mg/kg) or RDV (13 mg/kg). Plasma samples were taken for PK analysis at the following timepoints (h): -0.5, 0.5, 1, 3, 6, 8, 24. Animals were monitored continuously by veterinarians for any clinically relevant abnormalities during dosing and sample collection. PO solution and IV studies were performed by NCATS as described on the NCATS OpenData Portal with relevant

committee and regulatory approval. Data from all studies were (re-)analyzed using

PKSolver 2.0 and graphs were generated using GraphPad Prism 8.

Plasma pharmacokinetics

For capsule studies, plasma levels of GS-441524 were analyzed at Covance, Inc

(Princeton, NJ) on a fee-for-service basis using a liquid chromatography mass

spectrometry (LC-MS) assay previously described for quantification of GS-441524

following IV administration of RDV in NHP (12).

Author Contributions.

V.C.Y. analyzed data and wrote the manuscript. S.K., K.A., D.K.G., and J.J.A. provided

technical assistance. V.C.Y. and F.L.M. conceived and oversaw the study.

Acknowledgements.

This work was supported by the COVID-19 Early Treatment Fund (CETF). S.K. is

supported by the MD Anderson CPRIT Research Training Program Grant (RP170067)

and the Larry Deaven PhD Fellowship in Biomedical Sciences. We thank Gilead

Sciences for permission to use their assay established at Covance, Inc. to quantify GS-

441524 in plasma and NCATS for conducting PK studies.

Figure 1. Plasma concentrations of GS-441524 following a single oral dose of remdesivir or GS-441524 in dogs. (A) Head-to-head PK comparison following a single equimolar dose of remdesivir (black, 13.6 mg/kg) and GS-441524 (orange, 6.5 mg/kg) in male beagle dogs (N=1 per compound). Both compounds were administered in capsule form. Plasma concentrations of GS-441524 following compound administration are plotted for the following timepoints (h): 0.5, 1, 3, 6, 8, 24. A focused view of GS-441524 concentrations following oral administration of remdesivir is shown in top right corner. (B) Comparison of plasma concentrations of GS-441524 following oral administration as a solution (black, 5 mg/kg; N=3) or as a capsule (orange, 6.5 mg/kg; N=1). (C) Mean PK parameters following various routes of administration of GS-441524 and RDV. Raw values for GS-441524 dosed IV and PO dosed as a solution are adapted from NCATS OpenData Portal and have been re-calculated to match the

199 sampling timeframe of the capsule studies (T=0.5-24 h). All PK parameters were

200 calculated using PKSolver 2.0. In panels A and B, dotted lines correspond to EC_{50}

201 (bottom) and EC_{90} (top) values reported for GS-441524 in SARS-CoV-2-infected Calu3

202 cells (19).

Species, F%	Acyclovir	GS-441524
Human	10-20	15-30*
Monkey	3.7	8.3
Dog	54-90	85-93

Table 1. Oral bioavailabilities of acyclovir and GS-441524 in preclinical species

are similar. GS-441524 exhibits a similar pattern of F% as acyclovir across preclinical

species. F% for GS-441524 were obtained from the NCATS OpenData Portal and F%

for acyclovir were obtained from the FDA fact sheet on Zovirax (human), Laskin et al.

Clin. Pharm. (1983, monkey) (31) and Krasny et al. *J. Pharm. Exp. Ther.* (1981, dog)

(32). *Anticipated F% of GS-441524 in humans.

References.

1. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, Hohmann E, Chu HY, Luetkemeyer A, Kline S, Lopez de Castilla D, Finberg RW, Dierberg K, Tapson V, Hsieh L, Patterson TF, Paredes R, Sweeney DA, Short WR, Touloumi G, Lye DC, Ohmagari N, Oh M, Ruiz-Palacios GM, Benfield T, Fätkenheuer G, Kortepeter MG, Atmar RL, Creech CB, Lundgren J, Babiker AG, Pett S, Neaton JD, Burgess TH, Bonnett T, Green M, Makowski M, Osinusi A, Nayak S, Lane HC. 2020. Remdesivir for the Treatment of Covid-19 — Final Report. N Engl J Med https://doi.org/10.1056/NEJMoa2007764.

2. WHO Solidarity Trial Consortium. 2020. Repurposed Antiviral Drugs for Covid-19 - Interim WHO Solidarity Trial Results. N Engl J Med 1–15.

3. Wang Y, Zhang D, Du G, Du R, Zhao J, Jin Y, Fu S, Gao L, Cheng Z, Lu Q, Hu Y, Luo G, Wang K, Lu Y, Li H, Wang S, Ruan S, Yang C, Mei C, Wang Y, Ding D, Wu F, Tang X, Ye X, Ye Y, Liu B, Yang J, Yin W, Wang A, Fan G, Zhou F, Liu Z, Gu X, Xu J, Shang L, Zhang Y, Cao L, Guo T, Wan Y, Qin H, Jiang Y, Jaki T, Hayden FG, Horby PW, Cao B, Wang C. 2020. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. Lancet 395:1569–1578.

4. Mcguigan C, Devine KG, O'oonnor TJ, Galpin SA, Jeffries DJ, Kinchington D. 1990. Synthesis and evaluation of some novel phosphoramidate derivatives of 3'-azido-3'-deoxythymidine (AZT) as anti-HIV compoundsAntiviral Chemistry &

Chemotherapy.

5.   Yan VC, Muller FL. 2020. Advantages of the Parent Nucleoside GS-441524 over

Remdesivir for Covid-19 Treatment. ACS Med Chem Lett 11:1361–1366.

6.   Schaefer I-M, Padera RF, Solomon IH, Kanjilal S, Hammer MM, Hornick JL, Sholl

LM. 2020. In situ detection of SARS-CoV-2 in lungs and airways of patients with

COVID-19. Mod Pathol 33:2104–2114.

7.   Xu Y, Barauskas O, Kim C, Babusis D, Murakami E, Kornyeyev D, Lee G, Stepan

G, Perron M, Bannister R, Schultz BE, Sakowicz R, Porter D, Cihlar T, Feng JY.

2020. Off-target In Vitro Profiling Demonstrates that Remdesivir Is a Highly

Selective Antiviral Agent. Antimicrob Agents Chemother

https://doi.org/10.1128/aac.02237-20.

8.   Humeniuk R, Mathias A, Cao H, Osinusi A, Shen G, Chng E, Ling J, Vu A,

German P. 2020. Safety, Tolerability, and Pharmacokinetics of Remdesivir, An

Antiviral for Treatment of COVID-19, in Healthy Subjects. Clin Transl Sci

cts.12840.

9.   Yan VC, Muller FL. 2020. Captisol and GS-704277, but not GS-441524, are

credible mediators of remdesivir's nephrotoxicity. Antimicrob Agents Chemother

64:01920–1920.

10. European Medicines Agency. 2020. Remdesivir: Summary on compassionate use

EMA/178637/2020 - Rev. 1.

11. Food and Drug Administration. 2020. VEKLURY® (remdesivir).

12. Warren TK, Jordan R, Lo MK, Ray AS, Mackman RL, Soloveva V, Siegel D,

Perron M, Bannister R, Hui HC, Larson N, Strickley R, Wells J, Stuthman KS, Van

Tongeren SA, Garza NL, Donnelly G, Shurtleff AC, Retterer CJ, Gharaibeh D,

Zamani R, Kenny T, Eaton BP, Grimes E, Welch LS, Gomba L, Wilhelmsen CL,

Nichols DK, Nuss JE, Nagle ER, Kugelman JR, Palacios G, Doerffler E, Neville S,

Carra E, Clarke MO, Zhang L, Lew W, Ross B, Wang Q, Chun K, Wolfe L,

Babusis D, Park Y, Stray KM, Trancheva I, Feng JY, Barauskas O, Xu Y, Wong P,

Braun MR, Flint M, McMullan LK, Chen S-S, Fearns R, Swaminathan S, Mayers

DL, Spiropoulou CF, Lee WA, Nichol ST, Cihlar T, Bavari S. 2016. Therapeutic

efficacy of the small molecule GS-5734 against Ebola virus in rhesus monkeys.

Nature 531:381–385.

13. Sheahan TP, Sims AC, Graham RL, Menachery VD, Gralinski LE, Case JB, Leist

SR, Pyrc K, Feng JY, Trantcheva I, Bannister R, Park Y, Babusis D, Clarke MO,

Mackman RL, Spahn JE, Palmiotti CA, Siegel D, Ray AS, Cihlar T, Jordan R,

Denison MR, Baric RS. 2017. Broad-spectrum antiviral GS-5734 inhibits both

epidemic and zoonotic coronaviruses. Sci Transl Med 9:eaal3653.

14. Williamson BN, Feldmann F, Schwarz B, Meade-White K, Porter DP, Schulz J,

Van Doremalen N, Leighton I, Yinda CK, Pérez-Pérez L, Okumura A, Lovaglio J,

Hanley PW, Saturday G, Bosio CM, Anzick S, Barbian K, Cihlar T, Martens C,

Scott DP, Munster VJ, De Wit E. 2020. Clinical benefit of remdesivir in rhesus

macaques infected with SARS-CoV-2. Nature 585:273–276.

15. Tempestilli M, Caputi P, Avataneo V, Notari S, Forini O, Scorzolini L, Marchioni L,

Bartoli TA, Castilletti C, Lalle E, Capobianchi MR, Nicastri E, D'Avolio A, Ippolito

G, Agrati C. 2020. Pharmacokinetics of remdesivir and GS-441524 in two critically

ill patients who recovered from COVID-19. J Antimicrob Chemother dkaa239:1–4.

16.   Murakami E, Wang T, Babusis D, Lepist E-I, Sauer D, Park Y, Vela JE, Shih R,

Birkus G, Stefanidis D, Kim CU, Cho A, Ray AS. 2014. Metabolism and

Pharmacokinetics of the Anti-Hepatitis C Virus Nucleotide Prodrug GS-6620.

Antimicrob Agents Chemother 58:1943–1951.

17.   Lawitz E, Hill J, Marbury T, Hazan D, Gruener D, Webster L, Majauskas R,

Morrison R, DeMicco M, German P, Stefanidis D, Svaroskaia E, Arterburn S, Ray

A, Rossi S, McHutchinson J, Rodriguez-Torres M. 2012. GS-6620, A Liver-

Targeted Nucleotide Prodrug, Exhibits Antiviral Activity and Favorable Safety

Profile Over 5 Days in Treatment Naïve Chronic HCV Genotype 1 SubjectsEASL

47th Annual Meeting. EASL 47th Annual Meeting, Barcelona.

18.   Lo MK, Jordan R, Arvey A, Sudhamsu J, Shrivastava-Ranjan P, Hotard AL, Flint

M, McMullan LK, Siegel D, Clarke MO, Mackman RL, Hui HC, Perron M, Ray AS,

Cihlar T, Nichol ST, Spiropoulou CF. 2017. GS-5734 and its parent nucleoside

analog inhibit Filo-, Pneumo-, and Paramyxoviruses. Sci Rep 7:43395.

19.   Pruijssers AJ, George AS, Schä A, Baric RS, Denison MR, Sheahan TP. 2020.

Remdesivir Inhibits SARS-CoV-2 in Human Lung Cells and Chimeric SARS-CoV

Expressing the SARS-CoV-2 RNA Polymerase in Mice. Cell Rep 107940.

20.   Pedersen NC, Perron M, Bannasch M, Montgomery E, Murakami E, Liepnieks M,

Liu H. 2019. Efficacy and safety of the nucleoside analog GS-441524 for

treatment of cats with naturally occurring feline infectious peritoniti. J Feline Med

Surg 21:271–281.

21.   Murphy BG, Perron M, Murakami E, Bauer K, Park Y, Eckstrand C, Liepnieks M,

Pedersen NC. 2018. The nucleoside analog GS-441524 strongly inhibits feline

infectiousperitonitis (FIP) virus in tissue culture and experimental cat infection

studies. Vet Microbiol 219:226–233.

22. Dickinson PJ, Bannasch M, Thomasy SM, Murthy VD, Vernau KM, Liepnieks M,

Montgomery E, Knickelbein KE, Murphy B, Pedersen NC. 2020. Antiviral

treatment using the adenosine nucleoside analogue GS-441524 in cats with

clinically diagnosed neurological feline infectious peritonitis. J Vet Intern Med

jvim.15780.

23. Li Y, Cao L, Li G, Cong F, Li Y, Sun J, Luo Y, Chen G. 2021. Remdesivir

Metabolite GS-441524 E ff ectively Inhibits SARS-CoV ⬚ 2 Infection in Mouse

Models. J Med Chem https://doi.org/10.1021/acs.jmedchem.0c01929.

24. Lenze EJ, Mattar C, Zorumski CF, Stevens A, Schweiger J, Nicol GE, Miller JP,

Yang L, Yingling M, Avidan MS, Reiersen AM. 2020. Fluvoxamine vs Placebo and

Clinical Deterioration in Outpatients with Symptomatic COVID-19: A Randomized

Clinical Trial. JAMA 324:2292–2300.

25. Wang T, Babusis D, Park Y, Niu C, Kim C, Zhao X, Lu B, Ma B, Muench RC,

Sperger D, Ray AS, Murakami E. 2020. Species differences in liver accumulation

and metabolism of nucleotide prodrug sofosbuvir. Drug Metab Pharmacokinet

35:334–340.

26. Food and Drug Administration. 2002. ZOVIRAX® (acyclovir).

27. Levene DL. 1973. The absorption of potassium chloride: Liquid vs. tablet. Can

Med Assoc J 108:1480–1483.

28. Hammond JR, Stolk M, Archer RGE, McConnell K. 2004. Pharmacological

analysis and molecular cloning of the canine equilibrative nucleoside transporter 1.

Eur J Pharmacol 491:9–19.

29.   Agostini ML, Andres EL, Sims AC, Graham RL, Sheahan TP, Lu X, Clinton Smith

E, Brett Case J, Feng JY, Jordan R, Ray AS, Cihlar T, Siegel D, Mackman RL,

Clarke MO, Baric RS, Denison MR, Agostini CM, Gallagher T. 2018. Coronavirus

Susceptibility to the Antiviral Remdesivir (GS-5734) Is Mediated by the Viral

Polymerase and the Proofreading Exoribonuclease Downloaded from. MBio

9:ee00221-18.

30.   Mahase E. 2021. Covid-19□: Novavax vaccine efficacy is 86 % against UK

variant and 60 % against South African variant. BMJ 2021.

31.   Laskin OL. 1983. Clinical Pharmacokinetics of Acyclovir. Clin Pharmacokinet

8:187–201.

32.   Krasny HC, De Miranda P, Blum MR, Elion GB. 1981. Pharmacokinetics and

bioavailability of acyclovir in the dog. J Pharmacol Exp Ther 216:281–288.