The prevalence, predominance, and metabolic potentials of *Candidatus*

Prochlorobium terpiosii* in the coral-killing sponge, *Terpios hoshinota

Running title: Dominant cyanobiont in *Terpios hoshinota*

Yu-Hsiang Chen[1,2,3]*, Hsing-Ju Chen[3]*, Cheng-Yu Yang[3], Jia-Ho Shiu[3], Daphne Z.

Hoh[3,4,5], Pei-Wen Chiang[3], Wenhua Savanna Chow[3,4,5], Chaolun Allen Chen[3,4], Tin-Han

Shih[3], Szu-Hsien Lin[3], Chi-Ming Yang[3], James Davis Reimer[6,7], Euichi Hirose[6], Budhi

Hascaryo Iskandar[8], Hui Huang[9], Peter J. Schupp[10], Chun Hong James Tan[11,12], Hideyuki

Yamashiro[7], Ming-Hui Liao[3], Sen-Lin Tang[2,3,4,5#]

[1]Bioinformatics Program, Taiwan International Graduate Program, National Taiwan

University, Taipei, Taiwan

[2]Bioinformatics Program, Institute of Information Science, Taiwan International Graduate

Program, Academia Sinica, Taipei, Taiwan

[3]Biodiversity Research Center, Academia Sinica, Taipei, Taiwan

[4]Biodiversity Program, Taiwan International Graduate Program, Academia Sinica and

National Taiwan Normal University, Taipei, Taiwan

[5]Department of Life Science, National Taiwan Normal University, Taipei, Taiwan

[6]Department of Chemistry, Biology and Marine Science, Faculty of Science, University

of the Ryukyus, Nishihara, Okinawa, Japan

[7]Tropical Biosphere Research Center, University of the Ryukyus, Nishihara, Okinawa,

Japan

[8]Department of Fishery Resources Utilization, Faculty of Fisheries and Marine Science,

Bogor Agricultural University, Bogor, Indonesia

[9]Tropical Marine Biological Research Station in Hainan, Chinese Academy of Sciences,

Sanya, China

[10]Institute of Chemistry and Biology of the Marine Environment, University of

Oldenburg, Wilhelmshaven, Germany

[11]Faculty of Science and Marine Environment, Universiti Malaysia Terengganu, Kuala

Nerus, Terengganu, Malaysia

[12]Institute of Oceanography and Environment, Universiti Malaysia Terengganu, Kuala

Nerus, Terengganu, Malaysia

*Co-first authors

Corresponding author

Corresponding author information:

Dr. Sen-Lin Tang

Biodiversity Research Center, Academia Sinica

E-mail: sltang@gate.sinica.edu.tw

Address: A305, Interdisciplinary Research Building for Science and Technology, 128

Academia Road, Sec. 2, Nankang, Taipei 11529, Taiwan.

Tel: +886-2-27893863

Fax: +886-2-27890844

The authors declare no competing interests

Keywords: Coral-killing sponge, *Terpios hoshinota*, Cyanobacteriosponge, Symbiont,

Bacterial community

Abstract

Terpios hoshinota is a ferocious, space-competing sponge that kills a variety of stony corals by overgrowth. Outbreaks of this species have led to intense coral reef damage and declines in living corals on the square kilometer scale in many geographical locations. Our large-scale 16S rRNA metagenome study across three oceans revealed that the core microbiome of *T. hoshinota* included OTUs related to *Prochloron*, *Endozoicomonas*, *Pseudospirillum*, SAR116, *Magnetospira*, and *Ruegeria*. A *Prochloron*-like OTU was the most dominant cyanobacterium in *T. hoshinota* in the western Pacific Ocean, South China Sea, and Indian Ocean. The high-quality draft genome of the *Prochloron*-like cyanobacterium and our pigment analysis revealed that, unlike the previously described *Prochloron* species and their genomes, this bacterium had phycobiliproteins and phycobilins and lacked chlorophyll *b*. Furthermore, the phylogenetic analyses demonstrated that the bacterium was phylogenetically distinct to *Prochloron*, strongly suggesting that it should be a sister taxon to *Prochloron*; we therefore proposed this symbiotic cyanobacterium as a novel species under a new genus: *Candidatus* Prochlorobium terpiosii. In addition, we also characterize the metabolic potentials of the cyanobacterium in carbon and nitrogen cycling and propose a model for the interaction between *Ca*. Pb. terpiosii LD05 and *T. hoshinota*. This study builds the foundation for *T.*

73   *hoshinota*'s microbiome and paves a way for understanding the ecosystem, invasion

74   mechanism, and causes of outbreak of this coral-killing sponge.

Introduction

The coral-killing sponge, *Terpios hoshinota*, has received attention since its

outbreaks were detected in several coral reefs in the western Pacific Ocean, South China

Sea, and Indian Ocean [1-7]. The encrusting sponge can kill various stony corals by

overgrowth [8]. It grows up to 23 mm per month [9], and its fast growing and competitive

nature enables it to kill scleractinian corals quickly and at a high rate (30–80%) across

biogeographic regions [4, 5, 7, 10]. For instance, *T. hoshinota* overgrowth killed 30% of

corals in some coral reefs of Green Island, Taiwan in just one year [2]. Similarly, the

sponge jeopardizes coral reefs in numerous regions of Indonesia, Malaysia, Japan, India,

and the Great Barrier Reef [3, 10-12]. The gradual spread of *T. hoshinota* makes it a

serious threat to coral reefs. However, little is known about what causes its outbreaks.

Sponges are commonly known to harbor complex microbial communities, and

symbiotic microorganisms play vital roles in the development, health, and nutrient

acquisition of their hosts [13]. The microbe and host form an ecological unit called a

holobiont. In *T. hoshinota*, the sponge associates with a bacterial community of relatively

low diversity that is dominated by cyanobacteria [14]. Ultrastructural observations have

clearly shown that cyanobacteria are densely distributed in the sponge, contributing to

50% of the total cell volume [1, 14, 15]. The blackish color of *T. hoshinota* is also

attributed to cyanobacteria [14]. Accordingly, the sponge is called cyanobacteriosponge.

Several studies have shown that cyanobacteria play important roles in the sponge's

growth and competition with corals. For instance, high numbers of cyanobacteria can be

observed in the sponge's larvae [16, 17], suggesting that they are transmitted vertically

during embryogenesis [18]. On the other hand, *in situ* light shading was shown to

discontinue *T. hoshinota* expansion [19]. Even short-term shading can cause a long-term

decrease in the biomass of symbiotic cyanobacteria and lead to irreversible damage to *T.*

hoshinota [20]. Furthermore, one study found that, when *T. hoshinota* encountered a

variety of corals, the sponge formed a hairy tip structure packed with dense cyanobacteria

[8]. These results indicate that cyanobacteria are vital not only for *T. hoshinota* growth,

but also for its overgrowth of corals.

Though important, the identity and role of symbiotic cyanobacteria are still unclear.

In 2015, Yu *et al*. isolated and cultivated *Myxosarcina* sp. GI1, a baeocytous

cyanobacterium, from *T. hoshinota* at Green Island [21]. However, electron microscopy

has not identified vegetative cell aggregates of baeocyte in *T. hoshinota* [14-16].

Moreover, our previous study analyzed 16S ribosomal RNA gene sequences from Green

Island and found that the dominant cyanobacterium in *T. hoshinota* is a novel species

closely related to *Prochloron* sp. [14]. *Prochloron* is an obligate symbiont of some

ascidians; it carries chlorophyll *b* and lacks phycobilin, which is unusual in

cyanobacteria. However, a pigment analysis reported that the cyanobacteria in *T.*

hoshinota contained phycobilins but lacked chlorophyll *b*, which is typical of

cyanobacteria [15]. Hence, the identity and characteristics of the predominant

cyanobacterium in *T. hoshinota* remain uncertain, as does the question of whether its

predominance is universal. On the other hand, since cyanobacteria are attributed to the

sponge's health and invasion capacity, the ecological relationship between *T. hoshinota*

and its cyanobacteria needs to be determined.

Other bacteria in the *T. hoshinota* microbiome also contribute to holobiont functions.

Microbial community in sponge can be shaped by host-related factors, such as the

immune system and nutrient exchanges [22, 23]. Strict host selective pressure can

stabilize the microbial community [24]. On the other hand, the community can also be

determined by environmental factors, such as light availability, pH, and temperature.

Nevertheless, no study has investigated the microbiome of *T. hoshinota* under different

biogeographical backgrounds. Hence, this study investigated the bacterial community *T.*

hoshinota across different oceans to examine whether the bacterial community is stable

and host-specific or depends on geography.

Most sponge symbionts, including the dominant cyanobacteria in *T. hoshinota*, are

uncultivable. Metagenomic methods can be used to determine the microbial composition

in *T. hoshinota*, and knowing this composition will help us understand the ecosystem of

the *T. hoshinota* holobiont. In this study, we used 16S rRNA-based metagenomic analysis

to investigate the bacterial compositions and diversities of *T. hoshinota* samples from a

wide geographical range across the western Pacific Ocean, South China Sea, and Indian

Ocean. Despite the geographical variation in bacterial communities, *T. hoshinota* contains

a stable core community that accounts for most of the bacterial composition. Inside this

core community, we found that a single species predominated the cyanobacteria in *T.*

hoshinota from different locations. The draft genome of this predominant cyanobacterium

was reconstructed by whole-genome shotgun sequencing. The following genomic and

comparative genomic analyses elucidated the phylogenetic affiliation and taxonomy of

the dominant symbiotic bacterium in *T. hoshinota* and putative symbiotic interactions

between the bacterium and the host.

Materials and methods

Sample collection

Sixty-one sponge samples, summarized in Supplementary table 1, were collected

from 15 coastal sites in the western Pacific Ocean, South China Sea, and Indian Ocean.

The collection sites in the western Pacific Ocean included Okinoerabu (JPOKE), Bise

(JPBS), Miyako (JPMYK), Ishigaki (JPISG), Lyudao (TWLD), Lanyu (TWLY), Kenting

(TWKT), Guam (USGU), and the Great Barrier Reef (AUGBR). The collection sites in

the South China Sea included Yongxing (CNYX), Taiping (TWTP), Redang (MYRD),

and Pari (IDPR). Indian Ocean samples were collected from Faafu Atoll (MVFF).

DNA extraction

Before DNA isolation, sponge tissue was collected and placed in tubes. Samples

were then washed with 1 ml 1X TE buffer. For the sponge samples from JPBS, JPISG,

JPMYK, JPOKE, TWLD, TWLY, and USGU, total DNA was extracted using a DNeasy

Blood and Tissue kit (QIAGEN, Hilden, Germany) according to manufacturer's protocol.

For the other samples, total DNA was extracted by a modified CTAB method described in

163   our previous study [14]. After DNA isolation, the samples were stored at -80°C until

164   subsequent experiments.

16S rRNA gene amplification and multiplex tag sequencing

High throughput sequencing of the 16S rRNA hypervariable V6–V8 region was used

to characterize bacterial community diversity and composition. V6–V8 sequences were

amplified by PCR with forward primer 5´-AACGCGAAGAACCTTAC-3´and reverse

primer 5´-GACGGGCGGTGWGTRCA-3´. PCR mixtures contained 33.5 µL of sterilized

distilled water, 0.5 µL of 5 U TaKaRa Ex Taq (Takara Bio, Otsu, Japan), 5 µL of 10X Ex

Taq buffer, 4 µL of 10 mM dNTP, 1 µL of each primer at a concentration of 10 µM, and 5

µL template DNA in a total volume of 50 µL. PCR was programmed with an initial step

of 94°C for 5 min, 30 cycles of 94°C for 30 s, 52°C for 20 s, and 72°C for 30 s, followed

by a final step of 72°C for 10 min. PCR was performed again to add barcodes to the

amplicons. Each primer tag was designed with four extra nucleotides at the 5' end of both

primers. Unique tags were used for PCR barcoding to label each sample in the study.

PCR conditions were the same as that for the V6–V8 amplification, except that the

number of reaction cycles was reduced to five. The products were purified by 1.5%

agarose electrophoresis and QIAEX II Agarose Gel Extraction kit (QIAGEN, Hilden,

Germany) according to manufacturer's instruction. The quality of the purified products

was assessed using a Nanodrop spectrophotometer (Thermo Scientific, Vantaa, Finland).

Illumina sequencing and community analysis

DNA concentration was measured by a Quant-iT™ assay (Thermo Fisher Scientific,

USA). Equal pools of DNA were sent to Yourgene Bioscience (Taipei, Taiwan) and

sequenced using the MiSeq platform (Illumina, USA). Short and low-quality reads were

filtered by Mothur v.1.38.1 [25] to retain reads with an average quality score > 27 and

lengths of 365–451 base pairs (bp). Reads with homopolymer > 8 bp were excluded, and

chimera sequences from all samples were removed by USEARCH v8.1.1861 [26]. OTUs

were determined and classified using the UPARSE pipeline [27] with reference to the

Silva v128 database [28]. OTUs were defined using a 97% similarity threshold. OTUs

assigned to Archaea, Eukaryota, chloroplast, and mitochondria were removed.

Alpha and beta diversity indices were determined by QIIME 2 v2020.6 [29]. Alpha

diversity indices were calculated by Shannon diversity, Chao1 richness estimator, and

Faith's phylogenetic diversity. Dissimilarities in microbial community compositions

among samples were computed using Bray–Curtis distance dissimilarity, unweighted

UniFrac, and weighted UniFrac. The results were visualized by principal component

analysis (PCA) and dendrogram.

Electron microscopy

T. hoshinota samples at the sponge-coral border were collected by transmission

electron microscopy (TEM), and *T. hoshinota* was observed on rubber by scanning

electron microscopy.

Samples were prepared as described in our previous study [14]. They were fixed in

0.1 M phosphate buffer with 2.5% glutaraldehyde and 4% paraformaldehyde. The

samples were then washed with 0.1 M phosphate buffer for 15 min three times, then

immersed in 0.1 M phosphate buffer with 1% osmium tetroxide for 4 hr.

After being washed again three times, the samples for TEM were sequentially

dehydrated in acetone at 30%, 50%, 70%, 85%, 95%, and 100% for 20 min each time.

After being dehydrated, the samples were embedded in Spurr's resin and sectioned. The

sections were stained with 5% uranyl acetate in methanone and 0.5% lead citrate. The

stained samples were observed under a Philips CM-100 TEM.

Samples for SEM were sequentially dehydrated in 30% for 1 hr, 50% for 1 hr 70%

for 1 hr, 85% for 2 hr, 95% for 2 hr, 100% for 2 hr, and 100% for 12 hr. The samples were

dried in a Hitachi HCP-2, then gold coated with a Hitachi E-1010 and observed under an

FEI Quanta 200 SEM.

Pigment analysis by UPLC-QTOF-MS

Approximately 2.5 g of wet *T. hoshinota* tissue from each sample was centrifuged at

7,000 rpm for 30 s to remove excess water. Sponge tissues were placed in mortars and

ground with 4 ml of 100% acetone to homogenize the tissues. The homogeneous samples

were transferred into a 5 ml centrifuge tube and centrifuged at 7000 rpm at room

temperature for 30 s. The supernatant was transferred into a new 5-ml centrifuge tube.

The samples were covered with aluminum foil paper for being shaded and stored at 4°C

until pigment analysis.

Pigment content analysis was performed by UPLC-MS. The UPLC-MS system used

the Agilent 1290 Infinity II ultra-performance liquid chromatography (UPLC) system

(Agilent Technologies, Palo Alto, CA, USA) coupled online to the Dual AJS electrospray

ionization (ESI) source of an Agilent 6545 quadrupole time-of-flight (Q-TOF) mass

spectrometer (Agilent Technologies, Palo Alto, CA, USA). The samples were separated

using a Kinetex XB-C18 column (2.6 μm, 4.6 × 100 mm, Phenomenex, Torrance, CA,

USA) at 40°C. The chromatogram was acquired; mass spectral peaks were detected and

their waveform processed using Agilent Qualitative Analysis 10.0 software (Agilent,

USA).

Metagenome sequencing and assembly

Bacterial DNA and cDNA were used to reconstruct the cyanobacterial genome. For

DNA sequencing, *T. hoshinota* tissue was ground, and the samples were filtered through a

0.35 μm membrane. 2 x 10^3 cyanobacterial cells were purified from the samples using the

MoFlo XDP cell sorter based on fluorescence and forward scatter intensity and

transferring into two wells of a 96-well plate. The cells were centrifuged at 12,000 x g for

5 min at 4°C, and then lysed with denaturation solution containing 40 mM KOH and 90

mM EDTA in sterilized H_2O at 65°C for 3 min. The DNA was amplified with an REPLI-g

Mini Kit (QIAGEN, USA). The lysates were mixed with 40 μL master mix, which

contained 29 μL REPLI-g mini reaction buffer, 1 μL REPLI-g mini DNA polymerase, and

10 μL sterilized H_2O. After incubation at 30°C for 16 hr with a Thermo-cycler, the

reaction was stopped by heating the samples at 65°C for 3 min. Amplification was

performed twice. The amplified DNA was then purified by Microspin™ G-50 Columns.

Purified DNA was sent to Yourgene Bioscience (Taipei, Taiwan) and sequenced on a

MiSeq platform (Illumina, USA).

For cDNA sequencing. Total RNA was extracted from *T. hoshinota* tissues using

TRI   reagent   (Ambion,   Inc.)   according   to   the   manufacturer's   instructions.

MICROBExpress kits (Ambion, Inc.) were used to remove ribosomal RNA according to

the manufacturer's instructions with capture oligo. Eukaryotic mRNA was then removed

by oligo-dT. After the mRNA was reverse transcribed into cDNA, cDNA sequencing was

performed by NGS High Throughput Genomics Core on HiSseq platform (Illumina,

USA).

DNA and cDNA reads were merged, then trimmed and filtered by trimmomatic

v0.39 with the following parameters: ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10:3:

TRUE LEADING:10 TRAILING:10 SLIDINGWINDOW:5:15 MINLEN:50 [30]. The

processed reads were *de novo* assembled using metaSPAdes with k-mer sizes of 21, 33,

55, 77, and 99 without scaffolding. Binning was performed using VizBin [31], and the

cyanobacterial bin was confirmed by GTDB-TK taxonomy annotation [32].

Genome annotation

The genome of *Prochloron*, "Prochlorobium", and phylogenetically close

cyanobacteria were annotated using Prokka v1.13.7 with the 'usegenus' and 'rfam'

options [33]. The genomes were also annotated with KEGG functional orthologs (KO

numbers) and COGs [34, 35]. To annotate the KO numbers, protein sequences predicted

by Prodigal were blasted against the KEGG database using BlastKoala [36, 37]. The K

number annotation results were then used to reconstruct the transporter systems and

metabolic pathways using KEGG mapper [38]. To annotate COGs, the predicted protein

sequences were searched against the COG database by RPSblast with an e-value

threshold of 10^{-5} [39]. In addition, the transporter proteins were identified by searching

the putative protein sequences against TransportDB 2.0 (August 2019) using BLASTp

[40].

Phylogenetic analysis

To reconstruct a phylogenetic tree of the 16S rRNA genes, the sequences with high

identities to the 16S rRNA gene in LD05 genome were retrieved by searching the NCBI

nt database using BLASTn [39, 41]. The sequences were aligned by MUSCLE on

MEGA7 [42, 43]. A tree was then reconstructed using IQ-TREE v2.03 using automatic

model selection and 1,000 bootstraps [44]. The tree was visualized with iTOL v4 [45].

Genomes for each cyanobacteria species in GTDB database were downloaded and

120 single-copy gene protein sequences were used to reconstruct a tree [46]. Marker gene

protein sequences were identified, aligned, and concatenated by GTDB-TK v1.3.0 [32]. A

phylogenomic tree was then built using IQ-TREE v2.03 with automatic model selection

and 1,000 Ultrafast bootstraps. The tree was visualized with iTOL v4 [44, 45].

Availability of data and materials

All the raw data and genome were submitted under BioProject ID: PRJNA665642.

Results

Bacterial diversity in *Terpios hoshinota* samples from different oceans

To characterize the bacterial communities in coral-encrusting *T. hoshinota* across

different geographical backgrounds, samples were collected from 15 locations in the

Pacific Ocean, South China Sea, and Indian Ocean (Table S1). 16S ribosomal RNA

(rRNA) gene sequencing was used to determine the bacterial communities.

Statistical tests showed that operational taxonomic unit (OTU) Chao1 richness and

Faith's phylogenetic diversity (PD) indices were not significantly associated with the

oceanic regions from which the samples were collected (Chao1: $p = 0.23$, PD: $p = 0.30$)

(Fig. S1a and S1b). In terms of Shannon diversity, the indices were also not significantly

associated with the oceanic regions when the outliners were removed (Fig. S1c). Some

samples—e.g., TWKT, MYRD, and MVFF samples—from the same locations showed

high variation in Chao1 and Shannon indices. In contrast to the alpha diversity indices,

the similarities in microbial composition were associated with sample origins (Fig. S2).

PERMANOVA showed that samples from the same ocean were significantly more similar

than those from different oceans (Bray Curtis: pseudo-$F = 5.13514$, $p = 0.001$,

Unweighted UniFrac: pseudo-F = 1.60401, p = 0.01). When relative abundances were

taken into consideration, the effect of the sea on similarities increased (weighted UniFrac:

pseudo-F = 6.09022, p = 0.001).

Prochloron-like bacterium dominated _T. hoshinota_ bacterial communities across

different oceans

Microbial composition analysis revealed that _Cyanobacteria_ and _Proteobacteria_

accounted for most of the bacterial composition across the samples, ranging from

81.5–97.4% (Fig. 1, Fig. S3a, and Table S2). _Bacteroidetes_ and _Spirochaetae_ were

present across all the samples, though their abundances were only 0.34–6.29%.

Genus-level analysis showed that _Prochloron_-like OTUs were the most dominant and

present in all the samples (Fig. S3b). Moreover, _Prochloron_-like OTUs accounted for

most of the _Cyanobacteria_ in the samples (Table S3). In total, four OTUs were assigned

to _Prochloron_, but only one of them, referred to as Prochl-OTU01, was predominant

across the samples.

Along   with   Prochl-OTU01,   _Endozoicomonas_,   _Pseudospirillum_,   SAR116,

Magnetospira, and _Ruegeria_ OTUs were common and co-occurred across the samples,

forming a putative core microbiome. This core microbiome contributed about 80% of

bacterial abundance in each sample.

Structure of symbiotic *Prochloron*-like bacteria in *T. hoshinota*

To reveal the detailed structure of the *Prochloron*-like bacteria, a *T.*

hoshinota-encrusting abiotic substrate was observed using transmission electron

microscopy and scanning electron microscopy (Fig. 2). The *T. hoshinota* sponge, roughly

500 μm thick, were supported by the bundle of tylostyle spicules, and the spicules on the

surface were interlaced (Fig. 2a-b). Inside the body, spherical bacterial cells 4–6 μm in

diameter were widely distributed, and cells were observed in various stages of cell

division (Fig. 2c and 2f). A study published in 2015 isolated a baeocytous

cyanobacterium, *Myxosarcina* sp., from *T. hoshinota* [21]. However, no vegetative cell

aggregate of baeocytes could be observed in our results.

TEM results showed that these spherical cells contained thylakoids, typical

compartments inside cyanobacteria, and the arrangements were parietal (Fig. 2d–f). In

addition, phosphate granules and carboxysomes were found; gas vesicles were absent in

these cells, which is consistent with a previous description of *Prochloron*'s structure [47].

Genome assembly of a novel *Prochloron*-like species

The predominance of a *Prochloron*-like OTU in *T. hoshinota* from various locations

indicates the intimate association between bacterium and host. To reveal the identity and

characteristics of the *Prochloron*-like bacterium, a *Prochloron*-like genome, referred as to

LD05, was reconstructed by *de-novo* assembly of the metagenome from the sample at

Green Island, Taiwan. The metagenome-assembled genome (MAG) was 91.73%

complete, with 2.16% contamination (Table 1). A *Prochloron*-like 16S rRNA gene was

also successfully recovered from the genome, and the sequence was identical to that of

the predominant Prochl-OTU01 identified from the community survey. The analysis of

the GTDB-Tk demonstrated that the LD05 MAG was closest to *Prochloron didemni,* with

77.97% average nucleotide identity (ANI).

Recently, a putative *Prochloron* species genome called SP5CPC1 was recovered

from the metagenome of a sponge microbiome [48]. The ANI analysis found that the

LD05 genome shared 93.18% identity with the SP5CPC1 genome, which is below the

95% ANI cutoff for species delineation [49]. The LD05 shared a similar genome size, GC

ratio, and coding density with SP5CPC1 (Table S4). On the other hand, *P. didemni* P2-P4

had a much larger genome and lower GC ratio and coding densities. There was also a

discrepancy between LD05 and *P. didemni* P2-P4 based on the average amino acid

identity (AAI) and percentage of conserved proteins (POCP) analyses (Fig. S4). The

LD05 shared 91% AAI with SP5CPC1, but only 70% AAI with *P. didemni* P2–P4

genomes. On the other hand, the POCP between LD05 and *Prochloron* genomes ranged

from 49.8–52%, close to a proposed 50% POCP cutoff for the genera delineation [50].

Taken together, the LD05 and SP5CPC1 genomes were different species with high

similarity, and both were more distantly related to *P. didemni*.

Phylogenetic and functional genomic analyses revealed distinct characteristics of

LD05 and SP5CPC1 genomes compared to other *Prochloron* genomes

Although the LD05 and SP5CPC1 genomes had certain degrees of similarity to

Prochloron genomes, the phylogenetic, genomic, and functional genomic analyses

identified distinct characteristics between them and *Prochloron*, suggesting that they

belong to a different genus.

A phylogenetic tree of 120 cyanobacteria was constructed based on 120 single-copy

genes to determine phylogenic affiliation (Fig. 3a and Fig. S5). The tree demonstrated

that the LD05 and SP5CPC1 formed a clade that was sister to the clade containing *P.*

didemni. The larger clade encompassing the two clades was adjacent to the clade

containing *Synechocystis*, *Myxosarcina*, and other cyanobacteria. To provide an in-depth

view, homologs of the LD05 16S rRNA gene with high similarities were retrieved from

the NCBI database. A tree constructed by these 16S rRNA sequences depicted two

distinct clades, referred as to Clade I and Clade II (Fig. 3b). Clade II contained 16S rRNA

gene sequences from three genomes of *P. didemni* strains and other 16S rRNA sequences

from various ascidians. In contrast, Clade I mainly comprised sequences from sponge

holobionts, including SP5CPC1, LD05, and 16S rRNA sequences from a variety of other

sponges. Although many sequences in Clade I were assigned as *Synechocystis* and

Prochloron 16S rRNA sequences, we found that they actually shared higher identities

with the members of Clade I than *Synechocystis* or *Prochloron*. Moreover, the genome

phylogenomic analysis also revealed that the SP5CPC1 and LD05 are phylogenetically

distant from cultivated *Synechocystis* (Fig. 3a).

To characterize the functional differences between the bacteria in the two clades,

LD05, SP5CPC1, and *P. didemni* genomes were annotated with KEGG orthologs and

clusters of orthologous groups (COGs). Complete-linkage clustering of COG category

abundances demonstrated that the LD05 and SP5CPC1 were more functionally similar to

Microcystis groups than to *P. didemni* (Fig. S6).

Two hallmarks of *Prochloron* are the presence of chlorophyll *b* and absence of

phycobilins in the cell [51]. However, a previous study showed that the symbiotic

cyanobacteria in *T. hoshinota* had phycobilins based on microspectrophotometry results

[15], which is consistent with our finding that phycobilins synthase and the

phycobiliproteins genes were present in LD05 and SP5CPC1 genomes. The absence of

the chlorophyll *b* synthase gene in LD05 and SP5CPC1 genomes implied that the bacteria

lacked chlorophyll *b*. In addition, the failure to identify chlorophyll *b* in the ascidian

Trididemnum nubilum holobiont, which the PCR clone AO15 (DQ357958.1) in Clade I

was recovered from, implied that the bacterium also lacks chlorophyll *b*. In accordance

with the genomic analyses, our pigment analysis by LC-QTOF-MS also revealed the

presence of chlorophyll *a* and absence of chlorophyll *b* in *T. hoshinota* (Fig. 4). These

results indicated that the bacteria in Clade I had different light-harvesting systems from

Prochloron in Clade II.

In summary, the results indicate that the LD05 and SP5CPC1 were derived from a

common ancestor, phylogenetically different from the ancestor of *Prochloron*. More

importantly, the bacteria have photosynthetic machinery distinct from that of *Prochloron*.

Therefore, we propose that the bacteria in Clades I and II be classified into two different

genera and classify the species in Clade I as a novel genus "Prochlorobium". Moreover,

we classified LD05 as a novel species named *Candidatus* Prochlorobium terpiosii LD05

[Etymology: Gr. prep. and pref. *pro*, before, primitive; Gr. adj. *chlôros*, green; Gr. n. *bios*,

life; N.L. neut. n. *Prochlorobium*, primitive green life. N.L. gen. n. *terpiosii*, of *Terpios*, a

zoological scientific genus name], and SP5CPC1 was reclassified as *Candidatus*

Prochlorobium sp. SP5CPC1 (Fig. S7).

Metabolic features of the novel species *Ca.* Pb. terpiosii LD05

Ca. Pb. terpiosii LD05 possessed nearly all the genes required for photosynthesis,

carbon fixation, the tricarboxylic acid cycle (TCA), and glycolysis (Fig. 5). In addition,

genes related to sucrose metabolism—e.g., sucrose-6-phosphatase, sucrose synthase, and

sucrose phosphorylase—were identified. These genes were also present in the SP5CPC1

genome, but not in *P*. *didemni* genomes. For light-harvesting systems, like typical

cyanobacteria, the genes encoding phycobiliproteins—including phycoerythrin,

phycocyanin, and allophycocyanin—were recovered, except for *cpcC*, *cpcD*, and *cpeE*.

Phycobilin synthase genes were also detected in the genomes. On the other hand, the

genes encoding chlorophyll *a* synthesis were present, but the gene for chlorophyll *b*

synthase was absent.

Certain symbiotic cyanobacteria contain nitrogen fixation machinery and provide

their host with nitrogen sources. However, the *Ca*. Pb. terpiosii LD05 and SP5CPC1

lacked nitrogen fixation genes. The two genomes had genes for nitrate uptake and

assimilatory nitrate reduction pathways to convert nitrate into ammonium. The glutamine

synthetase/glutamate synthase (GS/GOGAT) pathway was also found in the genomes,

indicating that the ammonium derived from extracellular nitrate can be incorporated into

amino acids by this pathway. Additionally, the genomes also carried the complete gene

set for the urea transporter and urease. The bacteria may take in the urea from hosts and

convert it into ammonium. More importantly, the general L-amino acid transporters can

be identified in the genomes, which cannot be observed in *Prochloron* genomes. Hence,

the bacteria may have multiple methods for obtaining nitrogen resources from their host

and environment.

Many symbiotic bacteria in sponges have been shown to provide their hosts with

cofactors and vitamins such as vitamin B_{12} (cobalamin), which can only be synthesized

by prokaryotic organisms [22]. The *Ca*. Pb. terpiosii LD05 genome contained nearly

complete gene sets for the synthesis pathways of vitamin B_1 (thiamine), vitamin B_2

(riboflavin), vitamin B_7 (biotin), and vitamin B_{12}. The transporter for cobalt, a key

constituent of vitamin B_{12}, was also be identified. In contrast, the *Prochloron* genomes

lacked a cobalt transporter and many genes for vitamin B_{12} synthesis.

Previous studies have discovered a variety of secondary metabolites from the *T.*

hoshinota holobiont [52]; these metabolites may be produced by symbiotic bacteria [53].

Our analysis revealed the existence of multiple terpene synthesis clusters in *Ca*. Pb.

terpiosii LD05. Furthermore, the species also contains a gene encoding squalene cyclase,

suggesting that the bacterium can produce cyclic triterpenes.

Many symbiotic bacteria in sponge carry eukaryotic-like proteins (ELPs), which

may originate from eukaryotes [54]. The ELPs—e.g., leucine-rich repeat proteins (LRRs),

ankyrin repeat proteins (ARPs), WD-40 containing proteins (WD-40), and

tetratricopeptide repeat proteins (TPRs)—may be used to interact with hosts during

various biological processes. The genomic analysis demonstrated that *Ca*. Pb. terpiosii

LD05 carried these ELPs. However, some closely related free-living cyanobacteria also

possess ELPs. On the other hand, the ELPs were highly enriched across *Prochloron*

genomes.

Discussion

T. hoshinota is one of the most important biological threats to coral. Its associated

cyanobacteria and other symbiotic bacteria are essential for maintaining the sponge's

function. Nevertheless, the present study is the first to explore the composition and role of

T. hoshinota's microbiome. Our unprecedented large-scale survey of the bacterial

community in *T. hoshinota* based on 16S amplicon sequencing of samples from various

regions across the western Pacific Ocean, Indian Ocean, and South China Sea enabled us

to characterize the sponge's core microbiome.

We discovered a bacterium, closely related to *Prochloron*, that was dominant across

our sponge samples. The prevalence and predominance of this bacterium in *T. hoshinota*

highlight the strong relationship between bacteria and their hosts. Recovery of the

bacterial genome, followed by phylogenetic and functional analyses, demonstrated that

this cyanobacterium has a distinct phylogenetic affiliation and genomic and metabolic

characteristics from *Prochloron,* indicating that it belongs to a novel genus. We proposed

a new genus, "Prochlorobium", and refined the membership of *Prochloron*. Lastly, a

functional analysis was performed to understand the lifestyle of this novel bacterium in

the *T. hoshinota* holobiont. This study provides insights into the role of the

cyanobacterium in *T. hoshinota*, which will facilitate future studies to investigate the

ecosystem inside *T. hoshinota*, the causes of its outbreaks, and its invasion mechanism.

Biogeographical variation in the *T. hoshinota* microbiome and its core microbial

members

The microbial community in sponges can be shaped by several environmental

factors—e.g., light, nutrients, temperature, and pH—and may change in response to stress

[23, 24, 55, 56]. Environmental changes may force a sponge holobiont to alter its

microbial community to acclimatize [57]. On the other hand, host-related factors can also

be important for determining community composition. For example, previous studies

have shown that sponges have several immune receptors, such as NOD-like and Toll-like

receptors [13]. These receptors enable sponges to discriminate between symbionts and

prey. In order to live inside a sponge, microbes have to evade host immunity and

phagocytosis [58, 59]. Furthermore, metabolites and nutrients provided by hosts also

contribute to the community structure [60]. Some sponges show high stability in their

microbiome due to strict selective pressures from their host. For instance, *Ircinia* and

Hexadella sponges exhibit host-specific and stability in their associated bacterial

communities, despite large geographic distances between sampling sites [24, 61]. In

contrast, certain sponges, such as *Petrosia ficiformis,* harbor biogeographical-dependent

bacterial compositions [62].

Our survey of three oceans enabled us to determine whether bacterial communities

vary across different biogeographical backgrounds and experience strong selective

pressure from their host. Our dissimilarity analysis of the microbial composition of *T.*

hoshinota showed a weak correlation between sample location and bacterial composition.

The differences in microbial composition among *T. hoshinota* samples from different

locations may result from local acclimatization. Samples from the same oceanic regions

that share communities may equip their holobionts with similar metabolic functions to

cope with stress or increase fitness in certain environments.

However, despite the biogeographical variation, we still identified core bacterial

members that were present in all samples (Fig. S3). This core bacterial community

comprised only a few genera, but accounted for ~80% of the community across all

samples, indicating that it is vital to the holobiont. Host-related factors may help keep this

core microbiome stable because its members carry out core functions of the holobiont

[23]. The core members may contain the metabolic capability to utilize nutrients from the

sponge host's environment and play important roles in nutrient exchanges, such as sulfur,

carbon, and nitrogen cycling. This core group may be also responsible for defending

against predators and protecting the sponge symbionts from toxins and pathogens [23].

The most dominant OTU in the core microbiome was closely related to *Prochloron*,

a   genus   of   symbiotic   cyanobacteria   found   in   various   ascidians.   The

biogeographically-independent prevalence and predominance of this *Prochloron*-like

bacterium in *T. hoshinota* suggest that it has a crucial role in the holobiont.

Candidatus **Prochlorobium terpiosii gen. nov., sp. nov., a *Prochloron*-like bacterium**

prevalent in *T. hoshinota*

Although the genome of the *Prochloron*-like bacterium in *T. hoshinota* is similar to

that of *Prochloron*, deeper genomic, phylogenetic, and functional analyses, combined

with the recent recovery of another symbiont *Prochloron*-like bacterial genome in the

sponge, revealed that the two cyanobacteria are distinct, especially in terms of their

pigment contents and phylogenetic divergence. Hence, a new genus, "Prochlorobium",

was formally erected to distinguish these two groups.

Our 16S rRNA metagenomic analysis revealed that *Candidatus* Prochlorobium

terpiosii LD05 is the most dominant cyanobacterium within *T. hoshinota* specimens

collected across three different oceans. In order to determine whether *Ca*. Pb. terpiosii

LD05 is adapted to different locations, a small-scale microbial composition survey was

conducted in Taiwan and Japan using 16S rRNA PCR cloning. The results suggested that

the variations among the symbiotic *Ca*. Pb. terpiosii LD05 are small, and the sequences

from the same location could not be grouped in the resulting phylogenetic tree (Fig. S8),

indicating that there is no significant local adaptation.

Why does the dominant cyanobacterium species remain identical in *T. hoshinota*

samples across different oceans? The reason remains unclear. Recent studies suggest that

T. hoshinota larvae, which carry vertically transmitted cyanobacteria, may have a short

dispersal distances because they are more dense than the water, leading them to settle

rapidly after leaving their mother sponge [16-18]. In this circumstance, the symbiotic *Ca.*

Pb. terpiosii LD05 from various locations might accumulate genetic differences to adapt

to local environments. The absence of evident genetic alterations in our study may be the

result of the tight and stable symbiotic interactions between *Ca*. Pb. terpiosii LD05 and *T.*

hoshinota restricting genetic changes in the bacterium. Another possibility is that the

species becomes broadly dispersed through some other mechanism, such as via ocean

currents or transportation via vessels; this would enable *T. hoshinota* larvae to spread

quickly across the ocean with few genetic alterations. Evidence of recent *T. hoshinota*

outbreaks support this latter scenario [2].

Comparison between "Prochlorobium" and *Prochloron*

Comparative genomics between "Prochlorobium" and *Prochloron* enables us to infer

their evolutionary histories and respective relationship to their host. Previous studies have

shown that symbiotic bacteria usually have reduced genomes because certain genes erode

as symbiosis develops [63, 64]. Furthermore, a model of symbiont evolution has

proposed that during the evolutionary history of symbiosis large-scale pseudogenization

can happen during transitional events, such as a strict host association or vertical

transmission, leading to a sudden drop in coding density [64]. Eventually, the coding

density gradually bounces back due to deletion bias.

Comparing *Prochloron* and "Prochlorobium", we found that "Prochlorobium" had a

smaller genome but similar coding density to other phylogenetically closely related

cyanobacteria. *Prochloron*, on the other hand, had a similar genome size but lower coding

density. This difference may indicate that the transition toward the host-restricted

lifestyles occurred more recently in *Prochloron* than "Prochlorobium". The hypothesis

can be supported by our observation that *Prochloron* harbored highly-enriched

transposase genes as the elevation of mobile genetic element quantities, such as

transposons and insertion sequences, is thought to be associated with the recent transition

to a host-restricted symbiotic lifestyle [65].

Along with its unusual pigments, another hallmark of *Prochloron* is that is produces

patellamides, cytotoxic cyclic peptides. However, the synthesis gene cluster cannot be

found in *Ca*. Pb. terpiosii LD05 or SP5CPC1. On the contrary, the latter two genomes

have many terpene synthesis gene clusters and contain genes that encode the protein that

circulates terpene. Hence, "Prochlorobium" and *Prochloron* may produce distinct

secondary metabolites to increase the fitness of themselves or their entire holobiont.

Certain cyanobacteria contain genes for sucrose metabolism. Though the role of

sucrose in cyanobacteria remains understudied, several articles have shown that sucrose

can be utilized as a compatible solute, serve as signal molecule, or be used for glycogen

synthesis [66-69]. Interestingly, our genomic analysis found that the genes involved in

sucrose metabolism were present in *Ca*. Pb. terpiosii LD05 and SP5CPC1 but absent in

Prochloron. We hypothesize that "Prochlorobium" can use sucrose as an osmolyte to

cope with osmotic stress; this is supported by a previous finding that the genes related to

osmotic stress are enriched in sponge-associated bacterial genomes [22]. On the other

hand, we also found that *Ca*. Pb. terpiosii LD05 and SP5CPC1 contained osmoprotectant

transporter genes, which were not identified in *Prochloron* genomes. These results

indicate that the "Prochlorobium" species may live in environments with higher osmotic

stress, like places with high osmolarity or fluctuations in osmolarity, compared to

Prochloron. Alternatively, "Prochlorobium" and *Prochloron* may use different strategies

to deal with osmotic stress.

Another observation that drew our attention is that eukaryotic-like proteins (ELPs)

are highly-enriched in *Prochloron* but not "Prochlorobium". ELPs can be used to perform

protein-protein interactions, phagocytosis evasion, host cell binding, and interference

with the host ubiquitin system [70]. ELPs are also thought to be necessary for sponge

symbionts [8]. The differences in ELP quantities between *Prochloron* and

"Prochlorobium" may indicate that "Prochlorobium" uses an unidentified mechanism to

interact with their hosts. Another possibility is that the long-term symbiotic history of

"Prochlorobium" propelled bacteria to retain their essential ELPs and reduce unnecessary

ones.

Symbiotic interactions between *Ca*. Pb. terpiosii and *T. hoshinota*

Sponges harbor complex bacterial communities [13], and the symbiotic bacteria play

vital and diverse roles in the physiology and ecology of sponges. For instance, bacteria

can provide their hosts with various nutrients and recycle waste. Moreover, symbiotic

microorganisms can produce diverse secondary metabolites to compete with other

holobionts or defend from invasion by predators. The predominance of *Ca*. Pb. terpiosii

LD05 highlights its role in the *T. hoshinota* and the genome of *Ca*. Pb. terpiosii LD05

enables us to understand its metabolic potential, the symbiotic relationship, and even how

the holobiont overgrows corals.

Many sponges are mixotrophs and certain sponges can obtain carbon resources from

cyanobacteria [71]. Some sponges even acquire more than 50% of their energy demand

from the symbiotic cyanobacteria in the form of photosynthates [72]. Photosynthesis was

observed in the *T. hoshinota* holobiont, and its efficiency increased when the coral-killing

holobiont came into contact with coral; this mechanism may help sponges overgrow the

coral [73]. *Ca*. Pb. terpiosii LD05 is the only cyanobacterium in the core microbiome, and

it has been found to have the genes needed for photosynthesis. Hence, the bacterium may

be an important carbon source for *T. hoshinota* and facilitate competition with coral.

Ammonia secreted by sponges may serve as a nitrogen source for symbiotic bacteria.

Our analysis identified ammonium transporters and the GS-GOGAT pathway in *Ca*. Pb.

terpiosii LD05. Hence, the bacterium may recycle the ammonium from host. On the other

hand, the bacterium harbors the urea transporter and urease. Therefore, urea can be used

as an alternative nitrogen source. A previous study showed that the level of free amino

acids in *T. hoshinota*-living cyanobacteria increased when the holobiont encountered

coral. Our finding of an amino acid transporter in *Ca*. Pb. Terpiosii LD05 suggests that

the bacterium may benefit from "coral killing" by leaching amino acids or ammonium

that are released as the coral disintegrates.

Some microbes that live in sponges can store polyphosphate granules, which can

account for 25–40% of the total phosphate content in sponge tissue [74, 75]. These

polyphosphate granules can be used for energy storage and may enable the holobiont to

withstand periods of phosphate starvation. Our electron microscopy revealed

polyphosphate granules in *Ca*. Pb. terpiosii LD05. We also found genes encoding

polyphosphate kinase. Hence, *Ca*. Pb. terpiosii LD05 may play a pivotal role in

phosphate storage and cycling in the *T. hoshinota* holobiont. A previous study

demonstrated that the phosphate concentration in seawater was highly correlated with *T.*

hoshinota overgrowth [76], highlighting the importance of phosphate in the spread of the

T. hoshinota holobiont. Another study also proposed that *T. hoshinota* accumulates

nutrients to compete with corals [8]. We propose that *Ca*. Pb. terpiosii LD05 helps *T.*

hoshinota outcompete corals by serving as a carbon and phosphate storage. However,

how the phosphate concentration is correlated with *T. hoshinota* cover is unclear, as are

details about phosphate cycling among the host, *Ca*. Pb. terpiosii LD05, and other

symbiotic microbes.

Animals cannot synthesize essential vitamins, so symbiotic microorganisms are

thought to be important sources of essential vitamins for sponges. Our analysis of the *Ca*.

Pb. terpiosii LD05 genome identified the biosynthesis pathway for vitamin B_1, vitamin B_7,

and vitamin B_{12}. Thus, *Ca*. Pb. terpiosii LD05 may help maintain *T. hoshinota*'s health by

providing the holobiont with vitamins.

One of the leading questions in the study of *T. hoshinota* is how it kills corals.

Several mechanisms have been proposed. One argues that the sponge produces cytotoxic

allelochemicals to damage coral cells [77]. Another suggests that the sponge overgrows

corals and competes with them for nutrients [8]. Finally, the sponge may cause the coral's

microbial composition to change, causing it to malfunction [78]. These hypotheses are

not mutually exclusive. For the first, several cytotoxic compounds, including

nakiterpiosin, nakiterpiosinon, and terpiodiene, have been isolated from *T. hoshinota*

holobionts [79, 80]. Nakiterpiosin, and nakiterpiosinon are C-nor-D-homosteroids.

Previous studies showed that cyanobacteria can produce sterols by the cyclization of

squalene, a triterpene [81]. In the *Ca*. Pb. terpiosii LD05 genome, the synthesis pathway

of squalene and squalene cyclases were identified. Hence, the bacterium may be

responsible for the production of the toxins. These toxins may facilitate overgrowing

corals by damaging coral tissue directly or weakening the coral's defenses.

Conclusions

This study makes several fundamental discoveries about the bacterial community

associated with *T. hoshinota*. The core microbiome of *T. hoshinota* includes a

Prochloron-like bacterium, as well as *Endozoicomonas*, *Pseudospirillum*, SAR116,

Magnetospira, and *Ruegeria*. Our finding that the dominant cyanobacteria in *T. hoshinota*

is the same throughout the western Pacific Ocean, South China Sea, and Indian Ocean

suggests that this particular cyanobacterium is the obligate symbiont of the sponge. The

high-quality draft genome of this cyanobacterium was recovered, and the phylogenetic

and genomic analyses revealed that the genome should be classified as a novel species

under a new genus "Prochlorobium", close to *Prochloron*. The draft genome enables us to

understand the carbon and nitrogen metabolism of the bacterium, and the putative

symbiotic interaction between the bacterium and *T. hoshinota*. By identifying the core

microbiome and revealing the metabolic potential of the dominant cyanobacterium in *T.*

hoshinota, our conclusions will help future studies explore the detailed ecosystem inside

the holobiont, unveil how *Ca*. Pb. terpiosii LD05 contributes to the invasion and

coral-killing nature of *T. hoshinota*, and determine the cause of the outbreaks in the

future.

Acknowledgements

Y.H.C acknowledges the Taiwan International Graduate Program (TIGP) for its

fellowship towards his graduate studies. We thank Noah Last of Draft Editing for his

English language editing, Dr. Ker-yea Soong and Shih-Shou Fang for collecting *T.*

hoshinota samples, and the Green Island Marine Research Station for supporting the

research. We thank the electron microscope and flow cytometry divisions of IPMB,

Academia Sinica for their technical support and Chih-Yu Lin and Gong-Min Lin in the

Metabolomics Core Facility, Agricultural Biotechnology Research Center, Academia

Sinica for the technical support and for performing the UPLC-MS/MS analysis and

processing data.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by Biodiversity Research Center, Academia Sinica and the

National Science Council, Taiwan (NSC98-2321-B-001-025-MY3678106).

Authors' contributions

YHC, HJC, and SLT designed the study and prepared the manuscript. YHC and HJC

analyzed and interpreted the data. HJC, CYY, JHS, DZH, PWC, MHL, and WSC

collected and sequenced the samples. HJC, CYY, and JHS observed the electron

microscopy. HJC, THS, SHL, and CMY examined the pigments. CAC, JDR, EH, BHI,

HH, PJS, CHJT, and HY assisted with the sampling.

All authors read and approved the manuscript.

References

1.   Rützler K, Muzik K: **Terpios hoshinota, a new cyanobacteriosponge threatening**
Pacific reefs. *Recent advances in systematics and ecology of sponges Scientia*
Marina 1993.
2.   Liao M-H, Tang S-L, Hsu C-M, Wen K-C, Wu H, Chen W-M, Wang J-T, Meng P-J,
Twan W-H, Lu C-K: **The" Black Disease" of Reef-Building Corals at Green Island,**
Taiwan-Outbreak of a Cyanobacteriosponge. Terpios hoshinota (Suberitidae;
Hadromerida). *ZOOLOGICAL STUDIES-TAIPEI-* 2007, **46**(4):520.
3.   Fujii T, Keshavmurthy S, Zhou W, Hirose E, Chen C, Reimer J: **Coral-killing**
cyanobacteriosponge (Terpios hoshinota) on the Great Barrier Reef. *Coral Reefs*
2011, **30**(2):483-483.
4.   de Voogd NJ, Cleary DFR, Dekker F: **The coral-killing sponge Terpios hoshinota**
invades Indonesia. *Coral Reefs* 2013, **32**(3):755-755.
5.   Hoeksema BW, Waheed Z, de Voogd NJ: **Partial mortality in corals overgrown by**
the sponge Terpios hoshinota at Tioman Island, Peninsular Malaysia (South
China Sea). *B Mar Sci* 2014, **90**(4):989-990.
6.   Thinesh T, Jose PA, Hassan S, Selvan KM, Selvin J: **Intrusion of coral-killing sponge**
(Terpios hoshinota) on the reef of Palk Bay. *Curr Sci India* 2015,
109(6):1030-1032.
7.   Montano S, Chou WH, Chen CA, Galli P, Reimer JD: **First record of the coral-killing**
sponge Terpios hoshinota in the Maldives and Indian Ocean. *B Mar Sci* 2015,
91(1):97-98.
8.   Wang JT, Chen YY, Meng PJ, Sune YH, Hsu CM, Wei KY, Chen CA: **Diverse**
Interactions between Corals and the Coral-Killing Sponge, Terpios hoshinota
(Suberitidae: Hadromerida). *Zoological Studies* 2012, **51**(2):150-159.
9.   Plucerrosario G: **The Effect of Substratum on the Growth of Terpios, an**
Encrusting Sponge Which Kills Corals. *Coral Reefs* 1987, **5**(4):197-200.

10.   Yomogida M, Mizuyama M, Kubomura T, Reimer JD: **Disappearance and Return**
734   **of an Outbreak of the Coral-killing Cyanobacteriosponge Terpios hoshinota in**
735   **Southern Japan.** *Zoological Studies* 2017, **56**.

736   11.   Shi Q, Liu G, Yan HQ, Zhang HL: **Black Disease (Terpios hoshinota): A Probable**
737   **Cause for the Rapid Coral Mortality at the Northern Reef of Yongxing Island in**
738   **the South China Sea.** *Ambio* 2012, **41**(5):446-455.

739   12.   Madduppa H, Schupp PJ, Faisal MR, Sastria MY, Thoms C: **Persistent outbreaks of**
740   **the "black disease" sponge Terpios hoshinota in Indonesian coral reefs.** *Mar*
741   *Biodivers* 2017, **47**(1):149-151.

742   13.   Hentschel U, Piel J, Degnan SM, Taylor MW: **Genomic insights into the marine**
743   **sponge microbiome.** *Nat Rev Microbiol* 2012, **10**(9):641-U675.

744   14.   Tang SL, Hong MJ, Liao MH, Jane WN, Chiang PW, Chen CB, Chen CA: **Bacteria**
745   **associated with an encrusting sponge (Terpios hoshinota) and the corals**
746   **partially covered by the sponge.** *Environmental Microbiology* 2011,
747   **13**(5):1179-1191.

748   15.   Hirose E, Murakami A: **Microscopic anatomy and pigment characterization of**
749   **coral-encrusting black sponge with cyanobacterial symbiont, Terpios hoshinota.**
750   *Zoolog Sci* 2011, **28**(3):199-205.

751   16.   Wang JT, Hirose E, Hsu CM, Chen YY, Meng PJ, Chen CA: **A Coral-Killing Sponge,**
752   **Terpios hoshinota, Releases Larvae Harboring Cyanobacterial Symbionts: An**
753   **Implication of Dispersal.** *Zoological Studies* 2012, **51**(3):314-320.

754   17.   Hsu CM, Wang JT, Chen CA: **Larval release and rapid settlement of the**
755   **coral-killing sponge, Terpios hoshinota, at Green Island, Taiwan.** *Mar Biodivers*
756   2013, **43**(4):259-260.

757   18.   Nozawa Y, Huang YS, Hirose E: **Seasonality and lunar periodicity in the sexual**
758   **reproduction of the coral-killing sponge, Terpios hoshinota.** *Coral Reefs* 2016,
759   **35**(3):1071-1081.

760   19.   Soong K, Yang SL, Chen CA: **A Novel Dispersal Mechanism of a Coral-Threatening**
761   **Sponge, Terpios hoshinota (Suberitidae, Porifera).** *Zoological Studies* 2009,
762   **48**(5):596-596.

763   20.   Thinesh T, Meenatchi R, Pasiyappazham R, Jose PA, Selvan M, Kiran GS, Selvin J:
764   **Short-term in situ shading effectively mitigates linear progression of**
765   **coral-killing sponge Terpios hoshinota.** *Plos One* 2017, **12**(8).

766   21.   Yu CH, Lu CK, Su HM, Chiang TY, Hwang CC, Liu T, Chen YM: **Draft genome of**
767   **Myxosarcina sp. strain GI1, a baeocytous cyanobacterium associated with the**
768   **marine sponge Terpios hoshinota.** *Stand Genomic Sci* 2015, **10**:28.

22.   Webster NS, Thomas T: **The Sponge Hologenome**. *mBio* 2016,
7(2):e00135-00116.

23.   Pita L, Rix L, Slaby BM, Franke A, Hentschel U: **The sponge holobiont in a**
changing ocean: from microbes to ecosystems. *Microbiome* 2018, **6**(1):46.

24.   Pita L, Turon X, Lopez-Legentil S, Erwin PM: **Host rules: spatial stability of**
bacterial communities associated with marine sponges (Ircinia spp.) in the
Western Mediterranean Sea. *FEMS Microbiol Ecol* 2013, **86**(2):268-276.

25.   Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski
RA, Oakley BB, Parks DH, Robinson CJ *et al*: **Introducing mothur: Open-Source,**
Platform-Independent, Community-Supported Software for Describing and
Comparing Microbial Communities. *Appl Environ Microb* 2009,
75(23):7537-7541.

26.   Edgar RC: **Search and clustering orders of magnitude faster than BLAST.**
Bioinformatics 2010, **26**(19):2460-2461.

27.   Edgar RC: **UPARSE: highly accurate OTU sequences from microbial amplicon**
reads. *Nat Methods* 2013, **10**(10):996-+.

28.   Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J,
Ludwig W, Glockner FO: **The SILVA and "All-species Living Tree Project (LTP)"**
taxonomic frameworks. *Nucleic Acids Res* 2014, **42**(D1):D643-D648.

29.   Bolyen E, Rideout JR, Dillon MR, Bokulich N, Abnet CC, Al-Ghalith GA, Alexander
H, Alm EJ, Arumugam M, Asnicar F *et al*: **Reproducible, interactive, scalable and**
extensible microbiome data science using QIIME 2. *Nat Biotechnol* 2019,
37(8):852-857.

30.   Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina**
sequence data. *Bioinformatics* 2014, **30**(15):2114-2120.

31.   Laczny CC, Sternal T, Plugaru V, Gawron P, Atashpendar A, Margossian HH,
Coronado S, van der Maaten L, Vlassis N, Wilmes P: **VizBin - an application for**
reference-independent visualization and human-augmented binning of
metagenomic data. *Microbiome* 2015, **3**.

32.   Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH: **GTDB-Tk: a toolkit to classify**
genomes with the Genome Taxonomy Database. *Bioinformatics* 2020,
36(6):1925-1927.

33.   Seemann T: **Prokka: rapid prokaryotic genome annotation.** *Bioinformatics* 2014,
30(14):2068-2069.

34.   Tatusov RL, Galperin MY, Natale DA, Koonin EV: **The COG database: a tool for**
genome-scale analysis of protein functions and evolution. *Nucleic Acids Res*

2000, **28**(1):33-36.

35.   Kanehisa M, Goto S: **KEGG: Kyoto Encyclopedia of Genes and Genomes.** *Nucleic*
Acids Res 2000, **28**(1):27-30.

36.   Kanehisa M, Sato Y, Morishima K: **BlastKOALA and GhostKOALA: KEGG Tools for**
Functional Characterization of Genome and Metagenome Sequences. *J Mol Biol*
2016, **428**(4):726-731.

37.   Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ: **Prodigal:**
prokaryotic gene recognition and translation initiation site identification. *Bmc*
Bioinformatics 2010, **11**.

38.   Kanehisa M, Sato Y: **KEGG Mapper for inferring cellular functions from protein**
sequences. *Protein Sci* 2020, **29**(1):28-35.

39.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL:
BLAST plus : architecture and applications. *Bmc Bioinformatics* 2009, **10**.

40.   Elbourne LDH, Tetu SG, Hassan KA, Paulsen IT: **TransportDB 2.0: a database for**
exploring membrane transporters in sequenced genomes from all domains of
life. *Nucleic Acids Res* 2017, **45**(D1):D320-D324.

41.   Coordinators NR: **Database resources of the National Center for Biotechnology**
Information. *Nucleic Acids Res* 2018, **46**(D1):D8-D13.

42.   Edgar RC: **MUSCLE: a multiple sequence alignment method with reduced time**
and space complexity. *Bmc Bioinformatics* 2004, **5**:113.

43.   Kumar S, Stecher G, Tamura K: **MEGA7: Molecular Evolutionary Genetics**
Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 2016, **33**(7):1870-1874.

44.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler
A, Lanfear R: **IQ-TREE 2: New Models and Efficient Methods for Phylogenetic**
Inference in the Genomic Era. *Mol Biol Evol* 2020, **37**(5):1530-1534.

45.   Letunic I, Bork P: **Interactive Tree Of Life (iTOL) v4: recent updates and new**
developments. *Nucleic Acids Res* 2019, **47**(W1):W256-W259.

46.   Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA,
Hugenholtz P: **A standardized bacterial taxonomy based on genome phylogeny**
substantially revises the tree of life. *Nat Biotechnol* 2018, **36**(10):996-+.

47.   Whatley JM: **The fine structure of Prochloron.** *New Phytologist* 1977,
79(2):309-313.

48.   Podell S, Blanton JM, Oliver A, Schorn MA, Agarwal V, Biggs JS, Moore BS, Allen
EE: **A genomic view of trophic and metabolic diversity in clade-specific**
Lamellodysidea sponge microbiomes. *Microbiome* 2020, **8**(1):97.

49.   Richter M, Rossello-Mora R: **Shifting the genomic gold standard for the**

prokaryotic species definition. *Proc Natl Acad Sci U S A* 2009,
106(45):19126-19131.

50.   Qin QL, Xie BB, Zhang XY, Chen XL, Zhou BC, Zhou J, Oren A, Zhang YZ: **A**
proposed genus boundary for the prokaryotes based on genomic insights. *J*
Bacteriol 2014, **196**(12):2210-2215.

51.   Kuhl M, Behrendt L, Trampe E, Qvortrup K, Schreiber U, Borisov SM, Klimant I,
Larkum AW: **Microenvironmental Ecology of the Chlorophyll b-Containing**
Symbiotic Cyanobacterium Prochloron in the Didemnid Ascidian Lissoclinum
patella. *Front Microbiol* 2012, **3**:402.

52.   Mehbub MF, Perkins MV, Zhang W, Franco CMM: **New marine natural products**
from sponges (Porifera) of the order Dictyoceratida (2001 to 2012); a promising
source for drug discovery, exploration and future prospects. *Biotechnol Adv*
2016, **34**(5):473-491.

53.   Schorn MA, Jordan PA, Podell S, Blanton JM, Agarwal V, Biggs JS, Allen EE, Moore
BS: **Comparative Genomics of Cyanobacterial Symbionts Reveals Distinct,**
Specialized Metabolism in Tropical Dysideidae Sponges. *mBio* 2019, **10**(3).

54.   Burgsdorf I, Slaby BM, Handley KM, Haber M, Blom J, Marshall CW, Gilbert JA,
Hentschel U, Steindler L: **Lifestyle evolution in cyanobacterial symbionts of**
sponges. *mBio* 2015, **6**(3):e00391-00315.

55.   Lesser MP, Fiore C, Slattery M, Zaneveld J: **Climate change stressors destabilize**
the microbiome of the Caribbean barrel sponge, Xestospongia muta. *J Exp Mar*
Biol Ecol 2016, **475**:11-18.

56.   Morrow KM, Fiore CL, Lesser MP: **Environmental drivers of microbial community**
shifts in the giant barrel sponge, Xestospongia muta, over a shallow to
mesophotic depth gradient. *Environ Microbiol* 2016, **18**(6):2025-2038.

57.   Webster NS, Reusch TBH: **Microbial contributions to the persistence of coral**
reefs. *ISME J* 2017, **11**(10):2167-2174.

58.   Nguyen MT, Liu M, Thomas T: **Ankyrin-repeat proteins from sponge symbionts**
modulate amoebal phagocytosis. *Mol Ecol* 2014, **23**(6):1635-1645.

59.   Wiens M, Korzhev M, Krasko A, Thakur NL, Perovic-Ottstadt S, Breter HJ, Ushijima
H, Diehl-Seifert R, Muller IM, Muller WEG: **Innate immune defense of the sponge**
Suberites domuncula against bacteria involves a MyD88-dependent signaling
pathway - Induction of a perforin-like molecule. *J Biol Chem* 2005,
280(30):27949-27959.

60.   Fan L, Reynolds D, Liu M, Stark M, Kjelleberg S, Webster NS, Thomas T: **Functional**
equivalence and evolutionary convergence in complex communities of

microbial sponge symbionts. *P Natl Acad Sci USA* 2012, **109**(27):E1878-E1887.

61.   Reveillaud J, Maignien L, Eren AM, Huber JA, Apprill A, Sogin ML, Vanreusel A:
Host-specificity among abundant and rare taxa in the sponge microbiome. *Isme*
Journal 2014, **8**(6):1198-1209.

62.   Burgsdorf I, Erwin PM, Lopez-Legentil S, Cerrano C, Haber M, Frenk S, Steindler L:
Biogeography rather than association with cyanobacteria structures symbiotic
microbial communities in the marine sponge Petrosia ficiformis. *Frontiers in*
Microbiology 2014, **5**.

63.   Gao ZM, Wang Y, Tian RM, Wong YH, Batang ZB, Al-Suwailem AM, Bajic VB, Qian
PY: **Symbiotic Adaptation Drives Genome Streamlining of the Cyanobacterial**
Sponge Symbiont "Candidatus Synechococcus spongiarum". *Mbio* 2014, **5**(2).

64.   Lo WS, Huang YY, Kuo CH: **Winding paths to simplicity: genome evolution in**
facultative insect symbionts. *Fems Microbiol Rev* 2016, **40**(6):855-874.

65.   Moran NA, McCutcheon JP, Nakabachi A: **Genomics and Evolution of Heritable**
Bacterial Symbionts. *Annu Rev Genet* 2008, **42**:165-190.

66.   Kolman MA, Nishi CN, Perez-Cenci M, Salerno GL: **Sucrose in cyanobacteria: from**
a salt-response molecule to play a key role in nitrogen fixation. *Life (Basel)* 2015,
5(1):102-126.

67.   Curatti L, Giarrocco LE, Cumino AC, Salerno GL: **Sucrose synthase is involved in**
the conversion of sucrose to polysaccharides in filamentous nitrogen-fixing
cyanobacteria. *Planta* 2008, **228**(4):617-625.

68.   Blumwald E, Telor E: **Osmoregulation and Cell Composition in Salt-Adaptation of**
Nostoc-Muscorum. *Arch Microbiol* 1982, **132**(2):168-172.

69.   Desplats P, Folco E, Salerno GL: **Sucrose may play an additional role to that of an**
osmolyte in Synechocystis sp PCC 6803 salt-shocked cells. *Plant Physiol Bioch*
2005, **43**(2):133-138.

70.   Frank AC: **Molecular host mimicry and manipulation in bacterial symbionts.**
FEMS Microbiol Lett 2019, **366**(4).

71.   Taylor MW, Radax R, Steger D, Wagner M: **Sponge-associated microorganisms:**
evolution, ecology, and biotechnological potential. *Microbiol Mol Biol Rev* 2007,
71(2):295-347.

72.   Wilkinson CR: **Net primary productivity in coral reef sponges.** *Science* 1983,
219(4583):410-412.

73.   Wang JT, Hsu CM, Kuo CY, Meng PJ, Kao SJ, Chen CA: **Physiological**
Outperformance at the Morphologically-Transformed Edge of the
Cyanobacteriosponge Terpios hoshinota (Suberitidae: Hadromerida) when

913   **Confronting Opponent Corals.** *Plos One* 2015, **10**(6).

74. Zhang F, Blasiak LC, Karolin JO, Powell RJ, Geddes CD, Hill RT: **Phosphorus sequestration in the form of polyphosphate by microbial symbionts in marine sponges.** *P Natl Acad Sci USA* 2015, **112**(14):4381-4386.

75. Colman AS: **Sponge symbionts and the marine P cycle.** *P Natl Acad Sci USA* 2015, **112**(14):4191-4192.

76. Utami RT, Zamani NP, Madduppa HH: **Molecular identification, abundance and distribution of the coral-killing sponge Terpios hoshinota in Bengkulu and Seribu Islands, Indonesia.** *Biodiversitas Journal of Biological Diversity* 2018, **19**(6):2238-2246.

77. Bryan PG: **Growth rate, toxicity and distribution of the encrusting sponge Terpios sp.(Hadromerida: Suberitidae) in Guam, Mariana Islands.** *Micronesica* 1973, **9**(2):237-242.

78. Thinesh T, Meenatchi R, Lipton AN, Anandham R, Jose PA, Tang SL, Seghal Kiran G, Selvin J: **Metagenomic sequencing reveals altered bacterial abundance during coral-sponge interaction: Insights into the invasive process of coral-killing sponge Terpios hoshinota.** *Microbiol Res* 2020, **240**:126553.

79. Teruya T, Nakagawa S, Koyama T, Arimoto H, Kita M, Uemura D: **Nakiterpiosin and nakiterpiosinone, novel cytotoxic C-nor-D-homosteroids from the Okinawan sponge Terpios hoshinota.** *Tetrahedron* 2004, **60**(33):6989-6993.

80. Teruya T, Nakagawa S, Koyama T, Suenaga K, Uemura D: **Terpiodiene: A novel tricyclic alcohol from the Okinawan sponge Terpios hoshinota.** *Chem Lett* 2002(1):38-39.

81. Fagundes MB, Falk RB, Facchi MMX, Vendruscolo RG, Maroneze MM, Zepka LQ, Jacob-Lopes E, Wagner R: **Insights in cyanobacteria lipidomics: A sterols characterization from Phormidium autumnale biomass in heterotrophic cultivation.** *Food Res Int* 2019, **119**:777-784.

Figure Legends

FIG 1. Proportions of *T. hoshinota*-associated bacterial communities across oceans.

Proteobacteria (blue) and *Cyanobacteria* (green) were the dominant groups, ranging

from 81.5–97.4% of community. (a) Bacterial compositions across the western Pacific

Ocean, South China Sea, and Indian Ocean. Maps on the right show results in Okinawa,

Japan (b), and Taiwan (c). Different colors represent different bacterial phyla. Percentage

composition of phyla less than 0.05% are grouped as "Others."

FIG 2. Electron micrographs of *T. hoshinota* and associated cyanobacteria.

(a-c, SEM; d-f, TEM). (a) Sponge covering an abiotic substratum. The sponge's skeleton

was arranged in a tangential style, and the mesophyll was supported by a tract of spicules.

(b, c) Close view of the mesophyll. Numerous spheres composing the inside of the

mesophyll, and cross-matched with (d); their thylakoids identified the spheres as

cyanobacteria. The detailed structures of the cyanobacteria were observed from a single

cyanobacterial cell (e) and a dividing cell (f); they included the cell wall, thylakoid,

carboxysome (ca), phosphate granule (p), and nucleoid (n).

FIG 3. Molecular phylogenetic analyses of *Prochloron* and *Prochloron*-like bacteria.

(a) Pruned phylogenetic tree based on the concatenation of 120 single-copy gene protein

sequences. The complete phylogenetic tree including other cyanobacterial genera can be

found in Supplementary figure 5. The branches with Ultrafast bootstrap (UFBoot) value

>95% are highlighted with the red. The *Prochloron* genomes are labeled with yellow, and

LD05 and SP5CPC1 are labeled with pink. *Vampirovibrio chlorellavorus* C was used as

the outgroup. (b) A phylogenetic tree was constructed using the 16S rRNA gene by the

maximum-likelihood method with 1,000 bootstraps. The tree included 16S rRNA gene

sequences from SP5CPC1, LD05, *Prochloron* genomes—including *P. didemni* P2, *P.*

didemni P3, and *P. didemni* P4—and other related 16S rRNA gene sequences identified

by PCR cloning from environmental samples including. The 16S rRNA gene sequence of

Stanieria sp. NIES-3757 genome was used as an outgroup. The scare bar represents the

number of changes per nucleotide. Asterisk represents the genomes are available.

FIG 4. Extracted ion-chromatograms (EICs) of UPLC-QTOF-MS of the extracts of *T.*

hoshinota. Chlorophyll *a* analytical standard (a) and chlorophyll *b* analytical standard (b)

were used in the analysis. The analysis included two *T. hoshinota* samples. EIC of MS

spectra within the m/z value of 893.543 (a, c, e) and 907.522 (b, d, f). *a*: chlorophyll *a*

peak. *b*: chlorophyll *b* peak.

FIG 5. Metabolic potentials and putative nutrient cycling between *T. hoshinota* and *Ca.*

Pb. terpiosii LD05.

The bacterium may recycle nitrogen waste from host cells and provide them with

vitamins. The bacterium may also produce secondary metabolites as toxins to help the

sponge weaken coral tissues and facilitate overgrowth. On the other hand, the bacterium

may also store phosphate as an energy reservoir for the holobiont to prepare for the

competition with coral. Abbreviation: amt, ammonia transporter; NRT, nitrate transporter;

NR, nitrate reductase; NiR, ferredoxin--nitrite reductase; CA, carbonic anhydrase; PPK,

polyphosphate kinase; GS, glutamine synthetase; GOGAT, glutamine oxoglutarate

aminotransferase; RuBisCo, ribulose-1,5-bisphosphate carboxylase/oxygenase; pcyA,

phycocyanobilin:ferredoxin oxidoreductase; HO, heme oxygenase; pebA,

dihydrobiliverdin:ferredoxin oxidoreductase; pebB, phycoerythrobilin:ferredoxin

oxidoreductase. *The structures of nakiterpiosin and nakiterpiosinone are depicted based

on Shuanhu Gao et al (2012).

Figure 1

Figure 2

Figure 3

a

QS-4-48-99 sp003021505
Rubidibacter lacunae
Halothece salina
Spirulina major
ESFC-1 sp000380225
LD05
SP5CPC1
***Prochloron didemni* P4**
***Prochloron didemni* P3**
***Prochloron didemni* P2**
Limnothrix rosea
Geminocystis aponinum
Cyanobacterium stanieri
Stanieria cyanosphaera
Xenococcus sp000332055
Pleurocapsa sp000332195
Myxosarcina sp000756305
Gloeocapsa sp000332035
Aphanothece hegewaldii
Microcystis aeruginosa A
Hydrococcus minor
Cyanothece B sp000021825
Snowella sp003249095
Synechocystis sp000284455
Cyanothece A sp000021805
Crocosphaera watsonii

Tree scale: 0.1

b

Host

Stanieria sp. NIES-3757
LD05* | *Terpios hoshinota*

Clade I

Uncultured *Prochloron* sp. clone NA108 (DQ357945.1) | *Trididemnum clinides*
97
Uncultured *Synechocystis* sp. clone 29P35 (AY845228.1)
100   Uncultured *Synechocystis* sp. clone 29P18 (AY845227.1) | *Lendenfeldia chondrodes*
84   Uncultured *Synechocystis* sp. clone 36P1 (AY845229.1) | *Lamellodysidea herbacea*
Prochloron sp. SP5CPC1*
Uncultured *Prochloron* sp. clone AO105 (DQ357958.1) | *Trididemnum nubilum*
Uncultured bacterium isolate VG EF A19 (JN126199.1) | *Ectyoplasia ferox*
100   Uncultured bacterium isolate VG EF A02 (JN126186.1)
(Uncultured) *Synechocystis* sp. PLV1 (JX099359.1) | *Lissoclinum verrilli*
Uncultured bacterium clone AF1037 C19 (JN655231.1)
90   Uncultured bacterium clone AF1097 C30 (JX206500.1)
69   Uncultured bacterium clone AF1097 C21 (JX206495.1)
Uncultured bacterium clone AF1097 C8 (JX206482.1) | *Ircinia fasciculata*
Uncultured bacterium clone AF1097 C14 (JX206488.1)
Uncultured bacterium clone AF1097 C28 (JX206499.1)
Uncultured bacterium clone AF1097 C32 (JX206502.1)

Uncultured *Prochloron* sp. clone NA110 (DQ357960.1)
Uncultured *Prochloron* sp. clone HI5 (DQ357951.1)
59   Uncultured *Prochloron* sp. clone BO136 (DQ385852.1)
99   Uncultured *Prochloron* sp. clone HA96 (DQ357967.1)
Uncultured *Prochloron* sp. clone E11036 (KU511264.1)
Uncultured *Prochloron* sp. clone AO100 (DQ357955.1)

Clade II

100
Prochloron didemni P2*
78   Uncultured *Prochloron* sp. clone E11040 (KU511263.1)
Prochloron didemni P3*
Uncultured *Prochloron* sp. clone LI88 (DQ357969.1)
Uncultured *Prochloron* sp. clone LV5 (KU511266.1)
Uncultured *Prochloron* sp. clone Okinawa (KU511265.1)
Uncultured *Prochloron* sp. clone E11016 (KU511267.1)
Uncultured *Prochloron* sp. clone NA112 (DQ357961.1)
Prochloron didemni P4*
Uncultured *Prochloron* sp. clone HI10 (DQ357950.1)
Uncultured *Prochloron* sp. clone HW6 (DQ357949.1)
Uncultured *Prochloron* sp. clone AO101 (DQ357968.1)
52   Uncultured *Prochloron* sp. clone AK122 (DQ357953.1)
Uncultured *Prochloron* sp. clone TA113 (DQ357962.1)
87   Uncultured *Prochloron* sp. clone NA107 (DQ357966.1)
82   Uncultured *Prochloron* sp. clone AK125 (DQ357952.1)
Uncultured *Prochloron* sp. clone AK121 (DQ357954.1)

From various ascidians

Tree scale: 0.01

Bacteria isolated from sponges

Bacteria isolated from ascidians

Figure 4

Figure 5

Mesophyll of *Terpios hoshinota*