The excitatory/inhibitory balance of synaptic transmission

is impaired at higher extracellular Ca^{2+} concentrations in

***Scn1a$^{+/-}$* mouse model of Dravet syndrome**

Kouya Uchino[1,#], Hiroyuki Kawano[2,#], Yasuyoshi Tanaka[2,#], Yuna Adaniya[1], Ai Asahara[1],

Masanobu Deshimaru[2], Kaori Kubota[1], Takuya Watanabe[1], Shutaro Katsurabayashi[1,2,*],

Katsunori Iwasaki[1] and Shinichi Hirose[2,3]

[1]Department of Neuropharmacology, Faculty of Pharmaceutical Sciences, Fukuoka

University, Fukuoka, Japan, [2]Research Institute for the Molecular Pathogeneses of Epilepsy,

Fukuoka University, Fukuoka, Japan, [3]Department of Pediatrics, School of Medicine,

Fukuoka University, Fukuoka, Japan, [#]These authors contributed equally to this study.

*Correspondence should be addressed to:

Shutaro Katsurabayashi, Ph.D.

Department of Neuropharmacology, Faculty of Pharmaceutical Sciences,

Fukuoka University, 8-19-1 Nanakuma, Jonan-ku, Fukuoka 814-0180, Japan.

E-mail: shutarok@fukuoka-u.ac.jp

Tel: (+81) 92-871-6631 (ext. 6634), Fax: (+81) 92-863-0389

Number of text pages: 34

Number of words: 2464

Number of references: 29

Number of figures: 5

Number of tables: 0

ORCID numbers of first, corresponding, and senior authors: 0000-0003-1300-8469 (K.U),

0000-0001-7490-887X (H.K.), 0000-0002-9053-4342 (Y.T.), 0000-0002-8595-5820 (S.K.),

and 0000-0001-6796-3790 (S.H.)

Abstract

Dravet syndrome (DS) is an intractable form of childhood epilepsy that occurs in infancy.

More than 80% of all patients have a heterozygous abnormality in the *SCN1A* gene, which

encodes a subunit of Na^+ channels in the brain. However, the detailed pathogenesis of DS

remains unclear. This study investigated the synaptic pathogenesis of this disease in terms of

excitatory/inhibitory balance using a mouse model of DS. We show that excitatory

postsynaptic currents were similar between *Scn1a* knock-in neurons (*Scn1a*$^{+/-}$ neurons) and

wild-type neurons, but inhibitory postsynaptic currents were significantly lower in *Scn1a*$^{+/-}$

neurons. Moreover, both the vesicular release probability and the number of inhibitory

synapses were significantly lower in *Scn1a*$^{+/-}$ neurons compared with wild-type neurons.

There was no proportional increase in inhibitory postsynaptic current amplitude in response

to increased extracellular Ca^{2+} concentrations. Our study revealed that the number of

inhibitory synapses is significantly reduced in *Scn1a*$^{+/-}$ neurons, while the sensitivity of

inhibitory synapses to extracellular Ca^{2+} concentrations is markedly increased. These data

suggest that Ca^{2+} tethering in inhibitory nerve terminals may be disturbed following the

synaptic burst, likely leading to epileptic symptoms.

Introduction

Epilepsy is a pervasive and severe neurological disorder that affects 0.5%–1% of the

population, predominantly in childhood.[1] Epilepsy patients are generally treated with

antiepileptic drugs, but about 30% of patients have refractory epilepsy that does not respond

to such treatment.[2] The molecular mechanisms underlying the onset of refractory epilepsy

remain mostly unknown. Therefore, epilepsy is one of the most challenging diseases for

developing fundamental therapies and drugs. In addition, pediatric patients with intractable

epilepsy may develop complications, such as developmental disorders and mental retardation,

leading to anxiety about their social lives even in adulthood.[3,4]

We have previously identified more than 100 genetic abnormalities in Na$^+$ channels via

the genetic analysis of patients with intractable epilepsy resulting in mental retardation.[5,6]

Dravet syndrome (DS) is a severe form of epilepsy that develops in infancy and causes

frequent seizures, leading to severe encephalopathy and developmental disorders.

Heterozygous mutations in *SCN1A*, the gene encoding Na$_v$1.1 (the α subunit of voltage-gated

Na$^+$ channels), have been identified in more than 80% of patients with DS. However, it

remains unclear how this gene mutation causes intractable epilepsy. Recently, knock-in,

knock-out, and conditional knock-out mice with heterozygous mutations in the *Scn1a* gene

have been reported as mouse models of DS.[7-10] In parallel with the development of mouse

models, we established induced pluripotent stem cells from a DS patient with a heterozygous

nonsense mutation in the *SCN1A* gene.[11,12] We reported an apparent fragile mutation in the

maintenance of action potentials during sustained depolarizing stimulation of γ-aminobutyric

acid (GABA)ergic neurons. The common mechanism of these aforementioned studies was

that *SCN1A* mutations result in a loss of inhibitory neural function. This underlying

mechanism indicates that excitatory neural function is increased in the brain, making it highly

susceptible to epilepsy because of the suppression of inhibitory neural function in the central

nervous network.

In the brain, neurons connect via synapses and transmit information in a fine-tuned

manner to maintain central homeostasis. During epileptogenesis, the excitatory/inhibitory

(E/I) balance of the multicellular neural network is transiently or persistently disrupted,

resulting in overexcited synaptic activity.[13,14] Therefore, understanding epilepsy as a

"synaptic pathology" and elucidating synaptic recovery mechanisms may help to develop

fundamental epilepsy therapies and aid in drug discovery.

Results

Establishment of *Scn1a*-targeted mice

The targeting vector consisted of the neomycin-resistance cassette, of which the 5' and 3' ends

were connected to recombination arms derived from intron 4 to intron 7 (left arm) and intron

12 to intron 16 (right arm) of the mouse *Scn1a* gene, respectively (Figure 1A). After this was

introduced into mouse ES cells, 156 neomycin-resistant clones were isolated and screened for

targeted homologous recombination by PCR. Using primers corresponding to the neo gene

and intron 16, a 5.5-kb fragment that would emerge only from recombinant alleles was

observed in three clones. These candidates were further analyzed by Southern hybridization.

Two of them showed positive signals at 7.8 and 6.5 kb, which corresponded to the wild-type

and targeted alleles, respectively (Figure 1B, C). These two clones were regarded as having

desirable recombination and were used to establish the mutant mouse, in which exon 8 to

exon 12 of the *Scn1a* gene was heterogeneously replaced by the neomycin resistance cassette

(*Scn1a*$^{+/-}$ mouse). On establishment and after every mating, mouse specimens with

recombinant alleles were identified by PCR amplification of a 634-bp DNA fragment

spanning from intron 7 to the 5' FRT element.

Excitatory synaptic transmission of *Scn1a*$^{+/-}$ neurons

In the epileptic brain, changes in the E/I balance are exhibited as enhanced excitatory

synaptic transmission and/or reduced inhibitory synaptic transmission, and vice versa. We

first measured excitatory synaptic transmission. The EPSC amplitudes were not significantly

different between WT and *Scn1a*$^{+/-}$ neurons (Figure 2A, B). The mEPSC amplitudes were

also not significantly different between WT and *Scn1a*$^{+/-}$ neurons (Figure 2C, D). We then

analyzed the frequency of mEPSCs to assess the effects on presynaptic machinery, and found

no significant differences between WT and $Scn1a^{+/-}$ neurons (Figure 2E). These results

indicate that there are no changes in the excitatory synaptic transmission of $Scn1a^{+/-}$ neurons.

Inhibitory synaptic transmission of $Scn1a^{+/-}$ neurons

The absence of phenotypes in the excitatory synaptic transmission of $Scn1a^{+/-}$ neurons

(Figure 2) suggested that the inhibitory synaptic transmission of these neurons may be

reduced, in terms of the E/I balance between excitatory and inhibitory synaptic transmission

during epileptogenesis. Therefore, in the following experiment, we analyzed the inhibitory

synaptic transmission of $Scn1a^{+/-}$ neurons.

As predicted, IPSC amplitudes (i.e., inhibitory synaptic transmission elicited by action

potentials) were significantly lower in $Scn1a^{+/-}$ neurons compared with WT neurons (Figure

3A, B). In contrast, mIPSC amplitudes did not differ significantly between $Scn1a^{+/-}$ and WT

neurons (Figure 3C, D).

The finding that the amplitudes of IPSCs elicited by action potentials were reduced

while the amplitudes of mIPSCs remained unchanged suggests that presynaptic function,

rather than postsynaptic function, was affected in inhibitory synaptic transmission. To

investigate presynaptic function, we analyzed the frequency of spontaneous mIPSCs. There

were no significant differences in spontaneous mIPSC frequency between $Scn1a^{+/-}$ and WT

neurons (Figure 3E). These results suggest that inhibitory synaptic transmission in $Scn1a^{+/-}$

neurons may be reduced by action potential-dependent mechanisms.

Mechanisms of inhibitory synaptic transmission in $Scn1a^{+/-}$ neurons

Because there was no significant decrease in the frequency of mIPSCs in $Scn1a^{+/-}$ neurons, it

remained unclear whether presynaptic function was impaired in these cells. We therefore

measured the size of the RRP. There was no significant difference in RRP size between

$Scn1a^{+/-}$ and WT neurons (Figure 4A, B). However, the vesicular release probability (P_{vr}) of

synaptic vesicles was significantly lower in $Scn1a^{+/-}$ neurons compared with WT neurons

(Figure 4C).

Reduced inhibitory synaptic transmission may also indicate that the number of

inhibitory synapses is reduced. We therefore measured the numbers of inhibitory synapses in

WT and $Scn1a^{+/-}$ neurons using immunostaining. The number of GABAergic synapses was

significantly lower in $Scn1a^{+/-}$ neurons than in WT neurons (Figure 4D, E). These results

indicate that the decreased inhibitory synaptic transmission in $Scn1a^{+/-}$ neurons is likely

caused by a combination of a lower P_{vr} and a lower number of inhibitory synapses.

Mechanisms of depressed inhibitory synaptic vesicular release in $Scn1a^{+/-}$ neurons

The generation of action potentials depolarizes nerve terminals, and the subsequent influx of

extracellular Ca^{2+} into the nerve terminals causes activity-dependent neurotransmitter release.

Thus, the lower P_{vr} in the inhibitory synaptic transmission of $Scn1a^{+/-}$ neurons may be caused

by a lower sensitivity to extracellular Ca^{2+} concentrations ($[Ca^{2+}]_o$). We therefore examined

the changes in inhibitory synaptic transmission in response to higher or lower $[Ca^{2+}]_o$. There

was no proportional increase in IPSCs at higher calcium concentrations (4 mM and 8 mM) in

$Scn1a^{+/-}$ neurons (Figure 5A). To further understand Ca^{2+} sensitivity, the relative amplitudes

of evoked IPSCs were analyzed using Hill's equation, with IPSC at 8 mM as a reference.[21]

Compared with WT neurons, the graph of relative IPSC amplitude was shifted to the left in

$Scn1a^{+/-}$ neurons (Figure 5B), indicating that $Scn1a^{+/-}$ neurons are more sensitive to $[Ca^{2+}]_o$.

The K_d value calculated using Hill's equation is a dissociation constant. Its reciprocal is a

coupling constant, which can be approximated as the ion permeability in this case. In contrast,

the Hill coefficient indicates cooperativity with ions. The K_d values were reduced to about

one-third of WT values in $Scn1a^{+/-}$ neurons (WT: 0.60 ± 0.05 mM, $Scn1a^{+/-}$: 0.18 ± 0.10

mM), while Hill coefficients were higher compared with WT neurons (WT: 1.34 ± 0.15,

$Scn1a^{+/-}$: 2.03 ± 0.89). Thus, $Scn1a^{+/-}$ neurons had higher affinity and cooperativity with

$[Ca^{2+}]_o$ than WT neurons. These results indicate that the reduced inhibitory synaptic release

in $Scn1a^{+/-}$ neurons is not caused by low sensitivity to $[Ca^{2+}]_o$, but rather by high sensitivity

to $[Ca^{2+}]_o$.

E/I balance of *Scn1a*$^{+/-}$ neurons at high extracellular Ca^{2+} concentrations

Based on the high [Ca^{2+}]$_o$ sensitivity of the inhibitory synapses of *Scn1a*$^{+/-}$ neurons, we

hypothesized that *Scn1a* disruption might cause changes in the Ca^{2+} response of neurons.

Because epileptic seizures are caused by excessive Ca^{2+} influx into nerve terminals,[14] we

predicted that increasing [Ca^{2+}]$_o$ would induce a pseudo-epileptic condition and disrupt the

E/I balance even in the normal state. We therefore investigated the magnification of

excitatory and inhibitory synaptic transmission when [Ca^{2+}]$_o$ was changed from 2 mM to 8

mM. In excitatory neurons, there was no significant difference in the magnification of EPSCs

between WT and *Scn1a*$^{+/-}$ neurons (Figure 5C, D). This result is likely because EPSC

amplitudes increase in proportion to an increase in [Ca^{2+}]$_o$ in an activity-dependent manner.

Conversely, in inhibitory neurons, the IPSC amplitude magnification was significantly lower

in *Scn1a*$^{+/-}$ neurons than in WT neurons (Figure 5E, F). This finding indicates that IPSC

amplitudes were unable to increase in proportion to the increase in [Ca^{2+}]$_o$. This phenomenon

may be the result of significantly fewer synapses and increased [Ca^{2+}]$_o$ sensitivity at

inhibitory synapses in *Scn1a*$^{+/-}$ neurons. Combined with the finding of lower IPSC amplitude

at 2 mM [Ca^{2+}]$_o$, these results indicate that, for *Scn1a*$^{+/-}$ neurons, a higher [Ca^{2+}]$_o$ worsens the

E/I balance.

Discussion

E/I balance in the brains of $Scn1a^{+/-}$ mice

Neurons in the brain form networks that function by maintaining a reciprocal E/I balance.

When epilepsy develops, there is generally a temporary or persistent disruption of the E/I

balance.[13,14] We therefore predicted that the E/I balance would be altered by either an

increase in excitatory synaptic transmission or a decrease in inhibitory synaptic transmission.

In the present study, we did not observe any enhancement of excitatory synaptic transmission

at the single-neuron level in $Scn1a^{+/-}$ neurons. However, inhibitory synaptic transmission was

significantly lower in $Scn1a^{+/-}$ neurons than in WT neurons. The lack of any differences in

excitatory synaptic transmission may be because of the distribution of $Na_v1.1$. These subunits

have been reported to be expressed predominantly on the cell body of neurons,[22,23] but a

recent report revealed that $Na_v1.1$ is not expressed in excitatory neurons in the hippocampus.[8]

Although we did not confirm the expression of $Na_v1.1$ in the mouse model used in the present

study, our electrophysiological results strongly support the previous finding that $Na_v1.1$ is not

expressed in hippocampal excitatory neurons. It has also been reported that $Scn1a$-deficient

mice show abnormalities in inhibitory neuron function,[7] which is consistent with our results.

However, in *in vivo* conditions, the disruption of $Scn1a$ appears to cause abnormalities in

both excitatory and inhibitory neurons.[24] Thus, future studies need to examine neural

networks with mixed excitatory and inhibitory synaptic inputs.

Synaptic function in $Scn1a^{+/-}$ neurons

Although the RRP sizes at all synapses were similar between WT and $Scn1a^{+/-}$ neurons, the

number of GABAergic synapses was significantly lower in $Scn1a^{+/-}$ neurons. These results

suggest that the number of releasable synaptic vesicles per synapse is higher in $Scn1a^{+/-}$

neurons than in WT. We had hypothesized that the lower P_{vr} in $Scn1a^{+/-}$ neurons was caused

by reduced sensitivity to $[Ca^{2+}]_o$; however, the results were contrary to our expectations

(Figure 5B). Therefore, other mechanisms may affect the tethering system of Ca^{2+} influx into

nerve terminals. For example, synaptotagmin is a Ca^{2+} sensor for synaptic release. [25] It is thus

possible that the function of synaptotagmin was affected in the nerve terminals of $Scn1a^{+/-}$

neurons.

In general, epileptic seizures are caused by hyperexcitation of nerves, caused by the

influx of large amounts of Ca^{2+} from extracellular sources. It has recently been demonstrated

that a large amount of Ca^{2+} influx also occurs during epileptic seizures in animal models of

DS.[14] In the present study, we simulated epileptic seizures by increasing the $[Ca^{2+}]_o$, and

found that the E/I balance in $Scn1a^{+/-}$ neurons was more disrupted during pseudo-epileptic

seizures than during normal conditions. Although the mouse model used in this study only

had mutated *Scn1a*, which encodes $Na_v1.1$, there was an alteration in the Ca^{2+} response of

Scn1a$^{+/-}$ neurons. One reason for this may involve the expression of Ca^{2+} channels; it is

known that *Scn1a* disruption alters the expression of Na^+ channels as well as other ion

channels.[26] Therefore, disruption of the *Scn1a* gene may have caused changes in Ca^{2+}

channels, and thus altered Ca^{2+} responses in *Scn1a$^{+/-}$* neurons.

Susceptibility of *Scn1a$^{+/-}$* mice to seizures

When epilepsy is induced, it may be triggered by a small external stimulus. For example, the

firing of a spontaneous electrical potential, which generally has no physiological

consequences, may stimulate epileptogenic cells to induce bursts of action potential firings,

which then propagate through the neuronal network. Thus, even if a change in E/I balance

does not have severe consequences in the normal condition, visible disruption may occur in

the brains of *Scn1a$^{+/-}$* mice if they are experimentally subjected to a small number of external

stimuli (e.g., frequent stimulation or drug stimulation).

However, given that epilepsy tends to occur in juveniles, additional external stimulation

of juvenile synapses *in vitro* may be an effective way to detect small disruptions. At the

cellular level, intracellular Cl^- concentrations are high in the juvenile stage and low in the

mature stage.[27-29] That is, in juveniles, GABAergic input may cause Cl^- to leak out of cells

via $GABA_A$ receptors, resulting in cell depolarization. This developmental change may be

one of the mechanisms that contributes to seizure susceptibility. In conclusion, this study

suggests that mutations in the *Scn1a* gene can cause epileptic seizures, even under apparently

normal conditions.

Methods

Vector construction

The structure of the targeting vector is shown in Figure 1. Briefly, a 5.4 kb murine *Scn1a*

genomic fragment spanning from the end of exon 7 to the beginning of exon 5 (left arm) and

a 5.0 kb murine *Scn1a* genomic fragment spanning from the end of exon 13 to the beginning

of exon 16 (right arm) were amplified by polymerase chain reaction (PCR) from the bacterial

artificial chromosome (BAC) clone TRPCI23-201O18 (Advanced GenoTechs). The

accession number of the corresponding *Scn1a* gene reference sequence is NC_000068.7. The

resulting construct was modified to contain a lox71 sequence downstream of *Scn1a* exon 7,

and a neomycin-resistance (neo) cassette was added further downstream, which comprised

PPGK (the promoter sequence of the mouse phosphoglycerate kinase 1 [*Pgk*] gene), a

lox2272 sequence, and a *Pgk* polyadenylation sequence. The neo cassette was additionally

flanked by two yeast-derived flippase (FLP) recognition target sequences (FRTs) to allow the

removal of the drug-resistance cassette in a flippase-enabled background. The *Scn1a*

fragment was 5'-fused with a diphtheria toxin-A cassette, PCR subcloned from plasmid p03.

The full construct was inserted into the multiple cloning site of pBluescript II SK (+)

(Stratagene).

Culture and gene manipulation of mouse embryonic stem (ES) cells

Feeder-free mouse ES cells, which were TT2-KTPU8 strain (C57BL/6J × CBA) F1-derived

wild-type ES cells, were seeded at 6.0×10^6 cells per dish in collagen-coated 10-cm diameter

dishes containing Glasgow minimum essential medium (Thermo Fisher Scientific)

supplemented with 1% fetal calf serum, 14% knock-out serum replacement (Thermo Fisher

Scientific), and 1 kU/mL leukemia inhibitory factor (Thermo Fisher Scientific). After a 2-day

incubation at 37°C in 6.5% CO_2 to reach semi-confluence (1.6×10^7), the cells were washed

twice with phosphate-buffered saline, resuspended in 1.6 mL phosphate-buffered saline, and

divided into two 0.8-mL aliquots. Transfections were performed using a Gene Pulser

(Bio-Rad) and a 4-mm gap cuvette set at 0.8 kV/3 µF in the presence of a 20 µg

PvuI-linearized targeting vector. Transfected cells were expanded in G418-supplemented

medium (200 µg/mL) for 9 days. Cells with randomly inserted target vector DNA did not

survive during this time because non-homologous recombination failed to remove the

diphtheria toxin-A cassette. Of the approximately 500 colonies that developed, 192 were

isolated by pipetting under a microscope and were then cultured for 3 days. Next, two-thirds

of these cells were suspended in dimethyl sulfoxide-supplemented culture medium and

cryo-banked. The remaining cells were used for genomic DNA extraction, followed by PCR

and Southern analysis. The PCR primers RA/S and RA/A, which were used to amplify a

0.6-kb DNA fragment of hybridization probe, were designed according to the nucleotide

sequences of intron 13 and exon 13 of *Scn1a*, respectively (Figure 1B). The resulting DNA

fragments were used as the probe for Southern blot analysis following EcoRI-/PvuI-digestion

of genomic DNA, which hybridized with 7.8 kb and 6.5 kb fragments for the

non-recombinant and recombinant alleles, respectively (Figure 1C).

Animals

Experimental animals were handled under the ethical regulations for animal experiments of

the Fukuoka University experimental animal care and use committee and animal experiment

implementation. All animal protocols were approved by the ethics committee of Fukuoka

University (permit numbers: 1712128 and 1812092).

Pregnant ICR mice for the astrocyte (a type of glial cell) cultures were purchased from

CLEA Japan, Inc. (Catalogue ID: Jcl:ICR). Mice were individually housed in plastic cages

and kept in an animal house under the following conditions: room temperature 23°C ± 2°C,

humidity 60% ± 2%, and a 12-hour light/dark cycle (7:00 AM lights on), with free access to

water and food (CE-2, Nihon Crea).

Development of *Scn1a*$^{+/-}$ mice

Gene targeted clones of *Scn1a*-mutant ES cells were used to generate *Scn1a*-812neo mice.

We first removed two-cell stage zona pellucida cells from the embryos of the Institute for

Cancer Research (ICR) mouse line and immersed them in mutant ES cell suspension. After

an overnight incubation to enhance the aggregation of embryos and cells, chimeric embryos

were transplanted into the uteri of pseudopregnant females. From the resulting litter,

checker-coated chimeras were mated with C57BL/6J mice to produce F1 heterozygous

offspring with a single mutant *Scn1a* allele. The resulting mouse line containing the *Scn1a*

mutation was backcrossed with C57BL/6J mice over 10 generations to establish a congenic

strain. Mice with an *Scn1a* gene in which exon 8 to exon 12 was heterogeneously replaced

with a neomycin resistance gene were termed *Scn1a*$^{+/-}$ mice.

Genotyping

Genomic DNA was extracted from mouse tails by NaOH extraction. The PCR primers

Scn1ai7prS2 and FRT-RV, which were used to detect the recombinant allele, were designed

according to the nucleotide sequences of intron 7 of the *Scn1a* gene and FRT elements,

respectively. Using PCR, these primers only amplified a 0.6-kb DNA fragment when

genomic DNA containing the homologous recombination was used as the template. Newborn

313 mice were used for the functional analysis of primary neuronal cultures after PCR

314 verification.

Microisland cultures of astrocytes

317 We used a "coating stamp" to obtain uniformly sized cell adhesion areas, divided into dots

318 (squares with a side length of about 300 µm).[15,16] The coating stamp made it possible to

319 minimize variations in the size of each island of astrocyte culture. [15,16]

320  ICR mice immediately after birth (postnatal day [P]0–1) were used for astrocyte cultures.

321 Details of the astrocyte culture method are described in our previous paper.[15,16]

Autapse culture specimens of single neurons

324 We used autapse culture preparations, which are co-cultured preparations of dot-cultured

325 astrocytes and a single neuron, for the synaptic function analysis.[15-19] The axons of autapse

326 neurons form many synapses (autapses) to their cell bodies and dendrites, without any

327 synaptic contacts from other neurons. This feature allowed us to focus on the synapses of a

328 single neuron only, and to analyze the properties of synaptic function in more detail.

329  *Scn1a* knock-in neurons (*Scn1a*$^{+/-}$ neurons) were used in all experiments. For the culture

330 of single neurons, *Scn1a*$^{+/-}$ mice immediately after birth (P0–1) were decapitated to minimize

331 pain/discomfort and used; tails were harvested to confirm disruption of the *Scn1a* gene, and

PCR was used for genotyping. Neurons were enzymatically isolated from the hippocampus of

P0–1 *Scn1a$^{+/-}$* mice using papain (Worthington). Before plating neurons, the conditioned

medium of the microisland cultures was replaced with Neurobasal A medium (Thermo Fisher

Scientific) containing 2% B27 supplement (Thermo Fisher Scientific) and 1% GlutaMAX□I

supplement (Thermo Fisher Scientific). Cells were plated at a density of 1500 cells/cm^2 onto

the astrocyte microislands. Hippocampal neurons were used to analyze excitatory synaptic

transmission and striatal neurons were used to analyze inhibitory synaptic transmission. To

maintain the soluble factors from astrocytes, the medium was not changed during the culture

period. The incubation period was 13–16 days *in vitro*.

Electrophysiology

Synaptic analysis was performed using the patch-clamp method, where the membrane

potential was clamped at –70 mV. An evoked excitatory postsynaptic current (EPSC) or

inhibitory postsynaptic current (IPSC) was recorded in response to an action potential elicited

by a brief (2 ms) somatic depolarization pulse (to 0 mV) from the patch pipette under

voltage-clamp conditions. Action potentials triggering EPSCs or IPSCs were not significantly

different between wild-type (WT) and *Scn1a$^{+/-}$* neurons of the hippocampus and striatum,

respectively (Figure S1). Spontaneous miniature EPSCs (mEPSCs) and IPSCs (mIPSCs)

were recorded in the presence of a Na$^+$ channel inhibitor, 0.5 μM tetrodotoxin (FUJIFILM

Wako Pure Chemical Corporation), in the extracellular fluid. The readily releasable pool

(RRP) size was calculated as the total area of inward current elicited by transient

hyperosmotic extracellular fluid containing 0.5 M sucrose. The vesicular release probability

(P_{vr}) was calculated by dividing the evoked EPSC or IPSC area by the RRP area. Synaptic

responses were recorded at a sampling rate of 20 kHz and were filtered at 10 kHz. Data were

excluded from the analysis if a leak current of > 300 pA was observed. The data were

analyzed offline using AxoGraph X 1.2 software (AxoGraph Scientific).

Extracellular fluid (pH 7.4) was as follows (mM): NaCl 140, KCl 2.4, HEPES 10,

glucose 10, $CaCl_2$ 2, and $MgCl_2$ 1. Intracellular fluid (pH 7.4) for recording EPSCs was as

follows (mM): K-gluconate 146.3, $MgCl_2$ 0.6, ATP-Na_2 4, GTP-Na_2 0.3, creatine

phosphokinase 50 U/mL, phosphocreatine 12, EGTA 1, and HEPES 17.8. The intracellular

fluid for recording IPSCs was prepared by replacing K-gluconate with KCl in the

intracellular fluid for EPSCs. All experiments were performed at room temperature (20°C

–22°C).

Immunocytochemistry

Striatal neurons cultured for 14 days were fixed by being placed in 4% paraformaldehyde in

phosphate-buffered saline for 20 minutes. They were then incubated with primary antibodies

containing blocking solution (0.1% Triton X-100, 2% normal goat serum) at 4°C overnight.

The following primary antibodies were used: anti-microtubule-associated protein 2 (MAP2

(Cat. # 188 004); guinea-pig polyclonal, antiserum, 1:1000 dilution, Synaptic Systems) and

anti-vesicular GABA transporter (VGAT (Cat. # 131 002); rabbit polyclonal, affinity-purified,

1:2000 dilution, Synaptic Systems). Next, the cells were incubated in secondary antibodies

corresponding to each primary antibody (Alexa Fluor 488 for MAP2 and Alexa Fluor 594 for

VGAT, 1:400, Thermo Fisher Scientific) for 40 minutes at room temperature. The nuclei of

astrocytes and autaptic neurons were visualized by counterstaining with

4',6-diamidino-2-phenylindole (DAPI)-containing mounting medium (ProLong® Gold

Antifade Reagent with DAPI, Thermo Fisher Scientific). Only neurons with one nucleus were

analyzed when counting synapse numbers.

Image acquisition and quantification

Confocal images of autaptic neurons were obtained using a confocal microscope (LMS710,

Carl Zeiss) with a C-Apochromat objective lens (40×, NA 1.2, Carl Zeiss), with sequential

acquisition at high-resolution settings (1024 × 1024 pixels). The parameters of each image

were optimized for the z-stack setting (a depth of 9 μm with 0.5 μm steps) and pinhole setting

(1 Airy unit, 0.9 μm). The depth sufficiently covered the thickness of neurites in our autaptic

culture. The confocal microscope settings were identical for all scans in each experiment. A

single autaptic neuron was selected for each scan in a blinded manner, based on MAP2

fluorescence images. The sizes and numbers of synaptic puncta were detected using ImageJ

software. The procedures used to analyze the synaptic puncta have been published

previously.[19,20]

Statistical analysis

All data are expressed as the mean ± standard error. A lowercase n indicates the number of

neurons recorded, and an uppercase N indicates the number of cultures (lot number). Two

groups were compared using the Student's unpaired t-test. Statistical significance was

considered when $p < 0.05$.

Data accessibility statement

The data that support the findings of this study are available from the corresponding author

on reasonable request.

References

1. Sander JW, Shorvon SD. Epidemiology of the epilepsies. *J Neurol Neurosurg Psychiatry*.

1996; **61**: 433–43.

2. Dalic L, Cook MJ. Managing drug-resistant epilepsy: challenges and solutions.

Neuropsychiatr Dis Treat. 2016; **12**: 2605–2616.

3. Hirose S, Okada M, Kaneko S, Mitsudome, A. Are some idiopathic epilepsies disorders of

ion channels?: A working hypothesis. *Epilepsy Res*. 2000; **41**: 191–204.

4. Hirose S, Okada M, Yamakawa K, Sugawara T, Fukuma G, Ito M, et al. Genetic

abnormalities underlying familial epilepsy syndromes. *Brain Dev*. 2002; **24**: 211–22.

5. Hirose S, Mohney RP, Okada M, Kaneko S, Mitsudome A. The genetics of febrile seizures

and related epilepsy syndromes. *Brain Dev*. 2003; **25**: 304–12.

6. Hirose S, Mitsudome A, Okada M, Kaneko S; Epilepsy Genetic Study Group, Japan.

Genetics of idiopathic epilepsies. *Epilepsia*. 2005; **46**: 38–43.

7. Yu FH, Mantegazza M, Westenbroek RE, Robbins CA, Kalume F, Burton KA, et al.

Reduced sodium current in GABAergic interneurons in a mouse model of severe myoclonic

epilepsy in infancy. *Nat Neurosci*. 2006; **9**: 1142–9.

8. Ogiwara I, Miyamoto H, Morita N, Atapour N, Mazaki E, Inoue I, et al. Na$_v$1.1 localizes to

axons of parvalbumin-positive inhibitory interneurons: a circuit basis for epileptic seizures in

mice carrying an *Scn1a* gene mutation. *J Neurosci*. 2007; **27**: 5903–14.

9. Cheah CS, Yu FH, Westenbroek RE, Kalume FK, Oakley JC, Potter GB, et al. Specific

deletion of Na$_v$1.1 sodium channels in inhibitory interneurons causes seizures and premature

death in a mouse model of Dravet syndrome. *Proc Natl Acad Sci U S A*. 2012; **109**:

14646–51.

10. Han S, Tai C, Westenbroek RE, Yu FH, Cheah CS, Potter GB, et al. Autistic-like

behaviour in *Scn1a$^{+/-}$* mice and rescue by enhanced GABA-mediated neurotransmission.

Nature. 2012; **489**: 385–90.

11. Higurashi N, Uchida T, Lossin C, Misumi Y, Okada Y, Akamatsu W, et al. A human

Dravet syndrome model from patient induced pluripotent stem cells. *Mol Brain*. 2013; **6**: 19.

12. Tanaka, Y, Sone, T, Higurashi, N, Sakuma T, Suzuki S, Ishikawa M, et al. Generation of

D1-1 TALEN isogenic control cell line from Dravet syndrome patient iPSCs using

TALEN-mediated editing of the SCN1A gene. *Stem Cell Res*. 2018; **28**, 100–104.

13. Bonansco C, Fuenzalida M. Plasticity of hippocampal excitatory-inhibitory balance:

missing the synaptic control in the epileptic brain. *Neural Plast*. 2016; 8607038.

14. Brenet A, Hassan-Abdi R, Somkhit J, Yanicostas C, Soussi-Yanicostas N. Defective

excitatory/inhibitory synaptic balance and increased neuron apoptosis in a zebrafish model of

Dravet syndrome. *Cells*. 2019; **8**: 1199.

15. Kawano H, Katsurabayashi S, Kakazu Y, Yamashita Y, Kubo N, Kubo M, et al. Long-term

culture of astrocytes attenuates the readily releasable pool of synaptic vesicles. *PLoS One*.

2012; **7**: e48034.

16. Katsurabayashi S, Kawano H, Ii M, Nakano S, Tatsumi C, Kubota K, et al.

Overexpression of Swedish mutant APP in aged astrocytes attenuates excitatory synaptic

transmission. *Physiol Rep*. 2016; **4**: e12665.

17. Bekkers JM, Stevens CF. Excitatory and inhibitory autaptic currents in isolated

hippocampal neurons maintained in cell culture. *Proc Natl Acad Sci U S A*. 1991; **88**: 7834–8.

18. Rosenmund C, Stevens CF. Definition of the readily releasable pool of vesicles at

hippocampal synapses. *Neuron*. 1996; **16**: 1197–207.

19. Oyabu K, Takeda K, Kawano H, Kubota K, Watanabe T, Harata NC, et al. Presynaptically

silent synapses are modulated by the density of surrounding astrocytes. *J Pharmacol Sci*.

2020; **144**: 76–82.

20. Iwabuchi S, Kakazu Y, Koh JY, Harata NC. Evaluation of the effectiveness of Gaussian

filtering in distinguishing punctate synaptic signals from background noise during image

analysis. *J Neurosci Methods*. 2014; **223**: 92–113.

21. Rhee JS, Li LY, Shin OH, Rah J-C, Rizo J, Südhof TC, et al. Augmenting

neurotransmitter release by enhancing the apparent Ca^{2+} affinity of synaptotagmin 1. *Proc

Natl Acad Sci U S A*. 2005; **102**: 18664–9.

22. Westenbroek RE, Merrick DK, Catterall WA. Differential subcellular localization of the

RI and RII Na^{+} channel subtypes in central neurons. *Neuron*. 1989; **3**: 695–704.

23. Gong B, Rhodes KJ, Bekele-Arcuri Z, Trimmer JS. Type I and type II Na^{+} channel

α-subunit polypeptides exhibit distinct spatial and temporal patterning, and association with

auxiliary subunits in rat brain. *J Comp Neurol*. 1999; **412**: 342–52.

24. Gu F, Hazra A, Aulakh A, Žiburkus J. Purinergic control of hippocampal circuit

hyperexcitability in Dravet syndrome. *Epilepsia*. 2014; **55**: 245–55.

25. Fernández-Chacón R, Königstorfer A, Gerber SH, García J, Matos MF, Stevens CF, et al.

Synaptotagmin I functions as a calcium regulator of release probability. *Nature*. 2001; **410**:

41–9.

26. Shi X, He W, Guo S, Zhang B, Ren S, Liu K, et al. RNA-seq analysis of the SCN1A-KO

model based on CRISPR/Cas9 genome editing technology. *Neuroscience*. 2019; **398**: 1–11.

27. Luhmann HJ, Prince DA. Postnatal maturation of the GABAergic system in rat neocortex.

J Neurophysiol. 1991; **65**: 247–63.

28. Chen G, Trombley PQ, van den Pol AN. Excitatory actions of GABA in developing rat

hypothalamic neurones. *J Physiol*. 1996; **494**: 451–64.

29. Kakazu Y, Akaike N, Komiyama S, Nabekura J. Regulation of intracellular chloride by

cotransporters in developing lateral superior olive neurons. *J Neurosci.* 1999; **19**: 2843–51.

Acknowledgments

This work was supported by a KAKENHI Grant-in-Aid for Scientific Research (C) to S.K.

(No. 17K08328) from the Japan Society for the Promotion of Science, and the Science

Research Promotion Fund and The Fukuoka University Fund to S.H. (Nos. G19001 and

G20001), a grant for Practical Research Project for Rare/Intractable Diseases from Japan

Agency for Medical Research and development (AMED) to S.H. (Nos. 15ek0109038h0002

and 16ek0109038h0003), a KAKENHI Grant-in-Aid for Scientific Research (A) to S.H. (No.

15H02548), a KAKENHI Grant-in-Aid for Scientific Research (B) to S.H. (Nos. 20H03651,

20H03443 and 20H04506), the Acceleration Program for Intractable Diseases Research

utilizing Disease-specific iPS cells from AMED to S.H. (Nos. 17bm0804014h0001,

18bm0804014h0002, and 19bm0804014h0003), a Grant-in-Aid for the Research on

Measures for Intractable Diseases to S.H. (H31-Nanji-Ippan-010), Program for the Strategic

Research Foundation at Private Universities 2013-2017 from the Ministry of Education,

Culture, Sports, Science, and Technology (MEXT) to S.H. (No. 924), Center for Clinical and

Translational Research of Kyushu University Hospital to S.H. (No. 201m0203009 j0004). We

thank the members of our laboratory for their help. We also thank Bronwen Gardner, PhD,

from Edanz Group (https://en-author-services.edanz.com/ac) for editing a draft of this

manuscript.

527   Author contributions

528   K.U., H.K., Y.T., Y.A., and A. A. performed experiments and analyzed data; Y.T and M.D.

529   created the $Scn1a^{+/-}$ mouse model; S.K. conceived the study; K.K., T.W., K.I., and S.H.

530   interpreted the data; K.U. and S.K. wrote the manuscript with input from all authors.

532   Disclosure of conflicts of interest

533   None of the authors has any conflict of interest to disclose.

535   Ethical publication statement

536   We confirm that we have read the Journal's position on issues involved in ethical publication

537   and affirm that this report is consistent with those guidelines.

540   Figure legends

Figure 1. Targeting of the *Scn1a* gene in mouse embryonic stem cells.

(**A**) Strategy for *Scn1a* targeting. Upper, overall schematic view of the mouse *Scn1a* gene,

and a magnified view containing the targeted region from exons 8 to 12. Exons are shown as

numbered solid boxes, and introns are shown as horizontal lines. Middle, the organization of

the targeting vector. More detail is indicated in the Materials and Methods. Shaded and solid

triangles indicate the lox and flippase recognition target sequence (FRT) elements,

respectively. Regions to be recombined between genomic *Scn1a* and the targeting vector are

indicated by large Xs. Bottom, the targeted *Scn1a* allele that resulted from desirable

homologous recombination. (**B**) Comparison of the restriction patterns of wild-type and

targeted *Scn1a* alleles in Southern analysis. The position of the hybridization probe is

indicated by a solid bar. (**C**) A result of Southern hybridization for nine candidate clones. For

samples #1 and #4, both the 7.8- and 6.5-kb fragments were detected, which were derived

from the wild-type and homologously recombined *Scn1a* alleles.

Figure 2. No changes in the excitatory synaptic transmission of $Scn1a^{+/-}$ neurons.

(**A**) Representative traces of evoked excitatory postsynaptic currents (EPSCs) recorded from

a single autaptic hippocampal neuron of either wild-type (WT) or $Scn1a^{+/-}$ mice.

Depolarization artifacts caused by the generated action currents have been removed for

clarity of presentation. (**B**) Average amplitudes of the evoked EPSCs in neurons of WT or

$Scn1a^{+/-}$ mice (WT: 9.65 ± 0.62 nA, n = 104; $Scn1a^{+/-}$: 10.25 ± 0.69 nA, n = 102/N = 5

cultures). (**C**) Representative miniature EPSC (mEPSC) traces in neurons of either WT or

$Scn1a^{+/-}$ mice. (**D**) Average mEPSC amplitudes in neurons of either WT or $Scn1a^{+/-}$ mice

(WT: 39.47 ± 0.87 pA, n = 104; $Scn1a^{+/-}$: 38.15 ± 0.71 pA, n = 102/N = 5 cultures). (**E**)

Average mEPSC frequencies in neurons of either WT or $Scn1a^{+/-}$ mice (WT: 4.82 ± 0.52 Hz,

n = 104; $Scn1a^{+/-}$: 4.70 ± 0.56 Hz, n = 102/N = 5 cultures). Data were obtained from the same

autaptic neurons as in (**D**).

Figure 3. Changes in the inhibitory synaptic transmission of $Scn1a^{+/-}$ neurons.

(**A**) Representative traces of evoked inhibitory postsynaptic currents (IPSCs) recorded from a

single autaptic hippocampal neuron of either wild-type (WT) or $Scn1a^{+/-}$ mice.

Depolarization artifacts caused by the generated action currents have been removed for

clarity of presentation. (**B**) Average amplitudes of the evoked IPSCs in neurons of WT or

$Scn1a^{+/-}$ mice (WT: 7.94 ± 0.55 nA, n = 96; $Scn1a^{+/-}$: 6.18 ± 0.42 nA, n = 98/N = 5

cultures, *p < 0.05). (**C**) Representative miniature IPSC (mIPSC) traces in neurons of either

WT or $Scn1a^{+/-}$ mice. (**D**) Average mIPSC amplitudes in neurons of either WT or $Scn1a^{+/-}$

mice (WT: 26.94 ± 0.80 pA, n = 95; $Scn1a^{+/-}$: 26.70 ± 0.86 pA, n = 96/N = 5 cultures). (**E**)

Average mIPSC frequencies in neurons of either WT or $Scn1a^{+/-}$ mice (WT: 1.85 ± 0.25 Hz,

n = 95; $Scn1a^{+/-}$: 1.50 ± 0.21 Hz, n = 96/N = 5 cultures). Data were obtained from the same

autaptic neurons as in (**D**).

Figure 4. Changes in the number of inhibitory synapses in $Scn1a^{+/-}$ neurons.

(**A**) Representative traces of responses to 0.5 M hypertonic sucrose solution (10 s, black bar)

in an autapse neuron of either wild-type (WT) or $Scn1a^{+/-}$ mice. (**B**) The size of the readily

releasable pool (RRP) of autaptic neurons of either WT or $Scn1a^{+/-}$ mice (WT: 1.91 ± 0.12

nC, $n = 91$; $Scn1a^{+/-}$: 1.92 ± 0.12 nC, $n = 92/N = 5$ cultures). (**C**) Vesicular release probability

(P_{vr}) in single autaptic neurons of either WT or $Scn1a^{+/-}$ mice (WT: $29.99 \pm 2.31\%$, $n = 91$;

$Scn1a^{+/-}$: $24.01 \pm 1.66\%$, $n = 92/N = 5$ cultures, $*p < 0.05$). (**D**) Representative images of

autaptic neurons immunostained for the inhibitory nerve terminal marker vesicular GABA

transporter (VGAT; red) and the dendritic marker microtubule-associated protein 2 (MAP2;

green). Parts of the top row (scale bar 50 µm) are enlarged in the bottom row (scale bar 20

µm). (**E**) The number of VGAT–positive synaptic puncta in autaptic neurons of either WT or

$Scn1a^{+/-}$ mice (WT: 152.23 ± 7.91, $n = 70$, $Scn1a^{+/-}$: 117 ± 10.61, $n = 74/N = 8$ cultures, $*p <$

0.05).

Figure 5. Changes in extracellular Ca^{2+} concentration sensitivity of inhibitory synapses in

$Scn1a^{+/-}$ neurons.

(**A**) Inhibitory postsynaptic current (IPSC) amplitudes recorded in different extracellular

598 $[Ca^{2+}]_o$ (0.5, 1, 2, 4, 8 mM) with 1 mM Mg^{2+} held constant in wild-type (WT) and $Scn1a^{+/-}$

599 neurons (WT: $n = 35$, $Scn1a^{+/-}$: $n = 36/N = 9$ cultures). (**B**) Dose-response curves and Hill fits

600 for normalized IPSC amplitudes, recorded in different extracellular $[Ca^{2+}]_o$ (0.5, 1, 2, 4, 8

601 mM) with 1 mM Mg^{2+}, held constant in WT and $Scn1a^{+/-}$ neurons. IPSC amplitudes were

602 normalized to the IPSC amplitude at 8 mM $[Ca^{2+}]_o$. Original data were obtained from the

603 same cells as in (**A**). (**C**) Representative traces of evoked excitatory postsynaptic currents

604 (EPSCs) recorded at 2 mM $[Ca^{2+}]_o$ (black line) and 8 mM $[Ca^{2+}]_o$ (red line) from neurons of

605 either WT or $Scn1a^{+/-}$ mice. Depolarization artifacts caused by the generated action currents

606 have been removed for clarity of presentation. (**D**) Average magnification of evoked EPSCs

607 at 8 mM $[Ca^{2+}]_o$ based on EPSC amplitudes at 2 mM $[Ca^{2+}]_o$ in neurons of WT or $Scn1a^{+/-}$

608 mice (WT: 1.45 ± 0.07, $n = 62$, $Scn1a^{+/-}$: 1.48 ± 0.07, $n = 64/N = 11$ cultures). (**E**)

609 Representative traces of evoked IPSCs recorded at 2 mM $[Ca^{2+}]_o$ (black line) and 8 mM

610 $[Ca^{2+}]_o$ (red line) from neurons of either WT or $Scn1a^{+/-}$ mice. Depolarization artifacts caused

611 by the generated action currents have been removed for clarity of presentation. (**F**) Average

612 magnification of the evoked IPSCs at 8 mM $[Ca^{2+}]_o$ based on IPSC amplitudes at 2 mM

613 $[Ca^{2+}]_o$ in neurons of WT or $Scn1a^{+/-}$ mice (WT: 1.36 ± 0.09 mM, $n = 35$, $Scn1a^{+/-}$: $1.02 \pm$

614 0.04 mM, $n = 36/N = 9$ cultures, ***$p < 0.001$).

616 **Figure S1.** No changes in Na^+ current amplitudes in wild-type (WT) and $Scn1a^{+/-}$ neurons of

617   the hippocampus and striatum.

618   (**A**) Representative traces of Na$^+$ currents evoked by depolarizing pulses (–70 mV to 0 mV)

619   via patch pipette for 2 ms under voltage-clamp conditions. There were no differences in Na$^+$

620   currents between the WT and *Scn1a$^{+/-}$* neurons of the hippocampus and striatum. (**B**) Average

621   amplitudes of the Na$^+$ current in WT and *Scn1a$^{+/-}$* neurons of the hippocampus and striatum.

a

b

c