CRISPR interference identifies vulnerable cellular pathways with bactericidal

phenotypes in *Mycobacterium tuberculosis*.

Matthew B. McNeil[1,2#], Laura M Keighley[1], Josephine R. Cook[1], Chen-Yi Cheung[1] and

Gregory M. Cook[1,2]

1: Department of Microbiology and Immunology, University of Otago, New Zealand.

2: Maurice Wilkins Centre for Molecular Biodiscovery, University of Auckland, New

Zealand.

#Corresponding Author

Abstract

Mycobacterium tuberculosis remains a leading cause of death for which new drugs

are needed. The identification of drug targets has been advanced by high-throughput

and targeted genetic deletion strategies. Each though has limitations including the

inability to distinguish between levels of vulnerability, lethality and scalability as a

molecular tool. Using mycobacterial CRISPR interference in combination with

phenotypic screening we have overcome these individual issues to investigate

essentiality, vulnerability and lethality for 96 target genes from a diverse array of

cellular pathways, many of which are potential antibiotic targets. Essential genes

involved in cell wall synthesis and central cellular functions were equally vulnerable

and often had bactericidal consequences. Conversely, essential genes involved in

metabolism, oxidative phosphorylation or amino acid synthesis were less vulnerable

to inhibition and frequently bacteriostatic. In conclusion, this study provides novel

insights into mycobacterial genetics and biology that will help to prioritise potential

drug targets.

Introduction

Infections from *Mycobacterium tuberculosis* remain a leading cause of death[1]. *M. tuberculosis* is currently treated with a combination of four drugs for six months. However, the evolution and spread of drug resistance highlights a need for new drugs. Novel drugs with unique molecular targets that are (i) essential for growth, (ii) are highly vulnerable to inhibition (i.e. require low levels of inhibition to produce a bacterial phenotype)[2-4] and (iii) are lethal when inhibited are likely to have the greatest impact on reducing treatment times[5].

Transposon-saturation mutagenesis coupled to next-generation sequencing (i.e. Tn-seq) provides a high-throughput assessment of fitness costs associated with gene disruption. Tn-seq in *M. tuberculosis* has made significant contributions to the identification of essential genes [6-8]. Despite this, Tn-seq does not distinguish between bactericidal and bacteriostatic outcomes. Although improved statistical analysis has allowed for the identification of differences in genetic requirements between clinical isolates[9], there is also no standardized Tn-seq methodology to determine which genes are more vulnerable to inhibition[10]. Dual-control (DUC) switches that combine transcriptional repression and proteolysis of target genes overcomes issues associated with Tn-seq[3]. DUC switches can also partially knockdown target genes to generate hypomorphic strains for compound mechanism of action studies and for investigations into pathway vulnerability[2-4]. Despite this, the construction of DUC switches is resource intensive, prohibiting the parallel construction of DUC switches for multiple genes of interest. Notable exceptions include the large collaborative efforts of the PROSPECT screen[11].

Mycobacterial CRISPR interference (CRISPRi) is an alternative genetic platform that transcriptionally represses target gene expression[12-15]. CRISPRi plasmids are easy to

construct as (i) they only require the cloning of only a 20 bp segment of

complementarity to target genes (i.e. sgRNA sequences) and (ii) require only a single

transformation as CRISPRi plasmids also express the necessary catalytically dead

Cas9 (i.e. dCas9) [13,15]. We and others have shown that CRISPRi can (i) probe gene

essentiality, (ii) provide information on the lethality and (iii) can generate partial

knockdown strains[12-16]. Using mycobacterial CRISPRi, this current study has

phenotypically screened target genes from a diverse spectrum of cellular pathways in

M. tuberculosis to investigate variations in gene essentiality, vulnerability and lethality.

These results demonstrate that the majority of screened genes involved in cell wall

synthesis and core cellular functions (e.g. transcription and translation) are essential,

equally vulnerable to inhibition and bactericidal when inhibited. Conversely, the

majority of essential genes involved in metabolism or amino acid synthesis were

subject to buffering effects, requiring higher levels of repression to inhibit growth.

Furthermore, the inhibition of several genes previously defined as non-essential

inhibited bacterial growth suggesting a delay between target inhibition and the

necessary metabolic remodelling to facilitate growth. In conclusion, this study will help

to prioritise potential drug targets as well as reiterate the utility of CRISPRi in

investigating mycobacterial genetics and physiology.

Results

Mycobacterial CRISPRi to assess genetic essentially: Using CRISPRi we

investigated the phenotypes (i.e. growth and viability) of 96 genes (encoding 94

proteins and 2 ribosomal RNAs) from a diverse array of biological processes, many of

which are considered potential drug targets. Mycobacterial CRISPRi uses a type I

dCas9 from *Strepoccous thermophilus* that recognizes 15 permissible non-canonical

PAM sequences to repress target gene expression[15]. Both sgRNA and dCas9 are

expressed in response to the addition of anhydrotetracycline (ATc). There is a linear

relationship between the level of ATc added and the level of transcriptional repression

[13,16]. Two sgRNA sequences per target gene were selected and cloned (Table S1).

Some genes have only a single sgRNA due to cloning difficulties. Bacterial phenotypes

following target gene repression were screened in 96 well plates using an ATc gradient

(Figure 1). OD_{600} was determined 7-days post the addition of ATc and genes were

defined as either essential (i.e. <25% growth relative to no ATc control), growth

impairing (i.e. <50% growth relative to no ATc control) or non-essential (i.e. >50%

growth relative to no ATc control). Essentiality calls were compared with published Tn-

seq experiments as available in the mycobrowser database[6]. CFUs were determined

5-days post the addition of ATc as previous work demonstrated that this allows for the

detection of bacterial killing prior to the emergence of non-responsive CRISPRi

mutants[13,16]. Essential or growth impairing sgRNAs that produced ≥1 log_{10} reduction

in CFU/ml were defined as bactericidal, whilst <1 log_{10} reduction was

bacteriostatic[13,16].

Cell wall synthesizing genes are essential and bactericidal when inhibited: The

majority of sgRNAs targeting genes involved in cell wall synthesis in our screen were

essential (Figure 2A). Essentiality and non-essentiality (e.g. *rv3032* and *treS*) calls are

Figure 1: Experimental workflow for CRISPRi assessment of bacterial phenotypes. Cultures of *M. tuberculosis* mc^{2}6230 expressing CRISPRi plasmids that express a single sgRNA were maintained in 10 ml cultures. Cultures were diluted to an OD$_{600}$ of 0.01 in a deep well 96 well plate and then inoculated into 96 well assay plates that contain a dilution gradient of ATc to induce expression of both the dCas9 and sgRNA at a starting OD$_{600}$ of 0.005. Assays plates used to determine if sgRNAs inhibited bacterial growth were assessed after 7 days by determining the OD$_{600}$ relative to a no ATc control. Plates used to determine bacterial viability were assessed after 5 days by spotting for CFUs. Previous work has demonstrated that day 5 allows for the detection of bacterial killing prior to the emergence of non-responsive CRISPRi mutants[13,16]. Further details for experimental workflow are described in the materials and methods

Figure 2: Bacterial phenotypes following the transcriptional inhibition of genes involved in cell wall synthesis or core cellular processes. (A and B) Each sgRNA is designated by the target gene name and whether it is the first or second designed sequence (i.e. a or b). Gray shaded heatmap describes the results from previous TnSeq experiments[6] as presented in the mycobrowser database (https://mycobrowser.epfl.ch/) and whether it is essential (E), non-essential (NE) or whether it has a growth defect (D). X denotes genes for which there is no essentiality call. Pink shaded heatmap describes the predicted strength of PAM scores ranked between 1-15 based on previous studies by Rock et al[15]. Blue shaded heatmap describes the level of growth inhibition at increasing ATc concentrations relative to a no ATc control well. The level of inhibition is the mean of at least four replicate experiments. Orange shaded heatmap describes the log_{10} reduction in CFU/ml for each sgRNA between day 0 and day 5 at increasing concentrations of ATc. The log_{10} reduction in CFU/ml of inhibition is the mean of at least four replicate experiments.

consistent with previous experiments (Figure 2A)[6,17,18]. All genes involved in mycolic

acid synthesis (i.e. *kasA*, *inhA*, *hadA/B, pkS13, fadD32, mmpL3 and pptT*) were

essential and bactericidal when inhibited (Figure 2A). The majority of mycolic acid

synthesis genes caused a >3 log reduction in CFU/ml at the maximum ATc

concentration (i.e. 300 ng/ml) (Figure 2A). Genes involved in arabinogalactan

synthesis (i.e. *wecA*, *embA*, *dprE1* and *glf*) had either an essential or growth impairing

phenotype (Figure 2A). Inhibition of *wecA*, *embA* and *dprE1* had a weak bactericidal

phenotype (i.e. 1 log reduction in CFU/ml) at ATc concentrations >100 ng/ml (Figure

2A). Inhibition of *glf* had a stronger bactericidal phenotype (i.e. >2 log reduction in

CFU/ml at lower ATc concentrations) (Figure 2A). All genes in peptidoglycan synthesis

(i.e. *murE*, *murX*, *murI*, *alr*, *ddlA*) were essential for growth (Figure 2A). The inhibition

of *murE*, *murX* and *ddlA* had a bactericidal phenotype, whilst *murI* and *alr* were

bacteriostatic (Figure 2A). With the exception of *glgE* that had an essential-

bacteriostatic phenotype, all genes involved in glucan biosynthesis (i.e. *rv3032* and

treS) were individually non-essential (Figure 2A). The predicted efflux pump, *efpA*, had

an essential-bactericidal phenotype, whilst inhibition of the serine/threonine protein

kinase *pknA* had a growth impairing phenotype (Figure 2A). In conclusion, genes

involved in cell wall synthesis are predominately essential for growth, although there

is variation in whether their inhibition is bactericidal.

Core cellular processes are essential and have lethal phenotypes when

inhibited: The majority of sgRNAs targeting genes involved in core cellular processes

(i.e. transcription, translation, nucleotide metabolism, protein homeostasis and cell

division) were essential and consistent with previous Tn-seq experiments[6] (Figure 2B).

With the exception of *adoK*, genes involved in nucleotide biosynthesis (i.e. *pyrG* and

guaB2) were essential-bactericidal (Figure 2B). Both *gyrA* and *gyrB* were individually

essential, yet only the inhibition of *gyrB* had a bactericidal phenotype (Figure 2B).

Inhibition of DNA polymerase (i.e. *dnaN*) and RNA polymerase (i.e. *rpoB*) were

essential and bactericidal, with the phenotypically strongest sgRNAs having a >2 log

reduction in CFU/ml. Inhibition of the transcriptional terminator, *rho*, inhibited bacterial

growth but only at the highest ATc concentration (Figure 2B). Ribosomal RNAs (i.e.

16S-*rrs* and 23S-*rrl*) were essential for bacterial growth, but only the inhibition of *rrl*

resulted in reduced bacterial viability. The ribosomal protein *rpsA* was also essential

for growth, with the phenotypically strongest sgRNA reducing bacterial viability.

Similarly the transcriptional inhibition of genes encoding tRNA synthetase (i.e. *aspS*

and *leuS*), genes involved in protein homeostasis (i.e. *groES, clpC1, clpP1* and *clpP2*)

and cell division (i.e. *ftsZ*) were essential and had strong bactericidal phenotypes

(Figure 2B). The peptide deformylase encoded by *def* that was previously defined as

essential failed to inhibit bacterial growth following transcriptional repression (Figure

2B). In conclusion, genes involved in central cellular processes are essential for the

growth of *M. tuberculosis*, and frequently have a bactericidal phenotype.

Genes involved in metabolic process and oxidative phosphorylation are

generally bacteriostatic: Transcriptional inhibition of genes involved metabolic

processes or oxidative phosphorylation that had previously been defined as essential

resulted in growth inhibition (i.e. *fba*, *glcB*, *folP1*, *folC*, *atpE* and *atpB*) (Figure 3A and

B). However, only the inhibition of *glcB* and *atpE/B* significantly reduced bacterial

viability (Figure 3A and B). Some sgRNAs targeting genes previously defined as

essential failed to inhibit bacterial growth (e.g. *fum, menA, menD* and *birA*), whilst only

the strongest sgRNA targeting *trxB2* was able impair bacterial growth (Figure 3A and

B). Furthermore, some sgRNA targeting genes previously defined as non-essential

inhibited bacterial growth in our study (i.e. *eno*, *dlaT*, *mqo*, *mdh*, *icl1*, *pckA*, *ndh*, *qcrB*,

182
183
Figure 3: Bacterial phenotypes following the transcriptional inhibition of genes
involved in metabolic processes, oxidative phosphorylation or amino acid
synthesis. (A and B) Each sgRNA is designated by the target gene name and
whether it is the first or second designed sequence (i.e. a or b). Gray shaded heatmap
describes the results from previous TnSeq experiments[6] as presented in the
mycobrowser database (https://mycobrowser.epfl.ch/) and whether it is essential (E),
non-essential (NE) or whether it has a growth defect (D). Pink shaded heatmap
describes the predicted strength of PAM scores ranked between 1-15 based on
previous studies by Rock et al[15]. Blue shaded heatmap describes the level of growth
inhibition at increasing ATc concentrations relative to a no ATc control well. The level
of inhibition is the mean of at least four replicate experiments. Orange shaded
heatmap describes the log_{10} reduction in CFU/ml for each sgRNA between day 0 and
day 5 at increasing concentrations of ATc. The log_{10} reduction in CFU/ml of inhibition
is the mean of at least four replicate experiments.
198

ctaC and *ethA*). Whilst the majority were bacteriostatic the strongest *ndh* sgRNA was

bactericidal (Figure 3A and B). In conclusion, the transcriptional inhibition of many

genes involved in metabolic processes or oxidative phosphorylation inhibit or impair

bacterial growth, but have a largely bacteriostatic phenotype.

Genes involved in amino acid synthesis are essential but generally

bacteriostatic: Transcriptional inhibition of genes involved in amino acid biosynthesis

generally inhibited bacterial growth (i.e. *hisG*, *aroK*, *trpD/A/B*, *ilvB1/D*, *thrA*, *metX*,

lysA). These essential phenotypes are consistent with previous Tn-seq experiments

(Figure 3C)[6]. The majority were bacteriostatic with only *aroK* being bactericidal.

Although the sgRNA *metX*_a reduced bacterial viability at 33 ng/ml ATc (i.e. $1.02 \pm$

$0.92 \log_{10}$ reduction), this phenotype was not observed across other concentrations or

sgRNAs (Figure 3C). Only a single *argJ* sgRNA was able to inhibit bacterial growth at

the highest ATc concentration, whilst sgRNAs targeting *serB2* and *glnA* failed to inhibit

bacterial growth (Figure 3C)[6]. Non-essentiality calls for *ald* and *lysE* are consistent

with previous experiments (Figure 3C)[6]. In conclusion, the transcriptional inhibition of

genes involved in amino acid synthesis generally inhibit bacterial growth, but have a

bacteriostatic phenotype.

Comparison of sgRNA activity identifies pathways of altered vulnerability: To

identify cellular pathways more vulnerable to inhibition (i.e. requiring less ATc to inhibit

bacterial growth) all sgRNAs were grouped according to biological function and the

ATc MIC of each active sgRNA (i.e. that inhibited bacterial growth) was compared. To

account for potential variation between experiments, the ATc MIC of sgRNAs was

compared to the ATc MIC of the *mmpL3*_a sgRNA that was used as a positive

knockdown control in all experiments (Figure 4B). The majority of sgRNAs targeting

cell wall synthesis or core cellular process were as active as *mmpL3*_a (i.e. within a

log2 change) (Figure 4A). Several sgRNAs targeting cell wall synthesis, core process

and metabolic genes were more active (i.e. >-1 log2 change), representing pathways

of potentially increased vulnerability (Figure 4A-B, red symbols). Interestingly, *eno*,

which converts 2-phosphoglycerate to phosphoenolpyruvate as part of glycolysis, was

the most vulnerable of all assessed sgRNAs (Figure 4A-B). A significant proportion of

sgRNAs targeting metabolism, oxidative phosphorylation and amino acid synthesis

genes were less active than *mmpL3*_a (i.e. >1 log2 change), representing biological

pathways of potentially reduced vulnerability (Figure 4A-B, blue symbols). Dose

response curves for five sgRNAs of either increased or reduced vulnerability highlight

this shift in ATc MIC (Figure 4B).

235
236
Figure 4: Comparison of target vulnerability across cellular pathways. (A) Plotted
data is the average log2 fold change in ATc MIC compared to the *mmpL3*_a sgRNA
for each sgRNA that results in at least a growth impairment (i.e. <50% relative to no
ATc control) at the maximum ATc concentration. Data is the average fold change from
at least two experiments that individually included biological duplicates (i.e. at least 4
replicates). sgRNAs are grouped according to their predicted or validated cellular
function. Average sgRNA fold change values are presented in table S1. The average
log2 fold change for a cellular pathway is represented by a solid black line. (B) Dose
response curves for selected sgRNAs. Red and blue dots in A correspond to red and
blue lines in B. Dose response curves are the average of at least 4 replicate
experiments.

Discussion

There remains an urgent global need for novel therapeutic agents to treat both drug susceptible and resistant strains of *M. tuberculosis*. Biological pathways that are essential, highly vulnerable to inhibition and have a bactericidal outcome are likely to have the greatest clinical impact. High-throughput and targeted genetic deletion strategies have been useful in predicting gene essentiality. However, each strategy is subject to individual limitations that include an inability to determine target vulnerability, differentiation between lethal and static outcomes or issues with scalability. By coupling CRISPRi to assessments of growth and viability this study overcomes these individual limitations to identify variations in target vulnerability and consequences on cellular viability for a diverse array of biological pathways.

There was a strong correlation between genetic essentiality and CRISPRi mediated growth inhibition when targeting genes involved in cell wall synthesis or core cellular processes. The majority of cell wall synthesis or core cellular encoding genes were equally vulnerable to transcriptional inhibition and bactericidal. Genes involved in mycolic acid synthesis, DNA metabolism, transcription, translation, tRNA synthesis, protein homeostasis or cell division had the strongest bactericidal phenotype (i.e. both level of reduction and levels of ATc needed to induce killing). Inhibition of peptidoglycan and arabinogalactan biosynthesis had differing effects on cell viability depending on the gene being targeted, with bactericidal phenotypes generally being weaker. Due to the polar effects of CRISPRi[15,16] the repression of *murE* and *murX* is likely to repress multiple downstream genes within the *dcw* operon (including, *murD*, *murF*, *murG*, *ftsW* and *ftsQ*)[19]. Consequently, bactericidal phenotypes for *murE* and *murX* may not be reflective of single target gene repression. Similarly, the polar effects of CRISPRi suggest that transcriptional inhibition of both gyrase subunits, as would

be facilitated by the *gyrB* sgRNA on the *gyrBA* operon but not *gyrA* sgRNA, is needed for lethality. Furthermore, both sgRNAs targeting *rho* have low PAM scores (i.e. 9 and 11), which may explain the bacteriostatic phenotype compared to previously published bactericidal *rho*-DUC mutants[20]. Combined, the results of this study support the continued development of novel drugs that target proteins involved in cell wall synthesis and core cellular processes.

Genes involved in metabolic processes, oxidative phosphorylation or amino acid synthesis, were on average less vulnerable to transcriptional inhibition. Genes that inhibited bacterial growth were generally bacteriostatic, with the bactericidal exceptions of *glcB* (i.e. malate synthase)[21], *atpE/B* (i.e. ATP synthase operon)[16], *aroK* (i.e. shikimate biosynthesis) and *ndh* (i.e. NDH2 dehydrogenase). Recent studies have demonstrated that the consequences of inhibiting individual metabolic genes when using CRISPRi can be mitigated by metabolic buffering, to limit global effects on bacterial metabolism[22]. Consequently, several cell divisions may be needed to deplete enzyme products below critical levels. Previous studies in *E. coli* demonstrated that metabolic buffering resulted in time dependent delays in inhibition of bacterial growth, with only a small proportion of metabolic genes having rapid inhibitory phenotypes[22]. We hypothesize that metabolic buffering in *M. tuberculosis* may explain why many predicted essential metabolic, oxidative phosphorylation or amino acid synthesis genes fail to inhibit bacterial growth or have a significantly reduced vulnerability when targeted by CRISPRi. Metabolic buffering may also explain why *metX*, *thrA* and *ilvB1* were bacteriostatic, yet in previous studies deletion mutants had bactericidal phenotypes [2,23,24]. Consistent with this, previous observations of *metX* and *thrA* lethality are observed post 5-days, whilst our assay assessed viability at 5-days [2,23].

Media choice also plays an important role in the observation of phenotypes associated

with metabolic mutants [21,25,26]. Consequently, the use of OADC (i.e. glucose and oleic

acid as carbon sources) and exogenous pantothenic acid in our experiments to

supplement *M. tuberculosis* mc^{2}6230 (Δ*panCD*) may alter or mitigate the phenotypes

of some metabolic, oxidative phosphorylation or amino acid transcriptional

knockdowns. Furthermore, the lack of phenotype for sgRNAs targeting *birA* is likely

due to the presence of biotin in 7H9 media.

In our study, numerous previously defined non-essential genes had significant impacts

on bacterial growth. In contrast to the metabolic buffering that may mitigated the

phenotypes of essential genes, we hypothesize that a lack of metabolic buffering in

some cases may delay the necessary metabolic remodelling that is required to

facilitate growth in the absence of these non-essential genes. Consequently,

previously constructed knockouts or Tn-seq mutants of of *icl1*[27], *qcrB*[25], *ctaC*[25], *dlaT*[28]

and *ndh*[25] may reflect an adapted metabolic state that in our study are defined as

being essential for bacterial growth.

We acknowledge that variations in genetic vulnerability may be a result of variation in

CRISPRi transcriptional repression. It is also possible that differences in bacterial

phenotypes between sgRNAs (e.g. one static and one cidal) may be due to differences

in sgRNA efficacy. Previous studies have highlighted the influence of PAM sequence,

location of sgRNA within the genetic target and sgRNA sequence on dCas9

Streptococcus pyogenes activity[29-31]. Whilst we selected the strongest predicted PAM

sequence and sgRNA location does not influence activity of the dCas9$_{Sth1}$

orthologue[15], more work is required to determine the influence of sgRNA sequence on

dCas9$_{Sth1}$ activity. Off-target effects are also unlikely to be a confounding factor, as

previous work as suggested limited off-targeting for the dCas9$_{Sth1}$ orthologue[14].

Despite this, we speculate that variable levels of transcriptional repression are not a

confounding factor on our overall conclusions as CRISPRi studies in *E. coli* and *B.*

subtilis have demonstrated that genes involved in peptidoglycan synthesis and

translation are more vulnerable to inhibition than metabolic genes[31].

There is good correlation between the phenotypes of transcriptional inhibition and

many known antibiotics. For example, the bactericidal phenotype of isoniazid, a

cornerstone of current treatment regimens, correlates well with transcriptional

inhibition of its target *inhA*[32]. Interestingly, transcriptional inhibition of *rpoB* produced

a strong bactericidal phenotype, whilst rifampicin, a known inhibitor of RpoB, only

achieves bacterial killing when in significant excess of the MIC[32]. Whilst it is possible

that there are differences between the consequences of transcriptional and chemical

inhibition, these results suggest that rifampicin may be a poor inhibitor of RNA

polymerase *in vitro*. Combined, the results of our current study have provided novel

insights into the vulnerability and lethality of inhibiting different biological processes in

M. tuberculosis. These results will be a valuable resource for the mycobacterial

research community that will help to advance both basic biology and the advancement

of novel drug targets.

Methods

Bacterial strains and growth conditions: *Escherichia coli* MC1061 was grown in or on luria broth (LB) media or agar (1.5%) at 37°C and shaking at 200 rpm when required. *M. tuberculosis* strain mc²6230 (ΔpanCD, ΔRD1) [33-36], was grown and maintained in 7H9 liquid media or on 7H11 solid media supplemented with OADC (0.005% oleic acid, 0.5% bovine serum albumin, 0.2% dextrose, 0.085% catalase) and pantothenic acid (25 µg/ml) and incubated at 37°C. Liquid cultures were supplemented with 0.05% tyloxapol (Sigma) and grown with shaking at 140 rpm. *M. tuberculosis* strain mc²6230 is a BSL2 avirulent auxotroph that has been approved for use under BSL2 containment at the University of Otago. When necessary media was supplemented with Kanamycin at 50 µg/ml for *E. coil* and 25 µg/ml for *M. tuberculosis.* Anhydrotetracycline (ATc, Sigma) was solubilized in 70% ethanol and added to experiments at the stated concentrations.

Construction and transformation of CRISPRi plasmids: A 20-25 bp sequence downstream of permissible PAM sequences targeting the non-template strand of target genes of interest were identified[15]. For each target gene, the two sgRNAs based on PAM score were manually selected. Target sequences were ordered as oligos with GGGA and AAAC overhangs respectively (Table S1), and cloned into pJLR965 using BsmB1 and golden gate cloning as previously described[13]. Plasmids were cloned into *E. coli* and validated with sanger sequencing. Confirmed CRISPRi plasmids were electroporated into *M. tuberculosis* strain mc²6230 following previously established protocols[13,16].

CRISPRi phenotypic assessment of essentiality and viability: To determine the consequences of targeted gene repression on bacterial growth, phenotypic assays were performed as follows (Figure 1). *M. tuberculosis* mc²6230 strains containing

CRISPRi plasmids were grown and maintained in 7H9-supplemented media with KAN. Cultures were diluted initially to an OD_{600} of 0.1 in 7H9-supplemented media (+KAN) in a deep well 96 well plate. Cultures were diluted again to a final OD_{600} of 0.01 in a deep well 96 well plate. 96 well assay plates were prepared with a 3-fold dilution of ATc along the X-axis starting at 300 ng/ml of ATc in column 2 with a starting inoculum of OD_{600} 0.005 (Figure 1). This was achieved by adding 50 µl of 7H9-supplemented media (+KAN) to all wells of columns 3-11 except row A-H. 75 µl of 7H9-supplemented media (+KAN) containing the starting concentration of ATc (i.e. 600 ng/ml ATc) was added to column 2 except row A-H. ATc was diluted along the horizontal axis of the 96 well plate, transferring 25 µl between columns, down to column 10. Column 11 was used as a no ATc control. Columns 1 and 12 and rows A-H contained 100 µl of 7H9-supplemented media (+KAN) as contamination and background controls. Fifty µl of culture adjusted to an OD_{600} 0.01 was added to wells 2-11 of individual rows of a 96 well plate to achieve a starting OD_{600} of 0.005. Each row represents the ATc dilution gradient for a single *M. tuberculosis* mc^{2}6230 strain expressing a unique CRISPRi plasmids. All experiments included a non-targeting sgRNA (i.e. pJR965) and a *mmpL3_*a targeting sgRNA as a negative and positive essential-bactericidal control[13].

To assess gene essentially, duplicate plates were grown at 37°C without shaking for 7-days. OD_{600} was measured using a Varioskan-LUX microplate reader. The minimal inhibitory concentration (MIC) of ATc was determined using OD_{600} reads from duplicate plates relative to the growth of the no-ATc control, using a non-linear fitting of data to the Gompertz equation[37]. Targeted genes were defined as either essential (i.e. <25% growth relative to no ATc control), growth impairing (i.e. <50% growth relative to no ATc control) or non-essential (i.e. >50% growth relative to no ATc control). In this study, the vulnerability of genes to transcriptional inhibition is defined

by the ATc MIC, i.e. the lower the MIC the more vulnerable that gene is to CRISPRi

mediated transcriptional inhibition.

Duplicate 96 well assay plates were also set up to determine the consequences of

targeted gene repression on bacterial viability by taking colony forming units on day 0

and day 5. Previous work has demonstrated that day 5 allows for the detection of

bacterial killing prior to the emergence of non-responsive CRISPRi mutants[13,16].

Briefly, at day 0 hrs a 4-point ten-fold dilution of the 0.01 diluted culture was performed

in 7H9 base media (i.e. not supplemented), with 5 µl of each dilution spotted onto to

7H11-supplemented (+KAN) agar plates. At Day 5, approximately 100 µl of culture

was removed from columns 2-5 of each row and transferred to a new 96 well plate to

be diluted. A 4-point ten-fold dilution gradient was performed in 7H9 base media (i.e.

not supplemented), and 5 µl of each dilution was spotted onto to 7H11-supplemented

(+KAN) agar plates. Plates were incubated at 37°C for 4-5 weeks, at which point

colonies were counted. Essential genes with bacteriostatic consequences resulted in

no change in CFU/ml relative to 0 hrs, whilst bactericidal consequences produced at

least a 1 log reduction in CFU/ml relative to the day 0 inoculum.

408 **Acknowledgments**

409 This research was financially supported by the Maurice Wilkins Centre for Molecular

410 Biodiscovery, the Marsden Fund (Royal Society of New Zealand) (grant number

411 UOO1807). and the Health Research Council of New Zealand (grant number 20/459).

413 We have no conflicts of interest to declare.

References

WHO. Global Tuberculosis Report 2018. (2018).

Hasenoehrl, E. J., Rae Sajorda, D., Berney-Meyer, L., Johnson, S., Tufariello, J. M., Fuhrer, T., Cook, G. M., Jacobs, W. R., Jr. & Berney, M. Derailing the aspartate pathway of *Mycobacterium tuberculosis* to eradicate persistent infection. *Nat Commun* 10, 4215, (2019).

Kim, J.-H., O'Brien, K. M., Sharma, R., Boshoff, H. I. M., Rehren, G., Chakraborty, S., Wallach, J. B., Monteleone, M., Wilson, D. J., Aldrich, C. C., Barry, C. E., Rhee, K. Y., Ehrt, S. & Schnappinger, D. A genetic strategy to identify targets for the development of drugs that prevent bacterial persistence. *Proceedings of the National Academy of Sciences of the United States of America* 110, 19095-19100, (2013).

Li, W., Obregon-Henao, A., Wallach, J. B., North, E. J., Lee, R. E., Gonzalez-Juarrero, M., Schnappinger, D. & Jackson, M. Therapeutic potential of the *Mycobacterium tuberculosis* mycolic acid transporter, MmpL3. *Antimicrobial agents and chemotherapy* 60, 5198-5207, (2016).

Wei, J. R., Krishnamoorthy, V., Murphy, K., Kim, J. H., Schnappinger, D., Alber, T., Sassetti, C. M., Rhee, K. Y. & Rubin, E. J. Depletion of antibiotic targets has widely varying effects on growth. *Proceedings of the National Academy of Sciences* 108, 4176-4181, (2011).

DeJesus, M. A., Gerrick, E. R., Xu, W., Park, S. W., Long, J. E., Boutte, C. C., Rubin, E. J., Schnappinger, D., Ehrt, S., Fortune, S. M., Sassetti, C. M. & Ioerger, R. Comprehensive Essentiality Analysis of the Mycobacterium tuberculosis Genome via Saturating Transposon Mutagenesis. *mBio* 8, 1-17, (2017).

Sassetti, C. M., Boyd, D. H. & Rubin, E. J. Genes required for mycobacterial growth defined by high density mutagenesis. *Molecular Microbiology* 48, 77-84, (2003).

Sassetti, C. M., Boyd, D. H. & Rubin, E. J. Comprehensive identification of conditionally essential genes in mycobacteria. *Proceedings of the National Academy of Sciences of the United States of America* 98, 12712-12717, (2001).

Carey, A. F., Rock, J. M., Krieger, I. V., Chase, M. R., Fernandez-Suarez, M., Gagneux, S., Sacchettini, J. C., Ioerger, T. R. & Fortune, S. M. TnSeq of *Mycobacterium tuberculosis* clinical isolates reveals strain-specific antibiotic liabilities. *PLOS Pathogens* 14, e1006939, (2018).

Rock, J. Tuberculosis drug discovery in the CRISPR era. *PLOS Pathogens* 15, (2019).

Johnson, E. O., LaVerriere, E., Office, E., Stanley, M., Meyer, E., Kawate, T., Gomez, J. E., Audette, R. E., Bandyopadhyay, N., Betancourt, N., Delano, K., Da Silva, I., Davis, J., Gallo, C., Gardner, M., Golas, A. J., Guinn, K. M., Kennedy, S., Korn, R., McConnell, J. A., Moss, C. E., Murphy, K. C., Nietupski, R. M., Papavinasasundaram, K. G., Pinkham, J. T., Pino, P. A., Proulx, M. K., Ruecker, N., Song, N., Thompson, M., Trujillo, C., Wakabayashi, S., Wallach, J. B., Watson, C., Ioerger, T. R., Lander, E. S., Hubbard, B. K., Serrano-Wu, M. H., Ehrt, S., Fitzgerald, M., Rubin, E. J., Sassetti, C. M., Schnappinger, D. & Hung, D. T. Large-scale chemical-genetics yields new *M. tuberculosis* inhibitor classes. *Nature* 571, 72-78, (2019).

de Wet, T. J., Winkler, K. R., Mhlanga, M., Mizrahi, V. & Warner, D. F. Arrayed CRISPRi and quantitative imaging describe the morphotypic landscape of essential mycobacterial genes. *Elife* 9, (2020).

McNeil, M. & Cook, G. Utilization of CRISPR interference to validate MmpL3 as a drug target in *Mycobacterium tuberculosis. Antimicrob Agents Chemother* 63, (2019).

de Wet, T. J., Gobe, I., Mhlanga, M. M. & Warner, D. F. CRISPRi-Seq for the Identification and Characterisation of Essential Mycobacterial Genes and Transcriptional Units. *bioRvix*, (2018).

Rock, J. M., Hopkins, F. F., Chavez, A., Diallo, M., Chase, M. R., Gerrick, E. R., Pritchard, J. R., Church, G. M., Rubin, E. J., Sassetti, C. M., Schnappinger, D. & Fortune, S. M. Programmable transcriptional repression in mycobacteria using an orthogonal CRISPR interference platform. *Nature Microbiology* 2, 16274, (2017).

McNeil, M., Ryburn, H., Harold, L. K., Tirados, J. & Cook, G. Transcriptional Inhibition of the F1F0-Type ATP Synthase Has Bactericidal Consequences on the Viability of Mycobacteria. *Antimicrobial Agents and Chemotherapy* In press, (2020).

Stadthagen, G., Sambou, T., Guerin, M., Barilone, N., Boudou, F., Kordulakova, J., Charles, P., Alzari, P. M., Lemassu, A., Daffe, M., Puzo, G., Gicquel, B., Riviere, M. & Jackson, M. Genetic basis for the biosynthesis of methylglucose lipopolysaccharides in *Mycobacterium tuberculosis. J Biol Chem* 282, 27270-27276, (2007).

Lee, J. J., Lee, S. K., Song, N., Nathan, T. O., Swarts, B. M., Eum, S. Y., Ehrt, S., Cho, S. N. & Eoh, H. Transient drug-tolerance and permanent drug-resistance rely on the trehalose-catalytic shift in Mycobacterium tuberculosis. *Nat Commun* 10, 2928, (2019).

Munshi, T., Gupta, A., Evangelopoulos, D., Guzman, J. D., Gibbons, S., Keep, N. H. & Bhakta, S. Characterisation of ATP-dependent Mur ligases involved in the biogenesis of cell wall peptidoglycan in *Mycobacterium tuberculosis. PLoS One* 8, e60143, (2013).

Botella, L., Vaubourgeix, J., Livny, J. & Schnappinger, D. Depleting *Mycobacterium tuberculosis* of the transcription termination factor Rho causes pervasive transcription and rapid death. *Nature Communications* 8, (2017).

Puckett, S., Trujillo, C., Wang, Z., Eoh, H., Ioerger, T. R., Krieger, I., Sacchettini, J., Schnappinger, D., Rhee, K. Y. & Ehrt, S. Glyoxylate detoxification is an essential function of malate synthase required for carbon assimilation inMycobacterium tuberculosis. *Proceedings of the National Academy of Sciences* 114, E2225-E2232, (2017).

Donati, S., Kuntz, M., Pahl, V., Farke, N., Beuter, D., Glatter, T., Gomes-Filho, J. V., Randau, L., Wang, C. Y. & Link, H. Multi-omics Analysis of CRISPRi-Knockdowns Identifies Mechanisms that Buffer Decreases of Enzymes in E. coli Metabolism. *Cell Syst*, (2020).

Berney, M., Berney-Meyer, L., Wong, K.-W., Chen, B., Chen, M., Kim, J., Wang, J., Harris, D., Parkhill, J., Chan, J., Wang, F. & Jacobs, W. R. Essential roles of methionine and S-adenosylmethionine in the autarkic lifestyle of *Mycobacterium tuberculosis. Proceedings of the National Academy of Sciences*, 201513033, (2015).

Awasthy, D., Gaonkar, S., Shandil, R. K., Yadav, R., Bharath, S., Marcel, N., Subbulakshmi, V. & Sharma, U. Inactivation of the *ilvB1* gene in

Mycobacterium tuberculosis leads to branched-chain amino acid auxotrophy and attenuation of virulence in mice. *Microbiology (Reading)* 155, 2978-2987, (2009).

Beites, T., O'Brien, K., Tiwari, D., Engelhart, C. A., Walters, S., Andrews, J., Yang, H. J., Sutphen, M. L., Weiner, D. M., Dayao, E. K., Zimmerman, M., Prideaux, B., Desai, P. V., Masquelin, T., Via, L. E., Dartois, V., Boshoff, H. I., Barry, C. E., 3rd, Ehrt, S. & Schnappinger, D. Plasticity of the *Mycobacterium tuberculosis* respiratory chain and its impact on tuberculosis drug development. *Nat Commun* 10, 4970, (2019).

Ruecker, N., Jansen, R., Trujillo, C., Puckett, S., Jayachandran, P., Piroli, G. G., Frizzell, N., Molina, H., Rhee, K. Y. & Ehrt, S. Fumarase Deficiency Causes Protein and Metabolite Succination and Intoxicates Mycobacterium tuberculosis. *Cell Chem Biol* 24, 306-315, (2017).

Munoz-Elias, E. J. & McKinney, J. D. *Mycobacterium tuberculosis* isocitrate lyases 1 and 2 are jointly required for in vivo growth and virulence. *Nat Med* 11, 638-644, (2005).

Shi, S. & Ehrt, S. Dihydrolipoamide acyltransferase is critical for *Mycobacterium tuberculosis* pathogenesis. *Infect Immun* 74, 56-63, (2006).

Cui, L., Vigouroux, A., Rousset, F., Varet, H., Khanna, V. & Bikard, D. A CRISPRi screen in E. coli reveals sequence-specific toxicity of dCas9. *Nat Commun* 9, 1912, (2018).

Gasiunas, G., Young, J. K., Karvelis, T., Kazlauskas, D., Urbaitis, T., Jasnauskaite, M., Grusyte, M. M., Paulraj, S., Wang, P. H., Hou, Z., Dooley, S. K., Cigan, M., Alarcon, C., Chilcoat, N. D., Bigelyte, G., Curcuru, J. L., Mabuchi, M., Sun, Z., Fuchs, R. T., Schildkraut, E., Weigele, P. R., Jack, W. E., Robb, G. B., Venclovas, C. & Siksnys, V. A catalogue of biochemically diverse CRISPR-Cas9 orthologs. *Nat Commun* 11, 5512, (2020).

Hawkins, J. S., Silvis, M. R., Koo, B.-M., Peters, J. M., Osadnik, H., Jost, M., Hearne, C. C., Weissman, J. S., Todor, H. & Gross, C. A. Mismatch-CRISPRi Reveals the Co-varying Expression-Fitness Relationships of Essential Genes in Escherichia coli and Bacillus subtilis. *Cell Systems* 11, 523-535, (2020).

de Steenwinkel, J. E. M., de Knegt, G. J., ten Kate, M. T., van Belkum, A., Verbrugh, H. A., Kremer, K., van Soolingen, D. & Bakker-Woudenberg, I. A. J. M. Time-kill kinetics of anti-tuberculosis drugs, and emergence of resistance, in relation to metabolic activity of *Mycobacterium tuberculosis*. *Journal of Antimicrobial Chemotherapy* 65, 2582-2589, (2010).

Sambandamurthy, V. K., Wang, X., Chen, B., Russell, R. G., Derrick, S., Collins, F. M., Morris, S. L. & Jacobs, W. R., Jr. A pantothenate auxotroph of *Mycobacterium tuberculosis* is highly attenuated and protects mice against tuberculosis. *Nature Medicine* 8, 1171-1174, (2002).

Mayer, O., Jain, P., Weisbrod, T. R., Biro, D., Ho, L., Jacobs-Sera, D., Hatfull, G. F. & Jacobs, W. R., Jr. Fluorescent Reporter DS6A Mycobacteriophages Reveal Unique Variations in Infectibility and Phage Production in Mycobacteria. *J Bacteriol* 198, 3220-3232, (2016).

Dalecki, A. G., Haeili, M., Shah, S., Speer, A., Niederweis, M., Kutsch, O. & Wolschendorf, F. Disulfiram and Copper Ions Kill *Mycobacterium tuberculosis* in a Synergistic Manner. *Antimicrob Agents Chemother* 59, 4835-4844, (2015).

Jain, P., Hartman, T. E., Eisenberg, N., O'Donnell, M. R., Kriakov, J., Govender, K., Makume, M., Thaler, D. S., Hatfull, G. F., Sturm, A. W., Larsen,

M. H., Moodley, P. & Jacobs, W. R., Jr. phi(2)GFP10, a high-intensity fluorophage, enables detection and rapid drug susceptibility testing of *Mycobacterium tuberculosis* directly from sputum samples. *J Clin Microbiol* 50, 1362-1369, (2012).

37   Lamert, R. & Pearson, J. Susceptibility testing- accurate and reproducible minimum inhibitory concentration (MIC) and non-inhibitory concentration (NIC) values. *Journal of Applied Microbiology* 88, 784-790.