Nutritional redundancy in the human diet and its application in phenotype association studies

Xu-Wen Wang[1], Yang Hu[2], Giulia Menichetti[1,3], Francine Grodstein[4], Shilpa N. Bhupathiraju[1,2], Qi Sun[1,2,5], Xuehong Zhang[1,2], Frank B. Hu[1,2,5], Scott T. Weiss[1], Yang-Yu Liu[1,6]

[1]*Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02115, USA*

[2]*Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA*

[3]*Network Science Institute, Department of Physics, Northeastern University, Boston, MA 02115, USA*

[4]*Rush Alzheimer's Disease Center, Department of Internal Medicine, Rush Medical College, Rush University, Chicago, IL 60612, USA*

[5]*Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA*

[6]*Center for Artificial Intelligence and Modeling, The Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Champaign, IL 61801, USA*

Abstract

Studying human dietary intake may help us identify effective measures to treat or prevent many chronic diseases whose natural histories are influenced by nutritional factors. Here, by examining five cohorts with dietary intake data collected on different time scales, we show that the food intake profile varies substantially across individuals and over time, while the nutritional intake profile appears fairly stable. We refer to this phenomenon as 'nutritional redundancy' and attribute it to the nested structure of the food-nutrient network. This network enables us to quantify the level of nutritional redundancy for each diet assessment of any individual. Interestingly, this nutritional redundancy measure does not strongly correlate with any classical healthy diet scores, but its performance in predicting healthy aging shows comparable strength. Moreover, after adjusting for age, we find that a high nutritional redundancy is associated with lower risks of cardiovascular disease and type 2 diabetes.

Introduction

Human dietary intake fundamentally affects our nutrition, energy supply, and health. A better understanding of diet patterns can help us identify measures to prevent or treat health conditions and diseases such as obesity[1,2], type 2 diabetes[3,4], and cardiovascular disease[5,6]. Indeed, randomized trials have established the benefits[7] of the Mediterranean diet on clinical cardiovascular disease[8] and the Dietary Approaches to Stop Hypertension (DASH)[9] diet on blood

pressure control. To improve the diet quality, the 2020-2025 Dietary Guidelines for Americans recommended that consumers should select nutrient-dense foods and beverages, which can provide vitamins, minerals and other health-promoting components and have no or little added sugars, saturated fat, and sodium[10]. For example, intake of whole grains, legumes and vegetables and fruit is recommended to reduce ingredients, such as unhealthy fats, and excess sugar and sodium, which have been associated with many diseases such as coronary heart disease[11–13] and obesity, liver and other metabolic diseases[14–16].

Food is a complex combination of components that can be classified into nutrients and non-nutrients[17]. The preliminary step to developing a guide to healthy dietary guidance is nutrient profiling[18,19] --- the science of understanding which nutrients are present in a given food item. This has many potential applications, ranging from dietary guidance to nutrient labeling and the regulation of health claims[18]. The current nutrient profiling relies on some existing databases, such as the USDA's Food and Nutrient Database for Dietary Studies (FNDDS)[20,21], Frida[22], FooDB[23] and many other databases[24–26] (see **SI Sec. 1.1** for details). Those databases represent the most comprehensive efforts so far to integrate food composition data from specialized databases and experimental data.

Despite the various databases on food constituents, a comprehensive understanding of the food-nutrient relationship remains challenging. Human dietary patterns across the world could be drastically different[27]. For example, the Mediterranean diet is characterized by high consumption of olive oil, legumes and vegetables and lower consumption of non-fish meat products while the Western diet pattern is characterized by high consumption of highly processed meat[28–30], candy and sweets and low intakes of fruits, and vegetables. Although dietary patterns sometimes exhibit drastically divergent food choices, at a finer resolution, the underlying nutrient palettes could reveal a higher degree of similarity. This implies that despite the variability in food choice and intake, nutrient profiles can be highly stable across individuals and over time.

 Here, by analyzing human dietary intake data collected from different cohorts and on different time scales, we show that the food profile varies substantially across individuals and over time, while the nutritional profile is highly stable across individuals and over time. This phenomenon is strongly reminiscent of the phenomenon of functional redundancy observed in the human microbiome, i.e., the taxonomic profile is highly personalized while the gene composition (or functional profile) is highly conserved across individuals[31–33]. Hence, hereafter we will refer to this phenomenon as nutritional redundancy (NR). We find that NR cannot be simply explained by the fact that different foods contain similar nutrients. Instead, it is largely due to more sophisticated features (e.g., the highly nested structure) of the food-nutrient network --- a bipartite graph that connects foods to their nutrient constituents. The food-nutrient network also

enables us to quantify the level of NR for each diet assessment of any individual, i.e., the personal
NR. Interestingly, this personal NR measure does not strongly correlate with any classical healthy
diet scores, but its performance in predicting healthy aging shows comparable strength.
Moreover, after adjusting for age, we find that a high personal NR is associated with lower risks
of cardiovascular disease and type 2 diabetes. Hence, the concept of NR developed here may offer
us a new perspective on studying the human diet. The measure of personal NR could serve as a
powerful tool in nutrition science to modulate individual dietary patterns and study the
correlation with different health phenotypes.

Results

The phenomenon of population-level NR. To demonstrate the phenomenon of NR in the
human diet, we analyzed five comprehensive datasets with dietary data collected on different
time scales. (1) Diet-microbiome association study (DMAS)[34]: a longitudinal study of 34 healthy
individuals with dietary intake data and stool sample collected daily over 17 consecutive days.
Daily dietary intake data was collected using the automated self-administered 24-h (ASA24)
dietary assessment tool[35–37]. DMAS focused on 41 nutrients, and 9 food groups (e.g., grains, fruits,
vegetable, etc.) based on the Food and Nutrient Database for Dietary Studies (FNDDS) food
coding scheme[20]. Note that there are two outliers in DMAS, referred to as "shake drinkers", whose
reported diet consisted primarily of a nutritional meal replacement beverage. Those two outliers
were removed in our analysis. Also, in our analysis, we focused on those participants (in total
n=30) who have ASA24 data available for all the 17 days. (2) Nurse Healthy Study (NHS)[38,39]: NHS
began in 1976 when 121,700 female registered nurses, aged 30-55 years in the United States were
enrolled; the semiquantitative food frequency questionnaire (FFQ) was first administrated in 1980,
and then administrated approximately every four years. So far, we have up to eight time points
of FFQ data for NHS participants. In our analysis, we focused on those participants (in total
n=35,256) who have FFQ data available for all the eight time points. (3) Health Professionals
Follow-up Study (HPFS)[40,41]: HPFS is a prospective cohort of 51,529 men followed since 1986,
when participants initially ranged from 40-75 years. This all-male study was designed to
complement the all-female NHS/NHSII, and both were conducted identically. In our analysis, we
focused on those participants (in total n=1,7529) who have FFQ data available for all the 7 time
points. (4) Women Lifestyle Validation Study (WLVS)[37]: As a sub-study of the NHS and NHSII,
WLVS was designed to investigate the measurement-error structure associated with self-reported
dietary and physical activity assessments within one year (with up to four ASA24 records).
Women with a history of coronary heart disease, stroke, cancer, or major neurological disease
were excluded. Among the 796 enrolled participants, 692 completed at least one ASA24. In our
analysis, we focused on those WLVS participants with all four ASA24 records available (in total
n=216). (5) Men's Lifestyle Validation Study (MLVS)[42,43]: a one-year study nested in HPFS[40,41].

MLVS was designed to complement WLVS. In our analysis, we focused on those MLVS
participants with all of four ASA24 records available (in total n=451).

For those selected participants in each study, we first assessed the change in their food and
nutrient profiles (i.e., the relative abundances of food items and nutrients in their diet) over time.
The relative abundances of nutrients are calculated by converting the unit of each nutrient to
gram and then normalized by the total grams of all nutrients for an individual. We found that the
food profiles were highly dynamic for almost all individuals at different time scales: daily
(**Fig.1:a1**), monthly (**Fig.1:b1,c1**), and yearly (**Fig.1:d1, Fig.1:e1**). Moreover, the food profiles are
highly personalized[44], i.e., the intra-individual dissimilarity of the foods consumed over time is
significantly lower than the inter-individual dissimilarity of their consumed foods at both single
food (**Fig.S1a**) and food-group (**Fig.S1b**) level. By contrast, the nutrient profiles, as expected, were
highly conserved across different individuals and over the whole study time period for all five
studies (**Fig.1:a2-e2**), and were not highly personalized (**Fig.S1c**).

We observed that the most abundant food items consumed by individuals are sweetened
beverages and vegetables. This is partially due to the fact that the moisture content in these foods
is high, as well as their relevance as staple foods in different cuisines[45]. On the other hand, the
most abundant nutrients for all participants are carbohydrate, fat, and protein, confirming their
key role as dietary intake macronutrients, and representing broad families of chemical
compounds. Within the carbohydrate category, sugars and fiber are the driving factors.

To quantify the between-sample difference in food or nutrient profiles, we adopted the notion of
beta diversity from community ecology[46]. In particular, we used four different measures (Bray-
Curtis dissimilarity, root Jensen-Shannon divergence, Yue-Clayton distance, and negative
Spearman Correlation) to quantify the beta diversity. As shown in **Fig.S1**, the beta diversity of
nutritional profiles is significantly lower than that of food profiles at both single food (**Fig.S1d**)
and food-group (**Fig.S1e**) levels in terms of all the four measures of beta diversity for all the five
studies. While the beta diversity captured by food groups[47] is remarkably lower compared to the
food-level analysis, we found that it is still significantly higher than the beta diversity of
nutritional profiles.

Definition of personal NR. Consider a pool of N food items, which contains a collection of
M nutrients in total. The food profile $\boldsymbol{f}^{(v)} = \left[f_1^{(v)}, \cdots, f_N^{(v)}\right]$ of individual-v's diet assessment can
be directly related to its nutrient profile $\boldsymbol{n}^{(v)} = \left[n_1^{(v)}, \cdots, n_M^{(v)}\right]$ through the FNN (**Fig.2**). Here, $f_i^{(v)}$
(or $n_a^{(v)}$) represents the relative abundance of food-i (or nutrient-a) in the diet assessment of
individual-v. We define the FNN as a weighted bipartite graph connecting these foods to their
nutrients[5]. The FNN can be represented by an $N \times M$ incidence matrix $\boldsymbol{G} = (G_{ia})$, where a non-

negative value G_{ia} indicates the amount contributed by food-i to nutrient-a (see Fig.2a for the unit of each nutrient). The nutrient profile is given by $\boldsymbol{n}^{(v)} = c\boldsymbol{f}^{(v)}\boldsymbol{G}$, where $c = \left[\sum_{a=1}^{M}\sum_{i=1}^{N} f_i^{(v)} G_{ia}\right]^{-1}$ is a normalization constant.

A key advantage of the FNN is that it enables us to calculate the NR for each diet assessment of any individual, i.e., the within-sample or personal NR. In the ecological literature[32,33,48], the functional redundancy of a local community is interpreted as the part of its taxonomic diversity that cannot be explained by its functional diversity. Similarly, we can define the NR of a dietary assessment from a particular individual as the part of its food diversity (FD) that cannot be explained by its nutrient diversity (ND), i.e., NR = FD $-$ ND. Here we chose FD to be the Gini-Simpson index: GSI $\equiv 1 - \sum_{i=1}^{N} f_i^2 = \sum_{i=1}^{N}\sum_{j \neq i}^{N} f_i f_j$, representing the probability that randomly chosen two units of a individuals 's food profile (with replacement) belong to two different food groups; and ND is chosen to be the Rao's quadratic entropy $Q \equiv \sum_{i=1}^{N}\sum_{j \neq i}^{N} d_{ij} f_i f_j$, characterizing the mean nutritional distance between any two randomly chosen food items in the diet assessment[25,26]. Here $d_{ij} = d_{ji} \in [0, 1]$ denotes the nutritional distance between food-i and food-j. By definition, $d_{ii} = 0$ for $i = 1, \cdots, N$. For the sake of simplicity, to avoid major effects driven by the several orders of magnitude covered by nutrient amount in food[49], we compute d_{ij} as the (unweighted) Jaccard distance between the sets of nutrients within two food items (see **Methods** for definition). Note that with FD = GSI and ND = Q, we have NR = $\sum_{i=1}^{N}\sum_{j \neq i}^{N}(1 - d_{ij})f_i f_j$, naturally representing the nutrient similarity (or overlap) of two randomly chosen food items in any diet assessment.

The personal NR of each diet assessment is closely related to the phenomenon of population-level NR observed over a collection of diet assessments. Let's consider highly personalized food profiles from a population. There are two extreme cases: (i) Each food has its own unique nutrient content (**Fig.2b1**), hence $d_{ij} = 1$ for any $i \neq j$. In this case, for each individual we have FD = ND and NR = 0 (representing the lowest level of nutritional redundancy), and the nutrition profiles vary drastically across different individuals (**Fig.2c1**). (ii) All food share exactly the same nutrient contents (**Fig.2b3**), rendering $d_{ij} = 0$ for all i and j. In this case, for each individual we have ND = 0 and NR = FD (representing the highest level of nutritional redundancy), and the nutrition profiles are exactly the same for all individuals (**Fig.2c3**). These two extreme scenarios are of course unrealistic. In a more realistic intermediate scenario, the FNN has certain topological features such that different foods share a few common nutrients, but some foods are specialized to include some unique nutrients (**Fig.2b2**). In this case, the ND and NR of each individual's diet assessment can both be quite comparable, and the nutritional profiles can be highly conserved across individuals (**Fig.2c2**).

A reference FNN. To visualize the topological features of real FNNs, we constructed a
reference FNN based on USDA's Food and Nutrient Database for Dietary Studies (FNDDS) 2011-
2012[20], consisting of 7,618 foods and 65 nutrients and micronutrients. This reference FNN is
depicted as a bipartite graph, where for visualization purposes each food node represents one of
the nine highest-level food groups (based on the FNDDS food coding scheme) and each nutrient
node represents nutrient (**Fig.3a**). Note that the FNN associated with the diet assessment of any
individual can be considered as a particular subgraph of this reference FNN.

To characterize the structure of this reference FNN, we systematically analyzed its network
properties using the complete nutrient profile of all 7,618 foods. We first visualized its incidence
matrix (**Fig.3b**), where the presence (or absence) of a link connecting a food and a nutrient is
colored in green (or white), respectively. We found that some foods (e.g., bacon cheeseburger, hot
ham and cheese sandwich, corresponding to the leftmost columns in Fig.3b) contribute to almost
all of the nutrients, while some foods only include very few nutrients (e.g., sugar substitutes and
smart water, corresponding to the rightmost columns in Fig.3b). Moreover, we noticed that the
incidence matrix displays a highly nested structure, i.e., the nutrients of those food items (with
fewer distinct nutrients) in the right columns tend to be subsets of nutrients for those food items
(with more distinct nutrients) in the left columns. The nestedness of the FNN can be quantified
by utilizing the classical Nestedness metric based on Overlap and Decreasing Fill (NODF)
measure[50,51], and turns out to be much higher than expected by chance (see **Methods** for details).
We then calculated the nutritional distances among different food items, finding a unimodal
distribution with the peak centered around 0.25, indicating that most food items include very
similar nutrient components (**Fig.3c)**. Finally, we calculated the degree distributions of nutrient
nodes and food nodes, respectively. Here, the degree of a nutrient node in the FNN is just the
number of distinct foods that contain this nutrient. Similarly, the degree of a food node in the
FNN is the number of distinct nutrients it contains. We found that the degrees of food items
follow a Poisson-like distribution (**Fig.3d**), implying that different foods generally contain a very
similar number of nutrients. Nutrient degrees show a more extreme behavior, exhibiting a
probability density peak located at the higher end of the degree spectrum, indicating that the
majority of nutrients are different from zero in almost all foods collected in FNDDS, except a few
exceptions (**Fig.3e**).

FNN associated with each of the nine food groups and FNNs constructed by using other
databases, e.g., Frida[22] and Harvard food composition database (HFDB)[39], revealed very similar
network properties (see **Figs.S2, S3**). This confirms that the nested structure of the FNN is a
universal property across different databases, rather than determined by the nutrients' overlap
in USDA reference. In addition, we analyzed the FNN of 51 raw foods in HFDB, finding that the

network still displays high nestedness structure. However, NODF of this raw FNN is 0.573, which
is much lower than the original FNN (see **Fig.S4**).

We emphasize that the highly nested structure of the reference FNN is neither explained by the
presence of macronutrients, i.e., broad classes of chemicals such as carbohydrates, protein, and
fats that are key components of food and exhibit high degree, nor by the nutrient ontology used
to annotate the databases. First, as shown in Fig.3b, the incidence matrix of the FNN still displays
a highly nested structure even in the absence of high degree nutrients (the topmost green rows).
Second, FNN still shows highly nested structure after excluding nutrients without InChIKey[52],
effectively removing the first hierarchical level of the nutrient ontology, and also all those
nutrients that correspond to non-specific chemical mixtures (see **Fig.S5**). Third, if we randomize
the FNN but preserve the nutrient degree distribution, the randomized FNNs have much lower
nestedness than that of the real FNN, and the nutrient distances between different foods are
significantly increased (**Fig.S6**). Last but not least, we adopted tools from statistical physics[53] to
calculate the expected nestedness value and its standard deviation for an ensemble of
randomized FNNs in which the expected food and nutrient degree distributions match those of
the real FNN. We found that the expected nestedness of randomized FNNs is significantly lower
than that of the real FNN (one sample z-test yields $p_{\text{value}} < 10^{-5}$, see **Methods** for details).

Personal NR calculation based on dietary intake data. We calculated the personal
NR of diet assessments of those selected participants in the 4 studies: DMAS, WLVS, MLVS, NHS
and HPFS. First, we constructed a reference FNN based on the FNDDS to calculate the NR of
DMSA, WLVS and MLVS participants, and a reference FNN based on HFDB to calculate the NR
of NHS and HPFS participants (see **Methods** for details). Interestingly, we found that in all the 5
studies (DMAS, WLVS, MLVS, NHS and HPFS) NR~0.3 (**Fig.4**, purple boxes), suggesting that
nutritional redundancy and nutrient diversity are generally comparable in human diet.

The NR of the NHS at the population level displays a nonmonotonic decrease, indicating that the
diet patterns of those participants indeed have been adjusted. To better illustrate such a diet
pattern change, we projected the bipartite food-nutrient network (constructed from HFDB) into
the food space, resulting in a food similarity network (see **Fig.5a**). In this network, each node
represents a food item and a link connecting food item-i and item-j represents the unweighted
Jaccard similarity s_{ij} of their nutrient constituents (see **Methods**). Here, for visualization
purposes, only links with $s_{ij} \geq 0.85$ were retained. We found a clear modular structure in the
food similarity network, i.e., food items from the same food group form a densely connected
cluster or module (see **Fig.5a**), which is consistent with previous study that a food network based
on the foods' nutritional similarity displays separately clustered around animal-based foods and
plant-based foods at first, and fish and meats are separately clustered in animal-based food
cluster and grains, fruits, vegetables, nuts are separately clustered in plant-based food cluster[54].

Then, we examined the individual food similarity network of a particular NHS participant with the largest NR reduction from year 1984 to 2010 (see **Fig.5b,c**). We found the density of her food similarity network in 1984 is much higher than that in 2010, suggesting that this participant consumed foods with more overlapping nutrient constituents in 1984 (**Fig.5b**). Moreover, we found the most abundant food items in 2010 were water and yogurt, which do not connect with each other, indicating that she chose foods with more distinct nutrient constituents (**Fig.5c**).

Impact of FNN structure on personal NR. To identify key topological features of the FNN that determine the NR, we adopted tools from network science. In particular, we randomized the FNN using three randomization schemes, yielding three null models. Null-FNN-1: Complete randomization. We keep the number of foods (N) and number of nutrients (M) unchanged, but otherwise completely rewire the links between foods and nutrients. Null-FNN-2: Food-degree preserving randomization. We keep N, M, and the degree of each food node unchanged, but selects randomly the nutrients that link to each food. Null-FNN-3: Nutrient-degree preserving randomization. Here we keep N, M, and the degree of each nutrient unchanged, but select randomly the foods that link to each nutrient. Null-FNN-4: Nutrient-degree and food-degree preserving randomization. Here we keep N, M, and the degree of each nutrient and the degree of each food unchanged, but randomly rewire the links between food nodes and nutrient nodes. Then we recalculated the NR for each diet assessment (**Fig.4**). We found that for all the cohorts all the four null models yield much lower NR than that of the real FNN (**Fig.4**, purple boxes). This suggests that the real FNN must have certain topological features that determine the high NR in the human diet. Analyzing the network properties of those null models (**Fig.S6**), we found that those randomized FNNs all display lower nestedness and higher $\langle d_{ij} \rangle$ than those of the real FNN. Thus, the highly nested structure and low $\langle d_{ij} \rangle$ of the real FNN jointly contribute to the NR values observed in the human diet.

Impact of food composition on personal NR. To test if food composition plays an important role in determining NR, we randomized the food composition of each participant using three different randomization schemes, yielding three different null models. Then we recalculated NR for each sample (**Fig.4b**). We found that for each dietary intake record, if we preserve the food composition but randomly replace the food items by those present in the reference FNN, the resulting null model (Null-comp-1) always yields much higher NR than that of the original dietary record for ASA24 studies but is comparable for NHS and HPFS. This confirms that the food items present in each dietary record are not assembled at random but follow certain assembly rules. Interestingly, if we randomize the food compositions through random permutation of non-zero abundance for each dietary record across different food items (Null-comp-2) or for each food item across dietary records from different participants (Null-comp-3), we found that only Null-comp-3 did not significantly alter NR, suggesting that the

assembly rules are consistent across different participants. Taken together, results from the three
null composition models suggest that the nutritional redundancy in human diet is low and not
randomly selected.

Correlation between Personal NR and healthy diet scores. Adherence to healthy
diets has the potential to prevent disease and prolong life span. Many healthy diet scores have
been developed in the past. To test if the personal NR measure proposed here can be used to
quantify a healthy diet, we assessed the association between personal NR and the following four
existing healthy diet scores (see **SI Sec.2** for details): (i) Healthy Eating Index 2005 (HEI-2005)[55]:
a score that measures adherence to the USDA 2005 Dietary Guidelines for Americans. (ii)
Alternate Healthy Eating Index 2010 (AHEI-2010)[55]: a score that measures adherence to a diet
pattern based on foods and nutrients most predictive of disease risk in the literature.
(iii) Alternate Mediterranean Diet Score (AMED)[56]: a score adapted from form Mediterranean
Diet Score of Ref [57]. (iv) DASH Style Diet Score (DASH)[58]: a score capturing the characteristics
of the Dietary Approaches to Stop Hypertension diet. As shown in **Fig.6a**, we found that the NR
is positively correlated with AMED (Spearman correlation coefficient $\rho = 0.12$) and DASH ($\rho =$
0.08), but negatively correlated with HEI ($\rho = -0.16$) and AHEI ($\rho = -0.08$). But in all cases, the
Pearson correlation coefficients are weak, implying that personal NR should not be interpreted
as being synonymous with a healthy diet score. In other words, a heathy diet does not necessarily
have higher or lower nutritional redundancy.

Personal NR as an indicator of heathy aging. Then, we asked whether personal NR can
be an indicator of healthy aging --- an overall indicator we developed by combining measures of
physical function, cognitive function, mental health, and chronic diseases[59–61]. To address this
question, we focused on a subset of NHS participants (n=21,299) for whom we had all of these
four healthy aging components: healthy agers (n=3,491) and usual agers (n=17,808) through a
previous study[62]. The healthy agers are defined as those participants who survived beyond 65
years of age, with no history of chronic diseases, no reported memory impairment, no physical
disabilities, and intact mental health. The remaining participants who survived but did not
achieve good health in one or more domains were usual agers. The chronic disease domain used
to define usual agers includes cancer (other than nonmelanoma skin cancer), myocardial
infarction, coronary artery bypass surgery or percutaneous transluminal coronary angioplasty,
congestive heart failure, stroke, type 2 diabetes mellitus, kidney failure, chronic obstructive
pulmonary disease, Parkinson's disease, multiple sclerosis, and amyotrophic lateral sclerosis.
Other host factors that were collected in this previous study include age, education (registered
nurse, bachelor's degree, master, or doctorate), marital status (widowed, married, and
single/separated/divorced), median income from census tract (quintiles), BMI (<22, 22–24.9, 25–
29.9, and ≥30 kg/m^2), energy intake (quintiles of kcal/day), multivitamin use (yes/no), aspirin use

(<1, 1–2, or >2 tablets/week), pack-years of smoking (quintiles), and physical activity (quintiles of
MET-h/week) (see **Table S1** for details of those characteristics).

We first assessed the correlations between NR and these host factors. We found that BMI and
pack-years of smoking were negatively correlated with NR, while education, median income,
total energy intake, and physical activity were positively correlated with NR (**Fig.6b**). In all cases,
the Spearman correlation coefficients are very weak. To compare the prediction performance of
each of the four healthy diet scores (HEI-2005, AHEI-2010, AMED, DASH) with that of personal
NR in predicting healthy aging, we built a random forest classifier. In particular, we used features
(personal NR or one of the healthy diet scores, and the host factors mentioned above) collected
for those participants in 1998 to predict their heathy aging status in 2012 (see **SI Sec.5** for details).
We found that personal NR can achieve very similar error rate (i.e., the proportion of participants
that have been incorrectly classified by the model) and AUROC (area under the ROC curve) as
other four healthy diet scores (**Fig.6c,d**). Interestingly, we found NR can serve as a better indicator
than many other host factors (see **Fig.S7** for the importance ranking of those factors). We
emphasized that the healthy aging prediction does not rely on the particular NR definition or
classifier. We evaluated the performance of NR in healthy aging prediction using the Hill
number-based definition and another ensemble classifier: XGBoost[63], showing that both AUC and
error rate are robust to different Hill numbers and classifier (see **Figs.S8, S9** and **Methods** for
details).

We also performed the healthy aging prediction using data from a substudy of HPFS with 6,160
healthy agers and 11,534 usual agers[64]. Again, we used personal NR or one of the four healthy
diet scores in 1998 and other host factors to predict the healthy aging status. We found that NR
can also achieve very similar error rate (or AUROC) as other healthy diet scores in HPFS.
Moreover, the performance of NR in HPFS is comparable to that in NHS (**Fig.S10**).

The association between personal NR and the risks of type 2 diabetes and
cardiovascular disease. To further demonstrate the potential of the personal NR measure in
predicting well-defined phenotypes, we examined its associations with the risks of type 2
diabetes and cardiovascular disease in NHS and HPFS (see SI Sec.6 for detailed definitions of the
two disease outcomes). In particular, for each disease outcome, we used the age (months)- and
calendar year-stratified Cox proportional-hazard model to compute the hazard ratios and 95%
confidence intervals (CIs) of the disease according to tertiles of the NR for NHS participants from
1984 to 2014, and HPFS participants from 1986 to 2016.

For NHS participants, after adjusting for age, we observed the NR is associated with a lower risk
of the type 2 diabetes (see **Table 1**). In particular, those NHS participants whose NR values are at

tertile-2 and tertile-3 have a hazard ratio of 0.86 (95% CI: 0.80-0.93) and 0.78 (95% CI: 0.72-0.85),
respectively, with P for trend <0.001. To further check if this association is robust against many
other confounding factors, we also adjusted for total energy intake, race (white, African American,
Asian, others), BMI (<21.0, 21.0-22.9, 23.0-24.9, 25.0-26.9, 27.0-29.9, 30.0-32.9, 33.0-34.9, or ≥35.0
kg/m^2), smoking status (never smoked, past smoker, currently smoke 1-14 cigarettes per day, 15-
24 cigarettes per day, or ≥25 cigarettes per day), alcohol intake (0, 0.1-4.9, 5.0-9.9, 10.0-14.9, 15.0-
29.9, and ≥30.0 g/d), hypertension (yes, no), hypercholesterinemia (yes, no), multivitamin use (yes,
no), physical activity (quintiles), alternative healthy eating index, family history of diabetes,
postmenopausal hormone use (never, former, or current hormone use, or missing), and oral
contraceptive use. We found that those NHS participants whose NR values are at tertile-2 and
tertile-3 have a hazard ratio of 0.93 (95% CI: 0.86-1.01) and 0.93 (95% CI: 0.85-1.00), respectively,
with marginal P for trend = 0.09. For cardiovascular disease, we performed the same calculations.
Again, we observed that NR is associated with lower risk of the cardiovascular disease for NHS
participants (see **Table 1**). In particular, after adjusting for age (months) only, the P for trend <
0.001. After adjusting for a wide range of confounding factors, the P for trend = 0.006.

For HPFS participants, we observed similar results. For type 2 diabetes, after adjusting for age
(months) only, the P for trend < 0.001; after adjusting for a wide range of confounding factors, the
P for trend = 0.002 (see **Table 1**). For cardiovascular disease, after adjusting for age (months) only,
the P for trend = 0.004; after adjusting for a wide range of confounding factors, the P for trend =
0.04 (see **Table 2**).

For both disease outcomes and both cohorts, we also repeated the above calculations using
quintiles of the NR score. As shown in **Table S2**-**S3**, we found qualitatively very similar results.
372
Since NR is a part of FD and actually they are positively correlated (see **Fig.S11**), we wonder if
FD itself is associated with the risk of type 2 diabetes and cardiovascular disease. We performed
association analyses. Interestingly, for both NHS and HPFS participants, we found that, after
adjusting for a wide range of confounding factors, FD is not associated with lower risk of type 2
diabetes (see **Table S4**) or cardiovascular disease (see **Table S5**) at all. This result implies that the
association between NR and disease risks cannot be simply attributed to FD.

To understand the association between NR and the risk of type 2 diabetes and cardiovascular
disease in the two cohorts, we analyzed the food consumption pattern of each NR tertiles in those
two cohorts. We found that there is a consistent trend among the three NR tertiles for those
important food groups in NHS and HPFS (see **Fig.7**). For instance, abundances of Fruits,
Vegetables, Dairy, Cereal Grains are much higher for T3 (i.e., high-NR participants) than T2 and
T1; while abundances of Beverages are much lower for T3 than T2 and T1 in both NHS and HPFS.

This food consumption pattern might explain why NR is an indicator of low risk of type 2 diabetes
and cardiovascular disease.

Discussion

Through examining various human dietary intake datasets, we found that food profile varies
tremendously across individuals and over time, while the nutritional profile is highly conserved
across different individuals and over time. To quantify this nutritional redundancy, we
constructed the food-nutrient network --- a bipartite graph that connects foods to their nutrient
constituents. This food-nutrient network also allows us to assess the NR of any dietary assessment
from any individual. We found that this personal NR is not strongly correlated with any existing
healthy diet scores. We emphasize that, as the difference between the food diversity and the
nutrient diversity of a person's dietary assessment, the personal NR quantifies the nutrient
similarity (or overlap) of two randomly chosen food items in the diet assessment. Thus, a healthy
diet does not necessarily have higher or lower NR. Interestingly, we found that the personal NR
can be used to predict healthy aging with equally strong performance as those healthy diet scores.
Hence, the concept of personal NR offers us a completely new perspective on studying human
diet. Moreover, we examined its associations with the risks of type 2 diabetes and cardiovascular
disease in NHS (all female) and HPFS (all male). For both cohorts, we found a clear inverse
association between NR and the two phenotypes after adjusting for age. For HPFS, the inverse
association is observed even after adjusting for a wide range of confounding factors. Whether
these findings can lead to practical nutritional guidance warrant further interventional studies.

Since the personal NR measure is not strongly correlated with any classical healthy diet scores,
in principle we can combine the concepts of NR and those healthy diet scores to better capture
the total impact of diet on health outcomes. For instance, one can leverage the food-specific
subgraphs of the FNN (see **Fig.S2**) to calculate the NR of food groups contributing to each
component of a healthy diet score. This will enable us to define an NR-aware healthy diet score.
Systematically exploring this direction warrant dedicated efforts, which is beyond the scope of
the current work.

There are several limitations in our current framework of NR calculation. First, we did not
explicitly consider the nutrient difference between different food sources. We understand that
nutrient content and its fluctuations span several orders of magnitude, and different scaling
transformation, as well as different selections of nutrients, could modulate nutrient diversity and
redundancy across individuals[49]. We anticipate that incorporating this information in our NR
calculation will further improve the power of using NR to predict healthy aging or other disease
risks[65].

Second, the calculation of a personal NR relies on food intake measurements, e.g., ASA24 and FFQ, which are based on self-reported dietary intake questionnaires. We understand that such food intake measurements have inherent limitations, particularly measurement error related to poor recall, which can be overcome by the use of nutritional biomarkers that are capable of objectively assessing food consumption in different biological samples without the bias of self-reported dietary assessment[66]. Although nutritional biomarkers provide a more proximal measure of nutrient status than dietary intake, quantitatively studying NR using nutritional biomarkers is beyond the scope of the current study. We anticipate that our framework will trigger more research activities in this direction.

Third, due to the unmapped chemical complexity of food[26], the nutritional components listed in existing epidemiological databases represent only a small fraction of the several hundreds of thousands of molecules documented in food[67], many of which are bioactive substances contributed by the extensive secondary metabolism of plants, and substrates for the human gut microbiome[68,69]. Our NR calculation is based on the existing nutrient databases that are certainly incomplete. We anticipate that a more comprehensive food composition database will yield a more powerful measure of NR to predict healthy aging and other disease risks.

Forth, although we considered four null models of the bipartite food-nutrient network, systematically exploring other null models of this bipartite network (or its unipartite presentations) deserves dedicated efforts, which is beyond the scope of the current study. Hence, our analysis cannot entirely exclude the possibility of the presence of other network characteristics that are simpler than nestedness itself and can still be used to explain the NR values observed in the human diet.

Finally, food distance was measured using all nutrient components, which might introduce biases as some nutrients are correlated or redundant, such as fatty acids and amino acids. Although we have demonstrated that the FNN still shows a highly nested structure after excluding broad families of chemicals (**Fig.S5**), and that removing fat-related nutrient subtypes does not drastically alter NR (**Fig.S12**), more efforts are warranted to systematically address the potential impact of nutrition ontology on NR calculation.

METHODS

Food nutrient network. Consider a pool of N food items, which contains a collection of M nutrients in total. The food profile $f^{(v)} = \left[f_1^{(v)}, \cdots, f_N^{(v)}\right]$ of the diet record of individual-v can be directly related to its nutrient profile $n^{(v)} = \left[n_1^{(v)}, \cdots, n_M^{(v)}\right]$ through the FNN (Fig.2). Here, $f_i^{(v)}$

(or $n_a^{(v)}$) represents the relative abundance of food-i (or nutrient-a) in the diet record. And we

define the FNN as a weighted bipartite graph connecting these foods to their nutrients. The

FNN can be represented by an $N \times M$ incidence matrix $\boldsymbol{G} = (\boldsymbol{G_{ia}})$, where a non-negative value

$\boldsymbol{G_{ia}}$ indicates the amount contributed by food-i to nutrient-a (see Fig.2a for the unit of each

nutrient). The nutrient profile is given by $\boldsymbol{n}^{(v)} = \boldsymbol{cf}^{(v)} \boldsymbol{G}$, where $\boldsymbol{c} = \left[\sum_{a=1}^{M} \sum_{i=1}^{N} f_i^{(v)} G_{ia} \right]^{-1}$ is a

normalization constant.

Nestedness. Nestedness a classical concept in ecology, which is used to characterize the nested

structure of ecological systems, such as the species-site network (describing the distribution of

species across geographic locations), and the species-species interaction networks (e.g., host-

parasite, plant- pollinator interactions)[70–74]. In principle, an ecological system is said to be nested

if the items belonging to "smaller" elements (e.g., a small island containing few species, or a

specialist species with few interactions) tend to be a subset of the items belonging to "larger"

elements (e.g., a large island containing many species, or a generalist species with many

interacting partners). Mathematically, those ecological systems can be represented as bipartite

graphs with two types of nodes, e.g., sites and species, hosts and parasites, plants and pollinators,

etc. In this work, we focus on the food nutrient network of dietary, which is also a bipartite graph

with two types of nodes: foods and nutrients.

Numerical calculation. Consider a general bipartite graph with N type-1 nodes and M type-2

nodes. The structure of this bipartite graph can be represented by its $N \times M$ binary incidence

matrix $\boldsymbol{B} = (B_{ia})$, where $B_{ia} = 1$ if there is a link connecting the i-th type-1 node and the a-th

type-2 node, and 0 otherwise. Mathematically, nestedness can be defined as a property of the

incidence matrix $\boldsymbol{B}$. If there exists a permutation of rows and columns such that the set of links in

row-i contains the links in row-$(i + 1)$, and the set of links in column-a contains those in column-

$(a + 1)$, then B is a perfectly nested binary matrix. For example, consider the mainland and a series

of islands sorted according to their distances to the mainland. The mainland contains all the

species, the first island has a subset of species in the mainland, the second island has a subset of

species in the first island, etc.

The Nestedness metric based on Overlap and Decreasing Fill (NODF) nestedness metric is based

on two simple properties: decreasing fill and paired overlap. For any given bipartite graph with

incident matrix $\boldsymbol{B}$, the unweighted degree of i-th food node is $k_i = \sum_{a=1}^{M} B_{ia}$, and the unweighted

nutrient degree of a-th nutrient node is $k_a = \sum_{i=1}^{N} B_{ia}$. The number of nutrients shared by the i-th

food and the j-th food is given by $P_{ij} = \sum_{a=1}^{M} B_{ia} B_{ja}$. And the number of common foods that both

of a-th nutrient and b-th nutrient are included is given by $Q_{ij} = \sum_{i=1}^{N} B_{ia} B_{ib}$. Define $\tilde{P}_{ij} = 0$ if $k_i = $

k_j and $\tilde{P}_{ij} = P_{ij}/\min(k_i, k_j)$ otherwise. Then, define $\tilde{Q}_{ab} = 0$ if $k_a = k_b$ and $\tilde{P}_{ab} = P_{ab}/$

$\min(k_a, k_b)$ otherwise. The NODF is defined as

491
$$\text{NODF} = \frac{\sum_{i<j}^{N} \tilde{P}_{ij} + \sum_{a<b}^{M} \tilde{Q}_{ab}}{\frac{N(N-1)}{2} + \frac{M(M-1)}{2}}, \tag{1}$$

Theoretical approach. To theoretically analyze the nested structure of a given bipartite graph,

one can construct a grand canonical ensemble for this bipartite graph under the constraint that,

for the two types of nodes, the degree sequences in the ensemble match on average the empirical

ones[53]. This theoretical approach has two big advantages. First, constraining the ensemble's mean

degree sequence to be equivalent to the empirical one limits the possible effects of noisy data,

hence possible missing (false negative) or overrated (false positive) links can be dealt with

appropriately. Second, for this bipartite graph ensemble one can analytically derive the mean and

standard deviation of the distribution of any network property (such as the classical NODF

measure of nestedness) that can be analytically formulated in terms of the elements of the

bipartite adjacency matrix $\boldsymbol{B}$.

Nutritional distance measure. To avoid the influence of nutrient amount variability in foods,

we used the (unweighted) Jaccard index to quantify the nutritional distance between food item-

i and item-j:

506
$$d_{ij} = 1 - \frac{|G_i \cap G_j|}{|G_i \cup G_j|}, \tag{2}$$

507   where G_i represents the nutrients in food i. $d_{ij} = 0$ indicates that the food item-i and food item-j

508   share exactly the same nutrient constituents; $d_{ij} = 1$ means that they have totally different

509   nutrient constituents. The nutritional similarity between food item-i and item-j can be defined as

511
$$s_{ij} = 1 - d_{ij}. \tag{3}$$

Nutritional redundancy measure. In the main text, the nutritional redundancy (NR) is defined

as: $\text{NR}_\alpha = \text{FD}_\alpha - \text{ND}_\alpha$. FD_α is chosen to be the Gini-Simpson index: $\text{GSI} \equiv 1 - \sum_{i=1}^{N} p_i^2 =$

$\sum_{i=1}^{N} \sum_{j \neq i}^{N} p_i p_j$, representing the probability that two randomly chosen members of a subject's food

profile (with replacement) belong to two different food items; and ND_α is chosen to be the Rao's

quadratic entropy $Q \equiv \sum_{i=1}^{N} \sum_{j \neq i}^{N} d_{ij} p_i p_j$. Note that with $\text{FD}_\alpha = \text{GSI}$ and $\text{ND}_\alpha = Q$, we have $\text{NR}_\alpha =$

$\sum_{i=1}^{N} \sum_{j \neq i}^{N} (1 - d_{ij}) p_i p_j$, naturally representing the nutritional similarity (or overlap) of two

randomly chosen members in any subject's food profile. Here, we emphasize that various

taxonomic and functional diversity measurements can be used in NR definition, for example Hill

number.

Food diversity. Consider a subject' food profile of N foods with the relative abundance given by
a vector $\boldsymbol{f} = [f_1, \cdots, f_N]$. Hill introduced effective *number of species,* which assumes that the
taxonomic diversity (of order q) of a given subject with relative abundance profile $\boldsymbol{f} = [f_1, \cdots, f_N]$
is the same as that of an idealized subject of D equally abundant foods with relative abundance
profile $\tilde{\boldsymbol{f}} = [1/D, \cdots, 1/D]$.

$$\sum_{i=1}^{N} f_i^q = \sum_{i=1}^{D} (\tfrac{1}{D})^q = D^{1-q}, \tag{4}$$

529 This offers a parametric class of food diversity measures defined as follows:

$$\mathrm{FD}_q := \left(\sum_{i=1}^{N} f_i^q\right)^{\frac{1}{(1-q)}} \text{ for } q \neq 1. \tag{5}$$

531 And

$$\mathrm{FD}_1 := \lim_{q \to 1} \mathrm{FD}_q = \exp\left(-\sum_{i=1}^{N} f_i \log f_i\right). \tag{6}$$

533 Note that the Gini-Simpson index (GSI) used in the main text is related to FD_2 as follows:

$$\mathrm{GSI} := 1 - \sum_{i=1}^{N} f_i^2 = 1 - \frac{1}{\mathrm{FD}_2}. \tag{7}$$

535 **Nutritional diversity.** Consider a subject' food profile of N foods with the relative abundance
536 given by a vector $\boldsymbol{f} = [f_1, \cdots, f_N]$ and pair-wise nutritional distance matrix $\Delta = (d_{ij}) \in \mathbb{R}^{N \times N}$ with
537 $d_{ii} = 0$ for all $i = 1, \cdots, N$ and $d_{ij} = d_{ji} \geq 0$ for all $i \neq j$. Follow our former definition of functional
538 redundancy, we use the new pair-wise nutritional distance matrix to overcome the drawbacks of
539 original distance matrix

$$\Delta' = \left(d'_{ij}\right) = \begin{pmatrix} \lambda_1 & d_{12} & d_{13} & \cdots & d_{1N} \\ d_{21} & \lambda_2 & d_{23} & \cdots & d_{2N} \\ d_{31} & d_{32} & \lambda_3 & \cdots & d_{3N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ d_{N1} & d_{N2} & d_{N3} & \cdots & \lambda_N \end{pmatrix}, \tag{8}$$

541 where d_{ij} represents the original nutritional distance between food-i and j and $\lambda_i := \frac{\sum_{j \neq i}^{N} d_{ij}}{N-1}$ is the
542 average nutritional distance between food-i and all other foods. Note that, when different
543 species are equally distinct with a constant pairwise distance, λ is equal to this constant. The
544 final nutritional diversity of a subject can be defined as

$$\mathrm{ND}_q(Q') := D_q(Q') \cdot Q' = \left(\sum_{i=1}^{N} \sum_{j=1}^{N} \frac{d'_{ij}}{Q'} \left(f_i f_j\right)^q\right)^{\frac{1}{2(1-q)}} \cdot Q' \text{ for } q \neq 1, \tag{9}$$

546 and

$$\mathrm{ND}_1(Q') := D_1(Q') \cdot Q' = \exp\left[-\frac{1}{2} \sum_{i=1}^{N} \sum_{j=1}^{N} \frac{d'_{ij}}{Q'} f_i f_j \log\left(f_i f_j\right)\right] \cdot Q'. \tag{10}$$

548 **Nutritional redundancy.** In the literature of ecology, the functional redundancy (FR) of a sample
549 is often considered to be the part of the taxonomic diversity (TD) that cannot be explained by the
550 functional diversity (FD)[48]. Hence FR is typically defined to be the difference between TD and FD.

551 Similarly, we define the NR to be the difference between food diversity (FD) and nutritional
552 diversity (ND):

$$\text{NR} := \text{FD} - \text{ND}. \tag{11}$$

554 In the main text, we chose FD to be the Gini-Simpson index: $\text{GSI} \equiv 1 - \sum_{i=1}^{N} f_i^2 = \sum_{i=1}^{N} \sum_{j \neq i}^{N} f_i f_j$,
555 and ND is chosen to be the Rao's quadratic entropy $Q \equiv \sum_{i=1}^{N} \sum_{j \neq i}^{N} d_{ij} n_i n_j$. Hence,

$$\text{NR} = \text{GSI} - Q = 1 - \sum_{i=1}^{N} f_i^2 = \sum_{i=1}^{N} \sum_{j \neq i}^{N} (1 - d_{ij}) f_i f_j. \tag{12}$$

557 For the Hill number-based food diversity TD_q and nutritional diversity $\text{ND}_q(Q')$, and
558 define a parametric class of nutritional redundancy:

$$\text{NR}_q(Q') := \text{TD}_q - \text{ND}_q(Q'). \tag{14}$$

560 For $q \neq 1$,

$$\text{NR}_q(Q') := \left(\sum_{i=1}^{N} f_i^q \right)^{\frac{1}{(1-q)}} - \left(\sum_{i=1}^{N} \sum_{j=1}^{N} \frac{d'_{ij}}{Q'} (f_i f_j)^q \right)^{\frac{1}{2(1-q)}} \cdot Q'. \tag{15}$$

562 For $q = 1$,

$$\text{NR}_1(Q') := \exp\left(-\sum_{i=1}^{N} f_i \log f_i \right) - \exp\left[-\frac{1}{2} \sum_{i=1}^{N} \sum_{j=1}^{N} \frac{d'_{ij}}{Q'} f_i f_j \log(f_i f_j) \right] \cdot Q'. \tag{16}$$

Definition of disease outcomes. Type 2 diabetes was confirmed if at least one of the following
criteria by the National Diabetes Data Group[75] was met: (1) elevated plasma glucose levels (fasting
glucose ≥ **140** mg/dL or random glucose ≥ **200** mg/dL) with ≥ **1** classic symptoms (polydipsia,
polyuria, polyphagia, weight loss, or coma); (2) elevated plasma glucose on at least two occasions
(fasting glucose ≥ **140** mg/dL, random glucose ≥ **200** mg/dL, and/or glucose ≥ **200** mg/dL after
an oral glucose test) with no symptoms; and (3) hypoglycemic therapy with insulin or oral
medications. An updated cutoff of 126 mg/dL for fasting glucose was used for diagnoses after
1997 according to the American Diabetes Association diagnostic criteria[76].

The primary outcome measure was major Cardiovascular disease (CVD), which is defined as a
combined endpoint of non-fatal myocardial infarction, non-fatal stroke, or fatal CVD (fatal stroke,
fatal myocardial infarction, and other cardiovascular death). The secondary outcome measures
were assessed as following: (1) total CHD: defined as fatal CHD and nonfatal myocardial
infarction; (2) total stroke: all fatal and nonfatal stroke cases (ischemic, hemorrhagic, and
undetermined subtypes). When a participant (or family members of deceased participants)
reported an incident event, permission was requested to examine their medical records by

physicians who were blinded to the participant risk factor status. For each endpoint, the month
and year of diagnosis were recorded as the diagnosis date. Non-fatal events were confirmed
through review of medical records. Myocardial infarction was confirmed if the World Health
Organization criteria were met on the basis of symptoms plus diagnostic electrocardiogram
changes or elevated cardiac enzymes. If medical records were unavailable, we considered
myocardial infarctions probable when the participant provided additional confirmatory
information. Information on angina and coronary revascularization procedures (percutaneous
transluminal coronary angioplasty or coronary artery bypass grafting surgery) were self-reported,
and we included only events that occurred before a manifest cardiovascular event.
588
Strokes were confirmed if data in the medical records fulfilled the National Survey of Stroke
criteria requiring evidence of a neurological deficit with sudden or rapid onset that persisted
for >24h of until death[77]. We excluded cerebrovascular pathology due to infection, trauma, or
malignancy, as well as "silent" strokes discovered only by radiologic imaging. Radiology reports
of brain imaging (computed tomography or magnetic resonance imaging) were available in 89%
of those with medical records. We classified strokes as ischemic stroke (thrombotic or embolic
occlusion of a cerebral artery), hemorrhagic stroke (subarachnoid and intraparenchymal
hemorrhage), or stroke of probable/unknown subtype (a stroke was documented but the subtype
could not be ascertained owing to medical records being unobtainable).
598
Deaths were identified by reports of families, the U.S. postal authorities, and searches of the
National Death Index. The cause of death was assigned by physicians after review of medical
records and death certificate information. Follow-up for deaths was >98% complete. Fatal CVD
was defined as fatal CHD disease, fatal stroke, or fatal CVD. Fatal CHD was defined as ICD-9
(international classification of diseases, ninth revision) codes 410-412 and was considered
confirmed if fatal CHD was confirmed via medical records or autopsy reports or if CHD was
listed as the cause of death on the death certificate and there was prior evidence of CHD in the
medical records. We designated as probable those cases in which CHD was the underlying cause
on the death certificates, but no prior knowledge of CHD was indicated and medical records
concerning the death were unavailable. Similarly, we used ICD-9 codes 430-434 to define fatal
stroke and followed the same procedures to classify cases of confirmed or probable fatal stroke.
Lastly, fatal CVD was defined by ICD-9 codes 390-458.

Ethical approval statement. The study protocol was approved by the institutional review board
(IRB) of the Brigham and Women's Hospital, and the IRB allowed participants' completion of
questionnaires to be considered as implied consent. Written informed consent was obtained from
participants to release medical records documenting.

Data availability. The DMAS dataset is available from https://github.com/knights-
lab/dietstudy_analyses. The USDA Food Nutrient Database is available from
https://fdc.nal.usda.gov/download-datasets.html. The Women's lifestyle validation study, Men's
lifestyle validation study, Nurses' Health Study (NHS) and Health Professionals Follow-Up
Study (HPFS) are available under restricted access due to participant confidentiality and privacy
concerns. Access can be obtained by submitting an online request form. If the project is approved,
completion of a data use agreement, completion of CITI training demonstrating ethical training
in using human subjects' data, and a small provision of funds to support the computing system
will also be required. Harvard University Food Composition Database can be accessed at
https://regepi.bwh.harvard.edu/health/nutrition.html. FoodDB database can be accessed at
https://foodb.ca/downloads. Source data are provided with this paper.

Code availability. MATLAB Code (version R2016b) used in this work is available at
https://github.com/spxuw/Nutritional-redundancy or under Zenodo at
https://doi.org/10.5281/zenodo.7781521[78].

References

1. Richards, T. J., Patterson, P. M. & Tegene, A. Obesity and nutrient consumption: a rational
addiction? *Contemporary Economic Policy* **25**, 309–324 (2007).
2. Schönfeldt, H. C. & Gibson, N. Changes in the nutrient quality of meat in an obesity context.
Meat science **80**, 20–27 (2008).
3. Keen, H., Thomas, B. J., Jarrett, R. J. & Fuller, J. H. Nutrient intake, adiposity, and diabetes. *Br
Med J* **1**, 655–658 (1979).
4. Patti, M.-E. & Kahn, B. B. Nutrient sensor links obesity with diabetes risk. *Nature medicine* **10**,
1049–1050 (2004).
5. Milanlouei, S. *et al.* A systematic comprehensive longitudinal evaluation of dietary factors
associated with acute myocardial infarction and fatal coronary heart disease. *Nat Commun*
11, 6074 (2020).
6. Bellisle, F. Why should we study human food intake behaviour? *Nutrition, Metabolism and
Cardiovascular Diseases* **13**, 189–193 (2003).
7. Song, M. *et al.* Animal and plant protein intake and all-cause and cause-specific mortality:
results from two prospective US cohort studies. *JAMA Intern Med* **176**, 1453–1463 (2016).
8. Martínez-González, M. A. *et al.* Benefits of the Mediterranean diet: insights from the
PREDIMED study. *Progress in cardiovascular diseases* **58**, 50–60 (2015).
9. Chiu, S. *et al.* Comparison of the DASH (Dietary Approaches to Stop Hypertension) diet and a
higher-fat DASH diet on blood pressure and lipids and lipoproteins: a randomized controlled
trial1–3. *The American Journal of Clinical Nutrition* **103**, 341–347 (2016).

10. Dietary Guidelines for Americans, 2020-2025. 164.

11. Ulbricht, T. L. V. & Southgate, D. A. T. Coronary heart disease: seven dietary factors. *The*
lancet **338**, 985–992 (1991).

12. Hu, F. B. & Willett, W. C. Optimal diets for prevention of coronary heart disease. *Jama* **288**,
2569–2578 (2002).

13. Hu, F. B. *et al.* Dietary fat intake and the risk of coronary heart disease in women. *New*
England journal of medicine **337**, 1491–1499 (1997).

14. Kratz, M., Baars, T. & Guyenet, S. The relationship between high-fat dairy consumption and
obesity, cardiovascular, and metabolic disease. *European journal of nutrition* **52**, 1–24
(2013).

15. Fu, J.-H. *et al.* The effects of a fat-and sugar-enriched diet and chronic stress on
nonalcoholic fatty liver disease in male Wistar rats. *Digestive diseases and sciences* **55**,
2227–2236 (2010).

16. Oddy, W. H. *et al.* The Western dietary pattern is prospectively associated with nonalcoholic
fatty liver disease in adolescence. *American Journal of Gastroenterology* **108**, 778–785
(2013).

17. Chen, Y., Michalak, M. & Agellon, L. B. Focus: Nutrition and Food Science: Importance of
Nutrients and Nutrient Metabolism on Human Health. *The Yale journal of biology and*
medicine **91**, 95 (2018).

18. Drewnowski, A. & Fulgoni III, V. Nutrient profiling of foods: creating a nutrient-rich food
index. *Nutrition reviews* **66**, 23–39 (2008).

19. Garsetti, M., de Vries, J., Smith, M., Amosse, A. & Rolf-Pedersen, N. Nutrient profiling
schemes: overview and comparative analysis. *European journal of nutrition* **46**, 15–28
(2007).

20. Martin, C. L. *et al.* USDA Food and Nutrient Database for Dietary Studies 2011–2012:
Documentation and User Guide. Beltsville, MD: US Department of Agriculture.

21. Sebastian, R. S. *et al.* Flavonoid values for USDA survey foods and beverages 2007–2010.
Worldwide Web Site: Food Surveys Research Group (2016).

22. National Food Institute. Technical University of Denmark. *Frida Food data version 2 (DTU,*
2016); frida.fooddata.dk.

23. FooDB. Compounds http://foodb.ca/compounds (access 1 August 2019).

24. Neveu, V. *et al.* Phenol-Explorer: an online comprehensive database on polyphenol
contents in foods. *Database* **2010**, (2010).

25. Plumb, J. *et al.* Ebasis (bioactive substances in food information systems) and bioactive
intakes: Major updates of the bioactive compound composition and beneficial bioeffects
database and the development of a probabilistic model to assess intakes in europe.
Nutrients **9**, 320 (2017).

26. Barabási, A.-L., Menichetti, G. & Loscalzo, J. The unmapped chemical complexity of our diet.
Nat Food **1**, 33–37 (2020).

27. Cespedes, E. M. & Hu, F. B. *Dietary patterns: from nutritional epidemiologic analysis to*
national guidelines. *The American journal of clinical nutrition* vol. 101 899–900 (Oxford
University Press, 2015).

28. Micha, R., Wallace, S. K. & Mozaffarian, D. Red and processed meat consumption and risk of
incident coronary heart disease, stroke, and diabetes mellitus: a systematic review and
meta-analysis. *Circulation* **121**, 2271–2283 (2010).

29. Lustig, R. H. Processed food—an experiment that failed. *JAMA pediatrics* **171**, 212–214
(2017).

30. Menichetti, G., Ravandi, B., Mozaffarian, D. & Barabási, A.-L. Machine learning prediction of
food processing. *MedRxiv* 2021.05. 22.21257615 (2021).

31. The Human Microbiome Project Consortium. Structure, function and diversity of the
healthy human microbiome. *Nature* **486**, 207–214 (2012).

32. Louca, S. Function and functional redundancy in microbial systems. 8 (2018).

33. Tian, L. *et al.* Deciphering functional redundancy in the human microbiome. *Nat Commun*
11, 6217 (2020).

34. Johnson, A. J. *et al.* Daily Sampling Reveals Personalized Diet-Microbiome Associations in
Humans. *Cell Host & Microbe* **25**, 789-802.e5 (2019).

35. Subar, A. F. *et al.* The Automated Self-Administered 24-hour dietary recall (ASA24): a
resource for researchers, clinicians, and educators from the National Cancer Institute.
Journal of the Academy of Nutrition and Dietetics **112**, 1134–1137 (2012).

36. Kirkpatrick, S. I. *et al.* Performance of the Automated Self-Administered 24-hour Recall
relative to a measure of true intakes and to an interviewer-administered 24-h recall. *The*
American journal of clinical nutrition **100**, 233–240 (2014).

37. Yuan, C. *et al.* Validity of a Dietary Questionnaire Assessed by Comparison With Multiple
Weighed Dietary Records or 24-Hour Recalls. *Am J Epidemiol* **185**, 570–584 (2017).

38. Colditz, G. A. & Hankinson, S. E. The Nurses' Health Study: lifestyle and health among
women. *Nature Reviews Cancer* **5**, 388–396 (2005).

39. Colditz, G. A., Manson, J. E. & Hankinson, S. E. The Nurses' Health Study: 20-year
contribution to the understanding of health among women. *Journal of Women's Health* **6**,
49–62 (1997).

40. Chiuve, S. E. *et al.* Primary prevention of stroke by healthy lifestyle. *Circulation* **118**, 947
(2008).

41. Mukamal, K. J., Chiuve, S. E. & Rimm, E. B. Alcohol consumption and risk for coronary heart
disease in men with healthy lifestyles. *Archives of Internal Medicine* **166**, 2145–2150 (2006).

42. Mehta, R. S. *et al.* Stability of the human faecal microbiome in a cohort of adult men.
Nature microbiology **3**, 347–355 (2018).

43. Abu-Ali, G. S. *et al.* Metatranscriptome of human faecal microbial communities in a cohort
of adult men. *Nature microbiology* **3**, 356–366 (2018).

44. Hunter, D. J. *et al.* Variability in portion sizes of commonly consumed foods among a
population of women in the UnitedStates. *American journal of epidemiology* **127**, 1240–
1249 (1988).

45. Rutledge, K., Ramroop, T. & Boughdreau, D. Food Staple. *National Geographic.*
https://www. nationalgeographic. org/encyclopedia/food-staple/. Published February **22**,
(2011).

46. Legendre, P. & De Cáceres, M. Beta diversity as the variance of community data:
dissimilarity coefficients and partitioning. *Ecology letters* **16**, 951–963 (2013).

47. Rhodes, D. G., Adler, M. E., Clemens, J. C. & Moshfegh, A. J. What we eat in America food
categories and changes between survey cycles. *Journal of Food Composition and Analysis*
64, 107–111 (2017).
48. Pillar, V. D. *et al.* Functional redundancy and stability in plant communities. *Journal of*
Vegetation Science **24**, 963–974 (2013).
49. Menichetti, G. & Barabási, A.-L. Nutrient concentrations in food display universal behaviour.
Nature Food **3**, 375–382 (2022).
50. Almeida-Neto, M., Guimaraes, P., Guimaraes Jr, P. R., Loyola, R. D. & Ulrich, W. A consistent
metric for nestedness analysis in ecological systems: reconciling concept and measurement.
Oikos **117**, 1227–1239 (2008).
51. Ulrich, W., Almeida-Neto, M. & Gotelli, N. J. A consumer's guide to nestedness analysis.
Oikos **118**, 3–17 (2009).
52. Heller, S. R., McNaught, A., Pletnev, I., Stein, S. & Tchekhovskoi, D. InChI, the IUPAC
International Chemical Identifier. *J Cheminform* **7**, 23 (2015).
53. Payrató-Borràs, C., Hernández, L. & Moreno, Y. Breaking the Spell of Nestedness: The
Entropic Origin of Nestedness in Mutualistic Systems. *Phys. Rev. X* **9**, 031024 (2019).
54. Kim, S., Sung, J., Foo, M., Jin, Y.-S. & Kim, P.-J. Uncovering the Nutritional Landscape of
Food. *PLoS ONE* **10**, e0118697 (2015).
55. Chiuve, S. E. *et al.* Alternative dietary indices both strongly predict risk of chronic disease.
The Journal of nutrition **142**, 1009–1018 (2012).
56. Fung, T. T. *et al.* Mediterranean diet and incidence and mortality of coronary heart disease
and stroke in women. *Circulation* **119**, 1093 (2009).
57. Trichopoulou, A., Costacou, T., Bamia, C. & Trichopoulos, D. Adherence to a Mediterranean
diet and survival in a Greek population. *New England Journal of Medicine* **348**, 2599–2608
(2003).
58. Fung, T. T. *et al.* Adherence to a DASH-style diet and risk of coronary heart disease and
stroke in women. *Archives of internal medicine* **168**, 713–720 (2008).
59. Baumgartner, R. N. Body composition in healthy aging. *Annals of the New York Academy of*
Sciences **904**, 437–448 (2000).
60. Carmona, J. J. & Michan, S. Biology of healthy aging and longevity. *Revista de investigacion*
clinica **68**, 7–16 (2016).
61. Willcox, D. C., Scapagnini, G. & Willcox, B. J. Healthy aging diets other than the
Mediterranean: a focus on the Okinawan diet. *Mechanisms of ageing and development* **136**,
148–162 (2014).
62. Freitas-Simoes, T.-M., Wagner, M., Samieri, C., Sala-Vila, A. & Grodstein, F. Consumption of
Nuts at Midlife and Healthy Aging in Women. *Journal of Aging Research* **2020**, 1–7 (2020).
63. Chen, T. & Guestrin, C. XGBoost: A Scalable Tree Boosting System. *Proceedings of the 22nd*
ACM SIGKDD International Conference on Knowledge Discovery and Data Mining 785–794
(2016) doi:10.1145/2939672.2939785.
64. Zheng, Y. *et al.* Associations of Weight Gain From Early to Middle Adulthood With Major
Health Outcomes Later in Life. *JAMA* **318**, 255 (2017).
65. Giovannucci, E. *et al.* Intake of fat, meat, and fiber in relation to risk of colon cancer in men.
Cancer research **54**, 2390–2397 (1994).

66. Picó, C., Serra, F., Rodríguez, A. M., Keijer, J. & Palou, A. Biomarkers of Nutrition and Health:
New Tools for New Approaches. *Nutrients* **11**, (2019).
67. Menichetti, G. An AI pipeline to investigate the binding properties of poorly annotated
molecules. *Nature Reviews Physics* **4**, 359–359 (2022).
68. Balzerani, F. *et al.* Prediction of degradation pathways of phenolic compounds in the human
gut microbiota through enzyme promiscuity methods. *npj Systems Biology and Applications*
8, 24 (2022).
69. Kerimi, A., Kraut, N. U., da Encarnacao, J. A. & Williamson, G. The gut microbiome drives
inter-and intra-individual differences in metabolism of bioactive small molecules. *Scientific*
Reports **10**, 1–12 (2020).
70. Patterson, B. D. The principle of nested subsets and its implications for biological
conservation. *Conservation Biology* **1**, 323–334 (1987).
71. Patterson, B. D. & Atmar, W. Nested subsets and the structure of insular mammalian faunas
and archipelagos. *Biological journal of the Linnean society* **28**, 65–82 (1986).
72. Brualdi, R. A. & Sanderson, J. G. Nested species subsets, gaps, and discrepancy. *Oecologia*
119, 256–264 (1999).
73. Bascompte, J., Jordano, P., Melián, C. J. & Olesen, J. M. The nested assembly of plant–
animal mutualistic networks. *Proceedings of the National Academy of Sciences* **100**, 9383–
9387 (2003).
74. Rezende, E. L., Lavabre, J. E., Guimarães, P. R., Jordano, P. & Bascompte, J. Non-random
coextinctions in phylogenetically structured mutualistic networks. *Nature* **448**, 925–928
(2007).
75. Group, N. D. D. Classification and diagnosis of diabetes mellitus and other categories of
glucose intolerance. *diabetes* **28**, 1039–1057 (1979).
76. Gavin III, J. R., Alberti, K., Davidson, M. B. & DeFronzo, R. A. Report of the expert committee
on the diagnosis and classification of diabetes mellitus. *Diabetes care* **20**, 1183 (1997).
77. Walker, A. E. The national survey of stroke. Clinical findings. *Stroke.* **12**, I13-44 (1981).
78. Xu-Wen Wang. (2023). Code to compute nutritional redundancy.
https://doi.org/10.5281/zenodo.7781521

Acknowledgements. We thank Laura Sampson, Walter C. Willett, Rui Song, and Jorge Chavarro
for valuable discussions. Research reported in this publication was supported by grants
R01AI141529, R01HD093761, RF1AG067744, UH3OD023268, U19AI095219, and U01HL089856
from National Institutes of Health.

Author Contributions. Y.-Y.L. conceived and designed the project. X.-W.W. performed all the
numerical calculations and real data analyses. X.-W.W. and Y.H. performed the disease
association analyses. All authors analyzed and interpreted the results. X.-W.W. and Y.-Y.L. wrote
the manuscript. All authors edited the manuscript.
819

820 **Competing Interests Statements.** The authors declare no competing financial interests.
821 Correspondence and requests for materials should be addressed to Y.-Y.L.
822 (yyl@channing.harvard.edu).
823

Table 1. Hazard ratios (95% confidence intervals) of type 2 diabetes according to tertiles of NR in the Nurses' Health Study (NHS, 1984-2014) and Health Professionals Follow-Up Study (HPFS, 1986-2016).

NHS	T1	T2	T3	P for trend[1]
Cases/Person-year	1,436/312,411	1,245/312,509	1,133/312,388	
Age-adjusted model	1 (reference)	0.86(0.80,0.93)	0.78(0.72,0.85)	<0.001
Multivariable-adjusted model[2]	1 (reference)	0.93(0.86,1.01)	0.93(0.85,1.00)	0.0997
HPFS				
Cases/Person-year	684/147,746	502/148,000	502/148,085	
Age-adjusted model	1 (reference)	0.73(0.65,0.82)	0.73(0.65,0.82)	<0.001
Multivariable-adjusted model[2]	1 (reference)	0.77(0.69,0.87)	0.82(0.73,0.93)	0.0016

[1]P for trend was calculated using the median value of each tertiles (two-sided Chi-square test).
[2]Multivariable-adjusted model adjusted for age (years), ethnicity (white, African American, Asian, others), body mass index (<21.0, 21.0-22.9, 23.0-24.9, 25.0-26.9, 27.0-29.9, 30.0-32.9, 33.0-34.9, or ≥35.0 kg/m^2), smoking status (never smoked, past smoker, currently smoke 1-14 cigarettes per day, 15-24 cigarettes per day, or ≥25 cigarettes per day), alcohol intake (0, 0.1-4.9, 5.0-9.9, 10.0-14.9, 15.0-29.9, and ≥30.0 g/d), hypertension (yes, no), hypercholesterinemia (yes, no), multivitamin use (yes, no), physical activity (quintiles), alternative healthy eating index, family history of diabetes. In NHS, postmenopausal hormone use (never, former, or current hormone use, or missing) and oral contraceptive use were additionally adjusted.

Table 2. Hazard ratios (95% confidence intervals) of cardiovascular disease according to tertiles of NR in the Nurses' Health Study (NHS, 1984-2014) and Health Professionals Follow-Up Study (HPFS, 1986-2016).

NHS	T1	T2	T3	P for trend[1]
Cases/Person-year	1,477/325,684	1,324/325,688	1,250/325,419	
Age-adjusted model	1 (reference)	0.92(0.85,0.99)	0.85(0.79,0.92)	<0.001
Multivariable-adjusted model[2]	1 (reference)	0.94(0.87,1.02)	0.90(0.83,0.97)	0.0057
HPFS				
Cases/Person-year	1,358/145,486	1,285/145,662	1,290/145,726	
Age-adjusted model	1 (reference)	0.94(0.87,1.02)	0.89(0.83,0.96)	0.0041
Multivariable-adjusted model[2]	1 (reference)	0.97(0.89,1.04)	0.92(0.85,1.00)	0.0405

[1]P for trend was calculated using the median value of each tertiles (two-sided Chi-square test).

[2]Multivariable-adjusted model adjusted for age (years), ethnicity (white, African American, Asian, others), body mass index (<21.0, 21.0-22.9, 23.0-24.9, 25.0-26.9, 27.0-29.9, 30.0-32.9, 33.0-34.9, or ≥35.0 kg/m²), smoking status (never smoked, past smoker, currently smoke 1-14 cigarettes per day, 15-24 cigarettes per day, or ≥25 cigarettes per day), alcohol intake (0, 0.1-4.9, 5.0-9.9, 10.0-14.9, 15.0-29.9, and ≥30.0 g/d), hypertension (yes, no), hypercholesterinemia (yes, no), multivitamin use (yes, no), physical activity (quintiles), alternative healthy eating index, family history of myocardial infarction. In NHS, postmenopausal hormone use (never, former, or current hormone use, or missing) and oral contraceptive use were additionally adjusted.

Figure 1: Dietary food profiles are highly personalized while the nutrient profiles are highly stable. Rows represent results from selected participants of different cohorts: (**a**) DMAS (dietary intake data collected using ASA24 dietary assessment tool daily over 17 consecutive days); (**b**) WLVS (with four ASA24 records within one year); (**c**) MLVS (with four ASA24 records within one year); (**d**) NHS (with FFQ administrated every four years and with total eight time points). **(e)** HPFS (with FFQ administrated every four years and with total seven time points). Columns: (**1**) food profiles; (**2**) nutrient profiles. For DMAS, we plot the food and nutrient profiles of 30 participants who have ASA24 data available for all the 17 days. For the WLVS, MLVS and NHS datasets, we plot the food and nutrient profiles of 50 randomly chosen individuals. In the visualization of nutrient profiles, we only show the top-15 most abundant nutrients, while the remaining nutrients (after excluding amino acids, total fatty acids of saturated, monounsaturated, and polyunsaturated) are summarized as others.

Figure 2: Structure of the food-nutrient network is crucial for determining the nutrient redundancy of human diet. a: The food profiles vary drastically across individuals. **b1-b3:** Food-nutrient network is a bipartite graph that connect food items to their nutrient constituents, with edge weight representing the amount contributed by food-i to nutrient-a **b1:** Each food has a unique set of nutrient constituents. **b2:** Different foods share a few common nutrients; some foods are specialized to have some unique nutrients. **b3:** All foods share exactly the same set of nutrients. For each individual, its nutritional profile can be calculated from its food profile in (a) and the food-nutrient network in (b). **c1:** The nutritional profiles vary drastically across different individuals. **c2:** The nutritional profiles are highly conserved across different individuals. **c3:** The nutritional profiles are exactly the same across all individuals.

Figure 3: The food-nutrient network (FNN) has a nested structure. a: We used USDA's Food
and Nutrient Database for Dietary Studies (FNDDS) to construct the reference FNN, which
consists of 7,618 foods and 65 nutrients and micronutrients. This reference FNN is depicted as a
bipartite graph, where for visualization purposes each food node represents one of the nine
highest-level food groups (based on the FNDDS food coding scheme) and each nutrient node
represents a nutrient. **b:** The incidence matrix of the FNN, where the presence (or absence) of a
link between a food and a nutrient is colored in green (or white), respectively. NODF: Nestedness
metric based on Overlap and Decreasing Fill. **c:** The probability distribution of nutrient distances
(d_{ij}) among different food items. The bin size is 0.02. **d:** The food degree distribution. Here, the
degree of a food item is the number of distinct nutrients it contains. **e:** The nutrient degree
distribution. Here, the degree of a nutrient is the number of food items that contain this nutrient.
The 22 nutrients represent the amino acids and total fatty acids of saturated (S) monounsaturated
(M), and polyunsaturated (P) were not shown for visualization in panel a.

890
Figure 4: Topological features of the food-nutrient network and the human dietary pattern
contributes to the nutritional redundancy. a1-a5: The box plots of the nutritional redundancy
were calculated from the real FNN (purple box, n=30, 216, 451, 35,256 and 17,529 independent
subjects), as well as the randomized FNNs (other colored boxes, n=30, 216, 451, 35,256 and 17,529
independent subjects) using four different randomization schemes: Complete randomization
(Null-FNN-1); Food-degree preserving randomization (Null-FNN-2); Nutrient-degree
preserving randomization (Null-FNN-3); Food- and nutrient-degree preserving randomization
(Null-FNN-4). Here the degree of a nutrient is the number of foods that contain it, and the degree
of a food is the number of nutrients contained in it. **b1-b5:** The box plots of NR were calculated
from the real food composition (Real-comp, purple box, n=30, 216, 451, 35,256 and 17,529
independent subjects), as well as the randomized food compositions (other colored boxes, n=30,
216, 451, 35,256 and 17,529 independent subjects) using three different randomization schemes:
Randomized food assemblage generated by randomly choosing the same number of food items
from the food pool but keeping the food profile unchanged (Null-comp-1); Randomized food
abundance profiles through random permutation of non-zero abundance for each participant
across different foods (Null-comp-2); Randomized food abundance profiles through random
permutation of non-zero abundance for each food across different participants (Null-comp-3).
For the visualization purpose, we only showed four time points for each study. Boxes indicate
the interquartile range between the first and third quartiles with the central mark inside each box
indicating the median. Whiskers extend to the lowest and highest values within 1.5 times the
interquartile range.
912

Figure 5: Reference and personal food similarity networks. a: The reference food similarity network (FSN) was projected from the reference food-nutrient network constructed from the Harvard food composition database used in the NHS. The similarity between food-i and food-j was calculated using the unweighted Jaccard similarity index s_{ij}. Only links with $s_{ij} \geq 0.85$ were kept in the visualization. The color of each node represents the food group it belongs to. **b:** The personal FSN constructed for a particular NHS participant (ID #12137) using her FFQ data collected in 1984. This network is a subgraph of the reference FSN shown in panel-a, thus those nodes and links in global network but do not appear in the subnetwork are shown in white. **c:** The personal FSN of the same NHS participant constructed using her FFQ data collected in 2010. In panel-b and c, node sizes are proportional to the relative abundance of the food items and edge widths are proportional to the food similarities.

Figure 6: Nutrient redundancy serves as a potential metric to predict healthy aging in NHS.
a: Spearman correlation between the nutritional redundancy (NR) and existing healthy diet scores: Alternate Mediterranean Diet Score (AMED); Alternate Healthy Eating Index 2010 (AHEI-2010); Dash Style Diet Score (DASH); and Healthy Eating Index 2005 (HEI-2005). P values were calculated from two-sided t-test. **b:** Pearson correlation between NR and several host factors: body-mass index (BMI); education level; median income; energy intake level; Aspirin use; pack-years of smoking; physical activity. P values were calculated from two-sided t-test. **c:** Error rate of random forest classifier in the prediction of healthy aging status. **d:** AUROC of random forest classifier in prediction of healthy aging status. The participants are randomly spitted into 80% as the training set and the remaining 20% as the test set. The boxplot represents the performances of 200 independent splits. Boxes indicate the interquartile range between the first and third quartiles with the central mark inside each box indicating the median. Whiskers extend to the lowest and highest values within 1.5 times the interquartile range. All FDR-corrected P values were found using the paired and two-sided t-test. Significance levels: FDR-corrected p value <0.0001(****); <0.01(**); >0.05 (ns).

941
942 **Figure 7: Square root of median relative abundance of each food group for each tertile group**
943 **(T1-T3) based on the NR of participants in NHS (a) and HPFS (b).** Food group was defined in
944 Harvard food nutrition database and food groups presented in less than 3 tertile groups were not
945 shown.
946

Nutritional Redundancy in the Human Diet and its Application in Phenotype association studies

Supplemental Information

1. NR calculating using human dietary data

1.1 Reference FNN

1.1.1 USDA database

To construct the Food-Nutrient network, we downloaded the FNDDS 2011-2012 from the USDA database, which including 7,618 foods and 65 macronutrients. The USDA National Nutrient Database for Dietary Studies is the major source of food composition data in the United States. To be consistent with the version used in DMAS study, we chose the version 2011-2012 to construct reference Food-Nutrient network (FNN). This version includes 7,618 foods, which can be clarified into 9 highest items and the total number of nutrients is 65.

Remark 1: In calculating the nutritional profiles (Fig.1) as well as the nutritional redundancy (Fig.4), we excluded energy and water for the following considerations. First, energy does not have a unit of mass, and hence cannot be included in the nutritional profile where components represent relative abundances. Consequently, it cannot be used to calculate the nutritional redundancy either. Second, water was not considered as a nutrient in the Harvard Food Composition Database (HFDB). For consistency and comparison purposes, we also removed it from FNDDS when we calculate the nutritional profiles and nutritional redundancy from the DMAS, WLVS, MLVS data.

1.1.2 Frida database

The database Frida Food Data (frida.fooddata.dk) was created and published by the National Food Institute, Technical University of Denmark (DTU), including data on nutrient content of various foods. We used the version released at 08-02-2019, which includes 1,185 foods items and 198 nutrients.

1.1.3 Harvard Food Composition Database

We used the Harvard food composition table of year 2015 to construct a reference food-nutrient network to calculate the nutritional redundancy of participants from the NHS. The Harvard food composition table consists of 575 foods and 182 nutrients. To calculate the nutritional redundancy, we removed calories due to different unit and total protein, total fat and total sugar since those nutrients are overlapped with some sub-nutrients.

1.1.4 Other databases

Other databases, for example, FooDB, a database representing the most comprehensive effort to integrate food composition data from specialized databases and experimental data, has provided the information of 26,625 distinct bio-chemicals in foods. PhenolExplorer and eBasis have also resulted in wealth of information on food composition. Throughout our analysis, we focused on the nutrient level, rather than composition and compounds levels, thus we reference FNN are constructed by USDA and Frida databases.

1.2 Nutrition profile analysis

The nutritional composition of foods for DMAS was determined using ASA24-2016. Both of WLVS and MLVS composition of foods for DMAS was determined using ASA24-2012. ASA24 assigns nutrient information to foods using the USDA's Food and Nutrient Dietary Database (FNDDS). Subjects reported dietary intake as food records and entered their own dietary records directly into ASA24. To calculate the relative abundance of nutrient, we removed energy (unit is not in mass) and water (the water information is not included in NHS database, so we removed to be consistent).

1.3 Food Choice Analysis

Foods of DMAS, WLVS and MLVS were categorized according to their FNDDS food code and modification code as assigned by ASA24. The foods of NHS were mapped from serving data to the Harvard food composition table according to the food descriptions.

2. Healthy diet scores

2.1 HEI-2005

HEI-2005[1] is a score that measures adherence to the USDA 2005 Dietary Guidelines for Americans. The score range is 0 to 100. Each of the 12 components has a minimum score of zero and a maximum score of 5, 10 or 20. These components are: Total vegetables, Dark green & orange vegetables, Total fruit, Whole fruit, Total grains, Whole grains, Milk, Oils, Saturated fat, Sodium and SoFAAs.

2.2 AHEI-2010

AHEI-2010[1] is a score that measures adherence to a diet pattern based on foods and nutrients most predictive of disease risk in the literature. The minimum score = 0, maximum score = 110. Each of the 11 components has a minimum score of 0 and a maximum score of 10, as outlined in the table below. A score between the minimum and maximum is assigned on a continuous basis (except for sodium and alcohol). Those 11 components are: Vegetables, Fruit, Whole grains, Sugar-sweetened beverages and fruit juice, Nuts and legumes, Red meat and processed meat, Trans fat, Long-chain fats, Poly-unsaturated fatty acids, Sodium and Alcohol.

2.3 AMED

The components of AMED[2] are vegetables (excluding potatoes), fruits, nuts, whole grains, legumes, fish, ratio of monounsaturated to saturated fat, red and processed meats, and alcohol. The range of score is 0 to 9. The score criteria are: Intake above the FFQ-specific median intake received 1 point for vegetables, fruits, nuts, whole grains, legumes, fish, and M:S ratio; otherwise, they received 0 points. Red and processed meat consumption below the FFQ-specific median

received 1 point; otherwise, 0 points. Alcohol intake between 5 and 15 g/d for women and 10-25 g/d for men received 1 point; otherwise, 0 point.

2.4 DASH

This score was created to capture the characteristics of the Dietary Approaches to Stop Hypertension diet. The DASH[3] Components includes fruits, vegetables (excluding potatoes), nuts and legumes, low-fat dairy products, whole grains, sodium, sweetened beverages, red and processed meats. The score range is 8 to 40. The DASH Scoring Criteria is each food group first classified into FFQ-specific quintiles. For fruits, vegetables, nuts and legumes, low-fat dairy products, and whole grains, the score for that food group is the quintile ranking. i.e., quintile 1 is assigned 1 point and quintile 5, 5 points. For sodium, red and processed meats, and sweetened beverages, low intake is best. Therefore, the lowest quintile was given a score of 5 points and the highest quintile, 1 point.

3. Healthy aging prediction

We used two standard classifiers: RF (Random Forest, R package 'randomForest'[4]) and XGBoost (Extreme gradient boosting decision trees, R package 'xgboost'[5]) to predict the healthy aging status. The base learners of RF are decision trees. Each tree is a non-linear model constructed with many linear boundaries. A node in a decision tree is associated with a question asking about the data based on the value of a particular feature. XGBoost is a scalable end-to-end decision tree boosting system[5]. Unlike RF that applies the technique of bootstrap aggregating (i.e., bagging) to tree learners, the trees of a boosting system are built sequentially: each tree aims to reduce the error of its previous tree.

For hyperparameter tuning, we used the R package 'caret'[6] (Classification And REgression Training). The number of features randomly sampled as candidates at each split range from 1 to 15 and the number of trees to grow is fixed to 500 for RF. The parameter ranges for XGBoost are the following: (1) Learning rate (eta): 0.001, 0.01; (2) Number of features supplied to a tree (colsample_bytree): 0.4, 0.6, 0.8, 1.0; (3) The depth of the tree (max_depth): 4, 6, 8, 10, 100,000; (4) Maximum number of iterations (nrounds): 100, 1000; (5) Regularization (gamma): 0; (6) Minimum sum of instance weight (min_child_weight): 1; (7) Number of samples supplied to a tree (subsample): 0.5, 0.75, 1; (8) Number of trees to grow (ntree): 500.

To overcome the label imbalance issue, we used the downSample function in caret, which will randomly sample a data set so that all classes have the same frequency as the minority class. To compare the performances of NR and the four healthy diet scores, we split the data into 80% percentage of sample as training set and the remaining 20% samples as test set. For each splitting, we used one of NR, HEI-2005, AHEI-2010, AMED and DASH together with other confounding factors to train the model, then validate the classifier using the test set. We used the error rate, i.e., the proportion of participants that have been incorrectly classified by the model and AUC (area under the ROC curve) to quantify the performance.

4. Supplementary figures and tables

Figure S1: Nutritional profiles are highly conserved across individuals while food profiles are highly personalized. The Bray-Curtis dissimilarity (column-1), rJSD (rooted Jensen-Shannon divergence, column-2), Yue-Clayton distance (column-3) and 1-Spearman correlation (column-4) between the food profiles of the same individuals but different time points (intra-individual) and food profiles among different individuals (inter-individual) at single food level (a1-a4) and nine major food groups level (b1-b4) and nutrient profiles (c1-c4). The Bray-Curtis dissimilarity (column-1), rJSD (column-2), Yue-Clayton distance (column-3) and 1-Spearman correlation (column-4) between the food (or nutritional) profiles of different individuals and different time

points at the single food level (d1-d4) and nine major food group level (e1-e4). The Bray-Curtis dissimilarity between a pair of individuals, j and k is defined as: $BC_{jk} \equiv \frac{\sum_i |X_{ij} - X_{ik}|}{\sum_i (X_{ij} + X_{ik})}$. The rJSD dissimilarity is defined as: $D_{\text{rJSD}}(j,k) \equiv [\frac{D_{\text{KL}}(j,m) + D_{\text{KL}}(k,m)}{2}]^{1/2}$, in which $m \equiv \frac{j+m}{2}$ and $D_{\text{KL}}(j,k) \equiv \sum_{i \in S} X_{ij} \log \frac{X_{ij}}{X_{ik}}$ is the Kullback-Leibler divergence between j and k. The Yue-Clayton dissimilarity is defined as: $YC_{jk} \equiv \frac{\sum_i X_{ij} X_{ik}}{\sum_i \sum_i (X_{ij} - X_{ik})^2 + \sum_i (X_{ij} * X_{ik})}$. In all dissimilarity definitions, X_{ij} represents the relative abundance of food/nutrient i in individual j. We only choose 100 participants in NHS and HPFS due to computational complexity. The boxplot represents all pairwise dissimilarity. Boxes indicate the interquartile range between the first and third quartiles with the central mark inside each box indicating the median. Whiskers extend to the lowest and highest values within 1.5 times the interquartile range.

Figure S2: The food-specific subgraphs of the FNN display similar topological feature as the original FNN. Columns: (1) the incident matrix; **(2)** distribution of nutrient distance; **(3)** distribution of food degree; **(4)** distribution of nutrient degree. **Rows:** (**a**) Milk and Milk Products; (**b**) Meat_Poultry, Fish, and Mixtures; (**c**) Eggs; (**d**) Dry Beans, Peas, Other Legumes, Nuts, and Seeds; (**e**) Grain Products; (**f**) fruits; (**g**) Vegetables; (**h**) Fats, Oils, and Salad Dressings; (**i**) Sugars, Sweets, and Beverages.

Figure S3: Food-Nutrient networks (FNN) constructed from the Harvard food composition database (a) and the Frida database (b). Columns: (**1**) the incident matrix of FNN, where the presence (or absence) of a link between a food and a nutrient is colored in green (or pink), respectively. (**2**) The distribution of nutrient distances (d_{ij}) among different foods. The bin size is 0. (**3**) The distribution of food degrees. Here, the degree of a food item is the number of distinct nutrients it can provide. (**4**) The distribution of nutrient degrees. Here, the degree of a nutrient is the number of food items that contain this nutrient.

Figure S4: Raw food-Nutrient networks (FNN) constructed from the Harvard food composition database. We organized this matrix using the Nestedness Temperature Calculator to emphasize its nested structure[6].

Figure S5: Food-Nutrient networks (FNN) constructed from the FNDDS database after removing those nutrients that are not specific enough to have a SIMLES or InChIKey ID, e.g., sugar, total fat, protein, total fiber, etc. We organized this matrix using the Nestedness Temperature Calculator to emphasize its highly nested structure[6].

Figure S6: Comparison of structural properties between the real FNN and different randomized FNNs. **Columns**: FNN constructed from the USDA database **(a)**, the Harvard food composition database **(b)**, and the Frida database **(c). Rows: (1)** The nestedness based on the NODF measure of the real FNN (gray bar), as well as the randomized FNNs (colored bars) using four different FNN randomization schemes: Null-FNN-1, complete randomization; Null-FNN-2, Food-degree preserving randomization; Null-FNN-3, Nutrient-degree preserving randomization; Null-FNN-4, Food- and nutrient-degree preserving randomization. For each randomization scheme, 50 realizations were generated. **(2)** The distribution of nutrient distances (d_{ij}) between different foods calculated from the real FNN (gray lines) and the randomized FNNs (colored lines) using the same randomization schemes as in row (1). We generated 50 realizations for each randomization scheme, and the bin size is 0.02. All FDR-corrected P values were found using the paired and two-sided t-test. Significance level: FDR-corrected p value <0.0001(***).

Figure S7: Importance ranking of features quantified by their mean decreasing of accuracy. Here, we used the following features: Age, BMI (body mass index), ACT (physical activity), MDINC (median income from census tract), CALOR (energy intake), PKYR (pack-years of smoking), HIEDU (education), ASPD (aspirin use), MVT (multivitamin use), MARRIT (marital status), together with (**a**) NR (nutritional redundancy) or one of the healthy diet scores: (**b**) AMED; (**c**) AHEI; (**d**) DASH; and (**e**) HEI. The results were averaged over 100 different training/test splits. The error bars are almost invisible in those figures.

Fig. S8: Nutritional redundancy is robust in aging prediction with different Hill numbers. a: NR of NHS calculated with different definitions. GSI-Rao represents the NR calculated from Gini-Simpson index and Rao's quadratic entropy used in the main text. $q = 0,1,2,3$ represent the Hill-number based NR (see SI Sec.3.3). **b:** Error rate of the RF classifier in predicting the healthy aging status. **c:** AUROC of the RF classifier in predicting the healthy aging status. The participants are randomly spitted into 80% as the training set and the remaining 20% as the test set. The boxplot represents the performances of 200 independent splitting. Boxes indicate the interquartile range between the first and third quartiles with the central mark inside each box indicating the median. Whiskers extend to the lowest and highest values within 1.5 times the interquartile range.

Fig. S9: Nutritional redundancy is robust in predicting healthy aging with a different classifier.
a: Error rate of the XGBoost classifier in predicting the healthy aging status. **b:** AUROC of XGBoost in predicting the healthy aging status. The participants are randomly spitted into 80% as the training set and the remaining 20% as the test set. The boxplot represents the performances of 100 independent splitting. Boxes indicate the interquartile range between the first and third quartiles with the central mark inside each box indicating the median. Whiskers extend to the lowest and highest values within 1.5 times the interquartile range.

Figure S10: Nutrient redundancy serves as a potential metric to predict healthy aging in HPFS. a: Error rate of random forest classifier in the prediction of healthy aging status. **b:** AUROC of random forest classifier in prediction of healthy aging status. The participants are randomly spitted into 80% as the training set and the remaining 20% as the test set. The boxplot represents the performances of 100 independent splits. Boxes indicate the interquartile range between the first and third quartiles with the central mark inside each box indicating the median. Whiskers extend to the lowest and highest values within 1.5 times the interquartile range.

Figure S11: Spearman correlation between the food diversity (FD) and nutritional redundancy (NR) (or the normalized nutritional redundancy: nNR). DMAS (dietary intake data collected using ASA24 dietary assessment tool daily over 17 consecutive days); WLVS (with four ASA24 records within one year); MLVS (with four ASA24 records within one year); NHS (with FFQ administrated every four years and with total eight time points); HPFS (with FFQ administrated every four years and with total seven time points).

Figure S12: Removing fatty acids does not significantly alter NR. For participants in DMAS (a) and NHS (b, using 100 randomly selected subjects for simplicity), we calculated the Pearson correlation between their NR values calculated from the original food nutrient network (FNN) and the FNN with fatty acids-related nutrients (total fatty acids of saturated, total fatty acids of monounsaturated, and total fatty acids of polyunsaturated) removed. P-values were computed using two-sided Student's t-test.

Characteristics	Healthy agers ($n = 3,491$)	Usual agers ($n = 17,808$)
Mean age, years	58.1 (4.9)	62.8 (6.5)
Educational level (1992), % (n)		
Registered nurse	57 (1,999)	66 (11,841)
Bachelor's degree	25 (877)	21 (3,776)
Master or doctorate	18 (565)	13 (1,921)
Marital status (1996), % (n)		
Windowed	5 (182)	10 (1,792)
Married	86 (3,018)	82 (14,601)
Separated/divorced	7 (267)	8 (1,315)
Median neighborhood income, $		
<45302	16 (564)	21 (3,659)
45,302-55,373	17 (606)	20 (3,641)
55,373-66,363	20 (692)	20 (3,553)
66,364-82582	21 (746)	20 (3,506)
>82582	26 (873)	19 (3,414)
Body mass index, % (n)		
<22	26 (919)	16 (2,886)
22-24.9	35 (1,220)	27 (4,856)
25-29.9	30 (1,062)	35 (6,212)
≥ 30	9 (271)	22 (3,668)
Multivitamin use, % (n)	60 (2,108)	61 (10.921)
Aspirin use, tables per week, % (n)		
<1	67 (2,323)	60 (10,740)
1-2	8 (260)	7 (1,217)
>2	25 (869)	33 (5,656)
Pack-years of smoking		
0	54 (1,881)	48 (8,504)
0-4	13 (464)	11 (1,968)
5-20	20 (701)	20 (3,590)
21-112	12 (406)	20 (3,522)
Mean physical activity		
<3.6	11 (367)	21 (3,651)
3.7-9	16 (565)	21 (3,660)
9.1-16.7	21 (721)	21 (3,668)
16.8-30.4	24 (848)	20 (3,578)
>30.5	28 (986)	17 (3,214)
Energy intake		
<1,296	17 (601)	29 (3,318)
1,296-1,572	21 (737)	21 (3,683)
1,573-1,830	21 (716)	21 (3,754)
1,831-2,169	22 (748)	21 (3,724)
>2,170	19 (689)	18 (3,329)
Nutritional redundancy	0.331 (0.0791)	0.325 (0.0798)
AMED	4.814 (1.89)	4.48 (1.86)
AHEI	55.5 (11)	53.34 (10.77)
DASH	24.58 (4.49)	23.86 (4.48)
HEI	74.9 (9.16)	73.08 (10)

Table S1. Characteristics of a subset of NHS participants in 1998 for whom we have their aging status in 2012 from a previous study. Measures in this table were calculated among non-missing values (≤ 5 of data were missing). Data expressed as mean (SD) or percentage.

Table S2. Hazard ratios (95% confidence intervals) of type 2 diabetes according to quintiles of normalized nutritional redundancy (NR) in the Nurses' Health Study (1984-2014) and Health Professionals Follow-Up Study (1986-2016).

	Q1	Q2	Q3	Q4	Q5	P for trend[1]
			NHS			
Cases/Person-year	882/187,428	802/187,556	760/187,504	698/187,423	672/187,397	
Age-adjusted model	1 (reference)	0.90(0.82,0.99)	0.85(0.77,0.94)	0.75(0.68,0.83)	0.23(0.14,0.38)	<0.001
Multivariable-adjusted model[2]	1 (reference)	0.91(0.83,1.01)	0.92(0.83,1.02)	0.89(0.80,0.99)	0.64(0.39,1.06)	0.0827
			HPFS			
Cases/Person-year	419/88,643	351/88,705	317/88,792	274/88,848	327/88,844	
Age-adjusted model	1 (reference)	0.83(0.72,0.96)	0.75(0.65,0.87)	0.77(0.66,0.89)	0.26(0.14,0.50)	<0.001
Multivariable-adjusted model[2]	1 (reference)	0.88(0.76,1.02)	0.82(0.70,0.95)	0.88(0.76,1.03)	0.50(0.260.97)	0.0415

[1]P for trend was calculated using the median value of each quintile (two-sided Chi-square test).

[2]Multivariable-adjusted model adjusted for age (months), total energy intake, ethnicity (white, African American, Asian, others), body mass index (<21.0, 21.0-22.9, 23.0-24.9, 25.0-26.9, 27.0-29.9, 30.0-32.9, 33.0-34.9, or ≥35.0 kg/m²), smoking status (never smoked, past smoker, currently smoke 1-14 cigarettes per day, 15-24 cigarettes per day, or ≥25 cigarettes per day), alcohol intake (0, 0.1-4.9, 5.0-9.9, 10.0-14.9, 15.0-29.9, and ≥30.0 g/d), hypertension (yes, no), hypercholesterinemia (yes, no), multivitamin use (yes, no), physical activity (quintiles), alternative healthy eating index, total energy (quintiles), family history of myocardial infarction, postmenopausal hormone use (never, former, or current hormone use, or missing), and oral contraceptive use.

Table S3. Hazard ratios (95% confidence intervals) of cardiovascular disease according to quintiles of normalized nutritional redundancy (NR) in the Nurses' Health Study (1984-2014) and Health Professionals Follow-Up Study (1986-2016).

	Q1	Q2	Q3	Q4	Q5	P for trend[1]
			NHS			
Cases/Person-year	927/195,361	814/195,530	809/195,422	731/195,286	770/195,193	
Age-adjusted model	1 (reference)	0.90(0.82,0.99)	0.90(0.82,0.99)	0.84(0.76,0.92)	0.40(0.25,0.63)	<0.001
Multivariable-adjusted model[2]	1 (reference)	0.91(0.83,1.00)	0.93(0.84,1.02)	0.88(0.80,0.98)	0.53(0.33,0.86)	0.0096
			HPFS			
Cases/Person-year	848/87,283	774/87,340	735/87,409	765/87,432	811/87,411	
Age-adjusted model	1 (reference)	0.91(0.83,1.01)	0.86(0.78,0.95)	0.88(0.80,0.97)	0.58(0.38,0.88)	0.0106
Multivariable-adjusted model[2]	1 (reference)	0.93(0.84,1.03)	0.89(0.81,0.99)	0.91(0.83,1.01)	0.69(0.45,1.05)	0.0858

[1]P for trend was calculated using the median value of each quintile (two-sided Chi-square test).
[2]Multivariable-adjusted model adjusted for age (months), total energy intake, ethnicity (white, African American, Asian, others), body mass index (<21.0, 21.0-22.9, 23.0-24.9, 25.0-26.9, 27.0-29.9, 30.0-32.9, 33.0-34.9, or ≥35.0 kg/m^2), smoking status (never smoked, past smoker, currently smoke 1-14 cigarettes per day, 15-24 cigarettes per day, or ≥25 cigarettes per day), alcohol intake (0, 0.1-4.9, 5.0-9.9, 10.0-14.9, 15.0-29.9, and ≥30.0 g/d), hypertension (yes, no), hypercholesterinemia (yes, no), multivitamin use (yes, no), physical activity (quintiles), alternative healthy eating index, total energy (quintiles), family history of myocardial infarction, postmenopausal hormone use (never, former, or current hormone use, or missing), and oral contraceptive use.

Table S4. Hazard ratios (95% confidence intervals) of type 2 diabetes according to terciles of FD in the Nurses' Health Study (NHS, 1984-2014) and Health Professionals Follow-Up Study (HPFS, 1986-2016).

NHS	T1	T2	T3	P for trend[1]
Cases/Person-year	1,300/312,451	1,279/312,504	1,235/312,353	
Age-adjusted model	1 (reference)	0.98(0.91,1.06)	0.94(0.87,1.02)	0.1579
Multivariable-adjusted model[2]	1 (reference)	1.04(0.96,1.12)	1.01(0.93,1.09)	0.8529
HPFS				
Cases/Person-year	622/147,862	526/147,989	540/147,980	
Age-adjusted model	1 (reference)	0.84(0.75,0.95)	0.88(0.78,0.98)	0.0136
Multivariable-adjusted model[2]	1 (reference)	0.91(0.81,1.02)	0.99(0.88,1.12)	0.5966

[1]P for trend was calculated using the median value of each terciles (two-sided Chi-square test).

[2]Multivariable-adjusted model adjusted for age (years), ethnicity (white, African American, Asian, others), body mass index (<21.0, 21.0-22.9, 23.0-24.9, 25.0-26.9, 27.0-29.9, 30.0-32.9, 33.0-34.9, or ≥35.0 kg/m[2]), smoking status (never smoked, past smoker, currently smoke 1-14 cigarettes per day, 15-24 cigarettes per day, or ≥25 cigarettes per day), alcohol intake (0, 0.1-4.9, 5.0-9.9, 10.0-14.9, 15.0-29.9, and ≥30.0 g/d), hypertension (yes, no), hypercholesterinemia (yes, no), multivitamin use (yes, no), physical activity (quintiles), alternative healthy eating index, family history of diabetes. In NHS, postmenopausal hormone use (never, former, or current hormone use, or missing) and oral contraceptive use were additionally adjusted.

Table S5. Hazard ratios (95% confidence intervals) of cardiovascular disease according to terciles of FD in the Nurses' Health Study (NHS, 1984-2014) and Health Professionals Follow-Up Study (HPFS, 1986-2016).

NHS	T1	T2	T3	P for trend[1]
Cases/Person-year	1,438/325,560	1,339/325,704	1,274/325,527	
Age-adjusted model	1 (reference)	0.95(0.88,1.02)	0.92(0.85,0.99)	0.0307
Multivariable-adjusted model[2]	1 (reference)	0.98(0.91,1.06)	0.97(0.90,1.05)	0.4760
HPFS				
Cases/Person-year	1,374/145,508	1,265/145,700	1,294/145,667	
Age-adjusted model	1 (reference)	0.92(0.85,0.99)	0.92(0.86,1.00)	0.0315
Multivariable-adjusted model[2]	1 (reference)	0.95(0.88,1.02)	0.97(0.89,1.05)	0.3418

[1]P for trend was calculated using the median value of each terciles (two-sided Chi-square test).

[2]Multivariable-adjusted model adjusted for age (years), ethnicity (white, African American, Asian, others), body mass index (<21.0, 21.0-22.9, 23.0-24.9, 25.0-26.9, 27.0-29.9, 30.0-32.9, 33.0-34.9, or ≥35.0 kg/m^2), smoking status (never smoked, past smoker, currently smoke 1-14 cigarettes per day, 15-24 cigarettes per day, or ≥25 cigarettes per day), alcohol intake (0, 0.1-4.9, 5.0-9.9, 10.0-14.9, 15.0-29.9, and ≥30.0 g/d), hypertension (yes, no), hypercholesterinemia (yes, no), multivitamin use (yes, no), physical activity (quintiles), alternative healthy eating index, family history of myocardial infarction. In NHS, postmenopausal hormone use (never, former, or current hormone use, or missing) and oral contraceptive use were additionally adjusted.

References

1. Chiuve, S. E. *et al.* Alternative dietary indices both strongly predict risk of chronic disease.

   The Journal of nutrition **142**, 1009–1018 (2012).

2. Fung, T. T. *et al.* Mediterranean diet and incidence and mortality of coronary heart disease

   and stroke in women. *Circulation* **119**, 1093 (2009).

3. Fung, T. T. *et al.* Adherence to a DASH-style diet and risk of coronary heart disease and

   stroke in women. *Archives of internal medicine* **168**, 713–720 (2008).

4. Liaw, A. & Wiener, M. Classification and regression by randomForest. *R news* **2**, 18–22 (2002).

5. Chen, T. & Guestrin, C. XGBoost: A Scalable Tree Boosting System. *Proceedings of the 22nd

   ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* 785–794 (2016)

   doi:10.1145/2939672.2939785.

6. Kuhn, M. Building predictive models in R using the caret package. *Journal of statistical

   software* **28**, 1–26 (2008).