Mathematical model reveals that heterogeneity in the number of ion transporters regulates the fraction of mouse sperm capacitation

Alejandro Aguado-García[1,2,*], Daniel A. Priego-Espinosa[2,3,*], Andrés Aldana[4], Alberto Darszon[4,**], and Gustavo Martínez-Mekler[2,5,**]

[1]Instituto de Ciencias Básicas y Aplicadas, Universidad Autónoma del Estado de Morelos, Cuernavaca, México

[2]Instituto de Ciencias Físicas, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, México

[3]Department of Biology, University of Kentucky, Lexington, KY, USA

[4]Instituto de Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, México

[5]Centro de Ciencias de la Complejidad, Universidad Nacional Autónoma de México, CDMX, México

[*]These authors contributed equally to this work.

[**]Correspondence: G M-M: mekler@icf.unam.mx, A D: darszon@ibt.unam.mx

February 5, 2021

Abstract

Capacitation is a complex maturation process that mammalian sperm must undergo in the female genital tract to be able to fertilize an egg. This process involves, amongst others, physiological changes in flagellar beating pattern, membrane potential, intracellular ion concentrations and protein phosphorylation. Typically, in a capacitation medium, only a fraction of sperm achieve this state. The cause for this heterogeneous response is still not well understood and remains an open question. Here, one of our principal results is to develop a discrete regulatory network, with mostly deterministic dynamics in conjunction with some stochastic elements, for the main biochemical and biophysical processes involved in the early events of capacitation. The model criterion for capacitation requires the convergence of specific levels of a select set of nodes. Besides reproducing several experimental results and providing some insight on the network interrelations, the main contribution of the model is the suggestion that the degree of variability in the total amount and individual number of ion transporters among spermatozoa regulates the fraction of capacitated spermatozoa. This conclusion is consistent with recently reported experimental results. Based on this mathematical analysis, experimental clues are proposed for the control of capacitation levels. Furthermore, synergistic and interference traits that become apparent in the modelling among some components also call for future theoretical and experimental studies.

Author Summary

Fertilization is one of the fundamental processes for the preservation of life. In mammals sperms undergo a complex process during their passage through the female tract known as capacitation which enables them for fertilization. At the present time it is accepted from experimental observation, though not understood, is that only a fraction of the sperm is capacitated. In this work, by means of a network mathematical model for regulatory spermatozoa intracellular signaling processes involved in mice capacitation, we find that the variability in the distribution of the number of ion transporters intervenes in the regulation of the capacitation fraction. Experimental verification of this suggestion could open a line of research geared to

the regulation of the degree of heterogeneity in the number of ion transporters as a fertility control. The model also uncovers through in silico hyperactivation, loss of function and knockout of network elements, synergetic traits which again call for experimental verification.

1   Introduction

Sperm are highly specialized cells whose purpose is to reach, recognize and fuse with the egg. Their fundamental goal is to deliver the paternal genetic material to a female gamete. Despite lacking the machinery necessary for gene expression, these motile cells are capable of responding to changes in extracellular cues by means of intracellular electrochemical signals and phosphorylation cascades.

In the particular case of mammalian fertilization, sperm must reside inside the female genital tract for minutes to hours to complete a unique maturation process during which they acquire the adequate swimming modality to reach the egg at the right time and place, and the ability to fuse with it. In the early 50's, Austin and Chang identified the essential changes required for sperm to be able to fertilize eggs and called these processes "capacitation" [1, 2]. In 1957, *in vitro* fertilization (IVF) was performed with epididymal mice sperm exposed to a chemically defined medium (capacitation medium) [3]. In this procedure, multiple signaling events are involved which depend on a set of ion transporters located along sperm membranes, which are synthesized during spermatogenesis as they differentiate before becoming transcriptionally and translationally silent [4].

It is known that when murine sperm are exposed to a capacitation medium with the proper levels of $CaCl_2$, NaCl, $NaHCO_3$, KCl, cholesterol acceptors and metabolites, among others, only a fraction of the cell population is able to undergo a physiological secretory event known as acrosome reaction, which is necessary for gamete fusion. This reaction is proof that the sperm has become capacitated [5–7]. The causal relationships that originate the heterogeneous capacitation response remain poorly understood, furthermore the possible advantage of this heterogeneity remains speculative at present.

During mouse sperm capacitation, the membrane potential (V) changes from resting to hyperpolarized [4], the intracellular calcium concentration ($[Ca^{2+}]_i$) [8], pH (pH_i) [4] and chloride concentration ($[Cl^-]_i$) increase [9], whereas intracellular sodium concentration ($[Na^+]_i$) decreases [10]. Additionally, phosphorylation by PKA initiates within the first minutes of capacitation, whereas tyrosine phosphorylation starts at significantly later stages [4, 11, 12]. According to the molecular nature of each of these physiological changes, the signaling processes involved in capacitation occur at different time scales, (e.g. ion channel gating is in the subseconds whereas phosphorylation events can take from minutes to hours), hence the importance of asynchronous regulation. For the purposes of this paper, the hallmark of a capacitated state will be given in terms of the levels of the values reached by certain variables, namely, V, $[Ca^{2+}]_i$, pH_i and PKA.

Capacitation is a complex multifactorial transformation of the spermatozoa that takes place during their passage through a physical and chemical changing environment that regulates a network of linked intraflagellar biochemical and electrophysiological processes leading to a capacitated state. In order to contribute to a better comprehension of this phenomenon we construct, to our knowledge for the first time, a mathematical model for the dynamics of such an interacting regulatory network restricted to the early stages of capacitation. This is in itself one of the main contributions of this study. The model is firmly rooted on experimental evidence. It is a model where the components of the signaling regulatory network take discrete values. Given that, temporality in such a system is intrinsically asynchronous, we deal with it by working with a deterministic synchronous updating, subject to the inclusion of specific stochastic dynamics, that effectively introduces delays in a set of selected components of the network.

In this work we address the observation that typically only a fraction of murine sperm are capacitated *in vitro*, by looking into the heterogeneity in ion transporter numbers among sperm, acquired as a result of variable gene expression during spermatogenesis, exosome incorporation during epididymal maturation and proteolytic activity while capacitating [13–15]. With the model we can explore different levels of variability in the number of each functional ion transporter, which affects their relative number at the individual cell level, and investigate how the distribution properties of this variability influence the fraction of capacitated sperm [16]. This introduces another stochastic element in the modeling. The concept of such variability has been reported before [17–20] and is consistent with recent results indicating the presence of heterogeneous sperm population displaying differences in their $[Ca^{2+}]_i$ responses to external stimuli and in their membrane potential [21–23]. Our study provides us with a distribution parameter, the standard deviation, that can act as a control parameter on which

to focus future experimentation and search for physiological explanations. Additionally, the model exhibits the role of key components of the network and of their interrelations on capacitation. Within this framework, typical capacitation percentages can be controlled by single or double perturbations (either loss of function or overactivation) on selected signaling network elements, i. e. *in silico* mutations.

This paper is structured as follows. The results section has two main subsections, the first devoted to the building of the capacitation regulatory network, its topology and most importantly a mathematical model for its dynamics, the second centered on findings stemming from the model, in particular on the heterogeneous capacitation response and suggestions for capacitation control in terms of model parameters. After the discussion section, there is a methods section with detailed information and explanation on the construction of regulatory functions of the network and the conditions for the implementation of the reported simulations. Supplementary figures are included in the Appendix.

2 Results

2.1 Mathematical Model building

Experimental biological observations often reveal interrelations that can be integrated into an interacting web which we shall refer to as a "biological model". Here we present such a biological model for the regulatory pathways leading to capacitation. A further development is to quantify the interactions and build a corresponding mathematical model capable of a time dependent characterization. In our study, the mathematical model we chose is a logical network where the participating components take discrete values and time is also discrete. Central to this formulation is the construction of the regulatory relations which define the dynamics of the model.

In Figure 1 we show schematically the biological machinery involved in capacitation, which incorporates the main features mentioned in the introduction, with ion transporters, from both the midpiece and principal piece, along with the respective ions fluxes reported in [4], as well as early phosphorylation elements, which pertain to PKA activity [24].

From Figure1 we construct the network topology shown in Figure 2 by connecting the elements of interest for capacitation, based on molecular relationships established in the literature. The resulting network layout is a representation of the biological diagram (Figure 1) with the addition of auxiliary elements: those that help to represent explicitly the electromotive forces that generate ion fluxes through each type of ion transporter, those related to net ion fluxes, as well as recovery nodes. This is the starting point for our mathematical model for which time evolution will be dictated by the definition of regulatory rules that control the network dynamics.

2.1.1 Network dynamics

In order to characterize the dynamics of the signaling pathway related to the early capacitation response, we model each of the components and interactions in terms of a discrete dynamical network (DDN). The configuration of this kind of networks is given by a set of N discrete variables, $\{\sigma_1, \sigma_2, ..., \sigma_N\}$, which represent the dynamical state of all the network elements (nodes). The network configuration evolves in discrete time steps $t \in \mathbb{N}$ according to a set of dynamical rules that update the state of each node. One of most studied and used formalism within discrete dynamical networks is the Boolean network approximation. Such a network was proposed by Kauffman [25] as a modelling framework to study the dynamics of metabolic and genetic regulatory systems, with the underlying hypothesis that each particular temporal pattern can be associated to a cell phenotype. These type of models have been shown to be successful in the study of many biological networks, e. g. [26–30]. According to the Kauffman formalism, the dynamical state of a given node σ_i at time $t + 1$ is determined by the state of its own set of k regulators $\{\sigma_{i_1}, \sigma_{i_2}, ..., \sigma_{i_k}\}$ at time t as follows:

$$\sigma_i(t+1) = F_i[\sigma_{i_1}(t), ..., \sigma_{i_k}(t)] \tag{1}$$

where F_i is a regulatory function that assigns a value from a discrete state repertoire to a node σ_i. Note that each node σ_i has its own regulatory function F_i, which is built to capture qualitatively the inhibitory/activating character of the interactions of its particular regulator set. This function can be encoded through a truth table or expressed in terms of Boolean algebra equations [31]. In the most simple variant, the dynamical state of all nodes is updated synchronously by applying their respective regulatory function simultaneously at each iteration, (Eq. 1).

Figure 1: Representation of the capacitation machinery considered in this biological model. This figure is based on Ref. [4] Figure 6, with some modifications and additions. The gray bands represent the flagellum membrane. The gray semicircle represents internal calcium reservoirs. Inserted inside the light blue background (intracellular medium), we can see from left to right circles representing ion concentrations (green) and membrane potential (V, red). From left to right inserted inside the upper gray bands: cholesterol acceptor (ChAcc), sodium/calcium exchanger (NCX), calcium pump (PMCA), voltage/pH dependent calcium channel (CatSper), voltage/pH dependent potassium channel (Slo3), electroneutral but voltage dependent sodium/proton exchanger (sNHE). From up to down, inserted inside the gray semicircle (Redundant Nuclear Envelope, RNE): calcium dependent IP$_3$ Receptor calcium channel (IP$_3$R), calcium pump from the redundant nuclear envelope (SPCA). From left to right, inserted inside the inferior gray bands: electrogenic sodium/bicarbonate co-transporter (NBC), electroneutral chloride/bicarbonate exchanger (SLCAE), electrogenic chloride/bicarbonate exchanger (SLC26), chloride channel (CFTR/TMEM16A) and sodium channel (ENaC/TRP), soluble adenylate cyclase (sAC), (PDE) phosphodiesterase, (cAMP) cyclic adenosine monophosphate, (PKA) protein kinase A, (IP3) inositol triphosphate, Cap is a marker that represents the beginning of late capacitation. Due to scarcity of experimental information on chloride channels and sodium channels relevant to murine sperm capacitation, we modeled CFTR and ENaC only, which are the channels with more available experimental support.

Figure 2: Capacitation interaction network between elements considered in the biological model and additional auxiliary elements. Blue circles: extracellular ion concentrations. Yellow rectangles: gating of ion transporters. Pink circles: electromotive force of ion transporters ($\Delta\mu$, see methods 4.2.3) or flux of a given ion transporter. Light pink rectangles: total ion fluxes. Green circles: intracellular ion concentrations. Orange nodes: enzymes and molecules from cytosol. Red rectangle: joint node of select capacitation variables. Gray rectangles: auxiliary nodes for concentration and membrane potential recovery. Light pink triangle: membrane potential. Black arrows: positive regulation. Red arrows: negative regulation. Yellow arrows: dual regulation.

The Boolean model can be generalized to multi-state logical models, in which each node σ_i is allowed to take m states instead of only two:

$$\sigma_i(t) \in \tilde{e}_i = \{e_{i_1}, ..., e_{i_m}\} \tag{2}$$

where $\tilde{e}_i$ is the set of values σ_i can take, $e_{i_j} \in \mathbb{Z}$.

Most of the nodes in the capacitation network have two dynamical states. In the case of ion channels, the dynamical states portray: open (state 1) and closed (state 0). In the case of ion concentrations (Table 4), the dynamical states portray: high concentrations (state 1) and low concentrations (state 0). In the case of the activity of a given protein, the dynamical states portray: high activity (state 1) and low activity (state 0). In the case of current nodes, they have three dynamical states that represent the current direction: inward (state 1), null (state 0) and outward (state −1). In the case of membrane potential, the dynamical states portray: depolarized (state 1), resting (state 0) and hyperpolarized (state −1).

In general, there are many ways reported in the literature for building regulatory functions, e.g. truth tables with random outputs [25], Boolean threshold networks [32], majority rules [33], etc. The method for building regulatory functions in our model is *ad hoc*, related to thermodynamics and electrophysiological considerations explained in Sec. 4.2. The corresponding regulatory tables are shown in the supplementary files. Furthermore, the dynamical rules for integration nodes of ion fluxes and membrane potential were built in terms of an innovative scheme inspired in neural nets, explained in Sec. 2.1.4.

2.1.2 Multitemporal updating DDN

As pointed out in Sec. 1, it has been established experimentally that the time scales with which different nodes interact is diverse. In order to take this into account, in our model we define a time evolution equation in which we incorporate a stochastic interaction where each input of a given node is on average sampled at a different rate according to a biased Bernoulli processes (i. e. unfair coin tossing). The mean value of this process is calibrated to qualitatively approximate the operation rates of each kind of regulatory element shown in Sec. 4.3. With the above, we generalize the Kauffman multi-state model as follows:

$$\sigma_i(t+1) = F_i[B^t_{\tau_{i_1}}[\sigma_{i_1}(t)],...,B^t_{\tau_{i_k}}[\sigma_{i_k}(t)]], \tag{3a}$$

$$B^t_{\tau_{i_j}}\left[\sigma_{i_j}(t)\right] = \begin{cases} \sigma_{i_j}(t) & \text{if } b^t_{\tau_{i_j}} = 1 \\ 0 & \text{else} \end{cases} \tag{3b}$$

where $b^t_{\tau_{i_j}} \in \{0,1\}$ is a random variable from a biased Bernoulli process with mean $\tau_{i_j} \in [0,1]$, sampled at time t, and $\sigma_{i_j}(t) \in \tilde{e}_{i_j}$.

Given that PKA-dependent phosphorylation, cholesterol removal effects and calcium diffusion from the RNE to the principal piece, show characteristic times greater than other network processes (e. g. ion transport) [34, 35], we introduce a delay and persistence of their effects, a memory factor in the model i. e. they retain their value for a defined average number of steps. For these cases instead of equation 3b, the node evolution is given by:
for $t = 0$,

$$B^0_{\tau_{i_j}}\left[\sigma_{i_j}(0)\right] \equiv \sigma'_{i_j}(0) = \sigma_{i_j}(0), \tag{4}$$

and for $t > 0$,

$$B^t_{\tau_{i_j}}\left[\sigma_{i_j}(t)\right] \equiv \sigma'_{i_j}(t) = \begin{cases} \sigma_{i_j}(t) & \text{if } b^t_{\tau_{i_j}} = 1 \\ \sigma'_{i_j}(t-1) & \text{else} \end{cases}, \tag{5}$$

where $\sigma_{i_j}(t)'$ and $\sigma_{i_j}(t) \in \tilde{e}_{i_j}$. Notice that though the network model evolution is synchronous, the biological asynchronicity is dealt with in the model through this particular stochastic updating at the regulator interaction level.

2.1.3 Ion flux dynamics: Neural network scheme

In order to manage ion fluxes related to several types of transporters, we include ion flux integration nodes as auxiliary nodes that operate in a similar fashion as in neuron integrated signals coming from several inputs. Updating depends on whether a given threshold is surpassed. For theses cases, each total flux node σ_i evolves according to a regulatory function F_i that consists of a weighted sum of its regulators σ_{i_j}, as in the McCulloch&Pitts scheme [36]:

$$\sigma_i(t+1) = sgn\left[\sum_{j=1}^{k} w_{i_j}\sigma_{i_j}(t)\right] \tag{6}$$

wherein the *sgn* function discretizes the sum of regulators σ_{i_j} weighted by w_{i_j}, according to:

$$sgn(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0 \end{cases} \tag{7}$$

As in Sec. 2.1.2, we can generalize the neural network scheme to a multitemporal variant scheme as follows:

$$\sigma_i(t+1) = sgn\left[\sum_{j=1}^{k} w_{i_j}B^t_{\tau_{i_j}}[\sigma_{i_j}(t)]\right] \tag{8}$$

where the nodes σ_i now have a neuron-like regulatory function in which each regulator σ_{i_j} participates through a biased Bernoulli process. The values of the set of all weights of the network $\tilde{w} = \{w_{i_j}\}$ are based on biological evidence as it is shown in Sec. 4.3 .

2.1.4 Ion flux dynamics with heterogeneous ion transporter: Neural network scheme with weight variability at population level

In order to introduce variability in the total and relative number of ion transporters, we can generate populations of networks by modifying the set of weights $\tilde{w} = \{w_{i_j}\}$ of each network replicate. Every network r from the constructed population will have a set of weights

$\tilde{w}^r = \left\{ w_{i_j} G^r_{i_j}(1, D) \right\}$, where $G^r_{i_j}$ is a random number sampled from a Gaussian probability distribution $G(1, D)$ with mean 1 and standard deviation D, over an interval $[0, 2]$. In general, networks coming from such a population will have a different set of weights among them. It is at this stage that the second aforementioned stochasticity is included in the model. It acts on the values of the weights of regulators, on the links, not on the nodes themselves. Since, as shown in the methods section, each individual weight depends on the number of the corresponding ion transporter, the variability can be expressed in terms of number of ion transporters. We can extend this notion of variability to multitemporal-neuron-like regulatory functions F_i as follows:

$$\sigma^r_i(t+1) = sgn\left[\sum_{j=1}^{k} w_{i_j} G^r_{i_j}(1, D) B^t_{\tau_{i_j}}[\sigma_{i_j}(t)] \right] \tag{9}$$

With all these extensions of the Kauffman multi-state model and McCulloch&Pitts scheme, we have constructed a novel general updating scheme for the modeling of the early mouse sperm capacitation.

2.1.5 Ion flux integration nodes

Starting from Eq. 9 and adapting it to the particular case of ion flux integration nodes (I_{Ca}, J_H, I_{HCO3}, I_{Cl} and I_{Na}), we introduce auxiliary variables $\tilde{I}^r_i$, which integrate all the fluxes j of a particular ion i for a given sperm r. Each one of these auxiliary variables updates depending on the state of its particular ion transporter set at time t. In order to consider differences in operation rates among ion transporter I_{i_j}, we multiply each of them by a random variable $b^t_{\tau_j}$ from a biased Bernoulli process with mean τ_{i_j} at each time step t. Notice that this is a particular case of the scheme shown in Sec. 2.1.4. The parameter τ_{i_j} controls the average operation rate of I_{i_j}. The resulting dynamical rule is:

$$\tilde{I}^r_i(t+1) = sgn\left[\sum_{j=1}^{k} b^t_{\tau_{i_j}} w_{i_j} G^r_{i_j}(1, D) \Delta\mu_{i_j}(t) \right], \tag{10}$$

where $\Delta\mu_{i_j}$ is the electromotive force (emf) of flux j. $\Delta\mu_{i_j}$ will or will not contribute in the updating of $\tilde{I}^r_i$ at time $t+1$, depending on the random variable $b^t_{\tau_j}$, which can be either 0 or 1. Note that every spermatozoon r of the population has its own set of regulatory functions $\{F^r_i\}$ for the net ion fluxes $\tilde{I}^r_i$ which can differ from spermatozoon to spermatozoon. Also that regulator nodes $\Delta\mu_{i_j}(t)$ that represent emf associated with ion transporters j, and weights w_{i_j} would be the analog of effective conductances. The product of these terms gives an ion current expression based on Ohm's law. Regulator nodes $\Delta\mu_{i_j}(t)$ have a state repertoire defined by $\Delta\mu_{i_j}(t) \in \{-1, 0, 1\}$, where -1 corresponds to an emf that generates outward ion flux, 0 corresponds to null force, and 1 corresponds to an emf favoring ion influx. In Figure 3 we can see how a change in ion concentration x_i occurs through the net ion flux $\tilde{I}^r_i$ described above. Weights w_{i_j} and rates τ_{i_j} are reported in Sec. 4.3.

2.1.6 Membrane potential node

The membrane potential is a heavily linked core component of the network that in our model can take values from $\{-1, 0, 1\}$, which correspond respectively to hyperpolarized, resting and depolarized. For its time evolution we build a discrete version of the Hodgkin&Huxley equation [37] (shown in Eq. 25) through a discretizing partition defined by Eq. 27, such that the resting state is robust under small perturbations. As with the ion flux integration nodes, V is determined by a weighted sum of multiple sources: CatSper, Na$^+$-channel, Slo3, Cl$^-$-channel, the cotransporter NBC, the exchanger SLC26 and the leak current IL. The weight of the i-th contribution is given by its effective conductance. The last term of Eq. 11 is a consequence of Kirchhoff's law. Hence:

$$V^r(t+1) = \Theta\left[\sum_{j=V_1}^{V_k-1} z_j b^t_{\tau_j} w_j G^r_j(1, D) \Delta\mu_j(t) + w_{V_k} V^r(t) \right], \tag{11}$$

where w_{V_k} is the weight for the membrane potential at one time step before, and z_j is the charge sign of the reference ion in the current carried by transporter j. The latter quantity ensures that ion fluxes that are capable of modifying the membrane potential follow the typical convention used in electrophysiology, i.e. negative ion currents correspond to the influx of positive charges (e.g. CatSper, Na$^+$-channel), positive currents correspond to outward flux of positive charges (e.g. Slo3) or influx of negative charges (e.g. Cl$^-$-channel, NBC). The random

Figure 3: Diagram of the mechanism (time line) by which ion concentration x_i (bottom line) changes via net ion flux integration $\tilde{I}_i^r$ (middle line) determined by the weighted sum of all its linked electromotive forces (top line).

variable $b^t_{\tau_{i_j}}$ again is in charge of modeling different operation velocities of ion transporters. Weights w_j and velocities τ_j are reported in Sec. 4.3.

2.1.7 Operational definition of capacitation at the single-spermatozoon level

Most of the capacitation relevant variables are reported as isolated measurements of either population distributions or time series, i.e. there are very few available multivariate determinations in single spermatozoon [21, 38, 39]. This hinders the study of what changes occur simultaneously in a spermatozoon that reaches the capacitated state. Taking advantage of our modeling, we can set out explorations of the joint dynamics of changes involved in the early response. For this, we introduce an auxiliary node $\sigma_{\mathcal{J}}$, that integrates, at the single-cell level, the physiological changes associated with sperm capacitation, extrapolated from what is observed in bulk-cell measurements. For such a joint node, we consider as a hallmark for capacitation the following observable levels: hyperpolarized potential (state -1), high $[Ca^{2+}]_i$ (state 1), high pH_i (state 1) and high PKA activity (state 1). Though the levels of sodium and chloride are also determinant for capacitation, they are not included in $\sigma_{\mathcal{J}}$ due to the relative scarcity of experimental single-cell measurements.

Furthermore, in order to overcome the lack of sustained simultaneity in the capacitation conditions of individual sperm realizations, inherent to the stochastic nature of the network dynamics, we measure the joint dynamics with an observation window W. Thus, for a given spermatozoon r, its joint node $\sigma_{\mathcal{J}}^r(t)$ takes the state 1 at time t if the overlapping moving average with a window W of all its associated observables is above a threshold $\theta \in [0, 1]$. This means that $\sigma_{\mathcal{J}}^r(t)$ becomes active if along the preceding window W, each one of the above-mentioned variables is found in the state required for capacitation more than a certain fraction of time θ. If any of the variables fails, the joint node takes the state 0. In other words, $\sigma_{\mathcal{J}}^r(t)$ depends on the smoothed out values of our four selected variables, calculated at their respective window time-averages. The joint node threshold $\theta \in [0, 1]$ indicates what is the minimum percentage of time that the simultaneous conditions for capacitation needs to be fulfilled during an observation window. We define the dynamical rule of $\sigma_{\mathcal{J}}^r(t)$ as follows:

$$\sigma_{\mathcal{J}}^r(t|W, \theta) = \begin{cases} 1 & \text{if } \forall i \in \{V, Ca_i, pH_i, PKA\}, s_i \frac{1}{W} \sum_{\tau=1}^{W} \sigma_i^r(t-\tau) > \theta \\ 0 & \text{else} \end{cases} \tag{12a}$$

$$s_i = \begin{cases} -1 & \text{if } i = V \\ 1 & \text{else} \end{cases}, \tag{12b}$$

where s_i allocates the proper sign for the selected variable i. Summing up, the joint node tells us if, on average, the four selected variables were in their respective capacitation-associated states for a large enough fraction of time θ in an observation window W before the time t.

We will say that a single cell is capacitated if the steady state time-average of $\sigma_{\mathcal{J}}^r$ is above a threshold θ_c. Hence, we propose a capacitation criterion σ_{cap}^r in terms of the binary classifier Θ_c (Heaviside function), defined by:

$$\sigma_{\text{cap}}^r = \Theta_c \left[\frac{1}{L - t_r - 1} \sum_{t=t_r+1}^{L} \sigma_{\mathcal{J}}^r(t) \right] \tag{13a}$$

$$\Theta_c(x) = \left\{ \begin{array}{ll} 1 & \text{if } x \geq \theta_c \\ 0 & \text{else} \end{array} \right. , \tag{13b}$$

which is a discretization of the time average value of $\sigma_{\mathcal{J}}^r(t)$ during a post-stimulus observation time $L - t_r - 1$, after discarding a transient t_r required to reach steady state dynamics, where L is the total observation time. If $\sigma_{\text{cap}}^r = 1$, then the given sperm r will be considered capacitated.

2.1.8 Parameter summary

For the calibration and analysis of the dynamics of our network model, we can distinguish three kinds of parameters:

- Those that affect the individual (spermatozoon) network dynamics: w_{i_j}, a weight that determines the contribution (the influence) of regulator j on node i, ; τ_{i_j}, a kinetic parameter that sets a time scale that determines the influence rate of regulator j on node i.

- Those related to the network dynamics measurements that are instrumental for our capacitation operational definition (Sec. 2.1.7): W, an observation window of the moving average that sets the smoothing level of the joint node input variables; θ, a threshold of the observable average activity needed to turn on the joint node; θ_c, a threshold of the joint node average activity needed for a sperm to be classified as capacitated.

- Those that influence the spermatozoa network population: D, the standard deviation of the Gaussian distribution used to sample ion transporter counts.

2.2 Mathematical Model Findings

First we calibrate our model in order to qualitatively reproduce behaviors reported in the literature regarding electrophysiological determinations and early phosphorylation signals typically registered in sperm populations. Based on this systematic comparison we choose the values for the individual network dynamics parameters w_{i_j} and τ_{i_j} mentioned Sec. 2.1.8. Since in our work, we also look into blockage (loss-of-function mutants) by setting a mutated node to its least active state (0) and overactivation (gain-of-function mutants) by setting it to its most active state, we assess the compatibility between our simulations and the experimental results in unperturbed wild-type sperm (see Sec. 4.3) as well as mutants, when available.

Next, we investigate values of the joint node threshold θ, activity threshold θ_c, observation window W and variability D required to attain our operational definition of capacitation. Thereafter, we examine the effect of changes in the variability, taking D as a control parameter, on relevant variables for capacitation (Ca_i, pH_i, V and PKA) at the population level and explore ranges of parameter values required to reproduce the capacitation fraction reported in experiments. For a given set of such parameters we compare our simulation time averaged distributions of Ca_i, pH_i, V and PKA, for several values of D, for capacitated sperm subpopulations in contrast with non capacitated subpopulations. Finally, we illustrate how the capacitation fraction can be modified through perturbations in regulatory functions of the network, i.e. single and double *in silico* mutants. For all numerical simulations performed in this work, we used a population size of $N = 2 \times 10^3$ sperm, since from $N = 10^3$ onward the underlying behavior of numerical simulations does not change.

2.2.1 Model validation

To validate our model we test the effect of external stimulation on wild-type sperm (WT) and several loss-of-function (LOF) single mutants. Comparison of reported experimental measurements with their respective model counterpart was performed by means of the time series

averaged over a sperm population shown in Sec. S1.1. Results in absence of variability in the determination of the weight sets (w_{i_j}) that define the interactions in our network, $D = 0$, coincide with the variability case with $D = 0.25$. A summary of the response (for both cases) on WT conditions, coming from several stimuli (separate and joint increase in external bicarbonate and cholesterol acceptor) is shown in Table 1. The response with LOF cases of the select nodes under the joint stimulus with variability $D = 0.25$ is shown in table 2. All these tests contributed to the calibration of the model parameters.

Observables		Stimulus		
	HCO3$_e$		✓	✓
	ChAcc	✓		✓
	$\Delta_r^s\mathrm{Ca}_i$	$+^*$ $+$[40]	$+$ $+$[41]	$+$ $+$[8, 22]
	$\Delta_r^s\mathrm{Na}_i$	$-^*$ NA	$-$ NA	$-$ $-$[10]
	$\Delta_r^s\mathrm{pH}_i$	$+^*$ NA	$+$ $+$[42]	$+$ $+$[43]
	$\Delta_r^s\mathrm{V}$	0 0[42]	$+$ $+$[42]	$-$ $-$[44, 45]
	$\Delta_r^s\mathrm{Cl}_i$	0 NA	$+$ NA	$+$ $+$[9]
	$\Delta_r^s\mathrm{PKA}$	$+^*$ NA	$+$ NA	$+$ $+$[34]

Table 1: Summary of model validation on selected observables in wild-type networks. Each box contains a qualitative description of changes observed after a given stimulus (s) with respect to basal conditions (r): $\Delta_r^s(\sigma_i) \equiv \sigma_i^s - \sigma_i^r$. In the first column, only cholesterol acceptor (ChAcc) was added. In the second column, only bicarbonate was added. In the last column, both stimuli were added to the extracellular medium. For each box, the upper right corner corresponds to changes observed in time series averaged over a 2×10^3 simulated sperm population (e. g. Figure S2). Symbols $-$, 0 and $+$ refer to a decrease, no apparent change and an increase, respectively. In addition, lower corners contain references for that specific measurement, or NA if there is no available literature. Green indicates agreement between simulations and available experiments. The superscript $*$ refers to transient behavior.

| | LOF-network variant | | | | |
	CatSper	Slo3	Cl⁻-channel*	Na⁺-channel**	PKA
$\Delta_{\text{wt}}^{\text{lof}}\text{Ca}_i$	− / −[40]	0 / −[46]	0 / NA	0 / NA	− / −[41, 47]
$\Delta_{\text{wt}}^{\text{lof}}\text{Na}_i$	0 / NA	0 / NA	+ / NA	− / −[10]	+ / +[10]
$\Delta_{\text{wt}}^{\text{lof}}\text{pH}_i$	0 / NA	0 / NA	0 / −[48]	0 / NA	0 / NA
$\Delta_{\text{wt}}^{\text{lof}}\text{V}$	− / NA	+ / +[49, 50]	0 / +[9, 48]	− / −[51]	+ / +[52, 53]
$\Delta_{\text{wt}}^{\text{lof}}\text{Cl}_i$	0 / NA	+ / NA	0 / −[9]	0 / NA	0 / −[9]
$\Delta_{\text{wt}}^{\text{lof}}\text{PKA}$	− / NA	0 / 0[45]	0 / 0[54]	0 / NA	− / −[47]

(Row group label: Observables)

Table 2: Summary of model validation on selected LOF mutant networks. Each cell contains qualitative comparisons of the changes in a given observable, after a capacitating stimulus (see last column in Table 1, HCO$_e$+ChAcc), between sperm with a given blocked node and the reference wild-type ($\Delta_{\text{wt}}^{\text{lof}} = \sigma_i^{\text{lof}} - \sigma_i^{\text{wt}}$). The lower-left corner refers to experimental evidence changes, accompanied by references (if available), and the upper-right corner refers to changes corresponding to time series averaged over a 2×10^3 simulated sperm population with a variability of $D = 0.25$ in the distribution of ion transporters. Entries have the following meanings: NA accounts for cases where no literature is available for the particular observable under the specified node blockage; −, 0 and + correspond to a decrease, no apparent change and an increase with respect to the capacitation response in wild-type control. Colors refer to the level of agreement between experiment and simulations: full coincidence (green) and partial agreement (yellow). *The most likely candidates for Cl⁻-channels would be CFTR and/or TMEM16A. **The most likely candidates for Na⁺-channels would be ENaC and/or TRP.

2.2.2 How ion transporter numbers in single sperm impinge on the fraction of the population capacitated

Variability in the number of ion transporter counts is a known feature [55–57], here we explore the effect of modifying such variability on the capacitation response at the cell population level. For this we calculate the probability density of time-averaged variables among those most relevant for capacitation (our selected set V, Ca$_i$, pH$_i$ and PKA) during the capacitating stimulus for several variability distributions of weights w_{i_j} parametrized by their standard deviation D, shown in Figure 4. Given the connection between weights and the number of each ion transporter (Sec. 2.1.5, by modifying D, we are looking into the effect of changes in the variability of these numbers among sperm from the cell population modeled by a Gaussian distribution (Sec. 4.3).

Notice the following from Figure 4: The membrane potential probability density (red) widens drastically as a function of D, and becomes slightly bimodal at higher D values, showing a minor peak in positive values (centered around 0.1). Similarly, the calcium probability density (purple) becomes bimodal at higher D levels, with the second peak located at lower average Ca$_i$ values (centered around 0.37), however its upper boundary does not change with function of variability. Also, the pH$_i$ probability density (blue) widens drastically with the introduction of variability and changes from unimodal to bimodal. In contrast, PKA probability density (yellow) does not seem to be affected by the introduction of variability. D (Figure 4).

2.2.3 Capacitation criteria parameters that reproduce response heterogeneity

Here we explore the dependence of the capacitation percentage in population on the parameter values of the selected joint node time series: observation window W, capacitation time

Figure 4: Effect of variability in the probability density of time-averaged selected variables from stimulated 2×10^3-sperm populations. D is the numerical value of the standard deviation from the Gaussian distribution used to sample weight sets $\tilde{w}^r$ for each sperm r (Sec. 2.1.4).

overlap θ, signaling time averaged threshold θ_c (see (Eq. 12); and the population variability distribution width D of the amount of ion transporters. The first three parameters are instrumental for the determination of our operational definition of capacitation , they are model data detection quantities used to analyze the time-series generated by each individual sperm realization and intervene in the capacitation selection criteria proposed in Sec. 2.1.7; whereas D is at population level the standard deviation of the probability distribution of the set of weights $\tilde{w}^r = \left\{ w_{i_j} G_{i_j}^r (1, D) \right\}$ (see 2.1.4), and at single cell network dynamics it intervenes in the definition of the individual regulatory functions (Eq. 9). It is noteworthy to mention that variability D acts as a population control parameter.

Notice from the above that our operational capacitation definition contributes to: 1) discern stochastic fluctuations relevant to capacitation from non-relevant noise (parameter θ), 2) find what is the joint activity time interval of our selected variables (parameter W), 3) measure the average joint activity strength of our selected variables along observation time (parameter θ_c), and ultimately 4) allow us to discriminate between capacitated sperm and non-capacitated sperm.

After an extensive parameter value exploration we often encountered that for fixed values of θ: i) as the activity threshold θ_c is increased the maximum reachable capacitation fraction decreases, ii) there is global capacitation fraction maximum within a finite restricted range of W, iii) there are plateau of capacitation fraction as function of W and D with fixed θ_c.

In (Figure 5) we show for $\theta = 0.225$, variations of the capacitation fraction, as we modify D and W for several values of θ_c. When the latter parameter is 0.15, the figure exhibits a passage from non-capacitated population to capacitated, the peak of capacitation percentage is first attained around $D = 0.25$ with $W \approx 70$ and a plateau sets in for higher values of D within a window ranges size ($W \approx 10$ to $W \approx 200$), and capacitation fraction levels around $1/3$, typically reported in the literature [4] . This scenario is qualitatively retrieved in the Appendix figures S8 and S9 for different sets of parameter values. By adjustment of the joint node (θ) and joint activity (θ_c) thresholds, again, a well delimited region (plateau) can be achieved wherein several combinations of W and D values reproduce a capacitation fraction

similar to what is observed in bulk-cell experiments ($\approx 30\%$). For the remaining this paper we will be referring to the well delimited plateau region with parameter $\theta = 0.225$, $\theta_c = 0.15$, $W = 70$ $D = 0.25$.

Figure 5: Capacitated fraction (CF) in sperm populations, varying three parameters; observation window size (W, in iteration units), standard deviation of Gaussian variability (D) and activity threshold of joint node (θ_c). In all surfaces, each data point comes from 2×10^3-sperm sized populations with 2×10^4 iterations long simulations (5×10^3 iterations at resting state, 1.5×10^4 post stimulus iterations) and joint threshold $\theta = 0.225$. The classification of capacitated vs. non-capacitated sperm is applied on the last 10^4 time iterations of each individual sperm. The color bar represents the level of capacitation fraction. Note that the white zone corresponds to a capacitation level of $30 \pm 5\%$, which is close to the levels typically observed for *in vitro* capacitation in wild-type sperm. The chosen parameter set for the rest of the analyses ($D = 0.25, W = 70, \theta_c = 0.15, \theta = 0.225$) is indicated in purple (filled circle on the surface grid, filled diamond on the bottom heat map).

The point we want to highlight is the recurrence and robustness of this phenomenology. We have uncovered a procedure dependent on the degree of variability in the number of ion transporters, that leads to capacitation and that can regulate its fraction in population. Experimental corroboration of this model scenario is essential. Furthermore, the understanding of the underlying processes requires both experimental and theoretical work. On the theoretical side, the incorporation for the analysis of this "functional" transition of concepts, tools and methods for the study of physical and dynamical phase changes is enticing, on the experimental side stoichiometric considerations come to mind.

2.2.4 Histograms of time-averaged selected variables in sorted sperm sub-populations

After classifying sperm according to the above described criteria, we characterize for each sperm subpopulation (non-capacitated vs. capacitated) the distributions of time-averaged values of the selected variables: membrane potential (V, red), intracellular calcium (Ca$_i$,

purple), intracellular protons (pH$_i$, blue) and protein kinase A activity (PKA, yellow) in Figure 6. For Ca$_i$, pH$_i$ and PKA distributions, a slight increment is observed in their respective median values of capacitated sperm with respect to that of non capacitated sperm, whereas the width of those three distributions does not seem to change substantially. In contrast, V shows a decrease in the capacitated sperm distribution median value and width with respect to non capacitated sperm, both lower and upper bounds move to more negative values. There is a clearer separation between capacitated and non-capacitated sperm distributions, as compared to the other three variables. Interestingly, the introduction of variability induces bimodality in the distributions of membrane potential, calcium and pH$_i$ (Figure 4). These distribution shapes are reminiscent of those reported experimentally for membrane potential and calcium, measured by [22, 45].

Figure 6: Distributions of time-averaged variables relevant to capacitation in sperm population simulations under capacitating conditions. Sperm are sorted according to the classifier defined in Eq. 13 (Sec. 2.1.7). For each plot, the distributions of those sperm that did not reach the capacitated state (NC) are colored in gray. The capacitated fraction of sperm populations are labeled as C. The parameters used for the classification of sperm are: joint threshold $\theta = 0.225$, activity threshold $\theta_c = 0.15$, observation window $W = 70$ and variability $D = 0.25$. The distributions are calculated over 2×10^3-sperm populations. Each time series is averaged over a 1.5×10^4-iteration long capacitating stimulus, after discarding a 5×10^3-iteration long resting state. In order to compare the distributions, medians are indicated as dashed lines. These values are: V -0.040 (NC) and -0.122 (C), Ca$_i$ 0.444 (NC) and 0.454 (C), pH$_i$ 0.523 (NC) and 0.538 (C), finally PKA 0.434 (NC) and 0.440 (C).

Summing up, the capacitated subpopulation consists of sperm with higher hyperpolarization and moderate increases in Ca$_i$, pH$_i$ and PKA activity than in the non capacitated subpopulation. Notice that the probability densities reported in Figure 4, are compatible with the proportions of the capacitated and the non capacitated subpopulation in terms of the selected variable distributions shown in Figure 6 for D=0.25 .

2.2.5 Controlling capacitation fraction levels

In order to attain some insight on the relative importance of some proteins previously reported as relevant to capacitation, we analyze the response to changes in single and double mutants generated from a list of nodes of interest (Figure 7). The analysis is constrained to the proteins that are known to be expressed more specifically in the sperm flagella (sNHE, Slo3, CatSper, PKA) and those directly modifying electrochemical variables related to capacitation response (Na and Cl channels). In addition to the previously presented LOF scheme (Table

2), we explore another perturbation scheme: gain-of-function mutants (GOF), in which the node of interest is fixed to the most active state. After a capacitating stimulus, the resulting capacitation fraction in each mutant is depicted with a colored box. As reference, a population of WT sperm (unperturbed control) reaches approximately 30%d capacitation levels (green color delineates a $30 \pm 5\%$ region). The LOF-only and GOF-only scenarios are shown in the bottom-left and upper-right triangles, respectively, whereas mixed cases are contained in the upper-left quadrant. Notice that in the LOF-only mutant scenarios, the predominant effect is a general decrement of capacitation levels, except for individual ClC^{LOF}, NaC^{LOF} and their combinations.

Figure 7: Fraction of capacitated sperm determined by the model when one or two chosen nodes are modified: inactivated (LOF, state fixed to 0), over-activated (GOF, state fixed to 1). Results correspond to average values of the last 10^4 time steps of 1.5×10^4-iteration long simulations, performed over populations of 2×10^3 sperm, as in Figure 5. We used an observation window of size 70, Gaussian variability with D=0.25, joint node threshold $\theta = 0.225$ and activity threshold $\theta_c = 0.15$. The green zone indicates capacitation levels that are within the range of those typically observed for *in vitro* capacitation in wild-type sperm ($30 \pm 5\%$).

In the triangle of GOF-only network variants, the predominant effect is an increase of the capacitation levels, except for $CatSper^{GOF}$, ClC^{GOF} and NaC^{GOF} combinations.

The mixed LOF and GOF scenarios display a more complex pattern than the two above-mentioned types of mutants. More specifically, this kind of scenarios allow us to assess possible recovery or interference between nodes. For example, the model predicts that the over-activation of any of the selected nodes is not sufficient to recover from the capacitation fraction decrease produced by the blockage of either Slo3, CatSper or PKA. On the other hand, in a $sNHE^{LOF}$ background, Slo3 can totally recover the capacitation response, and even surpass the WT capacitation levels.

We remark that since the over stimulation of either Slo3 or PKA can maximize the capacitation levels in most of the combinations, they are predicted to be the strongest enhancers of capacitation. This is in agreement with the fact that they are strongly affecting membrane potential, a variable that is markedly different between capacitated and non-capacitated sub-

461   populations (Figure 6).

3   Discussion

The purpose of this paper is to increase our understanding of the early stages of murine capacitation within a systems biology approach. For this we build a mathematical model strongly based on experiment that can integrate the signaling processes related to this stage of capacitation with special attention on time dependent dynamical features. Novel contributions are implemented both at the mathematical and methodological levels impinging on the biological analysis. One of our main findings is that within our modeling we have identified a measurable quantity that regulates the percentage of sperm capacitation and hence may provide clues for a comprehension of this experimental observation. In the following, a listing of our principal contributions is summarized and discussed.

- On the model building: We construct a discrete, mostly deterministic regulatory network for early capacitation. The topology of the network is dictated by experimental knowledge, the elements only take discrete values, time is discrete and the regulatory functions that dictate the dynamics are also based on experiments. Novel propositions are considered in joint action nodes for ion fluxes and in the inclusion for some of the interactions of stochastic features that account for the presence of several time scales without the need of asynchronous updating, memory effects and population heterogeneity. Input from neural-network studies has been important. To our knowledge this is the first bottom to top capacitation model with the capability of incorporating and unraveling molecular mechanisms which intervene in the definition of regulatory relations that generate a dynamics for the study of temporal behavior. Previous modeling has been undertaken, which, with a top to bottom approach such as data-mining techniques focuses on determining the topology of prospective networks relevant to capacitation [58–62].

- On the operational definition of capacitation: The definition of a capacitated "state" within our modeling scheme, is crucial for our study. Keeping this in mind we proposed that the convergent attainment of certain levels of what we have called select set of nodes: PKA, Ca_i, Na_i, pH_i and V provides a good working definition. We are aware that as capacitation is a broad complex process, other variable could be considered. Such worthwhile efforts will be carried out in the future.

- On the connection between sperm heterogeneous capacitation response and the variability in population of the number of each flagellar ion transporter. A probability distribution of the amount of ion transporters in flagella is to be expected. Our contribution here is to show that within the model framework, changes in the standard deviation D (variability) of that distribution lead to modifications in the capacitation percentage. An extensive parameter exploration showed that this finding is recurrent and robust and can be attained for different parameter sets. As an example we consider the case of a parameter set for which the standard deviation $D = 0.25$ leads to about a third of capacitated sperm population. It is worthwhile pointing out that under these conditions, results with the value $D=0.25$ are consistent with the distributions of protein expression determined experimentally for other cells [19, 20]. At this value of D the time averaged distributions of Ca_i, pH_i and PKA favor higher values for capacitated sperm with respect to the non-capacitated population, whilst sperm membrane potential is at more negative hyperpolarized values (Sec 2.2.4), as physiologically expected. The characterization of this passage from a non-capacitated state to a capacitated state in terms of a variability control parameter is one of our main results.

  It is worth emphasizing that D allows for variations in the number and relation among ionic transporters which results, for a particular cell, in a combination of Ca_i, pH_i and V that dictates the activities of key capacitation enzymes such as PKA, determining if it reaches this maturational state.

  An in-depth analysis of this passage between different functional phases or states (i.e, non-capacitated/capacitated) with insight from other phase changes encountered in physics and nonlinear dynamics might well contribute to a better comprehension of the observed capacitation fraction. The unveiling from our modeling of a quantitative control parameter D for this qualitative transformation is experimentally suggestive, a corroboration of this prediction is called for. Results consistent with variations in the number of ion transporters modifying their relations leading to distinct $[Ca^{2+}]_i$, pH_i and V levels, and therefore functional capacities have been reported [21–23]. Also in this direction, recent findings suggest that proteolytic and biochemical processing of key proteins such CatSper 1, a subunit of CatSper, may contribute to explain why only few mammalian sperm out

of millions reach the fertilization site. This sperm exclusive multisubunit channel, in addition organizes $[Ca^{2+}]_i$ signaling nanodomains constituted by large protein complexes that are essential for sperm migration in the female tract. The orchestrated functioning of these complexes could filter out which sperm end up capacitated and able to travel and fertilize the egg [14, 15]. Therefore, observations documenting varying ionic transport component relations, as indicated in our modeling, may contribute to explain what selected characteristics allow sperm to complete their journey through the female reproductive tract and succeed in fertilizing the egg.

- On model experimental corroborations: To our knowledge this is the first dynamical model of the mammalian capacitation that reproduces experiments such as those reported in (Tab 1 and Tab 2). More specifically, it recovers 80 % of the selected set of results available to date (green cells), where the remaining 20 % is not in contradiction with the model (yellow cells). We cannot discard the possibility of missing components and mechanisms yet to be characterized by the field, e.g. ion channels like TMEM16A [63] and TRPs [64], which might complement the regulatory rules of our model and could help to increase agreement with experiments.

- On other model predictions: A result worthwhile highlighting is the single and double mutant analysis (Fig 7), which gives an insight on how the capacitation fraction levels can also be controlled through modification of specific nodes from the capacitation signaling network. Besides, it also exhibits hierarchies amongst key elements. Some cases of interest are extreme scenarios wherein the capacitation fraction either reaches levels of almost 100 % or is abolished.

  We draw special attention to the Slo3 gain of function case for which the model anticipates almost complete capacitation, and also to a noticeable capacitation increase with PKA gain of function. The model suggests that stimulating these network elements could lead to the design of experiments that would help in the study of assisted fertility.

  An intriguing finding is that the model variant with CatSper^{GOF} (Figure 7) predicts that capacitation levels are decreased with respect to the wild-type, furthermore a similar behavior is encountered in the CatSper^{LOF} counterpart. It is known that basal levels of intracellular calcium regulate the phosphorylation of several protein in the sperm flagella, however excessive intracellular calcium levels render the sperm flagella immotile [65]. In our simulations, we observe a decrease of capacitation fraction after sustained, high calcium levels (e.g. those produced by the overactive CatSper^{GOF} mutant), which suggests that extrusion mechanisms are important for the typical capacitation fractions. On the other hand, another important consequence of CatSper's sustained activity would be an increase in the depolarizing ion current that in turn would counteract the typical hyperpolarization response. This might explain the lower capacitation levels in our CatSper^{GOF} mutant simulations.

  Our model also predicts that the blockage of chloride channels leads to an increase of capacitation response (and vice versa); this might be indicative of the need of an additional Cl-related mechanism yet to be characterized. This result points in the opposite direction of previous experiments that assayed the effect of DPC, a general Cl-channel blocker, on hyperpolarization and AR of mouse sperm [9]. However, given the high inhibitor doses used in that study, and the low specificity of DPC, such contradiction has yet to be evaluated by the use of more specific drugs. Even though later works that used a CFTR specific inhibitor (inh-172) confirmed the hyperpolarization reduction reported with DPC [66], its ultimate effect on the whole capacitation process (e.g. by testing AR) remains to be characterized. In the regulation of intracellular chloride, many ion transporters participate forming intricate feedback loops. A plausible explanation of our model results is that CFTR and/or TMEM16A channels could be partially open in order to counter the charge generated by the electrogenic cotransporters of the WT sperm flagella. Inhibiting the intracellular chloride outward flux would contribute to a negative charge accumulation in the flagella hampering other required ion fluxes and leading to a more hyperpolarized membrane potential and increasing the capacitation fraction predicted by the model (and vice versa).

While emphasis is placed on the importance of capacitation regulation due to the variability in the number of ionic transporters, it is also worthwhile to highlight two findings within the model: confluence and synergy. The capacitation criterion implemented in the modeling requires the confluence within reduced time intervals of a particular behavior in the selected nodes, if only one of those nodes alters its behavior, capacitation is lost. Synergy manifests itself when considering the oversized response in the capacitation fraction under loss-of-function mutants and gain-of-function mutants when performed in pairs, as shown in the section 2.2.5.

As a final remark we would like to point out that the findings, conclusions and predictions of this study reflect a systemic, integrated analysis based on experimental information available.

4 Methods

In this subsection, we describe the building methods for regulatory functions of: ion concentrations, ion transport, emf nodes, auxiliary nodes and early phosphorylation, among others. We give a table with all initial conditions used in our simulations. Also, particular cases of ion flux nodes with modifiers are explained.

Due to the discrete formalism, we employ a discretization of Ohm's law, Nernst potential and electrochemical potential equations in order to calculate the regulatory functions according to physiological conditions. All parameters are reported in tables 4,5, and 6.

4.1 Initial conditions settings

Table 3 shows the discrete numerical values used in all our simulation as initial conditions.

4.2 Ad hoc regulatory functions

4.2.1 Intracellular and extracellular ion concentration nodes

Ion concentration nodes only consider two states: low concentration (state 0) and high concentration (state 1). Also, they have two regulators: the total flux of their corresponding ion and the intracellular concentration itself. Depending on the direction of the flux and intracellular concentration, there may be an increase, no change or a decrease in the concentration. For intracellular pH, sodium and bicarbonate, there is an extra regulatory node related to basal state recovery. Table 4 summarizes numeric values of voltage, extracellular and intracellular concentrations, used for the calculation of ion transporter flux direction and magnitude.

4.2.2 Gating nodes for ion channels

CatSper For the modeling of CatSper gating, we considered four regulators: voltage, intracellular calcium, pH_i and PKA. The following equations determine the effect of the first three regulators, as in [67] :

$$P_V(V, \mathrm{pH}_i) = \frac{1}{1 + e^{\frac{V_{pH}(\mathrm{pH}_i) - V}{\mathrm{K}}}} \tag{14a}$$

$$V_{\mathrm{pH}}(\mathrm{pH}_i) = A \frac{K_{\mathrm{pH}}^{n_H}}{\mathrm{pH}_i^{n_H} + K_{\mathrm{pH}}^{n_H}} + B \tag{14b}$$

$$P_{\mathrm{Ca}}(\mathrm{Ca}_i) = \frac{K_{\mathrm{Ca}}^{n_{\mathrm{Ca}}}}{\mathrm{Ca}_i^{n_{\mathrm{Ca}}} + K_{\mathrm{Ca}}^{n_{\mathrm{Ca}}}} \tag{14c}$$

where P_V is the opening probability of CatSper channels by influence of membrane potential and protons. Eq. 14a is a function used to model the sigmoid (conductance vs potential) G/V curve, as reported in [68]. Equation 14b (V_{pH}) models the pH_i-dependent shift of the half-activation voltage of Eq. 14a. P_{Ca} is the opening probability of CatSper channels by influence of calcium, where K_{pH} y K_{Ca} are empirical fitted parameters. Also, n_{Ca} and n_{pH} are cooperativity coefficients of calcium and pH_i, respectively. Finally, we multiply Eq. 14a and Eq. 14c in order to obtain the opening probability of CatSper channels, P_{CatSper}, by influence of the first three regulators. We discretize this final probability by a threshold value $max(P_V P_{Ca})/2$, where function max reaches the greatest possible value for the given regulator combination. We associate the state 0 (basal gating) to probabilities below this threshold value and state 1 (more permeable gating) to probabilities above of this threshold value. The only action of PKA in the gating consists in shifting the permeability to state 1 in spite of resting pH_i and resting membrane potential. Table 5 contains a summary of numerical values used in the gating equations.

Na$^+$-channel The sodium conductance increases (state 1) with respect to basal levels (state 0) when its only regulator ClC is closed (state 0).

Node	Name	Value
$\Delta\mu_{\mathrm{NCX}}$	Sodium/Calcium exchanger emf	0
$\Delta\mu_{\mathrm{PMCA}}$	Calcium pump emf from principal piece of flagellum	0
$\Delta\mu_{\mathrm{Cat}}$	CatSper emf	0
Ca_i	Intracellular calcium	0
Ca_e	Extracellular calcium	1
Na_i	Intracellular sodium	1
Na_e	Extracellular sodium	1
pH_i	Intracellular pH	0
pH_e	Extracellular pH	1
V	Membrane potential	0
Cat	CatSper gating	0
ICa	Calcium current	0
IH	Proton current	0
cAMP	Cyclic adenosine monophosphate	0
IP3	Inositol trisphosphate	0
$\Delta\mu_{\mathrm{SPCA}}$	Calcium pump from calcium reservoirs emf	0
$\Delta\mu_{\mathrm{IP3R}}$	IP3 channel emf	0
Slo3	Slo3 gating	0
K_e	Extracellular potassium	0
$\Delta\mu_{\mathrm{KSper}}$	Slo3 emf	0
NaC	NaC gating	0
ClC	ClC gating	0
$\Delta\mu_{\mathrm{NaC}}$	NaC emf	0
$\Delta\mu_{\mathrm{SLC26}}$	Electrogenic chloride/bicarbonate exchanger emf	0
$\Delta\mu_{\mathrm{ClC}}$	ClC emf	0
Cl_i	Intracellular chloride	0
Cl_e	Extracellular chloride	1
HCO_{3i}	Intracellular bicarbonate	0
HCO_{3e}	Extracellular bicarbonate	0
sAC	Soluble adenylate cyclase	0
PDE	General phosphodiesterases	0
PKA	Protein kinase A	0
INa	Sodium current	0
$\Delta\mu_{\mathrm{NBC}}$	Cootransporter sodium/bicarbonate emf	0
sNHE	Sodium/proton exchanger	0
$\Delta\mu_{\mathrm{sNHE}}$	Sodium/proton exchanger emf	0
$\Delta\mu_{\mathrm{SLCAE}}$	Electroneutral chloride/bicarbonate exchanger emf	0
IHCO3	Bicarbonate current	0
ICl	Chloride current	0
ChAcc	Cholesterol acceptor	0
$\Delta\mu_{\mathrm{pHR}}$	pH recovery emf	0
$\Delta\mu_{\mathrm{NaR}}$	Sodium recovery emf	0
$\Delta\mu_{\mathrm{HCO3R}}$	Bicarbonate recovery emf	0
$\Delta\mu_{\mathrm{Leak}}$	Leak current emf	0

Table 3: Summary of initial conditions used for all simulations, emf corresponds to electromotive force.

Cl⁻-channel The chloride conductance increases (state 1) with respect to basal levels (state 0) when its only regulator PKA is active (state 1).

Slo3 In order to choose whether the node will remain in basal level (state 0) or in a more permeable state (state 1), we used the gating equation of the supplementary material in [69] with the respective numerical constants. The discretization was done under the same criteria mentioned in Sec. 4.2.2. Under depolarized potential and high pH_i, Slo3 takes the state 1 if ChAcc or PKA or both are in state 1. Under high pH_i, if PKA and ChAcc are both in state 1, Slo3 can take the state 1 even in resting potential.

Node	State -1	State 0	State 1	Units	Reference
Ca_e	NA	0.01	2	mM	[8]
pH_e	NA	6.80	7.20	NA	[4]
ChAcc	NA	Absence	Presence	NA	[24]
Na_e	NA	0.01	130	mM	[10]
HCO_{3e}	NA	0.01	25	mM	[42]
Cl_e	NA	0.01	130	mM	[9]
K_e	NA	0.01	5	mM	[4]
Ca_i	NA	0.10	0.25	µM	[8]
pH_i	NA	6.50	6.80	NA	[4]
Na_i*	NA	10	14	mM	[10]
HCO_{3i}**	NA	5	15	mM	[42]
Cl_i	NA	30	40	mM	[9]
V	-60	-40	-20	mV	[4]

Table 4: Summary of numerical values of extracellular, intracellular ion concentrations and voltage used in calculations. (*) This value is an approximate extrapolation from [10]. (**) These values are a work hypothesis.

Constant	Numerical value
K	$30\,\mathrm{mV}$
K_{Ca}	$0.5\,\mathrm{µM}$
n_{Ca}	2
A	$79.7\,\mathrm{mV}$
K_{pH}	6.81
n_H	35.52
B	$8.35\,\mathrm{mV}$

Table 5: Numerical values of the constants used in CatSper gating equations. Extracted from [67]

4.2.3   Current nodes of ion channels

CatSper current   The magnitude and direction for this Ca^{2+} flux was estimated only considering the calcium gradient by means of the Nernst potential and Ohm's equation as follows:

$$E_{CatSper}(Ca_e, Ca_i) = \frac{RT}{2F} \ln\left(\frac{Ca_e}{Ca_i}\right) \tag{15a}$$

$$\Delta\mu_{CatSper} = V - E_{CatSper} \tag{15b}$$

$$I_{CatSper}(P_{CatSper}, V, E_{CatSper}) = P_{CatSper} N_{CatSper} g_{CatSper} \Delta\mu_{CatSper}, \tag{15c}$$

where R is the Molar gas constant, T is the temperature, F is the Faraday constant, and the 2 multiplying F in the denominator stands for the charge number of calcium ions, $P_{CatSper}$ is the output value of the gating node of CatSper, defined in Sec. 4.2.2, $N_{CatSper}$ is the number of channels and $g_{CatSper}$ is the CatSper unitary conductance. According to the combination of its regulator states ($Ca_e, Ca_i, P_{CatSper}, V$) with help of Eqs. 15a to 15c, we determined if the CatSper current will have an outward calcium current (state -1), a null current (state 0) or an inward calcium current (state 1). Table 6 shows the numerical values of constants used in Eqs. 15 through 18.

Na^+-channel current   To estimate the magnitude and direction of this ion flux, we considered the sodium gradient and we used the Na^+ Nernst potential and the Ohm equation as follows:

$$E_{NaC}(Na_e, Na_i) = \frac{RT}{F} \ln\left(\frac{Na_e}{Na_i}\right) \tag{16a}$$

Parameter	Numerical value	Parametrization considerations
$P_{K\mathrm{Na}}$	5	[70].
P_{Slo3}*	0.0035	After plugging maximal V and pH$_i$ values (Table 4) into model from [69].
N_{Slo3}	787	After fitting model from [69] to data from Fig3c in [71].
g_{Slo3}	90 pS	[72].
P_{ClC}*	1	
N_{ClC}	10	
g_{ClC}	8 pS	Considering CFTR conductance [73].
P_{HCO3Cl}	0.48	[74].
P_{NaC}*	1	
N_{NaC}	10	
g_{NaC}	5 pS	Considering ENaC conductance [75].
P_{CatSper}*	0.087	After plugging maximal V, maximal pH$_i$ and minimal Ca$_i$ values (Table 4) into Eq. 14.
N_{CatSper}	1026	After fitting model from [67] to data from [68] (Figs. 4b and c).
g_{CatSper}	0.5 pS	Assuming a low, sub-picosiemen Ca^{2+} conductance.
n_{NBC}	3	[42, 76].
n_{SLC26}	2	[66].

Table 6: Summary of the parameters used in current and energy equations of this section. (*) The values reported for these quantities, which portray the fraction of ion channels in open state, correspond to the state 1 in their discrete node counterpart. For N_{ClC} and N_{NaC}, their ion number values are a working hypothesis after taking into account that Slo3 and CatSper currents are dominant and other fluxes are of around one order of magnitude lower [72].

$$\Delta\mu_{\mathrm{NaC}} = V - E_{\mathrm{NaC}} \tag{16b}$$

$$I_{\mathrm{NaC}}(P_{\mathrm{NaC}}, V, E_{\mathrm{NaC}}) = P_{\mathrm{NaC}} N_{\mathrm{NaC}} g_{\mathrm{NaC}} \Delta\mu_{\mathrm{NaC}}, \tag{16c}$$

where P_{NaC} is the Na-channel gating node output value, N_{NaC} is the number of Na channels and g_{NaC} is the unitary conductance of Na channels. According to the combination of regulator states $(\mathrm{Na}_e, \mathrm{Na}_i, P_{\mathrm{NaC}}, V)$ we decided with the help of equations 16a to 16c if the Na-channel current will have an outward sodium current (state -1), a null current (state 0) or an inward sodium current (state 1).

Cl$^-$-channel current In order to obtain some notion of magnitude and direction of this ion flux, we considered a chloride/bicarbonate mixed current as in CFTR channel, and we used the Goldman-Hodgkin-Katz (GHK) voltage equation and the Ohm equation as follows:

$$E_{\mathrm{ClC}}(\mathrm{HCO3}_e, \mathrm{Cl}_e, \mathrm{HCO3}_i, \mathrm{Cl}_i) = \frac{RT}{F} \ln\left(\frac{P_{\mathrm{HCO3Cl}}\mathrm{HCO3}_i + \mathrm{Cl}_i}{P_{\mathrm{HCO3Cl}}\mathrm{HCO3}_e + \mathrm{Cl}_e}\right) \tag{17a}$$

$$\Delta\mu_{\mathrm{ClC}} = V - E_{\mathrm{ClC}} \tag{17b}$$

$$I_{\mathrm{ClC}}(P_{\mathrm{ClC}}, V, E_{\mathrm{ClC}}) = P_{\mathrm{ClC}} N_{\mathrm{ClC}} g_{\mathrm{ClC}} \Delta\mu_{\mathrm{ClC}}, \tag{17c}$$

where P_{ClC} is the Cl-channel gating node output value, N_{ClC} is the number of Cl channels and g_{ClC} is the unitary conductance of Cl channels. According to combination of regulatory states $(\mathrm{HCO3}_e, \mathrm{Cl}_e, \mathrm{HCO3}_i, \mathrm{Cl}_i, P_{\mathrm{ClC}}, V)$, we decided with help of equations 17a to 17c if current of ClC will have an outward chloride/bicarbonate current (state -1), a null current (state 0) or an inward chloride/bicarbonate current (state 1).

KSper   It is known that Slo3 is mainly responsible for the sperm potassium current named KSper [71]. However, unlike Slo1 channels, Slo3 selectivity for potassium is weak, this measured in heterologously expressed channels ($P_{K^+}/P_{Na^+} \sim 5$) [70], which would result in a reversal potential close to the resting potential. Even though it has been reported that the blockage or deletion of Slo3 abolishes the typically observed hyperpolarization in capacitated sperm [49, 50], the opening of a channel with those characteristics would never be able to generate hyperpolarizing currents. To overcome this apparent contradiction, we decided to model this current as a purely K^+-current from possible additional K^+ channels downstream of Slo3 activity and with higher selectivity. In order to obtain some notion of magnitude and direction for this current, we considered a potassium flux according to the K^+ Nernst potential and Ohm equation as follows:

$$E_{\text{KSper}}(K_e, K_i) = \frac{RT}{F} \ln \left(\frac{K_e}{K_i} \right) \tag{18a}$$

$$\Delta\mu_{\text{KSper}} = V - E_{\text{KSper}} \tag{18b}$$

$$I_{\text{KSper}} = P_{\text{Slo3}} N_{\text{Slo3}} g_{\text{Slo3}} \Delta\mu_{\text{KSper}}, \tag{18c}$$

where P_{Slo3} is the Slo3 gating node output value, N_{Slo3} is the number of Slo3 channels and g_{Slo3} is the unitary conductance of Slo3 channels. Taking into account the combination of regulator states ($K_e, K_i, P_{\text{Slo3}}, V$), for the KSper electromotive force we discretized equation 18b. The state of this node will be -1 if the gradient generates inward flux, 0 when there is no net flux, and 1 if the gradient causes outward flux.

4.2.4   Cotransporter and exchanger nodes

Electrogenic sodium/calcium exchanger   For the flux direction of sodium/calcium exchanger (NCX), we used an equation from appendix in [8] with the respective reported constants. We chose the state -1 for outward flux of sodium and influx of calcium, state 0 for null flux, state 1 for influx of sodium and outward flux of calcium.

Electroneutral sodium/proton exchanger   Estimation of the flux direction of the sperm-specific electroneutral sodium/proton exchanger (sNHE) was performed following energetic consideration for determining the most probable state of sNHE according to the regulator chemical gradients as follows:

$$\Delta G_{\text{NHE}} = \Delta\mu_{\text{Na}} - \Delta\mu_H = RT \ln \left(\frac{\text{Na}_i}{\text{Na}_e} \right) - RT \ln \left(\frac{H_i}{H_e} \right) \tag{19}$$

We assigned state -1 to the sodium outward flux coupled with protons influx, state 0 for null flux, state 1 for sodium influx coupled with protons outward flux. This node has a gating regulator that depends on voltage and cAMP. If the cytosolic cAMP concentration increases and the voltage is hyperpolarized then the flux is enhanced. If only one of the two regulators is present in the right state the flux is permitted. If none of these regulators are in the right state, the flux is forbidden.

Electrogenic sodium/bicarbonate cotransporter   We calculated the flux direction of sodium/ bicarbonate cotransporter (NBC), by means of the reversal potential and the following energetic consideration for determining the most probable state of NBC according to the regulator gradients and membrane potential:

$$E_{\text{NBC}} = \frac{RT}{(n_{\text{NBC}} - 1)F} \ln \left(\frac{\text{Na}_i \text{HCO}_{3i}^{n_{\text{NBC}}}}{\text{Na}_e \text{HCO}_{3e}^{n_{\text{NBC}}}} \right) \tag{20a}$$

$$\Delta\mu_{\text{NBC}} = V - E_{\text{NBC}} \tag{20b}$$

We assigned the state -1 to outward flux, state 0 for null flux and state 1 for influx of sodium and bicarbonate. The sodium/bicarbonate stoichiometry per individual transport event by this cotransporter, n_{NBC}, is reported in Table 6.

Electroneutral chloride/bicarbonate exchanger In order to obtain the ion flux direction of Cl^-/HCO_3^- electroneutral exchanger (SLCAE), we used the following energetic consideration for determining the most probable state of SLCAE according to the regulator gradients:

$$\Delta G_{\text{AE}} = \Delta\mu_{\text{HCO3}} - \Delta\mu_{\text{Cl}} = RT\ln\left(\frac{\text{HCO}_{3i}}{\text{HCO}_{3e}}\right) - RT\ln\left(\frac{\text{Cl}_i}{\text{Cl}_e}\right) \tag{21}$$

We assigned the state -1 to outward flux of chloride and influx of bicarbonate, state 0 for null flux, state 1 for influx of chloride and outward flux of bicarbonate.

Electrogenic chloride/bicarbonate exchanger The flux direction of electrogenic Cl^-/HCO_3^- exchanger (SLC26A3) was estimated using the following energetic consideration for determining the most probable state of SLC26 according to the regulator gradients and membrane potential:

$$\begin{aligned}
\Delta\mu_{\text{SLC26}} &= n_{\text{SLC26}}E_{\text{Cl}} - E_{\text{HCO3}} - (n_{\text{SLC26}} - 1)V \\
&= n_{\text{SLC26}}\frac{RT}{F}\ln\left(\frac{\text{Cl}_i}{\text{Cl}_e}\right) - \frac{RT}{F}\ln\left(\frac{\text{HCO}_{3i}}{\text{HCO}_{3e}}\right) - (n_{\text{SLC26}} - 1)V
\end{aligned} \tag{22}$$

We assigned state -1 to outward flux of chloride coupled with influx of bicarbonate, state 0 for null flux, state 1 for influx of chloride coupled with outward flux of bicarbonate. The stoichiometry of this exchanger, n_{SLC26}, is reported in Table 6.

4.2.5 Early phosphorylation nodes

The soluble adenylate cyclase (sAC) only has two regulators: intracellular calcium (Ca_i) and intracellular bicarbonate (HCO_{3i}). The sAC activates (state 1) if Ca_i or HCO_{3i} is in high concentration in the cytosol (state 1), otherwise sAC will be inactivate (state 0).

Phosphodiesterase (PDE) will increase its activity from basal (state 0) to high (state 1) if there is adenosine monophosphate in cytosol (state 1).

Cyclic Adenosine monophosphate (cAMP) has three regulators: itself, sAC and PDE. The cAMP will remain in its previous state if sAC and PDE both remain inactive. There will be production of cAMP (state 1) only while sAC is activated (state 1). There will be a decrease of cAMP (state 0) only while PDE is activated (state 1). There will be a decrease of cAMP (state 0) if PDE and sAC are both activated (state 1).

Protein kinase A (PKA) will remain activate (state 1) while there is cAMP in the cytosol (state 1) otherwise PKA will remain inactivated (state 0).

4.2.6 Auxiliary recovery nodes

When the recovery nodes of (pHR, NaR and HCO3R) turn on, they set their respective ion concentration (pH$_i$,Na$_i$ and HCO3$_i$) from the high (state 1) to their resting levels.

The leak current node (IL) is a nonspecific counterbalance mechanism to voltage changes, i. e. it tends to depolarize if V is hyperpolarized (state -1) and to hyperpolarize if V is depolarized (state 1), leading in both cases to resting values.

4.2.7 Miscellany of nodes

The inositol trisphosphate (IP3) node will increase its concentration in cytosol (state 1) while Ca_i remains high (state 1) and cAMP remains high (state 1). We discretized the continuous model proposed by [77], in which it is argued that IP3 is synthesized near the midpiece through a PLC isoform and can be regulated by calcium and cAMP.

IP3 sensitive calcium channels from calcium reservoirs (IP3R) node will promote a calcium current to cytosol (state 1) while IP3 remains high (state 1) and Ca_i remains low (state 0) in the cytosol.

Calcium pump from calcium reservoirs (SPCA) node will be activated (state 1) while there is a high intracellular calcium (state 1) promoting a calcium current from cytosol to the interior of calcium reservoirs.

Calcium pump from main piece of flagellum (PMCA) node will be activated (state 1) while there is high intracellular calcium concentration (state 1) promoting a current from cytosol to the outside of the cell. [8]

4.3 Ion transporter weights for ion flux integration and voltage regulators

Ion flux integration nodes (Eq. 10) play a fundamental role in the network dynamics. In order to ponder the importance of the different ion transporters contributing to an integration node we determine the weight w_j of each one of them.

For ion channel weight contributions, keeping in mind equations 15c, 16c, 17c and 18c, we look into total conductances η_j, evaluated from the average number of ion transporters N_j and unitary conductances g_j (both summarized in Table 6), and define them as follows:

$$\eta_j = N_j g_j \tag{23a}$$

$$w_j \equiv \frac{P_j \eta_j}{P_m \eta_m} = \frac{P_j g_j N_j}{P_m g_m N_m} \tag{23b}$$

where P_j, P_m are the maximum probability gating values, $w_j \in \mathbb{R}$ and index m corresponds to the ion channel having the highest total conductance, which we use as a rescaling factor, hence $w_j \leq 1$. For the above determinations, when available, we used data from patch-clamp measurements reported in the literature to estimate N_j, and we assumed such quantities to be representative of sperm populations.

Using as reference the maximal values for the main Ca^{2+} fluxes of the sperm flagellum, namely, CatSper (calculated from [67] and Eq. 15c), NCX and PMCA (calculated using [8] model), we distinguish three different levels, which are approximately interrelated by factors of 2. Within our coarse grained approach, we have preserved these ratios and implemented in general the assignment of: 1 to the lower currents, 2 to intermediate ones and 4 to higher currents.

Under these assumptions, based on experimental knowledge and the calibration underlying the results of tables 1 and 2 we assign the flux weights for the following ion transport integrators:

- I_{Ca}: $w_{CatSper}$=4, w_{NCX}=2, w_{PMCA}=1, w_{SPCA}=1, w_{IP3R}=4
- I_{Cl}: w_{ClC}=1, w_{SLCAE}=1, w_{SLC26}=2
- I_{Na}: w_{NaR}=2, w_{NaC}=4, w_{NBC}=1
- I_{HCO_3}: w_{ClC}=1, w_{NBC}=1, w_{SLCAE}=1, w_{SLC26}=1, w_{HCO3R}=1
- JH: w_{NHE}=1, w_{pHR}=2, w_{IHCO3}=1

Another pivotal integrating node is the membrane potential V calculated by Eq. 11, in this case besides some of the above w_j for ionic channel regulators we determine the weights for ion fluxes corresponding to the NBC co-transporter, the SLC26 exchanger and the leak current, as well as for the feedback from V at a previous time in a similar way and under the same assumptions implemented for the ion flux integrators, namely: w_{NBC}=1, w_{SLC26}=1, w_V=4 and w_{IL}=4.

Given an integrator i we shall further on label the w_j weights associated to it by w_{i_j}. In order to introduce variability in the amount of ion transporters present in the sperm populations, we multiply each of the w_{i_j} by a random variable that samples a normal distribution with average 1 and deviation D. This distribution was sampled over the interval $G_{i_j}^r(1, D) \in [0, 2]$ because negative conductances have no meaning in our modelling; furthermore, given that the center of the distribution is 1, in order to avoid any sampling bias while keeping symmetry, we fixed the interval up to 2. Thus, a random set of weights $\{w_{i_j} G_{i_j}^r(1, D)\}$ is assigned to each single sperm r at the beginning of the simulations. Note that the probabilistic element here introduced acts directly on network links through the stochastic modulation of the integrator weights, furthermore more and most importantly, given Eq. 23b it regulates the distribution of the numbers of ionic transporters per cell in the population and hence is experimentally challenging to determine.

We consider differences in operation velocity of various ion transporters, by using a random variable $b \in \{0, 1\}$ that samples a biased Bernoulli distribution at each iteration t, where τ_{i_j} is the distribution mean. If b is equal to 1 at time t, the ion transporter i_j will participate in the weighted sum with their respective ion flux output value. Otherwise, the ion transporter will have a null participation in the weighted sum at time t. The average of the biased Bernoulli distribution τ_{i_j} of each ion transporter i_j is related to velocity parameter v_{i_j} as follows:

$$\tau_{i_j} = 0.25 v_{i_j}, \text{ with } v_{i_j}^{-1} \in \{1, 2, 4, ..., 9\} \tag{24}$$

where 0.25 is a proportionality factor that enables capacitation time scales to be consistent with those reported in the literature (e. g. after $\approx$120 seconds of capacitating stimulus, PKA-dependent phosphorylation reaches maximal levels in boar spermatozoa [34, 78]), and is related to the maximum mean value that can be assigned to the Bernoulli distribution. A velocity equal to 1 corresponds to ion channels, a velocity equal to $1/2$ corresponds to exchangers and cotransporters, a velocity equal to $1/4$ corresponds to calcium pumps. The inverse velocities $(v_{i_j}^{-1})$ greater than 4 correspond to phosphorylation regulator nodes, ChAcc regulator node and calcium diffusion from RNE to the principal piece. These kinetic parameters reflect the flux given by the total number of each type of ion transporter in the flagellum. All parameters used in this section are reported in Table 6.

4.4 Thresholds of membrane potential node

Eq. 11 was deduced from a discrete version of Hodgkin-Huxley equation:

$$\frac{dV}{dt} = -\frac{1}{C_m} \sum_j^k \tilde{I}_j(t), \tag{25}$$

where C_m is the flagellum capacitance of the sperm and $\tilde{I}_j(t)$ is the flux of ion transporter j at time t. In general, the weighted sum in Eq. 11 can take values from the interval $x_i \in (l_{i_o}, l_{i_m})$ which can be partitioned into $p_j = (l_{i_{j-1}}, l_{i_j})$ and discretized as follows:

$$\Theta_i(x_i) = \begin{cases} e_{i_m} & \text{if } x_i \in p_m \\ . \\ . \\ . \\ e_{i_1} & \text{if } x_i \in p_1 \end{cases} \tag{26}$$

where $l_{i_j} \in (l_{i_o}, l_{i_m})$ and m is the number of states that σ_i can take. In particular, for the case of the membrane potential in Eq. 11, the function Θ discretizes *ad hoc* the output of the weighted sum as follows:

$$\Theta = \begin{cases} 1 & \text{if } 3 < x \\ 0 & \text{if } -2 \leq x \leq 3 \\ -1 & \text{if } x < -2 \end{cases}, \tag{27}$$

where $x = [\sum_{j=V_1}^{V_k-1} z_j b_{\tau_j}^t w_j G_j^r(1, D) \Delta\mu_j(t) + w_{V_k} V^r(t)]$, state 1 (depolarized), state 0 (resting) and state -1 (hyperpolarized).

4.5 Ion flux transporter modifiers

It is known that there are agonists and antagonists of the ion transporters involved in capacitation, and that they are modulated during this maturation process. To account for ion flux modulators, such as cholesterol acceptors (serum albumin), PKA-dependent phosphorylation and protein-protein interactions, we introduce functions that modify some of the interaction weights related to ion transporter nodes. This has the effect of changing the flux magnitude of the corresponding ion transporter by multiplying it by a factor.

Chloride fluxes   We determine the chloride flux integrator node $\tilde{I}_{\text{Cl}}$ regulation with:

$$\begin{aligned} \tilde{I}_{\text{Cl}}^r(t+1) = sgn \Big[&Q^t(\sigma_{\text{ChAcc}}) b_{\tau_{I_{\text{ClC}}}}^t w_{I_{\text{ClC}}} G_{I_{\text{ClC}}}^r(1, D) \Delta\mu_{I_{\text{ClC}}}(t) \\ &- 2P^t(\sigma_{\text{ClC}}, \sigma_{\text{ChAcc}}, \sigma_{\text{PKA}}) b_{\tau_{\text{SLC26}}}^t w_{\text{SLC26}} G_{\text{SLC26}}^r(1, D) \Delta\mu_{\text{SLC26}}(t) \\ &- Q^t(\sigma_{\text{ChAcc}}) b_{\tau_{\text{SLCAE}}}^t w_{\text{SLCAE}} G_{\text{SLCAE}}^r(1, D) \Delta\mu_{\text{SLCAE}}(t) \Big], \end{aligned} \tag{28}$$

where the chloride transporter modifier functions P^t and Q^t are defined as follows:

$$P^t = \begin{cases} 4 & \text{if } \sigma_{\text{ClC}} \wedge \sigma_{\text{PKA}} \wedge \sigma_{\text{ChAcc}} = 1 \\ 2 & \text{if } \sigma_{\text{ClC}} \wedge \sigma_{\text{PKA}} \wedge \neg\sigma_{\text{ChAcc}} = 1 \\ 1 & \text{else} \end{cases} \tag{29}$$

$$Q^t = \begin{cases} 4 & \text{if } \sigma_{\text{ChAcc}} = 1 \\ 1 & \text{else} \end{cases} \tag{30}$$

Bicarbonate fluxes We determine the bicarbonate flux integrator node $\tilde{I}_{\text{HCO3}}$ regulation:

$$
\begin{aligned}
\tilde{I}^r_{\text{HCO3}}(t+1) = sgn \Big[&Q^t(\sigma_{\text{ChAcc}}) b^t_{\tau_{I_{\text{ClC}}}} w_{I_{\text{ClC}}} G^r_{I_{\text{ClC}}}(1,D) \Delta\mu_{\text{IClC}}(t) \\
&+ P^t(\sigma_{\text{ClC}}, \sigma_{\text{ChAcc}}, \sigma_{\text{PKA}}) b^t_{\tau_{\text{SLC26}}} w_{\text{SLC26}} G^r_{\text{SLC26}}(1,D) \Delta\mu_{\text{SLC26}}(t) \\
&+ Q^t(\sigma_{\text{ChAcc}}) b^t_{\tau_{\text{SLCAE}}} w_{\text{SLCAE}} G^r_{\text{SLCAE}}(1,D) \Delta\mu_{\text{SLCAE}}(t) \\
&+ Q^t(\sigma_{\text{ChAcc}}) b^t_{\tau_{\text{NBC}}} w_{\text{NBC}} G^r_{\text{NBC}}(1,D) \Delta\mu_{\text{NBC}}(t) \Big],
\end{aligned}
\tag{31}
$$

where the bicarbonate transporter modifiers P^t and Q^t are defined in the same way of chloride.

Sodium fluxes We determine the sodium flux integrator node $\tilde{I}_{\text{Na}}$ regulation:

$$
\begin{aligned}
\tilde{I}^r_{\text{Na}}(t+1) = sgn \Big[&b^t_{\tau_{\text{NaR}}} w_{\text{NaR}} G^r_{\text{NaR}}(1,D) \Delta\mu_{\text{NaR}}(t) \\
&+ b^t_{\tau_{\text{INaC}}} w_{\text{INaC}} G^r_{\text{INaC}}(1,D) \Delta\mu_{\text{INaC}}(t) \\
&+ Q^t(\sigma_{\text{ChAcc}}) b^t_{\tau_{\text{NBC}}} w_{\text{NBC}} G^r_{\text{NBC}}(1,D) \Delta\mu_{\text{NBC}}(t) \Big],
\end{aligned}
\tag{32}
$$

where the bicarbonate transporter modifier Q^t is defined in the same way of chloride.

5 Acknowledgements

A.A-G, A.A and G. M-M thank the hospitality of the Physics Department of the École Normale Superérieure in Paris (ENS), G. M-M during a sabbatical leave, where part of this research was undertaken. A. D. thanks the hospitality of the Instituto Gulbenkian de Ciência, Oeiras, Portugal, during a sabbatical leave

6 Author contributions

Alejandro Aguado-García: Conceptualization, Investigation, Methodology, Formal Analysis, Software, Writing–Original Draft Preparation, Writing–Review & Editing.

Daniel A. Priego-Espinosa: Conceptualization, Investigation, Methodology, Formal Analysis, Software, Writing–Original Draft Preparation, Writing Review & Editing.

Andrés Aldana: Conceptualization, Investigation, Methodology.

Alberto Darszon: Conceptualization, Investigation, Methodology, Funding Acquisition, Validation, Supervision, Writing–review & Editing.

Gustavo Martínez-Mekler: Conceptualization, Investigation, Methodology, Funding Acquisition, Resources, Supervision, Writing-Original Draft Preparation, Writing–review & Editing.

7 Financial Disclosure Statement

All authors thank CONACyT for grant CB-2015-01 255914-F7. A.D. performed part of this work while carrying out a Sabbatical at the Instituto Gulbenkian de Ciência (IGC) supported by DGAPA/PASPA/ UNAM and IGC. A.A.A-G. thanks CONACyT for doctoral and mobility fellowships 509572. A.A-G. thanks CONACyT for doctoral fellowship 428858. G.M-M. acknowledges support from DGAPA/PASPA/UNAM for a sabatical leave at ENS, Paris. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

8 Competing interests

There are no competing interests.

References

[1] M. C. Chang. "Fertilizing Capacity of Spermatozoa deposited into the Fallopian Tubes". In: *Nature* 168.4277 (Oct. 1951), pp. 697–698. DOI: 10.1038/168697b0.

[2] C. R. Austin. "The 'Capacitation' of the Mammalian Sperm". In: *Nature* 170.4321 (Aug. 1952), p. 326. DOI: 10.1038/170326a0.

[3]   Y. Yoda, M. Yokoyama, and T. Hosi. "Studies on the fertilization of mouse eggs in vitro". In: *Japanese J. Anim. Reprod.* 16.4 (1971), pp. 147–151. DOI: 10.1262/jrd1955.16.147.

[4]   A. Darszon, T. Nishigaki, C. Beltran, and C. L. Treviño. "Calcium Channels in the Development, Maturation, and Function of Spermatozoa". In: *Physiol. Rev.* 91.4 (Oct. 2011), pp. 1305–1355. DOI: 10.1152/physrev.00028.2010.

[5]   C. R. Ward and B. T. Storey. "Determination of the time course of capacitation in mouse spermatozoa using a chlortetracycline fluorescence assay". In: *Dev. Biol.* 104.2 (1984), pp. 287–296. DOI: 10.1016/0012-1606(84)90084-8.

[6]   R. Yanagimachi. "Mammalian Fertilization". In: *Physiol. Reprod.* Ed. by E. Knobil and J. D. Neill. 2nd ed. New York, NY, USA: Raven Press, 1994, pp. 189–317.

[7]   R. Yanagimachi. "The movement of golden hamster spermatozoa before and after capacitation". In: *Reproduction* 23.1 (Oct. 1970), pp. 193–196. DOI: 10.1530/jrf.0.0230193.

[8]   G. Wennemuth, D. F. Babcock, and B. Hille. "Calcium Clearance Mechanisms of Mouse Sperm". In: *J. Gen. Physiol.* 122.1 (July 2003), pp. 115–128. DOI: 10.1085/jgp.200308839.

[9]   E. O. Hernández-González et al. "Involvement of Cystic Fibrosis Transmembrane Conductance Regulator in Mouse Sperm Capacitation". In: *J. Biol. Chem.* 282.33 (Aug. 2007), pp. 24397–24406. DOI: 10.1074/jbc.M701603200.

[10]   J. Escoffier, D. Krapf, F. Navarrete, A. Darszon, and P. E. Visconti. "Flow cytometry analysis reveals a decrease in intracellular sodium during sperm capacitation". In: *J. Cell Sci.* 125.2 (Jan. 2012), pp. 473–485. DOI: 10.1242/jcs.093344.

[11]   P. E. Visconti, J. L. Bailey, G. D. Moore, D. Pan, P. Olds-Clarke, and G. S. Kopf. "Capacitation of mouse spermatozoa. I. Correlation between the capacitation state and protein tyrosine phosphorylation." In: *Development* 121.4 (Apr. 1995), pp. 1129–1137.

[12]   P. E. Visconti et al. "Capacitation of mouse spermatozoa. II. Protein tyrosine phosphorylation and capacitation are regulated by a cAMP-dependent pathway." In: *Development* 121.4 (Apr. 1995), pp. 1139–1150.

[13]   P. Martin-DeLeon. "Epididymosomes: Transfer of Fertility-modulating Proteins to the Sperm Surface". In: *Asian J. Androl.* 17.5 (2015), pp. 720–725. DOI: 10.4103/1008-682X.155538.

[14]   J.-J. Chung, S.-H. Shim, R. A. Everley, S. P. Gygi, X. Zhuang, and D. E. Clapham. "Structurally Distinct Ca2+ Signaling Domains of Sperm Flagella Orchestrate Tyrosine Phosphorylation and Motility". In: *Cell* 157.4 (May 2014), pp. 808–822. DOI: 10.1016/j.cell.2014.02.056.

[15]   J. Y. Hwang et al. "Dual Sensing of Physiologic pH and Calcium by EFCAB9 Regulates Sperm Motility". In: *Cell* 177.6 (May 2019), pp. 1480–1494. DOI: 10.1016/j.cell.2019.03.047.

[16]   C. Stival, L. d. C. Puga Molina, B. Paudel, M. G. Buffone, P. E. Visconti, and D. Krapf. "Sperm Capacitation and Acrosome Reaction in Mammalian Sperm". In: *Sperm Acrosome Biog. Funct. Dur. Fertil.* Ed. by M. G. Buffone. Cham, Switzerland: Springer International Publishing, 2016. Chap. 5, pp. 93–106. DOI: 10.1007/978-3-319-30567-7_5.

[17]   W. J. Blake, M. KÆrn, C. R. Cantor, and J. J. Collins. "Noise in eukaryotic gene expression". In: *Nature* 422.6932 (Apr. 2003), pp. 633–637. DOI: 10.1038/nature01546.

[18]   A. Sigal et al. "Variability and memory of protein levels in human cells". In: *Nature* 444.7119 (Nov. 2006), pp. 643–646. DOI: 10.1038/nature05316.

[19]   M. R. Birtwistle, A. V. Kriegsheim, M. Dobrzyński, B. N. Kholodenko, and W. Kolch. "Mammalian protein expression noise: Scaling principles and the implications for knockdown experiments". In: *Mol. Biosyst.* 8.11 (2012), pp. 3068–3076. DOI: 10.1039/c2mb25168j.

[20] J. Beal. "Biochemical complexity drives log-normal variation in genetic expression". In: *Eng. Biol.* 1.1 (June 2017), pp. 55–60. DOI: 10.1049/enb.2017.0004.

[21] A. Romarowski et al. "A Specific Transitory Increase in Intracellular Calcium Induced by Progesterone Promotes Acrosomal Exocytosis in Mouse Sperm1". In: *Biol. Reprod.* 94.3 (2016), 63:1–63:12. DOI: 10.1095/biolreprod.115.136085.

[22] G. M. Luque et al. "Only a subpopulation of mouse sperm displays a rapid increase in intracellular calcium during capacitation". In: *J. Cell. Physiol.* 233.12 (2018), pp. 9685–9700. DOI: 10.1002/jcp.26883.

[23] L. C. P. Molina, S. Gunderson, J. Riley, P. Lybaert, A. Borrego-Alvarez, E. S. Jungheim, and C. M. Santi. "Membrane Potential Determined by Flow Cytometry Predicts Fertilizing Ability of Human Sperm". In: *Front. Cell Dev. Biol.* 7 (Jan. 2020), 387:1–387:12. DOI: 10.3389/fcell.2019.00387.

[24] L. C. Puga Molina, G. M. Luque, P. A. Balestrini, C. I. Marín-Briggiler, A. Romarowski, and M. G. Buffone. "Molecular Basis of Human Sperm Capacitation". In: *Front. Cell Dev. Biol.* 6 (July 2018), 72:1–72:23. DOI: 10.3389/fcell.2018.00072.

[25] S. Kauffman. "Metabolic stability and epigenesis in randomly constructed genetic nets". In: *J. Theor. Biol.* 22.3 (Mar. 1969), pp. 437–467. DOI: 10.1016/0022-5193(69)90015-0.

[26] L. Sánchez, J. Van Helden, and D. Thieffry. "Establishment of the dorso-ventral pattern during embryonic development of Drosophila melanogaster: A logical analysis". In: *J. Theor. Biol.* 189.4 (1997), pp. 377–389. DOI: 10.1006/jtbi.1997.0523.

[27] E. Balleza, E. R. Alvarez-Buylla, A. Chaos, S. Kauffman, I. Shmulevich, and M. Aldana. "Critical Dynamics in Genetic Regulatory Networks: Examples from Four Kingdoms". In: *PLoS One* 3.6 (June 2008). Ed. by S. Huang, e2456. DOI: 10.1371/journal.pone.0002456.

[28] J. Espinal, M. Aldana, A. Guerrero, C. Wood, A. Darszon, and G. Martínez-Mekler. "Discrete Dynamics Model for the Speract-Activated Ca2+ Signaling Network Relevant to Sperm Motility". In: *PLoS One* 6.8 (Aug. 2011). Ed. by C. V. Rao, e22619. DOI: 10.1371/journal.pone.0022619.

[29] J. G. T. Zañudo and R. Albert. "Cell Fate Reprogramming by Control of Intracellular Network Dynamics". In: *PLOS Comput. Biol.* 11.4 (Apr. 2015). Ed. by D. Thieffry, e1004193. DOI: 10.1371/journal.pcbi.1004193. arXiv: 1408.5628.

[30] B. C. Daniels et al. "Criticality Distinguishes the Ensemble of Biological Regulatory Networks". In: *Phys. Rev. Lett.* 121.13 (Sept. 2018), 138102:1–138102:6. DOI: 10.1103/PhysRevLett.121.138102.

[31] M. Huth and M. Ryan. "Binary decision diagrams". In: *Log. Comput. Sci. Model. Reason. about Syst.* 2nd ed. New York, NY, USA: Cambridge University Press, 2004. Chap. 6, pp. 358–413.

[32] J. G. T. Zañudo, M. Aldana, and G. Martínez-Mekler. "Boolean Threshold Networks: Virtues and Limitations for Biological Modeling". In: *Inf. Process. Biol. Syst.* Ed. by S. Niiranen and A. Ribeiro. Berlin, Heidelberg: Springer Berlin Heidelberg, 2011. Chap. 6, pp. 113–151. DOI: 10.1007/978-3-642-19621-8_6.

[33] C. Chaouiya, O. Ourrad, and R. Lima. "Majority Rules with Random Tie-Breaking in Boolean Gene Regulatory Networks". In: *PLoS One* 8.7 (July 2013). Ed. by J. Gomez-Gardenes, e69626. DOI: 10.1371/journal.pone.0069626.

[34] A. M. Salicioni, M. D. Platt, E. V. Wertheimer, E. Arcelay, A. Allaire, J. Sosnik, and P. E. Visconti. "Signalling pathways involved in sperm capacitation." In: *Soc. Reprod. Fertil. Suppl.* 65 (Jan. 2007), pp. 245–259.

[35] P. E. Visconti. "Understanding the molecular basis of sperm capacitation through kinase design." In: *Proc. Natl. Acad. Sci. U. S. A.* 106.3 (2009), pp. 667–668. DOI: 10.1073/pnas.0811895106.

[36] W. S. McCulloch and W. Pitts. "A logical calculus of the ideas immanent in nervous activity: The bulletin of mathematical biophysics". In: *Bull. Math. Biophys.* 5.4 (1943), pp. 115–133. DOI: 10.1007/BF02478259.

[37] A. L. Hodgkin and A. F. Huxley. "A quantitative description of membrane current and its application to conduction and excitation in nerve". In: *J. Physiol.* 117.4 (Aug. 1952), pp. 500–544. DOI: 10.1113/jphysiol.1952.sp004764.

[38] C. Sánchez-Cárdenas, M. R. Servín-Vences, O. José, C. L. Treviño, A. Hernández-Cruz, and A. Darszon. "Acrosome Reaction and Ca2+ Imaging in Single Human Spermatozoa: New Regulatory Roles of [Ca2+]i1". In: *Biol. Reprod.* 91.3 (Sept. 2014), 67:1–67:13. DOI: 10.1095/biolreprod.114.119768.

[39] K. Kerns, M. Zigo, E. Z. Drobnis, M. Sutovsky, and P. Sutovsky. "Zinc ion flux during mammalian sperm capacitation". In: *Nat. Commun.* 9.1 (Dec. 2018), 2061:1–2061:10. DOI: 10.1038/s41467-018-04523-y.

[40] J. Xia and D. Ren. "The BSA-induced Ca(2+) influx during sperm capacitation is CATSPER channel-dependent". In: *Reprod. Biol. Endocrinol.* 7.1 (Dec. 2009), 119:1–119:9. DOI: 10.1186/1477-7827-7-119.

[41] G. Orta, J. L. De La Vega-Beltran, X. D. Martín-Hidalgo, C. M. Santi, P. E. Visconti, and X. A. Darszon. "CatSper channels are regulated by protein kinase A". In: *J. Biol. Chem.* 293.43 (2018), pp. 16830–16841. DOI: 10.1074/jbc.RA117.001566.

[42] I. A. Demarco et al. "Involvement of a Na + /HCO Cotransporter in Mouse Sperm Capacitation". In: *J. Biol. Chem.* 278.9 (Feb. 2003), pp. 7001–7009. DOI: 10.1074/jbc.M206284200.

[43] Y. Zeng, J. a. Oberdorf, and H. M. Florman. "pH Regulation in Mouse Sperm: Identification of Na+-, Cl–, and[formula]Dependent and Arylaminobenzoate-Dependent Regulatory Mechanisms and Characterization of Their Roles in Sperm Capacitation". In: *Dev. Biol.* 173.2 (Feb. 1996), pp. 510–520. DOI: 10.1006/dbio.1996.0044.

[44] Y. Zeng, E. N. Clark, and H. M. Florman. *Sperm Membrane Potential: Hyperpolarization during Capacitation Regulates Zona Pellucida-Dependent Acrosomal Secretion.* 1995. DOI: 10.1006/dbio.1995.1304.

[45] J. Escoffier, F. Navarrete, D. Haddad, C. M. Santi, A. Darszon, and P. E. Visconti. "Flow Cytometry Analysis Reveals That Only a Subpopulation of Mouse Sperm Undergoes Hyperpolarization During Capacitation1". In: *Biol. Reprod.* 92.5 (May 2015), 121:1–121:11. DOI: 10.1095/biolreprod.114.127266.

[46] J. C. Chávez et al. "SLO3 K + Channels Control Calcium Entry through CATSPER Channels in Sperm". In: *J. Biol. Chem.* 289.46 (Nov. 2014), pp. 32266–32275. DOI: 10.1074/jbc.M114.607556.

[47] M. A. Nolan, D. F. Babcock, G. Wennemuth, W. Brown, K. A. Burton, and G. S. McKnight. "Sperm-specific protein kinase A catalytic subunit C 2 orchestrates cAMP signaling for male fertility". In: *Proc. Natl. Acad. Sci.* 101.37 (Sept. 2004), pp. 13483–13488. DOI: 10.1073/pnas.0405580101.

[48] W. M. Xu et al. "Cystic fibrosis transmembrane conductance regulator is vital to sperm fertilizing capacity and male fertility". In: *Proc. Natl. Acad. Sci.* 104.23 (June 2007), pp. 9816–9821. DOI: 10.1073/pnas.0609253104.

[49] J. C. Chávez et al. "Ion Permeabilities in Mouse Sperm Reveal an External Trigger for SLO3-Dependent Hyperpolarization". In: *PLoS One* 8.4 (Apr. 2013). Ed. by A. J. Travis, e60578. DOI: 10.1371/journal.pone.0060578.

[50] X.-H. Zeng, C. Yang, S. T. Kim, C. J. Lingle, and X.-M. Xia. "Deletion of the Slo3 gene abolishes alkalization-activated K+ current in mouse spermatozoa". In: *Proc. Natl. Acad. Sci.* 108.14 (Apr. 2011), pp. 5879–5884. DOI: 10.1073/pnas.1100240108.

[51] E. O. Hernández-González et al. "Sodium and Epithelial Sodium Channels Participate in the Regulation of the Capacitation-associated Hyperpolarization in Mouse Sperm". In: *J. Biol. Chem.* 281.9 (Mar. 2006), pp. 5623–5633. DOI: 10.1074/jbc.M508172200.

[52] P. Martínez-López et al. "Mouse sperm K+ currents stimulated by pH and cAMP possibly coded by Slo3 channels". In: *Biochem. Biophys. Res. Commun.* 381.2 (Apr. 2009), pp. 204–209. DOI: 10.1016/j.bbrc.2009.02.008.

[53] C. Stival et al. "Src Kinase Is the Connecting Player between Protein Kinase A (PKA) Activation and Hyperpolarization through SLO3 Potassium Channel Regulation in Mouse Sperm". In: *J. Biol. Chem.* 290.30 (July 2015), pp. 18855–18864. DOI: 10.1074/jbc.M115.640326.

[54] L. C. Puga Molina et al. "Essential Role of CFTR in PKA-Dependent Phosphorylation, Alkalinization, and Hyperpolarization During Human Sperm Capacitation". In: *J. Cell. Physiol.* 232.6 (June 2017), pp. 1404–1414. DOI: 10.1002/jcp.25634.

[55] A. A. Faisal, L. P. J. Selen, and D. M. Wolpert. "Noise in the nervous system". In: *Nat. Rev. Neurosci.* 9.4 (Apr. 2008), pp. 292–303. DOI: 10.1038/nrn2258.

[56] Z. Nusser, S. Cull-Candy, and M. Farrant. "Differences in synaptic GABA(A) receptor number underlie variation in GABA mini amplitude". In: *Neuron* 19.3 (1997), pp. 697–709. DOI: 10.1016/S0896-6273(00)80382-7.

[57] P. Caballero-Campo, M. G. Buffone, F. Benencia, J. R. Conejo-García, P. F. Rinaudo, and G. L. Gerton. "A Role for the Chemokine Receptor CCR6 in Mammalian Sperm Motility and Chemotaxis". In: *J. Cell. Physiol.* 229.1 (2014), pp. 68–78. DOI: 10.1002/jcp.24418.

[58] N. Bernabò, M. Mattioli, and B. Barboni. "The spermatozoa caught in the net: the biological networks to study the male gametes post-ejaculatory life". In: *BMC Syst. Biol.* 4.1 (Dec. 2010), 87:1–87:12. DOI: 10.1186/1752-0509-4-87.

[59] N. Bernabo, M. Mattioli, and B. Barboni. "Computational Modeling of Spermatozoa Signal Transduction Pathways: Just a Computer Game or a Reliable Tool in Studying Male Gametes Function?" In: *J. Comput. Sci. Syst. Biol.* 06.04 (2013), pp. 194–205. DOI: 10.4172/jcsb.1000117.

[60] N. Bernabò, L. Greco, A. Ordinelli, M. Mattioli, and B. Barboni. "Capacitation-Related Lipid Remodeling of Mammalian Spermatozoa Membrane Determines the Final Fate of Male Gametes: A Computational Biology Study". In: *Omi. A J. Integr. Biol.* 19.11 (2015), pp. 712–721. DOI: 10.1089/omi.2015.0114.

[61] Bernabò et al. "Cyclin–CDK Complexes are Key Controllers of Capacitation-Dependent Actin Dynamics in Mammalian Spermatozoa". In: *Int. J. Mol. Sci.* 20.17 (Aug. 2019), 4236:1–4236:17. DOI: 10.3390/ijms20174236.

[62] A. Taraschi et al. "Two-Player Game in a Complex Landscape: 26S Proteasome, PKA, and Intracellular Calcium Concentration Modulate Mammalian Sperm Capacitation by Creating an Integrated Dialogue—A Computational Analysis". In: *Int. J. Mol. Sci.* 21.17 (Aug. 2020), 6256:1–6256:11. DOI: 10.3390/ijms21176256.

[63] G. Orta, G. Ferreira, O. José, C. L. Treviño, C. Beltrán, and A. Darszon. "Human spermatozoa possess a calcium-dependent chloride channel that may participate in the acrosomal reaction". In: *J. Physiol.* 590.11 (June 2012), pp. 2659–2675. DOI: 10.1113/jphysiol.2011.224485.

[64] N. Mundt, M. Spehr, and P. V. Lishko. "TRPV4 is the temperature-sensitive ion channel of human sperm". In: *Elife* 7 (July 2018), e35853. DOI: 10.7554/eLife.35853.

[65] H. Tateno, D. Krapf, T. Hino, C. Sanchez-Cardenas, A. Darszon, R. Yanagimachi, and P. E. Visconti. "Ca2+ ionophore A23187 can make mouse spermatozoa capable of fertilizing in vitro without activation of cAMP-dependent phosphorylation pathways". In: *Proc. Natl. Acad. Sci.* 110.46 (Nov. 2013), pp. 18543–18548. DOI: 10.1073/pnas.1317113110.

[66] J. C. Chávez, E. O. Hernández-González, E. Wertheimer, P. E. Visconti, A. Darszon, and C. L. Treviño. "Participation of the Cl-/HCO3- Exchangers SLC26A3 and SLC26A6, the Cl- Channel CFTR, and the Regulatory Factor SLC9A3R1 in Mouse Sperm Capacitation1". In: *Biol. Reprod.* 86.1 (Jan. 2012), 14:1–14:14. DOI: 10.1095/biolreprod.111.094037.

[67] D. A. Priego-Espinosa, A. Darszon, A. Guerrero, A. L. González-Cota, T. Nishigaki, G. Martínez-Mekler, and J. Carneiro. "Modular analysis of the control of flagellar Ca2+-spike trains produced by CatSper and CaV channels in sea urchin sperm". In: *PLOS Comput. Biol.* 16.3 (Mar. 2020). Ed. by D. Umulis, e1007605. DOI: 10.1371/journal.pcbi.1007605.

[68] Y. Kirichok, B. Navarro, and D. E. Clapham. "Whole-cell patch-clamp measurements of spermatozoa reveal an alkaline-activated Ca2+ channel". In: *Nature* 439.7077 (Feb. 2006), pp. 737–740. DOI: 10.1038/nature04417.

[69] C. Yang, X.-H. Zeng, Y. Zhou, X.-M. Xia, and C. J. Lingle. "LRRC52 (leucine-rich-repeat-containing protein 52), a testis-specific auxiliary subunit of the alkalization-activated Slo3 channel". In: *Proc. Natl. Acad. Sci.* 108.48 (Nov. 2011), pp. 19419–19424. DOI: 10.1073/pnas.1111104108.

[70] M. Schreiber, A. Wei, A. Yuan, J. Gaut, M. Saito, and L. Salkoff. "Slo3, a Novel pH-sensitive K + Channel from Mammalian Spermatocytes". In: *J. Biol. Chem.* 273.6 (Feb. 1998), pp. 3509–3516. DOI: 10.1074/jbc.273.6.3509.

[71] B. Navarro, Y. Kirichok, and D. E. Clapham. "KSper, a pH-sensitive K + current that controls sperm membrane potential". In: *Proc. Natl. Acad. Sci.* 104.18 (May 2007), pp. 7688–7692. DOI: 10.1073/pnas.0702018104.

[72] X.-H. Zeng, B. Navarro, X.-M. Xia, D. E. Clapham, and C. J. Lingle. "Simultaneous knockout of Slo3 and CatSper1 abolishes all alkalization- and voltage-activated current in mouse spermatozoa". In: *J. Gen. Physiol.* 142.3 (Sept. 2013), pp. 305–313. DOI: 10.1085/jgp.201311011.

[73] B. Zerhusen, J. Zhao, J. Xie, P. B. Davis, and J. Ma. "A single conductance pore for chloride ions formed by two cystic fibrosis transmembrane conductance regulator molecules". In: *J. Biol. Chem.* 274.12 (1999), pp. 7627–7630. DOI: 10.1074/jbc.274.12.7627.

[74] D. Figueiras-Fierro et al. "Electrophysiological evidence for the presence of cystic fibrosis transmembrane conductance regulator (CFTR) in mouse sperm". In: *J. Cell. Physiol.* 228.3 (Mar. 2013), pp. 590–601. DOI: 10.1002/jcp.24166. arXiv: NIHMS150003.

[75] C. M. Canessa, L. Schild, G. Buell, B. Thorens, I. Gautschi, J.-d. Horisberger, and B. C. Rossier. "Amiloride-sensitive epithelial Na+ channel is made of three homologous subunits". In: *Nature* 367.6462 (Feb. 1994), pp. 463–467. DOI: 10.1038/367463a0.

[76] M. F. Romero and W. F. Boron. "Electrogenic Na + /HCO - 3 Cotransporters: Cloning and Physiology". In: *Annu. Rev. Physiol.* 61.1 (Mar. 1999), pp. 699–723. DOI: 10.1146/annurev.physiol.61.1.699.

[77] S. D. Olson, S. S. Suarez, and L. J. Fauci. "A Model of CatSper Channel Mediated Calcium Dynamics in Mammalian Spermatozoa". In: *Bull. Math. Biol.* 72.8 (Nov. 2010), pp. 1925–1946. DOI: 10.1007/s11538-010-9516-5.

[78] R. A. Harrison. "Rapid PKA-Catalysed Phosphorylation of Boar Sperm Proteins Induced by the Capacitating Agent Bicarbonate". In: *Mol. Reprod. Dev.* 67.3 (2004), pp. 337–352. DOI: 10.1002/mrd.20028.

Supplementary material

S1 Supplementary figures

S1.1 Time series used in model validation

Figure S1 shows the average, over sperm population, of the behavior of simulation time series for selected variables under external stimulation, in the absence of ion transporter weight variability. Three types of stimuli are considered: individual and joint addition of cholesterol acceptors and increase in the external medium of bicarbonate levels. The entries of Tab. 1 summarize qualitative trends displayed in this figure. Figure S2 shows that the corresponding behavior in the presence of ion transporter weight variability with $D = 0.25$ is unaltered, the figures almost overlap. Figs. S3 to S7 show for $D = 0.25$, a comparison between time series with a loss of function variable and the WT, both subject only to the joint external stimulation mentioned above. A summary of the trends observed in these figures is presented in Tab. 2 of Sec. 2.2.1.

Figure S1: Averaged time series of a select set of variables from a WT sperm population, without variability in their ion transporter weights, subject to external stimulation. In the simulations stimuli are introduced at time $t = 5 \times 10^3$ and consist of cholesterol acceptor-only (first column), bicarbonate-only (second column), or both (third column). The qualitative trends of each variable were used to validate our model and are summarized in Tab. 1 of Sec. 2.2.1 . Population size is 2×10^3 sperm.

S1.2 Joint node capacitation threshold exploration

Fig. S8 and Fig. S9 show additional capacitation fraction surfaces as shown in Fig. 5 for values of $\theta = 0.175$ and $\theta = 0.250$ respectively.

Figure S2: Averaged time series of a selected set of variables from a WT sperm population, with variability in ion transport weights sampled from a Gaussian distribution with standard deviation $D = 0.25$, under several stimuli. Simulations shown in the panels are performed as in Fig. S1. Notice the similarity of both figures, trends are preserved under the inclusion of the above mentioned variability.

Figure S3: Averaged time series with variability, under an external stimulation of the addition of cholesterol acceptors and higher bicarbonate levels in the extracellular medium, of a selected set of variables under a CatSper$^{\text{LOF}}$ network variant (left column) and for WT (right column). Under CatSper$^{\text{LOF}}$, membrane potential V hyperpolarizes, Ca$_i$ goes to basal levels, Na$_i$ decreases, whereas Cl$_i$, pH$_i$ and PKA activity increase. Population size is 2×10^3 sperm, the standard deviation used in introducing variability on ion transporter weights is $D = 0.25$.

Figure S4: Comparison of the averaged time series with variability, under an external stimulation of the addition of cholesterol acceptors and higher bicarbonate levels in the extracellular medium, of select nodes under loss of function (LOF) of CFTR/ClC (first column) and WT conditions (second column). Notice that under LOF membrane potential V hyperpolarizes, Na_i goes to basal levels, Ca_i, Cl_i, pH_i and PKA activity increase. Population size is 2×10^3 sperm, the standard deviation used for variability on ion transporter weights is $D = 0.25$.

Figure S5: Averaged time series with variability, under an external stimulation of the addition of cholesterol acceptors and higher bicarbonate levels in the extracellular medium, of select set of variables under NaC^{LOF} (first column) and WT conditions. Membrane potential V hyperpolarizes, Na_i decreases, Ca_i, Cl_i, pH_i and PKA activity increase. Population size is 2×10^3 sperm, the standard deviation used in introducing variability on ion transporter weights is $D = 0.25$.

Figure S6: Averaged time series of select set of variables with variability, under an external stimulation of the addition of cholesterol acceptors and higher bicarbonate levels in the extracellular medium, with $Slo3^{LOF}$ (first column) and WT conditions (second column). Membrane potential V depolarizes, Na_i decreases, Ca_i, Cl_i, pH_i and PKA activity increases. Population size is 2×10^3 sperm, the standard deviation used in introducing variability on ion transporter weights is $D = 0.25$.

Figure S7: Averaged time series with variability, under an external stimulation of the addition of cholesterol acceptors and higher bicarbonate levels in the extracellular medium, of select set of variables under PKA$^{\text{LOF}}$ (first column) and WT conditions (second column). Membrane potential V depolarizes, Na$_i$, Ca$_i$, Cl$_i$ and pH$_i$ increase, PKA activity goes to zero.. Population size is 2×10^3 sperm, the standard deviation used for introducing variability on ion transporter weights is $D = 0.25$.

Figure S8: Capacitated fraction (CF) in sperm populations for fixed joint threshold $\theta = 0.175$, varying three parameters; observation window size (W, in iteration units), standard deviation of Gaussian variability (D) and activity threshold of joint node (θ_c). In all surfaces, each data point comes from 2×10^3-sperm sized populations with 2×10^4 iterations long simulations (5×10^3 iterations at resting state and 1.5×10^4 post stimulus iterations). The classification of capacitated vs. non-capacitated sperm is applied on the last 10^4 time iterations of each individual sperm. The color bar represents the level of capacitation fraction. Note that the white zone corresponds to a capacitation level of $30 \pm 5\%$, which is close to the levels typically observed for *in vitro* capacitation in wild-type sperm.

Figure S9: Capacitation fraction (CF) surfaces in sperm populations, for fixed joint threshold $\theta = 0.250$, determined as in above figure S8

.