Predicted distribution of a rare and understudied forest carnivore: Humboldt martens

(*Martes caurina humboldtensis*)

Katie Moriarty[1], Joel Thompson[2], Matthew Delheimer[3], Brent Barry[4], Mark Linnell[5], Taal

Levi[6], Keith Hamm[7], Desiree Early[7], Holly Gamblin[8], Micaela Szykman Gunther[8], Jordan

Ellison[1], Janet S. Prevéy[9], Jennifer Hartman[10], Ray Davis[11]

[1] Western Sustainable Forestry, National Council for Air and Stream Improvement, Inc.,

Corvallis, Oregon, USA

[2] Pacific Northwest Region Data Resources Management, USDA Forest Service,

Joseph, Oregon, USA

[3] Pacific Southwest Research Station, USDA Forest Service, Placerville, California,

USA

[4] Confederated Tribes of the Grand Ronde, Grand Ronde, Oregon, USA

[5] Pacific Northwest Research Station, USDA Forest Service, Corvallis, Oregon, USA

[6] Department or Fisheries and Wildlife, Oregon State University, Corvallis, Oregon, USA

[7] Green Diamond Resource Company, Korbel, California, USA

[8] Department of Wildlife, Humboldt State University, Arcata, California, USA

[9] Pacific Northwest Research Station, USDA Forest Service, Olympia, Washington,

USA

[10] Rogue Detection Teams, Rice, Washington, USA

[11] Pacific Northwest Region Monitoring Program, USDA Forest Service, Corvallis,

Oregon, USA

Corresponding Author:

Katie Moriarty[1]

NCASI, 227 NW Third Street, Corvallis, OR, 97330, USA

Email address: kmoriarty@ncasi.org

Abstract

Background. A suite of mammalian species have experienced range contractions following European settlement and post-settlement development of the North American continent. For example, while North American martens (American marten, *Martes americana*; Pacific marten, *M. caurina*) generally have a broad range across northern latitudes, local populations have experienced substantial reductions in distribution and some extant populations are small and geographically isolated. The Humboldt marten (*M. c. humboldtensis*), a subspecies of Pacific marten that occurs in coastal Oregon and northern California, was recently designated as federally threatened in part due to its reduced distribution. To inform strategic conservation actions, we assessed Humboldt marten occurrence by compiling all known records from their range.

Methods. We compiled Humboldt marten locations since their rediscover to present (1,692 marten locations, 1996-2020). We spatially-thinned locations to 500-m to assess correlations with variables across contemporary Humboldt marten distribution (n=384). Using maximum entropy modeling (Maxent), we created distribution models with variables optimized for spatial scale; pre-selected scales were associated with marten ecology (50 to 1170 m radius). Marten locations were most correlated with abiotic factors (e.g., precipitation), which are unalterable and therefore uninformative within the context of restoration or management actions. Thus, we created variables to focus on hypothesized marten habitat relationships, including understory conditions such as predicted suitability of shrub species.

Results. Humboldt marten locations were positively associated with increased shrub cover (salal (*Gautheria shallon*), mast producing trees), increased pine (*Pinus sp*) overstory cover and precipitation at home-range spatial scales, areas with low and high amounts of canopy cover and slope, and cooler August temperatures. Unlike other recent literature on the species, we found little evidence that Humboldt marten locations were associated with old growth structural indices, perhaps because of a potential mismatch in the association between this index and

shrub cover. As with any species distribution model, there were gaps in predicted distribution

where Humboldt martens have been located during more recent surveys, for instance the

southeastern portion of Oregon's coast range. Conservation efforts and our assessment of

potential risks to Humboldt marten populations would benefit from additional information on

range extent, population sizes, and fine-scale habitat use. Like many rare and lesser-known

species, this case study provides an example of how limited information can provide differing

interpretations, emphasizing the need for study-level replication in ecology.

Key words: California, distribution model, habitat relationships, Humboldt marten, *Martes*

caurina humboltensis, Maxent, Oregon, Pacific marten

Introduction

Modeling predicted distributions of rare or declining species can direct conservation

efforts, but creating accurate predictions is challenging for a multitude of reasons.

Contemporary location information may associate the species with conditions that were

unaffected by prior agents of population decline, but not with favored characteristics

where the species resided prior (Caughley 1994). For instance, bison (*Bison bison*)

were historically widely distributed throughout western North America with herd sizes up

to tens of millions of individuals (Shaw 1995), but species distribution models based on

their contemporary distribution might suggest that they are associated with conditions

present in Yellowstone National Park, such as extremely cold winters and thermal

geysers. Instead, in this example, a logical conclusion is that bison in this area received

less persecution. Nonetheless, a range-wide and broad-scale contemporary species

distribution model ignorant of their biology and the causes of bison declines would

misrepresent predicted habitat; creating more thermal geysers would not equate to

more bison.

For populations that have declined or are currently declining, modeling species

distributions may be complicated by additional factors – for example, contemporary

locations may not include all areas where a species could physically colonize in the

future, if the species is constrained by mobility or time (Barry et al. In review) or barriers

that were not in place prior to the species' decline (e.g., human development; Brown et

al. 1996). Species may also require resources at multiple spatial scales, during different

seasons, or for different durations of time – a female desert bighorn sheep (*Ovis*

canadensis nelsoni) requires high nutritional-value vegetation during spring when

providing for young (fine spatial extent, short duration), a water source within her range

during the heat of the summer (large spatial extent, short duration), and immediate

cover from predation year-round (fine spatial extent, long duration; e.g., Gedir et al.

2020; Hoglander et al. 2015). As such, limited observations in space or time may miss

the conditions necessary for population persistence. Despite challenges for

understudied species (Raphael & Molina 2007), spatial models can predict conditions of

suitability that allow for species' occurrence, which could identify areas of potential

occupancy that are unknown due to lack of survey effort (Sofaer et al. 2019).

Modeling habitat and predicted distribution of North American marten populations

(American marten, *Martes americana*; Pacific marten, *M. caurina*) is particularly

challenging, due to presumed selection of both broad-scale landscape features (1st

order selection sensu Johnson 1980) and fine-scale features within home ranges (4th

order selection; e.g., Minta et al. 1999). Martens are small-bodied (600-1300g)

carnivores that occupy large home ranges relative to their body size and circumnavigate

the perimeter of their territory often (e.g., every 4-5 days, Balharry 1993; Moriarty et al.

2017). To fulfill some life history requirements, martens select fine-scale forest elements

that facilitate foraging, daily resting and seasonal denning activities, and allow them to

avoid predators (Andruskiw et al. 2008; Cheveau et al. 2013; Joyce 2013; Shirk et al.

2012). Simultaneously, fulfillment of other life history requirements (e.g., size and

positioning of home ranges relative to intraspecific competitors or potential mates)

requires selection for broader-scale forest attributes such as cohesive and complex

vegetation cover (Hargis et al. 1999; Potvin et al. 2000); as such, martens have been

highlighted as a species benefiting from multi-scale habitat selection models (Bissonette

& Broekhuizen 1995; Shirk et al. 2014; Wasserman et al. 2010).

Humboldt martens (*M. c. humboldtensis*) are a subspecies of Pacific marten that

historically occurred throughout coastal forests of northern California and Oregon.

Contemporary range-wide surveys suggest that distribution of Humboldt martens is

substantially reduced compared to historical representations (Moriarty et al. 2019;

Zielinski et al. 2001). Consequently, Humboldt martens were listed as Endangered

under the state of California's Endangered Species Act (CDFW 2019) and as

threatened under the federal Endangered Species Act as a "coastal distinct population

segment of Pacific martens" (USFWS 2020). Evaluating current and future Humboldt

marten distribution is urgently needed for conservation planning, yet modeling their

distribution is constrained by a paucity of ecological information (e.g., population sizes

and extents) and apparent non-stationary associations with vegetation among

populations. For example, Humboldt martens have been associated with a diversity of

forest types across their distribution, including mature forests with high canopy cover,

mature forests with dense shrub understories (Slauson et al. 2007), young forests (<80

years old) with modest canopy cover and relatively small diameter trees with dense

shrub cover (Eriksson et al. 2019; Moriarty et al. 2019), and forests in serpentine soils

with sparse canopy cover and dense shrub cover. A dense and spatially-extensive

shrub layer appears to be an attribute shared by each of these studies (Eriksson et al.

2019; Gamblin 2019; Moriarty et al. 2019; Slauson et al. 2007).

A previous range-wide Humboldt marten distribution model Slauson et al.

(2019b) emphasized a strong correlation between Humboldt marten occurrence and an

"old-growth structural index" (OGSI) variable, which is a composite index of factors

considered common to old-growth forests in the region, including density of large live

trees, snags and downed wood, and diversity of tree sizes (Ohmann 2012). However,

the model relied on modest number of detections from 1996–2010 with poor coverage

outside a portion of northern California (USFWS 2019). Since 2010, we initiated large-

scale surveys for Humboldt martens that greatly increased the spatial extent and

number of Humboldt marten detections in both California and Oregon (e.g., Barry 2018;

Gamblin 2019; Linnell et al. 2018; Moriarty et al. 2019). Further, recent research efforts

appear to contradict the finding that Humboldt martens are strongly associated with old-

growth forests, particularly in Oregon (e.g., Eriksson et al. 2019), suggesting a potential

mismatch in previously-predicted associations between OGSI and Humboldt marten

distribution. Such a mismatch could lead to a "wicked problem" by focusing

management or restoration in areas that would not benefit the species (Gutiérrez 2020).

For example, a landscape connectivity model for Humboldt martens determined both

"habitat cores" and connectivity between cores based almost exclusively on OGSI

values (Schrott & Shinn 2020).

Here, our objective was to create a contemporary range-wide model of predicted

Humboldt marten distribution facilitated by including recent location data collected from

broad-scale randomized surveys throughout the historic range, combined with more

recent and accurate vegetation layers (e.g., shrub layers). Our goal was to predict

factors contributing to probabilities of Humboldt marten spatial use and to highlight

areas for future surveys and conservation efforts.

Materials & Methods

Study Area

Humboldt martens are thought to occur in four populations in California and

Oregon, represented as the Central Coastal Oregon, Southern Coastal Oregon,

California-Oregon Border, and Northern Coastal California populations (USFWS 2019;

Fig. 1). Data were collected in Oregon in both near-coastal and montane areas (Oregon

Coast Range), where dominant forest types include Sitka spruce (*Picea sitchensis*) on

the coast and western hemlock (*Tsuga heterophylla*) (Franklin & Dyrness 1973). Sitka

spruce zones were mild (annual average temperature 10-11 °C), averaging 200-300 cm

of annual precipitation interspersed with regular fog and cloud cover. Western hemlock

zones, often co-dominated by Douglas-fir (*Pseudotsuga menziesii*), were are also wet

(150-300 cm annual precipitation) and somewhat cooler (7-10 °C) environments but

experience drier summers (6-9% of total precipitation) with fairly extensive summer fog

and low cloud cover (Dye et al. 2020). Surveys in California similarly occurred in both

coastal and montane areas (Klamath Mountains, California Coast Range), with

dominant forest types including redwood (*Sequoia sempervirens*) along the coast and

Douglas-fir or mixed-conifer in the mountains (Whittaker 1960). These areas included a

mix of coniferous and hardwood forest with a spatially-extensive shrub understory and

often received substantial precipitation (100-300 cm annual precipitation) with cooler (7-

10 °C) environments but with drier summers dominated with fog and low cloud moisture

(Rastogi et al. 2016). Dominant forest types include tanoak (*Notholithocarpus*

densiflora) in the southeastern portion of the northern California study area. Common

conifer species included western hemlock, Port Orford cedar (*Chamaecyparis*

lawsoniana), and western redcedar (*Thuja plicata*). Common hardwood species

included chinquapin (*Chrysolepis chrysophylla*), red alder (*Alnus rubra*), bigleaf maple

(*Acer macrophyllum*), and Pacific madrone (*Arbutus menziesii*). Dominant shrub

species throughout the range includes salal *(Gautheria shallon)*, evergreen huckleberry

(Vaccinium ovatum), Pacific rhododendron *(Rhododendron macrophyllum)*, and red

huckleberry *(V. parvifolium)*.

Marten locations

We incorporated verifiable, contemporary, spatially-referenced Humboldt marten

(hereafter, marten) locations collected during 1996-2020 into our model. We removed

marten locations from our model sample that occurred in areas that disturbance (fire or

harvest) between the data of observation and the date represented by our vegetation

layers (2016). If there were multiple locations within the 500-m cell, we spatially-thinned

locations prioritizing those in the order of: (1) rest and den locations from telemetry

studies in the Central Coastal Oregon population (Linnell et al. 2018) and the Northern

Coastal California population (Delheimer et al. In press; PSW 2019); (2) locations from

scat dog detection surveys (Rogue Detection Teams, Rice, WA; Conservation Canines,

University of Washington, Seattle, WA) from both Oregon populations (detailed methods

within Moriarty et al. 2018; Moriarty et al. 2019); and (3) locations from baited camera

and/or track plate surveys from all four populations in Oregon (Barry 2018; Moriarty et

al. 2019) and California (Gamblin 2019; Slauson et al. 2012). We thinned locations to

one within a 500-m cell, attempting to maximize distances between locations and

induce spatial independence for modeling (Kramer-Schadt et al. 2013). We used

presence-only data to incorporate telemetry-based information and detection dog

surveys in areas that outside the geographic extent of baited and lured detection/non-

detection surveys and because data on survey locations without detections and

detection histories were not available for older surveys (e.g., <2014).

For the data in which the authors were responsible, our protocols were reviewed

by the USDA Forest Service Research and Development or Humboldt State University's

Institutional for Use and Care Committee. We received approval for marten captures

and treatment in the Central Coast (USDA FS R&D 2015-002). We received a waiver

for camera trapping in Oregon by the same committee (USDA FS R&D 2017-005) and

camera surveys in the Border population were under Humboldt State University

16/17.W.05-A. We obtained Scientific Take Permits for hair snares and samples

collected through the Oregon Department of Fish and Wildlife (ODFW 119-15, 128-16,

033-16, 109-19, 107-20). Other verified survey data were provided by the US Fish and

Wildlife Service with no additional information.

Modeling approach

Our overall modeling approach incorporated our collated Humboldt marten locations,

biotic and abiotic predictor variables, and randomly-generated background points (n =

10,000). We used a minimum convex polygon around Humboldt marten locations

buffered by 10 km for the model training and testing (Fig. 1b). We chose a 10 km buffer

because this was approximately the upper quartile of daily marten movement (Moriarty

et al. 2017). We projected our model to available vegetation data from Gradient Nearest

Neighbor (GNN) 2016 data supplied by the Landscape Ecology, Modeling, Mapping and

Analysis lab (Bell et al. 2021; Bell et al. 2020), which included the coastal and Klamath

level-3 eco-provinces (U.S. Environmental Protection Agency 2013). We applied a mask

to only include areas typed for forested vegetation, which removed urban areas and

water, similar to Davis et al. (2016). We summarized the range, average, and standard

deviation for each variable at used marten locations, within the minimum convex

polygon (MCP) boundary within a 10-km buffer around known marten locations, and the

overall study area (Table 1, Fig. 1).

Biotic variables

We incorporated biotic variables into our models depicting forest structure and forest

composition (e.g., tree and shrub species). We acquired variables representing forest

structure through the 2016 version of GNN (Ohmann & Gregory 2002) and included

forest age, canopy cover, and OGSI. We depicted forest age as the basal area-

weighted age based on field-recorded or modeled ages of dominant and codominant

trees. Canopy cover percent was calculated using methods in the Forest Vegetation

Simulator (Crookston & Stage 1999). The OGSI variable is a composite index ranging

from 0-100 that is calculated from 4 elements: density of large diameter live trees per

hectare, density of large diameter snags per hectare, percentage of downed wood

greater than 25 cm in diameter, and an index of tree diameter diversity computed from

tree densities in different diameter classes. For live trees and snags, "large diameter" is

dependent on forest type and is defined for twelve vegetative zones, each with a unique

minimum diameter threshold (50-100 cm for live trees, 50-75 cm for snags). Detailed

methods for calculating the OGSI variable can be found in Davis et al. (2015); see

Supplemental information (Item S1) for more information on integration of the OGSI

variable into our model.

We created a variable called percent pine, which was a the combination of

percent total basal area of shore pine (*Pinus contorta*), Jeffery pine (*P. jefferii*), and

knobcone pine (*P. attenuata*) from GNN. We modeled this variable because martens

have been found with sparse shore pine communities in the Oregon Central Coast

population (Eriksson et al. 2019; Linnell et al. 2018), and areas with serpentine soils,

which are characterized by sparse tree cover of Jeffery and knobcone pines, stunted

tree growth, and dense shrub understories (Harrison et al. 2006; Kruckeberg 1984;

Safford et al. 2005). Serpentine soils have been associated with martens in northern

California (Slauson et al. 2019b). We visually inspected the serpentine soil layer created

by the US Fish and Wildlife Service with the percent pine layer (Schrott & Shinn 2020),

confirming overlap inland and inclusion of coastal areas.

263  Humboldt martens have been associated with dense shrub cover throughout

264 their range (Moriarty et al. 2019; Slauson et al. 2007); salal (*Gautheria shallon*) and

265 evergreen huckleberry (*Vaccinium ovatum*) appear particularly important, as the berries

266 of each occur in Humboldt marten diets and also provide food for marten prey items

267 (Eriksson et al. 2019; Moriarty et al. 2019). We modeled salal and evergreen

268 huckleberry occurrence and combined model outputs to create a shrub cover variable

269 (shrub), which was the sum of the probabilities of species occurrence of both respective

270 shrub rasters (Prevéy et al. 2020a; Prevéy et al. 2020b). We modeled our shrub cover

271 variable using Maxent, version 3.3 (Phillips & Dudík 2008) with detailed methods

272 published elsewhere (Prevéy et al. 2020a; Prevéy et al. 2020b). Briefly, these models

273 used locations from 10 sources, including FIA plots, USDA Forest Service Ecology and

274 Bureau of Land Management plots, iNaturalist, Oregon Flora Project, Consortium of

275 Pacific Northwest Herbaria, and the Global Biodiversity Information Forum. We related

276 locations to contemporary (1981-2010) bioclimatic variables from the AdaptWest project

277 (Wang et al. 2016) to depict the probability of species occurrence (1-100%), or expected

278 habitat quality, for each species (Prevéy et al. 2020a; Prevéy et al. 2020b). The

279 evergreen huckleberry model has not been published previously, but we created it using

280 the same methods as the published salal model (Prevéy et al. 2020a; Prevéy et al.

281 2020b).

282  Our final biotic variable (mast) represented hardwood tree and shrub species

283 that produce nuts, seeds, buds, or fruits eaten by wildlife. American marten population

284 numbers appear correlated with mast in hardwood forests of New York (Jensen et al.

285 2012); similarly, fishers (*Pekania pennanti*) appear associated with mast producing-

hardwoods in California (Townsend 2019). It is unclear whether Humboldt martens are

similarly associated, but their diet appears dominated by berries and animals that feed

on berries and seeds such as passerines and chipmunks (*Tamias sp.*) (Eriksson et al.

2019; Manlick et al. 2019; Slauson & Zielinski 2017). Here, our variable consisted of the

percent of basal area within 2016 GNN comprised of tanoak (*Notholithocarpus*

densiflorus), giant chinquapin (*Castanopsis chrysophylla*), coastal live oak (*Quercus*

agrifolia), canyon live oak (*Q. chrysolepis*), and California bay (*Umbellularia californica*).

Abiotic variables

We incorporated abiotic variables into our model including those depicting temperature

(°C), precipitation (cm), cloud cover (%), coastal proximity, percent slope, and

topographic position index. We used 30-year normal PRISM variables of Average

Annual Precipitation converted to cm and Maximum Temperature in August available at

an 800-m scale (1981-2010, PRISM Climate Group, Oregon State University,

http://prism.oregonstate.edu, created 10/17/2019). We explored annual data for

temperature (2010-2018), but the available 4 km resolution produced artifacts in the

model.

We created a coastal low cloud/fog variable for both the mean and interannual

standard deviation of coastal low cloudiness from May through September for 22 years

(1996–2017). The 4-km resolution coastal low cloudiness record was satellite derived

(NASA/NOAA Geostationary Environmental Satellite Imager measurements) as

described by Clemesha et al. (2016). We quantified coastal low cloudiness as the

percent of time that low clouds were present relative to the number of corresponding

valid half-hourly observations in a 24-hour day. We optimized and validated retrieved

data using coastal airports.

We created models with the variable Coastal Proximity, which uses PRISM data

and combines coastal proximity and temperature advection influenced by terrain (Daly

et al. 2003) modified for the western United States (Daly et al. 2008). We derived

percent slope and topographic position index from US Geological Survey digital

elevation models. Topographic position index is an indicator of slope position and

landform category; it is the difference between the elevation at a single cell and the

average elevation of the user-defined radius around that cell (Jenness 2006).

Scale optimization

We optimized spatial scales for each variable included in our models. We smoothed

variables using the extract function in package *raster* in R (Himjmans 2020; R Core

Team 2020) with a radius of 50 m, 270 m, 742 m, and 1170 m. Our smallest scale (50

m, 0.81 ha) provided localized conditions matching likely vegetation and location

accuracy. We assumed 270 m (20 ha) was approximately the size of a Humboldt

marten core use area, similar to optimized scales of vegetation characteristics used in

predicting conditions for marten rest structures elsewhere in California (Tweedy et al.

2019). We estimated home ranges for individuals with >30 locations and <400 sq m

expected location error (shifted locations were provided in Data S2; Lindenmayer &

Scheele 2017). Female Humboldt marten home range estimates averaged 0.8 km^2 (80

ha) in the Central Coast (n = 6 females, Linnell et al. 2018) and 2.4 km^2 (260 ha) using

a comparable minimum convex polygon estimate in northern California (n = 9 females,

PSW 2019). The average of all known telemetered Humboldt marten females was 1.73

km^2 (173 ha, Table S1) so we used the scale of 742 m (174 ha) as a representation of

female home range size. Our broadest scale was based on the largest size of a

Humboldt marten male home range (1170 m, 428 ha, Table S1), assuming a male

would overlap multiple females and could be interpreted as the smallest unit of

population level selection (Linnell et al. 2018; PSW 2019). We used univariate linear

models (glm) using our training location data (1) and a random background sample of

9,600 points within the MCP (0) with a binomial distribution. Similar to prior examples

(Cushman and McGarigal 2002, McGarigal et al. 2016, Wasserman et al. 2010), we

selected the scale for each variable that had the most extreme, and thus the most

predictive, coefficient. We visually inspected the fit of each spatial scale using boxplots

(Figs. S1, S2, S3).

We visually evaluated whether our final variables were similar between all marten

locations, thinned marten locations, surveyed locations without detections (non-

detection), and random locations.

Predicted distribution

We used MaxEnt modeling software v3.4.1 (Phillips et al. 2006) to estimate the

probability of Humboldt marten presence within the modeling regions (Merow et al.

2013). MaxEnt uses a machine learning process to develop algorithms that relate

environmental conditions at documented species presence locations to that of the

surrounding background environment in which they occurred (Elith et al. 2011; Phillips &

Dudík 2008). We excluded variables with highly correlated predictors (|Pearson

coefficient| > 0.5), selecting the variable that was most interpretable for managers

(Table S2). During this process, we considered the variance inflation (Table S3), which

allows for evaluation of correlation and multicollinearity. Variance inflation factors equal

to 1 are not correlated and factors greater than 5 are highly correlated as determined by

$(1/(1-R_i^2))$, where R_i^2 is squared multiple correlation of the variable i (Velleman &

Welsch 1981).

Within each model iteration, we selected the bootstrap option with 10 replicates,

random seed, and 500 iterations. We trained our models using a random subset of 75%

of presence locations and tested these using the remaining 25% with logistic output. We

used the default of 10,000 random background samples. We varied the response

functions to include linear, product, and quadratic features. We selected the "auto

features" option for all runs, which allows Maxent to further limit the subset of response

features from those selected by retaining only those with some effect.

Maxent provides two metrics to determine importance of input variables in the

final model: percent contribution and permutation importance. Percent

contribution measures the increase in likelihood associated with modeling rules based

on each environmental variable and then divides this by the total gain in likelihood to

calculate percent of the total gain associated with each variable (Elith et al. 2011).

Maxent keeps track of which environmental variables are contributing to fitting the

model, and at each step its algorithm increases the gain of the model by modifying the

coefficient for a single feature; the program assigns the increase in the gain to the

environmental variable(s) upon which that the feature depends (Phillips 2006). Percent

contribution converts these increases to percentages at the end of the training process

(Phillips 2006). Halvorsen (2013) produced simulation results suggesting percent

contribution can be more informative with uncorrelated environmental variables. This

metric is often used to assess variable significance (e.g., Warren et al. 2014).

Permutation importance randomizes the values for each environmental variable

between the presences and background points to make that variable uninformative and

then measures the resulting drop in area under the curve (AUC; ability to separate

known presences from background points). The bigger the drop, the more important

that variable was in the overall quality of the model and the resultant data are

normalized to percentages (Elith et al. 2011). The permutation importance measure

depends only on the final Maxent model, not on the path used to obtain it. Searcy &

Shaffer (2016) suggest that permutation importance provides better variable

assessment when models and variables are correlated.

Species distribution maps were produced from all models using the maximum

training sensitivity plus specificity threshold, which minimizes both false negatives and

false positives. We evaluated the AUC statistic to determine model accuracy and fit to

the testing data (Fielding & Bell 1997). The AUC statistic is a measure of the model's

predictive accuracy, producing an index value from 0.5 to 1, with values close to 0.5

indicating poor discrimination and a value of 1 indicating perfect predictions (Elith et al.

2006). We assessed variables using response curves, variable contributions, and

jackknife tests.

Because over-parameterized models tend to underestimate habitat availability

when transferred to a new geography or time period, we used selection methods

suggested by Warren & Seifert (2011). Maxent provides the option of reducing

overfitting with a regularization multiplier that can be altered by the user to apply a

penalty for each term included in the model (β regularization parameter) to prevent

overcomplexity or overfitting (Merow et al. 2013; Morales et al. 2017). A higher

regularization multiplier will reduce the number of covariates in the model, becoming

more lenient with an increased sample size (Merow et al. 2013). We did not include

model replicates, an option in the interface, to output the required data (lambda file) and

set output to logistic. We altered the Regularization Multiplier from 0.5 to 4 for each 0.5

increment (e.g., Radosavljevic & Anderson (2014).

We ranked candidate models using Akaike's Information Criterion corrected for

small sample sizes (AIC_c; Burnham & Anderson 2002). We calculated AIC_c using the

formula $2k - 2lnL + (2k(k+1))/(n - k - 1)$. We calculated log likelihood (lnL) by summing

the values in all raster cells of the output model, subtracting the log of each cell by the

log value of that sum, and extracting and summing the values at the occurrence points

(Warren & Seifert 2011). We defined *k* as the number of rows in the lambda files with

non-zero lambda values and we considered *n* the number of thinned marten locations.

We considered the model with the lowest AIC_c value to be the top model, calculated the

ΔAIC_c value (i.e., change in AIC_c value from the top model) for all other models, and

considered models with $\Delta AIC_c < 2$ to be competitive models.

For our top model, we generated predicted-to-expected (P/E) ratio curves for our

model using only the testing data to evaluate its predictive performance, which was

based on the shape of the curves, a continuous Boyce index (Boyce et al. 2002), and

Spearman rank statistics. We used the predicted-to-expected curve to inform our

suitability thresholds following Hirzel et al. (2006), including predicted unsuitable (P/E

and confidence intervals 0-1), marginal (P/E > 1 but overlapping confidence intervals),

and suitable (P/E and confidence intervals > 1).

Results

Locations

We compiled 1,692 Humboldt marten location records collected between 1996-2020

(baited stations = 542, detection dog = 263, VHF telemetry = 831, roadkill = 15, other =

41 locations). After we spatially-thinned contemporary locations to no more than one per

500 m^2, 384 locations remained and were generally evenly spread among regions in

proportion to their area - Central Coastal Oregon *(n* = 77 locations, 6% of the

designated Extant Population Area)*,* Southern Coastal Oregon (*n* = 77 locations, 37% of

the EPA), California-Oregon Border (*n* = 33 locations, 3% of the EPA), and Northern

Coastal California (*n* = 192 locations, 54% of the EPA) populations, but also included 5

locations that did not occur within the boundaries of these designated populations

(USFWS 2019). Location types included den or rest structure locations (18%),

genetically-verified scats or telemetry locations (32%), and baited camera or track plate

locations (50%).

Our thinned sample had similar medians and data distributions to all

contemporary locations with the exception of mast and precipitation, where the medians

were slightly lower for our thinned sample. Similarly, surveyed locations without marten

detections and random locations were comparable with the most notable different

between the medians for salal (Table 1, Fig. 3). Differences between the non-detection

and random locations were likely due to clustered sampling efforts (Fig. 1b).

Distribution modeling

Our model included 8 variables with <60% correlation (Table S2) with variance inflation

factors <5 (Table S3). All variables in our model had an optimized spatial scale at our

largest extent (1,170 m) except OGSI (50 m), suggesting these variables were most

closely matched to martens at the home range scale or fine-scale. Our top model had a

Regularization Multiplier of 1.5. The top model, in order of percent contribution, included

a positive relationship with salal (23.3%), percent pine (22.5%), average annual

precipitation (21.6%), canopy cover (18.7%), and mast (5.4%) followed by a negative

relationship with average maximum August temperature (4.7%), percent slope (2.7%),

and OGSI (1.1%, Table 2). The permutation importance was similar with the top four

variables highly contributing - but with a slightly modified order of percent pine (30.3%),

average annual precipitation (25.3%), canopy cover (20.2%), and salal (15.5%) for the

top variables (Table 2). The OGSI variable contributed least for both metrics.

We interpreted Maxent's univariate response curves, but have provided the

marginal plots as a supplemental (Fig. S4). Marten locations were correlated with both

low and high amounts of canopy cover and percent slope (quadratic response, Fig. 3).

We suspect these could be biologically correlated in that extensive flat areas in the

Central Coast also have low tree canopy cover, but high shrub density. Moderate

amounts of canopy cover (e.g., 5-50%) appeared to be negatively correlated with

marten locations. Predicted distribution was positively correlated with predicted salal

distribution with some likelihood of a threshold at high values (Fig. 3), percentage of

pine (Fig. 3), average annual precipitation (Fig. 3), and mast (Fig. 3). We observed a

negative correlation between marten locations and August temperature (Fig. 3). We

observed a slight negative relationship between marten locations and OGSI (Fig. 3).

Our final combined model predicted versus expected curve placed unsuitable

area <14%, suitable areas of 15-30%, and predicted highly suitable at >30% (Fig. 4)

with an AUC value on the test data at 92.1%. The model depicted southern Oregon and

northern California as having the largest extent for marten distribution, including areas

south of the current known population (Fig. 5).

Discussion

Our distribution model both predicts areas where Humboldt marten populations are

known to occur and identifies areas of potential occurrence outside of known population

extents. However, as with all models, there are limitations associated with our

predictions, and a clear assessment of these constraints is critical for model results to

be accurately used to inform management decisions (Sofaer et al. 2019). For example,

abiotic variables such as precipitation or slope are predictive of habitat use but may be

uninformative in a management framework because they cannot be altered.

Nonetheless, abiotic variables such as temperature and precipitation provide alternative

information that can be used for contemporary climatic baselines and future climate

scenarios. However, establishing patterns of causality is challenging, and it is also

plausible that these abiotic variables are linked to occurrence because remnant

Humboldt marten populations happen to occur in wet and steep regions. Further,

interpretation of model variables requires consideration of biological (e.g., biotic and

abiotic) processes within a spatial context, and to achieve this, we included predictors

that provided spatially-accurate depictions of habitat use at the most predictive spatial

scale (McGarigal et al. 2016). The culmination of these efforts resulted in a predicted

distribution map and hypotheses regarding resources correlated with Humboldt

martens.

The role of biotic interactions in shaping the distribution of species has been

reported (e.g., Forchhammer et al. 2005, Guisan and Thuiller 2005). However, evidence

of the importance of biotic variables alongside abiotic variables for predicting

distributions at larger spatial scales has been largely lacking (e.g., Wisz et al. 2013).

Here, we show that inclusion of certain abiotic variables in species distribution models

can improve model predictions markedly, providing support for the role of biotic

interactions in shaping species distributions at the landscape-scale (Wisz et al. 2013).

Occurrence of pine and canopy cover were two biotic variables with high model

contributions. We interpret the combination of the pine-dominated vegetation type with

the quadratic response of canopy cover as copacetic; that is areas with pine dominant

forest types typically have lower canopy cover and require adequate access to light for

growth, including areas with relatively low canopy cover (Augusto et al. 2003). In these

areas, shrub cover, particularly salal and mast-producing shrubs, may be more

prevalent. High shrub cover appears to be a consistent component of Humboldt marten

habitat in both California (Slauson & Zielinski 2009, Slauson et al. 2007) and Oregon

(Moriarty et al. 2019). Predicted marten distribution also was related to increased

canopy cover, a finding consistent with other research indicating that martens seek

areas with relatively high cover (Bissonette et al. 1997, Hargis et al. 1999). In the

univariate responses, canopy cover was quadratic, suggesting a relatively low

correspondence with moderate levels of canopy cover. More information is needed to

understand if there is an avoidance of thinned forests with moderate levels of canopy

cover similar to other Pacific marten literature (Moriarty et al. 2015; Moriarty et al. 2016).

Additional information quantifying amounts and extent of shrub and canopy cover are

needed to better understand how martens interact with vegetation at fine spatial scales,

the combination of low canopy cover with a dense understory layer of shrub and mast

producing species or a high level of canopy cover represent achievable conditions that

can guide management or target restoration.

Alternatively, biotic variables such as percent pine may not offer similar

guidance. Here, they may act as "proxy" variables to conditions where martens currently

persist. The influence of percent pine in our predicted Humboldt marten model is rooted

in a biological truth, as well-drained or serpentine soils prohibit extensive canopy

closure thus promoting extensive shrub growth. However, the presence of pine species

is most likely an artifact of co-occurrence with dense shrub cover. Our interpretation

concerns are especially salient for historically-occupied areas that no longer share

similar attributes, such as previously extensive shrub fields with succession into

homogenous but not yet structurally diverse forests.

Biotic variables influencing predicted Humboldt marten distribution in our model

were consistent with previous literature with some exceptions, most notably forest age

and old growth characteristics. Forest age was not a strong predictor in any model

iteration, nor was OGSI. We incorporated OGSI into our model, despite its meager

contributions to various model iterations (<5% contribution), because of its high

predictive contribution in the Slauson et al. (2019b) model. Within our model, the

predicted relationship between Humboldt marten distribution and higher OGSI values

was not only weak but negative – higher OGSI values could be interpreted as less

suitable for Humboldt marten use (Supplemental Item 1). We suspect the difference

between our model and the Slauson et al. (2019) model in regard to OGSI was due to

our survey efforts being greatly increased in geographic scope and located martens in a

greater diversity of vegetation types with widely differing ages. For example, the

Slauson et al. (2019) model included 8 marten locations in Oregon where our model

included 167 marten locations with survey efforts randomly or systematically distributed

(Moriarty et al. 2018; Moriarty et al. 2019). This resulted in a relatively even proportion

of locations throughout the distribution of the Humboldt marten, including telemetry

locations in younger forests previously identified as unsuitable (Zielinski et al. 2001).

The Slauson et al. (2019b) model had an advantage of using non-detection data and

our predictive response curves were similar in both models – both slopes were relatively

flat or decreasing with relatively broad confidence intervals. The most obvious

difference between models is geographic scope and extent, which is a challenge for

non-stationarity.

Although our model indicated that Humboldt marten locations are negatively

associated with OGSI, we are not suggesting an avoidance of old forest and

characteristics associated with resting, denning or foraging. Rather, OGSI appears to be

a poor predictor for Humboldt marten distribution when considering the entire range,

perhaps because of a potential mismatch in the association between OGSI and shrub

cover. For example, while mature Douglas fir-dominated forests inhabited by Humboldt

martens in California may have a substantial shrub component (Slauson et al. 2007),

much of coastal Oregon and California is dominated by areas of mature western

hemlock, a shade-tolerant species that is strongly associated with reduced shrub cover

(Kerns & Ohmann 2004). When examining the individual contributions of OGSI and

marten locations in a model only with the components of OGSI, downed wood was the

most influential variable (Supplemental Item S1). In other words, Humboldt marten

distribution appears to be more strongly associated with shrub cover and understory

conditions rather than moderately old forests with a density of larger trees or snags –

with the mismatch being that some areas where Humboldt martens occur (e.g., mature

Douglas fir forest; Slauson et al. 2007) are characterized by both older forest conditions

and substantial shrub cover, while other areas (e.g., serpentine or coastal pine forests;

Eriksson et al. 2019, Moriarty et al. 2019) are characterized by substantial shrub cover,

but not older forest conditions. Contemporarily, throughout the range, occurrence does

not appear to be correlated with higher OGSI values, which suggests a potential

overemphasis of the OGSI variable in the Slauson et al. (2019) model – particularly in

Oregon with few marten locations to base inference. Applications of the Slauson model

to Oregon was an extrapolation beyond the range of the data (Romesburg 1981;

Rosenbaum 2010).

Both models have challenges in interpretation from non-stationary effects. Non-

stationarity in this context describes species-habitat relationships that are variable

across time or space and can fundamentally alter interpretation of model output. Here,

we were unable to account for temporal non-stationarity (e.g., winter, summer) because

the location data seldom had locations surveyed in both seasons. We might expect

habitat selection to be more critical in the summer when martens are rearing kits and

mating, similar to findings in Martin et al. (In press), but we could not evaluate such

influence. It would be deeply flawed to assume that our model, and any other range-

wide model, could accurately address the local variation of predictive variables between

populations at fine spatial scales. Similarly, models are based on data collected to as of

summer 2020 and we suspect there are more martens east in the Oregon coast (e.g., in

the vicinity of Ashland, Applegate based on new locations) but our model would not

predict such areas due to sparse location information. Future endeavors could develop

site-specific models, ideally using telemetry data that are biologically linked with fitness

(e.g., long-lived adult female rest and den structures) to address predicted habitat.

Understanding habitat was not our goal with this exercise; nonetheless, our modeling

exercise provides opportunities to test the influence of these predictive covariates with

independent data.

Abiotic factors such as increased precipitation, proximity to the coast, and cool

temperatures could be an indication of vegetation growth or may be indicative of a

physiological temperature threshold. Precipitation, an unalterable condition but

potentially a factor that may be altered in a changing climate, was one of the top 3

predictive variables, fulfilling a high proportion of the model contribution (>20%) in all

model simulations. Range limit theorems have long postulated the importance of

elevation, altitude, and weather establishing the upper limits of species distributions

(e.g., Darwin 1859). If these variables are causally linked to marten occurrence, a

plausible mechanism is that cooler wetter conditions are related to increased dense

vegetation, which probably aids martens in avoiding predators. Predation was

documented as the single greatest limitation to a small and endangered fisher

population (Sweitzer et al. 2015) and is a plausible hypothesis for Humboldt marten

distribution limitation. Coupled with additional food from berries and small animals that

eat berries (e.g., birds, rodents), these areas may create exceptional, yet typically

uncharacteristic, habitat (Eriksson et al. 2019). A second synergistic mechanism aligns

with Humboldt marten persistence in areas with relatively stable climates (e.g., a

backwards velocity evaluation of past change) and areas that may remain relatively

stable in future climate predictions (Belote et al. 2018; Carroll et al. 2015). Evidence for

this hypothesis would align with observations of martens associated with cooler

temperatures associated with higher precipitation within their range limits. The global

distributions of "boreal" martens (*Martes caurina, M. americana, M. martes, M. zibellina,*

M. melampus) are north of 45 degrees latitude, and these species reside in areas with

deep snow cover for a large portion of the year (Krohn et al. 2004; Suffice et al. 2020).

The thermoneutral zone of martens has yet to be explored. Generally, species range

limits appear highly influenced by abiotic factors (e.g., Normand et al. 2009), although

there have been several situations where biotic factors such as vegetation, competition,

or predation appear to influence distributions (Louthan et al. 2015; Sirén & Morelli

2020). Here, we would predict that the stable cool and wet climate in near-coastal

forests may provide Humboldt martens with a less energetically-expensive temperature

gradient.

This modeling effort can be integrated into the framework proposed in the

Humboldt marten conservation assessment and strategy, which described a 3-prong

approach for Humboldt marten conservation: (1) protect existing populations; (2) re-

establish populations in areas with habitat; and (3) restore or focus management efforts

to improve habitat (Slauson et al. 2019a). A fundamental foundation of protecting

existing populations includes identifying population locations and geographic extents,

describing population sizes, and monitoring changes in populations over time. Our

model predicts habitat outside of the currently understood boundaries of Humboldt

marten populations; prioritizing survey effort in areas that are predicted to be habitat but

are of unknown marten occupancy would be an important first step in protecting existing

populations. A stratified-random approach, expending survey effort in both predicted

suitable and unsuitable areas, would best evaluate marten occurrence. Incorporating

non-invasive survey methods (e.g., remote cameras), coupled with genetic analyses,

further presents an opportunity to effectively survey species that are cryptic and occur at

low densities over large spatial and temporal scales, and to robustly answer questions

concerning animal distribution, abundance and demographics (Pauli et al. 2010, Lamb

et al. 2019).

Consideration of areas where Humboldt martens do not appear to occur in

contemporary times, but our model predicts as suitable habitat, is exemplary of

interpretation that requires careful evaluation of biological processes. For example, our

model predicts marten distribution in the northern and southern extremes of our

modeling region, including the Tillamook State Forest in northern Oregon coast and the

King Range National Conservation Area in California, respectively. While the Tillamook

State Forest had higher amounts of the abiotic variables of our model (e.g., high

precipitation amounts, cool temperatures, and coastal proximity), there were fewer

areas predicted with forest vegetation conditions there (Supplemental File 1).

Alternatively, the suitability of the unoccupied southern modeling extent (King Range

National Conservation Area) was primarily driven by biotic factors (e.g., high canopy

cover, high amount of masting trees and shrubs; Supplemental Fig. x). Given the

importance of these biotic factors in our model, and that the southern portion represents

a significant portion of the historical range of the Humboldt marten (Zielinski et al. 2001),

it may provide marten habitat that is currently unoccupied. While suitability as proxy for

population viability in a given area has been questioned broadly (Stephens et al. 2015),

potential but unoccupied habitat should be areas considered for future survey efforts.

While our model predicts habitat in apparently unoccupied areas that could

eventually be considered for population re-establishment and restoration, it also

underscored our lack of reliable knowledge on critical and consistent components of

Humboldt marten habitat, and we suggest it is premature to pursue actions such as

population re-establishment without first refining our understanding of marten habitat

requirements. This effort and prior explorations of vegetation associations exemplify the

vast range of vegetation conditions used by Humboldt martens (Eriksson et al. 2019;

Moriarty et al. 2019; Slauson et al. 2007). Yet while some vegetation conditions are

certainly required for species' population persistence, vegetation conditions may not act

as a categorical surrogate for habitat.

Although species' distributions can be influenced by very apparent biotic (e.g.,

vegetative) and abiotic (e.g., temperature thresholds) conditions, they may also be

strongly influenced by less-apparent factors such as interspecific interactions with

predator or competitor species (Siren 2020). As an example, spotted owls (*Strix*

occidentalis) closely align with old-growth forest conditions which have been

characterized with relatively high accuracy (Davis et al. 2016), yet spotted owl

population viability is dramatically decreased with presence of barred owls (*S. varia*)

due to competition, predation, and predation risk (Diller et al. 2016; Dugger et al. 2016;

Wiens et al. 2014). Although few examples exist for carnivores, a recent evaluation

suggests that while lynx (*Lynx lynx*) distributions are closely-tied to deep snow, the

influence of reducing bobcat (*L. rufus*) competition was stronger than the influence of

snow itself (Siren 2020). A directed research effort would be necessary to understand

the relative importance of vegetation structures, vegetation types, prey, predation, and

competition for Humboldt marten persistence.

We developed a range-wide species distribution model based on extensive

survey coverage and modern vegetation and climatic variables. Although these results

provide predictions for habitat components, describing habitat would be best informed

by measures of survival and fecundity rates. At present, managers have conflicting

sources of evidence that can lead to confusion in interpretating of Humboldt marten

ecology. In this paper, we provided models that are complementary, but not similar to

other models (Schrott & Shinn 2020; Slauson et al. 2019b), particularly in regard to

interpretation of OGSI. Instead of interpreting differences between models as a conflict,

we posit this as evidence of the conservation challenge described by Caughley (1994)

and in our bison example. We lack enough information regarding where Humboldt

martens resided historically to compare with our contemporary distribution (Loehle

2020), and we are generally ignorant of population densities, causal associations of

population declines, and population limitations. Such an understanding is essential to

describe expectations of future range (Brown et al. 1996). Each available case study

provides an imperfect piece of evidence, and the lack of consistency among studies is

suggestive of imperfect knowledge of what components constitute Humboldt marten

habitat. To avoid "wicked problems" from differing views for rare species conservation

(e.g., Gutiérrez 2020; Jones et al. 2020), amassing information collaboratively with a

goal of prospective meta-analyses and study-level replication will be essential (Facka &

Moriarty 2017; Nichols et al. 2019).

Conclusions

Based on our modeling and an evaluation of available evidence, we conclude that the

most consistent range-wide characteristic with Humboldt marten distributions are

vegetation associations with extensive dense shrub cover or complex understory

vegetation, which may ultimately be an association with increased food availability or

predation escape cover. An understanding of the strength of these interactions and

factors that limit populations will ultimately be needed for making wise conservation

decisions. Moving forward with conservation decisions with incomplete knowledge

necessitates an adaptive management framework.

Acknowledgements

The desire and decision to request an updated model incorporating newer presence

and habitat information was from the Oregon Humboldt Marten stakeholder group,

which is facilitated by the U.S. Fish and Wildlife Service in Oregon. Non-invasive marten

surveys were conducted by Pacific Northwest and Southwest Research Stations,

Oregon State University, Humboldt State University, Green Diamond Resource

Company, NCASI, the Siuslaw, Rogue-Siskiyou, and Six Rivers National Forests,

Hancock Forest Management, Weyerhaeuser, Oregon Department of Forestry, and the

Confederated Tribes of Siletz Indians of Oregon. Detection dog surveys were completed

by Rogue Detection Dog Teams and the former group within Conservation Canines,

University of Washington. Considerable aid with field logistics, vehicles, housing, and

equipment was provided by the U.S. Fish and Wildlife Service, Salem District BLM,

USFS Rogue River-Siskiyou and Siuslaw National Forests, Weyerhaeuser, Hancock

Forest Management, and USFS Region 6 Regional Office. We obtained private land

access or surveys were completed by trained staff within the ownership for all randomly

selected survey points – thanks to Weyerhaeuser, Hancock Forest Management,

Starker Forests, and Roseburg Timber for access or data. Reviews by Drs. D. Miller and

J. Verschuyl, B. Hollen, our anonymous peer reviewers, and the Associate Editor

improved previous versions of this manuscript. Extreme thanks to all field crew leaders

(S. Smythe, M. Linnell, B. Peterson, G. W. Watts, J. Bakke, C. Shafer, K. Kooi, and M.

Penk) and team members (E. Anderson, D. Baumsteiger, A. Benn, J. Buskirk, B.

Carniello, M. Cokeley, S. Hart, P. Iacano, A. Kornak, T. McFadden, E. Morrison, A.

Palmer, T. Peltier, N. Palazzotto, S. Roon, S. Riutzel, C. Scott, K. Smith, R. Smith, T.

Stinson, M. Williams, B. Woodruff, and K. Wright). Any use of trade, firm, or product

names is for descriptive purposes only and does not imply endorsement by the U.S.

Government.

Funding

Marten survey efforts were funded by the Oregon State University Fish and Wildlife

Habitat in Managed Forests Research Program, NCASI, Coos Bay-Bureau of Land

Management (BLM), the USDA Forest Service (USFS) Siuslaw and Rogue-Siskiyou

National Forests and Six Rivers National Forest, the Oregon Forestry Industry Council,

U.S. Fish and Wildlife Service (Arcata office), and Humboldt State University Sponsored

Programs Foundation. NCASI, USFS Pacific Northwest and Southwest Research

Stations funded genetic confirmation of scats, remote camera data processing and

management, and provided external support. Telemetry data were funded by the

Siuslaw National Forest, Green Diamond Resource Company, and U.S. Fish and

Wildlife Service.

Figure 1. Our study area and modelling region for Humboldt martens (*Martes caurina humboldtensis*) included all of coastal Oregon and northern California.

We modeled Humboldt marten predicted distributions in forested lands (Panel A, green mask) in 2 ecoregions [left]. We created a minimum convex polygon of known locations buffered by 10-km (hatched area). We compiled 1,692 known marten locations (icon

762   color) from 5,143 surveyed sites with non-detections in light gray, collected during 1996-

763   2020 (panel B). We spatially thinned locations to approximately 500m apart, prioritizing

764   den and rest locations and resulting in 384 locations (black dots, panel C).

Figure 2. We investigate the range of variables in our thinned dataset compared to all marten locations and detection/non-detection data.

To provide the range of values observed in this study, we depict boxplots for the variables in the top model showing the thinned marten data (Marten), all marten known

771   locations (Marten_DB), non-detected but surveyed locations (Non-detection), and

772   random locations within the minimum convex polygon (9,600 random locations).

Figure 3. We depict predicted relationships between Humboldt marten locations and each of the variables within our final model.

Here, each curve is the predicted probability of presence with no conflicting influence of potentially correlated variables. Humboldt marten locations were correlated with both low and high amounts of canopy cover and percent slope (quadratic response).

Predicted distribution was positively correlated with predicted salal (*Gaultheria shallon*)

distribution, percentage of pine, precipitation, and mast. We observed a negative

correlation between marten locations and August temperature. We observed a slight

negative relationship between marten locations and the old growth structural index.

Percent contribution and permutation importance values were reported in Table 2. The

curves reveal the mean response (black) and standard deviation (gray) for 10 replicate

Maxent runs.

Figure 4. Our predicted suitable transitions for Humboldt marten (*Martes caurina humboldtensis*) range.

We present mean predicted vs. expected curve (solid black line) from our model replicates, showing 95-percent confidence intervals (gray-shaded vertical bars). The P/E = 1 threshold is where the curve crosses the random chance line (horizontal orange line), and the blue dashed vertical lines are the 95-percent confidence intervals. We used the predicted-to-expected curve to inform our suitability thresholds following Hirzel et al. (2006), including predicted unsuitable (P/E and confidence intervals 0-1), marginal (P/E > 1 but overlapping confidence intervals), and suitable (P/E and confidence intervals > 1; map depicted in Fig. 5).

Figure 5. We display our modeled predicted range for Humboldt marten (*Martes caurina humboldtensis*).

For predicted range, we followed Hirzel et al. (2006) with predicted versus expected

ratios transitioning between predicted highly suitable (green), suitable (orange), and

marginal or not predicted suitable (gray). Marten location information was displayed

(black dots). We zoomed to population extents to provide increased visual resolution

within the Central Oregon Coast (Panel 3a), South coast (Panel 3b), and northern

California (Panel 3c).

Table 1. Data ranges, means, and standard deviations for the model region, the contemporary Humboldt marten
distribution, and at Humboldt marten locations.
We depict individual layer statistics within our Humboldt marten (*Martes caurina humboldtensis*) model region in coastal

Oregon and northern California. We display the variable, optimized spatial scale with a radius in meters, value range from

the coastal ecoregions, means and standard deviation (SD) for the model region, minimum convex polygon around all

known marten locations (MCP), and values from spatially thinned marten locations (n = 384), our layer source, and a

description of that variable. We only considered variables with < 60% correlation in our final model (Table S2).

Variable	Scale	Value Range	Model Region (Mean ± SD)	Minimum convex polygon (Mean ± SD)	Marten Locations (Mean ± SD)	Source	Description
Forest age, years	270	0 – 712	95.5 ± 43	104.3 ± 49.4	109.8 ± 69.6	2016 GNN	Basal area weighted stand age based on field recorded or modeled ages of dominant/codominant trees
Canopy cover (%)	1170	2 – 99	65.9 ± 13	66.4 ± 14	71.3 ± 18.6	2016 GNN	Canopy cover percentage of all live trees
Coastal proximity	50	2 – 700	511.7 ± 193.1	516.3 ± 203.1	361.8 ± 197.9	PRISM	Optimal path length from the coastline accounting for terrain blockage (Daly et al. 2008)
Diameter diversity index	1170	26 – 811	433.9 ± 103	437.6 ± 111.7	459.4 ± 123.6	2016 GNN	Diameter diversity index - measure of stand structure based on tree densities in diff. DBH classes (x100)

Percent downed wood	270	0 – 797	69.3 ± 54.7	70.9 ± 50	68.5 ± 60.1	2016 GNN (created)	Created within GNN to estimated percentage of large downed wood, a component of OGSI
Salal	1170	0 – 100	35.7 ± 30.9	50.7 ± 32.3	72.7 ± 17.8	Prevéy	Probability of *Gautheria shallon* species occurrence (Prevéy et al. 2020)
Masting vegetation	1170	0 – 72	5.9 ± 7.4	5.2 ± 6.7	9.3 ± 9	2016 GNN	Percent of stand basal comprised of tanoak (*Notholithocarpus densiflorus*; LIDE), giant chinquapin (*Castanopsis chrysophylla*; CHCH), coastal live oak (*Quercus agrifolia*; QUAG), canyon live oak (*Quercus chrysolepis*; QUCH), and California bay (*Umbellularia californica*; UMCA) (mast producing evergreen hardwoods, indicator of prey abundance)
Old growth structural index	50	0 – 100	32.7 ± 15.8	33.2 ± 16.1	33.8 ± 16.9	2016 GNN	Old-growth structure index based on abundance of large live trees, snags, down wood, and ddi
Percent pine	1170	0 – 94	1.2 ± 3.5	1.5 ± 4.5	10.9 ± 20.1	2016 GNN	Percent of pixel basal area comprised of shore pine (*Pinus contorta*; PICO), Jefferey pine (*Pinus jeffreyi*; PIJE) and knobcone pine (*Pinus attenuata*; PIAT). We use this as an indicator of serpentine and coastal dune environments.
Percent slope	1170	0 – 74	33.8 ± 10.9	36.2 ± 10.6	31.7 ± 15.8	USGS DEM	Percent slope in degrees

Precipitation	1170	13 – 198	66.9 ± 27	70 ± 30.1	102.4 ± 30.5	2016 GNN	Average annual precipitation 1981-2010 (inches)
Large snag density	742	0 – 48	4.9 ± 4.3	5.8 ± 4.6	6.9 ± 4.9	2016 GNN (created)	Created within GNN to estimated density of large snags, a component of OGSI
Temperature (August max)	1170	8 – 24	16.5 ± 2.3	16.1 ± 1.7	16.4 ± 1.7	PRISM	Average annual maximum temperature 1981-2010 (Celcius).
Topographic position index	270	-149 – 174	0.7 ± 26.7	1.1 ± 28.8	-0.3 ± 28.6	USGS DEM	Topographic position index - difference of cell elevation with mean of all cells w/in 450 m radius
Large tree density	1170	0 – 47	3.2 ± 3.5	4.4 ± 4.2	5.2 ± 5.9	2016 GNN (created)	Created within GNN to estimated density of large trees, a component of OGSI
Huckleberry	1170	2 – 99	32.7 ± 24.6	39.1 ± 26	42.7 ± 27.2	Prevéy	Probability of species occurrence for *Vaccinium ovatum* (created)

820 **Table 2. We show the percent contribution and permutation importance from our top Maxent model.**

821 We ordered variables by their percent contribution and report the optimized spatial scale (focal radius in meters), the

822 univariate response type, and whether the univariate dependent plots were generally positively or negatively correlated

823 with Humboldt marten (*Martes caurina humboldtensis*) locations.

Variable	Scale	Response	Univariate Relationship	Percent contribution	Permutation importance
Salal	1170	Quadratic	+	23.3	15.5
Percent pine	1170	Product	+	22.5	30.3
Precipitation_30 year average	1170	Product	+	21.6	25.3
Canopy cover	1170	Quadratic	+	18.7	20.2
Mast	1170	Product	+	5.4	1.3
August temperature_30 year average	1170	Linear	-	4.7	2.3
Percent slope	1170	Quadratic	-	2.7	4.4
Old growth structural index	50	Linear	-	1.2	0.7

Supplemental Information Captions

Supplemental Data S1. We provide marten and random locations with associated modeled values at multiple scales.

We provide raw data from thinned Humboldt marten (*Martes caurina humboldtensis*) and random locations within the known area (minimum convex polygon). We extracted data at 4 spatial scales (radius = 50m, 270m, 742m 1170m) associated with Humboldt marten biology (e.g., Table S1) and ran univariate general linear models to select the most predictive spatial scale.

Supplemental Data S2. We provide marten locations from two radio telemetry studies for the purpose of estimating home range size.

We provide raw data from Humboldt marten (*Martes caurina humboldtensis*) locations collected within the Northern California (VHF telemetry) or Oregon Central Coast (GPS telemetry) regions with methods described elsewhere (Delheimer et al. In press; Linnell et al. 2018). We used only data from individuals with greater than 30 locations and with location error estimated at less than 400 sq m to estimate range size (Table S1). We used range size to inform spatial scale smoothing extents for modeled variables. Because Humboldt martens are state endangered in California and federally threatened, we were not comfortable revealing location data (Lindenmayer & Scheele 2017). For each marten, we selected either added (heads) or subtracted (tails) the individual marten's location by a random value (range = 50,000 – 100,000 m) such that all location data for a marten were similar spatially for the purpose of estimating range size but not relatable to the true location where the marten was monitored.

Supplemental Item S1. Exploration of the Old Growth Structural Index and

Humboldt martens (*Martes caurina humboldtensis*)

Because the old growth structural index has been interpreted as a variable indicative of

Humboldt marten predicted locations in other models, but not within our exploration, we

explore the variable in detail and explain nuances to why there might be differences.

Supplemental Figure S1, S2. We display boxplots for all considered biotic

variables with predicted associations with Humboldt martens.

To provide the range of values observed in this study, we depict boxplots for the biotic

variables in the top model showing the thinned marten data and random locations

(25/marten location; 9,600 random locations) for each spatial scale (radius = 50m,

270m, 742m 1170m) associated with Humboldt marten biology (e.g., Table S1,

described in methods).

Supplemental Figure S3. We display boxplots for all considered abiotic variables

with predicted associations with Humboldt martens.

To provide the range of values observed in this study, we depict boxplots for the abiotic

variables in the top model showing the thinned marten data and random locations

(25/marten location; 9,600 random locations) for each spatial scale (radius = 50m,

270m, 742m 1170m) associated with Humboldt marten biology (e.g., Table S1,

described in methods).

Supplemental Figure S4. We depict predicted relationships (marginal plots) between Humboldt marten locations and each of the variables within our final model.

Here, each curve is the predicted probability of presence given the other variable responses. Predicted distribution of Humboldt martens were correlated with increasing canopy cover, percent pine, precipitation, temperature, mast, percent salal (*Gaultheria shallon*) distribution, and the old growth structural index. We observed a negative correlation between marten locations percent slope. Percent contribution and permutation importance values were reported in Table 2. The curves reveal the mean response (black) and one standard deviation (gray) for 10 replicate Maxent runs.

Supplemental Table S1. Humboldt marten home range sizes by sex and region.

We compiled home range size information for Humboldt martens in the Central Coast of Oregon and Northern California to base our scale optimization for variables in this distribution model. Data from northern California were extracted from unpublished data (PSW 2019) and data from the Central Oregon Coast were in the supplemental material within Linnell et al. (2018).

Supplemental Table S2. We evaluated Pearson correlation coefficients for inclusion in our distribution model.

We evaluated Pearson correlation coefficients and restricted variables with correlation >0.6 to guide variables in a Humboldt marten (*Martes caurina humboldtensis*) distribution model. For our final model, we selected canopy cover over the diameter

diversity index. We selected OGSI over tree age or diameter diversity index due to its

use in prior Humboldt marten range wide models. We selected salal over huckleberry

because of salal's presumed structural use (Eriksson et al. 2019, Moriarty et al. 2019).

Supplemental Table S3. We evaluated Variance Inflation Factors, which reveal

correlation and multicollinearity, to guide variables in a Humboldt marten (*Martes*

***caurina humboldtensis*) distribution model.**

The absolute value of variance inflation factors equal to 1 are considered not correlated

and values greater than 5 are highly correlated as determined by Velleman & Welsch

(1981). Here, diameter diversity index conflicted with canopy cover and we used

canopy cover in our final model for the ease of interpretation and use in a management

context.

Literature cited

Andruskiw M, Fryxell JM, Thompson ID, and Baker JA. 2008. Habitat-mediated variation in predation risk in the American marten. *Ecology* 89:2273-2280.

Augusto L, Dupouey J-L, and Ranger J. 2003. Effects of tree species on understory vegetation and environmental conditions in temperate forests. *Annals of Forest Science* 60:823-831.

Balharry D. 1993. Social organization in martens: an inflexible system. *Symposia of the Zoological Society of London* 65:321-345.

Barry BR. 2018. Distribution, habitat associations, and conservation status of Pacific fisher (*Pekania pennanti*) in Oregon Thesis. Oregon State University.

Barry BR, Moriarty KM, Green DS, Hutchinson RA, and Levi T. In review. Integrating detection dogs, camera traps, and recovery trajectories into species distribution models.

Bell DM, Acker SA, Gregory MJ, Davis RJ, and Garcia BA. 2021. Quantifying regional trends in large live tree and snag availability in support of forest management. *Forest Ecology and Management* 479:118554. https://doi.org/10.1016/j.foreco.2020.118554

Bell DM, Gregory MJ, and Davis R. 2020. Gradient nearest neighbor map data quality summary: GNN-2020. In: Service UF, editor: USDA Forest Service and Oregon State University.

Belote RT, Carroll C, Martinuzzi S, Michalak J, Williams JW, Williamson MA, and Aplet GH. 2018. Assessing agreement among alternative climate change projections to

inform conservation recommendations in the contiguous United States. *Scientific Reports* 8:1-13.

Bissonette JA, and Broekhuizen S. 1995. Martes populations as indicators of habitat spatial patterns: the need for a multiscale approach. In: Lidicker WZ, ed. *Landscape Approaches in Mammalian Ecology and Conservation*. Minneapolis, Minnesota, USA: University of Minnesota Press, 95-121.

Boyce MS, Vernier PR, Nielsen SE, and Schmiegelow FKA. 2002. Evaluating resource selection functions. *Ecological Modelling* 157:281-300. http://dx.doi.org/10.1016/S0304-3800(02)00200-4

Brown JH, Stevens GC, and Kaufman DM. 1996. The geographic range: size, shape, boundaries, and internal structure. *Annual Review of Ecology and Systematics* 27:597-623.

Carroll C, Lawler JJ, Roberts DR, and Hamann A. 2015. Biotic and climatic velocity identify contrasting areas of vulnerability to climate change. *PLoS ONE* 10:e0140486.

Caughley G. 1994. Directions in conservation biology. *Journal of Animal Ecology* 63:215-244.

CDFW. 2019. Title 14: OAL Matter Number: 2019-0201-02 Endangered Status for Humboldt marten, tri-colored blackbird, fisher southern Sierra ESU. California Department of Fish and Wildlife; 18 Mar 2019 Amendment.

Cheveau M, Imbeau L, Drapeau P, and Belanger L. 2013. Marten space use and habitat selection in managed coniferous boreal forests of eastern Canada. *The Journal of Wildlife Management*. 10.1002/jwmg.511

Clemesha RE, Gershunov A, Iacobellis SF, Williams AP, and Cayan DR. 2016. The

northward march of summer low cloudiness along the California coast.

Geophysical Research Letters 43:1287-1295.

Crookston N, and Stage A. 1999. Percent canopy cover and stand structure statistics

from the forest vegetation simulator. USDA Forest Service, Rocky Mountain

Research Station. Gen. Tech. Rep. RMRS-GTR-24.

Daly C, Halbleib M, Smith JI, Gibson WP, Doggett MK, Taylor GH, Curtis J, and

Pasteris PP. 2008. Physiographically sensitive mapping of climatological

temperature and precipitation across the conterminous United States.

International Journal of Climatology: A Journal of the Royal Meteorological

Society 28:2031-2064.

Daly C, Helmer EH, and Quiñones M. 2003. Mapping the climate of Puerto Rico,

Vieques and Culebra. *International Journal of Climatology: A Journal of the Royal*

Meteorological Society 23:1359-1381.

Darwin C. 1859. *The origin of species. 6th edition*: John Murray, London.

Davis R, Ohmann J, Kennedy R, Cohen W, Gregory M, Yang Z, Roberts H, Gray A, and

Spies T. 2015. Northwest Forest Plan – the first 20 years (1994-2013): status

and trends of late-successional and old-growth forests USDA Forest Service:

Portland, OR, USA.

Davis RJ, Hollen B, Hobson J, Gower JE, and Keenum D. 2016. Northwest Forest

Plan—the first 20 years (1994–2013): status and trends of northern spotted owl

habitats. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific

Northwest Research Station. General Technical Report PNW-GTR-929. p 54.

Delheimer MS, Moriarty KM, Slauson KM, Roddy AM, Early DA, and Hamm KA. In

press. Comparative reproductive ecology of two subspecies of Pacific marten

(*Martes caurina*) in California. *Northwest Science.*

Diller LV, Hamm KA, Early DA, Lamphear DW, Dugger KM, Yackulic CB, Schwarz CJ,

Carlson PC, and McDonald TL. 2016. Demographic response of northern spotted

owls to barred owl removal. *The Journal of Wildlife Management* 80:691-707.

Dugger KM, Forsman ED, Franklin AB, Davis RJ, White GC, Schwarz CJ, Burnham KP,

Nichols JD, Hines JE, and Yackulic CB. 2016. The effects of habitat, climate, and

Barred Owls on long-term demography of Northern Spotted Owls. *The Condor*

118:57-116.

Dye AW, Rastogi B, Clemesha RE, Kim JB, Samelson RM, Still CJ, and Williams AP.

2020. Spatial patterns and trends of summertime low cloudiness for the Pacific

Northwest, 1996–2017. *Geophysical Research Letters* 47:e2020GL088121.

Elith J, Graham CH, Anderson RP, Dudík M, Ferrier S, Guisan A, Hijmans RJ,

Huettmann F, Leathwick JR, Lehmann A, Li J, Lohmann LG, Loiselle BA, Manion

G, Moritz C, Nakamura M, Nakazawa Y, Overton JM, Peterson AT, Phillips SJ,

Richardson K, Scachetti-Pereira R, Schapire RE, Sobero´n J, Williams S, Wisz

MS, and Zimmermann NE. 2006. Novel methods improve prediction of species'

distributions from occurrence data. *Ecography* 29:129-151.

Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE, and Yates CJ. 2011. A statistical

explanation of MaxEnt for ecologists. *Diversity and Distributions* 17:43-57.

Eriksson CE, Moriarty KM, Linnell MA, and Levi T. 2019. Biotic factors influencing the

unexpected distribution of a Humboldt marten (Martes caurina humboldtensis)

population in a young coastal forest. *PLoS ONE* 14:e0214653.

10.1371/journal.pone.0214653

Facka AN, and Moriarty KM. 2017. An approach to foster a new generation of broad-

scale collaboration within the *Martes* Working Group. In: Zalewski A, Aubry KB,

O'Mahony D, Birks JDS, and Proulx G, eds. *The Martes complex in a new*

millennium: Mammal Research Institute; Polish Academy of Sciences.

Fielding AH, and Bell JF. 1997. A review of methods for the assessment of prediction

errors in conservation presence/absence models. *Environmental Conservation*

24:38-49.

Franklin JF, and Dyrness CT. 1973. Natural vegetation of Oregon and Washington.

USDA Forest Service General Technical Report, Pacific Northwest Forest and

Range Experiment Station.

Gamblin HE. 2019. Distribution and habitat use of a recently discovered population of

Humboldt martens in California Master of Science. Humboldt State University.

Gedir JV, Cain III JW, Swetnam TL, Krausman PR, and Morgart JR. 2020. Extreme

drought and adaptive resource selection by a desert mammal. *Ecosphere*

11:e03175.

Gutiérrez R. 2020. Invited commentary: when a conservation conflict comes full circle —

the spotted owl conflict is a wicked problem. *Journal of Raptor Research* 54:337-

348.

Halvorsen R. 2013. A strict maximum likelihood explanation of MaxEnt, and some

implications for distribution modelling. *Sommerfeltia* 36:1-132.

Hargis CD, Bissonette JA, and Turner DL. 1999. The influence of forest fragmentation

and landscape pattern on American martens. *Journal of Applied Ecology* 36:157-

172.

Himjmans RJ. 2020. raster: Geographic data analysis and modeling. In: 3.3-13 Rpv,

editor. https://CRAN.R-project.org/package=raster.

Hirzel AH, Le Lay G, Helfer V, Randin C, and Guisan A. 2006. Evaluating the ability of

habitat suitability models to predict species presences. *Ecological Modelling*

199:142-152.

Hoglander C, Dickson BG, Rosenstock SS, and Anderson JJ. 2015. Landscape models

of space use by desert bighorn sheep in the Sonoran Desert of southwestern

Arizona. *The Journal of Wildlife Management* 79:77-91.

Jenness J. 2006. Topographic Position Index (tpi_jen. avx) extension for ArcView 3. x,

v. 1.3 a. Jenness Enterprises (accessed May 2019.

Jensen PG, Demers CL, Mcnulty SA, Jakubas WJ, and Humphries MM. 2012. Marten

and fisher responses to fluctuations in prey populations and mast crops in the

northern hardwood forest. *Journal of Wildlife Management* 76:489-502.

Johnson DH. 1980. The comparison of usage and availability measurements for

evaluating resource preference. *Ecology* 61:65-71. doi:10.2307/1937156

Jones GM, Gutiérrez R, Block WM, Carlson PC, Comfort EJ, Cushman SA, Davis RJ,

Eyes SA, Franklin AB, and Ganey JL. 2020. Spotted owls and forest fire:

Comment. *Ecosphere* 11.

Joyce MJ. 2013. Space use behavior and multi-scale habitat selection of American

marten (*Martes americana*) in northeastern MinnesotaMaster of Science Thesis.

University of Minnisota.

Kerns BK, and Ohmann JL. 2004. Evaluation and prediction of shrub cover in coastal

Oregon forests (USA). *Ecological Indicators* 4:83-98.

Kramer-Schadt S, Niedballa J, Pilgrim JD, Schröder B, Lindenborn J, Reinfelder V,

Stillfried M, Heckmann I, Scharf AK, Augeri DM, Cheyne SM, Hearn AJ, Ross J,

Macdonald DW, Mathai J, Eaton J, Marshall AJ, Semiadi G, Rustam R, Bernard

H, Alfred R, Samejima H, Duckworth JW, Breitenmoser-Wuersten C, Belant JL,

Hofer H, and Wilting A. 2013. The importance of correcting for sampling bias in

MaxEnt species distribution models. *Diversity and Distributions* 19:1366-1379.

10.1111/ddi.12096

Krohn W, Hoving C, Harrison DJ, Phillips DM, and Frost H. 2004. *Martes* foot-loading

and snowfall patterns in eastern North America: implications to broad-scale

distributions and interactions of mesocarnivores. In: Harrison DJ, Fuller AK, and

Proulx G, eds. *Martens and fishers (Martes) in human-altered environments: an*

international perspective. New York, New York, USA: Cornell University Press,

115-134.

Lindenmayer D, and Scheele B. 2017. Do not publish. *Science* 356:800-801.

Linnell MA, Moriarty K, Green DS, and Levi T. 2018. Density and population viability of

coastal marten: a rare and geographically isolated small carnivore. *PeerJ*

6:e4530 - '4521 pg. 10.7717/peerj.4530

Loehle C. 2020. Quantifying species' geographic range changes: conceptual and

statistical issues. *Ecosphere* 11.

Louthan AM, Doak DF, and Angert AL. 2015. Where and when do species interactions

set range limits? *Trends in Ecology & Evolution* 30:780-792.

Manlick PJ, Petersen SM, Moriarty KM, and Pauli JN. 2019. Stable isotopes reveal

limited Eltonian niche conservatism across carnivore populations. *Functional*

Ecology https://doi.org/10.1111/1365-2435.13266.

Martin ME, Moriarty KM, and Pauli JN. In press. Scale and season influence the

resource selection of a snow-adapted forest carnivore, the Pacific marten.

Landscape Ecology.

McGarigal K, Wan HY, Zeller KA, Timm BC, and Cushman SA. 2016. Multi-scale habitat

selection modeling: a review and outlook. *Landscape Ecology*:1-15.

10.1007/s10980-016-0374-x

Merow C, Smith MJ, and Silander JA. 2013. A practical guide to MaxEnt for modeling

species' distributions: what it does, and why inputs and settings matter.

Ecography 36:1058-1069. 10.1111/j.1600-0587.2013.07872.x

Minta SC, Kareiva PM, and Curlee AP. 1999. Carnivore research and conservation:

learning from history and theory. In: Clark TW, Curlee AP, Minta SC, and Kareiva

PM, eds. *Carnivores in ecosystems: the Yellowstone experience*. London, United

Kingdom: Yale University Press, 323-404.

Morales NS, Fernández IC, and Baca-González V. 2017. MaxEnt's parameter

configuration and small samples: are we paying attention to recommendations?

A systematic review. *PeerJ* 5:e3093. 10.7717/peerj.3093

Moriarty KM, Epps CW, Betts MG, Hance DJ, Bailey JD, and Zielinski WJ. 2015.

Experimental evidence that simplified forest structure interacts with snow cover

to influence functional connectivity for Pacific martens. *Landscape Ecology*

30:1865-1877. 10.1007/s10980-015-0216-2

Moriarty KM, Epps CW, and Zielinski WJ. 2016. Forest thinning for fuel reduction

changes movement patterns and habitat use by Pacific marten. *The Journal of*

Wildlife Management 80:621-633. 10.1002/jwmg.1060

Moriarty KM, Linnell MA, Chasco B, Epps CW, and Zielinski WJ. 2017. Using high-

resolution short-term location data to describe territoriality in Pacific martens.

Journal of Mammalogy 98:679-689.

Moriarty KM, Linnell MA, Thornton JE, and Watts III GW. 2018. Seeking efficiency with

carnivore survey methods: a case study with elusive martens. *Wildlife Society*

Bulletin 42:403-413.

Moriarty KM, Verschuyl J, Kroll AJ, Davis R, Chapman J, and Hollen B. 2019.

Describing vegetation characteristics used by two rare forest-dwelling species:

Will established reserves provide for coastal marten in Oregon? *PLoS ONE*

14:e0210865. 10.1371/journal.pone.0210865

Nichols JD, Kendall WL, and Boomer GS. 2019. Accumulating evidence in ecology:

Once is not enough. *Ecology and Evolution* 9:13991-14004.

Normand S, Treier UA, Randin C, Vittoz P, Guisan A, and Svenning JC. 2009.

Importance of abiotic stress as a range-limit determinant for European plants:

insights from species responses to climatic gradients. *Global Ecology and*

Biogeography 18:437-449.

Ohmann JL, and Gregory MJ. 2002. Predictive mapping of forest composition and

structure with direct gradient analysis and nearest-neighbor imputation in coastal

Oregon, USA. *Canadian Journal of Forest Research* 32:725-741.

Phillips S. 2006. A brief tutorial on Maxent. AT&T Research. Retrieved March 2019 from

American Museum of Natural History Website.

Phillips SJ, Anderson RP, and Schapire RE. 2006. Maximum entropy modeling of

species geographic distributions. *Ecological Modelling* 190:231-259.

Phillips SJ, and Dudík M. 2008. Modeling of species distributions with Maxent: new

extensions and a comprehensive evaluation. *Ecography* 31:161-175.

Potvin F, Belanger L, and Lowell K. 2000. Marten habitat selection in a clearcut boreal

landscape. *Conservation Biology* 14:844-857.

Prevéy JS, Parker LE, and Harrington CA. 2020a. Projected impacts of climate change

on the range and phenology of three culturally-important shrub species. *PLoS*

ONE 15:e0232537. https://doi.org/10.1371/journal.pone.0232537

Prevéy JS, Parker LE, Harrington CA, Lamb CT, and Proctor MF. 2020b. Climate

change shifts in habitat suitability and phenology of huckleberry (*Vaccinium*

membranaceum). *Agricultural and forest meteorology* 280:107803.

PSW. 2019. Humbodlt marten data summary and report for the northern coastal

California population: distribution and population parameter estimates. USDI Fish

and Wildlife Service, USDA Forest Service Pacific Southwest Research Station.

p 33.

R Core Team. 2020. R: a language and environment for statistical computing.

https://www.R-project.org/. Vienna, Austria: R Foundation for Statistical

Computing.

Radosavljevic A, and Anderson RP. 2014. Making better Maxent models of species

distributions: complexity, overfitting and evaluation. *Journal of Biogeography*

41:629-643. https://doi.org/10.1111/jbi.12227

Raphael MG, and Molina R. 2007. *Conservation of rare or little-known species:*

biological, social, and economic considerations. Washington D.C., USA: Island

Press.

Rastogi B, Williams AP, Fischer DT, Iacobellis SF, McEachern K, Carvalho L, Jones C,

Baguskas SA, and Still CJ. 2016. Spatial and temporal patterns of cloud cover

and fog inundation in coastal California: Ecological implications. *Earth*

Interactions 20:1-19.

Romesburg HC. 1981. Wildlife science: gaining reliable knowledge. *Journal of Wildlife*

Management 45:293-313.

Rosenbaum PR. 2010. Design sensitivity and efficiency in observational studies.

Journal of the American Statistical Association 105:692-702.

10.1198/jasa.2010.tm09570

Schrott GR, and Shinn J. 2020. A landscape connectivity analysis for the coastal marten

(*Martes caurina humboldtensis*). Arcata, CA.

https://www.fws.gov/arcata/shc/marten/#report: USDI Fish and Wildlife Service. p

123.

Searcy CA, and Shaffer HB. 2016. Do ecological niche models accurately identify

climatic determinants of species ranges? *The American Naturalist* 187:423-435.

Shaw JH. 1995. How many bison originally populated western rangelands? *Rangelands*

17:148-150.

Shirk AJ, Raphael MG, and Cushman SA. 2014. Spatiotemporal variation in resource

selection: insights from the American marten (*Martes americana*). *Ecological*

Applications 24:1434-1444.

Shirk AJ, Wasserman TN, Cushman SA, and Raphael MG. 2012. Scale dependency of

American marten (*Martes americana*) habitat relations. In: Aubry KB, Zielinski

WJ, Raphael MG, Proulx G, and Buskirk SW, eds. *Biology and conservation of*

martens, sables, and fishers: a new synthesis. Ithaca, New York, United States:

Comstock Publishing Associates, 269-283.

Siren AP. 2020. Interacting effects of climate and biotic factors on mesocarnivore

distribution and snowshoe hare demography along the boreal-temperate

ecotoneDissertation. University of Massachusetts, Amherst.

Sirén AP, and Morelli TL. 2020. Interactive range-limit theory (iRLT): An extension for

predicting range shifts. *Journal of Animal Ecology* 89:940-954.

Slauson KM, Baldwin JA, and Zielinski WJ. 2012. Occupancy estimation and modeling

in *Martes* research and monitoring. In: Aubry KB, Zielinski W, Raphael MG,

Proulx G, and Buskirk SW, eds. *Biology and conservation of martens, sables,*

and fishers: a new synthesis. Ithaca, New York, USA: Comstock Publishing

Associates, 343-370.

Slauson KM, Schmidt GA, Zielinski WJ, Detrich PJ, Callas RL, Thrailkill J, Devlin-Craig

B, Early DA, Hamm KA, Schmidt KN, Transou A, and West CJ. 2019a. A

conservation assessment and strategy for the Humboldt marten in California and

Oregon In: U.S. Department of Agriculture FS, and Station PSR, editors. Arcata,

California: U.S. Department of Agriculture, Forest Service; Pacific Southwest

Research Station PSW-GTR-260. p 124.

Slauson KM, and Zielinski WJ. 2017. Seasonal specialization in diet of the Humboldt

marten (*Martes caurina humboldtensis*) in California and the importance of prey

size. *Journal of Mammalogy* 98:1697-1708. 10.1093/jmammal/gyx118

Slauson KM, Zielinski WJ, and Hayes JP. 2007. Habitat selection by American martens

in coastal California. *Journal of Wildlife Management* 71:458-468.

Slauson KM, Zielinski WJ, LaPlante DW, and Kirk TA. 2019b. A landscape suitability

model for the Humboldt marten (*Martes caurina humboldtensis*) in coastal

California and coastal Oregon. *Northwest Science*.

Sofaer HR, Jarnevich CS, Pearse IS, Smyth RL, Auer S, Cook GL, Edwards Jr TC,

Guala GF, Howard TG, and Morisette JT. 2019. Development and delivery of

species distribution models to inform decision-making. *BioScience* 69:544-557.

Stephens PA, Pettorelli N, Barlow J, Whittingham MJ, and Cadotte MW. 2015.

Management by proxy? The use of indices in applied ecology. *Journal of Applied*

Ecology 52:1-6.

Suffice P, Cheveau M, Imbeau L, Mazerolle MJ, Asselin H, and Drapeau P. 2020.

Habitat, climate, and fisher and marten distributions. *The Journal of Wildlife*

Management 84:277-292.

Sweitzer RA, Thompson CM, Green RE, Barrett RH, and Purcell KL. 2015. Survival of

fishers in the Southern Sierra Nevada region of California. *Journal of*

Mammalogy:gyv177.

Townsend AM. 2019. Tree squirrels and fishers in northern California: the effects of

masting hardwoods on stand use Master's of Science Thesis. Humboldt State

University.

Tweedy PJ, Moriarty KM, Bailey JD, and Epps CW. 2019. Using fine scale resolution

vegetation data from LiDAR and ground-based sampling to predict Pacific marten

resting habitat at multiple spatial scales. *Forest Ecology and Management*

452:117556.

U.S. Environmental Protection Agency. 2013. Level III ecoregions of the continental

United States. EPA- National Health and Environemental Effects Research

Laboratory: Corvallis, Oregon, USA.

USFWS. 2019. Species Status Assessment for the coastal marten (*Martes caurina*),

Version 2.1. Arcata, CA, USA: US Fish and Wildlife Service, Region 8. p 141.

USFWS. 2020. Endangered and threatened wildlife and plants; threatened species

status for coastal distinct population segment of the Pacific marten with a Section

4(d) rule *Federal Register* 85:63806-63831.

Velleman PF, and Welsch RE. 1981. Efficient computing of regression diagnostics. *The*

American Statistician 35:234-242. 10.2307/2683296

Wang T, Hamann A, Spittlehouse D, and Carroll C. 2016. Locally downscaled and

spatially customizable climate data for historical and future periods for North

America. *PLoS ONE* 11:e0156720.

Warren DL, and Seifert SN. 2011. Ecological niche modeling in Maxent: the importance

of model complexity and the performance of model selection criteria. *Ecological*

Applications 21:335-342.

Warren DL, Wright AN, Seifert SN, and Shaffer HB. 2014. Incorporating model

complexity and spatial sampling bias into ecological niche models of climate

change risks faced by 90 C alifornia vertebrate species of concern. *Diversity and*

Distributions 20:334-343.

Wasserman TN, Cushman SA, Schwartz MK, and Wallin DO. 2010. Spacial scaling and

multi-model Inference in landscape genetics: *Martes americana* in northern

Idaho. *Landscape Ecology* 25:1610-1612.

Whittaker RH. 1960. Vegetation of the Siskiyou mountains, Oregon and California.

Ecological Monographs 30:279-338.

Wiens JD, Anthony RG, and Forsman ED. 2014. Competitive interactions and resource

partitioning between northern spotted owls and barred owls in western Oregon.

Wildlife Monographs 185:1-50.

Wisz MS, Pottier J, Kissling WD, Pellissier L, Lenoir J, Damgaard CF, Dormann CF,

Forchhammer MC, Grytnes JA, and Guisan A. 2013. The role of biotic

interactions in shaping distributions and realised assemblages of species:

implications for species distribution modelling. *Biological Reviews* 88:15-30.

Zielinski WJ, Slauson KM, Carroll CR, Kent CJ, and Kudrna DG. 2001. Status of

American martens in coastal forests of the Pacific states. *Journal of Mammalogy*

82:478-490.

1243

1244