1 **Ancient genomes reveal hybridisation between extinct short-faced bears and the extant**
2 **spectacled bear (*Tremarctos ornatus*)**
3

4 Alexander T Salis[1#*], Graham Gower[1,2], Blaine W. Schubert[3], Leopoldo H. Soibelzon[4], Holly
5 Heiniger[1], Alfredo Prieto[5], Francisco J. Prevosti[6,7], Julie Meachen[8], Alan Cooper[9], Kieren J.
6 Mitchell[1*]
7

[1]Australian Centre for Ancient DNA (ACAD), School of Biological Sciences, University of Adelaide, South
Australia 5005, Australia

[2]Lundbeck GeoGenetics Centre, GLOBE Institute, University of Copenhagen, Copenhagen 1350, Denmark

[3]Center of Excellence in Paleontology and Department of Geosciences, East Tennessee State University
(ETSU), Johnson City, Tennessee 37614, USA

[4]División Paleontología de Vertebrados, Museo de La Plata, 1900 La Plata, Argentina

[5]Centro de Estudios del Hombre Austral: Instituto de la Patagonia, Universidad de Magallanes, Punta Arenas,
Chile

[6]Museo de Ciencias Antropológicas y Naturales, Universidad Nacional de La Rioja (UNLaR), La Rioja,
Argentina

[7]Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)

[8]Anatomy Department, Des Moines University, Des Moines, IA, USA

[9]South Australian Museum, Adelaide, South Australia 5000, Australia

Lead contact

*Corresponding author(s): A.T.S. (alexander.t.salis@gmail.com), and K.J.M. (kieren.mitchell@adelaide.edu.au)
23

Summary:

Two genera and multiple species of short-faced bear from the Americas went extinct during
or toward the end of the Pleistocene, and all belonged to the endemic New World subfamily
Tremarctinae [1-7]. Two of these species were giants, growing in excess of 1,000 kg [6, 8, 9],
but it remains uncertain how these extinct bears were related to the sole surviving short-faced
bear: the spectacled bear (*Tremarctos ornatus*). Ancient mitochondrial DNA has recently
suggested phylogenetic relationships among these lineages that conflict with interpretations
based on morphology [1, 10-12]. However, widespread hybridisation and incomplete lineage
sorting among extant bears mean that the mitochondrial phylogeny frequently does not reflect
the true species tree [13, 14]. Here we present ancient nuclear genome sequences from
representatives of the two extinct short-faced bear genera, *Arctotherium* and *Arctodus*. Our
new data support a third hypothesis for the relationships among short-faced bears, which
conflicts with existing mitochondrial and morphological data. Based on genome-wide D-
statistics, we suggest that the extant spectacled bear derives substantial ancestry from
Pleistocene hybridisation with an extinct short-faced bear lineage, resulting in a discordant
phylogenetic signal between the mitochondrion and portions of the nuclear genome.

Results and Discussion:

The spectacled bear (*Tremarctos ornatus)* is the only extant species of short-faced bear (Tremarctinae), a once diverse subfamily endemic to the Americas. This subfamily also includes many species that became extinct during the Pleistocene, including the Florida cave bear (*Tremarctos floridanus*), two species of North American short-faced bears (*Arctodus* spp. [3, 4]), and as many as five species of South American short-faced bears (*Arctotherium* spp. [2, 6]), one of which (*Arctotherium wingei*) has recently been discovered as far north as the Yucatan of Mexico [5]. Notably, the genera *Arctodus* and *Arctotherium* both included giant (>1,000kg) forms [8, 9] — *Arctodus simus* and *Arctotherium angustidens*, respectively — and based on morphology it was hypothesised that these genera were closely related [1, 6, 10, 11]. However, recently published mitochondrial DNA data suggested that *Arctotherium* was most closely related to the extant spectacled bear, to the exclusion of North American *Arctodus* [12]. While this result supported the convergent evolution of giant bears in North and South America, the mitochondrial genome does not always reflect the true relationships among species [e.g. 15, 16-19]. Importantly, discordance between mitochondrial and nuclear loci has been previously noted in bears, and has been attributed to a combination of stochastic processes and the rapid evolution of bears [13], as well as hybridisation between species [13, 14, 20-25]. To further resolve the evolutionary history of short-faced bears, we applied ancient DNA techniques to retrieve and analyse whole genome data from both *Arctodus* and *Arctotherium.*

Ancient DNA (aDNA) was extracted and sequenced from three *Arctodus simus* specimens: one each from placer mines at Sixty Mile Creek (ACAD 438; Canadian Museum of Nature; CMN 42388) and Hester Creek (ACAD 344; Yukon Government; YG 76.4) in the Yukon Territory, Canada; and one from Natural Trap Cave in Wyoming, USA (ACAD 5177; University of Kansas; KU 31956). We also analysed one specimen of *Arctotherium* sp. from Cueva del Puma, Patagonia, Chile (ACAD 3599; complete right femur, no. 32104, Centro de Estudios del Hombre Austral, Instituto de la Patagonia, Universidad de Magallanes). The *Arctotherium* specimen was previously dated to $12,105 \pm 175$ cal yBP (Ua-21033) [26], while two of the *Arctodus* specimens have been dated: ACAD 438 at $47,621 \pm 984$ cal yBP (TO-2699) [27] and ACAD 5177 at $24,300 \pm 208$ cal yBP (OxA-37990) (Table S1). The *Arctotherium* specimen has yielded mitochondrial aDNA in a previous studies [12], however, here we shotgun sequenced this specimen, along with the three *A. simus* specimens, at much

greater depth in order to reconstruct nuclear genome sequences. Mapping our new
sequencing data from these specimens to the giant panda (*Ailuropoda melanoleuca*) reference
genome (LATN01) yielded average depths of coverage between 0.12x to 5.9x for the *A.*
simus specimens and 3.9x for the *Arctotherium* specimen (Table S3). We compared these
new genomic data to previously published genomes from all extant species of bear (Table
S2): spectacled bear, giant panda, brown bear (*Ursus arctos*), American black bear (*U.*
americanus), Asian black bear (*U. thibetanus*), polar bear (*U. maritimus*), sloth bear (*U.*
ursinus), and sun bear (*U. malayanus*).
81

Phylogenetic analyses on a concatenated dataset of genome-wide SNPs revealed
relationships within Ursinae that were consistent with previous genomic studies: *U.*
americanus, U. maritimus, and *U. arctos* formed a monophyletic clade sister to a clade
consisting of *U. thibetanus*, *U. malayanus*, and *U. ursinus* [13, 14]. In contrast, within short-
faced bears (Tremarctinae) we recovered strong support for a close relationship between the
spectacled bear and the North American short-faced bear (*Arctodus simus*) to the exclusion of
the South American *Arctotherium* (Figure 1A, Figure S2). This result conflicts with the
mitochondrial tree, which instead supports a clade comprising *Arctotherium* and *Tremarctos*
ornatus to the exclusion of *Arctodus simus* [12] (Figure 1B). As the radiation of bears is
thought to have occurred rapidly during the Miocene - Pliocene transition, it is possible that
this discordance could be explained by incomplete lineage sorting (ILS) [28], a process
whereby pre-existing genetic variation in an ancestral species is randomly inherited and fixed
in descendant species [29, 30]. Alternatively, given the observed propensity of bears for
hybridisation [e.g. 13, 14, 20-22, 25, 31], mitochondrial/nuclear discordance within short-
faced bears may instead result from gene flow between *Tremarctos* and either *Arctodus* or
Arctotherium.
98

100
Figure 1: Phylogenetic relationships among ursids **A.** Maximum likelihood tree based on nuclear
SNPs constructed in RAxML. Branch labels represent bootstrap support percentages. For
RAxML tree with all individuals analysed see Figure S2. **B.** Bayesian phylogeny based on full
mitochondrial genomes adapted from Mitchell, et al. [12]. Blue bars represent 95% highest
posterior density interval on node ages. Branch labels represent BEAST posterior support
values.

To test for potential phylogenetic discordance across our short-faced bear genomes, we

constructed phylogenetic trees from 500 kb non-overlapping windows (n = 2622) across the

85 largest autosomal scaffolds of the giant panda reference genome (LATN01). Trees created

from roughly 70% of windows agreed with the results from our genome-wide concatenated

dataset (Topology 1; *i.e. Tremarctos + Arctodus*; Figure 2B & S3). However, approximately

30% of windows instead supported the mitochondrial tree topology (Topology 2; *i.e.*

Tremarctos + Arctotherium; Figure 2B & S3), while the third possible topology where the

two extinct genera form a clade — *Arctodus + Arctotherium —* was rejected for over 95% of

windows. The frequencies of the three possible tree topologies are difficult to explain as a

result of ILS, which we would expect to result in a more even representation of the two

"minority" topologies (*i.e.* Topologies 2 and 3). Our results therefore suggest that

introgression may be the most likely explanation for the observed phylogenetic discordance.

Consequently, we calculated D-statistics [32, 33] using our concatenated genome-wide SNPs

in order to identify signals of hybridisation between the bear species in our dataset.

Figure 2: A. The three possible short-faced bear (Tremarctinae) tree topologies. **B.** Discordance visualisation using DiscoVista from 2622 500 kb genomic fragments. The x-axis represents topologies tested and the y-axis the proportion of fragments that support the topology, with >80% bootstrap support used to define strong support. For more comprehensive tests of phylogenetic placements see Figure S3. **C.** Divergence time estimates (TMRCA) of *Tremarctos, Arctodus, and Arctotherium* (Node 2) and for sister species (Node 1) of the two most common short-faced bear topologies.

Consistent with previous studies [i.e. 14], our D-statistics revealed compelling evidence for hybridisation between: Asian black bears and all North American ursine bears (including the polar bear); sun bears and North American ursine bears; and Asian black bear and sun bear (Table S4). In contrast, we did not obtain any significantly non-zero values for D-statistics calculated using our two extinct short-faced bear genomes, any member of Ursinae, and the panda outgroup (Table 1). This result suggests that no gene flow occurred between *Arctodus* or *Arctotherium* and the ancestors of any modern ursine bear, and also demonstrates a lack of any discernible reference bias in the ancient genomic data (which would result in asymmetrical allele sharing with the reference). Thus, it appears *Arctodus* and *Arctotherium* did not hybridise with brown and black bears in the Americas during the late Pleistocene, even though the distribution of *Arctodus* overlapped with both ursines, and *Arctotherium* may have encountered them in Mexico or Central America [5]

Contrary to previous studies, our D-statistics revealed signals consistent with gene flow between the spectacled bear and members of Ursinae (Table 1 & S5), suggesting the possibility that *Tremarctos* hybridised with ancestors of either the brown bear or American

black bear during the Pleistocene. This signal is surprising given the deep divergence
between ursine and short-faced bears, having split approximately 10 million years ago (mya)
[12, 14, 28]. However, in support of this hypothesis, offspring between spectacled bear and
Asiatic black bear have resulted from hybridisation in zoos, although whether these hybrids
were fertile remains unknown [34]. Importantly, members of *Tremarctos* and the ancestors of
modern American black bears had overlapping distributions throughout the Pleistocene in
North America [4, 10], meaning that hybridisation may have occurred when the two lineages
were less divergent and reproductive barriers had had less time to evolve.

In addition to evidence for hybridisation between *Tremarctos* and ursine bears, we also
recovered convincing evidence for hybridisation between *Arctotherium* and *Tremarctos*
(Table 1). These results are consistent with a model where the divergence between *Arctodus*
and *Tremarctos* occurred in North America after the ancestors of *Arctotherium* dispersed
southwards into South America, with subsequent hybridisation between *Tremarctos* and
Arctotherium. This interpretation is supported by the presence of *Arctodus* and *Tremarctos*
(and absence of *Arctotherium*) in the late Pliocene fossil record of North America [3, 4, 7,
10]. The fossil record further suggests that contact between *Tremarctos* and *Arctotherium*
occurred during the late Pleistocene, when representatives of *Arctotherium* were distributed
as far north as the Yucatan of Mexico [5], providing an opportunity for hybridisation.

D-statistic: D(H1, H2, H3, Giant Panda)	D	Stderr	Z-Score	BABA	ABBA	nSNPs
D(*T. ornatus* (Chaparri), *Arctodus*, *Arctotherium*)	0.3116	0.009559	32.6*	41219	21633	6071021
D(*T. ornatus* (Nobody), *Arctodus*, *Arctotherium*)	0.3112	0.009462	32.889*	41187	21637	6070446
D(*T. ornatus* (Chaparri), *T. ornatus* (Nobody), *Arctodus*)	0.0206	0.022575	0.911	1375	1320	6393667
D(*T. ornatus* (Chaparri), *T. ornatus* (Nobody), *Arctotherium*)	0.0172	0.021533	0.801	1227	1186	6258343
D(*T. ornatus* (Nobody), *Arctotherium* sp., *Ursus arctos*)	0.2079	0.008878	23.418*	11177	7329	6184195
D(*T. ornatus* (Chaparri), *Arctotherium* sp., *Ursus arctos*)	0.2074	0.009242	22.442*	11185	7342	6185696
D(*T. ornatus* (Nobody), *Arctodus simus*, *Ursus arctos*)	0.2152	0.009651	22.302*	11172	7214	6317724
D(*T. ornatus* (Chaparri), *Arctodus simus*, *Ursus arctos*)	0.2131	0.010012	21.281*	11191	7260	6319318
D(*Arctodus simus*, *Arctotherium* sp., *Ursus malayanus* (Klaus))	0.0215	0.00854	2.523	9662	9254	6010428
D(*Arctodus simus*, *Arctotherium* sp., *Ursus americanus*)	0.0225	0.009168	2.453	9661	9236	6025070
D(*Arctodus simus*, *Arctotherium* sp., *Ursus malayanus* (Anabell))	0.0211	0.009128	2.308	9691	9291	6009103
D(*Arctodus simus*, *Arctotherium* sp., *Ursus maritimus* (PB1))	0.0187	0.009374	1.99	9677	9323	6019285
D(*Arctodus simus*, *Arctotherium* sp., *Ursus arctos*)	0.0167	0.009247	1.802	9717	9398	6033406
D(*Arctodus simus*, *Arctotherium* sp., *Ursus maritimus* (PB9))	0.016	0.009483	1.683	9752	9446	6044167
D(*Arctodus simus*, *Arctotherium* sp., *Ursus ursinus*)	0.016	0.009584	1.666	9608	9306	6012634
D(*Arctodus simus*, *Arctotherium* sp., *Ursus thibetanus*)	0.0157	0.009487	1.652	9644	9346	6019681

If the ancestors of the spectacled bear hybridised with *Arctotherium* somewhere in the

American mid-latitudes during the migration of *Tremarctos* into South America, then signals

of gene flow between members of these two genera could date to the latest Pleistocene or

earliest Holocene, when spectacled bears are thought to have migrated into South America [6,

35, 36]. To test this hypothesis, we estimated divergence times among the three short-faced

bear lineages for all 500 kb windows from the largest 40 scaffolds corresponding to either

Topology 1 (n = 980) or Topology 2 (n = 413) and summarised the results (Figure 2c). The

age of the most recent common ancestor (TMRCA) of *Tremarctos, Arctodus, and*

Arctotherium was similar irrespective of topology (Topology 1: 3.6 mya; Topology 2: 3.6

mya), as was the subsequent divergence between the remaining two lineages (Topology 1: 3.1

mya; Topology 2: 3.1 mya). Assuming that members of *Tremarctos* migrated southward no

earlier than the latest Pleistocene, our results superficially appear to be incompatible with late

Pleistocene/Holocene hybridisation between *Tremarctos* and *Arctotherium*. The fossil record

suggests two ways these observations may be explained.

Late Pleistocene fossil data indicate that the ancestors of the spectacled bear are likely

to have encountered *Arctotherium* individuals from Mexico, Central America, and/or northern

South America, which were comparable in size and diet to the spectacled bear [1, 5, 37] and

which may have represented a different *Arctotherium* species from the Chilean specimen

sequenced in the present study [1, 6, 12, 26]. Indeed, throughout the Pleistocene a number of

Arctotherium species have been described across South and Central America, with putative

species ranging from gigantic in the early-mid Pleistocene to relatively small in the late

Pleistocene [1, 2, 6, 9]. If the ancestors of our sampled Patagonian *Arctotherium* specimen

diverged from those of more northerly *Arctotherium* species during the Pliocene or early

Pleistocene, then our molecular dating results remain consistent with hybridisation being the

primary driver of phylogenetic discordance in our genomic data. Alternatively, hybridisation

between *Tremarctos* and *Arctotherium* could have occurred in Central America during the

Pleistocene. *Tremarctos* and *Arctotherium* have both been recorded in Central American cave

deposits [5, 38], however, the extent of occupation by both genera in the region is unknown,

and conceivably Central America represents a contact zone between the genera throughout the

Pleistocene where hybridisation may have occurred.

An alternative interpretation of our phylogenetic results is that Topology 2 (*Tremarctos*

+ *Arctotherium*), which is supported by the mitochondrion and ~30% of our nuclear genome

windows, is the pre-hybridisation tree. Recently, Li, et al. [39] suggested that under scenarios

involving substantial gene flow the predominant phylogenetic signal across the genome may

not reflect the pre-hybridisation tree. If this were the case for short-faced bears, the majority

of support for Topology 1 would actually result from extensive hybridisation between

Arctodus and *Tremarctos* in North America. Li, et al. [39] contend that the phylogenetic

signal of the pre-hybridisation tree may be enriched in regions of low recombination,

especially on the X-chromosome. In order to test this hypothesis, we identified panda

scaffolds corresponding to the ~40 Mb recombination cold-spot on the X-chromosome

highlighted by Li, et al. [39] and produced phylogenetic trees for each 500 kb window along

this region (Figure S4). Interestingly, the majority of these fragments supported Topology 2

(*Tremarctos + Arctotherium*), the same topology as the mitochondrial phylogeny but

contrasting with the majority of autosomal scaffolds.

Unlike felids [e.g. 40, 41, 42], a high-quality reference assembly and linkage map does not exist for any bear species, meaning scaffolds pertaining to high and low recombination areas of the genome could not be identified. Unfortunately, this currently makes it impossible to further explore the possibility that Topology 2 (*Tremarctos* + *Arctotherium*) may reflect the pre-hybridisation short-faced bear tree, rather than Topology 1 (*Tremarctos* + *Arctodus*). In the absence of a linkage map, sequencing aDNA from either the extinct *Tremarctos floridanus* or more northerly *Arctotherium* populations will be key to further resolving the evolutionary history of short-faced bears, though this will be challenging given that the core range of these species lies in the lower-latitudes where aDNA preservation is less reliable. For now we conclude that the weight of evidence supports a closer relationship between the spectacled bear and the extinct short-faced bears from North America (*Arctodus*) rather than South America (*Arctotherium*). In any case, our genomic data imply extensive hybridisation occurred between the spectacled bear and one of the extinct short-faced bear lineages. These results contribute to the growing consensus that hybridisation is widespread among carnivoran groups generally [13, 14, 39, 43].

Acknowledgements:

We would like to thank the following institutions for allowing access to specimens: Canadian Museum of Nature, University of Kansas Natural History Museum, Yukon Government, Centro de Estudios del Hombre Austral, Instituto de la Patagonia, Universidad de Magallanes. In addition, we are grateful to the following individuals who helped in the collection and identification of specimens and/or provided laboratory support: Grant Zazula (Yukon Territorial Government, Palaeontology Program, Canada), Fabiana Martin (Universidad de Magallanes, Chile), Jeremy Austin (University of Adelaide, Australia) and Sarah Bray (University of Adelaide, Australia). We would like to thank the Wyoming BLM and permit number PA-13-WY-207. This research was funded by an Australian Research Council Laureate Fellowship awarded to AC (FL140100260), U.S. National Science Foundation grant (EAR/SGP#1425059) awarded to JM and AC, and Agencia Nacional de Promoción Científica y Técnica' (ANPCyT, Argentina) (PICT 2015–966) awarded to FJP.

Author Contributions:

Conceptualization, A.T.S., A.C. and K.J.M.; Methodology, A.T.S., G.G., A.C., and K.J.M.; Investigation, A.T.S., H.H., and K.J.M.; Writing – Original Draft, A.T.S. and K.J.M.; Writing – Review & Editing, G.G., B.W.S., L.H.S., H.H., A.P., F.J.P., J.M., and A.C.; Funding

Materials and Methods:

Sampling

Analyses were performed on three bone samples identified as *Arctodus simus* and one sample identified as *Arctotherium* sp. (Table S1). The *Arctotherium* specimen ACAD 3599, had previously be radiocarbon dated, as well as one of the *A. simus* specimens (ACAD 438), a further *A. simus* specimen was radiocarbon dated at the Oxford Radiocarbon Accelerator Unit of the University of Oxford. All radiocarbon dates were calibrated with the either the IntCal13 curve [44] or the SHCal13 curve [45] using OxCal 4.4 [46] (Table S1).

Sample preparation and extraction

All pre-PCR steps (*i.e.,* extraction, library preparation) were conducted in purpose-built ancient DNA clean-room facilities at the University of Adelaide's Australian Centre for Ancient DNA (ACAD). Potential surface contamination on each sample was reduced by UV irradiation for 15 min each side, followed by abrasion of the exterior surface (c. 1 mm) using a Dremel tool and a disposable carborundum disk. The sample was then pulverised using a metallic mallet. Approximately 100 mg of powder was extracted using an in-house silica-based extraction protocol adapted from Dabney, et al. [47] optimised for the recovery of small fragments. the powder was digested first in 1 mL 0.5 M EDTA for 60 min, followed by an overnight incubation in 970 µL fresh 0.5 M EDTA and 30 µL proteinase K (20 mg/ml) at 55°C. The samples were centrifuged and the supernatant mixed with 13 mL of a modified PB buffer (12.6 mL PB buffer (Qiagen), 6.5 µL Tween-20, and 390 µL of 3M Sodium Acetate) and bound to silicon dioxide particles, which were then washed two times with 80% ethanol. The DNA was eluted from silica particles with 100 µL TE buffer.

Library preparation

Double-stranded Illumina libraries were constructed following the protocol of Meyer, et al. [48] from 25 µL of DNA extract. In addition, all samples underwent partial uracil-DNA glycosylase (UDG) treatment [49] to restrict cytosine deamination, characteristic of ancient DNA, to terminal nucleotides. A short round of PCR using PCR primers complementary to the library adapter sequences was performed to increase the total amount of DNA and add full-length Illumina sequencing adapters. Cycle number was determined via rtPCR and each library split into 8 separate PCR reactions to minimise PCR bias and maintain library complexity. Each PCR of 25 µL contained 1× HiFi buffer, 2.5 mM MgSO4, 1 mM dNTPs,

295   0.5 mM each primer, 0.1 U Platinum Taq Hi-Fi polymerase and 3 μL DNA. The cycling
296   conditions were 94 °C for 6 min, 8–10 cycles of 94 °C for 30 s, 60 °C for 30 s, and 72 °C for
297   40 s, followed by 72 °C for 10 min. Following PCR, replicates were pooled and purified using
298   AxyPrep™ magnetic beads, eluted in 30 μL H2O quantified on TapeStation (Agilent
299   Technologies).
300
301   *Sequencing*
302   Libraries were initially pooled and sequenced on an Illumina NextSeq using 2 x 75 bp PE
303   (150 cycle) High Output chemistry. For deeper sequencing, libraries were diluted to 1.5 nM
304   and each was run on one lane of an Illumina HiSeq X Ten using 2 x 150 bp PE (300 cycle)
305   chemistry, except for ACAD 438 which was run on two lanes of an Illumina HiSeq X Ten.
306
Data processing
Demultiplexed sequencing reads were processed through Paleomix v1.2.12 [50]. Within
Paleomix, raw reads were filtered, adapter sequences removed, and pair-end reads merged
using ADAPTER REMOVAL v2.1.7 [51], trimming low quality bases (<Phred20 --
minquality 4) and discarding merged reads shorter than 25 bp (--minlength 25). Read quality
was   visualised   before   and   after   adapter   trimming   using   fastQC   v0.11.5
(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/)   to   ensure   efficient   adapter
removal. Reads were mapped to the Panda ASM200744v1 genome [52] with BWA v0.7.15
using the mem algorithm [53]. Reads with mapping Phred scores less than 25 were removed
using   SAMtools   1.5   [54]   and   PCR   duplicates   were   removed   using   "paleomix
rmdup_collapsed"   and   MARKDUPLICATES   from   the   Picard   package
(http://broadinstitute.github.io/picard/). Indel realignment was performed using GATK [55]
and damage profiles assessed using MapDamage v2.0.8 [56] (Figure S1).
320
321   Sequencing reads were downloaded from the European Nucleotide Archive for all
322   extant bear species (Table S2) [14, 21, 24, 57, 58] and processed using the same pipeline as
323   for the ancient samples.
324
Phylogenetic analysis
Indexed VCF files were created for each BAM file using mpileup, part of the SAMtools
package v0.1.19 [54], and the call and index functions as a part of the BCFtools package
v0.1.19. Parallel v2010622 [59] was used to process each BAM file in parallel. BCFtools was

then used to filter SNPs within 3 bp of an indel (--SnpGap 3). The 85 largest scaffolds of the

Panda reference genome were renamed as chromosomes (chr1–85) in each VCF file using

BCFtools annotate. Biallelic variants in VCF files were converted to random pseudohaploid

variants in eigenstrat format for the 85 largest scaffolds using vcf2eig (part of eig-utils;

https://github.com/grahamgower/eig-utils) including monomorphic (-m) and singleton (-s)

sites, and excluding transitions (-t). Eigenstrat formatted files were then converted to PHYLIP

files using eig2phylip (part of eig-utils; https://github.com/grahamgower/eig-utils). A

supermatrix tree was then created in RAxML v8.2.4 [60] using the rapid bootstrapping

algorithm (-f a) and using the GTRCAT model of substitution with ascertainment correction

(-m ASC_GTRCAT) with 100 bootstrap replicates (-#100) and using the Felsenstein

ascertainment correction (--asc-corr=felsenstein) based on the number of invariant sites

(calculated from the total ungapped length of the largest 85 scaffolds of the Panda reference

genome minus the length of the alignment).

Discordance analysis using DiscoVista

The eigenstrat files were broken down into non-overlapping 500kb sliding windows using

eigreduce (part of eig-utils; https://github.com/grahamgower/eig-utils). We only used the

higher coverage A. simus sample (ACAD 344) in these analyses. For each window a PHYLIP

file and tree were created as described above. The frequency and support of different tree

topologies was then summarised and visualised using DiscoVista [61], using bootstrap values

of 80 as the cutoff for strong support. Topologies tested included: 1) the inclusion of

Arctotherium with ursine bears; 2) the inclusion of *Arctodus* with ursine bears; 3) the

inclusion of *Tremarctos* with ursine bears; 4) any combination of tremarctine bears included

with ursine bears; 5) the monophyly of Tremarctinae; 6) monophyly of *Tremarctos* and

Arctodus; 7) monophyly of *Tremarctos* and *Arctotherium*; and 8) monophyly of *Arctotherium*

and *Arctodus*.

D-statistics

To test for signals of gene flow within Tremarctinae and between tremarctine and ursine

lineages we used D-statistics as implemented by Admixtools [62] in admixr [63]. We only

used the higher coverage *A. simus* sample (ACAD 344) in this analysis. The giant panda was

used as outgroup and block jack-knife procedure used to test for significant departures from

zero ($|Z|>3$). D-statistics within Tremarctinae were calculated in the form D(*Arctodus*,

Tremarctos, *Arctotherium*, panda) and for detecting gene flow between Tremarctinae and

Ursinae in the form D(U1, U2, T1, panda), where T1 is any short-faced bear and U1 and U2

any ursine individual. D-statistics were also performed to detect gene flow within Ursinae (as

per Kumar, et al. 2017), using either the giant panda or spectacled bear as outgroup. To

account for the possibility of a reference bias in ancient samples, within Tremarctinae D-

statistics were recalculated using the Asiatic black bear as outgroup (Table S5).

Molecular dating

Divergence times were estimated for each 500kb fragment from the discordance analysis

using MCMCtree, part of the PAML package v4.8a [64], using the topology from the ML tree

produced in the discordance analysis as the input tree. Four calibrations were used to calibrate

the phylogeny:

1. The crown-age of Ursidae (*i.e.* the divergence of the giant panda lineage) was

constrained to between 11.6 and 23 million years ago (mya) based on the presence of

Kretzoiarctos [65], a putative ailuropodine, in the middle Miocene and the assumption

that early Miocene *Ursavus* representatives are likely ancestral to modern ursids [10].

2. The divergence of Tremarctinae and Ursinae was constrained to between 7 and 13

mya based on the presence of putative early tremarctine bears (*e.g. Plionarctos*) in the

Late Miocene/Early Pliocene [66].

3. The common ancestor of all sampled ursine bears was constrained to between 4.3 and

6 mya based on the occurrence of *Ursus minimus* [14, 67].

4. The divergence of polar and brown bears was constrained to between 0.48 and 1.1

mya based on previous nuclear estimates [21, 24, 25].

The JC +G substitution model with 5 discrete gamma categories was used with

autocorrelated-rates, also known as the geometric Brownian diffusion clock model. Uniform

priors for node ages using the birth-death (BD) process were used [λ_{BD} = 1 (birth-rate), μ_{BD} =

1 (death-rate), and ρ_{BD} = 0.1 (sampling fraction for extant species)]. A gamma-Dirichlet

distribution was used for the prior on rate with an α shape parameter of 2 (diffuse prior). The

σ_i^2 prior was defined as a diffuse gamma-Dirichlet distribution (2,2). MCMC tree runs were

performed with a burn-in of 10000, and a sample size of 10000, sampling every ten iterations.

Median node ages were then averaged for each tree topology.

Low-recombining region of X-chromosome

Scaffolds of the panda ASM200744v1 reference genome [52] corresponding to low

recombination regions of the X chromosome were identified by mapping all scaffolds to the

recombination cold-spot of the X chromosome of the domestic cat (FelCat5) using minimap2.

Default parameters were used, meaning the alignment lacked base-level precision (to account

to phylogenetic distance between giant panda and the domestic cat). Only scaffolds larger

than 500kb and with greater than 100 kb of segments mapping to the low recombining region

of the domestic cat X-chromosome were retained, resulting in 15 scaffolds linked to the low

recombination region of the X-chromosome. A maximum-likelihood phylogenetic tree and

gene-tree discordance analysis were performed on these 15 scaffolds as described above for

the genome-wide dataset.

References:

1.   Soibelzon, L.H., Tonni, E.P., and Bond, M. (2005). The fossil record of South
American short-faced bears (Ursidae, Tremarctinae). J. South Am. Earth Sci. *20*, 105-
113.

2.   Soibelzon, L.H. (2004). Revisión sistemática de los Tremarctinae (Carnivora, Ursidae)
fósiles de América del Sur. Rev. Mus. Argent. Cienc. Nat *6*, 107-133.

3.   Kurtén, B. (1966). Pleistocene bears of North America. 1. Genus *Trematctos*, spectacled
bears. Acta Zool. Fenn. *115*, 1-120.

4.   Kurtén, B., and Anderson, E. (1980). Pleistocene Mammals of North America, (New
York: Columbia University Press).

5.   Schubert, B.W., Chatters, J.C., Arroyo-Cabrales, J., Samuels, J.X., Soibelzon, L.H.,
Prevosti, F.J., Widga, C., Nava, A., Rissolo, D., and Erreguerena, P.L. (2019). Yucatan
carnivorans shed light on the Great American Biotic Interchange. Biol Lett *15*,
20190148.

6.   Prevosti, F.J. (2018). Evolution of South American Mammalian Predators During the
Cenozoic: Paleobiogeographic and Paleoenvironmental Contingencies, 1st ed. 2018.
Edition, (Cham: Springer International Publishing).

7.   Kurtén, B. (1967). Pleistocene bears of North America. 2. Genus *Arctodus*, short-faced
bears. Acta Zool. Fenn. *117*, 1-60.

8.   Christiansen, P. (1999). What size were *Arctodus simus* and *Ursus spelaeus* (Carnivora:
Ursidae)? Ann. Zool. Fenn. *36*, 93-102.

9.   Soibelzon, L.H., and Schubert, B.W. (2011). The largest known bear, *Arctotherium*
angustidens, from the early Pleistocene Pampean region of Argentina: with a discussion
of size and diet trends in bears. J. Paleontol. *85*, 69-75, 67.

10.   McLellan, B., and Reiner, D.C. (1994). A review of bear evolution. Bears Their Biol.
Manag. *9*, 85-96.

11.   Trajano, E., and Ferrarezzi, H. (1995). A Fossil Bear from Northeastern Brazil, with a
Phylogenetic Analysis of the South American Extinct Tremarctinae (Ursidae). J.
Vertebr. Paleontol. *14*, 552-561.

12.   Mitchell, K.J., Bray, S.C., Bover, P., Soibelzon, L., Schubert, B.W., Prevosti, F., Prieto,
A., Martin, F., Austin, J.J., and Cooper, A. (2016). Ancient mitochondrial DNA reveals
convergent evolution of giant short-faced bears (Tremarctinae) in North and South
America. Biol. Lett. *12*, 20160062.

13.   Kutschera, V.E., Bidon, T., Hailer, F., Rodi, J.L., Fain, S.R., and Janke, A. (2014).
Bears in a forest of gene trees: Phylogenetic inference is complicated by incomplete
lineage sorting and gene flow. Mol. Biol. Evol. *31*, 2004-2017.

14.   Kumar, V., Lammers, F., Bidon, T., Pfenninger, M., Kolter, L., Nilsson, M.A., and
Janke, A. (2017). The evolutionary history of bears is characterized by gene flow across
species. Sci. Rep. *7*, 46487.

15.   Funk, D.J., and Omland, K.E. (2003). Species-Level Paraphyly and Polyphyly:
Frequency, Causes, and Consequences, with Insights from Animal Mitochondrial DNA.
Annu. Rev. Ecol. Evol. Syst. *34*, 397-423.

16.   Leache, A.D., and McGuire, J.A. (2006). Phylogenetic relationships of horned lizards
(*Phrynosoma*) based on nuclear and mitochondrial data: evidence for a misleading
mitochondrial gene tree. Mol. Phylogenet. Evol. *39*, 628-644.

17.   Wallis, G.P., Cameron-Christie, S.R., Kennedy, H.L., Palmer, G., Sanders, T.R., and
Winter, D.J. (2017). Interspecific hybridization causes long-term phylogenetic
discordance between nuclear and mitochondrial genomes in freshwater fishes. Mol.
Ecol. *26*, 3116-3127.

18.   Toews, D.P., and Brelsford, A. (2012). The biogeography of mitochondrial and nuclear
discordance in animals. Mol. Ecol. *21*, 3907-3930.

19.   Gompert, Z., Forister, M.L., Fordyce, J.A., and Nice, C.C. (2008). Widespread mito-
nuclear discordance with evidence for introgressive hybridization and selective sweeps
in *Lycaeides*. Mol. Ecol. *17*, 5231-5244.

20.   Barlow, A., Cahill, J.A., Hartmann, S., Theunert, C., Xenikoudakis, G., Fortes, G.G.,
Paijmans, J.L.A., Rabeder, G., Frischauf, C., Grandal-d'Anglade, A., et al. (2018).
Partial genomic survival of cave bears in living brown bears. Nat. Ecol. Evol. *2*, 1563-
1570.

21.   Cahill, J.A., Green, R.E., Fulton, T.L., Stiller, M., Jay, F., Ovsyanikov, N., Salamzade,
R., St. John, J., Stirling, I., Slatkin, M., et al. (2013). Genomic evidence for island

population conversion resolves conflicting Theories of polar bear evolution. PLoS
Genet. *9*, e1003345.

22. Cahill, J.A., Stirling, I., Kistler, L., Salamzade, R., Ersmark, E., Fulton, T.L., Stiller, M.,
Green, R.E., and Shapiro, B. (2015). Genomic evidence of geographically widespread
effect of gene flow from polar bears into brown bears. Mol. Ecol. *24*, 1205-1217.

23. Hailer, F. (2015). Introgressive hybridization: brown bears as vectors for polar bear
alleles. Mol. Ecol. *24*, 1161-1163.

24. Liu, S.P., Lorenzen, E.D., Fumagalli, M., Li, B., Harris, K., Xiong, Z.J., Zhou, L.,
Korneliussen, T.S., Somel, M., Babbitt, C., et al. (2014). Population genomics reveal
recent speciation and rapid evolutionary adaptation in polar bears. Cell *157*, 785-794.

25. Hailer, F., Kutschera, V.E., Hallstrom, B.M., Klassert, D., Fain, S.R., Leonard, J.A.,
Arnason, U., and Janke, A. (2012). Nuclear genomic sequences reveal that polar bears
are an old and distinct bear lineage. Science *336*, 344-347.

26. Martin, F.M., Prieto, A., Morello, F., Prevosti, F., and Borrero, L. (2004). Late
Pleistocene megafauna at Cueva del Puma, Pali-Aike Lava Field, Chile. Curr. Res.
Pleistocene *21*, 101-103.

27. Harington, C.R., Naughton, D., Dalby, A., Rose, M., and Dawson, J. (2003). Annotated
Bibliography of Quaternary Vertebrates of Northern North America, (Toronto:
University of Toronto Press).

28. Krause, J., Unger, T., Nocon, A., Malaspinas, A.S., Kolokotronis, S.O., Stiller, M.,
Soibelzon, L., Spriggs, H., Dear, P.H., Briggs, A.W., et al. (2008). Mitochondrial
genomes reveal an explosive radiation of extinct and extant bears near the Miocene-
Pliocene boundary. BMC Evol. Biol. *8*, 220.

29. Nichols, R. (2001). Gene trees and species trees are not the same. Trends Ecol. Evol.
16, 358-364.

30. Maddison, W.P. (1997). Gene trees in species trees. Syst. Biol. *46*, 523-536.

31. Cahill, J.A., Heintzman, P.D., Harris, K., Teasdale, M.D., Kapp, J., Soares, A.E.R.,
Stirling, I., Bradley, D., Edwards, C.J., Graim, K., et al. (2018). Genomic evidence of
widespread admixture from polar bears into brown bears during the last ice age. Mol.
Biol. Evol. *35*, 1120-1129.

32. Green, R.E., Krause, J., Briggs, A.W., Maricic, T., Stenzel, U., Kircher, M., Patterson,
N., Li, H., Zhai, W.W., Fritz, M.H.Y., et al. (2010). A Draft Sequence of the Neandertal
Genome. Science *328*, 710-722.

33. Durand, E.Y., Patterson, N., Reich, D., and Slatkin, M. (2011). Testing for ancient
admixture between closely related populations. Mol. Biol. Evol. *28*, 2239-2252.

34. Mondolfi, E., and Boede, E. (1981). A hybrid of a spectacled bear (*Tremarctos ornatus*)
and an asiatic black bear (*Selenarctos thibetanus*) born at the Maracay zoological park,
Venezuela. Mem. Soc. Cienc. Nat. La Salle *41*, 143-148.

35.   Stucchi, M., Salas-Gismondi, R., Baby, P., Guyot, J.-L., and Shockey, B.J. (2009). A
511   6,000 year-old specimen of a spectacled bear from an Andean cave in Peru. Ursus *20*,
512   63-68.

513   36.   García-Rangel, S. (2012). Andean bear *Tremarctos ornatus* natural history and
514   conservation. Mammal Rev. *42*, 85-119.

515   37.   Figueirido, B., and Soibelzon, L.H. (2010). Inferring palaeoecology in extinct
516   tremarctine bears (Carnivora, Ursidae) using geometric morphometrics. Lethaia *43*,
517   209-222.

518   38.   Czaplewski, N.J., Krejca, J., and Miller, T.E. (2003). Late quaternary bats from Cebada
519   Cave, Chiquibul cave system, Belize. Caribb. J. Sci. *39*, 23-33.

520   39.   Li, G., Figueiro, H.V., Eizirik, E., and Murphy, W.J. (2019). Recombination-aware
521   phylogenomics reveals the structured genomic landscape of hybridizing cat species.
522   Mol. Biol. Evol. *36*, 2111-2126.

523   40.   Menotti-Raymond, M., David, V.A., Roelke, M.E., Chen, Z.Q., Menotti, K.A., Sun, S.,
524   Schaffer, A.A., Tomlin, J.F., Agarwala, R., O'Brien, S.J., et al. (2003). Second-
525   generation integrated genetic linkage/radiation hybrid maps of the domestic cat (*Felis
526   catus*). J. Hered. *94*, 95-106.

527   41.   Menotti-Raymond, M., David, V.A., Lyons, L.A., Schaffer, A.A., Tomlin, J.F., Hutton,
528   M.K., and O'Brien, S.J. (1999). A genetic linkage map of microsatellites in the domestic
529   cat (*Felis catus*). Genomics *57*, 9-23.

530   42.   Li, G., Hillier, L.W., Grahn, R.A., Zimin, A.V., David, V.A., Menotti-Raymond, M.,
531   Middleton, R., Hannah, S., Hendrickson, S., Makunin, A., et al. (2016). A high-
532   resolution SNP array-based linkage map anchors a new domestic cat draft genome
533   assembly and provides detailed patterns of recombination. G3 (Bethesda) *6*, 1607-1616.

534   43.   Li, G., Davis, B.W., Eizirik, E., and Murphy, W.J. (2016). Phylogenomic evidence for
535   ancient hybridization in the genomes of living cats (Felidae). Genome Res. *26*, 1-11.

536   44.   Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck,
537   C.E., Cheng, H., Edwards, R.L., Friedrich, M., et al. (2013). Intcal13 and Marine13
538   radiocarbon age calibration curves 0-50,000 years cal BP. Radiocarbon *55*, 1869-1887.

539   45.   Hogg, A.G., Hua, Q., Blackwell, P.G., Niu, M., Buck, C.E., Guilderson, T.P., Heaton,
540   T.J., Palmer, J.G., Reimer, P.J., Reimer, R.W., et al. (2013). Shcal13 Southern
541   Hemisphere Calibration, 0-50,000 Years Cal Bp. Radiocarbon *55*, 1889-1903.

542   46.   Ramsey, C.B. (2009). Bayesian analysis of radiocarbon dates. Radiocarbon *51*, 337-
543   360.

544   47.   Dabney, J., Knapp, M., Glocke, I., Gansauge, M.-T., Weihmann, A., Nickel, B.,
545   Valdiosera, C., García, N., Pääbo, S., Arsuaga, J.-L., et al. (2013). Complete
546   mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from
547   ultrashort DNA fragments. Proc. Natl. Acad. Sci. U. S. A. *110*, 15758-15763.

548   48.   Meyer, M., Kircher, M., Gansauge, M.T., Li, H., Racimo, F., Mallick, S., Schraiber,
549   J.G., Jay, F., Prufer, K., de Filippo, C., et al. (2012). A high-coverage genome sequence
550   from an archaic Denisovan individual. Science *338*, 222-226.

551   49.   Rohland, N., Harney, E., Mallick, S., Nordenfelt, S., and Reich, D. (2015). Partial
552   uracil-DNA-glycosylase treatment for screening of ancient DNA. Philos. Trans. R. Soc.
553   Lond. B. Biol. Sci *370*, 20130624.

554   50.   Schubert, M., Ermini, L., Sarkissian, C.D., Jonsson, H., Ginolhac, A., Schaefer, R.,
555   Martin, M.D., Fernandez, R., Kircher, M., McCue, M., et al. (2014). Characterization of
556   ancient and modern genomes by SNP detection and phylogenomic and metagenomic
557   analysis using PALEOMIX. Nat. Protoc. *9*, 1056-1082.

558   51.   Schubert, M., Lindgreen, S., and Orlando, L. (2016). AdapterRemoval v2: rapid adapter
559   trimming, identification, and read merging. BMC Res. Notes *9*, 88.

560   52.   Hu, Y., Wu, Q., Ma, S., Ma, T., Shan, L., Wang, X., Nie, Y., Ning, Z., Yan, L., Xiu, Y.,
561   et al. (2017). Comparative genomics reveals convergent evolution between the bamboo-
562   eating giant and red pandas. Proc. Natl. Acad. Sci. U. S. A. *114*, 1081-1086.

563   53.   Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-
564   Wheeler transform. Bioinformatics *25*, 1754-1760.

565   54.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
566   Abecasis, G., Durbin, R., and Genome Project Data Processing, S. (2009). The sequence
567   alignment/map format and SAMtools. Bioinformatics *25*, 2078-2079.

568   55.   McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A.,
569   Garimella, K., Altshuler, D., Gabriel, S., Daly, M., et al. (2010). The Genome Analysis
570   Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.
571   Genome Res. *20*, 1297-1303.

572   56.   Jonsson, H., Ginolhac, A., Schubert, M., Johnson, P.L.F., and Orlando, L. (2013).
573   mapDamage2.0: fast approximate Bayesian estimates of ancient DNA damage
574   parameters. Bioinformatics *29*, 1682-1684.

575   57.   Zhao, S.C., Zheng, P.P., Dong, S.S., Zhan, X.J., Wu, Q., Guo, X.S., Hu, Y.B., He,
576   W.M., Zhang, S.N., Fan, W., et al. (2013). Whole-genome sequencing of giant pandas
577   provides insights into demographic history and local adaptation. Nat. Genet. *45*, 67-
578   U99.

579   58.   Miller, W., Schuster, S.C., Welch, A.J., Ratan, A., Bedoya-Reina, O.C., Zhao, F.Q.,
580   Kim, H.L., Burhans, R.C., Drautz, D.I., Wittekindt, N.E., et al. (2012). Polar and brown
581   bear genomes reveal ancient admixture and demographic footprints of past climate
582   change. Proc. Natl. Acad. Sci. U. S. A. *109*, E2382-E2390.

583   59.   Tange, O. (2011). Gnu parallel-the command-line power tool. The USENIX Magazine
584   *36*, 42-47.

585   60.   Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-
586   analysis of large phylogenies. Bioinformatics *30*, 1312-1313.

61. Sayyari, E., Whitfield, J.B., and Mirarab, S. (2018). DiscoVista: Interpretable
visualizations of gene tree discordance. Mol. Phylogenet. Evol. *122*, 110-115.

62. Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck,
T., Webster, T., and Reich, D. (2012). Ancient admixture in human history. Genetics
192, 1065-1093.

63. Petr, M., Vernot, B., and Kelso, J. (2019). admixr-R package for reproducible analyses
using ADMIXTOOLS. Bioinformatics *35*, 3194-3195.

64. Yang, Z. (2007). PAML 4: phylogenetic analysis by maximum likelihood. Mol. Biol.
Evol. *24*, 1586-1591.

65. Abella, J., Alba, D.M., Robles, J.M., Valenciano, A., Rotgers, C., Carmona, R.,
Montoya, P., and Morales, J. (2012). *Kretzoiarctos* gen. nov., the oldest member of the
giant panda clade. PLoS ONE *7*, e48985.

66. Tedford, R.H., and Martin, J. (2001). *Plionarctos*, a tremarctine bear (Ursidae:
Carnivora) from western North America. J. Vertebr. Paleontol. *21*, 311-321.

67. Vangengeim, E.A., Vislobokova, I., and Sotnikova, M. (1998). Large Ruscinian
Mammalia in the territory of the former Soviet Union. Stratigr. Geol. Correl *6*, 368-382.
603