Protein glycosylation is essential for SARS-CoV-2 infection

Aitor Casas-Sanchez[1,2]*, Alessandra Romero-Ramirez[1,2]§, Eleanor Hargreaves[3]§, Edward I Patterson[1,2], Grant L Hughes[1,2], Tobias Zech[3], Álvaro Acosta-Serrano[1,2]

[1]Department of Vector Biology and [2]Department of Tropical Disease Biology, Liverpool School of Tropical Medicine, UK

[3]Department of Molecular and Cellular Physiology, University of Liverpool, UK

§ Equal contribution

*Corresponding author: aitor.casas-sanchez@lstmed.ac.uk

Keywords: SARS-CoV-2, COVID-19, coronavirus, *N*-glycosylation, *N*-glycan, viral infection.

Abstract

SARS-CoV-2 extensively *N*-glycosylates its surface spike (S) proteins. This post-

translational modification is essential to modulate protein conformation and host cell

invasion. Each S monomer can be modified with up to 22 *N*-glycans. To meet the high

demand of protein glycosylation during virus replication, SARS-CoV-2 upregulates the

expression of host *N*-glycosylation genes. Although a substantial amount of detail is

known about the structure of S protein *N*-glycans, the role of *N*-glycosylation in SARS-

CoV-2 infection remains largely undetermined. Here, we investigated the essentiality

of the host *N*-glycosylation pathway and viral *N*-glycans for SARS-CoV-2 infection.

When either monkey or human cells were preincubated with glycosylation inhibitors,

including FDA-approved iminosugars, virus infection was significantly reduced. This

infection phenotype was confirmed after RNAi knockdown of several glycosylation

genes. In addition, enzymatic deglycosylation of whole viral particles confirmed that

accessible oligosaccharides on the SARS-CoV-2 surface are essential for host cell

infection. Altogether, we show evidence that the normal functioning of the host *N*-

glycosylation machinery is essential not only for SARS-CoV-2 to infect, but also to

produce new functional virions. These findings open the door for developing new

approaches targeting *N*-glycosylation against COVID-19.

Introduction

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), responsible for the COVID-19 pandemic, is a positive single-stranded RNA betacoronavirus composed of four structural proteins that contain the viral genome[1]. The SARS-CoV-2 surface spike glycoprotein has been extensively studied and is the target of most vaccines and treatments under development due to its exposure, immunogenicity and essential role in infection. It is well established that the main route for SARS-CoV-2 to infect the host cell is via attachment to the receptor angiotensin-converting enzyme 2 (ACE-2) through the spike's receptor-binding domain (RBD), followed by membrane fusion and invasion[2,3]. Both spike and ACE-2 are modified by the addition of a variable number of *N*-glycans. While ACE-2 contains seven occupied glycosylation sites[4], the homotrimeric spike protein has been shown to be heavily *N*-glycosylated with up to 66 oligosaccharides (22 per monomer) when recombinantly expressed in human HEK293 cells[5,6]. Cryo-electron microscopy has confirmed the presence of such glycans in real SARS-CoV-2 virions[7]. Furthermore, it has been recently suggested that spike *N*-glycans may play critical roles including proper protein folding and conformational changes[8], successful recognition and attachment to ACE-2 and masking certain epitopes to avoid antibody neutralization[3,5]. Most of these, however, are inferred from computational models or resemblance to other viruses. There is little direct evidence of the essentiality of spike *N*-glycans for SARS-CoV-2 infection. So far, only the use of pseudotyped vesicular stomatitis virus (VSV) expressing the SARS-CoV-2 spike as a proxy for the study of spike-driven host cell invasion has served useful for this purpose[9]. By mutating *N*-glycosylation sites, it has been shown that the absence of glycans on Asn^{331} and Asn^{343}, both within the RBD, resulted in a drastic reduction of pseudovirus infection[10]. In addition, knocking out *MGAT1* gene, which encodes for *N*-acetylglucosaminyl-transferase, an essential enzyme in the formation of hybrid and complex *N*-glycans, also reduced invasion of pseudotyped VSV suggesting that hybrid and complex spike *N*-glycans may be of importance for this process[11]. Although SARS-CoV-2 cDNA clones have been available for months[12,13], no study has yet used real virions to provide direct evidence.

The role of *N*-glycosylation, the main eukaryote protein post-translational modification, in the context of a SARS-CoV-2 infection remains unknown to date. The *N*-glycosylation pathway starts in the endoplasmic reticulum (ER) membrane where the

universal precursor $Glc_3Man_9GlcNAc_2$-dolichol-P is assembled in a stepwise manner.

The glycan portion is then co-translationally added to an acceptor asparagine residue

–within the sequon N-X-S/T; X≠P[14]– on a nascent polypeptide by STT3, the catalytic

subunit of the oligosaccharyl-transferase (OST) complex[15]. The attached glycan is

then modified by ER α-glucosidases I and II which remove the terminal glucose

residues thus allowing the glycoprotein to enter into the calnexin/calreticulin quality

control pathway and subsequent export to the Golgi apparatus[16]. In the Golgi, the

oligomannosidic *N*-glycan gets trimmed by the action of several α-mannosidases and

its structure can be further modified into a hybrid- or complex-type glycan thanks to

the action of MGAT1[17]. The resulting glycan structure is highly variable depending on

cell type and species. There are small molecules that can specifically inhibit enzymes

along the *N*-glycosylation pathway, especially the α-glucosidase I. Although many of

these compounds can be toxic, some are FDA-approved and used to treat human

diseases such as type I Gauche disease (Miglustat[18]) or diabetes (Miglitol[19],

Acarbose[20]). Others like Celgosivir, are in an advanced trial phase for their broad

antiviral activity[21]. Previous drug screening studies have shown the efficacy of some

of these and similar compounds against SARS-CoV-2[11,22-24].

In this study, we investigated the role of the host *N*-glycosylation pathway for SARS-

CoV-2 infection *in vitro*. We found that by inhibiting key *N*-glycosylation steps using

either chemical or genetic approaches, or by direct enzymatic deglycosylation, host

cell infection is drastically inhibited. Thus, glycosylation of viral proteins represents a

promising target against SARS-CoV-2 infection.

Results

N-glycosylation inhibitors reduce SARS-CoV-2 infection *in vitro*

We first hypothesized that the partial or complete inhibition of the host *N*-glycosylation pathway may interfere with the normal development of a SARS-CoV-2 infection. This drug-driven inhibition may either impede the normal processing and assembly of new virions, hamper their capacity to infect further host cells and/or prevent the successful recognition of cells presenting overall *N*-glycosylation defects, including key surface receptors such as ACE-2. We tested 10 compounds known to inhibit key steps along the *N*-glycosylation pathway, at concentrations considered to be at a low or high end of the effective range. As inhibitors of the early pathway in the ER, we used tunicamycin that prevents the formation of the *N*-glycan precursor, and NGI-1 that inhibits the ER's oligosaccharyltransferase catalytic subunit STT3. Further in the pathway, we inhibited ER α-glucosidase I using the iminosugars Miglitol, N-butyl-deoxynojirimycin (Miglustat), 1-deoxynojirimycin (DNJ), N-Nonyldeoxynojirimycin (NN-DNJ), Acarbose, Swainsonine and Celgosivir. In addition, we tested the efficacy of Deoxymannojirimycin (DMM) as an inhibitor of Golgi α-mannosidase I. Pre-incubation of either Vero E6 green monkey kidney epithelial cells or HEK293^{ACE-2} (lentiviral overexpression of human ACE-2) human kidney cells with most inhibitors prior to SARS-CoV-2 challenge (1 to 16 hours before infection), resulted in a significant reduction in infection compared to untreated cells (**Figure 1**). The same protective effect was observed when the inhibitors were added alongside the virus and up to 6 hours after infection (not shown). We determined the relative proportion of infection by scanning whole wells and measuring the area occupied by infected (α-spike-positive) cells in relation to the total (nuclei DAPI-stained) (**Figure 1a-c**). For selected inhibitors, we measured the proportion of cells showing cytopathic effects (CPE) as a result of the infection using crystal violet staining (**Figure 1d**) and quantified virus titres in cell supernatants by plaque assay (**Figure 1e**). While untreated controls displayed most cells as infected in clusters, both early inhibitors of the *N*-glycosylation pathway (i.e., tunicamycin and NGI-1) provided the highest protection rates against the virus in Vero (97.3% and 97%) and HEK cells (92.3% and 92%), showing only few isolated infected cells (**Figure 1a,b**). Tunicamycin and NGI-1 also protected cells against CPE (94% and 99.5%) (**Figure 1d**) and at high doses led to undetectable levels of infectious particles in cell supernatants, in a dose-dependent manner (**Figure 1e**). Inhibitors of

α-glucosidase I proved to be less effective and presented a greater variability. The

most effective ones (i.e., Miglustat, NN-DNJ and Celgosivir) showed reduced overall

infections in Vero and HEK cells, smaller clusters of infected cells, CPE protection and

moderate virus titre reduction in cell supernatants (**Figure 1**). Inhibition of the Golgi's

α-mannosidase I using DMM had no effect at preventing infection in Vero cells and a

slightly reduced infection in HEK cells. Viability assays using AlamarBlue were run in

parallel for all inhibitors to rule out poor metabolism or toxicity as the main causes for

infection decrease (**Figure 1e**).

Figure 1. Chemical inhibition of the host *N*-glycosylation pathway reduces SARS-CoV-2 infection *in vitro*. (**a**) Representative whole-well scans of confluent Vero E6 monolayers treated with either high, medium or low dose (see below) of several *N*-glycosylation inhibitors and infected with SARS-CoV-2 (MOI=0.05, 24 hours). Cells immuno-stained using anti-spike antibodies (green); DMSO untreated control; scale bar 5 mm. (**b**) Representative images of either Vero E6 or HEK293[ACE-2] cells in 'a' pre-treated with a high dose of *N*-glycosylation inhibitor and infected with SARS-CoV-2 (MOI=0.05, 24 hours for Vero; MOI=0.1 for 24 hours for HEK293). Cells stained with anti-Spike (grey channel) and merged (green) with DNA counterstain DAPI (blue); uninfected and DMSO untreated controls included scale bars 200μm. (**c**) Quantification of the percentage of infected (spike-positive) cells in 'a' normalized to DMSO controls (dashed horizontal line) of either Vero E6 or HEK293[ACE-2] cells pre-treated with inhibitors (at low, medium and high doses) and infected with SARS-CoV-2 (MOI=0.05, 24 hours). Bars indicate the mean values; dots indicate individual values; error bars represent standard deviation; two technical replicates; two independent experiments performed. Inhibitor concentrations low, medium to high (μM): tunicamycin (0.001, 0.01, 0.05); NGI-1 (0.001, 5, 75); DNJ (0.01, 60, 120); Miglitol (0.01, 50, 200); Acarbose (0.01, 100, 300); Miglustat (0.01, 50, 100); Celgosivir, Swainsonine, NN-DNJ and DMM (0.01, 25, 100). (**d**) Percentage of dead Vero E6 cells (area ratio of unstained cells with crystal violet) pre-treated with glycosylation inhibitors after SARS-CoV-2 infection (MOI=0.001, 48 hours). Line represents mean values, dots indicate individual values, error bars indicate standard deviation; two technical replicates, two independent experiments performed. (**e**) Concentration of infective viral particles (pfu/ml; in red) in supernatants of Vero E6 cells pre-treated with several inhibitors and infected with SARS-CoV-2 (MOI=0.001 for 48 hours), determined by plaque assay in Vero E6 cells. Toxicity percentage measured by AlamarBlue assay plotted alongside in grey. Dots indicate the mean values; error bars represent standard deviation; two technical replicates from two independent experiments.

<u>Genetic ablation of the *N*-glycosylation pathway reduces SARS-CoV-2 infection</u>

To validate the drug inhibition results, we used siRNAs to knock down genes encoding

for key enzymes in the *N*-glycosylation pathway prior to infection with SARS-CoV-2

(**Figure 2**). We knocked down 1) the OST's catalytic subunit STT3 (isoforms A, B and

both) to abolish transfer of *N*-glycan precursors to proteins, directly resembling the

inhibitory effects of NGI-1 and indirectly those of tunicamycin; 2) the GANAB catalytic

subunit of the ER α-glucosidase I to prevent the deglucosylation of *N*-glycans already

attached to proteins (emulating the effect that the α-glucosidase I inhibitors should

produce); and 3) the Golgi's MGAT1 enzyme to stop the formation of hybrid and

complex *N*-glycans. As expected, full knockdown of *STT3* (si*STT3A+B*) led to an

average infection reduction of ~55.4% in Vero and ~54.2% in HEK cells (**Figure 2a-**

c). To dissect out the contribution of each STT3 isoform, individual knockdowns were

also performed separately. While si*STT3-A* showed a profound inhibitory effect in HEK

cells (52% reduction), it did not result in significant virus reduction in Vero. On the

contrary, si*STT3-B* reduced the infection in Vero down to 48% but did not protect HEK

cells from infection. Reduction of *GANAB* levels further down in the pathway,

consistently resulted in less infections in both Vero and HEK cells, although the effect

was not as pronounced as seen in si*STT3* (41.4% and 42.5% respectively). Lastly,

si*MGAT1* had no effect in Vero but provided some protection to HEK cells (58.8%

infection). Knockdown levels were determined for each group and biological replicate

by immunoblotting using specific antibodies (**Figure 2d**).

a

	anti-Spike	DAPI	Merge
siNT			
siSTT3-A			
siSTT3-B			
siSTT3-A+B			
siGANAB			
siMGAT1			

b

Vero E6 — anti-Spike, Merge; HEK293^{ACE-2} — anti-Spike, Merge

Uninfected, siNT, siSTT3-A, siSTT3-B, siSTT3-A+B, siGANAB, siMGAT1

c

Vero E6

HEK293^{ACE-2}

Normalized infection (%)

siNT, siSTT3-A, siSTT3-B, si-STT3A+siSTT3B, siGANAB, siMGAT1

d

Figure 2. Genetic ablation of host *N*-glycosylation genes reduces SARS-CoV-2 infectivity.
(**a**) Representative whole-well scans of confluent Vero E6 monolayers transfected either with siNT
(negative control) or siRNAs for the knockdown of the *N*-glycosylation genes siSTT3-A, siSTT3-B,
siSTT3A+B, siGANAB and siMGAT1, and infected with SARS-CoV-2 (MOI=0.05, 24 hours). Cells
immunostained with anti-spike antibodies (green), counterstained with DAPI (purple) and merged;
scale bar 5 mm. (**b**) Representative images of either Vero E6 or HEK293^{ACE-2} siRNA cells infected
with SARS-CoV-2 (MOI=0.05 for 24 hours for Vero; MOI=0.1 for 24 hours for HEK293). Cells
immune-stained with anti-spike (gray channel) and merged (green) with DAPI counterstain (blue);
scale bars 200 μm. (**c**) Quantification of the percentage of infected (spike-positive) cells in 'a'
normalized to siNT controls of either Vero E6 or HEK293^{ACE-2} cells infected with SARS-CoV-2
(MOI=0.05, 24 hours for Vero; MOI=0.1 for 24 hours for HEK293). Bars indicate the mean values;
dots indicate individual values, error bars represent standard deviation; asterisks indicate
significance ($p<0.001$); three technical replicates, four independent experiments performed. (**d**)
Western blotting of protein lysates from Vero cells comparing siNT control cells with either
siSTT3-A, siSTT3-B, siGANAB or siMGAT1 cells probing with specific antibodies against the
knocked down protein, and with anti-GAPDH as loading control; apparent molecular weight
markers annotated on the left.

254
255
256

<u>SARS-CoV-2 *N*-glycans are essential for infection</u>
Having shown that the normal functioning of the host *N*-glycosylation machinery is
important for SARS-CoV-2 infection, we next wanted to determine whether viral *N*-
glycans alone also played a critical role. We hypothesized that glycoproteins in virions
assembled in cells with inhibited *N*-glycosylation may acquire abnormal *N*-
glycosylation profiles and consequently become less infectious. To test this
hypothesis, we used as model virions produced by Vero E6 cells treated with 5 μM
NGI-1. This treatment was previously observed to greatly reduce the proportion of
infected cells while allowing the formation of fewer infective viral particles. As
determined by plaque assay, supernatants from 5 μM NGI-1-treated cells contained
on average 392-fold less infective viral particles than untreated control cells (**Figure
3a**). However, quantification by qRT-PCR of total viral RNA (i.e., infective and non-
infective particles) in those same supernatants showed only a reduction of 82-fold in
NGI-1-treated cells compared to the control (**Figure 3b**). This difference between total
and infective particles (4.8-fold) may account for virions incapable of infecting host

cells as a result of altered and/or missing protein *N*-glycans. To validate this
hypothesis, the same supernatants were used to infect either untreated or 5 μM NGI-
1-treated Vero cells, normalizing the virus inoculum using RT-qPCR quantification
data. Despite having provided cells with virtually an identical amount of viral RNA, cells
infected with virions produced by NGI-1-treated cells still presented 4.4-fold less
infection, evidencing once more the existence of a sub-population of virions unable to
infect subsequent host cells (**Figure 3c**). As expected, NGI-1 treatment protected from
infection and made it even more pronounced when treated cells were infected with
NGI-1-derived virions. To confirm that the absence of *N*-glycans in certain viral
proteins renders viral particles non-infective, we performed *N*-glycan enzymatic
release on intact SARS-CoV-2 control virions prior to infection. Virions were purified
from cell supernatants by size exclusion and incubated with Peptide-N-Glycosidase F
(PNGase-F), which removes all types of *N*-linked oligosaccharides from glycoproteins.
Under this condition, the enzyme is only expected to cleave accessible *N*-glycans from
surface exposed viral glycoproteins (i.e., spike protein). Protein deglycosylation in
PNGase-F-treated virus lysate was confirmed by lectin blotting probing with either
concanavalin A (ConA) or wheat germ agglutinin (WGA) (**Figure 3d**), with overall lane
fluorescence intensity reductions of 57% and 70% respectively (**Figure 3e**). Some
visible protein bands (e.g., 120 kDa) also experienced a down shift in apparent
molecular weight due to the loss of *N*-glycans. After inoculation, Vero cells incubated
with PNGase-F-treated purified virions (PNGase-F +) remained uninfected, while cells
inoculated with purified virions incubated in the absence of the enzyme (PNGase-F -)
reached normal infection levels (~22%) comparable to those in cells inoculated with
purified virus only (29% infection) (**Figure 3f,g**). Control non-purified virions produced
a slightly higher average infection rate of 42%, proving that both steps of purification
and incubation with PNGase-F's reaction buffer had little impact on virion infectivity.
Supernatants of infected cells were collected to measure the amount of newly released
virions. Both total viral RNA quantification (**Figure 3h**) and plaque formation (**Figure
3i**) confirmed that cells infected with PNGase-F-treated virions did not develop a
SARS-CoV-2 infection.
302
303
304

Figure 3. Surface exposed viral *N*-glycans are essential for infection. (**a**) SARS-CoV-2 titres (pfu/mL) measured by plaque assay in supernatants of Vero E6 infected cells (MOI=0.001, 48 hours) either treated with 0.1, 5, 75 μM NGI-1 or untreated (0); asterisks indicate significance ($p<0.001$). (**b**) Total SARS-CoV-2 RNA (arbitrary units) detected in supernatants in 'a' by RT-qPCR. (**c**) Proportion of Vero E6 spike-positive cells either treated with 0, 0.1, 5 or 75 μM NGI-1 infected with virus either from untreated (DMSO) or 5 μM NGI-1-treated cell supernatant, at MOI=0.05 for 24 hours normalized using total viral RNA data. (**d**) Lectin blotting of SARS-CoV-2-infected Vero E6 supernatant (Control), purified and PNGase F-treated for 5 hours at 37°C (+PNGase F), purified untreated (- PNGase F) or purified only (Direct) lysates. Membrane probed with ConA-FITC (green), WGA-rhodamine (red) and nigrosine-stained (white) afterwards for loading control. (**e**) Fluorescence intensity quantification of entire lanes in 'd' normalized to controls. (**f**) Representative images of Vero E6 cells infected with supernatants in 'd' (MOI=0.05, 24 hours), immuno-stained with anti-spike antibodies (green) and DAPI counterstain (blue); scale bar 200μm. (**g**) Quantification of the proportion of infected cells in 'f'. (**h**) Total SARS-CoV-2 RNA in supernatants produced in 'f' quantified by RT-qPCR. (**i**) SARS-CoV-2 titres measured in supernatants produced in 'f' by plaque assay.

Discussion

N-glycosylation, the main eukaryote protein post-translational modification, plays an essential role in cell physiology. *N*-glycans can modulate protein folding, stability, antigenicity and intermolecular interactions. Many human pathogenic viruses, including SARS-CoV-2, *N*-glycosylate proteins involved in fundamental processes[25,6]. SARS-CoV-2 surface homotrimeric spike proteins, essential for cell host recognition and invasion, are decorated with up to 22 *N*-glycans per monomer. It also *N*-glycosylates the highly abundant nucleocapsid phosphoprotein (N) that envelops the viral genetic material[26]. Although there is little evidence of *N*-glycosylation occurring in other SARS-CoV-2 proteins, most are likely to be modified with *N*-glycans since many contain canonical *N*-glycosylation sequons with enough predicted potential. Previous studies have predicted that spike *N*-glycans may modulate protein folding and multimerization[8], conformational changes, binding to the ACE-2 human receptor and immunological shielding[5]. Also, it has been shown using recombinant proteins and pseudotyped viruses that mutations in certain *N*-glycosylation sites of the spike lead to reduced infectivity[10], and that treatment with kifunensine or MGAT1 knockout to prevent the formation of complex *N*-glycans reduces pseudovirus infection[11]. However, the effects of altering *N*-glycosylation in SARS-CoV-2 virions remain undetermined.

Here we investigated for the first time the essentiality of the host *N*-glycosylation pathway and SARS-CoV-2 protein *N*-glycosylation in relation to infection, and whether these could be potentially exploited to treat COVID-19. We hypothesized that interfering with the host *N*-glycosylation pathway may negatively impact the SARS-CoV-2 infection outcome because 1) newly synthesized viral proteins not properly *N*-glycosylated could misfold or be non-functional; 2) newly assembled virions with *N*-glycosylation defects could become non-infective; 3) host cell receptors (e.g., ACE-2) presenting *N*-glycosylation defects could impede successful viral invasion of cells. We found that the inhibition of key steps within the host *N*-glycosylation pathway using known inhibitors greatly reduced SARS-CoV-2 infection and replication *in vitro,* in both green monkey Vero E6 and human HEK293 kidney epithelial cells. Most inhibitors reduced the overall number of infected cells, the number of cells presenting cytopathic effect, and the viral load produced by these cells. We validated these results by knocking down the expression of genes encoding for key enzymes in the *N*-

glycosylation pathway, including some the inhibitor targets (i.e., STT3 and GANAB).
The most effective infection inhibition was achieved when altering the early *N*-
glycosylation pathway, by either directly preventing the addition of the *N*-glycan
precursor $Glc_3Man_9GlcNAc_2$ to the nascent polypeptide (with NGI-1[27] and STT3
RNAi), or indirectly by blocking the formation of GlcNAc-dolichol phosphate using
tunicamycin. Interestingly, *STT3* isoform-specific knock down resulted in opposed
phenotypes in Vero and HEK cells. The knock down of *STT3-A*, responsible for the
co-translational glycosylation of most acceptor sites, only reduced viral infection in
human HEK cells but showed no impact in Vero cells despite an effective knock down
being achieved. STT3-B however, the isoform glycosylating those sites omitted by
STT3-A or close to the polypeptide ends in a post-translational manner, showed the
most pronounced effect in Vero cells that we could not replicate in HEK cells, likely
due to inefficient knock down. A similar phenotype for siSTT3-B was reported in Lassa
virus-infected cells[28]. Double knock down of *STT3-A* and *STT3-B* always resulted in
protection rates against infection similar to the effect caused by the most effective
isoform-specific knock down in cell type respectively. Moreover, we found that the
further downstream the *N*-glycosylation pathway is halted, the lesser the impact on
virus infectivity. Inhibition of the ER α-glucoside I, which removes the terminal α-
glucose residue from protein *N*-glycans, using either inhibitors or knock down of its
catalytic subunit GANAB, also resulted in reduced infectivity although not as high as
in early pathway inhibition. Not all α-glucosidase I inhibitors tested here were found to
be equally effective. While treatment with the iminosugars Miglustat, Celgosivir and
NN-DNJ led to a strong reduction of infection, CPE and viral load, others such as
Miglitol, DNJ, Acarbose or Swainsonine barely reduced the overall infection while
slightly protecting cells against CPE. Lastly, alterations later in the pathway either
using DMM to inhibit the Golgi's α-mannosidase I, or knocking down MGAT1 to
prevent the formation of hybrid and complex glycan types, resulted in no significant
infectivity decrease in Vero cells, and in a mild infection reduction in HEK cells.
Variations in the *N*-glycosylation profiles of both host cell and generated virions may
account for these differences observed between cell types.
384
The most potent inhibitors here used greatly reduced the overall number of infected
cells regardless the treatment timing, hours before, during or after infection. Unlike

untreated samples which displayed most infected cells in clusters, probably formed by
the spread of the virus from an infected cell to neighbouring ones, inhibitor-treated
samples showed discrete individual infected cells instead. These did also present
higher amounts of trapped spike protein, suggesting viral replication occurs but virions
cannot either assemble or egress properly for neighbour cells to become infected.
Experiments involving variable MOIs supported this hypothesis as the number of
infected cells in inhibitor-treated samples was proportional to the initial viral inoculum,
while untreated cells rapidly became fully infected. In addition, we hypothesized that
cells with inhibited *N*-glycosylation may be producing virions with glycosylation defects
that render them non-infective to further cells. We validated this hypothesis by
analyzing the supernatant of infected cells treated with a sub-inhibitory dose of NGI-1
and comparing the number of infective virions with the total amount of viral genetic
material. We found that indeed, these inhibitor-treated cells were producing a
significant population of virions uncapable of infecting further host cells, unlike
untreated cells which produced infective virions only. Further validation involved
enzymatic deglycosylation of intact virions, which rendered completely non-infective
viruses, evidencing the essential role of, at least, surface SARS-CoV-2 *N*-glycans for
infection.

It is expected that the blockage of the host *N*-glycosylation pathway and the
consequent removal or alteration of virus *N*-glycan structures have such a profound
impact in virus infectivity. Our results confirmed the predicted important roles of *N*-
glycans in relation to viral glycoprotein maturation, virion assembly and host cell
recognition and invasion. In addition, it likely has a remarkable impact because SARS-
CoV-2 has been shown to require upregulating genes along the *N*-glycosylation
pathway[29], especially those related with OST and glucosidase activities, likely to
ensure enough *N*-glycosylation capacity for the increased viral glycoprotein
production. The evidence generated in this study opens the door to further research
on developing ways to exploit the *N*-glycosylation inhibition and SARS-CoV-2 *N*-
glycans to treat COVID-19. Some of the inhibitors used here (i.e., Miglustat, Miglitol
and Acarbose) are FDA-approved drugs widely used to treat different unrelated
human diseases and could be considered for repurposing if found effective in trials,
alone or in combination. Celgosivir is in an advanced trial phase and could also be
considered. Also, drug discovery approaches may focus on identifying better and safer

OST inhibitors, especially against STT3-B, as a way to target viral *N*-glycosylation

more specifically. On the other hand, we have demonstrated that SARS-CoV-2

surface *N*-glycans are essential for infection. New potential treatments may consist of

blocking such glycans using antibodies or lectins to reduce and neutralize the spread

of the infection[30]. Concerns are currently arising about new SARS-CoV-2 variants that

could potentially escape treatments under development or vaccine-derived antibody

neutralization. Approaches targeting essential *N*-glycans may offer an advantage over

protein-based ones as *N*-glycosylation is a more conserved, independent protein

modification less favourable to mutation. In summary, the study of *N*-glycosylation in

relation to viral diseases may offer new and exciting opportunities to fight, not only

SARS-CoV-2, but further virus infections.

Methods

Cell culture and SARS-CoV-2 infection

Vero E6 cells (C1008; green African monkey kidney cells) were obtained from Public Health England and cultured in DMEM medium (Lonza) supplemented with 10% fetal bovine serum (FBS) (Gibco) and 50 μg/mL gentamicin (Sigma) at 37°C and 5% CO_2. HEK293[ACE-2] cells (lentiviral expression of human angiotensin-converting enzyme 2, source) were cultured in DMEM supplemented with 10% FBS, 1mM sodium pyruvate (Gibco) and 2 μg/mL puromycin (Invivogen) at 37°C and 5% CO_2. Cells were regularly passaged (passage 2 to 30) by trypsinization when >80% confluence was reached. SARS-CoV-2 isolate SARS-CoV-2/human/Liverpool/REMRQ0001/2020, isolated from naso-pharyngeal swab from a patient, was passaged four times in Vero E6 cells[31]. Experimental cells were infected by incubation with SARS-CoV-2 (passage 4 in Vero E6) containing 2×10^7 pfu/mL in DMEM 2% FBS for 30 minutes at 37°C and 5% CO_2 with variable MOIs (see below).

N-glycosylation inhibitor assays

Cells were pre-treated for 2 to 16 hours before infection with either DMSO or several N-glycosylation inhibitors (from DMSO stock solutions, DMSO final concentration never exceeding 2%) in DMEM 2% FBS. Inhibitors used: Tunicamycin (Sigma), NGI-1 (Sigma), DNJ (Cambridge Bioscience), Miglitol (Stratech Scientific), Miglustat (Cambridge Bioscience), Acarbose (Fisher Scientific), NN-DNJ (Cambridge Bioscience), Celgosivir (Sigma), DMM (Tebu-Bio) and Swainsonine (Cambridge Bioscience). Cells were then infected by incubation with SARS-CoV-2 P4 supernatant with the corresponding inhibitor for 30 minutes at 37°C and 5% CO_2, at either MOI=0.001 (Vero E6 for plaque assay), MOI=0.05 or MOI=0.1 (Vero E6 and HEK293[ACE-2] for immunofluorescence assay, respectively). Cells were washed and left in DMEM 2% FBS and N-glycosylation inhibitor at 37°C and 5% CO_2 for 48 hours (Vero E6 for plaque assay) or 24 hours (Vero E6 and HEK293[ACE-2] for immunofluorescence). For toxicity assay, cells in 96-well plates were treated with variable concentrations of inhibitors in DMEM 2% FBS in triplicate and after 24 hours, AlamarBlue 10X solution (Invitrogen) was added and incubated for 4 hours at 37°C before measuring well fluorescence intensity at 590 nm using a FLUOstar Omega plate reader (BMG Labtech)

siRNA knockdowns

A total of 2×10^5 Vero E6 or HEK293^{ACE-2} cells were seeded in 6-well plates 24 hours

before transfection with one of the following siRNAs pools (Qiagen SMARTpool):

siSTT3A (catalogue number GS3703), siSTT3B (GS201595), siGANAB (catalogue

number: GS23193) or siMGAT1 (GS84061). Double knockdown of STT3A and STT3B

was performed by transfecting both siSTT3A and siSTT3B pools. Non-targeting

control (siNT) was obtained from Dharmacon. 4µL of siRNA were combined with 6µL

Lipofectamine RNAiMax (Invitrogen) in 200µL Opti-MEM (Gibco) and incubated for 20

minutes at RT before being added dropwise to the cells. 4 hours after transfection,

cells were flushed out with fresh medium. For siGANAB and siMGAT1, 2 hits of siRNA

over a 120-hour incubation period were performed. For siSTT3A, siSTT3B and

siSTT3A+siSTT3B, 3 hits of siRNA over a 168-hour period were needed. The day

before optimum knockdown (96 hours for 2 hits, 144 hours for 3 hits), cells were

trypsinized and 2×10^4 cells for each condition were seeded onto 96-well optical plates

(Greiner) in quadruplicate and infected the following day. Infections were carried out

by incubation with SARS-CoV-2 P4 supernatant at either MOI=0.05 (Vero E6) or

MOI=0.1 (HEK293^{ACE-2}) for 30 minutes at 37°C, 5% CO_2, washing and incubation in

DMEM 2% FBS 37°C, 5% CO_2 for 24 hours.

Plaque assay

Vero E6 were cultured in DMEM 10% FBS in 24-well plates and allowed to reach

confluency overnight. Cells were incubated with 0.1 mL serially diluted supernatant

containing SARS-CoV-2 for 30 minutes at 37°C and 5% CO_2 before being discarded.

Cells were then washed in DMEM 2% FBS and left in 1.5mL overlay (DMEM 2%FBS,

1.2% Avicel) at 37°C and 5% CO_2 for 72 hours. Cells were fixed in formaldehyde for

30 minutes, washed in water and stained with 0.1% crystal violet solution (in 20%

ethanol) for manual plaque counting.

PNGase-F glycan enzymatic release

0.5mL SARS-CoV-2 P4 supernatant containing 2×10^7 pfu/mL was purified using an

Amicon Ultra (MWCO 100kDa) column (Merck) by centrifugation and washing in PBS

at 2,000xg. 18uL purified supernatant (containing 9×10^5 pfu) were combined with

recombinant glycerol-free PNGase-F and Glycobuffer 2 (New England Biolabs)

following the manufacturer's protocol for non-denaturing digestion, and incubated at

37°C for 5 hours. Parallel controls included incubation of purified virions without

PNGase-F and without both PNGase-F and buffer. After incubation, samples were

either directly used for infection or combined with RIPA buffer supplemented with

protease inhibitors for blotting.

<u>RT-qPCR SARS-CoV-2 quantification</u>

Total RNA was extracted from supernatants of SARS-CoV-2-infected Vero E6 cells

using the RNeasy kit (Qiagen) following the manufacturer's protocol. 4 μL purified RNA

were subsequently used for RT-qPCR reaction using the Luna Universal One-Step

RT-qPCR kit (New England Biolabs) and 0.4μM of each primer: CoV2-N-F

(CACATTGGCACCCGCAATC) and CoV2-N-R (GAGGAACGAGAAGAGGCTTGAC).

Reactions were set and analyzed in an Agilent Mx3005P machine as follows: 55°C for

10 minutes, 95°C for 1 minute, 40 cycles of 95°C for 10 seconds followed by 60°C for

30 seconds (SYBR green measure), with final melting curve from 55°C to 95°C.

Samples were quantified using the ΔCt method.

<u>Immunofluorescence assay</u>

Cells cultured in optical plates were fixed in 4% formaldehyde (Pierce) for 30 minutes,

washed in PBS and permeabilized in 0.5% saponin for 10 minutes. Cells were then

blocked in 1% bovine serum albumin (BSA) for 45 minutes and further incubated in

blocking solution containing anti-SARS-CoV-2 spike protein (S) subunit 1 (Abcam)

1:1000 dilution for 1 hour at 37°C. Blocking solution was then incubated for 1 hour at

RT containing DAPI 1μg/mL and anti-mouse IgG Alexa Fluor 488 Plus-conjugated

(Thermofisher). Cells were washed and left in PBS for imaging. Imaging was carried

out using a Zeiss LSM800 confocal laser scanning microscope by tile-scanning entire

wells or plates conserving identical settings between comparable samples. Images

were analyzed using Fiji (ImageJ) by thresholding to select positive signals only and

measuring area coverage of DAPI channel (total number of cells) and mean intensity

of the Alexa Fluor channel (infected cells).

<u>Western blotting</u>

Cell lysates were obtained at either 120-hours (2 hits) or 168-hours (3 hits). Cells were

washed in cold PBS 3 times and incubated in RIPA buffer with 1% phosphatase and

protease inhibitor cocktail (ThermoScientific) for 10 minutes on ice before scrapping,

centrifuging at 15,000 rpm for 10 minutes at 4°C and collecting the supernatants.

Protein concentration was determined using Precision Red (Cytoskeleton) and 60μg

of protein were separated using 4-15% SDS-PAGE gels (BioRad) and transferred onto

nitrocellulose membrane (GE) using a BioRad Trans-Blot Turbo Transfer system.

Membranes were blocked in either 5% milk or 2% BSA in Tris-buffered saline, 0.1%

Tween-20 (TBS-T) for 30 minutes at RT and incubated with their corresponding

primary antibodies at 4°C overnight. Primary antibodies used: anti-STT3A

(HPA030735, Cambridge Bioscience), anti-STT3B (A15574, Universal Biologicals),

anti-GANAB (A13851, Universal Biologicals), anti-MGAT1 (CSB-PA013773ESR1Hu,

2B Scientific) and anti-GAPDH (10494-1-AP, ProteinTech). Blots were then washed 3

times under agitation for 5 minutes in TBS-T before incubation with horseradish

peroxidase (HRP)-conjugated anti-rabbit IgG secondary antibody (Cell Signalling) for

1 hour at RT. After washing, blots were visualized on a ChemiDoc™ Touch Gel

Imaging System (BioRad) using enhanced chemiluminescence substrate (BioRad).

GAPDH was utilized as loading control for all blots.

Lectin blotting

Cell lysates of siRNA and *N*-glycosylation inhibitor-treated cells were obtained as

described above. 60μg protein was separated in 12.5% SDS-PAGE gels, transferred

onto nitrocellulose membranes and blocked in 1% BSA PBS overnight at 4°C. Blots

were incubated in 1% BSA containing both concanavalin A (ConA) FITC-conjugated

(1 μg/mL) and wheat germ agglutinin (WGA) rhodamine-conjugated (1μg/mL) (Vector

Labs) for 30 minutes at RT. After extensive washing in PBS 0.05% Tween-20,

membranes were imaged using a Thermofisher iBright FL1500 system.

Financial support

This work was supported in part by the Liverpool School of Tropical Medicine Director's Catalyst Fund awarded to ACS and the UKRI-BBSRC COVID rolling fund (BB/V017772/1) awarded to AAS, ACS, EIP and GLH. GLH was supported by the BBSRC (BB/T001240/1 and V011278/1), a Royal Society Wolfson Fellowship (RSWF\R1\180013), the NIH (R21AI138074), UKRI (20197), and the NIHR (NIHR2000907). GLH is affiliated to the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Emerging and Zoonotic Infections at University of Liverpool in partnership with Public Health England (PHE), in collaboration with Liverpool School of Tropical Medicine and the University of Oxford. GLH is based at LSTM. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, the Department of Health or Public Health England. Confocal imaging facilities at LSTM were funded by a Wellcome Trust Multi-User Equipment Grant (104936/Z/14/Z).

Author Contributions

ACS and AAS conceptualized and designed the study. ACS, ARR, EH, TZ performed the experiments. ACS, ARR, EH, EIP, GLH, TZ, AAS analyzed, curated and validated data. ACS wrote the paper with input from all the authors.

Competing interests

The authors declare no competing interests.

References

1. Lu, R. *et al.* Genomic characterisation and epidemiology of 2019 novel
coronavirus: implications for virus origins and receptor binding. *Lancet* **395,**
565–574 (2020).
2. Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2
and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181,** 271–280.e8
(2020).
3. Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2
Spike Glycoprotein. *Cell* (2020). doi:10.1016/j.cell.2020.02.058
4. Shajahan, A. *et al.* Comprehensive characterization of N- and O- glycosylation
of SARS-CoV-2 human receptor angiotensin converting enzyme 2.
Glycobiology (2020). doi:10.1093/glycob/cwaa101
5. Watanabe, Y., Allen, J. D., Wrapp, D., McLellan, J. S. & Crispin, M. Site-
specific glycan analysis of the SARS-CoV-2 spike. *Science* eabb9983 (2020).
6. Shajahan, A., Supekar, N. T., Gleinich, A. S. & Azadi, P. Deducing the N- and
O-glycosylation profile of the spike protein of novel coronavirus SARS-CoV-2.
Glycobiology **30,** 981–988 (2020).
7. Yao, H. *et al.* Molecular Architecture of the SARS-CoV-2 Virus. *Cell* (2020).
8. Casalino, L. *et al.* Beyond Shielding: The Roles of Glycans in the SARS-CoV-2
Spike Protein. *ACS Cent Sci* **6,** 1722–1734 (2020).
9. Dieterle, M. E. *et al.* A replication-competent vesicular stomatitis virus for
studies of SARS-CoV-2 spike-mediated cell entry and its inhibition. *Cell Host*
Microbe (2020). doi:10.1016/j.chom.2020.06.020
10. Li, Q. *et al.* The Impact of Mutations in SARS-CoV-2 Spike on Viral Infectivity
and Antigenicity. *Cell* (2020). doi:10.1016/j.cell.2020.07.012
11. Yang, Q. *et al.* Inhibition of SARS-CoV-2 viral entry upon blocking N- and O-
glycan elaboration. *Elife* **9,** 450 (2020).
12. Thao, T. T. N. *et al.* Rapid reconstruction of SARS-CoV-2 using a synthetic
genomics platform. *Nature* 1–27 (2020). doi:10.1038/s41586-020-2294-9
13. Xie, X. *et al.* An Infectious cDNA Clone of SARS-CoV-2. *Cell Host Microbe* **27,**
841–848.e3 (2020).
14. Gavel, Y. & Heijne, von, G. Sequence differences between glycosylated and
non-glycosylated Asn-X-Thr/Ser acceptor sites: implications for protein
engineering. *Protein Eng* **3,** 433–442 (1990).
15. Helenius, A. & Aebi, M. Roles of N-linked glycans in the endoplasmic
reticulum. *Annu Rev Biochem* **73,** 1019–1049 (2004).
16. Herscovics, A. Importance of glycosidases in mammalian glycoprotein
biosynthesis. *Biochim Biophys Acta* **1473,** 96–107 (1999).
17. Schachter, H. *The joys of HexNAc. The synthesis and function of N- and O-*
glycan branches. Glycoconjugate journal **17,** 465–483 (Kluwer Academic
Publishers-Plenum Publishers, 2000).
18. Cox, T. M. *et al.* The role of the iminosugar N-butyldeoxynojirimycin (miglustat)
in the management of type I (non-neuronopathic) Gaucher disease: a position
statement. *J Inherit Metab Dis* **26,** 513–526 (2003).
19. Scott, L. J. & Spencer, C. M. Miglitol: a review of its therapeutic potential in
type 2 diabetes mellitus. *Drugs* **59,** 521–549 (2000).
20. Campbell, L. K., White, J. R. & Campbell, R. K. Acarbose: its role in the
treatment of diabetes mellitus. *Ann Pharmacother* **30,** 1255–1262 (1996).
21. Durantel, D. Celgosivir, an alpha-glucosidase I inhibitor for the potential
treatment of HCV infection. *Curr Opin Investig Drugs* **10,** 860–870 (2009).

22. Marcello, A. *Repurposing of Miglustat to inhibit the coronavirus Severe*
Acquired Respiratory Syndrome SARS-CoV-2. bioRxiv 1–17 (2020).
doi:10.1101/2020.05.18.101691
23. Williams, S. J. & Goddard-Borger, E. D. α-glucosidase inhibitors as host-
directed antiviral agents with potential for the treatment of COVID-19. *Biochem*
Soc Trans (2020).
24. Clarke, E. C., Nofchissey, R. A., Ye, C. & Bradfute, S. B. The iminosugars
celgosivir, castanospermine and UV-4 inhibit SARS-CoV-2 replication.
Glycobiology (2020). doi:10.1093/glycob/cwaa091
25. Watanabe, Y. *et al.* Vulnerabilities in coronavirus glycan shields despite
extensive glycosylation. *Nat Commun* **11,** 2688–10 (2020).
26. Supekar, N. T. *et al.* SARS-CoV-2 Nucleocapsid protein is decorated with
multiple N- and O-glycans. *bioRxiv* 2020.08.26.269043 (2020).
27. Puschnik, A. S. *et al.* A Small-Molecule Oligosaccharyltransferase Inhibitor
with Pan-flaviviral Activity. *Cell Rep* **21,** 3032–3039 (2017).
28. Zhu, S. *et al.* Comprehensive Interactome Analysis Reveals that STT3B Is
Required for N-Glycosylation of Lassa Virus Glycoprotein. *J. Virol.* **93,**
e01443–19 (2019).
29. Oommen, A., Cunningham, S. & Joshi, L. Transcriptomic Analysis of
Respiratory Tissue and Cell Line Models to Examine Glycosylation Machinery
during SARS-CoV-2 Infection. *Viruses* **13,** (2021).
30. da Silva, L. C. N. *et al.* Exploring lectin-glycan interactions to combat COVID-
19: lessons acquired from other enveloped viruses. *Glycobiology* (2020).
doi:10.1093/glycob/cwaa099
31. Patterson, E. I. *et al.* Methods of Inactivation of SARS-CoV-2 for Downstream
Biological Assays. *J Infect Dis* **222,** 1462–1467 (2020).
675