Functional ultrasound imaging of the spreading activity following optogenetic stimulation of the rat visual cortex

Authors: M. Provansal[1], G. Labernede[1], C. Joffrois[1], A. Rizkallah[1], R. Goulet[1], M. Valet[1], W. Deschamps[1], U. Ferrari[1], A. Chaffiol[1], D. Dalkara[1], J.A. Sahel[1,2,3], M. Tanter[4], S. Picaud[1], G. Gauvain*[1], F. Arcizet*[1].

- * Equal contribution

[1]Sorbonne Université, INSERM, CNRS, Institut de la Vision, F-75012 Paris, France; [2]Department of Ophthalmology, The University of Pittsburgh School of Medicine, Pittsburgh, PA 15213, United States; [3]Department of Ophthalmology and Vitreo-Retinal Diseases, Fondation Ophtalmologique Rothschild, F-75012 Paris, France; and [4]Physics for Medicine Paris, INSERM, CNRS, Ecole Supérieure de Physique et de Chimie Industrielles (ESPCI Paris), Paris Sciences et Lettres (PSL) Research University, 75012 Paris, France.

Optogenetic stimulation of the primary visual cortex (V1) is a promising therapy for sight restoration, but it remains unclear what total cerebral volume is activated after surface stimulation. In this study, we expressed the red-shifted opsin ChrimsonR in excitatory neurons within V1 in rats, and used the fine spatial resolution provided by functional ultrasound imaging (fUS) over the whole depth of the brain to investigate the brain response to focal surface stimulation. We observed optogenetic activation of a high proportion of the volume of V1. Extracellular recordings confirmed the neuronal origin of this activation. Moreover, neuronal responses were even located in deep layers under conditions of low irradiance, spreading to the LGN and V2, consistent with a normal visual information process. This study paves the way for the use of optogenetics for cortical therapies, and highlights the value of coupling fUS with optogenetics.

Introduction

Optogenetics has revolutionized investigation of the central nervous system[1], providing hope

for the treatment of a number of conditions, including deafness[2,3] or vision loss[4]. Optogenetic

therapy is already widely applied to retinal cells to restore vision for *in vivo* light application[5–

11], and is currently being assessed in two different clinical trials[12,13]. However, different

approaches, targeting cells other than those of the retina, are required for diseases causing

degeneration of the optic nerve (e.g. glaucoma) and for advanced retinal degeneration (late

AMD). For such conditions, direct stimulation of the primary visual cortex (V1) is a promising

alternative strategy. Indeed, high performance rates have been reported for the detection of

forms, with great accuracy, by blind human patients with cortical implants[14], and cortical

electrical prostheses have been shown to elicit visual percepts and to alter visual behavior in

nonhuman primates (NHP)[15]. Successes have been achieved with implantable devices, but

this approach nevertheless has a number of serious drawbacks: invasive surgery, signal

degradation over time, and a lack of cell-type specificity. In this respect, optogenetic therapy

stimulating V1 at the cortical surface could potentially afford similar benefits, but with less

invasive administration, stable expression over time and precise genetic targeting of the

appropriate cell population. Two of the key aspects of this strategy are the activation of

cortical layer IV neurons, as these are the first cells to receive visual information from the

visual thalamus[16] and the propagation of activity to other visual structures, which would favor

the generation of visual percepts.

Layer IV is located deep in the cortex (>1 mm in NHP). It is therefore a major challenge to

read and write neuronal activity, to demonstrate the efficacy of stimulation. Red-shifted

opsins are very good tools for neuronal stimulation, as they make it possible to use lower

light power to activate deeper neurons than blue-sensitive opsins, whilst also making it

possible to use higher light intensity safely[17–19]. Electrophysiological recordings can report

neuronal activity with unmatched spatiotemporal resolution, but over a very small spatial

area[20]. Conversely, techniques such as optical imaging and fMRI have been coupled with

optogenetics to report activity throughout the brain, but at the expense of a loss of both

spatial and temporal resolution[21,22]. Like fMRI, ultrafast functional ultrasound imaging (fUS)

can provide brain-sized maps of neurovascular activity changes, with a high spatiotemporal

resolution (100 μm x 100 μm, 1Hz) even in deep structures (up to 1.5 cm)[23]. This technique

has been used to investigate sensory processing in anesthetized[24], awake[24–26] and asleep[27]

rodents, and in behaving primates[28,29]. fUS imaging and electrophysiological recordings can,

therefore, be used to describe the dynamics of local neuronal activity accurately whilst

scanning neurovascular activity over the entire brain. We used these two techniques to

determine whether the optogenetic stimulation of V1 at the cortical surface could induce

neuronal activity even in deep cortical layers, and initiate the propagation of information to

other visual structures.

We show here that red light stimulation at the cortical surface can activate visual neurons

localized in deep cortical layers without triggering a thermal hemodynamic response and

toxicity. We were also able to follow the propagation of this information in other visual brain

structures (i.e. LGN and V2). More generally, this work shows that fUS imaging has the

potential to provide a clear, brain-wide mesoscopic view of the neuronal activities resulting

from optogenetic stimulation.

Results

fUS imaging of V1 optogenetic activation in rats. We used the optogenetic actuator

ChrimsonR for visual restoration in the cortex, because of its red-shifted opsin properties[17]

and because it was already being used for visual restoration in the primate retina[11] and had

given promising results in clinical trials[13]. We maximized the optogenetic activation of V1 by

using the CaMKII promoter to ensure expression limited to the excitatory neurons of V1.

Indeed, a ubiquitous promoter might lead to the silencing of pyramidal neurons through the

recruitment of inhibitory neurons. ChrimsonR was fused to the fluorescent reporter tdTomato

to facilitate the visualization of transfected areas. Following preliminary screening, we used

the AAV9-7m8 mutated viral capsid to express ChrimsonR in the V1 neurons of Long-Evans

rats (Fig. 1a). The mean rate of neuronal transfection was 5.5% over all cortical layers (Fig.

S1). ChrimsonR expression was not restricted to the soma, but spread to the axons and

dendrites (Fig. 1a). We assessed the efficacy of the optogenetic stimulation of V1 cortical

neurons, by using fUS imaging to measure brain activity in a large proportion of the brain:

from AP -3.5 mm to AP -8 mm, the zone in which most of the early visual system areas are

located. Activity in the cortical layers was assessed following either direct stimulation of the

contralateral eye with a white LED (58 mW.cm^{-2}), or stimulation at 595 nm delivered with an

optic fiber placed at the surface of the transfected or non-transfected V1 areas (7 mW, ~140

mW.mm^{-2} at the brain surface) (Fig. 1b). We chose to use durations and magnitudes of

parameters similar to those previously used[24,30] for stimulation of the eye or cortex (2 s at 4

Hz or 20 Hz for stimulation of the eye and cortex, respectively, separated by a 13 s period of

darkness. This cycle was performed 20 times). For eye stimulation (Fig. 1b, left), we first

imaged V1 with a single imaging plane at AP -7.5 mm and constructed an activation map

including only pixels displaying significant CBV (cerebral blood volume) responses (p<0.05

with Bonferroni-Holm correction, Wilcoxon signed-rank test, one-tailed, relative to baseline

activity). We detected strong activation in both the ipsilateral and contralateral superior

colliculi (SC) (ipsilateral, n=121/488 activated pixels; contralateral, n=134/421 activated

pixels), but almost no activation in the ipsilateral and contralateral V1 areas (ipsilateral,

n=8/634 activated pixels; contralateral, n=9/370 activated pixels). The lack of response in

both V1 areas and the strong signal in both SC may reflect the retinotectal nature of most

rodent retinal outputs[16] or an effect of anesthesia[31]. An increase in CBV was already clearly

visible on single-cycle responses (gray dashed lines), as illustrated for the significant pixel

(#14-92) in the ipsilateral V1 area (insert). Following direct stimulation of the contralateral

eye, the mean response (black curve) peaked 2 seconds after the two-second stimulation

represented by the patch in gray (mean: 19.8 ± 18.3%). For cortical stimulation (Fig. 1b,

center), we observed a broad activation of the ipsilateral V1 area (n=310/634 activated

pixels). The activation spread out of the V1 area at each border (medial and ventral) with V1

projections onto other visual areas. Single-cycle responses of the same example pixel showed larger, less variable increases in CBV variation than for stimulation of the contralateral eye (optogenetic, mean: 65.8 ± 32.6%, $p<0.0001$, Mann-Whitney). A previous study showed that blue light delivery to the brain could itself generate non-specific changes in CBV [30]. We therefore performed a control stimulation by locating the optic fiber at the surface of the non-injected V1 area, which did not express ChrimsonR-tdT (Fig. 1b, right). We used the same light stimulation parameters for this control as before. We detected no significant CBV responses in the non-injected V1 area under such optogenetic stimulation conditions.

We further characterized the V1 activation volume generated by contralateral eye stimulation or by stimulation of the transfected V1 area, by imaging all the planes containing V1 (from AP -6 to -8 mm, Fig. 1c-d). As shown in Fig. 1b, direct eye stimulation induced CBV responses mostly in the SC areas, but very little activation was observed in the two V1 areas (mean: 0.05 ± 0.12%). Strikingly for the same animal shown (as shown in Fig. 1c), direct optogenetic stimulation of the injected V1 area generated significantly stronger CBV responses, with a mean active volume of 37% (range: 0.5 to 75.9%). The percentage mean active injected V1 area, over all animals, was higher for optogenetic cortical stimulation than for direct eye stimulation (optogenetic, mean: 16.2 ± 17.8 %, Wilcoxon signed-rank test, one-tailed, $p=0.001$, Fig. S1). Neurovascular activity and ChrimsonR expression were distributed similarly along the AP axis and their amplitudes were correlated (Fig. 1d, Fig. S1). Again, our control experiments demonstrated that direct optogenetic stimulation of a non-transfected cortical area resulted in no activation. The findings for these control conditions indicate that the parameters we used for optogenetic stimulation at the cortical surface did not generate CBV variations in areas into which the virus was not injected (mean: 0 ± 0%). These observations indicate that the optogenetic light stimulation used here does not trigger a vascular response detectable by fUS imaging. Consequently, with fUS imaging, we were

able to visualize the entire volume of an optogenetically evoked response resulting from focal

stimulation within the primary visual cortex.

We then sought to confirm that the observed changes in blood volume following optogenetic

stimulation at the surface of V1 were indeed due to an increase in neuronal activity, and not

to indirect factors, such as heating. We therefore performed electrophysiological recordings

of V1 during the stimulation of the contralateral eye with white light (200 ms, 1 Hz, 100

cycles, 58 mW.cm^{-2}) or of the injected V1 area with light at 595 nm (200 ms, 1 Hz, 100

cycles, 140 mW.mm^{-2}). We used the Spyking Circus algorithm[32] to sort the multi-unit

recordings, to obtain single-cell responses. In total, we recorded a population of 171 neurons

from nine animals expressing ChrimsonR in V1. These neurons displayed several distinctive

patterns of activity under both direct eye and optogenetic stimulation conditions (Fig. 1e). We

plotted the spike density function (SDF) of four V1 neurons for both direct eye stimulation

(black lines) and optogenetic conditions (red lines), to highlight the diversity of these activity

patterns. Based on the profiles of visual and optogenetic responses, we classified neurons

into four different groups: neurons responding to both visual and optogenetic stimulation (V +

O neurons, n=13, 7.6%, Wilcoxon signed-rank test, two-tailed, p<0.01, between baseline: [-

100 0] ms, and stimulus presentation window: [0 200] ms), to visual stimulation only (V

neurons, n=20, 11.7%), to optogenetic stimulation only (O neurons, n=81, 47.4%), and non-

responsive neurons ('None' neurons, n=57, 33.3%). Most of the neurons responding to visual

stimulation displayed phasic responses, with an ON response occurring after the start of

stimulation followed, in some cases, by an OFF response. By contrast, the neurons

responding to optogenetic stimulation displayed a unique ON response. We characterized

the neuronal activation further, by comparing the onset latencies and durations of the V1

responses for both direct eye and optogenetic stimulation conditions, for V, O and V+O

neurons (n=109, Fig. 1f). As observed for the representative neurons shown in Fig. 2a, the

onset latencies of V1 responses were significantly shorter after optogenetic stimulation

(mean: 1.76 ± 3.14 ms, n=76 units) than after stimulation of the contralateral eye (mean:

41.24 ± 16.61 ms, n=33 units, Mann-Whitney, p<0.0001). These very short response latencies for optogenetic stimulation are consistent with the direct activation of transfected neuronal cell bodies, bypassing all retinal synapses, by contrast to natural visual signal transmission. We also analyzed the duration of visual/optogenetic responses, to determine whether V1 neurons presented transient or sustained activity, according to the type of stimulation. For direct eye stimulation, the duration of neuronal responses was tuned on a single ensemble, with a mean duration of activation of 28.09 ± 17.05 ms (n=33). By contrast, in optogenetic conditions, two subgroups emerged: neurons displaying transient and sustained responses. Based on these results, we defined two subsets of neurons: transient (durations < 51 ms, n=33) and sustained (durations > 197 ms, n=22) neurons. The remaining neurons (n=17) had intermediate response durations. The transient responses may result from inhibitory cortical feedback from interneurons or a difference in voltage-gated channel properties between neuronal subtypes leading to the silencing of these neurons. We found no significant difference in the depth distribution of transient and sustained neurons (Fig. 1g), revealing an absence of correlation between neuronal response patterns and potential decreases in stimulus intensity with tissue depth; transient and sustained neurons were recorded within the same cortical layers (L2/3 to L6), suggesting a direct activation of cortical neurons by optogenetic stimulation at the cortical surface. We then assessed the specificity of optogenetic activation, by performing electrophysiological recordings on naive animals. In the animals in which no injection was performed (Fig. S1, n=3 animals), almost all the recorded neurons (n=187/188) displayed a total absence of response to optogenetic stimulation; the only responsive neuron had a very low response rate (3.89 Hz) relative to its baseline activity (1.71 Hz, p = 0.0014). In this experiment, most neurons displayed a visual response when the contralateral eye was stimulated (n=80/118), whereas another group of neurons (n=37/118) did not respond to either visual or optogenetic stimulation. Thus, by recording single-cell activities in transfected areas of V1, we were able to demonstrate that optogenetic light stimulation at the surface of the cortex triggered both an increase in cerebral blood volume, as shown by fUS imaging, and direct neuronal activation. We can

191 therefore conclude that the fUS variations we observed reflected the optogenetic activation of

192 V1 neurons. As in fUS imaging, we found that a larger number of neurons responded to

193 optogenetic stimulation than to visual stimulation (Fig. 1e), consistent with the broader

194 activation of areas in response to optogenetic stimulation than following direct contralateral

195 eye stimulation and fUS imaging.

Neuronal and neurovascular optogenetic sensitivity

As both single-cell recording and fUS imaging can be used to monitor optogenetic neuronal

activation, we decided to assess the sensitivities of these two approaches. We first imaged

the same single V1 plane (AP -7.5 mm) by fUS while optogenetically stimulating the surface

of the ChrimsonR-expressing V1 area with various irradiances (from 1.2 to 106 mW.mm^{-2}).

The size of the active volume within the injected V1 area increased with irradiance, with a

major increase between 36 mW.mm^{-2} and 70 mW.mm^{-2} and a slight decrease at 106

mW.mm^{-2} (Fig. 2a). One also found that the contralateral SC was slightly activated at

irradiances above 1.2 mW.mm^{-2}. For each irradiance, we then calculated the mean CBV

variation over all significant pixels in the V1 area and over all trials (Fig. 2b, left panel). For

each animal, we normalized these values against those obtained for the highest irradiance

(106 mW.mm$^{-2)}$. Normalized CBV variation peaked 4 s after the start of stimulation. Peak

CBV values increased as a function of irradiance. No CBV variation was recorded for the

lowest irradiance (1.2 mW.mm^{-2}). Difference in peak CBV values relative to that for the

lowest irradiance tested started to become significant from 6 mW.mm^{-2} onwards (1.2

mW.mm^{-2}, mean: 0 ± 0; 6 mW.mm^{-2}, mean: 0.36 ± 0.34, Wilcoxon signed-rank test, one-

tailed, $p<0.05$). For each irradiance, we then calculated the percentage normalized activated

ipsilateral V1 volume (Fig. 2b, right panel) in all animals (n=9). We again observed an

increase in the normalized active volume of V1 with irradiance. Difference with respect the

value obtained at the lowest irradiance became significant from 6 mW.mm^{-2} onwards (1.2

mW.mm^{-2}, mean: 0 ± 0%; 6 mW.mm^{-2}, mean: 22.3 ± 39.0%, Wilcoxon signed-rank test, one-

tailed, $p<0.05$). We then investigated whether the variation of CBV responses observed with

irradiance was related to the number of neurons recruited. The SDF from a single-unit

example was determined for the different irradiances; the amplitude of the neuronal response

increased with irradiance from 8.2 ± Hz for 6 mW.mm^{-2} to 36.6 ± 73 Hz for 106 mW.mm^{-2}

(Fig. 2a). We generalized this analysis by plotting the cumulative distribution of the mean

maximal firing rates for each irradiance (Fig. 2c, left panel). The cumulative curves reached a

plateau for lower maximal firing rates with decreasing irradiance. We noted a significant

difference in distribution between 1.2 mW.mm^{-2} and irradiances of 6 mW.mm^{-2} and above

(1.2 mW.mm^{-2}, median: 0 Hz; 6 mW.mm^{-2}, 3.8 Hz, Kolmogorov-Smirnov test, $p<0.0001$),

demonstrating that electrophysiological recordings and fUS imaging had equivalent

sensitivities. Finally, in our total population of V1 neurons, the proportion of responsive

neurons increased with irradiance (from 28/92 units at 6 mW.mm^{-2} to 71/92 units at 106

mW.mm^{-2}, Fig. 2c, right panel). Interestingly, the depth distribution of the activated neurons

did not change with increasing irradiance (Fig. 2c, right panel). It was, therefore, possible to

activate neurons from deep cortical layers even at very low irradiances.

Spread of optogenetic activation to downstream and upstream visual areas. The results

described above relate to optogenetic activation in V1 with the optic fiber placed at the

surface of the primary visual cortex. We then investigated whether the activity initiated in V1

spread to other visual structures up- or downstream. This aspect is important for optogenetic

therapies for the restoration of cortical vision because it would demonstrate the propagation

of visual information favoring the generation of visual percepts. For the downstream pathway,

LGN terminals end in V1 at the depth of cortical layer IV, whereas cortical V1 layer VI sends

feedback connections to the LGN. We thus hypothesized that injecting AAV9-7m8-CaMKII-

ChrimsonR-tdT into V1 would increase ChrimsonR expression in at least one of these two

categories of fibers and that the optogenetic activation of V1 would lead to direct activation of

the LGN via LGN terminals in V1, or to an indirect activation of the LGN through feedback

connections in V1 cortical layer VI. Histological analyses confirmed that some ChrimsonR

expression occurred in the LGN, but we were unable to identify any ChrimsonR-expressing

cell bodies in the LGN, suggesting that only retrograde fibers originating from cortical V1

layer VI expressed this opsin (Fig. S2). We imaged the planes containing the LGN (AP -3.5

to AP -5.5 mm). Figure 3a shows the fUS imaging planes at AP -5 mm for the direct eye and

optogenetic stimulation conditions. We noted a slight activation of the visual cortex following

visual stimulation (see Fig. 1b), and a strong activation of the ipsilateral LGN (n=85/225

active pixels) when the contralateral eye was stimulated with the white LED, suggesting that

this kind of stimulation was more appropriate for LGN and SC activations than for the visual

cortex. Indeed, in the 11 animals (Fig. 3b), the mean percentage of the LGN volume visually

activated was 20.5 ± 13.7% which is much greater than the volume activation obtained for V1

(less than 1%). When the injected V1 surface was stimulated with the optic fiber (Fig. 3a),

CBV responses increased significantly in the ipsilateral LGN, but with a much smaller

number of activated pixels (n=12/225 pixels) than for visual stimulation. We also performed a

control optogenetic stimulation, in which we stimulated the non-injected V1 area. We

observed no activation of the ipsilateral LGN, confirming that, on stimulation of the injected

hemisphere, ipsilateral LGN activation was due to optogenetic activation of LGN terminals in

V1 or feedback from the V1 area (Fig. S2). In the 11 animals tested, the active LGN volumes

for visual stimulation were larger than those for optogenetic stimulation (visual, mean: 20.5 ±

13.7%; optogenetic, mean: 6.5 ± 12.3%, Wilcoxon signed-rank test, p=0.0068). In addition,

for the five animals tested by optogenetic stimulation of the non-transfected V1 area, we

detected no significant responsive pixels (Fig. 3b, mean: 0 ± 0%, n=5 animals). We also

performed single-cell recordings in the LGN, to demonstrate that the neurovascular

activations imaged by fUS in the LGN coincided with the activation of LGN neurons (Fig. 3c).

We recorded a total of 153 neurons in the LGN. Only two units responded to both visual and

optogenetic stimulation; seven units were responsive only to optogenetic stimulation, and 40

units were responsive only to visual stimulation (n=7 animals). The vast majority of LGN

neurons were unresponsive to both types of stimulation. The distribution of onset latencies

following direct eye stimulation was broader for LGN than for V1 single units (Fig. 3d, mean:

47.88 ± 23.55 ms, n=42 units, and see Fig. 1f). This may reflect the recording of both cells

activated by retinal ganglion cells and cells retrogradely activated by the V1 area, which have

higher latencies. Following optogenetic stimulation at the surface of the V1 area, the onset

latencies for LGN neurons were shorter than those following visual stimulation (mean: 9.86 ±

4.95 ms, *n*=7 units, Mann-Whitney, *p*<0.05). However, these response latencies were greater

than those recorded in V1 (*n*=42 units. Mann-Whitney, *p*<0.0001, see Fig 1f for V1 units).

This result suggests that the optogenetically activated LGN single units recorded here were

activated indirectly, by retrograde fibers from V1 cortical layer VI, as suggested by the

histological data.

We also investigated whether V1 optogenetic stimulation could spread to the direct upstream

visual area toward which V1 projects: the V2 area. We obtained fUS activation maps in

different imaging planes (AP -6 to AP -8 mm) in which the entire V2 area was present after

either direct stimulation of the contralateral eye or optogenetic stimulation at the surface of

the V1 area. We show representative activation maps from single imaging planes (AP -8.5

mm) in Figure 4a. Visual stimulation of the contralateral eye led to almost no activation of the

V2 area on the same side as the injected V1. By contrast, optogenetic stimulation of the

transfected V1 area led to a stronger activation of the ipsilateral V2, mostly within the ventral

part of V2. As previously reported, within the different imaging planes, we also observed

strong activation in the lower parts of V1 and V2 containing the axons. We next quantified the

active volume over all imaging planes for the V2 area, for all 10 animals (Fig. 4b). As for the

V1 area, visual stimulation resulted in only weak activation of V2 (mean: 0.2 ± 0.3% of

activated volume). Averaged activation volumes were significantly larger for optogenetic

stimulation than for visual stimulation (mean: 5.6 ± 8.0% activated volume, *n*=10 animals,

Wilcoxon signed-rank test, one-tailed, *p*<0.05). Control stimulation of the non-injected V1

area confirmed the specificity of the spread of optogenetic activation from V1 to V2, as no

activation was observed in the contralateral V2 area (Fig. 4b, Fig. S2).

Electrophysiological recordings within V2 confirmed that the fUS variations we observed in

V2 were consistent with direct neuronal activation (Fig. 4c). We recorded V2 single units

responding to visual and optogenetic stimulation. As previously described, we characterized

the onset latency of these neurons, comparing the values obtained with those for V1 single

units (Fig. 4d). Similar visual latencies were recorded for V1 and V2 single units (V1, mean:

41.24 ± 16.61 ms, n=33 units (see Fig. 1f); V2, mean: 39.43 ± 9.557 ms, n=14 units).

Interestingly, optogenetic onset latencies were higher for V2 single units than for V1 (V1,

mean: 1.76 ± 3.14 ms, n=76 units. V2, mean: 62.78 ± 77.44 ms, n=9 units, Mann-Whitney,

p<0.0001), suggesting that these neurons were activated indirectly by ChrimsonR-expressing

V1 neurons. Two V2 neurons presented particularly long optogenetic ON latencies, possibly

due to a difference in the microcircuits involved. We checked that this variability did not bias

the latency delay for V2 and V1 neurons after optogenetic stimulation, by performing the

same statistical test on a V2 data set restricted to the remaining seven fast V2 units. The

difference in ON latency between V1 neurons and these fast V2 neurons was conserved

(fast V2 units, mean: 12.2 ± 9.5 ms, n=7 units, Mann-Whitney, p<0.0001), suggesting that all

V2 neurons were activated indirectly after the onset of stimulation. This conclusion is

supported by the lack of ChrimsonR expression in V2 on brain slices from the animals used

to record these single units. However, these V2 latencies were shorter than V2 latencies for

natural eye stimulation, demonstrating that they were genuinely produced by V1 optogenetic

activation.

Discussion

Coupling of optogenetics with fUS

In this study, we demonstrate that optogenetic activation can be detected by fUS imaging.

Rungta and colleagues[30] indicated that blue light from the tip of the optic fiber *per se* could

induce neurovascular responses in naïve mice, at irradiances higher than 2 mW ~ 18

mW.mm^{-2}. Another recent study reported hemodynamic responses following blue light

illumination of the retrosplenial cortex in Thy-ChR2 mice or the illumination of specific

neuronal subpopulations of the superior colliculus in other transgenic lines[26]. The authors

explained that they used a lower light power and irradiance (0.3 mW ~1.5 mW.mm^{-2}) to

prevent non-specific activation. By contrast, we used an AAV-mediated strategy to express

ChrimsonR in V1 neurons, resulting in a lower density of opsin-expressing cells than in

transgenic mice lines[33]. This generalizes the possibility of recording neurovascular

optogenetic activation due to a small number of transfected neurons. Moreover, we show

here that stimulation of the control non-injected hemisphere does not induce a vascular

response, thereby demonstrating the specificity of ChrimsonR-elicited optogenetic activity.

Single-cell recordings confirmed the neuronal optogenetic activation correlated with these

CBV variations. We also used a higher wavelength (595 nm) for illumination than Rungta and

coworkers, but with comparable values of power and irradiance. These results indicate that

red light can be used even at high power, in protocols combining the optogenetic activation

of red-shifted opsins and fUS imaging. Importantly, electrophysiological recordings from the

control animals showed no neuronal activation, suggesting negligible thermal effects with the

use of 595 nm light under the parameters used here. Indeed, we included these parameters

in the heat propagation model of Stujenske[34], and recorded a very local (<1 mm-diameter

sphere from the tip of the optic fiber) increase in temperature, by 0.2°C, which is too small to

alter neuronal firing rates. Finally, we characterized the dose-response dependence of our

recorded optogenetic activations. We obtained equivalent sensitivities for optogenetic

responses measured by fUS imaging and by electrophysiological recordings.

The use of fUS imaging to analyze optogenetic responses has the advantage that it could

potentially provide a mesoscopic view of the activated area. Unlike electrophysiological

recordings, which extract information very locally[20], and calcium imaging, which is dependent

on both expression of the calcium indicator and the penetration of blue light through the

tissue[35], it can display information at a brain-wide spatial scale. The coupling of fMRI with

optogenetics meets these criteria, but with a lower spatial resolution[21,22,36] (of the order of a

millimeter per pixel). By contrast, fUS imaging can detect brain activity at a submesoscopic

resolution (100 x 100 µm²), with less cumbersome equipment than for fMRI.

Spread of the activity

We found that optogenetic neurovascular activation was well correlated with ChrimsonR

expression in the AP axis. We also noted that optogenetic V1 stimulation spread beyond the

borders of V1, as indicated by the activation of both the LGN and V2 on both fUS and

electrophysiology. This spread of activity may be due to fibers located at the base of V1,

connected to the LGN and expressing ChrimsonR (Fig. S1). These fibers may generate the

neurovascular activity detected ventrally to V1 on our activation maps. Furthermore, the

mean neuronal transfection rate was 5.5% in the coronal plane displaying the highest

fluorescence signal, whereas a larger volume of V1 was activated in this plane. A first

explanation for this may be the spread of neuronal activation to other V1 cells, although we

recorded a rather homogeneous set of short-onset latencies. Alternatively, neurovascular

activations may be broader *per se* than neuronal responses. Some studies of rat olfactory

bulb glomeruli have provided evidence of a close overlap between capillary blood flow and

neuronal activity[37], whereas others have reported a mismatch between the areas of

neurovascular and neuronal activation[38,39]. Specifically for V1, a lack of correlation between

BOLD signals and spiking activity has been observed in cats[40], and a lack of correlation

between single-vessel hemodynamic responses and calcium imaging signals has been found

in cats and rats[41]. Importantly, we show here that the neurovascular and neuronal activities

initiated in V1 spread to the LGN and V2. This propagation of visual information is important

for optogenetic cortical vision restoration therapy, because it favors the generation of visual

percepts.

Cortical visual restoration

We detected neurovascular and neuronal responses, even at low irradiance (6 mW.mm^{-2}),

with no modification of the depth distribution of the activated neurons. One recent study[42]

reported that the stimulation of deep V1 layers (>1.5 mm) in non-human primates with

electrical prostheses elicited behavioral responses. Our ability to activate neurons in deep

cortical layers highlights the potential value of red-shifted opsin ChrimsonR for optogenetic

cortical vision restoration strategies. A key element of visual restoration is the induction of

neuronal responses with characteristics matching those resulting from a natural visual

stimulus[5–11]. We show here that the firing rates induced by optogenetic stimulation were

similar to those induced by visual stimulation. Optogenetically activated neurons had very

short latencies, of the order of 1-2 ms. Theoretically, this is sufficient for the encoding of

natural images into optogenetic stimulations at a temporal resolution matching the resolution

of the natural visual signal. Moreover, those optogenetic onset latencies were quite

homogeneous, despite being obtained for neurons located in different layers. These results

demonstrate that we can induce a signal that does not lose its temporal resolution with

depth.

Finally, the feasibility of inducing visual percepts by optogenetic cortical vision restoration

remains to be demonstrated. In species with a more complex hierarchical organization of

cortical visual areas, such as nonhuman primates, a few studies have shown behavioral

effects due to the optogenetic stimulation of higher visual areas. Jazayeri[43] and coworkers

reported that saccades following fixation tasks were shifted towards the receptive field of the

region of V1 optogenetically activated following the fixation point offset. Ju[44] and coworkers

demonstrated the successful detection of optogenetic percepts in a saccade task. Our ability

to detect a spread of activity from V1 to other visual areas is consistent with these behavioral

studies indicating perception. Here, we used a single optic fiber for optogenetic stimulation.

Replacing this device with a more complex stimulation system, such as arrays of micro-

LEDs, might make it possible to generate percepts more complex than phosphenes and to

develop discrimination behavioral tasks to assess their perception.

Materials and methods

Animals

All animal experiments and procedures were approved by the Local Animal Ethics Committee (registration number 2018032911282465) and performed in accordance with European Directive 2010/63/UE. We used wild-type male Long-Evans rats (Janvier Laboratories), nine weeks old at the time of viral injection. Rats were maintained under a reverse 12-hour light/12-hour dark cycle, with ad libitum access to food and water, except during surgery and electrophysiological recordings.

AAV production

The AAV9-7m8-CaMKII-ChrimsonR-tdT vector was packaged as previously described, by the triple transfection method, and purified by iodixanol gradient ultracentrifugation[45]. The AAV9-7m8-CaMKII-ChrimsonR-tdT vector was titered by qPCR with SYBR Green[46] (Thermo Fisher Scientific). The titer used in this study was 4.39×10^{12} vg.mL^{-1}.

Immunostaining and confocal imaging

Following electrophysiological recordings, rats were euthanized, and their brains were extracted and fixed by overnight incubation in 4% paraformaldehyde (100496, Sigma-Aldrich) at 4°C. Brains were then cryoprotected in 30% sucrose (84097, Sigma-Aldrich) and 50 μm sagittal slices were cut with a microtome (HM450, Microm). The slices with the most intense tdT fluorescence from each brain were selected for further immunohistochemistry and imaging. Cryosections were permeabilized by incubation with 0.5% Triton X-100 in PBS for 1 h at room temperature and were then incubated in blocking buffer (PBS + 1% BSA + 0.1% Tween 20) for 1 hour at room temperature. Samples were incubated overnight at 4°C with monoclonal anti-NeuN antibody (1:500; Mouse, MAB377, Merck Millipore), in a 50% dilution of blocking buffer + 0.5% Triton X-100. Secondary antibodies conjugated with Alexa Fluor dyes (1:500; Molecular Probes) and DAPI (1:1000, D9542, Merck Millipore), were incubated with the samples for 1 hour at room temperature. An Olympus FV1000 laser-scanning confocal microscope with a 20x or 40x objective (UPLSAPO 20XO, NA: 0.85) was used to acquire images of brain sections.

Viral injections

Viral injections were performed in aseptic conditions with a digital small-animal stereotaxic instrument (David Kopf Instruments). Ear bars were covered with xylocaine to ensure that the animals felt no pain. Rats were anesthetized in a sealed box containing gaseous isoflurane (5%), and maintained under anesthesia in the stereotaxic frame (25% ketamine, 10%

medetomidine and 65% saline injected intraperitoneally) for the entire surgical procedure,
and animal body temperature was maintained at 37°C with a heating pad. Buprenorphine
was injected subcutaneously to reduce inflammation, and Lubrithal was applied to the eyes
to prevent them from drying out. Xylocaine, 70% ethanol and Vetedine were applied
successively to the scalp before incision, to minimize pain and maintain sterile conditions.
Cranial sutures were cleaned to remove connective tissue, by applying H_2O_2, to facilitate
localization of the injection coordinates. We injected a total volume of 1.2 µL of viral
suspension unilaterally into rates, via two injection tracks, at a flow rate of 50-75 nL/min, with
a 5 or 10 µL microsyringe (Hamilton) equipped with a microinjector (Sutter Instrument) and
controller (World Precision Instruments). The coordinates for viral injection were +2.8 / +3.2
mm from midline (M-L axis), -6.5 / -7.5 mm from Bregma (A-P axis) and 1.6-1.35-1.1 mm
ventral to the skull surface (D-V axis), based on the 2004 edition of the Paxinos and Franklin
rat brain atlas. Viral efflux was prevented by leaving the needle in a 1.8 mm ventral position
for two minutes before beginning the injection, with a three-minute interval left between
injections before complete withdrawal of the needle from the cortex. After surgery, rats were
brought round from anesthesia with a subcutaneous injection of Antisédan (0.15 mL).

In vivo **electrophysiological recordings**

Bilateral craniotomies were performed with a digital small-animal stereotaxic instrument
(David Kopf Instruments), at least 30 days after viral injection, to allow time for opsin
expression. Ear bars were covered with xylocaine to prevent pain. Rats were anesthetized in
a sealed box containing gaseous isoflurane (5%) and maintained under anesthesia in the
stereotaxic frame (25% ketamine, 10% medetomidine and 65% saline injected
intraperitoneally) for the entire surgical procedure, and electrophysiological recordings were
taken with body temperature maintained at 37°C with a heating pad. Buprenorphine was
injected subcutaneously to reduce inflammation, and Lubrithal was applied to the eyes to
prevent them from drying out. Cranial sutures were cleaned to remove connective tissue, by
applying H_2O_2, to reveal the injection tracks. Parietal bones were removed by drilling
rectangular flaps and gently moving the bone away from the dura mater, exposing the cortex
from 3 to 8.5 mm from Bregma (A-P axis), to cover the injection tracks. The dura was then
gently removed. During drilling, the skull was regularly cooled with PBS, and once the cortex
was exposed, it was protected from dehydration by the regular application of cortex buffer.
After surgery, electrophysiological recordings were performed with 16-channel electrodes
(A1x16-5 mm-50-703-OA16LP) coupled with a 400 µm-core fiber (Thorlabs M79L005 Fiber
Cable, MM, 400 µm 0.39NA, FC/PC to 1.25 mm ferrule, 0.5 m) at various positions close to
the injection sites. Electrophysiological data were acquired with MC RACK software. For

visual stimulation, a white LED (Thorlabs MNWHL4 Mounted LED, 5V, 60 mW/cm^2) was

placed 15 cm in front of the eye on the contralateral side of the cranial window.

Acquisition protocol for electrophysiological recordings

For optogenetic stimulation, we connected the optic fiber (reference above) to a light source

(Thorlabs M595F2 (fiber coupled LED @595 nm), Ø400 µm, 150 µW/cm^2) delivering light at

the ChrimsonR excitation wavelength (595 nm). We targeted the transfected region of V1 by

imaging tdT fluorescence with a Micron IV imaging microscope (Phoenix Research

Laboratories) before recordings. The fiber was placed on the surface of the cortex while the

electrode was inserted in the tissue. Both stimulations consisted of 100 repeats of 200 ms

flashes at 1 Hz. The onset of the flashes was aligned with electrophysiological data, with

Clampex 9.2 software. We used several different irradiances of light at 595 nm in this study.

Power at the fiber tip was measured with a power meter (Thorlabs, PM100D), by placing the

fiber tip in contact with the sensor. The irradiance corresponding to each power was

calculated as previously described[30].

Irradiance (mW.mm^{-2})	Power (mW)
142	7.1
140	7.0
106	5.3
70	3.5
36	1.8
14	0.7
6	0.3
1.2	0.06

487 Table 1: 595 nm light irradiances and powers used in this study

Spike sorting

Offline spike sorting of the electrophysiological recordings (linear 16-channel electrodes) was
performed with the SpyKING CIRCUS package[32]. Raw data were first high-pass filtered (>
300 Hz) and spikes were detected when a filtered voltage trace crossed the threshold.
Automatic cluster extraction was performed and candidate clusters were curated. Refractory
period violations (< 2 ms, >1% violation) and noisy spike shapes led to cluster deletion. Spike
templates with coordinated refractory periods in the cross-correlogram together with similar
waveforms led to the merging of cluster pairs.

Electrophysiological data analysis

All electrophysiological data were extracted and analyzed with a custom-made Matlab script.
Responsive units were defined as those displaying a significant difference in neuronal activity
between the pre-stimulation period (averaged 100 ms before stimulus onset) and the
stimulation interval (averaged 200 ms following stimulus onset, Wilcoxon signed-rank test,
$p<0.01$). For each unit, responses are represented as the spike density function (SDF), which
was calculated from the mean peristimulus time histogram (PSTH, bin size: 1 ms, 100 trials)
smoothed with a Gaussian filter (2 ms SD).

Calculation of latencies and response durations

The latency of the units displaying activation was defined as the first time point at which the
SDF crossed the value of the baseline plus 2SD and remained higher than this value for at
least 10 ms. Conversely, the offset of activation was defined as the first time point after
latency that the SDF crossed back below the value of the baseline plus 2SD and remained
lower than this value for at least 10 ms. Not all active neurons from a given population met
these criteria, accounting for the slight difference between the number of active neurons and
the number of latencies presented here.

Generation of functional ultrasound images

fUS imaging was performed as previously described[24], with a linear ultrasound probe (128
elements, 15 MHz, 110 µm pitch and 8 mm elevation focus, Vermon; Tour, France) driven by
an ultrafast ultrasound scanner (Aixplorer, Supersonic Imagine; Aix-en-Provence, France).

Acquisition protocol for fUS imaging

3D fUS acquisitions were performed after craniotomy and electrophysiological recordings, as
previously described. When optogenetic responses were observed, the position of the optic
fiber on the surface of the cortex was kept unchanged until control acquisitions were
performed, in which the fiber was moved to the other hemisphere. The cortex buffer on the

surface of the cortex dried out, and 1 cm^3 of ultrasound coupling gel was placed between the

cortex and the linear ultrasound probe. Acquisition protocols consisted of 20 stimulation

blocks, each consisting of 13 s of rest followed by 2 s of stimulation. For visual stimulation,

the white LED used for electrophysiological recordings was kept in the same position, and

the 2 s stimulation consisted of eight repeats of 125 ms flashes at 4 Hz. For optogenetic

stimulation, the 2 s stimulation consisted of 40 repeats of 25 ms flashes at 20 Hz.

Acquisitions were performed on coronal planes from 3 mm to 8.5 mm from Bregma (AP

axis), with a 0.5 mm increment corresponding to the thickness of the imaging plane.

529 Building activation maps

For each pixel, we averaged, for each block, the intensity of Doppler power at baseline (2 s

before stimulus onset) and during a response window (4 s after stimulus onset). The signals

were then compared in one-tailed Wilcoxon signed-rank tests. Only pixels with p-values <

4.03×10^{-6} (corresponding to a global p-value < 0.05 with Bonferroni correction) were

considered significant. On the maps, CBV during the response window is presented as a

percentage of the baseline value. Region of interest (ROI) as V1, V2, SC and LGN were

determined for each imaging plane, based on the Matt Gaidica rat brain atlas.

539 References

1. Deisseroth, K. Optogenetics: 10 years of microbial opsins in neuroscience. *Nat. Neurosci.* **18**,

1213–1225 (2015).

2. Wrobel, C. *et al.* Optogenetic stimulation of cochlear neurons activates the auditory pathway and

restores auditory-driven behavior in deaf adult gerbils. *Sci. Transl. Med.* **10**, (2018).

3. Dieter, A., Keppeler, D. & Moser, T. Towards the optical cochlear implant: optogenetic approaches

for hearing restoration. *EMBO Mol. Med.* **12**, e11618 (2020).

4. Roska, B. & Sahel, J.-A. Restoring vision. *Nature* **557**, 359–367 (2018).

5. Bi, A. *et al.* Ectopic Expression of a Microbial-Type Rhodopsin Restores Visual Responses in Mice

with Photoreceptor Degeneration. *Neuron* **50**, 23 (2006).

6. Lagali, P. *et al.* Light-activated channels targeted to ON bipolar cells restore visual function in

retinal degeneration. *Nature neuroscience* vol. 11 https://pubmed.ncbi.nlm.nih.gov/18432197/

(2008).

7. Gaub, B. M. *et al.* Restoration of visual function by expression of a light-gated mammalian ion

channel in retinal ganglion cells or ON-bipolar cells. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E5574–

E5583 (2014).

8. Khabou, H. *et al.* Noninvasive gene delivery to foveal cones for vision restoration. *JCI Insight* **3**,

(2018).

9. Sengupta, A. *et al.* Red-shifted channelrhodopsin stimulation restores light responses in blind

mice, macaque retina, and human retina. *EMBO Mol. Med.* **8**, 1248–1264 (2016).

10.   Chaffiol, A. *et al.* A New Promoter Allows Optogenetic Vision Restoration with Enhanced

Sensitivity in Macaque Retina. *Mol. Ther.* **25**, 2546–2560 (2017).

11.   Gauvain, G. *et al.* Optogenetic therapy: High spatiotemporal resolution and pattern

recognition compatible with vision restoration in non-human primates. *bioRxiv*

2020.05.17.100230 (2020) doi:10.1101/2020.05.17.100230.

12.   Allergan. *Phase I/IIa, Open-Label, Dose-Escalation Study of Safety and Tolerability of*

Intravitreal RST-001 in Patients With Advanced Retinitis Pigmentosa (RP).

https://clinicaltrials.gov/ct2/show/NCT02556736 (2020).

13.   GenSight Biologics. *A Phase 1/2a, Open-Label, Non-Randomized, Dose-Escalation Study to*

Evaluate the Safety and Tolerability of GS030 in Subjects With Retinitis Pigmentosa.

https://clinicaltrials.gov/ct2/show/NCT03326336 (2020).

14.   Beauchamp, M. S. *et al.* Dynamic Stimulation of Visual Cortex Produces Form Vision in

Sighted and Blind Humans. *Cell* **181**, 774-783.e5 (2020).

15.   Bosking, W. H., Beauchamp, M. S. & Yoshor, D. Electrical Stimulation of Visual Cortex:

Relevance for the Development of Visual Cortical Prosthetics. *Annu. Rev. Vis. Sci.* **3**, 141–166

(2017).

16.   Seabrook, T. A., Burbridge, T. J., Crair, M. C. & Huberman, A. D. Architecture, Function, and

Assembly of the Mouse Visual System. *Annu. Rev. Neurosci.* **40**, 499–538 (2017).

17.   Klapoetke, N. C. *et al.* Independent optical excitation of distinct neural populations. *Nat.*

Methods **11**, 338–346 (2014).

18.   Mager, T. *et al.* High frequency neural spiking and auditory signaling by ultrafast red-shifted

optogenetics. *Nat. Commun.* **9**, 1750 (2018).

19.   Marshel, J. H. *et al.* Cortical layer-specific critical dynamics triggering perception. *Science* **365**,

(2019).

20.   Buzsáki, G., Anastassiou, C. A. & Koch, C. The origin of extracellular fields and currents —

EEG, ECoG, LFP and spikes. *Nat. Rev. Neurosci.* **13**, 407–420 (2012).

21.   Liang, Z. *et al.* Mapping the Functional Network of Medial Prefrontal Cortex by Combining

Optogenetics and fMRI in Awake Rats. *NeuroImage* **117**, 114–123 (2015).

22.   Chen, X. *et al.* Mapping optogenetically-driven single-vessel fMRI with concurrent neuronal

calcium recordings in the rat hippocampus. *Nat. Commun.* **10**, (2019).

23.   Macé, E. *et al.* Functional ultrasound imaging of the brain. *Nat. Methods* **8**, 662–664 (2011).

24.   Gesnik, M. *et al.* 3D functional ultrasound imaging of the cerebral visual system in rodents.

Neuroimage **149**, 267–274 (2017).

25.   Macé, É. *et al.* Whole-Brain Functional Ultrasound Imaging Reveals Brain Modules for

Visuomotor Integration. *Neuron* **100**, 1241-1251.e7 (2018).

26.   Brunner, C. *et al.* A Platform for Brain-wide Volumetric Functional Ultrasound Imaging and

Analysis of Circuit Dynamics in Awake Mice. *Neuron* **108**, 861-875.e7 (2020).

27.   Bergel, A., Deffieux, T., Demené, C., Tanter, M. & Cohen, I. Local hippocampal fast gamma

rhythms precede brain-wide hyperemic patterns during spontaneous rodent REM sleep. *Nat.*

Commun. **9**, (2018).

28.   Dizeux, A. *et al.* Functional ultrasound imaging of the brain reveals propagation of task-

related brain activity in behaving primates. *Nat. Commun.* **10**, (2019).

29.   Blaize, K. *et al.* Functional ultrasound imaging of deep visual cortex in awake nonhuman

primates. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 14453–14463 (2020).

30.   Rungta, R. L., Osmanski, B.-F., Boido, D., Tanter, M. & Charpak, S. Light controls cerebral

blood flow in naive animals. *Nat. Commun.* **8**, 14191 (2017).

31.   Gao, Y.-R. *et al.* Time to wake up: Studying neurovascular coupling and brain-wide circuit

function in the un-anesthetized animal. *NeuroImage* **153**, 382–398 (2017).

32.   Yger, P. *et al.* A spike sorting toolbox for up to thousands of electrodes validated with ground

truth recordings in vitro and in vivo. *eLife* **7**,.

33.   Arenkiel, B. R. *et al.* In vivo light-induced activation of neural circuitry in transgenic mice

expressing channelrhodopsin-2. *Neuron* **54**, 205–218 (2007).

34.   Stujenske, J. M., Spellman, T. & Gordon, J. A. Modeling the Spatiotemporal Dynamics of Light

and Heat Propagation for In Vivo Optogenetics. *Cell Rep.* **12**, 525–534 (2015).

35.   Emiliani, V., Cohen, A. E., Deisseroth, K. & Häusser, M. All-Optical Interrogation of Neural

Circuits. *J. Neurosci.* **35**, 13917–13926 (2015).

36.   Gerits, A. *et al.* Optogenetically-induced behavioral and functional network changes in

primates. *Curr. Biol. CB* **22**, 1722–1726 (2012).

37.   Chaigneau, E., Oheim, M., Audinat, E. & Charpak, S. Two-photon imaging of capillary blood

flow in olfactory bulb glomeruli. *Proc. Natl. Acad. Sci.* **100**, 13081–13086 (2003).

38.   Iadecola, C., Yang, G., Ebner, T. J. & Chen, G. Local and propagated vascular responses

evoked by focal synaptic activity in cerebellar cortex. *J. Neurophysiol.* **78**, 651–659 (1997).

39.   Iadecola, C. The neurovascular unit coming of age: a journey through neurovascular coupling

in health and disease. *Neuron* **96**, 17–42 (2017).

40.   Kayser, C., Kim, M., Ugurbil, K., Kim, D.-S. & König, P. A Comparison of Hemodynamic and

Neural Responses in Cat Visual Cortex Using Complex Stimuli. *Cereb. Cortex* **14**, 881–891 (2004).

41.   O'Herron, P. *et al.* Neural correlates of single-vessel haemodynamic responses in vivo. *Nature*

534, 378–382 (2016).

42.   Chen, X., Wang, F., Fernandez, E. & Roelfsema, P. R. Shape perception via a high-channel-

count neuroprosthesis in monkey visual cortex. *Science* **370**, 1191–1196 (2020).

43.   Jazayeri, M., Lindbloom-Brown, Z. & Horwitz, G. D. Saccadic eye movements evoked by

optogenetic activation of primate V1. *Nat. Neurosci.* **15**, 1368–1370 (2012).

44.   Ju, N., Jiang, R., Macknik, S. L., Martinez-Conde, S. & Tang, S. Long-term all-optical

interrogation of cortical neurons in awake-behaving nonhuman primates. *PLoS Biol.* **16**, e2005839

(2018).

45.   Choi, V. W., Asokan, A., Haberman, R. A. & Samulski, R. J. Production of recombinant adeno-

associated viral vectors. *Curr. Protoc. Hum. Genet.* **Chapter 12**, Unit 12.9 (2007).

46.   Aurnhammer, C. *et al.* Universal real-time PCR for the detection and quantification of adeno-

associated virus serotype 2-derived inverted terminal repeat sequences. *Hum. Gene Ther.*

Methods **23**, 18–28 (2012).

Acknowledgments

We thank Pierre Pouget, Frédéric Chavane, Brice Bathellier, Kévin Blaize, Sara Cadoni and Diep
Nguyen for their useful discussions about the data. We thank Julie Dégardin for her help with
surgery, Baptiste Lefebvre for assistance with spike sorting and Hicham Serroune for technical
assistance with fUS imaging. This work was supported by the European Research Council (ERC)
Synergy Grant Scheme (ERC Grant Agreement 610110).

Competing financial interests

M.T. is cofounder and S.P. and M.T. are shareholders of ICONEUS. D.D. and S.P. are consultants for
Gensight Biologics.

Author contributions

M.P., S.P., J-A.S., M.T., G.G., and F.A. designed the study; M.P. and D.D. designed viral vectors; M.P.
and W.D. produced viral vectors; M.P., G.L., A.R., R.G. and M.V. performed intracortical injections;
M.P., G.L. and C.J. performed electrophysiological recordings and fUS acquisitions; G.L. performed
histological experiments; M.P., U.F., A.C., G.G., F.A. and S.P. analyzed data; M.P., G.G. and F.A.
constructed the figures; M.P., A.C., G.G., F.A. and S.P. wrote the manuscript.

Figure 1 | V1 neurovascular and neuronal activations resulting from cortical surface stimulation in rats. (a) Left: 2.5x imaging (top) and 20x confocal imaging of a brain section showing the localization of ChrimsonR in V1. White dashed lines delimit the cortical surface and border between V1 and the white matter. Right: close-up (40x) of the area delimited by the yellow lines in the previous image, showing two ChrimsonR-expressing neurons. Scale bars: 500 µm, 200 µm, 20 µm. (b) fUS activation maps obtained after visual stimulation of the contralateral eye (left), optogenetic stimulation of the ipsilateral V1 area expressing ChrimsonR (middle) and control optogenetic stimulation of the uninjected contralateral V1 area (right), from a single imaging plane (AP -7.5 mm) from the same animal. White dashed lines delimit the V1 and SC areas. Colored pixels indicate significant CBV variation (Wilcoxon signed-rank test with Bonferroni-Holm correction). Right insets: patterns of single-pixel activation. Gray lines represent single-trial activity (n=20) and the black line represents the mean CBV variation. Colored patches indicate light stimulation (duration: 2 s) (c) fUS activation maps for visual and optogenetic activation, for all imaging planes (AP -6 mm to AP -8 mm) in which V1 (delimited by white dashed lines) is present, from the same animal (animal #2). (d) Left: percentage of V1 activated during optogenetic stimulation for each fUS imaging plane, for all animals (n=10). Right: AP distribution of the ChrimsonR expression area on brain sections. (e) Spike density function (SDF) of typical V1 single units during visual (black lines) or optogenetic (red lines) activation. Four subpopulations of neurons were identified: double-responsive (n_{V+O}=13), responsive to the visual (n_V=20) or optogenetic stimulus only (n_O=81), non-responsive (n_{None}=57). (f) Left: ON response latencies for visual (n=33 units) and optogenetic (n=76 units) stimulation (Mann-Whitney, p<0.0001). Right: ON response durations for visual (n=33 units) and optogenetic stimulation (n=72 units). (g)

Depth profile of transient (ON duration<51 ms, n=33 units) and sustained (ON duration>197 ms, n=22 units) neurons activated by optogenetics. (c-d) Scale bar: 2 mm. The irradiance used for optogenetic and control stimulation was 140 mW.mm^{-2}.

Figure 2 | Neuronal and neurovascular optogenetic sensitivity. (a) Top: fUS activation maps of a single imaging plane (AP -7.5 mm) from animal #4 at different irradiances. Scale bar: 2 mm. White dashed lines delimit the V1 and SC areas. Bottom: SDF from a typical V1 single unit during optogenetic stimulation for 200 ms (gray patch) at different irradiances. (b) Left panel: mean CBV variation over all animals (n=9) for each irradiance. of the values for each animal were normalized against those obtained at 106 mW.mm^{-2}. Error bars represent the standard deviation. The colored patch corresponds to the 2 s optogenetic stimulation. Right panel: normalized active volumes of the ipsilateral V1 area for each irradiance. Open circles are individual values, bars represent the mean and the standard deviation (n=9 animals, Wilcoxon signed-rank test, one-tailed, p<0.05). (c) Left: cumulative distribution of the mean maximal firing rates of all recorded V1 single units (n=92) during optogenetic stimulation, for each irradiance. Right panel, top: depth profile of V1 single units activated (black squares) or not (gray squares) by the optogenetic stimulation, for each irradiance (n=92 single units). Bottom: percentage of active (black) and non-responsive (gray) neurons for each set of conditions.

Figure 3 | Spread of optogenetic activation to the LGN. (a) fUS activation maps obtained during visual stimulation of the contralateral eye (top) and optogenetic stimulation of the ChrimsonR-expressing ipsilateral V1 area (bottom) from a single imaging plane in which the LGN is present (AP -5 mm) from the same animal. White dashed lines delimit the ipsilateral LGN. Scale bar: 2 mm. (b) Percentage active volume of the LGN after visual, optogenetic or control stimulation. For both visual and optogenetic stimulations, we show the volumes of the LGN ipsilateral to the injection, whereas, for control stimulations, the volume of the contralateral LGN is shown. Open circles represent mean values over all imaging planes for each animal (visual and optogenetic stimulation, n=11, Wilcoxon signed-rank test, one-tailed, p=0.0068, n=5 for control, Mann-Whitney test, one-tailed, p=0.0288), bars represent the mean for all animals. (c) SDF of a typical LGN single unit responding to visual (black line) and optogenetic (red line) stimulation of the ipsilateral V1 area. (d) ON latencies of V1 and LGN single-unit responses to visual (V1, n=33 units, LGN, n=42 units) or optogenetic stimulation of V1 (V1, n=29 units, LGN, n=7 units. Mann-Whitney test, p<0.05). The irradiance used for optogenetic and control stimulation was 142 mW.mm^{-2}.

Figure 4 | Spread of the optogenetic activation to V2. (a) fUS activation maps obtained during visual stimulation of the contralateral eye (top) and optogenetic stimulation of the ChrimsonR-

expressing ipsilateral V1 area (bottom) and control stimulation of the uninjected contralateral area from
a single imaging plane (AP -7.5 mm) containing the V2 area. White dashed lines delimit the ipsilateral
V2 area. Scale bar: 2 mm. (b) Percentage active volume of V2 after visual, optogenetic or control
stimulation. For both visual and optogenetic stimulation, we show the volumes of V2 ipsilateral to the
injection, whereas, for the control, we shown the volume of the contralateral V2. Open circles
represent mean values over all imaging planes for each animal (visual and optogenetic, n=1, Wilcoxon
signed-rank test, one-tailed, p=0.002, n=7 for control, Mann-Whitney test, one-tailed, p=0.0004), bars
represent the mean over all animals. (c) SDF of a typical V2 single unit responding to visual (black
line) and optogenetic (red line) stimulation of the ipsilateral V1 area. (d) ON latencies of V1 and V2
single-unit responses to visual (V1, n=33 units, V2, n=14 units) or optogenetic stimulation of V1 (V1,
n=76 units, V2, n=8 units). The irradiance used for optogenetic and control stimulation was 140
mW.mm^{-2}.