The landscape of high-affinity human antibodies against intratumoral antigens

Goran Rakocevic[1], Irina Glotova[1], Ines de Santiago[1], Berke Cagkan Toptas[1], Milena Popovic[1], Milos Popovic[1], Dario Leone[1], Andrew L. Stachyra[1], Raphael Rozenfeld[1], Deniz Kural[1,*], Daniele Biasci[1,*]

[1] Totient Inc. 1 Alewife Center Suite 120, Cambridge, MA 02140

* Corresponding authors: daniele.biasci@totient.bio, deniz.kural@totient.bio

Abstract

High expression of immunoglobulin transcripts is often detected in human tumours and correlates with favourable clinical outcomes across different cancer types. However, the antigens recognized by such immunoglobulins (Ig) and their biological significance remain largely unknown. We computationally inferred the paired sequence of thousands of clonally expanded Ig using bulk RNA sequencing data from solid tumors in The Cancer Genome Atlas (TCGA). After expressing 283 Ig as recombinant antibodies in mammalian cells, we individually tested them against a library of twenty thousands full-length human proteins and an additional library of six thousand structurally intact membrane proteins, thus identifying their candidate target antigens. Using surface plasmon resonance we then confirmed 16 high-affinity antibodies that bind to their targets with K_D in the nanomolar range. Our work provides insights into the antigens that drive B cell responses in human cancers, while also obtaining fully human, high-affinity recombinant antibodies against them.

Main

Transcripts encoding immunoglobulin light and heavy chains are often detected in solid tumors across different cancer types[1], but their functional relevance remains unclear. Certain characteristics of the intratumoral Ig repertoire (e.g., transcripts abundance, clonality and number of detectable somatic mutations) have been associated with favourable clinical outcomes, such as

longer overall survival and response to immune checkpoint inhibitors[1-3]. Moreover, the presence

of intratumoral plasma cells and ectopic germinal centers, which are key components of the

antibody selection and production machinery[4], has been associated with longer overall survival

and immunotherapy response[5-8]. Despite these observations, the contribution of intratumoral Ig

to immune responses against cancer remains largely unknown.

The main obstacle to the functional characterization of intratumoral Ig is our limited

understanding of their target antigens. Previous studies have demonstrated that sequences of

individual Ig chains can be reconstructed using bioinformatics methods from bulk RNA

sequencing data (RNA-Seq) generated by large-scale cancer genomics efforts[1,3]. Compared to

specialized B-cell receptor (BCR) sequencing or single-cell sequencing, bulk RNA-seq has the

significant advantage of being readily available for thousands of clinically annotated tumor

samples[9]. However, previous studies have been limited to in-silico analysis: they did not attempt

to pair heavy and light Ig chains, nor to express the resulting sequences as complete antibodies,

two key steps required to experimentally identify their target antigens.

In this work we assembled and paired thousands of intratumoral Ig chains using legacy bulk

RNA-Seq data from TCGA, one of the most comprehensive genomic studies of human cancer to

date[9] (Figure 1). We individually performed gene synthesis, mammalian expression, and

purification for 283 pairs of Ig chains, obtaining high quality, fully human recombinant

antibodies in most cases. We then individually screened each antibody against two large

collections of recombinant proteins, covering the vast majority of the human proteome, in order

to obtain the most likely binding targets. In selected cases, we confirmed binding using surface

plasmon resonance (SPR) in order to characterize the binding kinetics. Our results show that

fully functional antibodies can be obtained from legacy bulk tumour RNA-Seq, without the need

for specialized BCR or single-cell sequencing. Using this approach, we were able to identify the

target antigens of several high-affinity Ig expressed in human tumors.

Results

Identification of dominant Ig sequences in the TCGA dataset

It has been observed that in some tumor samples the Ig repertoire is particularly clonal: a small

set of Ig transcripts are expressed at high levels and account for the vast majority of all the reads

attributable to Ig genes in that sample[1,3]. We speculated that this phenomenon, which probably

results from selected B cell clones winning the competition for the limited access to T follicular

helper (Tfh) cells in the germinal center (GC) reaction[4], could be used to correctly pair heavy and

light Ig chains from bulk RNA-Seq data, at least in some cases. Accordingly, we computationally

assembled Ig chain sequences expressed in each TCGA tumor sample and we calculated the

associated Berger–Parker index[10] in order to identify samples with particularly dominant

sequences. We observed that a small, but not insignificant, proportion of the analyzed samples

expressed highly dominant Ig sequences (Supplementary Figure 2).

Dominant Ig sequences can be correctly paired from bulk RNA-Seq in-silico

As a next step, we wanted to assess whether the dominance information captured by the

Berger–Parker index could be used to correctly pair heavy and light chains from bulk RNA-Seq

data. In order to do so, we used RNA-Seq data from single B cells[11] to simulate bulk RNA-Seq

samples for which the expression level, sequence, and correct pairing of all Ig chains was known.

Using this approach, we established that for samples with dominance scores larger than 0.382 we

could identify and correctly pair the dominant Ig chains with high accuracy (true positive rate

0.46, false positive rate 0.1, AUC=0.757; Figure 2). We used this result to select 191 paired

intratumoral Ig sequences from our TCGA analysis for expression in mammalian cells and

further characterization (Figure 3). As a control, we also selected for expression and

characterization 92 sequences below our chosen dominance score cutoff.

In-silico paired Ig sequences can be expressed at high levels in mammalian cells

We performed gene synthesis for 283 paired Ig sequences and attempted expression of the

corresponding antibody proteins in mammalian cells (HEK293). For each candidate, we replaced

the heavy constant region with a standard human IgG1 sequence in order to facilitate subsequent

detection and screening. We successfully expressed 275 recombinant antibodies out of 283 with

high yield (mean yield 2.35 mg/150mL; Supplementary Table S1). After expression, each

antibody was purified using protein A, stored in PBS at a standard concentration of 1ug/uL and

used for subsequent target antigen screening.

Identification of target antigens using high-throughput proteomics

Having successfully expressed the intratumoral Ig we paired in-silico, we decided to use the resulting recombinant antibodies to attempt identification of their target antigens. To this aim, we used high throughput proteomics to screen individual antibodies against a large collection of wild-type human proteins. We screened 173 antibodies against a collection of about twenty thousand human proteins, each represented in duplicate on the surface of a protein array, covering about 80% of the known human proteome[12]. While protein arrays are a cost effective way to test antibodies against a large number of potential targets, this technology is not designed to display membrane proteins in their correct conformation. Complex membrane proteins are unlikely to fold correctly when immobilized on the surface of the array, outside or their native membrane context. For this reason, we used high-throughput fluorescence-activated cell sorting (HT-FACS) to screen 92 antibodies against a library of six thousand structurally-intact membrane proteins expressed in human HEK0293T cells[13]. As a result of these screenings, we were able to identify high confidence hits for 84 of our antibody candidates screened using protein arrays (48%) and 21 of candidates screened using HT-FACS (23%). Targets recognized by our in-silico paired intratumoral Ig included well known cancer-specific antigens[14] (NY-ESO-1, MAGEA3, GAGE2A, DLL3) as well as immunomodulatory molecules expressed in the tumor microenvironment (ANXA1, TGFBI, C4BPB; Figure 4 and Supplementary tables 2, 4 and 5).

Intratumoral Ig bind their target antigens with high-affinity

As a next step, we decided to independently confirm the interaction between selected intratumoral Ig and their putative target antigens by using surface plasmon resonance (SPR)[15]. For each intratumoral Ig sequence, we characterized the binding affinity of the corresponding recombinant antibody to its putative target antigen, having sourced the antigen independently of the vendor used for the high-throughput proteomics screening. We confirmed 19 antibody-antigen interactions (Supplementary Table S2). We observed that the recombinant antibodies derived from intratumoral Ig sequences bind to their target antigen with very high-affinity, with K_{D} in the low nanomolar range (Figure 4 and Supplementary Table S2). When we compared our fully human antibodies against commercial antibodies obtained from rabbits (a

model organism known to generate very high-affinity antibodies[16,17]) after immunisation with the

same antigen, we found no significant differences in their binding affinity (Figure 4).

Epitope mapping for recombinant antibodies derived from intratumoral Ig

Having demonstrated that our in-silico paired intratumoral Ig bind to their target antigens with

high-affinity, we decided to assess whether it would be possible to identify their epitope, at least

in principle. We selected one corresponding recombinant antibody and performed epitope

mapping using hydrogen/deuterium exchange mass spectrometry (HDX-MS)[18]. Using this

technique, we were able to identify the putative binding region of the antibody on its designated

target (C4BPB; Figure 5). We observed that the most likely epitope overlaps with the C4BPB

binding site for protein S[19], which is crucial for its biological function[20]. Although additional

functional studies would be required to clarify whether or not this particular antibody is able to

interfere with the binding between C4BPB and protein S, our workflow shows that in principle it

is possible to identify the target antigen and the corresponding epitope for in-silico paired

intratumoral Ig, thus gaining further insights into their biological function.

Discussion

Despite the striking correlation between expression of Ig transcripts and favourable clinical

outcomes in human tumors, the functional role of intratumoral Ig remains substantially unknown.

In this study, we demonstrated for the first time that in-silico pairing of intratumoral Ig can be

used to obtain fully functional antibodies from legacy tumor RNA sequencing data. Moreover,

we showed that it is possible to use high-throughput proteomics techniques to identify their

target antigens, characterize their binding kinetics, and map the corresponding epitopes. These

steps are crucial to enable further functional characterization studies. Interestingly, we have

demonstrated that highly clonal intratumoral Ig are selected to bind not only cancer-specific

antigens (NY-ESO-1, MAGEA3, GAGE2A, DLL3), but also wild-type proteins expressed in the

tumor microenvironment (ANXA1, TGFBI, C4BPB). Despite being directed against

non-mutated self-antigens, these Ig bind to their target with very high-affinity, similarly to

antibodies obtained by immunizing a different species with the same antigen. The importance of

this observation is twofold: on one hand, it highlights the extent to which peripheral tolerance

might be compromised during immune responses in cancer-affected tissues; on the other hand, it

suggests that high-affinity fully human antibodies against human proteins can be obtained by
sequencing Ig transcripts expressed in tissues affected by chronic inflammation. While we have
not identified the target antigens for many of the antibodies produced in this study, it should be
noted that our antigen screening only considered non-mutated proteins. The remaining orphan
antibodies could be explained by errors in our in-silico pairing method. Alternatively, it is
possible that some of these orphan antibodies bind to antigens for which we could not screen in
this study, including neoantigens specific for particular patients, or non-protein antigens such as
glycans[20,21]. Advancements in antigen screening methods might in the future allow the
deorphaning of additional candidates. Despite these limitations, this study generated the largest
collection of individually screened fully human intratumoral Ig generated so far, paves the way
to improve our understanding of their functional role during anti-tumor immune responses, and
suggests a novel way to extract immunological insights from legacy RNA sequencing data.

Methods

Ig reconstruction. For the purpose of reconstruction of Ig chain sequences from RNA
sequencing data we implemented a computational workflow following the same basic idea
underlying previously described approaches [11,22]. We reduced the number of sequencing reads not
originating from Ig transcripts through three filtering steps. First, we selected only the reads that
either map to the Ig genes, or fail to map using Kallisto (version 0.44) [23] and Gencode [24] (version
22) protein coding sequences as the reference. Second, we mapped the remaining reads to the
full human genome (version GRCh38, [24,25]), and retained only the reads that mapped to the Ig
loci, or failed to map at all. In order to extract the selected reads and revert to FASTQ format we
used Sambamba [26] view (version 0.5.9) and Samtools [27] Fastq (version 1.8) commands. Third, we
depleted the reads originating from viral and bacterial sources by running kraken2[28] and retained
the unclassified reads. We assembled the retained reads using Trinity RNA Seq assembler
(version 2.8.4)[29] with custom parameters "--no_normalize_reads", "--max_chrysalis_cluster_size
100", and "--max_reads_per_graph 5000000". We then mapped the assembled sequences
germline V, D, and J gene regions using IgBLAST[28,30] (version 1.13). Sequences with high V
gene match scores (cutoff = 100) were kept as putative Ig chains.

Ig reconstruction performance evaluation. To evaluate the performance of this workflow and evaluate the possibility of identifying which heavy and light chain form a pair in cases where there is a strongly expressed clone we generated a synthetic benchmark using single cell sequencing data for sample PW2 from Lindman et.al.[11]. We estimated the distribution of Ig reads in TCGA data and found it to closely follow a lognormal model (Supplementary Figure 1). We then generated a set of 1000 synthetic samples. We generated each sample by sampling 25 numbers from this distribution and assigning them as the total number of Ig reads to 25 random B cells from the PW2 data. We then randomly selected the corresponding number of RNA sequencing reads from those mapped to Ig chains in each of the 25 B cells. To this mix we added 10 million reads randomly selected from a bulk RNA sequencing sample (TCGA-04-1348-01A), to act as background. In evaluation, we considered the reconstruction to be a success if all of the CDR regions (CDR1, CDR2, and CDR3) for both the heavy and light chain of the top clone were correct and the pair was correctly selected as the most abundant Ig.

Ig quantification, pairing, and somatic hypermutation analysis. For each processed TCGA sample we used Kallisto to quantify the amount of RNA sequencing reads originating from each of the putative Ig chain transcripts. We separately calculated the *Berger-Parker dominance index*[10] for both the heavy and light Ig chains, as the proportion of reads attributed to the most common chain of the corresponding type. We then calculated the dominance score for each sample as the geometric mean of the heavy and light chain Berger-Parker indices. Using the results of the synthetic benchmark we determined the threshold on this score (>0.382) above which we expect the sequence reconstruction to be accurate, and the top (most abundant) heavy and light chain to form a paired Ig. For each such Ig we assigned the putative isotypes by mapping the constant portions of the heavy chains (sections of the assembled sequences post the annotated J regions) to the Gencode v22 reference immunoglobulin C sequences. We used the IgBlast mappings to the germline segment to identify the mutated positions in each chain. For every position as defined by the Martin numbering scheme[31] we calculated the mutation frequency across all of the reconstructed immunoglobulin chains.

Processing of TCGA RNA sequencing data. TCGA BAM files were retrieved and processed in Cancer Genomics Cloud[32] (CGC). We applied sambamba-sort to sort the BAM files and Samtools fastq to convert them to FASTQ files, followed by Salmon quant in mapping-based

mode (GENOME v27 (GRCh38.p10) transcript assembly was used) with GC bias correction. We

obtained TPM values for 10,533 tumor samples and 730 normal samples across 33 TCGA cancer

types (Supplementary Table S3).

Statistical analysis of antigen arrays. Raw signal intensities (mean F635 values) were

correlated for background signal (mean B635 values) using the function background from R

package limma[33], with the correction method set to 'normexpr' with 'mle' parameter estimation

strategy. The replicate spots for the same target clone were summarized by calculating the

geometric mean followed by logarithmic transformation. We noticed that a number of spots

reported high signal intensities across multiple arrays likely due to non-specific binding. To

counter this issue we subtracted the mean signal value across all of the 172 arrays for each target

clone. We then centered and scaled the signal intensities and performed multiple testing

corrections by calculating the false discovery rate according to the Benjamini-Hochberg

method[33,34]. We used a stringent q-value cutoff of 0.01 in order to control the number of false

positive hits.

Antibody synthesis. The variable region of the antibodies were recombined with a constant

region of human IgG class I using AbAb's recombinant platform (Absolute Antibody Ltd,

Oxford, UK). The antibodies were expressed into HEK293 cells using the Absolute Antibody

transient expression system and purified by one-step affinity chromatography. We performed

Quality Control (QC) analysis to assess concentration, total amount, level of endotoxin, and

sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) (Supplementary Table

S1).

HuProt™ Protein arrays. We tested each candidate antibody using the human proteome

microarray HuProt™ 4.0, a comprehensive library of GST-tagged recombinant human protein

expressed in yeast. Briefly, the HuProt arrays were blocked with 5% BSA/1×TBS-T at room

temperature for 1 h. Then antibodies were incubated at 4 °C overnight, washed and labelled with

secondary antibodies prior detection.

Membrane Proteome Array (MPA) specificity testing. Membrane Proteome Array (MPA) is

Integral Molecular's cell-based array of ~6,000 human membrane proteins, each expressed in

live unfixed cells in separate wells of a 384-well plate[13]. In this study, the MPA was expressed in
HEK-293T cells 36h prior to testing. Each MAb was fluorescently labeled and added to the MPA
at a concentration optimized for the best signal-to-background ratio for target detection using an
independent immunofluorescence titration curve against membrane-tethered protein A. Binding
was measured by Intellicyt iQue3. Each 384-well plate contained positive (Fc-binding) and
negative (empty vector) controls to ensure plate-by-plate data validity. Hits were validated by
flow cytometry with serial dilutions of antibody, and the target identity was confirmed by
sequencing.

Surface Plasmon Resonance (SPR) assay. A Biacore 8K instrument, exploiting the SPR
principle, was used to calculate the equilibrium dissociation constant (K_D). Selected antibodies
(ligand) were individually immobilized to a sensor chip coated with Protein A to ensure the
correct orientation of the antibody while the antigens (analyte) were injected within the flow
stream at 30 µl/min. We used 90s association time for each experiment with 600s dissociation
time. Data analysis was performed using the R package *pbm*[35]. Sensorgram raw data time series
measurements were downloaded from the Biacore 8K instrument, and fitted to an appropriate
observation model from the pbm package using non-linear least squares parameter estimation
techniques. Models were selected parsimoniously; those which could adequately explain the data
using fewer parameters were generally preferred. In a small number cases, selected concentration
curves were excluded from the fitting procedure due to either instrumental measurement
anomalies, or apparent statistical outliers. Examples of qualifying anomalies included refractive
index bulk shift discontinuities[36] at the transition between the association and dissociation
measurement phases.

Epitope mapping. Hydrogen deuterium exchange mass spectrometry (HDX-MS) was used to
determine the epitope recognized by the antibodies. Linear peptides spanning the target antigen
length were incubated in a deuterium containing buffer in presence or absence of the relevant
antibody to observe differential exchange with hydrogen at the putative binding sites.

Data availability
The raw RNA sequencing reads used in this study are generated by the TCGA Research
Network: https://www.cancer.gov/tcga, and available through the TCGA data portal. All relevant

processed data are included as Source Data linked to this paper.

Code availability

255 The code for generation of the synthetic data used in the benchmarks, statistical analyses of the
256 arrays, and plots is from the authors available upon a reasonable request.

Authors contributions

GR, IDS, and BCT developed the sequence reconstruction workflow and generated the Ig chain
sequences. GR performed the performance evaluation of the workflow and analysed the protein
array data. ALS prepared supplementary figures depicting sensorgram plots for SPR assay
experiments. Milena P. performed statistical analysis comparing K_D values of human vs.
commercially available rabbit-derived antibodies. IG prepared the main figures. Milos P. and IG
processed the epitope mapping and generated the corresponding visualization. DB and DK
designed research and DB oversaw the implementation. GR, DB, DAL and DK prepared the
manuscript. All authors reviewed the final manuscript.

Competing interests

267 GR, IG, BCT, Milena P, Milos P, ALS, RR, DB and DK are participants in Totient's Equity
268 Incentive Plan. GR, IDS, BCT and DB are coinventors on patents assigned to Totient, Inc.

References

1.   Hu, X. *et al.* Landscape of B cell immunity and related immune evasion in human cancers.

Nat. Genet. **51**, 560–567 (2019).

2.   Zhao, J. *et al.* Author Correction: Immune and genomic correlates of response to anti-PD-1

immunotherapy in glioblastoma. *Nat. Med.* **25**, 1022 (2019).

3.   Bolotin, D. A. *et al.* Antigen receptor repertoire profiling from RNA-seq data. *Nat.*

Biotechnol. **35**, 908–911 (2017).

4.   De Silva, N. S. & Klein, U. Dynamics of B cells in germinal centres. *Nat. Rev. Immunol.* **15**,

137–148 (2015).

5. Helmink, B. A. *et al.* B cells and tertiary lymphoid structures promote immunotherapy

response. *Nature* **577**, 549–555 (2020).

6. Cabrita, R. *et al.* Author Correction: Tertiary lymphoid structures improve immunotherapy

and survival in melanoma. *Nature* **580**, E1 (2020).

7. Petitprez, F. *et al.* B cells are associated with survival and immunotherapy response in

sarcoma. *Nature* **577**, 556–560 (2020).

8. Siliņa, K. *et al.* Germinal Centers Determine the Prognostic Relevance of Tertiary

Lymphoid Structures and Are Impaired by Corticosteroids in Lung Squamous Cell

Carcinoma. *Cancer Res.* **78**, 1308–1320 (2018).

9. Weinstein, J. N. *et al.* The Cancer Genome Atlas Pan-Cancer analysis project. *Nature*

Genetics vol. 45 1113–1120 (2013).

10. Berger, W. H. & Parker, F. L. Diversity of Planktonic Foraminifera in Deep-Sea Sediments.

Science vol. 168 1345–1347 (1970).

11. Lindeman, I. *et al.* BraCeR: B-cell-receptor reconstruction and clonality inference from

single-cell RNA-seq. *Nat. Methods* **15**, 563–565 (2018).

12. Jeong, J. S. *et al.* Rapid identification of monospecific monoclonal antibodies using a

human proteome microarray. *Mol. Cell. Proteomics* **11**, O111.016253 (2012).

13. Tucker, D. F. *et al.* Isolation of state-dependent monoclonal antibodies against the

12-transmembrane domain glucose transporter 4 using virus-like particles. *Proc. Natl. Acad.*

Sci. U. S. A. **115**, E4990–E4999 (2018).

14. Chen, Y.-T. *et al.* Genomic cloning and localization of CTAG, a gene encoding an

autoimmunogenic cancer-testis antigen NY-ESO-1 to human chromosome Xq28.

Cytogenetic and Genome Research vol. 79 237–240 (1997).

15. Oliveira, L. C., Lima, A. M. N., Thirstrup, C. & Neff, H. F. *Surface Plasmon Resonance Sensors: A Materials Guide to Design, Characterization, Optimization, and Usage*. (Springer, 2019).

16. Weber, J., Peng, H. & Rader, C. From rabbit antibody repertoires to rabbit monoclonal antibodies. *Exp. Mol. Med.* **49**, e305 (2017).

17. Peng, H. *et al.* Mining Naïve Rabbit Antibody Repertoires by Phage Display for Monoclonal Antibodies of Therapeutic Utility. *J. Mol. Biol.* **429**, 2954–2973 (2017).

18. Masson, G. R. *et al.* Recommendations for performing, interpreting and reporting hydrogen deuterium exchange mass spectrometry (HDX-MS) experiments. *Nat. Methods* **16**, 595–602 (2019).

19. Fernández, J. A. & Griffin, J. H. A protein S binding site on C4b-binding protein involves beta chain residues 31-45. *J. Biol. Chem.* **269**, 2535–2540 (1994).

20. Rezende, S. M., Simmonds, R. E. & Lane, D. A. Coagulation, inflammation, and apoptosis: different roles for protein S and the protein S–C4b binding protein complex. *Blood* vol. 103 1192–1201 (2004).

21. Dube, D. H. & Bertozzi, C. R. Glycans in cancer and inflammation — potential for therapeutics and diagnostics. *Nature Reviews Drug Discovery* vol. 4 477–488 (2005).

22. Upadhyay, A. A. *et al.* BALDR: a computational pipeline for paired heavy and light chain immunoglobulin reconstruction in single-cell RNA-seq data. *Genome Med.* **10**, 20 (2018).

23. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

24. Harrow, J. *et al.* GENCODE: the reference human genome annotation for The ENCODE

Project. *Genome Res.* **22**, 1760–1774 (2012).

25. Schneider, V. A. *et al.* Evaluation of GRCh38 and de novo haploid genome assemblies

demonstrates the enduring quality of the reference assembly. *Genome Res.* **27**, 849–864

(2017).

26. Tarasov, A., Vilella, A. J., Cuppen, E., Nijman, I. J. & Prins, P. Sambamba: fast processing

of NGS alignment formats. *Bioinformatics* **31**, 2032–2034 (2015).

27. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,

2078–2079 (2009).

28. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.

Genome Biol. **20**, 257 (2019).

29. Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a

reference genome. *Nature Biotechnology* vol. 29 644–652 (2011).

30. Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin variable

domain sequence analysis tool. *Nucleic Acids Res.* **41**, W34–40 (2013).

31. Abhinandan, K. R. & Martin, A. C. R. Analysis and improvements to Kabat and structurally

correct numbering of antibody variable domains. *Mol. Immunol.* **45**, 3832–3839 (2008).

32. Lau, J. W. *et al.* The Cancer Genomics Cloud: Collaborative, Reproducible, and

Democratized-A New Paradigm in Large-Scale Computational Research. *Cancer Res.* **77**,

e3–e6 (2017).

33. Silver, J. D., Ritchie, M. E. & Smyth, G. K. Microarray background correction: maximum

likelihood estimation for the normal-exponential convolution. *Biostatistics* **10**, 352–363

(2009).

34. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*

(Methodological) vol. 57 289–300 (1995).

35. Davies, J. Protein Binding Models [R package pbm version 1.1.0]. (2018).

36. Schasfoort, R. B. M. Chapter 1:Introduction to Surface Plasmon Resonance. in *Handbook of*

Surface Plasmon Resonance 1–26 (2017).

Figures

Fig. 1 | Ig reconstruction and characterization protocol.

a. Steps of the computational workflow which starts from raw RNA sequencing data of tumor samples as the input, removes the reads mapped to non-Ig transcripts, reconstructs the Ig chain sequences, and outputs the paired sequences for which both chains satisfy the dominance threshold. **b**. Steps of the experimental workflow aimed at expressing the reconstructed Ig as recombinant antibodies, screening for their target antigens using two different human protein libraries, and confirming the result using surface plasmon resonance (SPR).

Fig. 2 | Evaluation of reconstruction performance on synthetic data.

a. Distribution of correctly and incorrectly reconstructed samples depending on the dominance score (geometric mean of the Berger-Parker indices for heavy and light Ig chains) estimated by the workflow. Above the threshold of 0.382 the top Ig for 90% of the synthetic samples were correctly reconstructed. **b**. ROC curve for the evaluation at different values of the dominance score. The red cross marks the dominance score of 0.382 selected as the point with the highest true positive rate (0.46) where the false positive rate is below 0.1.

Fig. 3 | Immunoglobulin isotypes, SHM, and cancer types

a. Distribution of the reconstructed Ig across TCGA cancer types. **b**. Distribution of amino acid lengths of IgH CDR3 regions. **c**. Number of selected Ig by putative isotype. **d.** and **e**. Mean SHM rate for each position in the Martin numbering scheme across the heavy and light chains in the selected set of Ig. For each chain, per amino-acid mutation rates were estimated by mapping the chains to germline sequence constructs by IgBlast and numbered following the Martin numbering scheme. For chains where multiple amino acids map to the same Martin number we used the mean value across those amino acids.

Fig. 4 | Distribution of K_D values for antibodies derived from intratumoral Ig and

commercially available antibodies against the same antigens.

a. Distribution of empirically determined K_D values for human-derived antibodies (Abs) and

their rabbit-derived counterparts for the same antigens, showing no statistically significant

difference using a paired t-test. As Abs have been tested in multiple experiments, to calculate the

p-value, we first averaged the $-\log10(K_D)$ across the experiments with a specific analyte

(different source of antigen), then paired each of these averages from human-derived Abs with

averages from rabbit-derived Abs for the same analyte and applied paired t-test. **b**. K_D values for

human and rabbit-derived antibodies presented in a. When multiple experiments have been

performed for a human or rabbit Ab, we calculated the average K_D from these experiments. For

detailed information about each experiment consult Supplementary Table S2. **c**. Representative

sensorgrams of SPR-determined antibody-antigen interaction for CYC214 (anti-C4BPB

antibody), for **d**. CYC066 (anti-MAGEA3 antibody) and for **e.** CYC168 (ant-TGFBI antibody).

Solid colored lines indicate raw data observed by Biacore 8K instrument, and overlaid solid

black lines indicate the fit result estimated using the model. For additional experimental details

and sensograms, see Supplementary Figure S3.

Fig. 5 | Epitope mapping results for C4BPB show overlap with protein S binding site.

HDX-MS was used to measure the level of deuterium (D) uptake by C4BPB alone or in presence

of CYC214 antibody. **a.** Relative D uptake difference per residue (blue shading) across the entire

protein surface. **b**. Details of the protein region containing the known binding site[20] for protein S.

High D uptake difference (dark blue) is detected in the region containing the binding site for

protein S, thus suggesting that CYC214 might be disrupting the interaction between C4BPB and

protein S. Each C4BPB fragment is measured up to three times and each residue can be covered

by one or more overlapping fragments: the uptake difference per residue was calculated using the

mean of the uptake differences of the fragments covering the residue.

Supplementary figures

Figure S1 | Distribution of Ig reads mapping to different chains in TCGA data follows a lognormal model.

The plot represents the distribution of the read counts mapped to each reconstructed Ig chain in the TCGA dataset. The data closely match a lognormal distribution with the estimated parameters mean=5.78, standard deviation=1.52.

Figure S2 | Distribution of Berger-Parker dominance score across different cancer types in the TCGA data

The plot represents the distribution of Berger-Parker dominance scores of the heavy and the light chains reconstructed from tumour samples in the TCGA dataset.

Figure S3 | Surface plasmon resonance assay sensorgram plots providing antibody-antigen equilibrium dissociation constants

Sensorgrams plots depict the SPR response of the Biacore 8K instrument in relative units (RU) vs. time, as commercially sourced antigens are flowed across surface-captured antibodies during an initial 90 s measurement interval, and then permitted to dissociate without antigen flowing, during a subsequent 600 s measurement interval. Antigens are prepared across a range of two-fold serially diluted concentrations, and within a given set of concentration curves comprising a single measurement attempt, data from all concentrations are fitted simultaneously using non-linear least squares regression to arrive at a single unifying K_D estimate (or a pair of K_D estimates, when a biphasic binding model provides a better fit) valid for that measurement attempt. Solid colored lines indicate raw data observed by the apparatus, and overlaid solid black lines indicate the fit result estimated using the model; a closer match between the two indicates a

better fit, with correspondingly superior explanatory power. In some cases, indicated by the absence of an overlaid solid black regression curve, data corresponding to one or more concentration curves may be discarded from the input to the regression model, if the result leads to a significantly better fit across the remaining concentration curves. Parameter estimates provided by the fit model are listed in the table at the top of each page, while lettered panels enumerate cases in which multiple replicate measurements were attempted for a given antibody-antigen pairing.

Figure S4 | Binding scores for top 5 hits from protein array screening.
Plots depict the scaled binding scores from top 5 hits from each of the protein arrays. Binding scores were calculated by fitting random effects models to the signal intestines, weighting each target by its mean rank across the arrays (to eliminate common promiscuous binders) and scaled by taking the z scores (Methods section). Asterisks mark hits with Benjamini and Hochberg false discovery rate < 0.01.

Figure S5 | Membrane proteome-wide specificity testing of recombinant antibodies. Each antibody was screened for binding against Integral Molecular's Membrane Proteome Array, consisting of ~6,000 human membrane proteins in their native state in unfixed cells. Antibody binding was detected by flow cytometry, and hits were defined as resulting in binding signals more than 3 standard deviations higher than background. Validated targets are shown in blue.

Supplementary tables

Table S1 | Antibody expression and antigen screening summary.
Table S2 | Biacore results.
Table S3 | Gene expression table TCGA (Salmon, TPM, no other normalization).
Table S4 | Protein array results.
Table S5 | Membrane proteome-wide screening results

a

Tumor samples
from patients

Immunoglobulin (Ig) reads
selected from bulk RNAseq

Light chains   Heavy chains

...   ...

Assembled Ig
chain sequences

Paired Ig
chain sequences

b

Antibody expression

Whole human
proteome screening

Human membrane
proteome screening

Hits confirmation with
SPR (Biacore)

Confirmed target
antigens

Fig. 1 l Antibody reconstruction and characterization protocol.

Fig. 2 I Evaluation of reconstruction performance on synthetic data.

Fig. 3 | Immunoglobulin isotypes, SHM, and chain pairing in the selected set of antibodies.

b

Antigen	Analyte	Human-derived Ab	K_D (Human-derived Ab), nM	Rabbit-derived Ab	K_D (Rabbit-derived Ab), nM
ANXA1	APrEST71575	CYC188	7.333	HPA011271	7.629
ANXA1	APrEST71576	CYC188	4.986	HPA011271	4.879
DLL3	9749-DL-050	CYC077	7.618	AB4315	8.002
GAGE2A	AR51161PU-S	CYC014	8.259	HPA043232	8.166
GAGE2A	AR51161PU-S	CYC025	8.583	HPA043232	9.178
MAGEA3	ab99958	CYC066	8.442	ab223162	9.784
TGFBI	TP300411	CYC168	7.777	HPA008612	6.481
TXLNA	TP310301	CYC086	6.957	HPA045383	6.764

Fig. 4 | Distribution of KD values for Totient antibodies and commercially available antibodies against the same antigens.

Fig. 5 | Epitope mapping results for C4BPB show overlap with protein S binding site.