1

Trophic resources of the edaphic mesofauna: a worldwide

review of the empirical evidence

Víctor Nicolás Velazco[1,2], Leonardo Ariel Saravia[3], Carlos Eduardo

Coviella[1,2], Liliana Beatriz Falco[1,2]

[1] Departamento de Ciencias Básicas, Universidad Nacional de Luján

(Argentina), [2] Instituto de Ecología y Desarrollo Sustentable (UNLu –

CONICET), [3] Instituto de Ciencias, Universidad Nacional de General

Sarmiento (Argentina)

Corresponding author:

Dr. Carlos E. Coviella

carlosecoviella@yahoo.com

voice: +541134456580

fax: +542323425795

22
23
24

25 Author contributions: LF and LS originally formulated the idea, NV, CC

26 developed the methodology, NV, CC, LF, and LS analyzed the data

27 and wrote the manuscript.

Abstract

Ecosystem sustainable use requires reliable information about structure and functioning. Accurate knowledge of trophic relations is central for the understanding of ecosystem dynamics, essential to develop sustainable use practices. This review gathers the information available worldwide about the trophic resources within the soil mesofauna. From over 130000 hits of the initial search on soil mesofauna, only a total of 78 published works relate particular species, genera, and families to particular trophic resources, the majority of them dealing with soils of the Paleartic region. Thirteen trophic resource categories were identified for six of the main orders of soil Acari and Collembola, using a combination of eight different methodologies. Out of 2717 records relating specific taxonomic groups to their trophic resources, 515 mention saprophytic fungi as a food resource, 381 cite nematodes, 308 mention bacteria, 286 litter and 256 cite mychorrizae. The available information is also highly skewed, for 68.16% of all the available information comes from Acari, and within these, 44.5% correspond just to Sarcoptiformes. For Collembola, the information available in the literature is much scarce the majority of it from Arthropleona. This review shows the general lack of information relating species, genera, and families of the soil mesofauna to specific trophic resources, with Collembolans largely unexplored in this regard. It also highlights the studies mostly come from European soils, with the use of trophic resources by the mesofauna of majority of the soils in most of the world still largely unknown.

53   Keywords: trophic web, soil sustainable use, soil mesofauna, Acari,

54   Collembola

Introduction

Only in the last few decades, sustainable use of the soil has come sharply into focus. The Ecosystem Millenium Assessment (2005), the report on Global Biodiversity and Ecosystem Services (IPBES 2019), and the recent State of Knowledge of Soil Biodiversity (FAO 2020) clearly state the importance of the soil ecosystem and the central role soil biodiversity plays. However, critical information is lacking on edaphic biodiversity and interaction webs for most soils.

The view about the edaphic biota has changed dramatically in recent years. Much of the work on taxonomic descriptions, richness, and biodiversity estimation, and microenvironments (Anderson 1978; Petersen and Luxton 1982), highlights the fragility and need for conservation of soils, and soil biodiversity (FAO 2020; Walker 1992). Gradually, the focus has shifted towards the study of the functioning of the soil ecosystem, in which the interactions between the biota and the edaphic environment affect its structure and functioning in a hierarchical manner, regulating the functioning of the edaphic ecosystem. (Barrios 2007; Behan-Pelletier and Newton 1999; Brussaard 1997; P. Lavelle et al. 2006) The relationships of species richness with ecosystem dynamics allow establishing causal relationships between the characteristics of the organisms present and the processes and services of the ecosystem (Hooper et al. 2005; Martín-López et al. 2007) in natural as well as in managed environments. (Wall et al. 2012).

The soil mesofauna is mainly constituted by microarthropods belonging to the Acari and Collembola subclasses that inhabit the upper soil horizons (Martinez and Narciso 2009), where complex communities are developed and maintained. These communities are influenced by a high microhabitats heterogeneity and the number of food resources available (Anderson 1977,

1978; Lavelle and Spain 1994), which provide a wide and varied set of ecological niches (Nielsen et al. 2010; Wallwork 1958a). The mesofauna, through its trophic relationships, contributes to the edaphic functioning through the processes of fragmentation of organic matter, the dynamics of the nutrient cycle, the transport of microflora propagules, and the regulation of microflora and microfauna populations that affect primary production. (Brussaard 1997; Cragg and Bardgett 2001; Lavelle 1996; Wall et al. 2012; FAO 2020).

The study of the flow of energy and matter in the soil ecosystem, groups edaphic organisms into functional guilds to understand the relationships with the functioning of the ecosystem. However, this approach leaves aside the understanding of how changes in the community influence the ecosystems (Thompson et al. 2012; Wall and Moore 1999). Thompson et. Al. (2012) suggest addressing this problem from the point of view of the trophic networks linking the flow of matter and energy and the diversity of a community. Thus, by studying the structure of the trophic networks, we can get a better understanding of the role of biodiversity in the functioning of the ecosystem. The methods used to obtain empirical evidence on trophic relationships of soil animals are generally used in both aquatic and terrestrial communities (JM Nielsen et al. 2018; Ruess and Chamberlain 2010; Tiunov 2007), but information about the latter is remarkably scarce in comparison.

Knowing the resources that microarthropods groups use as food is difficult due to their size and their cryptic environment. Furthermore, the observation of what an organism consumes at any given time does not indicate its potential diet (Nielsen et al. 2018). In this sense, the recognition of the food resources that are part of the diet of soil microarthropods must be based on empirical evidence, which constitutes the first step to establish the interactions between trophic species *sensu* Briand and Cohen (1984) and

the trophic resources that describe the trophic webs. (Briand and Cohen

1984; van Straalen 1998; Walter, Kaplan and Permar 1991).

Despite the increasing amount of descriptive works and lists of taxonomic

groups, the information available worldwide is still largely fragmentary and

incomplete, and the taxonomic resolution varies considerably between and

within published works. The same happens with trophic relationships of the

different taxonomic groups: information is still scarce. This data is necessary

to elucidate these relationships to analyze and build webs of interactions that,

in turn, could be used to assess the stability and conservation status of these

ecosystems. (Barrios 2007; Briones 2014; Thakur et al. 2020).

Thus, the objectives of this review are to gather all the information currently

available regarding the trophic resources used by mites and collembolans

that inhabit soils, and establish trophic relationships and potential diets of

these soil microarthropods.

Materials and methods

<u>Empirical evidence</u>

The recognition of the food resources that are part of the diet of soil

microarthropods is based on the empirical evidence provided by the

bibliography. These evidences are: [A] observations of consumption events

in microcosm experiments where feeding preferences are studied (Hubert et

al. 2001), selection of foods (Schneider 2005), physiological phenomena

related to feeding (Saur and Ponge 1988) or studying trophic interactions

(Schneider and Maraun 2009). [B] The observation of **intestinal contents**

assumes that what is actually observed is what is consumed (Hartenstein

1962; Schneider et al. 2004; Wallwork 1958a). [C] Morphology and

functioning **of mouthparts** related to food handling and processing (Buryn

and Brandl 1992; Kaneko 1988; Wallwork 1958b). [D] **Molecular analysis**

of intestinal content (Heidemann et al. 2014; King et al. 2008; Read et al.

2006). [E] Analysis of digestive **enzyme activity** (OC Nielsen 1962; Siepel

and Ruiter-Dijkman 1993). [F] Analysis of carbon and nitrogen **stable**

isotopes (Potapov, Tiunov, and Scheu 2019; Tiunov 2007). And, [G] the

study of **fatty acids** lipid profile as dietary markers (Pollierer, Scheu, and

Haubert 2010; Ruess and Chamberlain 2010).

General review publications are also included in this review, provided that

they collect information on the use of trophic resources from publications not

readily available.

Bibliography search and database construction

A systematic search was carried out through Google Scholar using keywords

related to the taxonomic groups of interest (mesofauna - collembola - acari),

trophic relationships (trophic - trophic interactions - predation - diet - soil -

food - soil food web), the empirical evidence (gut content - stable isotopes -

food preferences - digestive - enzymes - lipids - fatty acids), and their

combinations. The search yielded a total of 138,331 hits. This search was

then successively refined, to take into account only those publications that

provided evidence of the use of at least one trophic resource by soil mites

and/or springtails, and was further restricted to publications that refer to

specific levels of taxonomic resolution. The final database yielded a total of

78 citations (Electronic Supplementary Material, ESM I).

For the elaboration of the database from this search (ESM II), the taxonomic

information (Mayr 1996) and trophic resources were obtained, from the

papers and their published appendices (Thakur et al. 2020). To deal with the

heterogeneous information provided by the empirical evidence, it was

necessary to group the trophic resources to summarize their trophic and

ecological characteristics in the edaphic ecosystem. Once the trophic

resources were defined, associations were made between the taxa and

those resources.

Additional information can be found in the Electronic Supplementary

Material.

<u>Data analysis</u>

The records were then analyzed by counting the associations between the

taxa, the methods used and the trophic resources identified. We described

trophic resources, their number of records, and their relationships with the

different taxonomic levels. Then, the breakdown of each taxon was carried

out within the taxonomic categories and their relationship with trophic

resources. We calculated the proportions of the different methods and their

relationship with the taxonomic level and trophic resources.

Finally, we estimated the potential importance of resources on the diet for

the main orders of Acari and Collembola using the proportion of mentions

between trophic resources and taxa included in such orders, that is, the sum

of the proportion of mentions of a resource concerning the total mentions for

a particular taxon. For example, if a species has ten records of the

consumption of the same trophic resource, it is likely that the selection of this

food item is a reflection of the use of the resource. Then if we gathered the

information available from different species of the same genus, and the food

items consumed by them, then we could assign it to the potential diet for the

genus. Similarly, the resource use of the genera within a family can be

thought of as reflecting the potential diet of the family.

That is, the diet of higher taxonomic hierarchies will be constituted by the

sum of the resources used by the lower taxonomic levels.

The calculations, graphs, and tables were prepared using Excel and R.

Results

Only 21 articles from the 78 selected publications (ESM I) mention the countries in which the studies were carried out. Of these, 2 records belong to the Neotropical region, 3 to the Neartic region, 1 to the Australian region, 1 to the Eastern region, and the remaining 14 are all located in the Palearctic region. This is important to highlight because the species strongly vary depending on the biogeographic region it shows how incomplete still is the available information on this topic. It is also evident that knowledge about the trophic resources of Acari and Collembola was developed mainly in European countries and much work remains to be done worldwide. Only 49 of the publications mention the environments in which the studies were conducted. These authors analyzed 33 temperate forests, 3 tropical forests, 10 grasslands, 10 agroecosystems, and 1 desert. Regarding other environmental characteristics, 23 authors provide taxonomic information of the soil, 18 report on the pH, 10 mention the type of hummus. Other descriptions, environmental measures, and contributions of each bibliographic source can be found in ESM I.

Focusing on the use of trophic resources, a total of 2717 records were obtained from the empirical evidence (ESM II). To the species level, 127 records were Collembola species and 307 records were Acari species. The latter contributed 1848 of the 2717 total records (68.02%). At the genus level, 550 records were found, 21 genera belonging to Collembola and 115 genera belonging to Acari, out of a total of 53 and 234 genera of Collembola and Acari identified adding up genera and species within genera. The family level adds up to 240 records with 50 different families out of the total of 142 families identified in this review.

<u>Methods for identification of trophic resources used</u>

The method of observations in laboratory tests provide the main empirical

evidence from the database with 1082 records (Figure 1).

Figure 1. Methods used in the literature to assign trophic resources to
soil mesofauna taxa. Colors within columns refer to the number of
different taxonomic levels for which each method assigned at least one
resource. Assay: laboratory tests and observations. Isotopes: stable
isotopes. Gut: intestinal content. Enzymes: digestive enzymes. Mouth:
mouthparts morphology. Lipid: lipid profile. Mo. Det: molecular detection
of intestinal content. Reviews: general reviews by other authors.
232

Only 1.3% of the records describe feeding events for taxonomic levels higher

than family. Mesofauna families constitute 17% of the observations in

laboratory tests, with 85% of the total corresponding to the Prostigmata

suborder, within which 27 different families are mentioned. In turn, 35 of the

records correspond to the Prostigmata suborder that mentions 5 different

superfamilies (see ESM II). The evidence offered by observations in

laboratory tests adds up to 83% for species and genera. The orders

Mesostigmata (55.6%), Sarcoptiformes (26.7%) and Arthropleona (11.4%)

are the ones with the higher presence in the literature.

Stable isotopes follow in importance adding up to 500 records. From these,

75% of the records mention species. Genera and family add up to 22% of

the records, and the remainder of the taxa mentioned accumulate 3%. The

observation of the intestinal content reaches 17.6% of the records, in which

the species taxonomic level corresponds to 80% of the total records and the

genus level to 20%. For this method there were no records associated with

the order Mesostigmata, Acari. (see ESM II). The activity of digestive

enzymes (231 records) in all cases reports down to the species level. For

this empirical evidence the authors worked with the orders Sarcoptiforme

(Acari), and Arthropleona and Symphypleona (Collembola). Intestinal

contents analyses with molecular techniques for the detection of DNA is a

tool of recent development and represents 1.1% of the total records.

Figure 2 shows the number of records for each trophic resource, and the

empirical evidence that detects their relationships with them.

Laboratory tests mention the consumption of all thirteen trophic resources

described in the available literature. The mention of nematodes is the single

most important (225 records). Laboratory methods are the ones that most

often mention collembola, mites, invertebrate larvae, invertebrate eggs, and

enchytraeids as food resources. This method rarely mentions the

consumption of humus and carrion.

The detection of digestive enzymes (Figure 2) assigns 24% and 20% of its

contribution to saprophytic soil fungi and mycorrhizae, respectively. Litter

and plant tissue participate together with 33% and bacteria are mentioned

17.75% of the times. The study of the lipid profile of 22 species of Collembola

and 12 species of Acari, allowed assigning bacteria (33.6%), saprophytic

fungi (27.5%), leaf litter (17.6%,) and nematodes (16%) as the most

important trophic resources.

Figure 2. The number of records in the literature assigning each one of the
13 trophic resources to a mesofauna taxon as shown in Table 1. Colors in
the columns refer to the method used to assign those trophic resources to a
particular taxon. Methods as in Fig. 1.
274

Similar to laboratory tests, stable isotopes were associated with each and all

of the trophic resources, saprophytic fungi are the main resource mentioned

(114 times) followed by litter (93 mentions), bacteria (59 mentions), and

nematodes (54 mentions).

From highest to lowest, the number of mentions of resources provided by the

intestinal content analyses was: saprophytic fungi > mycorrhizae > bacteria

> litter > humus > plant tissue > nematodes > carrion > collembola - mites >

invertebrate eggs.

<u>Use of trophic resources by soil mesofauna</u>

Table 1 shows the trophic resources identified divided into 13 categories.

Trophic Resource	Description	Number of consumer records			
		Total allocations by trophic resource, absolute values (percentage values). (2717 records)[1]	FAMILY	GENERA	SPECIE
Saprophytic fungi	They are ubiquitous soil fungi that break down organic matter.[c]	515 (18,9)	16	93	396
Nematodes	Soil nematodes and protozoa, tardigrades, rotatory and other edaphic microfauna.[e]	381 (14)	45	115	211
Bacteria	They include bacteria with enormous autotrophic and heterotrophic capacities.[b]	308 (11,3)	9	63	227
Litter	Dead plant tissue accumulated in the soil with different degrees of fragmentation and decomposition.[a]	286 (10,5)	4	41	232
Mycorrhizae	Symbiotic fungal hyphae with plant roots. [i]	256 (9,4)	8	39	205
Plant tissue	Includes non-vascular plants (mosses, lichens, etc.), live roots, and seedlings.	174 (6,4)	1	15	154
Springtails	Juvenile and adult springtails.	171 (6,3)	47	42	73
Mites	Soft-bodied, non-sclerotic, unshielded, or juvenile mites.[f]	141 (5,2)	40	29	62
Larvae	Soft-bodied invertebrate larvae.	127 (4,7)	23	33	66
Humus	Complex and amorphous organic matter with a high degree of decomposition: debris, fecal pellets, etc.[d]	110 (4)	4	17	88
Invertebrate eggs	Invertebrate eggs found by predators.[g]	102 (4)	23	26	49
Enchytraeids	Anatomically homogeneous, soft-bodied oligochaete annelids.[c]	84 (3)	12	23	46
Invertebrate carrion	Animal tissue, molts, invertebrate corpses, etc.[h]	62 (2,3)	8	13	40

[1]Supplementary material II [2]Supplementary material I [a]B. Berg y McClaugherty 2008 [b]Clark 1971 [c]Warcup 1971 [d]Ponge 1991 [e]Persson et al. 1980 [f]Krantz et al. 2009 [g]Chernova et al. 2007 [h]Rusek 1998 [i]Schneider et. al. 2005

Table 1 Basic description of trophic resources. Total allocations by trophic resource,
absolute values (percentage values). Number of mentions of a trophic resource for the level
of taxonomic resolution. [1]Supplementary material II [2]Supplementary material I [a]Berg &
McClaugherty 2008 [b]Clark 1971 [c]Warcup 1971 [d]Ponge 1991 [e]Persson et al. 1980 [f]Krantz &
Walter 2009 [g]Chernova et al. 2007 [h]Rusek 1998 [i]Schneider et al. 2005

It can be seen that the main resource mentioned corresponds to saprophytic

fungi (18.9%) followed by nematodes (14%), bacteria (11.3%), and litter

(10.5%), the records for mites, collembolans, enchytraeidae, larvae, and

eggs accumulate 625 mentions (23.2%) and the lowest participation is for

invertebrate carrion (2.3%). Table 1 shows in bold the number of records (in

number and percentage) identified for each trophic resource, as well as the

participation of each taxonomic level for which a resource was identified.

Thus, for example, 515 total records (18.9%) of consumption of saprophytic

fungi were identified, 16 of which were identified at the family level, 93 at the

genus level, and 396 at the species level.

Interestingly, for the family level, the most numerous records correspond to

trophic resources such as collembola, nematodes, larvae, enchytraeids, and

mites, typical resources of predatory microarthropods that together add up

to 190 records. In the same way, the predation detected at the genus level

totals 268 records (48.82%). For the species taxonomic level, the main

singular resource mentioned is saprophytic fungi with 396 records (21.42%),

while total predation represents 27.42% (507 records).

Table 2 provides relevant information about how the consumption of the

same trophic resources is composed in the six orders mentioned in the

literature (3 orders of Collembola and 3 orders of Acari), identified according

to whether they are mentioned at the family, genus, or species level. The

number of times that each taxon is mentioned in the literature at the level of

family, genus, or species is first identified, and the column below shows the

mentions of resources consumed according to the records published for that

taxon. Within the family taxonomic category, 23 families were mentioned in

the order Mesostigmata (Acari). For the identity and more information about

the families, genera and species mentioned in Table 2, see ESM II.

Trophic Resource	FAMILY						GENERA						SPECIE					
	M (23)	S (77)	T (31)	A (8)	N (1)	Sy (2)	M (64)	S (151)	T (19)	A (43)	N (2)	Sy (8)	M (79)	S (223)	T (5)	A (114)	N (1)	Sy (12)
Saprophytic fungi	5	66	9	7	1	1	5	116	8	35	1	6	2	145	3	86	0	6
Nematodes	21	31	27	7	0	1	56	45	8	26	0	2	62	58	0	33	0	2
Bacteria	4	57	6	8	1	1	4	92	5	31	2	3	3	107	0	50	1	5
Litter	3	42	1	6	1	2	3	79	0	27	1	6	2	97	0	48	0	6
Mycorrhizae	1	54	6	7	0	1	1	83	6	33	0	2	0	91	3	58	0	2
Plant tissue	3	43	2	7	0	1	3	72	1	18	0	5	2	92	0	24	0	7
Springtails	15	2	25	4	0	0	35	2	11	7	0	0	50	2	1	7	0	0
Mites	14	2	25	2	0	0	35	2	5	4	0	0	47	2	1	4	0	0
Larvae	14	3	12	2	0	0	38	3	0	4	0	0	54	3	0	3	0	0
Humus	2	27	3	7	1	1	2	43	0	16	2	1	0	44	0	26	1	1
Invertebrate eggs	16	2	12	1	0	0	36	2	0	2	0	0	42	2	0	1	0	0
Enchytraeids	14	2	7	0	0	0	34	2	0	0	0	0	40	2	0	0	0	0
Invertebrate carrion	1	18	2	4	1	1	1	23	0	12	1	1	1	23	0	11	1	1

322 Table 2. Nested boxes represent the main orders of Acari and Collembola. In parentheses the
323 total number of taxa for the taxonomic category. Within the cells, the number of taxa associated
324 with the trophic resource is shown.
325

At all the taxonomic levels considered, the order Sacroptiformes (Acari) is

mentioned consuming mainly saprophytic fungi, bacteria and mycorrhizae.

The records for the order Trombidiformes (Acari), show them mainly as

predators, with 27 families consuming nematodes, 25 families consuming

springtails, the same number for mites, and 12 families using invertebrate

eggs and larvae. The use of litter, plant tissue and hummus were only

associated with 1, 2 and 3 families of Trombidiformes respectively.

For Collembola, the information available on the trophic resources used is

much scarce. At the family level, the order Arthropleona is mentioned in only

8 records, Neelipleona in only one record, and Symphypleona in only two

records.

Considering the mentions at the genus level, the order Mesostigmata (Acari)

is mentioned consuming nematodes in 56 of the 64 genera compiled for this

order. Likewise, 38 genera consume invertebrate larvae, 35 genera consume

mites or collembola, and 34 genera make use of the enchytraeids as a food

resource. This information indicates predation as the main feeding strategy

for this order of Acari. Saprophytic fungi, bacteria, plant tissue, and litter were

only mentioned by 5, 4, 3 and 3 genera respectively.

The information available at the species level for the order Symphypleona

(Collembola), registered mentions for 12 different species, 7 of them

provided evidence of consumption of plant tissue, 6 species consume

saprophytic fungi and leaf litter, 5 consume bacteria and two species were

associated with nematodes.

For Arthropleona (Collembola) the information at the species level indicates

that they mainly consume saprophytic fungi with 86 of the 114 species

records registered for this order mentioning this resource. Mycorrhizae,

bacteria, and litter follow in importance. Nematodes are mentioned as a

resource for 33 species. Adding up all the records at the three taxonomic

levels (family, genus and species), the order Sarcoptiformes (Acari) accounts

for 44.5% of the total records. Table 2 indicates that the literature mentions

trophic resources for 77 families, 151 genera, and 223 species within this

order. Ordering the information collected at the highest level of taxonomic

resolution (species), the mentions of trophic resources for this order are:

fungi > saprophytes > bacteria > litter > plant tissue > mycorrhizae. The

nematode resource was associated with 58 species, hummus with 44

species, and carrion was recorded as a resource for 23 species. The

associations that correspond to typical predator resources do not exceed 3

associations with species, except for nematodes.

Figure 3 is a bipartite diagram in which the thickness of each line gives

visually information about the proportion in which the resources are

associated with the main orders of microarthropods, according to the detail

provided in Tables 1 and 2. The saprophytic fungus resource is the main

constituent of the Arthropleona diet, along with bacteria, mycorrhizae and

litter. Trombidiformes have a diet mainly based on microarthropods and

nematodes and to a lesser extent saprophytic fungi.

Figure 3. Bipartite graph showing the use of trophic resources by the main orders of Acari and Collembola. Upper nodes: Acari and Collembola orders as in Table 2. Neelipleona with only 8 records excluded. Lower nodes: trophic resources. The thickness and intensity of the lines give an idea of the proportion of mentions in the available literature about their use of trophic resources. Resource names as in Tables 1 and 2.

The Sarcoptiformes diet is mainly of saprophytic fungi, leaf litter, bacteria, and plant tissue. Nematodes are the main items mentioned as a secondary resource. Mesostigmata have a basically predatory diet, consisting mainly of nematodes, microarthropods, eggs, and worm-shaped prey. For the order Symphypleona, the diet is, from greater to lesser number of associations: plant tissue > saprophytic fungi > bacteria > litter.

Discussion

Until the 1960s, soil fauna considered only worms, and terrestrial ecologists considered the rest of the soil fauna as a "black box" of decomposers and detritivores (Briones, 2014). The information gradually collected over the following decades regarding the edaphic mesofauna is reaching the point

where it is possible to focus on integrative works. However, the available

information is still scarce and mostly restricted to the most numerous or

conspicuous groups of the soil mesofauna.

This review gathers the information currently available regarding the use of

trophic resources by the edaphic mesofauna. This information, together with

other characteristic features of the soil mesofauna, will allow building a food

web interactions network, that could result in a better understanding of the

structure and functioning of the edaphic biota (FAO 2020). Trophic networks

will, in turn, allow for comparing the state of different soils, or the same soils

under different intensities of anthropic impact (Pankhurst, Doube, and Gupta

1997; Pey et al. 2014; Thompson et al. (2012).

Estimating the diet of organisms provides the foundation for understanding

trophic interactions by characterizing how trophic resources are used by

consumers within the soil ecosystem. Furthermore, it provides much-needed

information on their trophic niches, their nutritional physiology, and the flows

of matter and energy throughout the soil ecosystem (JM Nielsen et al. 2018;

Potapov et al. 2020). The empirical characterization of trophic interactions is

challenging due to the spatial and temporal complexity of feeding patterns,

and the limitations of the methods to identify and quantify the components of

the diet (JM Nielsen et al. 2018; Pankhurst, Doube, and Gupta 1997; Walter,

Kaplan, and Permar 1991).

For soil microarthropods, the evidence provided by laboratory work results

in the most relevant approximation to know how these animals use trophic

resources, their feeding behavior, their food preferences, and their

development and growth in controlled environments. The results presented

here show the importance of this method to better understand the use of

trophic resources by these cryptic soil organisms. The method used to relate

the isotopic signature of organisms and their resources is a recently

developed tool that is useful for detecting the importance and the changes in time or space of assumed trophic relationships, which means that we have to know in advance if the trophic relationship exists. This method can analyze a large number of species and an important variety of resources (Potapov, Tiunov, and Scheu 2019). The use of methods that allow tracing biomolecules along the food chain is advancing towards the development of strategies to interpret soil bulk volume analyzes. (Kühn, Schweitzer, and Ruess 2019; Ruess et al. (2007).

The trophic resources used by soil microarthropods can be determined in a complementary way by using different methods (Potapov et al. 2020; Walter, Kaplan, and Permar 1991). The most used resources are saprophytic fungi, nematodes, and bacteria. If we associate them with their nutritional characteristics, these trophic resources are rich in molecules with great nutritional value as determined in the dietary routes labeled by fatty acids, stable isotopes, or the ubiquitous enzymatic pathways of carbohydrases that hydrolyze nonstructural polysaccharides. (OC Nielsen 1962; Potapov, Tiunov, and Scheu 2019; Ruess and Chamberlain 2010).

We found that a large proportion of the resources are defined as taxonomic categories of species, genera, and families, which would be important to estimate the diets of higher taxonomic groups. The available information for low taxonomic levels could be used as a reference to address the problem of what the use of soil resources will be like by higher level taxonomic groups (Bedano 2007).

This information must be interpreted with caution, however, because within a taxonomic category, each species could apply different strategies to the problem of exploiting resources (Lavelle and Spain 2001; Moore, Walter, and Hunt 1988), as many taxa in the available literature are considered to be general consumers. We allowed a closer look at the complexity of the

interaction networks between the main orders of Acari and Collembola and

the trophic resources present in the edaphic environment. Furthermore, this

information suggests that the edaphic ecosystem might have a high level of

homeostasis because of the high number of redundant species that exist in

the edaphic system (Hättenschwiler, Tiunov, and Scheu 2005).

Moreover, the results presented here provide valuable new information

about the different feeding strategies of the main groups of the soil

mesofauna. At the same time, the still quite scant information available in this

regard must be highlighted. Of the total number of publications retrieved in

the bibliographic search, we found only 78 works providing information on

the relationships between taxa of the edaphic mesofauna and their trophic

resources. Even more scarce is the geographic information, with only 21

papers mentioning the country of origin of the data. The great majority of

studies are from countries of the Palearctic region focusing on temperate

forest soils. Likewise, it is illustrative to note that most of the works have been

carried out in European countries.

It is also to be noted that the information available for Acari and Collembola

is strongly asymmetric. Out of the total of 2717 records retrieved in the

bibliographic search, 1848 (68.16%) correspond to Acari, and within this

number, the majority (44.5%) correspond to Sarcoptiformes.

Finally, it is necessary to call attention to the need for more studies on the

trophic relationships of the soil mesofauna. Out of a total of approximately

9000 described species of Collembola (Bellinger et al. 2020), it was only

possible to find trophic information for just 127 soil inhabitant species. For

Acari, out of the approximately 58,000 species described (Schmidt 2020), it

was only possible to find references of trophic relationships for 307 soil

species.

Further deepening on the knowledge of functional and trophic relationships

of the soil the fauna would allow for a better and more precise evaluation of

the functioning of the edaphic ecosystem, the protection of the ecosystem

services that the soil mesofauna provides, and the sustainable use of the

soil.

Declarations

This work was partially funded by a scholarship from Consejo Nacional de

Ciencia y Tecnología (CONICET), Universidad Nacional de Luján, and

Instituto de Ecología y Desarrollo Sustentable (INEDES), Argentina. The

authors declare no conflicts of interest. All the data for this work not shown

in here, is available as Electronic Supplementary Material and referred to in

the text when appropriate.

Conflict of Interest: The authors declare they have no conflict of interest.

Bibliography

Anderson JM (1978) A method to quantify soil-microhabitat complexity and its application to a study of soil animal species diversity. Soil Biol. Biochem. 10, 77–78. https://doi.org/10.1016/0038-0717(78)90014-7

Anderson JM (1977) The Organization of Soil Animal Communities. Ecol. Bull. 25, 15–23.

Barrios E (2007) Soil biota, ecosystem services and land productivity. Ecol. Econ. 64, 269–285. https://doi.org/10.1016/j.ecolecon.2007.03.004

Bedano JC (2007) El rol de la mesofauna edáfica en la evaluación de la calidad del Suelo, In: De La Biología de Los Suelos a La Agricultura. Universidad Nacional de Río Cuarto, pp. 247–258.

Behan-Pelletier V, Newton G (1999) Linking soil biodiversity and ecosystem function. The taxonomic dilemma. BioScience 2. https://doi.org/10.2307/1313540.

Bellinger PF, Christiansen KA, Janssens F (2020). Internet resource: http://www.collembola.org. Last updated: November 30, 2020. Last accessed: December 23, 2020.

Berg B, McClaugherty C (2008) Plant litter: decomposition, humus formation, carbon sequestration, 2nd. ed. ed. Springer, Berlin.

Briand F, Cohen JE (1984) Community food webs have scale-invariant structure. Nature, Letters to Nature 307, 264–267.

Briones MJI (2014) Soil fauna and soil functions: a jigsaw puzzle. Front. Environ. Sci. 2, 1–22. https://doi.org/10.3389/fenvs.2014.00007

Brussaard L (1997) Biodiversity and Ecosystem Functioning in Soil. Ambio. 26, 563–570.

Buryn R, Brandl R (1992) Are the morphometrics of chelicerae correlated with diet in mesostigmatid mites (Acari)? Exp. Appl. Acarol. 14, 67–82. https://doi.org/10.1007/BF01205353

Chernova NM, Bokova AI, Varshav EV, Goloshchapova NP, Savenkova

YuYu (2007) Zoophagy in Collembola. Entomol. Rev. 87, 799–811.

https://doi.org/10.1134/S0013873807070020

Clark FE (1971) Bacterios del Suelo, in: Burges, A., Raw, F. (Eds.), Biología

del Suelo, Chapter 2. Ediciones Omega, Barcelona, pp. 27–68.

Cragg RG, Bardgett RD (2001) How changes in soil faunal diversity and

composition within a trophic group influence decomposition processes. Soil

Biol. Biochem. 33, 2073–2081. https://doi.org/10.1016/S0038-

0717(01)00138-9

FAO, ITPS, GSBI, SCBD, EC. (2020) State of knowledge of soil biodiversity.

Status, challenges and potentialities, Report 2020. Rome, FAO.

https://doi.org/10.4060/cb1928en

Hartenstein R (1962) Soil Oribatei. I. Feeding Specificity among Forest Soil

Oribatei (Acarina). Ann. Entomol. Soc. Am. 55, 202–206.

https://doi.org/10.1093/aesa/55.2.202

Hättenschwiler S, Tiunov AV, Scheu, S, (2005) Biodiversity and Litter

Decomposition in Terrestrial Ecosystems. Annu. Rev. Ecol. Evol. Syst. 36,

191–218. https://doi.org/10.1146/annurev.ecolsys.36.112904.151932

Heidemann K, Ruess L, Scheu S, Maraun M, (2014) Nematode consumption

by mite communities varies in different forest microhabitats as indicated by

molecular gut content analysis. Exp. Appl. Acarol. 64, 49–60.

https://doi.org/10.1007/s10493-014-9807-x

Hooper DU, Chapin FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton

JH, Lodge DM, Loreau M, Naeem S, Schmid B, La HS, Symstad AJ,

Vandermeer J, Wardle DA (2005) Effects of biodiversity on ecosystem

functioning: A consensus of current knowledge. Ecol. Monogr. 75, 33.

https://doi.org/10.1890/04-0922

Hubert J, Žilová M, Pekár, S (2001) Feeding preferences and gut contents

of three panphytophagous oribatid mites (Acari: Oribatida). Eur. J. Soil Biol.

37, 197–208. https://doi.org/10.1016/S1164-5563(01)01083-4

IPBES (2019) Global assessment report on biodiversity and ecosystem

services of the Intergovernmental Science-Policy Platform on Biodiversity

and Ecosystem Services. Brondizio ES, Settele J, Díaz S, Ngo HT (eds).

IPBES secretariat, Bonn, Germany.

Kaneko N (1988) Feeding habits and cheliceral size of oribatid mites in cool

temperate forest soils in Japan. Rev. Écologie Biol. Sol 25, 353–363.

King RA, Read DS, Traugott M, Symondson WOC (2008) Molecular analysis

of predation: A review of best practice for DNA-based approaches. Mol. Ecol.

17, 947–963. https://doi.org/10.1111/j.1365-294X.2007.03613.x

Krantz GW, Walter DE (Eds.) (2009) A manual of acarology, 3rd ed. Texas

Tech University Press, Lubbock, Texas.

Kühn J, Schweitzer K, Ruess L (2019) Diversity and specificity of lipid

patterns in basal soil food web resources. PLOS ONE 14, e0221102.

https://doi.org/10.1371/journal.pone.0221102

Lavelle P (1996) Diversity of soil fauna and ecosystem function. Biol.

International 33, 3–16.

Lavelle P, Decaëns T, Aubert M, Barot S, Blouin M, Bureau F, Margerie P,

Mora P, Rossi JP (2006) Soil invertebrates and ecosystem services. Eur. J.

Soil Biol. 42, S3–S15. https://doi.org/10.1016/j.ejsobi.2006.10.002

Lavelle P, Spain AV (2001) Soil Ecology, 2nd. ed. Kluwer Academic

Publisher. Springer.

Martínez PA, Narciso EN (2009) Mesofauna. In: Momo, F.R., Falco, L.B.

(Eds.), Biología y Ecología de la Fauna del Suelo. Imago Mundi, Buenos

Aires, p. 186.

Martín-López, B, González JA, Díaz S, Castro I, García-Llorente M (2007)

Biodiversidad y bienestar humano: el papel de la diversidad funcional.

Ecosistemas 16, 69–80.

Mayr E (1996) What Is a Species, and What Is Not? Philos. Sci. 63, 262–

277. https://doi.org/10.1086/289912

Moore JC, Walter, DE, Hunt, HW (1988) Arthropod Regulation of Micro- and

Mesobiota in Below-Ground Detrital Food Webs. Annu. Rev. Entomol. 33,

419–435. https://doi.org/10.1146/annurev.en.33.010188.002223

Nielsen CO (1962) Carbohydrases in Soil and Litter Invertebrates. Oikos 13,

200–2015. https://doi.org/10.2307/3565085

Nielsen JM, Clare EL, Hayden B, Brett MT, Kratina P (2018) Diet tracing in

ecology: Method comparison and selection. Methods Ecol. Evol. 9, 278–291.

https://doi.org/10.1111/2041-210X.12869

Nielsen UN, Osler GHR, Campbell CD, Neilson R, Burslem DFRP, van der

Wal R (2010) The Enigma of Soil Animal Species Diversity Revisited: The

Role of Small-Scale Heterogeneity. PLoS ONE 5, e11567.

https://doi.org/10.1371/journal.pone.0011567

Pankhurst C, Doube BM, Gupta VVSR (1997) Biological Indicators of Soil

Health. CAB INTERNATIONAL.

Persson T, Bååth E, Clarholm M, Lundkvist H, Söderström BE, Sohlenius B

(1980) Trophic Structure, Biomass Dynamics and Carbon Metabolism of Soil

Organisms in a Scots Pine Forest. Ecol. Bull. 32, 419–459.

Petersen H, Luxton, M (1982) A Comparative Analysis of Soil Fauna

Populations and Their Role in Decomposition Processes. Oikos 39, 288.

https://doi.org/10.2307/3544689

Pey B, Nahmani J, Auclerc A, Capowiez Y, Cluzeau D, Cortet J, Decaëns T,

Deharveng L, Dubs F, Joimel S, Briard C, Grumiaux F, Laporte MA, Pasquet

A, Pelosi C, Pernin C, Ponge JF, Salmon S, Santorufo L, Hedde M (2014)

Current use of and future needs for soil invertebrate functional traits in

community ecology. Basic Appl. Ecol. 15, 194–206.

https://doi.org/10.1016/j.baae.2014.03.007

Pollierer MM, Scheu S, Haubert D (2010) Taking it to the next level: Trophic

transfer of marker fatty acids from basal resource to predators. Soil Biol.

Biochem. 42, 919–925. https://doi.org/10.1016/j.soilbio.2010.02.008

Ponge JF, (1991) Food resources and diets of soil animals in a small area of

Scots pine litter. Geoderma 49, 33–62. https://doi.org/10.1016/0016-

7061(91)90090-G

Potapov AM, Pollierer MM, Salmon S, Šustr V, Chen TW (2020)

Multidimensional trophic niche approach: gut content, digestive enzymes,

fatty acids and stable isotopes in Collembola. bioRxiv 2020.05.15.098228.

https://doi.org/10.1101/2020.05.15.098228

Potapov AM, Tiunov AV, Scheu S (2019) Uncovering trophic positions and

food resources of soil animals using bulk natural stable isotope composition:

Stable isotopes in soil food web studies. Biol. Rev. 94, 37–59.

https://doi.org/10.1111/brv.12434

Read DS, Sheppard SK, Bruford MW, Glen DM, Symondson WOC (2006)

Molecular detection of predation by soil micro-arthropods on nematodes.

Mol. Ecol. 15, 1963–1972. https://doi.org/10.1111/j.1365-

294X.2006.02901.x

Ruess L, Chamberlain PM (2010) The fat that matters: Soil food web analysis

using fatty acids and their carbon stable isotope signature. Soil Biol.

Biochem. 42, 1898–1910. https://doi.org/10.1016/j.soilbio.2010.07.020

Ruess L, Schütz K, Migge-Kleian S, Häggblom MM, Kandeler E, Scheu S

(2007) Lipid composition of Collembola and their food resources in

deciduous forest stands—Implications for feeding strategies. Soil Biol.

Biochem. 39, 1990–2000. https://doi.org/10.1016/j.soilbio.2007.03.002

Rusek J, (1998) Biodiversity of Collembola and their functional role in the

ecosystem. Biodivers. Conserv. 7, 1207–1219.

https://doi.org/10.1023/A:1008887817883

Saur É, Ponge J (1988) Alimentary studies on the Collembolan

Paratullbergia callipygos using transmission electron microscopy.

Pedobiologia 31, 355–379.

Schmidt KH (2020) Internet resource: http://miteresearch.org. Last

accessed: December 23, 2020.

Schneider K (2005) Feeding biology and diversity of oribatid mites (Oribatida,

Acari). PhD dissertation.

Schneider K, Maraun M (2009) Top-down control of soil microarthropods -

Evidence from a laboratory experiment. Soil Biol. Biochem. 41, 170–175.

https://doi.org/10.1016/j.soilbio.2008.10.013

Schneider K, Renker C, Maraun M, (2005) Oribatid mite (Acari, Oribatida)

feeding on ectomycorrhizal fungi. Mycorrhiza 16, 67–72.

https://doi.org/10.1007/s00572-005-0015-8

Schneider K, Renker C, Scheu S, Maraun M, (2004) Feeding biology of

oribatid mites: a minireview. Phytophaga 14, 247–256.

Siepel H, Ruiter-Dijkman EM (1993) Feeding guilds of oribatid mites based

on their carbohydrase activities. Soil Biol. Biochem. 25, 1491–1497.

https://doi.org/10.1016/0038-0717(93)90004-U

Thakur MP, Phillips HRP, Brose U, De Vries FT, Lavelle P, Loreau M,

Mathieu J, Mulder C, Van der Putten WH, Rillig MC, Wardle DA, Bach EM,

Bartz MLC, Bennett JM, Briones MJI, Brown G, Decaëns T, Eisenhauer N,

Ferlian O, Guerra CA, König-Ries B, Orgiazzi A, Ramirez KS, Russell DJ,

Rutgers M, Wall DH, Cameron EK (2020) Towards an integrative

understanding of soil biodiversity. Biol. Rev. 95, 350–364.

https://doi.org/10.1111/brv.12567

Thompson RM, Brose U, Dunne JA, Hall RO Jr, Hladyz S, Kitching RL,

Martinez ND, Rantala H, Romanuk TN, Stoufer DB, Tylianakis JM (2012)

Food webs: reconciling the structure and function of biodiversity. Trends

Ecol. Evol. 27, 689–697. http://dx.doi.org/10.1016/j.tree.2012.08.005

Tiunov AV (2007) Stable isotopes of carbon and nitrogen in soil ecological

studies. Biol. Bull. 34, 395–407.

https://doi.org/10.1134/S1062359007040127

van Straalen NM (1998) Community structure of soil arthropods as a

Bioindicator of Soil Healt, in: Pankhurst, C., Doube, B.M., Gupta, V.V.S.R.

(Eds.), Biological Indicators of Soil Health. CAB INTERNATIONAL, U.K., pp.

235–264.

Walker BH (1992) Biodiversity and ecological redundancy. Conserv. Biol. 6,

18–23. https://doi.org/10.1046/j.1523-1739.1992.610018.x

Wall DH, Bardgett RD, Behan-Pelletier V, Herryck JE, Jones HT, Ritz K, Six

J, Strong DR, Van der Putten WH (2012) Soil ecology and ecosystem

services, Journal of Chemical Information and Modeling. Oxford University

Press.

Wall DH, Moore JC (1999) Interactions underground. Soil biodiversity,

mutualism, and ecosystem processes. BioScience 49, 109–117.

https://doi.org/10.2307/1313536

Wallwork JA (1958) Notes on the Feeding Behaviour of Some Forest Soil

Acarina. Oikos 9, 260–271. https://doi.org/10.2307/3564770

Walter DE, Kaplan DT, Permar TA, (1991) Missing links: a review of methods

used to estimate trophic links in soil food webs. Agric. Ecosyst. Environ. 34,

399–405. https://doi.org/10.1016/0167-8809(91)90123-F

Warcup JH (1971) Hongos del Suelo, In: Burges, A., Raw, F. (Eds.), Biología

del Suelo. Chapter 3. Ediciones Omega, Barcelona, pp. 27–68.